RNAi-based screen for pigmentation in *Drosophila melanogaster* reveals regulators of brain dopamine and sleep

Samantha L Deal[1,2,#], Danqing Bei[2,3], Shelley B Gibson[2,3], Harim Delgado-Seo[2,4], Yoko Fujita[2,3], Kyla Wilwayco[2,3], Elaine S Seto[5,6,7], Shinya Yamamoto[1,2,3,4,8,*]

[1] Program in Developmental Biology, Baylor College of Medicine (BCM), Houston, TX 77030, USA

[2] Jan and Dan Duncan Neurological Research Institute (NRI), Texas Children's Hospital (TCH), Houston, TX 77030, USA

[3] Department of Molecular and Human Genetics, BCM, Houston, TX 77030, USA

[4] Department of Neuroscience, BCM, Houston, TX 77030 USA

[5] Department of Neurology, TCH, Houston, TX 77030, USA

[6] Epilepsy Center, TCH, Houston, TX 77030, USA

[7] Department of Neurophysiology, TCH, Houston, TX 77030, USA

[8] Development, Disease Models & Therapeutics Graduate Program, BCM, Houston, TX 77030, USA

[#]Current Address: Department of Neuroscience, University of Pennsylvania, Philadelphia, Pennsylvania, USA

*Corresponding Author: Shinya Yamamoto

Correspondence: yamamoto@bcm.edu

Summary

The dopaminergic system has been extensively studied for its role in behavior in animals as well as human neuropsychiatric and neurological diseases. However, we still know little about how dopamine levels are tightly regulated *in vivo*. To identify novel regulators of dopamine, we utilized *Drosophila melanogaster* cuticle pigmentation as a readout, where dopamine is a precursor to melanin. We measured dopamine from genes known to be critical for cuticle pigmentation and performed an RNAi-based screen to identify new regulators of pigmentation. We found 153 potential pigmentation genes, which were enriched for conserved homologs and disease-associated genes as well as developmental signaling pathways and mitochondria-associated proteins. From 35 prioritized candidates, we found 10 caused significant reduction in head dopamine levels and one caused an increase. Two genes, *clueless* and *mask (multiple ankyrin repeats single KH domain),* upon knockdown, reduced dopamine levels in the brain. Further examination suggests that Mask regulates the transcription of the rate-limiting dopamine synthesis enzyme, *tyrosine hydroxylase*, and its knockdown causes dopamine-dependent sleep phenotypes. In summary, by studying genes that affect cuticle pigmentation, a phenotype seemingly unrelated to the nervous system, we were able to identify several genes that affect dopamine metabolism as well as a novel regulator of behavior.

Keywords: *Drosophila melanogaster*, dopamine, pigmentation, RNAi, screen, *mask*, sleep

Introduction

Dopamine is a conserved neurotransmitter that functions in the nervous system to regulate a variety of behaviors, including learning & memory, locomotion, reward, and sleep. In humans, disruptions in dopamine have been associated with neurological and neuropsychiatric disorders, including addiction, depression, sleep disorders, and schizophrenia[1–3]. In addition, one of the hallmarks of Parkinson's disease is early degeneration of dopaminergic neurons, and Parkinson's patients are often given a dopamine precursor, l-dihydroxyphenylalanine (L-DOPA), as a method for treating symptoms [4,5]. Most genetic research on dopamine biology has focused on dopaminergic signaling and transport; however, changes in dopamine levels can have a great impact on behavioral outcomes [6–8]. Thus, identifying genes that affect dopamine levels is critical to understanding and potentially treating dopamine-associated disorders.

Dopamine synthesis is a conserved process, whereby the amino acid tyrosine is converted into L-DOPA via the rate-limiting enzyme, Tyrosine hydroxylase (TH) [9]. Then, L-DOPA is converted into dopamine via the enzyme Dopa decarboxylase (Ddc) [10]. In mammals, dopamine is degraded via oxidation or methylation using the enzymes Monoamine oxidase (MAO) and Catechol-O-methyltransferase (COMT), respectively [11]. In invertebrates, dopamine degradation goes through β-alanylation via the β-alanyl amine synthase (encoded by the *ebony* gene in *Drosophila melanogaster*) and acetylation via the acetyltransferase (encoded by the *speck* gene in *Drosophila*) [12]. However, insects also convert dopamine into melanin to form proper cuticle pigment and structure. In fact, changes in genes that are involved in dopamine metabolism also change the cuticle color [13–15]. In other words, knockdown of *TH* (or the *pale (ple)* in *Drosophila*) or *Ddc* leads to a paler cuticle, and mutations or knockdown of *ebony* or *speck* leads to a darker cuticle.

Historically, *Drosophila* has been used to study dopamine biology in multiple contexts,

including learning and memory (16,17). Over the years, an extensive genetic toolkit has been

generated in the fruit fly, making it a great system for dissecting dopaminergic neural circuitry and

signaling (18–20). In addition to the fly's contribution to understanding dopaminergic circuitry, this

model organism helped to identify and investigate the cellular functions of Parkinson disease's

genetic risk factors, *PINK1* and *PRKN* (i.e. *Pink1* and *parkin* in the fruit fly) (21). The conservation

of systems in neural biology along with the efficiency of working in *Drosophila* has made it a great

system for large-scale screening of novel genes. Previously, a large-scale RNAi screen (11,619

genes, ~89% of genome) was performed to identify genes that affect mechanosensory organs

and cuticle formation in the *Drosophila* thorax (22). In this screen, the researchers found 458

genes that affected cuticle color upon knockdown, but no further validation was performed.

Here, we utilize the fruit fly cuticle to identify novel regulators of dopamine. We start by

systematically examining classical pigmentation genes for their effect on dopamine in the head

and brain. From this we found that genes that are involved in dopamine synthesis are the only

genes with expected effects on dopamine; genes involved in degradation of dopamine either have

no effect or unexpectedly show reduced dopamine. We go on to characterize 458 genes identified

from the RNAi screen by Mummery-Widmer et al. (2009). We tested 330 of them with independent

RNAi lines and found 153 genes (~46%) with consistent cuticle pigmentation defects.   We

examine 34 prioritized genes hits for changes in dopamine in the head and brain. From this

analysis, we found two genes, *mask* and *clu*, that reduced brain dopamine levels upon knockdown

in dopaminergic neurons. Lastly, we used molecular biology and sleep behavioral studies to show

that *mask* appears to alter brain dopamine through regulation of TH transcription.

Results

Drosophila pigmentation genes affect dopamine in unexpected ways

To systematically test if changes in pigmentation genes affect dopamine as theorized in the field, we characterized loss-of-function alleles of genes involved in dopamine metabolism (Fig 1A) for pigmentation phenotypes and dopamine levels in the head and brain. The enzymes required for dopamine synthesis, TH and Ddc, are essential for survival (23,24). Thus, we took advantage of established RNAi lines and UAS/GAL4 system (25–28). It is known that reduction of *TH* or *Ddc* level or activity causes pale pigmentation and reduced dopamine (9,10,14). We validated our method by knocking down *TH* using a verified UAS-RNAi (29), which causes a strong pale cuticle phenotype (Figs 1B and 1-C), reduction in head dopamine by ~60% (Fig 1E), and reduction in brain dopamine by ~32% (Fig 1E'). *Ddc* knockdown causes a weaker effect than *TH* RNAi in the cuticle (Figs 1B and 1,D) and in the head (~35% dopamine reduction, Fig 1E), with no significant effect on dopamine in the brain (Fig 1E'). This may be due to the strength of the RNAi and/or enzyme level requirements (since TH is the rate-limiting enzyme).

Figure 1

Fig 1. Manipulation of pigmentation enzymes have unexpected effects on dopamine. (A) *Drosophila* dopamine metabolism pathway. Gene names encoding pigmentation enzymes are in blue and metabolites are in black. (B-D) Thorax phenotypes observed upon knockdown of Tyrosine hydroxylase (TH, also known as Pale) (C) and Dopa decarboxylase (Ddc) (D) compared to control (B). (E-E') Dopamine levels from the head (E) and brain (E') upon knockdown of *TH* and *Ddc*. Pigmentation phenotypes observed mutant alleles in the thorax (F-L) and the body (F'-L') (Arrow in I' indicates the darker region characteristic of *speck* mutants). *T2A-GAL4* alleles for *black*, *speck*, and *tan* adult brain expression with dopaminergic neurons (DAN) staining in magenta (DANs) (M-P). (Q) Dopamine in the head of classical pigmentation mutants, *black[1]*, *ebony[1]*, *straw[1]*, and *yellow[1]*. (R) Dopamine levels in dissected brains of classical pigmentation mutants. Ordinary one-way ANOVA test with Dunlett multiple comparisons test. *=$p<0.05$, **=$p<0.01$, ***=$p<0.001$, ****=$p<0.0001$. Error bars show standard deviation (SD).

To test genes implicated in dopamine degradation and cuticle development (pigmentation

and structural rigidity), we examined known mutant alleles. These alleles cause darker or paler

cuticles (Figs 1F-1L and, 1F'-1L') depending on their role in melanin metabolism. Unexpectedly,

black, *speck*, *straw* and *yellow* mutants reduce head dopamine levels (Fig 1Q), while *ebony* and

tan mutants had no significant effect (Fig 1Q). Next, we checked the brain expression patterns of

pigmentation enzymes. Based on our own expression analysis (Figs 1M-1P) and previously

reported expression data (30,31), most are expressed in the fly brain, but their expression does

not appear to overlap with DANs. However, we found no significant effect on brain dopamine in

these pigmentation genes (Fig 1R).

Together this data shows that changes in dopamine synthesis cause a paler cuticle, which

reflects a reduction in dopamine, but changes in dopamine metabolism enzymes downstream of

dopamine synthesis have pigmentation phenotypes that do not correlate with dopamine level

differences.

132   RNAi-based cuticle pigmentation screen

Mining a previously published genome-wide scale RNAi screen (22), we identified 458

genes with pigmentation defects. To check if these phenotypes can be reproduced, we located

718 additional UAS-RNAi lines corresponding to 330 genes. 426 lines came from the National

Institute of Genetics in Mishima, Japan (32), and 292 came from the Transgenic RNAi Project

(TRiP) at Harvard Medical School (25). 153 genes caused pigmentation defects in at least one

additional RNAi line, showing a validation of 46.4% from the previous screen (Fig 2A).

Figure 2

Fig 2. RNAi pigmentation screen reveals a broad spectrum of phenotypes and classes of

genes (A) Flow chart depicting screening process. (B) Examples of the phenotypes observed

from the RNAi-based pigmentation screen. (C) Protein-protein interaction network for 153 high

confidence pigmentation gene hits from the RNAi-screen using STRING network analysis.

Strength of line indicates confidence or strength of data support. Genes with no known

interactions are not displayed. (D-F) Pie charts for the 153 pigmentation gene hits, including

strength of phenotype (D), phenotype color (E), and conservation (F).

Our RNAi-based secondary screen produced a spectrum of phenotypes (Fig 2B), but the

phenotypes strengths (mild, moderate, or strong) were equally represented (mild = 38%,

moderate = 33%, strong = 29%, Fig 2D). Pigmentation color changes were classified in three

primary categories, pale, dark, or mix (a combination of pale and dark in a single thorax), which

were not equally represented (Fig 2E). Dark cuticle phenotypes were the most common (56%),

with pale and mixed cuticle phenotypes seen in 28% and 16% of genes, respectively.

We performed a protein-network analysis (Fig 2C) and found a group of proteins

previously linked to cuticle pigmentation as well as clusters related to RNA processing, ubiquitin

proteasome system, mitochondria-associated proteins, and signaling pathway genes, Hippo and

EGF (epidermal growth factor) signaling in particular. GO enrichment analysis highlighted terms

associated with cuticle pigmentation, protein stability, and several signaling pathways (S1 Fig).

We also observed that 85% of our genes were conserved between *Drosophila* and human, an

enrichment from the 68% of conserved genes in the original screen (Fig 2F and S2 Fig).

In summary, the cuticle pigmentation screen identified a spectrum of pale, dark and mixed

phenotypes in 153 genes, showing a validation of ~50% from the original screen. Close

examination of these hits highlights conserved molecular processes.

Many pigmentation genes isolated have a neurological

consequence

To investigate if our gene candidates have a neurological consequence, we first

checked if their homologs are associated with a neurological disease. We found that 51.2% of the

fly gene homologs are associated with a disease, 78% of those have at least one disease with

neurological phenotypes (Table 1 and S2 Fig), such as intellectual disability, autism spectrum

disorder, ataxia, epilepsy, and cerebral atrophy.

Table 1. Fly pigmentation genes identified from this screen with neurological disease

associated homologs in human.

Fly Gene	Human Homologs with Neurological Phenotypes (DIOPT Score, Max=18)	Notable Neurological Phenotypes
AP-1sigma	AP1S2 (12), AP1S1 (11), AP2S1 (3)	ataxia, DD, seizures
ATP7	ATP7A (13), ATP7B (13)	dystonia, neurodegeneration, seizures, tremor
Atu	LEO1 (10)	ASD
byn	TBXT (11), TBX19 (10)	neural tube defects, seizures
CG16758	PNP (14)	ataxia, behavioral problems, spastic diplegia, tremor
CG4328	LMX1B (10)	spina bifida
CG43689	MYT1L (7)	DD, ID
CG4679	PTCD3 (14)	cerebral atrophy, myoclonus, psychomotor delay
CG5902	AAMERCR1 (13)	cerebral atrophy, early motor delay, myoclonus
CG8728	PMPCA (15)	ataxia
ClC-c	CLCN3 (12), CLCN4 (11), CLCNKA (3)	ID, language delay, seizures, structural abnormalities
ctrip	TRIP12 (10)	delayed speech, ID, psychomotor delay
Cyp301a1	CYP24A1 (5), CYP27B1 (4), CYP27A1 (4)	ataxia, motor delay, psychiatric symptoms, seizures
CysRS-m	CARS2 (15)	cerebral atrophy, epilepsy, psychomotor delay
Ddc	DDC (14), HDC (4)	dystonia, psychomotor delay, sleep disturbance
Dgp-1	GTPBP2 (3)	ataxia, dystonia, ID, psychomotor delay, seizures
Dscam2	DSCAM (10)	ASD
Egfr	ERBB4 (10), ERBB2 (10), ERBB3 (9)	ALS, axonal neuropathy, DD, seizures
Gbeta13F	GNB1 (15), GNB4 (13), GNB2 (12), GNB5 (3)	ADHD, CMT, DD, ID, language delay

gig	TSC2 (13)	brain structural abnormalities, seizures
Hrb98DE	HNRNPA2B1 (12), HNRNPA1 (9)	ALS, dementia, motor neuropathy
Itpr	ITPR1 (14), ITPR3 (11)	ataxia, cerebellar atrophy, motor delay
Khc	KIF5C (12), KIF5A (12)	ALS, psychomotor delay, seizures, spastic paraplegia
Larp7	LARP7 (10)	anxiety, DD, ID, unstable gait
LeuRS-m	LARS2 (15)	cognitive impairment, DD, dystonia, tremor, seizures
mask	ANKRD17 (10)	DD, ID, motor delay, speech delay
mbt	PAK1 (3), PAK3 (3), PAK2 (3)	ID, macrocephaly, seizures, speech delay
Membrin	GOSR2 (13)	cerebral atrophy, DD, seizures
Mp	COL18A1 (6)	ataxia, seizures
Mpcp2	SLC25A3 (14)	motor delay
ND-13B	NDUFA5 (13)	ASD
ND-42	NDUFA10 (14)	DD, strutural abnormalities
Npc2b	NPC2 (7)	behavioral problems, DD, dementia, dystonia, seizures
per	PER3 (8), PER2 (7)	sleep abnormalities
PIG-A	PIGA (15)	seizures, structural abnormalities
ple	TH (15)	ataxia, dystonia, Parkinsonism, motor and sleep delay
Prosap	SHANK2 (10), SHANK3 (8)	ASD, DD, schizophrenia, seizures
Raf	BRAF (14), RAF1 (11)	DD, seizures
Rbp9	ELAVL2 (12), ELAVL3 (10)	ASD
rdx	SPOP (10)	DD, ID, speech delay
robo1	ROBO3 (8)	structural abnormalities

SCOT	OXCT1 (15)	coma, psychomotor delay
sesB	SLC25A4 (12)	motor delay, hyperreflexia
slp1	FOXG1 (6), FOXI1 (3), FOXJ1 (3), FOXC1 (3)	dystonia, motor delay, sleep and structural abnormalities
Smurf	ITCH (4), NEDD4L (3)	ID, psychomotor delay, seizures
Sod3	SOD1 (4)	ALS, spastic tetraplegia
Sos	SOS1 (14), SOS2 (12)	cognitive impair, learning disabilities
Spn38F	SERPINC1 (4), SERPINI1 (4)	cerebral atrophy, dementia, seizures
stv	BAG3 (9)	abnormal gait, behavioral problems, neuropathy
tws	PPP2R2A (13)	ataxia
Usp2	USP8 (4)	behavioral problems
Usp7	USP7 (13)	ID, psychomotor delay, seizures, structural abnormalities

Table 1. Homologs were defined as a Drosophila Integrative Orthology Prediction Tool (DIOPT,

version 8.0) score greater than or equal to three. Neurological disease associations were based

on Online Mendelian Inheritance in Man (OMIM)_clincal disease phenotypes and high-confident

ASD genes listed in Simons Foundation Autism Research Initiative (SFARI) gene database (class

S, 1, 2 or 3). ASD=Autism Spectrum Disorder, DD=Developmental Delay, ID=Intellectual

Disability, ALS=Amytrophic Lateral Sclerosis, CMT=Charcot-Marie-Tooth disease.

Since dopamine is known to regulate locomotion and sleep (22,33), we screened for

locomotion and sleep defects using the Drosophila Activity Monitor (DAM) (34). We tested 274

RNAi lines from the NIG, TRiP, and VDRC RNAi collections (35–37) corresponding to the

validated 152 gene hits. We scored the lines for total locomotion, total sleep, sleep bout length,

and sleep latency during the night and day (Fig 3). There was a broad range in the RNAi lines for

each of these phenotypes (Figs 3C). We classified RNAi lines "outliers" if they were two standard

deviations from the mean for a given phenotype (locomotion: 23 lines, total sleep: 20 lines, sleep

bout length: 21 lines, sleep latency: 25 lines). While total sleep and locomotion strongly

overlapped with each other, sleep latency and sleep bout length did not overlap with the other

sleep phenotypes (Fig 3D). We assessed if certain gene categories (>10 genes) showed any

distinct behavioral patterns (Fig 2C). There was no change in total locomotion or total sleep in the

categories assessed (Fig 3F and 3G). However, each categories showed a distinct pattern in

sleep bout length and latency (Fig 3F and 3H). This could indicate a distinction in how different

gene functions could affect sleep. Since genes that affect sleep can often show distinct

phenotypes that do not necessarily correspond to multiple phenotypes or times of day (38), any

gene with one phenotype during the day or night was classified as an "outlier". This produced

sixty lines, corresponding to 50 genes.

Figure 3

202
203

Fig 3. *Drosophila* Activity Monitor (DAM) screen identifies behavior "outliers" and

highlights distinct phenotypes for gene categories. (A-B) Representative activity and sleep

data measured in a neutral RNAi line (*mCherry RNAi*, A) and a behavioral "outlier" (*Raf RNAi*,

B) knocked down in DANs (*TH-GAL4*). (C) Bar graph showing all RNAi lines tested for 24hr

locomotion and total sleep during the 12hr light period, where light pink/blue represents fly lines

±1SD away from the mean, and dark pink/blue are fly lines ±2SDs away from the mean (a.k.a.

"outliers"). (D) Venn diagram of "outliers" for the four phenotypes examined (sleep latency, sleep

bout length, locomotion, and total sleep). (E-H) Behavioral analysis of lines in three gene

categories captured in the pigmentation screen, including locomotion (E), sleep latency (F),

sleep bout duration (G), and total sleep (H). Individual t-tests were performed to assess if genes

in a category were significantly different to all other genes tested, and only those with a p<0.10

are reported, with *=p<0.05.

Taken together, 77 of our 132 conserved genes are likely to have a neurological function,

because they are implicated human neurological disease (52 conserved genes) or were classified

as a behavioral "outliers" (45 conserved genes).

221 **11 genes alter dopamine levels in the fly head**

To prioritize our gene list for dopamine measurement, we identified conserved genes (132

conserved/152 pigmentation genes). Then, we prioritized these based on whether knockdown of

these genes showed a strong pigmentation effect (38 genes), a behavioral effect in the DAM (45

genes), and/or a human neurological disease association (52 genes). 72% (95/132 conserved

genes) were represented in one of these categories, and 29% (39/132) were represented in at

least two (Fig 4A). We classified genes in two or more categories as priority hits and tested them

for changes in dopamine using High Performance Liquid Chromatography (HPLC). We repeated

previous results showing that knocking down *TH* or overexpressing a *TH* cDNA using *TH-GAL4*

significantly reduces or increases head dopamine without effecting serotonin (Figs 4B and 4C)

(29).

Figure 4

Fig 4. Eleven genes alter head dopamine upon knockdown. (A) Venn diagram of the

categories used to prioritize 39 genes (in yellow). (B) Representative chromatograms for

dopamine (DA) and serotonin (5-HT) upon knockdown or overexpression of TH in Dopaminergic

cells (*TH-GAL4*). (C) Quantification of DA and 5-HT from chromatograms. (D-E) HPLC analysis

on prioritized genes for UAS-RNAi lines from NIG (D) and TRiP (E). Pie charts for all genes for

cuticle color (F) and trend in dopamine (G). (H) XY plot comparing pigmentation phenotype to

dopamine levels. For (C) Brown-Forsythe ANOVA test performed with Dunnett's T3 multiple

comparisons test. For (D) and (E) ordinary One-way ANOVA test with Dunnett's multiple

comparisons test. **=p<0.01, ***=p<0.001, ****=p<0.0001. Error bars represent SD.

We found that 11 of 35 prioritized genes (note: some not tested due to lethality or stock

issues) significantly altered head dopamine levels (Figs 4D and 4E). Since control NIG lines

showed about ~80% dopamine levels compared to the TRiP collection (S3 Fig), we compared

the DA levels to control lines appropriate for each collection (e.g. UAS-lacZ RNAi). Unlike cuticle

pigmentation, where 65% of genes caused a dark cuticle (excludes mixed cuticle phenotype),

most genes trended toward a reduction in dopamine (71%, 25/35, Figs 4F and 4G). There was

no correlation between the cuticle color and the dopamine level in the fly head (Fig 4H).

In summary, our HPLC analysis on fly heads revealed 11 genes that significantly affect

dopamine levels. We saw a trend in reduction in dopamine across all lines tested, and we

observed no correlation between cuticle color and dopamine level.

Brain study reveals *mask* and *clueless* as potential regulators

of brain dopamine

To determine which of the 11 genes that affect dopamine in the head have consequences

on the brain, we examined the expression of four of them in the brain using available T2A-GAL4

lines (39–41). We found that several overlapped with somewhere between 25% - 100% of DANs

(S5 Fig, Figs 5B and 5G). In publicly available single-cell mRNA sequencing datasets, ten of our

gene hits (all except *Spn38F*) show expression in the brain with varying overlap with TH

expressing DANs (42,43).

Figure 5

Fig 5. Brain dopamine analysis identifies *mask* and *clu* as novel regulators of dopamine

levels in the fly brain (A) Total dopamine in the adult fly brain upon knockdown of *clu*. (B)

clu::GFP expression pattern in the adult brain (B-E, B'-E'). (F) Total dopamine in the adult fly

brain upon knockdown of *mask*. (G) *mask::GFP* expression pattern in the adult brain (H-J, H'-

J'). (K) qRT-PCR of TH upon *mask* knockdown with the recombined DAN (*TH, R58-GAL4*)

driver. (L) Quantification of total TH protein level through immunofluorescence in DAN clusters

upon knockdown of *mask*. (M) Quantification of TH positive (TH+) neurons in DAN clusters

upon knockdown of *mask*. Ordinary one-way ANOVA was done with Dunnett's multiple

comparisons test (F, I, N). Error bars represent SD. Scale bars are 50 μm.

The *TH-GAL4* line commonly used for dopamine studies misses a large portion of DANs,

particularly in the PAM cluster (~87 DANs missed/~142 total DANs per hemisphere) (44), making

it not ideal for quantifying total changes in brain dopamine. We recombined the *TH-GAL4* with

another driver that hits a large portion of the PAM cluster neurons (*GMR58E02-GAL4 or R58-*

GAL4) (18) generating a recombined line (labelled *TH,R58-GAL4*) that hits ~95% of DANs (S4

Fig).

We microdissected adult fly brains and measured dopamine for our 11 gene hits. Most of

them showed no change in brain dopamine (S5 Fig), but *mask* (*multiple ankyrin repeats single*

KH domain) and *clueless* (*clueless*) knockdown reduced total brain dopamine for two independent

RNAi lines (Figs 5A and 5F). Protein trap lines (45,46) for both Clueless and Mask show broad

expression patterns in the brain (Figs 5B-5EJ-5M and 5G-5J). When we focused on DAN clusters

(e.g. PPL1 and PPM2), Clueless protein appears to be enriched in DANs (Figs 5C-5E and 5C'-

5E'), whereas Mask shows a broad and equal expression throughout most cells in the brain (Figs

5H-5J and 5H'-5J').

In summary, our fly brain analysis revealed two genes that (18%, 2/11) significantly altered

brain dopamine.

mask regulates Tyrosine Hydroxylase and alters sleep in a

dopamine-dependent manner.

To test if *clu* and *mask* knockdown could be due to DAN loss, we quantified the number

of DANs in several clusters (44). There was no difference in neuron number for every cluster

examined (Fig 5M and S6 Fig). Since our pigmentation gene study revealed that only the

manipulation in TH reduced brain dopamine (Fig 1), we examined if knocking down *clu* or *mask*

affects *TH* RNA or protein level. Knockdown of *mask* led to a significant reduction in *TH* mRNA

level for one RNAi (50% reduction), with the other RNAi trending (20-30% reduction, Fig 5K).

Upon examining TH levels in different DAN clusters, two clusters showed significant reduction in

TH protein level, with other clusters showing a trend in TH protein reduction (PAL and PPM2/3,

Fig 5L and S6 Fig). To assess behavioral consequences, we performed additional sleep analysis

on *mask*. Dopamine is a wake-promoting agent, and complete loss of dopamine in the brain

significantly increases sleep (47,48). Knocking down *TH* using the *TH-GAL4* did not significantly

alter total levels of sleep during light or dark periods (S7 Fig), which may be due to the selective

expression of the GAL4 or the partial reduction of dopamine (Fig 4C). However, there was a

significant increase in sleep in the two hours before light onset, which is classified as light

anticipation. Similarly, knockdown of *mask* with the TH-GAL4 showed a highly consistent

reduction in light anticipation (Fig 6A, 6B, 6B', and S8 Fig). When we feed the flies L-DOPA, the

effect is no longer seen (Fig 6C, S8 Fig). Additionally, in *Drosophila* caffeine's effects on sleep

are mediated by dopamine through a point that is upstream of L-DOPA, potentially via TH activity

(49). Based on this, we suspected that *mask* knockdown would ameliorate the effects of caffeine

on sleep, since *mask* appears to reduce the synthesis of *TH*. We saw that caffeine's effect on

total sleep and sleep in the dark is ameliorated upon *mask* knockdown (Fig 6D, 6E, and S8 Fig).

 Figure 6

Fig 6. *mask* knockdown reduces TH mRNA and protein and causes sleep phenotypes

that align with reduction in TH levels (A) 24-hr sleep graph (sleep binned in 30-minute

intervals) upon mask knockdown using the TH-GAL4. (B) Quantification of total sleep

throughout the 12-hr dark period upon mask knockdown (B') Quantification of light anticipation

upon mask knockdown. (C) Light anticipation in mask knockdown flies with L-DOPA feeding. (D-

E) Caffeine effects on sleep upon mask knockdown – (D) Total Sleep and (E) Sleep during the

12-hr dark period. For samples with more than one experimental, ordinary one-way ANOVAs

were performed with Dunnett's multiple comparisons. (B, B', C). For samples with one

experimental, individual t-tests were performed (D,E). *=p<0.05, **p<0.01, ****p<0.0001. Error

bars represent SD.

Follow up studies on *clueless* showed that *clueless* knockdown increases TH RNA level

by about 1.5-2X, with no effect on TH protein level or TH+ neurons (S9 Fig), and while there

was an increase in sleep across several RNAi lines, there is no light anticipation phenotype.

Feeding TH>clu RNAi flies L-DOPA was unable to rescue the *clueless* effects on sleep (S9 Fig),

suggesting that the effects on dopamine and TH may be a secondary effect. Additionally, the

antioxidant NACA was also unable to rescue the sleep defects seen upon *clueless* knockdown.

In conclusion, *mask* appears to reduce total dopamine levels and sleep by affecting

transcription of TH, while *clueless* is acting through an alternative mechanism.

Discussion

Validation of 148 novel cuticle pigmentation genes

Our RNAi screened confirmed 153 genes that were previously implicated in *Drosophila*

pigmentation – 148 of these genes have not been previously associated with dopamine biology.

Because we observed a strong enrichment for genes involved in pigmentation and tyrosine

metabolism (S1 Fig), our method for screening and analysis effectively identified genes that play

critical roles in the biological process of interest.

While prior research has compared genes that overlap with different types of screening

strategies (e.g. chemical mutagenesis vs. RNAi), most RNAi phenotypes from large-scale screens

have not been validated with alternative RNAi lines or collections. The original source RNAi

screen used the lines available at the VDRC stock center (35,37). Alternative sources for RNAi

lines (i.e. TRiP and NIG) start with different backgrounds, vectors, hairpin lengths, insertion

methods, and/or target different areas of the gene (36,50). This allow researchers to overcome

the off-targeting and weak silencing seen with RNAi analysis, but to our knowledge, there have

been no screens or meta-analyses to systematically test consistency across lines from different

sources. In this study, we found that ~46% (153/330) of genes with a pigmentation phenotype

showed a phenotype in an alternative RNAi, and 87% of those phenotypes showed agreement in

pigmentation color, highlighting that pigmentation as a phenotype is relatively consistent (S10

Fig).

Interestingly, 85% of our pigmentation genes had at least one homolog, whereas genes

tested in the original RNAi screen list only had ~68% conservation (1.25-fold enrichment, S2 Fig).

Even though pigmentation of the cuticle through dopamine and its synthesis enzymes, produced

through TH and Ddc, is an invertebrate-specific phenomenon (12), the enrichment for conserved

genes suggests that this phenotype is relevant for some conserved biological pathway and

mammalian biology. We propose that cuticle pigmentation in insects may be a "phenolog" (51) of

certain neurological traits in mammals

Protein-protein interaction network highlights unexpected gene associations

When we investigated what kind of gene categories were found in the pigmentation

screen, we saw clusters of genes that are associated with ubiquitin-proteasome system (UPS),

RNA processing, mitochondria, and developmental signaling pathways. Each of these processes

could be relevant to dopamine biology. For example, there is some evidence that the UPS system

regulates TH levels in the mammals since inhibition of the UPS system has been shown to

increase the TH protein in a PC12 rat cell line (52,53). In addition, changes in RNA processing

could indicate changes in the dopamine synthesis enzymes TH and Ddc since these enzymes

undergo RNA processing to create two different isoforms (54). The two TH isoforms differ in tissue

expression and kinetics(55). In mammals, post-transcriptional trafficking of TH mRNA and

translational regulation has been suggested to be important for its function (56).

We identified 12 genes that are known to function at the mitochondria, mostly in the

electron transport chain (*ND-13B*, *ND-42*, *COX5A*), mitochondrial mRNA translation (*mRpL47*,

mRpL48, *CG4679*), and mitochondrial protein localization (*CG8728*, *clu*, *mask*, *Mpcp2*). It is well-

known that the mitochondria have a key role in dopamine metabolism in mammals since MAO

acts in the mitochondria to degrade dopamine. However, there is no known MOA homolog in

flies(12). Interestingly, the metabolites for oxidation and methylation of dopamine (DOPAC and

HVA) have been observed in flies (57–60). In addition, mutants for enzymes thought to be the

primary regulators of dopamine degradation (*ebony*, *tan*, and *speck*) (61,62) did not have any

effect on dopamine in the brain (Fig 1), suggesting an alternative process regulates dopamine

degradation in the brain. It is plausible that mitochondria could be involved in this process.

The last major group of genes identified from our screen was developmental signaling,

specifically EGF signaling and Hippo signaling. To date, neither EGF nor Hippo signaling have an

established connection with dopamine biology in *Drosophila*. We may have pulled these genes

out is because the process of forming the dorso-central thorax is developmentally regulated (63–

65). However, we did not identify core components of the other key pathways, such as Notch or

Hedgehog, suggesting that EGF and Hippo are more specifically involved in this pigmentation

process.

Cuticle pigmentation phenotypes may not reflect changes in

dopamine

The field has generally assumed that a darker cuticle reflects an elevated level of

dopamine (15,66). However, when we measured dopamine from degradation enzyme mutants

using HPLC, we saw either no effect on dopamine level or reduced level of dopamine in the head.

In addition, when we measured dopamine from the pigmentation genes from the screen, a darker

cuticle did not reflect elevated levels of dopamine in the head. While we were able to increase

dopamine by 3-5 fold upon TH cDNA overexpression in the head (Fig 4C), this phenomenon was

not observable in the brain. One reason could be that there is a selective pressure or a strong

feedback mechanism that prevents elevated levels of dopamine in the nervous system. Since

oxidized dopamine could be toxic (67), this could be a mechanism to prevent cell toxicity.

Regardless, changes in cuticle pigmentation do not necessarily reflect changes in dopamine level,

so assumptions should not be made that increase in melanin levels in the fly corresponds to

increased levels of dopamine.

Upon seeing alterations of dopamine in the head for some of the classic pigmentation

genes, we next wanted to test if this corresponded to changes in the brain. In the case of

established dopamine metabolism genes, we only saw a change in dopamine levels upon

reduction of TH. When we looked at expression for some of these genes, we saw that all the

genes we were able to assess (*black*, *speck*, *tan*) were expressed in the brain, but their expression

did not overlap well with DANs. It may be that dopamine degradation does not happen in the

neurons since it was hypothesized to occur in the glia (31,68). However, the expression does not

seem to show a broad glial expression using the reagents we had access to. For the 11 genes

from the pigmentation RNAi screen that showed altered levels of dopamine, only 2 (18%) of them

showed an effect on dopamine in the brain. Hence, head dopamine levels should not be used as

a proxy for brain dopamine levels.

clu and _mask_ reduces brain dopamine levels through different

mechanisms

417   Once we identified that knockdown of *clueless* and *mask* both reduced dopamine in the

418 brain without affecting the number of DANs, we hypothesized that they may be acting through TH

419 since our initial dopamine metabolism studies showed that only changes in TH significantly

420 affected dopamine levels in the brain.

421   *mask* encodes a scaffolding molecule that functions in multiple contexts (69–71). In this

422 study, *mask* knockdown reduced both *TH* mRNA and protein levels, suggesting it acts on TH

423 transcription. Follow-up behavior analysis, showed that *mask* knockdown in DANs led to a highly

424 consistent change in sleep. Specifically, *mask* knockdown reduced the number of flies that

425 anticipated light. While fruit flies typically start to wake before light onset (72), loss of mask in

426 DANs suppresses this anticipation behavior. Supporting this mechanism, we saw a similar

427 phenotype with the TH RNAi and the DAN driver (TH>TH RNAi, S7 Fig). Dopamine is a wake-

428 promoting molecule, and if dopamine synthesis normally increases before light onset as a means

429 of waking the fly, reduced levels of TH could put strain on this mechanism.

430   When we fed *mask* knockdown flies the precursor for dopamine, L-DOPA, this abolished

431 the effect on light anticipation. We suspect that feeding L-DOPA does not override all sleep-

432 dependent effects because clueless sleep phenotypes remained when we fed them L-DOPA (S9

433 Fig). There is evidence that in the fruit fly caffeine affects sleep through TH. Specifically, Nall et

434 al. showed that caffeine's effect on sleep is absent in TH mutant flies, even if L-DOPA is given

435 (49). Since *mask* knockdown effects TH mRNA levels, we tested if mask knockdown would

436 suppress the effects of caffeine. We found that the reduction in total sleep and sleep during the

437 dark period normally seen upon caffeine administration were absent upon mask knockdown. This

438 further supports that mask alters TH levels.

439   While we do not have any direct evidence for how *mask* may be altering TH levels, *mask*

440 has been implicated in multiple biological pathways that were highlighted from our screen,

including EGF signaling (e.g. *Egfr*, *Sev*, Raf), Hippo signaling (e.g. *hpo*, *wts*, *mats*), and

mitochondrial biology (73–76). Egf and Hippo signaling have well-known transcriptional effects

(77,78), but whether they may regulate TH transcription directly or indirectly in this context need

further investigation. It is also possible that *mask* is acting in several of these pathways at once,

which causes it to have a strong additive effect that produces a significant change in dopamine

in the whole brain, while knockdown of individual pathway genes may have a more restricted role

(e.g. *Egfr*, *Sos*, *hpo*).

In addition to *mask*, we also saw that knockdown of *clueless* leads to significant reduction

in dopamine. However, when we explored the behavioral consequence of *clueless* knockdown, it

did not show the same effect on sleep as *mask* or *TH* knockdown. In addition, the sleep phenotype

was not rescued by L-DOPA, and molecular biology showed that *clueless* knockdown leads to an

increase in TH mRNA with no change in TH protein. Taken together, this indicates that the effect

on dopamine from *clueless* knockdown may be a secondary consequence. *clueless* encodes a

ribonucleoprotein thought to act at the outer mitochondrial membrane to promote proper formation

of protein complexes including mitophagy proteins like Pink1 and Parkin (79,80). It also interacts

with the ribosome and translocase complex at the mitochondrial membrane (79,81) suggesting it

acts as a regulator for mitochondrial protein translation and import. Knockdown of *clueless*

showed no effect on TH protein level, but since this is a mitochondrial protein, it may be more

likely that the effect on dopamine is through the previously proposed roles of mitochondria in

regulating dopamine levels.

In conclusion, while both genes cause a reduction in dopamine, *mask* appears to regulate

dopamine by altering TH levels, possibly through regulation of TH mRNA. On the other hand,

clueless effects on dopamine appear to be a secondary consequence, with a behavioral effect

that is independent of dopamine. Additional analysis on changes in enzymatic activity as well as

potential downstream targets would help identify how these genes are acting.

466 Some regulators of dopamine levels may act locally

When we examined the expression of some of our pigmentation genes in the brain (S11

Fig), we found that many of them do not overlap with all DANs. Indication that measuring global

changes in dopamine may not be best for determining effects on DA for many of these genes.

They may require targeted dopamine measurements, perhaps through a fixed or live reporter

(82,83). It is also possible to assess changes in dopamine using behavior, but this would require

detailed assessment for each gene since behaviors are often cluster specific (84).

In conclusion, our screen of identified novel pigmentation genes, a subset of which were

identified as novel regulators of dopamine *in vivo*. Unbiased forward genetic screens in model

organisms are powerful ways to identify unanticipated links between distinct biological pathways

and provide new molecular handles to study the functional connections between them. Application

of such strategies to the regulation of dopamine levels will likely continue to identify novel factors,

some of which will impact our understanding of human neurological and neurodevelopmental

diseases.

480 Resource Availability

481   All data generated or analyzed during this study are included in this published article and its

482   supplementary information files. This study did not generate any new unique reagents.

483 Acknowledgements

We thank the Bloomington *Drosophila* Stock Center (USA), the National Institutes Genetics Fly

Stock Center (Japan), *Vienna* Drosophila RNAi Center (Austria) and Dr. Hugo Bellen for providing

useful fly stocks and reagents for this project. We would like to thank Drs. Jonathan Andrews,

Hugo Bellen, Brigitte Dauwalder, Herman Dierick, Linsday Goodman, Oguz Kanca, Kartik

Venkatachalam, Michael Wangler, Sheng Zhang and Huda Zoghbi for useful suggestions,

discussions and advice on the overall project. This work was supported by startup funds to SY

from the Jan and Dan Duncan Neurological Research Institute at Texas Children's Hospital and

the Department of Molecular and Human Genetics at Baylor College of Medicine. Confocal

microscopy at BCM was supported in part by the Intellectual and Developmental Disabilities

Research Center (IDDRC, grant number U54HD083092) from the Eunice Kennedy Shriver

National Institute of Child Health & Human Development.

Authors Contributions

496 S.Y. and S.L.D. conceived the experiments and wrote the manuscript. S.Y., E.S.S., D.B., and

497 S.L.D. conducted the pigmentation screen and DAM behavior analysis screen. S.L.D. and K.W.

498 conducted the pigmentation mutant analysis experiments. S.L.D., S.B.G, H.D-S, Y.F., and K.W.

499 conduced the HPLC analysis. S.L.D. and Y.F. performed confocal imaging. S.L.D. conducted all

500 data analysis. All authors participated in the critical analysis of the manuscript.

Declarations of Interests

502 The authors declare no competing interests.

Materials and Methods

Fly Maintenance

505 Flies (*Drosophila melanogaster*) strains and crosses were reared at 25°C on a 12:12 Light/Dark

506 (LD) program unless stated otherwise. Flies were fed a molasses-based food source. Different

507 temperatures were used to achieve different levels of gene knockdown or overexpression, which

508   is documented for each experiment. Flies were transferred to a new vial 0-2 days post-eclosion

509   (dpe) for age-appropriate analysis. Experiments were run on 3-7 dpe flies unless otherwise stated.

Fly Stocks and Crosses

Mutant and transgenic trains were obtained from Bloomington *Drosophila* Stock Center (BDSC,

https://bdsc.indiana.edu/),   Vienna   Drosophila   Research   Center   (VDRC,

https://www.viennabiocenter.org/vbcf/vienna-drosophila-resource-center/),   the   Japanese

National Institute of Genetics (https://shigen.nig.ac.jp/fly/nigfly/), or were gifts from scientists in

the field or generated in house (S1 Table – S5 Table).

For the cuticle screen *pnr-GAL4* (85) was crossed to the specified UAS-RNAi line (see S1 Table)

at 29°C, 25°C, and/or 18°C and examined after pigmentation is completed (>1 dpe). For the

behavior and HPLC analysis *TH-GAL4* females (also called *ple-GAL4*, BDSC stock #8848) were

crossed to the specified RNAi line (see S2 Table). For all HPLC analysis, the controls were

specific to the tested UAS-RNAi line (86). Thus, the TRiP collection of RNAi lines are compared

to the *UAS-mCherry(mCh) RNAi* line (BDSC #35785) and the NIG collection of RNAi lines as well

as the VDRC RNAi lines are compared to the *UAS-lacZ RNAi* line. For the HPLC analysis

performed on brains, we recombined the *TH-GAL4* with another DAN driver *GMR58E02-GAL4*

(BDSC #41347), which expresses in a large group DANs in the PAM cluster (27). These flies are

referred to as *TH,R58-GAL4* throughout the article. The *TH,R58-GAL4* line was also used for

qPCR and TH protein analysis via immunohistochemistry. For gene expression analysis *UAS-*

mCh::nls (BDSC #38424) females were collected and crossed with respective T2A-GAL4 lines

(see S5 Table). Female and males were both imaged at ~5-7dpe. For the follow up sleep studies,

the *UAS-RNAi* males were crossed with *TH-GAL4* virgins for the experimental. For the controls

w[1118] females were crossed to males from either the *TH-GAL4* and *UAS-RNAi* lines

independently.

Notum Dissection and Imaging

For thorax analysis and dissection, males and females were both selected, though differences were not generally observed, and females are shown here. Flies that were 3-5 dpe were collected and placed in 70% EtOH. Notum dissection and imaging was previously published (87). In summary, the thorax was dissected by removing the legs, abdomen, and head. Then the thorax was cut such that a hole was on the ventral side to allow solution to pass into the thorax. These dissected thoraces were placed in 10% KOH for 10 minutes at 90°C on a heat block. Then, the KOH was removed and replaced with 70% EtOH solution. They were mounted on a slide prepared with tape on two sides in mounting media (50% glycerol, 50% EtOH). The thoraces were imaged on a stereo microscope (Leica MZ16) using OPTRONICS® MicroFIRE camera. The images are z-stack brightfield images taken and collapsed using extended depth of field in Image-Pro Plus 7.0 and In-Focus (Version 1.6).

Brain Expression Analysis

Brain dissections and imaging were performed as previously reported (88). In summary, the brains were dissected at 3-7 dpe in ice cold PBS and then fixed in 4% PFA in 0.5% PBST for 20 minutes. Then, they were washed with a quick wash in 0.5% PBST followed by three 10-20-minute washes in 0.5% PBST while on a rotator. The samples were placed in primary antibody solution [anti-Tyrosine Hydroxylase 1:500, PelFreez Biologicals, rabbit, P40101; anti-elav 1:100, Developmental Studies Hybridoma Bank, rat, 7E8A10 in solution (5% Normal Donkey Serum, 0.1% NaN_3 in 0.5% PBST)] at 4°C for 3 days. Then, the samples were washed with a quick wash in 0.5% PBST followed by three 10-20-minute washes in 0.5% PBST while on a rotator. After the last wash, the samples were placed in a secondary antibody solution [anti-Rabbit 1:200 (Thermo Fisher Sci., Alexa-647, A-21208); anti-rat 1:200 (Thermo Fisher Sci., Alexa-488, A-27040) in

solution (5% Normal Donkey Serum, 0.1% NaN$_3$ in 0.5% PBST)] for two hours at room

temperature. The samples were washed again with one quick wash followed by three 10-20

minutes washes in 0.5% PBST on a rotator. Then, the brains were mounted in Vectashield

mounting media and imaged on a confocal microscope (Zeiss LSM 710 or Zeiss LSM 880). All

images shown in this manuscript are Z-projection images that were generated using the ZEN

software (Zeiss).

RNAi-based Pigmentation Screen

We identified pigmentation gene hits from a primary screen performed in Jurgen Knoblich's lab

using the Vienna Drosophila Resource Center collection of RNAi lines by accessing their public

RNAi screen database (https://bristlescreen.imba.oeaw.ac.at/start.php). The original screen was

performed on 20,262 RNAi lines, encompassing 11,619 genes (~82% of protein-coding genes)

(22). We selected genes that showed a cuticle pigmentation score for any RNAi line tested (gene

scores ranged from two to ten). To validate pigmentation defects, we selected genes that had

RNAi lines within the Fly National Institute of Genetics (NIG, 220 genes, 426 RNAi lines) or the

Harvard Transgenic RNAi Project (TRiP, 221 genes, 292 RNAi lines) collections. Each RNAi line

was tested at 29°C and 25°C. If there was no phenotype no further testing was performed. If it

was lethal, we tested it at 18°C.

Classification of Cuticle Phenotypes

Phenotypes were scored using a qualitative scoring system. This system used *UAS-TH RNAi* and

UAS-ebony RNAi, some of the strongest regulators of cuticle color, as a baseline for the "strong".

Then, the other lines were placed along the spectrum as mild, moderate, or strong. Phenotypes

were scored by two independent observers and differences were settled by a third independent

observer. If there was variability with one line or if there was variability amongst lines, this might

appear as mild-moderate, moderate-strong, or mild-strong. In these cases, the strongest

phenotype observed was documented as the recorded phenotype for a given gene.

580 Protein-protein interaction network and gene ontology

581 analysis

The protein-protein interaction network was generated using STRING (search tool for recurring

instances of neighboring genes; https://string-db.org/). This database includes protein-protein

interactions that are physical interactions and functional associations. For our data set we

included these sources of interactions: gene neighborhoods (genes that are found close together

across species) (89), curated databases (publicly available databases of protein interactions),

and experimental evidence (known complexes and pathways from curated sources) (90,91). Any

genes that did not interact with other genes from our screen were not included in the interaction

network. This database was last accessed on 05/05/2023. The lines between genes represents

the confidence in the interactions, which is based on complex algorithms that are dependent on

the source.

The gene ontology analysis was generated using GOrilla (Gene Ontology enRIchment anaLysis

and visuaLizAtion tool, https://cbl-gorilla.cs.technion.ac.il/). The 153 pigmentation genes hits were

included as target genes, and the background genes were the entire VDRC RNAi collection

included in the Mummery-Widmer et al. 2009 screen. Only the GO terms with a p-value greater

than 0.001 and a fold enrichment of greater than three were included in the analysis. Redundant

terms were removed using Revigo (http://revigo.irb.hr/) with a stringency of 0.7. Those that have

a fold enrichment greater than five were included in S1 Figure.

599 Human Neurological Disease Gene Classification

Homologs for each fly gene were identified using DIOPT (DRSC Integrative Ortholog Prediction

Tool, v8.0, https://www.flyrnai.org/diopt) (92). Genes were classified as orthologs if they had a

DIOPT score≥3. The human genes were scored as neurological disease-causing genes if they

had an OMIM (Online Mendelian Inheritance of Man, last accessed 01/01/2023,

https://www.omim.org/) disease association with any documented neurological phenotype (93).

Genes were also classified as human neurological disease genes if they were in the Simons

Simplex Collection of Autism Spectrum Disorder gene list (last accessed 06/01/2022,

https://gene.sfari.org/) and had a score equal to or less than 3 (1, 2, or 3) and/or syndromic (S)

(94).

609 Behavior Analysis using the *Drosophila* Activity Monitor

610 Behavior screen

For the behavior screen, the crosses were set in a 12:12 LD chamber at 25°C and transferred

every 2-3 days to increase the number of progenies. Upon eclosion, flies were transferred to 29°C

and kept for 2-3 more days before testing. Individual tubes (PPT5x65 Polycarbonate, Trikinetics

Inc, USA) appropriate for use with the *Drosophila* Activity Monitor (DAM, Model DAM2 for 5mm

tubes, TriKinetics Inc, USA) were loaded with approximately ½" worth of molasses-based food.

The end of the tube with food was sealed by placing the vial into Paraplast® (Sigma-Aldrich) wax

three times and allowing it to dry. Once they were dry, individual male flies were placed into each

vial, the vial was placed into the DAM recording chamber, and it was sealed using a 5mm tube

cap (CAP5, TriKinetics Inc, USA) with a hole stuffed with cotton. For each genotype, 16 individual

males were run, except in a few cases where we were unable to get that many flies. In which

case, as many living males as possible were run. The flies were loaded into a 12:12 LD chamber

at 29°C and monitored for 3 days. Then, the flies were removed and dead flies were eliminated

from analysis.

Locomotion and sleep analysis was run on a 24-hour period, which started at the first onset of

light after the flies were placed in the chamber (i.e. 18-24 hours after being placed in the chamber).

Locomotion was calculated based on the number of beam crossings over the full 24 hours, the

12 hours of light, and the 12 hours of dark. Sleep was classified as 5 minutes without any beam

crossings. Total Sleep was quantified as total time spent sleeping over the 12 hours of light and

the 12 hours of dark. Sleep latency was quantified as the amount of time after light onset or dark

onset before a 5-minute period of sleep. Sleep Bout Length was quantified as the average length

of sleep bouts (>5-minute period of sleep) during the 12 hours of light and 12 hours of dark.

"Outliers" were classified as lines that were more than two standard deviations from the mean of

all lines for any of the phenotypes scored (24-hour locomotion, 12-hour light locomotion, 12-hour

dark locomotion, 12-hour light total sleep, 12-hour dark total sleep, 12-hour light sleep bout length,

12-hour dark sleep bout length, sleep latency during the light, or sleep latency during the dark).

Mask follow up behavior studies

For these studies, similar tools and settings were used with these core exceptions. Upon eclosion

flies were transferred and kept at 25°C in group housing. For the behavior without drugs, individual

males were placed in tubes with sucrose food (2% agar + 5% sucrose) at 3-6 dpe and behavioral

analysis was collected and analyzed for the following five days. Sleep was then averaged across

the five days for each individual fly before statistical analysis was performed. For the behavior

with drugs, similar conditions were followed, except data was collected on the 2-6 days after they

were loaded into behavior tubes to allow the drug to take effect. The following drug doses were

used for their respective experiments (L-DOPA: 3mg/mL L-DOPA + 12.5 µg/mL carbidopa;

caffeine: 0.5 mg/mL caffeine; NACA: 40 µg/mL N-acetylcysteine amide antioxidant). Drugs were

dissolved directly into the sucrose food except for carbidopa which was dissolve in H_2O at 1:50x

concentration and then added into the sucrose food.

High Performance Liquid Chromatography (HPLC) analysis

Sample Preparation

For the HPLC on fly heads and brain, we adapted and modified the protocol previously published (95,96). Crosses were reared at 29°C and flies were transferred into new tubes 0-2 days post eclosion (dpe). They remained at 29°C until they were 3-5 dpe. Female heads were collected by anesthetizing flies with CO_2 and cutting their heads off with a razor blade. Samples were collected between ZT04-ZT08. The heads were placed into 60 µL of 50 mM citrate acetate (pH=4.5) and either used for analysis that day or frozen at -20°C. Five heads were used per sample, and ~10 samples were run per genotype. The day of HPLC analysis, the samples were thawed and grounded using a pestle for 30 seconds (Cordless Pestle Motor and Fisherbrand™ Disposable Pellet Pestle for 1.5mL tube). Then the samples were spun down at 13,000 rpm for 10 minutes. The supernatant was removed and placed into a new vial. 10 µL from each sample was used for the Bradford Protein Analysis Assay. The rest of the sample solution (~40-50 µL) was loaded into a vial (300 µL Polypropylene Sample Vials with 8mm Snap Caps) for HPLC analysis.

For HPLC analysis on fly brains, flies were collected 3-7 dpe. Then, 10 fly brains (5 male, 5 female) were dissected in ice cold PBS. Right after dissection, the brains were transferred to 60uL of ice cold 50mM citrate acetate (pH=4.5). Samples were frozen at -20°C and ran within 4 weeks of collection. The day of HPLC analysis, the samples were homogenized similarly to the fly heads, though for the Protein Assay, 20 µL of sample was used. Then, the remaining solution was loaded for HPLC analysis.

Bradford Protein Analysis Assay

The Bio-Rad Bradford Assay was used for colorimetric scoring of total protein. See product details

for full description (Bio-Rad Protein Assay Kit I #5000001). Briefly, the dye reagent was diluted

1:4 in MilliQ H_2O. Then, the solution was filtered through a 0.22 µm SFCA Nalgene filter. 10 µL

of each protein sample (standard or fly sample) was placed into single wells of a 96-well plate.

Then, 200 µL of diluted 1:4 dye reagent was added to each well. The samples rested for 30-45

minutes and then absorbance was measured using BMG Labtech FLUOstar OPTIMA microplate

reader. Protein measurements for each sample were calculated based on standards ran on the

same plate.

HPLC Information

The HPLC used is an Antec® Scientific product with a LC110S pump, SYSTEC OEM MINI

Vacuum Degasser, AS110 autosampler, a SenCell flow cell with salt bridge reference electrode

in the Decade Lite. Data was collected and processed using DataApex Clarity™ chromatography

software. The column used is chosen to work well for neurotransmitters (Acquity UPLC BEH C18

Column, 130Å, 1.7 µm, 1 mm X 100 mm with Acquity In-Line 0.2 µm Filter). The mobile phase

was a 6% Acetonitrile mobile phase optimized for our samples (74.4 mg $NA_2EDTA \cdot 2H_20$, 13.72

mL 85% w/v phosphoric acid, 42.04 g citric acid, 1.2 g OSA, 120 mL acetonitrile, H_20 up to 2L,

pH=6.0 using 50% NaOH solution). The mobile phase was degassed for 10 minutes using the

Bransonic® Ultrasonic Bath before being loaded into the HPLC machine.

The standards for HPLC were made by generating master stocks of 100 mM dopamine and 10

mM serotonin diluted in MilliQ H_2O, which were kept at 4°C. The day of HPLC analysis, the master

stocks were diluted to produce standards ranging from 5-100 nM for 5-HT and 5-1000 nM for

dopamine. Sample concentrations of dopamine (Sigma-Aldrich, Cat#H8502) and serotonin

(Sigma-Aldrich, Cat#H7752) were calculated based on standards run in the same batch.

Standards were compared to standards run on other days to assess for overall performance.

RNA expression analysis

Flies were raised at 29°C until 5-7 dpe, then heads were collected by cutting them off with a razor blade. Twenty heads were collected per sample with a mixture of males and females and placed directly on dry ice. Samples were kept on dry ice or at -80°C until RNA isolation. For RNA isolation samples were homogenized in 200µL of TRIzol®. Then 800µL of TRIzol® was added and the samples were mixed by pipetting. The samples incubated for 5 minutes at room temperature and were then spun down at 12,000 rpm for 10 minutes at 4°C. The supernatant was removed and 200µL of chloroform was added. The samples were mixed by shaking and then incubated at room temperature for 3 minutes. Then, they were spun at 10,000 rpm for 15 minutes at 4°C, and the aqueous layer was collected and placed in a new tube. 500µL of isopropanol was added and the samples were left to incubate for 10 minutes at room temperature. The samples were spun at 10,000 rpm for 10 minutes at 4°C and the supernatant was removed. The pellet was washed with 1mL of 75% EtOH and centrifuged at 5,000 rpm for 5 minutes. All EtOH was removed and the pellet dried at room temperature for 10 minutes. Then, the pellet was dissolved in 100µL of H_2O. RNA content and quality was measured using DeNovix DS-11 Fx spectrophotometer/fluorometer. cDNA reverse transcription was performed according to the iScript™ reverse transcriptase Bio-Rad kit and 2 µL of RNA sample was added in a 10 µL reaction for both the reverse and no reverse transcriptase reaction. Samples were measured for content and quality and then they were diluted to 100ng/µL. qPCR was performed using Bio-Rad iQ SYBR Green Supermix and measured on Bio-Rad CFX96™ Real-Time System. The TH primers used Forward: 5'-ATGTTCGCCATCAAGAAATCCT-3' and Reverse: 5'-GGGTCTCGAAACGGGCATC-3', and the control primers were for Rpl32 were Forward: 5'-ATGCTAAGCTGTCGCACAAATG-3' and Reverse: 5'-GTTCGATCCGTAACCGATGT-3'. For each sample, two technical replicates were run for the reverse transcriptase reaction, and the final quantity was based on the average of the

717   two. If the no reverse transcriptase reaction produced a product that was a Cq within ten, the

718   sample was discarded.

DAN and TH Quantification

Samples were stained and imaged like the brain expression analysis, except the primarily

antibody solution included an Elav antibody [anti-Tyrosine Hydroxylase 1:500, PelFreez

Biologicals, rabbit, P40101; anti-Elav 1:100, Developmental Studies Hybridoma Bank, rat,

7E8A10 in solution (5% Normal Donkey Serum, 0.1% NaN_3 in 0.5% PBST)]. Confocal imaging

was performed as described above. Once the samples were imaged, anterior, posterior, and

whole brain projections were generated that included all fluorescent signals from the anterior (PAL

and PAM clusters) and posterior (PPL1, PPM2, PPM3, PPL2ab) neuron clusters of interest. The

number of neurons per cluster (excluding the PAM cluster) were counted for each sample. PPM2

and PPM3 clusters were counted together. For TH protein quantification samples were assessed

using ImageJ. A boundary was drawn around each individual cluster and quantified. Then, the

local background for that cluster was subtracted from the cluster to give the fluorescence for the

given sample. Each hemisphere was treated as a separate sample.

Statistical Analysis

Statistical analysis was performed using Graphpad Prism 10. Unless otherwise stated, all data

was subjected to a ROUT outlier test, where all outliers were removed, and then a one-way

ANOVA was performed. Each of the experimental samples were compared to the control for the

given samples. In the cases where there were only two samples (i.e. control and experimental),

a t-test was performed.

For the HPLC screen on heads, the samples were normalized to the controls run on that day. The

TRiP collection was normalized to *UAS-mCh RNAi* control and the NIG collection was normalized

to the *UAS-lacZ RNAi* control.

References

1. Radwan B, Liu H, Chaudhury D. The role of dopamine in mood disorders and the

associated changes in circadian rhythms and sleep-wake cycle. Brain Res [Internet].

2019 Jan;1713:42–51. Available from: https://pubmed.ncbi.nlm.nih.gov/30481503/

2. Grace AA. Dysregulation of the dopamine system in the pathophysiology of

schizophrenia and depression. Nat Rev Neurosci [Internet]. 2016 Jan;17(8):524–32.

Available from: https://pubmed.ncbi.nlm.nih.gov/27256556/

3. Nutt DJ, Lingford-Hughes A, Erritzoe D, Stokes PRA. The dopamine theory of addiction:

40 years of highs and lows. Nat Rev Neurosci [Internet]. 2015 Jan;16(5):305–12.

Available from: https://pubmed.ncbi.nlm.nih.gov/25873042/

4. Church FC. Treatment Options for Motor and Non-Motor Symptoms of Parkinson's

Disease. Biomolecules [Internet]. 2021 Jan;11(4). Available from:

https://pubmed.ncbi.nlm.nih.gov/33924103/

5. Armstrong MJ, Okun MS. Diagnosis and Treatment of Parkinson Disease: A Review.

JAMA [Internet]. 2020 Jan;323(6):548–60. Available from:

https://pubmed.ncbi.nlm.nih.gov/32044947/

6. López-Cruz L, Miguel NS, Carratalá-Ros C, Monferrer L, Salamone JD, Correa M.

Dopamine depletion shifts behavior from activity based reinforcers to more sedentary

ones and adenosine receptor antagonism reverses that shift: Relation to ventral striatum

DARPP32 phosphorylation patterns. Neuropharmacology [Internet]. 2018 Jan;138:349–

59. Available from: https://pubmed.ncbi.nlm.nih.gov/29408363/

7.   Steiner H, Kitai ST. Unilateral striatal dopamine depletion: time-dependent effects on

763   cortical function and behavioural correlates. Eur J Neurosci [Internet]. 2001;14(8):1390–

764   404. Available from: https://pubmed.ncbi.nlm.nih.gov/11703467/

765   8.   Rohwedder A, Wenz NL, Stehle B, Huser A, Yamagata N, Zlatic M, et al. Four

766   Individually Identified Paired Dopamine Neurons Signal Reward in Larval Drosophila.

767   Curr Biol [Internet]. 2016 Jan;26(5):661–9. Available from:

768   https://pubmed.ncbi.nlm.nih.gov/26877086/

769   9.   Budnik V, White K. Genetic dissection of dopamine and serotonin synthesis in the

770   nervous system of Drosophila melanogaster. J Neurogenet [Internet]. 1987 Jan;4(6):309–

771   14. Available from: https://www.tandfonline.com/doi/abs/10.3109/01677068709167191

772   10.   Livingstone MS, Tempel BL. Genetic dissection of monoamine neurotransmitter synthesis

773   in Drosophila. Nature [Internet]. 1983;303(5912):67–70. Available from:

774   https://pubmed.ncbi.nlm.nih.gov/6133219/

775   11.   Meiser J, Weindl D, Hiller K. Complexity of dopamine metabolism. Cell Commun Signal

776   [Internet]. 2013;11(1). Available from: https://pubmed.ncbi.nlm.nih.gov/23683503/

777   12.   Yamamoto S, Seto ES. Dopamine dynamics and signaling in Drosophila: an overview of

778   genes, drugs and behavioral paradigms. Exp Anim [Internet]. 2014;63(2):107–19.

779   Available from: https://pubmed.ncbi.nlm.nih.gov/24770636/

780   13.   Wittkopp PJ, True JR, Carroll SB. Reciprocal functions of the Drosophila yellow and

781   ebony proteins in the development and evolution of pigment patterns. Development

782   [Internet]. 2002;129(8):1849–58. Available from:

783   https://pubmed.ncbi.nlm.nih.gov/11934851/

784   14.   True JR, Edwards KA, Yamamoto D, Carroll SB. Drosophila wing melanin patterns form

785   by vein-dependent elaboration of enzymatic prepatterns. Curr Biol [Internet]. 1999

786   Jan;9(23):1382–91. Available from: https://pubmed.ncbi.nlm.nih.gov/10607562/

787   15.   Spana EP, Abrams AB, Ellis KT, Klein JC, Ruderman BT, Shi AH, et al. speck, First

788   Identified in Drosophila melanogaster in 1910, Is Encoded by the Arylalkalamine N-

789   Acetyltransferase (AANAT1) Gene. G3 (Bethesda) [Internet]. 2020 Jan;10(9):3387–98.

790   Available from: https://pubmed.ncbi.nlm.nih.gov/32709620/

791   16.   Gervasi N, Tchénio P, Preat T. PKA dynamics in a Drosophila learning center:

792   coincidence detection by rutabaga adenylyl cyclase and spatial regulation by dunce

793   phosphodiesterase. Neuron [Internet]. 2010 Jan;65(4):516–29. Available from:

794   https://pubmed.ncbi.nlm.nih.gov/20188656/

795   17.   Heisenberg M, Borst A, Wagner S, Byers D. Drosophila mushroom body mutants are

796   deficient in olfactory learning. J Neurogenet [Internet]. 1985;2(1):1–30. Available from:

797   https://pubmed.ncbi.nlm.nih.gov/4020527/

798   18.   Jenett A, Rubin GM, Ngo TTB, Shepherd D, Murphy C, Dionne H, et al. A GAL4-driver

799   line resource for Drosophila neurobiology. Cell Rep [Internet]. 2012 Jan;2(4):991–1001.

800   Available from: https://pubmed.ncbi.nlm.nih.gov/23063364/

801   19.   Frighetto G, Zordan MA, Castiello U, Megighian A, Martin JR. Dopamine Modulation of

802   Drosophila Ellipsoid Body Neurons, a Nod to the Mammalian Basal Ganglia. Front

803   Physiol [Internet]. 2022 Jan;13. Available from:

804   https://pubmed.ncbi.nlm.nih.gov/35492587/

805   20.   Xie T, Ho MCW, Liu Q, Horiuchi W, Lin CC, Task D, et al. A Genetic Toolkit for Dissecting

806   Dopamine Circuit Function in Drosophila. Cell Rep. 2018 Apr 10;23(2):652–65.

807   21.   Clark IE, Dodson MW, Jiang C, Cao JH, Huh JR, Seol JH, et al. Drosophila pink1 is

808   required for mitochondrial function and interacts genetically with parkin. Nature [Internet].

809   2006 Jan;441(7097):1162–6. Available from: https://pubmed.ncbi.nlm.nih.gov/16672981/

810   22.   Mummery-Widmer JL, Yamazaki M, Stoeger T, Novatchkova M, Bhalerao S, Chen D, et

811   al. Genome-wide analysis of Notch signalling in Drosophila by transgenic RNAi. Nature

[Internet]. 2009 Jan;458(7241):987–92. Available from:

https://pubmed.ncbi.nlm.nih.gov/19363474/

23.   Pendleton RG, Rasheed A, Sardina T, Tully T, Hillman R. Effects of Tyrosine

Hydroxylase Mutants on Locomotor Activity in Drosophila: A Study in Functional

Genomics. Vol. 32, Behavior Genetics. 2002.

24.   Wright TRF, Bewleyz GC, Sherald3 AF. THE GENETICS OF DOPA DECARBOXYLASE

IN DROSOPHILA MELANOGASTER. 11. ISOLATION AND CHARACTERIZATION OF

DOPA-DECARBOXY LASE-DEFICIENT MUTANTS AND THEIR RELATIONSHIP TO

THE a-METHYL-DOPA-HYPERSENSITIVE MUTANTS. Vol. 8, Genetics. 1976.

25.   Ni JQ, Liu LP, Binari R, Hardy R, Shim HS, Cavallaro A, et al. A Drosophila resource of

transgenic RNAi lines for neurogenetics. Genetics [Internet]. 2009 Jan;182(4):1089–100.

Available from: https://pubmed.ncbi.nlm.nih.gov/19487563/

26.   Pan Y, Li W, Deng Z, Sun Y, Ma X, Liang R, et al. Myc suppresses male-male courtship

in Drosophila. EMBO J [Internet]. 2022 Jan;41(7). Available from:

https://pubmed.ncbi.nlm.nih.gov/35167135/

27.   Riemensperger T, Issa AR, Pech U, Coulom H, Nguyễn MV, Cassar M, et al. A single

dopamine pathway underlies progressive locomotor deficits in a Drosophila model of

Parkinson disease. Cell Rep [Internet]. 2013 Jan;5(4):952–60. Available from:

https://pubmed.ncbi.nlm.nih.gov/24239353/

28.   Brand H, Perrimon N. Targeted gene expression as a means of altering cell fates and

generating dominant phenotypes. 1993.

29.   Bayersdorfer F, Voigt A, Schneuwly S, Botella JA. Dopamine-dependent

neurodegeneration in Drosophila models of familial and sporadic Parkinson's disease.

Neurobiol Dis [Internet]. 2010 Jan;40(1):113–9. Available from:

https://pubmed.ncbi.nlm.nih.gov/20211259/

30. Hovemann BT, Ryseck RP, Walldorf U, Störtkuhl KF, Dietzel ID, Dessen E. The

Drosophila ebony gene is closely related to microbial peptide synthetases and shows

specific cuticle and nervous system expression. Gene [Internet]. 1998 Jan;221(1):1–9.

Available from: https://pubmed.ncbi.nlm.nih.gov/9852943/

31. Richardt A, Rybak J, Störtkuhl KF, Meinertzhagen IA, Hovemann BT. Ebony protein in

the Drosophila nervous system: optic neuropile expression in glial cells. J Comp Neurol

[Internet]. 2002 Jan;452(1):93–102. Available from:

https://pubmed.ncbi.nlm.nih.gov/12205712/

32. Yamamoto-Hino M, Yoshida H, Ichimiya T, Sakamura S, Maeda M, Kimura Y, et al.

Phenotype-based clustering of glycosylation-related genes by RNAi-mediated gene

silencing. Genes to Cells. 2015 Jun 1;20(6):521–42.

33. Riemensperger T, Isabel G, Coulom H, Neuser K, Seugnet L, Kume K, et al. Behavioral

consequences of dopamine deficiency in the Drosophila central nervous system. Proc

Natl Acad Sci U S A. 2011 Jan;108(2):834–9.

34. Chiu JC, Low KH, Pike DH, Yildirim E, Edery I. Assaying locomotor activity to study

circadian rhythms and sleep parameters in Drosophila. Journal of Visualized

Experiments. 2010;(43).

35. Dietzl G, Chen D, Schnorrer F, Su KC, Barinova Y, Fellner M, et al. A genome-wide

transgenic RNAi library for conditional gene inactivation in Drosophila. Nature [Internet].

2007 Jan;448(7150):151–6. Available from: https://pubmed.ncbi.nlm.nih.gov/17625558/

36. Ni JQ, Zhou R, Czech B, Liu LP, Holderbaum L, Yang-Zhou D, et al. A genome-scale

shRNA resource for transgenic RNAi in Drosophila. Nat Methods [Internet]. 2011

Jan;8(5):405–7. Available from: https://pubmed.ncbi.nlm.nih.gov/21460824/

37. Vissers JHA, Manning SA, Kulkarni A, Harvey KF. A Drosophila RNAi library modulates

Hippo pathway-dependent tissue growth. Nature Communications 2016 7:1 [Internet].

2016 Jan;7(1):1–6. Available from: https://www.nature.com/articles/ncomms10368

38.   Shafer OT, Keene AC. The Regulation of Drosophila Sleep. Vol. 31, Current Biology. Cell

Press; 2021. p. R38–49.

39.   Diao F, Ironfield H, Luan H, Diao F, Shropshire WC, Ewer J, et al. Plug-and-play genetic

access to drosophila cell types using exchangeable exon cassettes. Cell Rep [Internet].

2015 Jan;10(8):1410–21. Available from: https://pubmed.ncbi.nlm.nih.gov/25732830/

40.   Lee PT, Zirin J, Kanca O, Lin WW, Schulze KL, Li-Kroeger D, et al. A gene-specific T2A-

GAL4 library for Drosophila. Elife [Internet]. 2018 Jan;7. Available from:

https://pubmed.ncbi.nlm.nih.gov/29565247/

41.   Kanca O, Zirin J, Garcia-Marques J, Knight SM, Yang-Zhou D, Amador G, et al. An

efficient CRISPR-based strategy to insert small and large fragments of DNA using short

homology arms. VijayRaghavan K, VijayRaghavan K, editors. Elife [Internet]. 2019

Nov;8:e51539. Available from: https://doi.org/10.7554/eLife.51539

42.   Janssens J, Aibar S, Taskiran II, Ismail JN, Gomez AE, Aughey G, et al. Decoding gene

regulation in the fly brain. Nature [Internet]. 2022 Jan;601(7894):630–6. Available from:

https://pubmed.ncbi.nlm.nih.gov/34987221/

43.   Li H, Janssens J, de Waegeneer M, Kolluru SS, Davie K, Gardeux V, et al. Fly Cell Atlas:

A single-nucleus transcriptomic atlas of the adult fruit fly. Science (1979). 2022 Mar

4;375(6584).

44.   Mao Z, Davis RL. Eight different types of dopaminergic neurons innervate the Drosophila

mushroom body neuropil: Anatomical and physiological heterogeneity. Front Neural

Circuits. 2009 Jan;3(JUL):5.

45.   Morin X, Daneman R, Zavortink M, Chia W. A protein trap strategy to detect GFP-tagged

proteins expressed from their endogenous loci in Drosophila [Internet]. Available from:

www.pnas.orgcgidoi10.1073pnas.261408198

46.   Buszczak M, Paterno S, Lighthouse D, Bachman J, Planck J, Owen S, et al. The

888   carnegie protein trap library: A versatile tool for drosophila developmental studies.

889   Genetics. 2007 Mar;175(3):1505–31.

890   47.   Voet M Van Der, Harich B, Franke B, Schenck A. ADHD-associated dopamine

891   transporter, latrophilin and neurofibromin share a dopamine-related locomotor signature

892   in Drosophila. Mol Psychiatry. 2016 Jan;21(4):565–73.

893   48.   Kume K, Kume S, Park SK, Hirsh J, Jackson FR. Dopamine is a regulator of arousal in

894   the fruit fly. Journal of Neuroscience. 2005 Jan;25(32):7377–84.

895   49.   Nall AH, Shakhmantsir I, Cichewicz K, Birman S, Hirsh J, Sehgal A. Caffeine promotes

896   wakefulness via dopamine signaling in Drosophila. Sci Rep. 2016 Jan;6.

897   50.   Ni JQ, Markstein M, Binari R, Pfeiffer B, Liu LP, Villalta C, et al. Vector and parameters

898   for targeted transgenic RNA interference in Drosophila melanogaster. Nature Methods

899   2008 5:1 [Internet]. 2007 Jan;5(1):49–51. Available from:

900   https://www.nature.com/articles/nmeth1146

901   51.   McGary KL, Park TJ, Woods JO, Cha HJ, Wallingford JB, Marcotte EM. Systematic

902   discovery of nonobvious human disease models through orthologous phenotypes. Proc

903   Natl Acad Sci U S A. 2010 Apr 6;107(14):6544–9.

904   52.   Nakashima A, Ohnuma S, Kodani Y, Kaneko YS, Nagasaki H, Nagatsu T, et al. Inhibition

905   of deubiquitinating activity of USP14 decreases tyrosine hydroxylase phosphorylated at

906   Ser19 in PC12D cells. Biochem Biophys Res Commun [Internet]. 2016 Jan;472(4):598–

907   602. Available from: https://pubmed.ncbi.nlm.nih.gov/26969276/

908   53.   Kawahata I, Ohtaku S, Tomioka Y, Ichinose H, Yamakuni T. Dopamine or biopterin

909   deficiency potentiates phosphorylation at (40)Ser and ubiquitination of tyrosine

910   hydroxylase to be degraded by the ubiquitin proteasome system. Biochem Biophys Res

911   Commun [Internet]. 2015 Jan;465(1):53–8. Available from:

912   https://pubmed.ncbi.nlm.nih.gov/26225746/

54.   Birman S, Morgan B, Anzivino M, Hirsh J. A novel and major isoform of tyrosine

hydroxylase in Drosophila is generated by alternative RNA processing. Journal of

Biological Chemistry. 1994 Jan;269(42):26559–67.

55.   Vié A, Cigna M, Toci R, Birman S. Differential regulation of Drosophila tyrosine

hydroxylase isoforms by dopamine binding and cAMP-dependent phosphorylation. J Biol

Chem [Internet]. 1999 Jan;274(24):16788–95. Available from:

https://pubmed.ncbi.nlm.nih.gov/10358021/

56.   Gervasi NM, Scott SS, Aschrafi A, Gale J, Vohra SN, Macgibeny MA, et al. The local

expression and trafficking of tyrosine hydroxylase mRNA in the axons of sympathetic

neurons. RNA [Internet]. 2016 Jan;22(6):883–95. Available from:

https://pubmed.ncbi.nlm.nih.gov/27095027/

57.   Freeman A, Pranski E, Miller RD, Radmard S, Bernhard D, Jinnah HA, et al. Sleep

fragmentation and motor restlessness in a Drosophila model of Restless Legs Syndrome.

Current Biology [Internet]. 2012 Jan;22(12):1142. Available from:

/pmc/articles/PMC3381864/

58.   Chaudhuri A, Bowling K, Funderburk C, Lawal H, Inamdar A, Wang Z, et al. Interaction of

Genetic and Environmental Factors in a Drosophila Parkinsonism Model. The Journal of

Neuroscience [Internet]. 2007 Jan;27(10):2457. Available from:

/pmc/articles/PMC6672491/

59.   Wang Z, Ferdousy F, Lawal H, Huang Z, Daigle JG, Izevbaye I, et al. Catecholamines up

integrates dopamine synthesis and synaptic trafficking. J Neurochem [Internet]. 2011

Jan;119(6):1294–305. Available from: https://pubmed.ncbi.nlm.nih.gov/21985068/

60.   Zhang YQ, Friedman DB, Wang Z, Woodruff E, Pan L, O'Donnell J, et al. Protein

expression profiling of the Drosophila fragile X mutant brain reveals up-regulation of

monoamine synthesis. Molecular and Cellular Proteomics [Internet]. 2005 Jan;4(3):278–

90. Available from: http://www.mcponline.org/article/S1535947620326864/fulltext

61.   Wicker-Thomas C, Hamann M. Interaction of dopamine, female pheromones, locomotion

and sex behavior in Drosophila melanogaster. J Insect Physiol. 2008 Oct;54(10–

11):1423–31.

62.   Takahashi A. Pigmentation and behavior: potential association through pleiotropic genes

in Drosophila. Vol. 88, Genes Genet. Syst. 2013.

63.   Letizia A, Bario R, Campuzano S. Antagonistic and cooperative actions of the EGFR and

Dpp pathways on the iroquois genes regulate Drosophila mesothorax specification and

patterning. Development [Internet]. 2007 Jan;134(7):1337–46. Available from:

https://pubmed.ncbi.nlm.nih.gov/17329358/

64.   Lu J, Wang Y, Wang X, Wang D, Pflugfelder GO, Shen J. The Tbx6 Transcription Factor

Dorsocross Mediates Dpp Signaling to Regulate Drosophila Thorax Closure. Int J Mol Sci

[Internet]. 2022 Jan;23(9). Available from: https://pubmed.ncbi.nlm.nih.gov/35562934/

65.   Zeitlinger J, Bohmann D. Thorax closure in Drosophila: involvement of Fos and the JNK

pathway. Development [Internet]. 1999 Jan;126(17):3947–56. Available from:

https://pubmed.ncbi.nlm.nih.gov/10433922/

66.   Massey JH, Akiyama N, Bien T, Dreisewerd K, Wittkopp PJ, Yew JY, et al. Pleiotropic

Effects of ebony and tan on Pigmentation and Cuticular Hydrocarbon Composition in

Drosophila melanogaster. Front Physiol. 2019 Jan;10(MAY):518.

67.   Zhang S, Wang R, Wang G. Impact of Dopamine Oxidation on Dopaminergic

Neurodegeneration. ACS Chem Neurosci [Internet]. 2019 Jan;10(2):945–53. Available

from: https://pubs.acs.org/doi/abs/10.1021/acschemneuro.8b00454

68.   Davla S, Artiushin G, Li Y, Chitsaz D, Li S, Sehgal A, et al. AANAT1 functions in

astrocytes to regulate sleep homeostasis. Elife [Internet]. 2020 Jan;9:1–48. Available

from: https://pubmed.ncbi.nlm.nih.gov/32955431/

69.   DeAngelis MW, McGhie EW, Coolon JD, Johnson RI. Mask, a component of the Hippo

964   pathway, is required for Drosophila eye morphogenesis. Dev Biol. 2020 Aug 1;464(1):53–

965   70.

966   70.   Zhu M, Zhang S, Tian X, Wu C. Mask mitigates MAPT- and FUS-induced degeneration

967   by enhancing autophagy through lysosomal acidification. Autophagy. 2017 Nov

968   2;13(11):1924–38.

969   71.   Martinez D, Zhu M, Guidry JJ, Majeste N, Mao H, Yanofsky ST, et al. Mask, the

970   Drosophila ankyrin repeat and KH domain-containing protein, affects microtubule stability.

971   J Cell Sci. 2021 Oct 1;134(20).

972   72.   Dubowy C, Sehgal A. Circadian rhythms and sleep in Drosophila melanogaster.

973   Genetics. 2017 Jan;205(4):1373–97.

974   73.   Sidor CM, Brain R, Thompson BJ. Mask proteins are cofactors of Yorkie/YAP in the

975   Hippo pathway. Curr Biol [Internet]. 2013 Jan;23(3):223–8. Available from:

976   https://pubmed.ncbi.nlm.nih.gov/23333315/

977   74.   Sansores-Garcia L, Atkins M, Moya IM, Shahmoradgoli M, Tao C, Mills GB, et al. Mask is

978   required for the activity of the Hippo pathway effector Yki/YAP. Curr Biol [Internet]. 2013

979   Jan;23(3):229–35. Available from: https://pubmed.ncbi.nlm.nih.gov/23333314/

980   75.   Smith RK, Carroll PM, Allard JD, Simon MA. MASK, a large ankyrin repeat and KH

981   domain-containing protein involved in Drosophila receptor tyrosine kinase signaling.

982   Development [Internet]. 2002;129(1):71–82. Available from:

983   https://pubmed.ncbi.nlm.nih.gov/11782402/

984   76.   Zhu M, Li X, Tian X, Wu C. Mask loss-of-function rescues mitochondrial impairment and

985   muscle degeneration of Drosophila pink1 and parkin mutants. Hum Mol Genet [Internet].

986   2015 Jan;24(11):3272–85. Available from: https://pubmed.ncbi.nlm.nih.gov/25743185/

77.   Lusk JB, Lam VYM, Tolwinski NS. Epidermal Growth Factor Pathway Signaling in

Drosophila Embryogenesis: Tools for Understanding Cancer. Cancers (Basel) [Internet].

2017 Jan;9(2). Available from: /pmc/articles/PMC5332939/

78.   Oh H, Irvine KD. Yorkie: the final destination of Hippo signaling. Trends Cell Biol

[Internet]. 2010 Jan;20(7):410. Available from: /pmc/articles/PMC2919348/

79.   Sen A, Kalvakuri S, Bodmer R, Cox RT. Clueless, a protein required for mitochondrial

function, interacts with the PINK1-Parkin complex in Drosophila. Dis Model Mech

[Internet]. 2015 Jan;8(6):577–89. Available from:

https://pubmed.ncbi.nlm.nih.gov/26035866/

80.   Wang ZH, Clark C, Geisbrecht ER. Drosophila clueless is involved in Parkin-dependent

mitophagy by promoting VCP-mediated Marf degradation. Hum Mol Genet [Internet].

2016 Jan;25(10):1946–64. Available from: https://pubmed.ncbi.nlm.nih.gov/26931463/

81.   Sen A, Cox RT. Clueless is a conserved ribonucleoprotein that binds the ribosome at the

mitochondrial outer membrane. Biol Open [Internet]. 2016 Jan;5(2):195–203. Available

from: https://pubmed.ncbi.nlm.nih.gov/26834020/

82.   Inagaki HK, De-Leon SBT, Wong AM, Jagadish S, Ishimoto H, Barnea G, et al.

Visualizing Neuromodulation In Vivo: TANGO-Mapping of Dopamine Signaling Reveals

Appeptite Control of Sugar Sensing. Cell [Internet]. 2012 Jan;148(3):583. Available from:

/pmc/articles/PMC3295637/

83.   Sun F, Zhou J, Dai B, Qian T, Zeng J, Li X, et al. Next-generation GRAB sensors for

monitoring dopaminergic activity in vivo. Nat Methods [Internet]. 2020 Jan;17(11):1156.

Available from: /pmc/articles/PMC7648260/

84.   Kasture AS, Hummel T, Sucic S, Freissmuth M. Big Lessons from Tiny Flies: Drosophila

melanogaster as a Model to Explore Dysfunction of Dopaminergic and Serotonergic

Neurotransmitter Systems. Int J Mol Sci [Internet]. 2018 Jan;19(6). Available from:

https://pubmed.ncbi.nlm.nih.gov/29914172/

85.   Calleja M, Herranz H, Estella C, Casal J, Lawrence P, Simpson P, et al. Generation of

1014   medial and lateral dorsal body domains by the pannier gene of Drosophila. Development

1015   [Internet]. 2000;127(18):3971–80. Available from:

1016   https://pubmed.ncbi.nlm.nih.gov/10952895/

1017   86.   Kennerdell JR, Carthew RW. Heritable gene silencing in Drosophila using double-

1018   stranded RNA. Nat Biotechnol [Internet]. 2000;18(8):896–8. Available from:

1019   https://pubmed.ncbi.nlm.nih.gov/10932163/

1020   87.   Yamamoto S, Charng WL, Rana NA, Kakuda S, Jaiswal M, Bayat V, et al. A mutation in

1021   EGF repeat-8 of notch discriminates between serrate/jagged and delta family ligands.

1022   Science (1979). 2012 Nov 30;338(6111):1229–32.

1023   88.   Marcogliese PC, Deal SL, Andrews J, Harnish JM, Bhavana VH, Graves HK, et al.

1024   Drosophila functional screening of de novo variants in autism uncovers damaging

1025   variants and facilitates discovery of rare neurodevelopmental diseases. Cell Rep. 2022

1026   Mar 15;38(11).

1027   89.   Snel B, Lehmann G, Bork P, Huynen MA. STRING: a web-server to retrieve and display

1028   the repeatedly occurring neighbourhood of a gene [Internet]. Vol. 28, Nucleic Acids

1029   Research. 2000. Available from: http://www.bork.embl-heidelberg.de/STRING

1030   90.   Szklarczyk D, Gable AL, Nastou KC, Lyon D, Kirsch R, Pyysalo S, et al. The STRING

1031   database in 2021: Customizable protein-protein networks, and functional characterization

1032   of user-uploaded gene/measurement sets. Nucleic Acids Res. 2021 Jan 8;49(D1):D605–

1033   12.

1034   91.   Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, et al. The STRING

1035   database in 2023: protein-protein association networks and functional enrichment

1036   analyses for any sequenced genome of interest. Nucleic Acids Res. 2023 Jan 6;51(1

1037   D):D638–46.

92.   Hu Y, Flockhart I, Vinayagam A, Bergwitz C, Berger B, Perrimon N, et al. An integrative

approach to ortholog prediction for disease-focused and other functional studies. BMC

Bioinformatics. 2011 Aug 31;12.

93.   Hamosh A, Amberger JS, Bocchini C, Scott AF, Rasmussen SA. Online Mendelian

Inheritance in Man (OMIM®): Victor McKusick's magnum opus. Am J Med Genet A. 2021

Nov 1;185(11):3259–65.

94.   Abrahams BS, Arking DE, Campbell DB, Mefford HC, Morrow EM, Weiss LA, et al.

SFARI Gene 2.0: A community-driven knowledgebase for the autism spectrum disorders

(ASDs). Mol Autism. 2013;4(1).

95.   Cichewicz K, Garren EJ, Adiele C, Aso Y, Wang Z, Wu M, et al. A new brain dopamine-

deficient Drosophila and its pharmacological and genetic rescue. Genes Brain Behav.

2017 Mar 1;16(3):394–403.

96.   Hardie SL, Hirsh J. An improved method for the separation and detection of biogenic

amines in adult Drosophila brain extracts by high performance liquid chromatography. J

Neurosci Methods. 2006 Jun 15;153(2):243–9.

1053

1054 Supplemental Information

1055 Document S1. Figures S1-S10

1056 Table S2. Excel file containing data too large to fit in a PDF Table S1-S5

1057

S1 Fig. GO term enrichment analysis performed on the pigmentation screen hits.

S2 Fig. Comparison between genes identified as hits from our pigmentation screen to

those tested in the original thorax screen (Mummery-Widmer et al., 2009).

S3 Fig. Comparison of dopamine levels between lacZ RNAi and mCh RNAi lines in the fly

head.

S4 Fig. Validation of recombined GAL4 lines used for dopamine measurement in the brain.

S5 Fig. Brain data for prioritized genes with no change in brain dopamine.

S6 Fig. Additional clusters of TH positive neurons upon *mask* knockdown show trend or

little effect.

S7 Fig. Sleep behavior analysis upon TH GAL4 driven TH knockdown.

S8 Fig. Additional sleep analysis from TH-GAL4 driven *mask* knockdown. (

S9 Fig. Sleep behavior analysis of *clueless* knockdown highlights a different pathway

than mask.

S10 Fig. Comparison of RNAi lines from the pigmentation screen

S11 Fig. Expression analysis for other prioritized candidate genes that did not show any

significant change in total head dopamine.

S1 Table. Pigmentation screen results by RNAi line

S2 Table. Drosophila Activity Monitor results by RNAi line

S3 Table. HPLC results by RNAi line

S4 Table. Prioritized gene list for HPLC

S5 Table. All RNAi lines used with stock center information