RNAi-based screen for cuticle pigmentation in *Drosophila melanogaster* reveals novel regulators of brain dopamine

Samantha L Deal[1,2], Danqing Bei[2,3], Shelley B Gibson[2,3], Harim Delgado-Seo[2,4], Yoko Fujita[2,3],

Kyla Wilwayco[2,3], Elaine S Seto[5,6,7], Shinya Yamamoto[1,2,3,4,8,#]

[1] Program in Developmental Biology, Baylor College of Medicine (BCM), Houston, TX 77030, USA

[2] Jan and Dan Duncan Neurological Research Institute (NRI), Texas Children's Hospital (TCH), Houston, TX 77030, USA

[3] Department of Molecular and Human Genetics, BCM, Houston, TX 77030, USA

[4] Department of Neuroscience, BCM, Houston, TX 77030 USA

[5] Department of Neurology, TCH, Houston, TX 77030, USA

[6] Epilepsy Center, TCH, Houston, TX 77030, USA

[7] Department of Neurophysiology, TCH, Houston, TX 77030, USA

[8] Development, Disease Models & Therapeutics Graduate Program, BCM, Houston, TX 77030, USA

#Corresponding Author: Shinya Yamamoto

 E-mail: yamamoto@bcm.edu; Phone: 832-824-8119; Fax: N/A

Keywords: *Drosophila melanogaster*, dopamine, pigmentation, RNAi, screen, *clueless*, *mask*

Abstract

The dopaminergic system has been extensively studied for its role in behavior in various animals as well as human neuropsychiatric and neurological diseases. However, we still know little about how dopamine levels are tightly regulated *in vivo*. In order to identify novel regulators of dopamine levels, we utilized *Drosophila melanogaster* cuticle pigmentation as a readout, where dopamine is used as a precursor to melanin. First, we measured dopamine from classical mutants of genes that are known to be important for cuticle pigmentation to understand the relationship between dopamine levels and cuticle color. We then performed an RNAi-based screen to identify novel regulators of cuticle pigmentation. We identified 153 genes that were not only enriched for conserved homologs and disease-associated genes, but were unexpectedly enriched for multiple developmental signaling pathways and mitochondria-associated proteins. Upon measuring dopamine from 35 prioritized candidates from this cohort, we found 10 that caused significant reduction in dopamine in the head while one caused an increase. While most of these 11 genes are expressed in the fly brain, only two of them, *clu* and *mask,* altered dopamine levels specifically in the brain upon knockdown in dopaminergic cells. Interestingly, *mask* may act as a hub of dopamine regulation since it is associated with three nodes found in our screen (Receptor tyrosine kinase/EGF signaling, Hippo signaling, and mitochondrial dynamics). Further examination suggests that Mask likely acts on dopamine synthesis by regulating transcription of the rate-limiting dopamine synthesis enzyme, *tyrosine hydroxylase*. In conclusion, this screen identified 11 new regulators of dopamine levels and provides molecular handles to investigate how dopamine levels are controlled *in vivo*, as well as revealed an unexpected relationship between fly pigmentation genes, developmental signaling and human neurological and neurodevelopmental disease genes.

Introduction

Dopamine is a conserved neurotransmitter that functions in the nervous system to regulate variety of behaviors, including learning & memory, locomotion, reward, and sleep. In humans, disruptions in dopamine have been associated with neurological and neuropsychiatric disorders, including addiction, depression, sleep disorders, and schizophrenia (Grace 2016; Nutt et al. 2015; Radwan et al. 2019). In addition, one of the hallmarks of Parkinson's disease is early degeneration of dopaminergic neurons, and Parkinson's patients are often given a dopamine precursor, l-dihydroxyphenylalanine (L-DOPA), as a method for treating symptoms (Church 2021; Armstrong and Okun 2020). Most genetic research on dopamine biology has focused on dopaminergic signaling and transport; however, changes in dopamine levels can have a great impact on behavioral outcomes (López-Cruz et al. 2018; Steiner and Kitai 2001; Rohwedder et al. 2016). Thus, identifying genes that affect dopamine levels is critical to understanding and potentially treating dopamine-associated disorders.

Dopamine synthesis is a conserved process, whereby the amino acid tyrosine gets converted into L-DOPA via the rate-limiting enzyme, Tyrosine hydroxylase (TH) (Budnik and White 1987). Then, L-DOPA is converted into dopamine via the enzyme Dopa decarboxylase (Ddc) (Livingstone and Tempel 1983), which is also an important enzyme for the biosynthesis of serotonin. In mammals, dopamine can be degraded either via oxidation or methylation using the enzymes Monoamine oxidase (MAO) and Catechol-O-methyltransferase (COMT), respectively (Meiser et al., 2013). In invertebrates, the primary methods for dopamine degradation are thought to be β-alanylation via the β-alanyl amine synthase (encoded by the *ebony* gene in *Drosophila melanogaster*), and acetylation via the acetyltransferase (encoded by the *speck* gene in *Drosophila*) (Yamamoto and Seto 2014). However, insects also convert dopamine into melanin to form the proper cuticle pigment and structure. In fact, changes in genes that are involved in dopamine metabolism lead to changes in cuticle color (Spana et al. 2020; True et al. 1999;

Wittkopp et al. 2002). In other words, knockdown of *TH* (known as the *pale (ple)* gene in

Drosophila) or *Ddc* leads to a paler cuticle, and mutations or knockdown of *ebony* or *speck* leads

to a darker cuticle.

Historically, *Drosophila* has been used to study dopamine biology in multiple contexts.

Early establishment of behavioral assays in the fly identified that the dopaminergic neurons

(DANs) that innervate the mushroom body are critical for learning and memory (Gervasi et al.

2010; Heisenberg et al. 1985). Over the years, an extensive genetic toolkit has been generated

in the fruit fly, making it a great system for dissecting dopaminergic neural circuitry and signaling

involved in many behavioral contexts (Jenett et al. 2012; Frighetto et al. 2022; Xie et al. 2018). In

addition to the fly's contribution to understanding dopaminergic circuitry, this model organism has

also been used to identify and investigate the cellular functions of Parkinson disease's genetic

risk factors such as *PINK1* and *PRKN* (i.e. *Pink1* and *parkin* in the fruit fly, Clark et al. 2006). The

conservation of systems and mechanisms in neural biology along with the efficiency of working in

Drosophila has made it a great system for large-scale screening of novel genes involved in a

phenotype of interest. Previously, a large-scale RNAi screen (11,619 genes, ~89% of genome)

was performed to identify genes that affect mechanosensory organs and cuticle formation in the

Drosophila thorax (Mummery-Widmer et al. 2009). In this screen, the researchers found 458

genes that affected cuticle color upon knockdown, but no further validation was performed.

In this manuscript, we attempted to utilize the fruit fly cuticle to identify novel regulators of

dopamine. We started by systematically examining classical pigmentation genes for their effect

on dopamine in the head and brain. From this we found that genes that are involved in dopamine

synthesis are the only genes with expected effects on dopamine; genes involved in degradation

of dopamine either have no effect or unexpectedly show reduced dopamine. We next went on to

characterize the 458 genes identified from the RNAi screen by Mummery-Widmer et al. (2009).

We tested 330 of them with independent RNAi lines and found 153 genes (~46%) with consistent

cuticle pigmentation defects. After prioritizing these genes based on their cuticle effects, potential

links to human diseases and their roles in the brain, we examined their effects on dopamine in

the head and brain. From this analysis, we found two genes, *mask* and *clu*, that upon knockdown

significantly reduce dopamine in the brain, and further show that *mask* affects mRNA and protein

levels of TH.

Results

Drosophila pigmentation genes affect dopamine in unexpected ways

To systematically test if changes in pigmentation genes affect dopamine as theorized in

the field, we characterized loss-of-function alleles of genes involved in dopamine metabolism

(Figure 1A) for pigmentation phenotypes and dopamine levels in the head and brain. The

enzymes required for dopamine synthesis, TH and Ddc, are essential for survival (Pendleton et

al. 2002; Wright et al. 1976). Thus, we took advantage of established RNAi lines (Ni et al. 2009;

Pan et al. 2022; Riemensperger et al. 2013) and UAS/GAL4 system (Brand and Perrimon 1993)

to examine their effects. It is well documented and accepted that reduction of *TH* or *Ddc* levels or

activity causes pale pigmentation and reduction of dopamine levels (True et al. 1999; Livingstone

and Tempel 1983; Budnik and White 1987). By knocking down *TH* using a UAS-RNAi transgene

that have been previously validated using a *TH-GAL4* driver (Bayersdorfer et al. 2010), the effect

of which is highlighted by the strength of the pale cuticle phenotype (Figure 1B-C), we found total

head dopamine levels to be reduced by ~60% (Figure 1E) and brain dopamine levels by ~32%

(Figure 1E'). On the other hand, *Ddc* knockdown causes a weaker effect than *TH* RNAi in the

cuticle (Figure 1B,D) and in the head (~35% dopamine reduction, Figure 1E), and we did not

observe a significant effect on dopamine in the brain (Figure 1E'). This may be due to the strength

of the RNAi and/or enzyme level requirements (since TH is known to be the rate-limiting enzyme

with a more restricted expression pattern).

To test the genes that have been implicated in dopamine degradation and cuticle

development (pigmentation and structural rigidity), we examined mutants that are thought to be

null or strong hypomorphic alleles. These alleles cause darker or paler cuticle formation (Figure

1F-1L, 1F'-1L') depending on their role in the melanin formation. While we expected to see

elevated levels of dopamine or possibly no effect for most of these alleles, *black*, *speck*, *straw*

and *yellow* mutants caused a reduction in dopamine in the head (Figure 1Q). Mutants for *ebony*

and *tan* on the other hand had no significant effect on head dopamine levels (Figure 1Q). In order

to identify if any of these alleles are likely to affect brain dopamine levels, we first asked if any of

them show expression in the brain using publicly available T2A-GAL4 gene trap lines (Diao et al.

2015; Lee et al. 2018; Kanca et al. 2019). Based on our own expression analysis (Figure 1 M-P)

and previously reported expression data (Hovemann et al., 1998; Richardt et al., 2002), several

of these genes are expressed in the fly brain but their expression does not appear to overlap with

DA neurons. When we measured effects of these mutants on brain dopamine levels (Figure 1R),

we found that mutations in these pigmentation genes do not significantly affect brain dopamine

levels.

Together this data shows that while changes in dopamine synthesis causes a paler cuticle,

which reflects a reduction in dopamine in the head and brain, changes in dopamine metabolism

enzymes downstream of dopamine synthesis cause pigmentation phenotypes that do not

correlate with head or brain dopamine level differences in adulthood.

RNAi-based cuticle pigmentation screen

From mining a previously published genome-wide scale RNAi screen (Mummery-Widmer

et al. 2009), we identified 458 genes with pigmentation defects that were discovered using

pannier(pnr)-GAL4, which is expressed in the dorsocentral nota. In order to check if these

phenotypes can be reproduced using independent reagents, we located 718 additional UAS-RNAi

lines for 330 of these genes from alternative sources. 426 of these lines were from the National

Institute of Genetics in Mishima, Japan (Yamamoto-Hino et al. 2015), and 292 of them came from

the Transgenic RNAi Project (TRiP) at Harvard Medical School (Ni et al. 2009). Upon screening

these RNAi's using the same *pnr-GAL4* driver, we found that 153 genes caused pigmentation

defects in at least one additional RNAi line, showing a validation of 46.4% of the hits for the

previous screen (Figure 2A).

Our RNAi-based pigmentation secondary screen produced a broad spectrum of

phenotypes (Figure 2B). When we classified these phenotypes into three categories, mild,

moderate, or strong, we found that these three categories were equally represented (mild = 38%,

moderate = 33%, strong = 29%, Figure 2D). When we scored for pigmentation color changes, we

saw three primary categories, pale, dark, or mix (a combination of pale and dark in a single

thorax). These categories were not equally represented (Figure 2E), with dark cuticle phenotypes

representing the majority (56%) of genes. Pale cuticle effects were found in 28% of genes, and

those with a mixed phenotype only represented 16%.

In order to classify what categories of genes cause pigmentation effects in our cohort, we

performed a protein-network analysis (Figure 2C). We found a group of proteins that have been

previously linked to cuticle pigmentation as expected, but we also found clusters related to RNA

processing, ubiquitin proteasome system, mitochondria-associated proteins, and signaling

pathway genes, Hippo and EGF (epidermal growth factor) signaling in particular. GO enrichment

analysis also highlighted terms associated with cuticle pigmentation, protein stability, and several

signaling pathways (Supplemental Figure 1). In addition to this, we also observed that 85% of our

genes were conserved between *Drosophila* and human (57.9% of genes are conserved between

the two species (Yamamoto et al. 2023), an enrichment from the 68% of fly genes conserved in

genes tested the original RNAi screen (Figure 2F, Supplemental Figure 2).

In summary, the cuticle pigmentation screen identified a broad spectrum of pale, dark and

mixed phenotypes in 152 genes, showing that for cuticle color ~50% of RNAi hits can be validated.

Close examination of these hits highlighted that many of them are conserved and implicated in

unexpected molecular processes, such as developmental signaling.

Many pigmentation genes isolated have a neurological consequence

To assess if any of the genes identified as hits from our pigmentation genes have a neurological consequence, and thus might cause an effect on dopamine in the brain, we checked if any of the homologs have a neurological disease association. We found that 51.2% of the fly genes with a human homolog are associated with some type of disease and of those disease-causing genes 78% have at least one disease that leads to a neurological phenotype (Table 1, Supplemental Figure 2). Among the neurological phenotypes identified, there were a broad spectrum of symptoms noted, including intellectual disability, autism spectrum disorder, ataxia, epilepsy, and cerebral atrophy.

In addition to classifying neurological consequences in homologous human genes, we directly assessed neurological consequences in the fly. Since dopamine is known to regulate locomotion and sleep (Riemensperger et al. 2011; Pendleton et al. 2002), we screened for locomotion and sleep defects using the Drosophila Activity Monitor (DAM; Chiu et al. 2010). We tested 274 RNAi lines from the NIG, TRiP, and VDRC RNAi collections (Dietzl et al. 2007; Ni et al. 2011; Vissers et al. 2016) corresponding to 152 genes that showed consistent pigmentation defects. These lines were driven using *TH-GAL4* and scored for changes in total locomotion, total sleep, sleep bout length, and sleep latency during the night and day (Figure 3). There was a broad range in the RNAi lines for each of these phenotypes (Figure 3E-H). Since we tested different RNAi lines from multiple sources that were made using different techniques, we focused on RNAi lines that behaved as "outliers" by identifying hits that were two standard deviations from the mean for a given phenotype. Each phenotype showed a similar number of "outliers" (locomotion: 23 lines, total sleep: 20 lines, sleep bout length: 21 lines, sleep latency: 25 lines). While total sleep and locomotion strongly overlapped with each other, sleep latency and sleep bout length did not overlap very well with the other sleep phenotypes (Figure 3D). Genes that affect sleep can often show distinct phenotypes that do not necessarily correspond to multiple phenotypes or times of day (Shafer and Keene 2021), so any gene with one phenotype during the day or night was

classified as an "outlier". Sixty lines, which corresponded to 50 genes, were classified as an

"outlier" for at least one phenotype.

Taken together, we found that 77 of our 132 conserved genes are likely to have a

neurological function, either because they are implicated in a human neurological disease (52

genes) or have been classified as behavioral "outlier" in our locomotion and sleep analysis (45

genes).

Dopamine levels in the fly head were significantly altered in 11 genes

To prioritize our gene list for dopamine measurement, we identified which of our genes

were conserved (130 conserved/152 pigmentation genes). Then, we prioritized these based on

whether knockdown of these genes showed a strong pigmentation effect in the cuticle (38 genes),

a behavioral effect in the DAM (45 genes), and/or a human neurological disease association (52

genes). About 72% of the genes (95/132 conserved genes) were represented in one of these

categories, and 29% (39/132 conserved genes) were represented in at least two (Figure 4A). We

classified any gene that fit into at least two of these categories as a priority hit and tested them

for changes in dopamine in the head using High Performance Liquid Chromatography (HPLC).

Using our system, we can repeat previous results that show that knocking down *TH* using an

RNAi or overexpressing a *TH* cDNA using *TH-GAL4* significantly reduces or increases head

dopamine without any effect on serotonin (Figure 4B,C, Bayersdorfer et al. 2010).

When we examined the 35 prioritized genes (note: some genes were not tested due to

lethality or stock issues), we found that 11 of them (not including *TH*) significantly alter head

dopamine levels (Figure 4D,E). We found that NIG lines showed about ~80% dopamine levels

compared to the TRiP collection (Supplemental Figure 3), so we decided to compare the DA

levels to neutral control lines appropriate for each collection (e.g. UAS-lacZ RNAi) to assess the

relative decrease or increase of dopamine. Hits from both collections show similar patterns

compared to the controls. First, unlike what we saw in cuticle pigmentation, where 65% of genes

caused a dark cuticle (excludes mixed cuticle phenotype for simplicity), most genes trended

toward a reduction in dopamine (71%, 25/35, Figure 4F,G). Among the 11 genes that significantly

alter dopamine, 10 of them (91%) significantly reduced dopamine and only one gene increased

dopamine. In addition, there was no direct correlation between the cuticle color and the dopamine

level in the fly head (Figure 4H).

In summary, our HPLC analysis on fly heads revealed 11 genes that significantly affect

dopamine levels. We saw a trend in reduction in dopamine across all lines tested as well as those

with significant changes, and we observed no correlation between cuticle color and dopamine

level, which is similar to what we observed in the classical pigmentation genes (Figure 1).

Brain analysis reveals *mask* and *clu* as potential regulators of dopamine in the

brain

To determine which of the 11 genes that affect dopamine in the head can have

consequences on the brain, we examined the expression of four of them in the brain using

available T2A-GAL4 lines (Diao et al. 2015; Lee et al. 2018; Kanca et al. 2019). We found that

each of these genes show some expression in the brain. Two of these genes show very broad

expression (*Bsg and ClC-c,* Figure 5A,B) and appear to overlap with most if not all DANs. Two

others showed a slightly more selective neuronal expression (*PIG-A* and *rk*, Figure 5C,D), and

they seem to overlap with somewhere between 30-50% of DANs. In addition, when we explored

publicly available single-cell mRNA sequencing datasets, ten of our gene hits (all except *Spn38F*)

show some expression in the brain with varying amount of overlap with TH expressing DANs

(Janssens et al. 2022; Li et al. 2022).

In order to test for changes in brain dopamine levels, we wanted to generate a line that

hits most if not all DANs. The *TH-GAL4* line commonly used for dopamine studies and in our

dopamine head analysis misses a large portion of DANs, especially in the PAM cluster (~87 DANs

missed/~142 total DANs per hemisphere; Mao & Davis 2009), making it not ideal for quantifying

total changes in dopamine. To get around this, we recombined the *TH-GAL4* with another driver

that hits a large portion of the PAM cluster neurons (*GMR58E02-GAL4 (R58-GAL4)*; Jenett et al.

2012) generating a recombined line (labelled *TH,R58-GAL4*) that hits ~95% of DANs

(Supplemental Figure 4).

Using two different RNAi lines per gene, we microdissected adult fly brains and measured

dopamine for each of these 11 gene hits. Most of them did not show a consistent change in

dopamine levels in the brain (Figure 5E-M), but *mask* (*multiple ankyrin repeats single KH domain*)

and *clu* (*clueless*) knockdown showed reduction in total brain dopamine for two independent RNAi

lines (Figure 5N,S). While we do not have a gene trap for expression analysis for either of these

genes, we obtained protein trap lines (Buszczak et al. 2007; Morin et al. 2001), allowing us to

examine where the protein is expressed within the brain. Both Clueless and Mask are found in

many cells in the brain (Figure 5). When we focused on some of the DAN clusters (e.g. PPL1 and

PPM2), we found Clueless protein to be enriched in DANs (Figure 5P-R, P'-R'). Mask on the other

hand shows a broad and equal level of expression throughout most cells in the brain (Figure 5T-

W, U'-W').

In summary, our fly brain analysis revealed that while many of these genes are expressed

in DANs, only two of them (18%, 2/11) significantly altered dopamine in the brain, highlighting that

changes in head dopamine do not necessarily reflect changes in brain dopamine.

***mask* regulates Tyrosine Hydroxylase transcript and protein levels**

First, we assessed whether reduction in dopamine caused by *clu* and *mask* knockdown

could be due to DAN loss. We quantified the number of DANs in major clusters (excluding the

PAM cluster where there is more variation in the number of DANs) based on immunostaining

against TH (Mao & Davis 2009). There was no difference in neuron number for any cluster

examined for either *clu* or *mask* knockdown (Figure 6D, Supplemental Figure 6). Next, since our

pigmentation gene study (Figure 1) revealed that the only enzyme in the dopamine metabolism

pathway that shows a significant effect on brain dopamine level upon manipulation is TH, we

examined if knocking down either *clu* or *mask* affects *TH* RNA or protein level. Knockdown of *clu*

showed an increase in TH RNA level of about 1.5-2X depending on the RNAi (Figure 6A) with no

effect on TH protein level (Figure 6C, Supplemental Figure 6). Interestingly, knockdown of *mask*

led to a significant reduction in *TH* RNA level for one RNAi (50% reduction), with the other RNAi

trending (20-30% reduction, Figure 6A). Upon performing immunostaining and examining TH

levels in different DAN clusters, two clusters showed significant reduction in TH protein level and

other clusters showed a trend in TH protein reduction (PAL and PPM2/3, Figure 6C, Supplemental

Figure 6). In conclusion, *mask* appears to reduce total dopamine levels by affecting transcription

of TH, while *clu* is acting through an alternative mechanism.

Discussion

Validation of 147 novel cuticle pigmentation genes through independent RNAi lines

Our RNAi screened confirmed 153 genes that were previously implicated in *Drosophila* pigmentation – 148 of these genes have not been previously associated with dopamine biology. Because we observed an enrichment for genes involved in pigmentation and tyrosine metabolism (Supplemental Figure 1), our method for screening and analysis effectively identified genes that play critical roles in the biological process of interest.

While prior research has compared genes that overlap with different types of screening strategies (e.g. chemical mutagenesis vs. RNAi), most RNAi phenotypes from large-scale screens have not been extensively validated with alternative RNAi lines or collections. For example, in Yamamoto et al, they identified 11 genes on the X chromosome with pigmentation defects from an EMS (ethyl methanesulfonate) mutagenesis screen; however, of the 91 X-chromosome genes identified from the RNAi screen only two of them overlapped between these screens (Yamamoto et al. 2014). Both of those genes, *ATP7* and *Hr4*, were validated in our analysis. The original RNAi screen used the lines available at the VDRC stock center, which includes hairpin RNAi constructs (averaging ~325 base pairs) that are either random transposon mediated insertions or targeted recombinase mediated insertions (Dietzl et al. 2007; Vissers et al. 2016). Alternative sources for RNAi lines (i.e. TRiP and NIG) start with different backgrounds for their transgenesis, different vectors for their insertion, different hairpin lengths, different insertion methods, and/or target different areas of the gene (Ni et al. 2011, 2007). These differences allow researchers to overcome the off-targeting seen with RNAi analysis by testing alternative lines, but to our knowledge, there have been no screens or meta-analyses to systematically test consistency across lines from different sources. In this analysis, we found that ~46% (153/330) of genes with a pigmentation phenotype showed a phenotype in an alternative RNAi. If there were more lines available for a given gene, we were more likely to see a phenotype. The genes validated had an

average of 2.5 additional lines available, and the genes with no pigmentation phenotype had an

average of 1.8 additional lines (Supplemental Figure 7). However, when we examined the number

of positive hits in genes with more than one RNAi line, there was very little decrease in the number

of positive hits (from ~77% agreement between 2 lines to ~63% agreement between 4 lines),

suggesting that more RNAi lines may not be necessary to identify positive phenotypes. We also

found 87% agreement in terms of the pigmentation color change between lines, highlighting that

pigmentation as a phenotype is relatively consistent (Supplemental Figure 7).

Interestingly, 85% of our pigmentation screen genes had at least one human homolog,

whereas the fly genome has ~60% conservation (1.42 fold enrichment) and genes tested in the

original RNAi screen list only had ~68% conservation (1.25 fold enrichment, Supplemental Figure

2). Similarly, there is an enrichment for disease-causing genes (51%/29%, 1.76 fold enrichment).

While pigmentation of the cuticle using dopamine produced through TH and Ddc is an

invertebrate-specific phenomenon (Yamamoto and Seto 2014), this enrichment for conserved

genes suggests that this phenotype is relevant for some conserved biological pathway and

mammalian biology. We propose that cuticle pigmentation in insects can be considered as a

"phenolog" (McGary et al. 2010) of certain neurological traits in mammals, a theory that requires

a more comprehensive investigation. Interestingly about 78% of genes with human homologs

from this screen are associated with a human disease with some type of neurological or

neurodevelopmental phenotype. This is further substantiated by the enrichment for certain GO

terms seen in our hits.

Protein-protein interaction network analysis highlights unexpected gene

associations between cuticle pigmentation genes

When we investigated what kind of genes were specifically found from the pigmentation

screen, we found some interesting trends, specifically clusters of genes that are associated with

ubiquitin-proteasome system (UPS), RNA processing, mitochondria, and developmental signaling

pathways. A number of mutations in the UPS have been linked to Parkinson's disease, but most

prevailing theories revolve around its effect on α-synuclein levels (Lim and Tan 2007; McKinnon

et al. 2020). However, there is some evidence that the UPS system regulates TH levels and

therefore dopamine levels in the mammalian system. For example, inhibition of the UPS system

has been shown to increase the TH protein levels in a PC12 rat cell line (Kawahata et al. 2015;

Nakashima et al. 2016). At least one TH phosphorylation site has been shown to be conserved

(Davis et al. 2007), and it is possible that the UPS is a conserved mechanism for regulating TH

levels and thus dopamine.

Changes in RNA processing could be indicative of changes in the dopamine synthesis

enzymes TH and Ddc since these enzymes undergo RNA processing to create two different

isoforms (Birman et al. 1994). One isoform is specific to the cuticle/epidermal and the other is

specific to the central nervous system. The two TH isoforms differ quite a bit in their kinetics, with

the epidermal isoform being generally more enzymatically active (Vié et al. 1999). In mammals,

post-transcriptional trafficking of TH mRNA and translational regulation has been suggested to be

important for its function. For example, TH mRNA gets localized to the axons for more direct

regulation of TH and dopamine at the axon terminal (Gervasi et al. 2016). In addition, short-term

stress can lead to elevated levels of TH mRNA and protein that often does not correlate with

transcriptional expression (Tank et al. 2008). It is also possible RNA processing factors regulate

dopamine levels in a non-specific manner. Global changes in RNA processing could lead to

cellular stress, which can significantly affect dopamine levels and dopaminergic signaling

(Imperato et al. 1989; Spanos et al. 2013). This may be dependent on the type of stressors since

while we observe a number of genes may be associated with cellular stress either directly in

through their molecular function from our screen (e.g. *sesB*, *Raf*, *Vps39*) (Kroemer et al. 2010;

Rezatabar et al. 2019; Takáts et al. 2014; Terhzaz et al. 2010; Xu et al. 2017), we did not identify

core components of the well-known stress pathway genes (e.g. JNK, PERK, GCN3). However, it

is important to note that activation of ASK1, a stress signaling integrator, is sufficient cause

darkening of the cuticle that is accompanied by increased expression of TH in Ddc (Sekine et al.

2012, 2011). Hence, our screen may not have reached saturation and there are still a number of

novel cuticle pigmentation genes that awaits discovery.

We identified 12 genes that are known to function at the mitochondria, mostly in the

electron transport chain (*ND-13B*, *ND-42*, *COX5A*), mitochondrial mRNA translation (*mRpL47*,

mRpL48, *CG4679*), and mitochondrial protein localization (*CG8728*, *clu*, *mask*, *Mpcp2*). It is well-

known that the mitochondria have a key role in dopamine metabolism in mammals since MAO

acts in the mitochondria to degrade dopamine. However, there is no known MOA homolog in flies,

and the primary mechanisms of dopamine degradation are thought to be acetylation and β-

alanylation that has not been linked to the mitochondria (Yamamoto and Seto 2014). Interestingly,

the metabolites for oxidation and methylation of dopamine (DOPAC and HVA) have been

observed in flies (Freeman et al. 2012; Chaudhuri et al. 2007; Wang et al. 2011; Zhang et al.

2005). In addition, we found mutants for enzymes that have been thought to be the primary

regulators of dopamine degradation (*ebony*, *tan*, and *speck*) (Wicker-Thomas and Hamann 2008;

Takahashi 2013) did not have any effect on dopamine in the brain (Figure 1) suggesting there is

either an alternative process regulating dopamine degradation in the brain. Researchers have

shown that pathogenic variants in *PINK1* and *PRKN* (genes that regulate mitochondrial

maintenance) can cause hereditary forms of Parkinson's Disease, and the drug MPTP (1-methyl-

4-phenyl-1,2,3,6-tetrahydropyridine) can lead to environmentally induced Parkinson's Disease,

by inhibiting Complex I activity and killing DANs (William Langston et al. 1983; Kitada et al. 1998;

Valente et al. 2004; Nicklas et al. 1985). It has been theorized that DANs are vulnerable because

of their structure or the potential of dopamine oxidizing within the cell (Burbulla et al. 2017), but

there is also some evidence that dopamine synthesis may be occurring at the mitochondria (Wang

et al. 2009). Our results show pigmentation can be affected by mitochondrial changes in the fruit

fly, and provides a molecular handle to study an unexplored novel degradation or synthesis

mechanism for dopamine at the mitochondria which may have relevance to human health.

388   The last major group of genes identified from our screen as a regulator of cuticle

pigmentation was developmental signaling, specifically EGF signaling and Hippo signaling. These

clusters of genes were not only clear from our network analysis, but were also identified based

on GO terms enrichment analysis (Supplementary Figure 1). To date, neither EGF nor Hippo

signaling have an established connection with dopamine biology in *Drosophila*. One reason we

may have pulled these genes out is because the process of forming the dorso-central thorax is

developmentally regulated. The thorax requires extensive remodeling from the wing imaginal disc

for formation, which utilizes many different key developmental signaling pathways (Letizia et al.

2007; Lu et al. 2022; Zeitlinger and Bohmann 1999). However, we did not identify core

components of the other key pathways, such as Notch or Hedgehog, suggesting that EGF and

Hippo are more specifically involved in this pigmentation process. We know that both EGF and

Hippo signaling are important to the development of the nervous system, and in mice they have

shown that both pathways can specifically affect the development of the dopaminergic system

(Yoon and Baik 2013). Since the development of most neural circuit-systems requires reiteration

for proper development (Pan and Monje 2020), it is plausible that EGF or Hippo signaling could

directly or indirectly regulate dopamine levels. Since we did not see an effect on dopamine in the

adult heads, this suggests that this may be developmentally specific. Overall, our pigmentation

screen identified unexpected pathways that could each be a potential area of pursuit in dopamine

metabolism since none have been explored in this context.

Cuticle pigmentation phenotypes do not necessarily reflect changes in dopamine

409   Biochemical analysis and pigmentation studies have established the enzymes involved in

dopamine metabolism in the fruit fly, but we are not aware of a research study that systematically

analyzed dopamine levels of classical pigmentation genes. The field has generally assumed that

a darker cuticle reflects an elevated level of dopamine (Massey et al. 2019; Spana et al. 2020).

However, when we measured dopamine from degradation enzyme mutants using the state-of-

the-art HPLC system, we saw either no effect on dopamine level or reduced level of dopamine in

the head. In addition, when we measured dopamine from the pigmentation genes from the screen,

a darker cuticle did not reflect elevated levels of dopamine in the head. While we were able to

increase dopamine by 3-5 fold upon TH cDNA overexpression in the head (Figure 4C), this

phenomenon was not observable in the brain. This result is a bit puzzling, considering that the

GAL4/UAS overexpression system typically allows one to overexpress a gene of interest well

outside normal conditions (often 20-30 fold mRNA expression, Goentoro et al. 2006). One reason

could be that there is a selective pressure or a strong feedback mechanism that prevents elevated

levels of dopamine in the nervous system. Since oxidized dopamine could be toxic (Zhang et al.

2019), this could be a mechanism to prevent cell toxicity. Another possibility is that disrupting

dopamine levels leads to structural changes that in turn lead to unstable dopamine. In the cuticle

melanin is not the only final product of dopamine; it is also used to make different types of

sclerotins (Sugumaran and Barek 2016). These sclerotins provide additional structural integrity to

the cuticle, so in theory disrupting this structural integrity could perhaps lead to increased

dopamine degradation, though we do not have evidence for this. Regardless, changes in cuticle

pigmentation do not necessarily reflect changes in dopamine level, so assumptions should not be

made that increase in melanin levels in the fly corresponds to increased level of dopamine. Also,

we found that it is generally more common to see a reduction in dopamine than an increase in

both the head and the brain of the fly, indicating there may be a regulatory mechanism to prevent

the increase of dopamine *in vivo*, which warrants further investigation.

Upon seeing alterations of dopamine in the head for some of the classic pigmentation

genes, we next wanted to test if this corresponded to changes in the brain. In the case of

established dopamine metabolism genes, we only saw a change in dopamine levels upon

reduction of TH. None of the enzymes suspected to be involved in dopamine degradation caused

a change in dopamine in the brain as whole-body null or strong loss-of-function mutants. When

we looked at expression for some of these genes, we saw that all the genes we were able to

assess (*black*, *speck*, *tan*) were expressed in the brain, but their expression was limited and did

not seem to overlap with many DANs. It may be that dopamine degradation does not happen in

the neurons since it was hypothesized to occur in the glia (Richardt et al. 2002; Davla et al. 2020).

However, the expression does not seem to show a broad glial expression using the reagents we

had access to. Taken together, this suggests that there is some unknown mechanism for

dopamine degradation in the fly brain. For the 11 genes from the pigmentation RNAi screen that

showed altered levels of dopamine, we assessed dopamine levels in the brain. Only 2 (18%) of

them showed an effect on dopamine in the brain even though several other genes are expressed

in DANs (ranging between 25-100% of DANs examined). This emphasizes what we found when

we observed in the pigmentation mutants that dopamine in the head does not necessarily reflect

a change in the brain. Hence, head dopamine levels should not be used as a proxy for brain

dopamine levels, which is sometimes used in the literature to study Parkinson disease and other

dopamine related disorders using *Drosophila* (Islam et al. 2012; Shimizu et al. 2020; Knight et al.

2014).

***clu* and *mask* reduce brain dopamine levels through different mechanisms**

Once we identified that knockdown of *clu* and *mask* both reduced dopamine in the brain

without affecting the number of DANs, we hypothesized that they may be acting through TH since

our initial dopamine metabolism studies showed that only changes in TH significantly affected

dopamine levels in the brain. While these genes could also act on Ddc or other aspect of

dopamine homeostasis, we decided to focus on TH since this is the rate-limiting enzyme for

dopamine (Levitt et al. 1965) and due to reagent availability.

clu encodes a ribonucleoprotein that is thought to act at the outer mitochondrial membrane

to promote proper formation of different complexes that are crucial to mitochondrial function,

including mitophagy proteins likePink1 and Parkin (Sen et al. 2015; Wang et al. 2016). It also

interacts with the ribosome and translocase complex at the mitochondrial membrane (Sen et al.

2015; Sen and Cox 2016) suggesting it may act as a general regulator for mitochondrial protein

translation and import. In fact, *clu* alleles have been show to exhibit reduced ATP levels, reduced

mitochondrial membrane potential and elevated levels of reactive oxygen species (Sen et al.

2013; Yang et al. 2022). Knockdown of *clu* showed no effect on TH protein level, but since this is

a mitochondrial protein, it may be more likely that the effect on dopamine is through the previously

proposed mechanisms related to unknown roles of mitochondria to regulate dopamine levels. One

study proposed that dopamine synthesis can take place at the mitochondria and was regulated

through phosphorylation of TH (Wang et al. 2009). Hence, it is possible that while total TH protein

level is unaffected, post-translational status or activity of TH may be affected upon *clu* knockdown.

Unfortunately, we were unable to obtain an antibody that recognized phosphorylated *Drosophila*

TH from fly heads to test this hypothesis. Though others have been able to stain for

phosphorylated TH when its activity is at its peak, so it may be feasible to try additional antibodies

using alternative conditions (Davis et al. 2007). Considering that we observed elevated levels of

TH mRNA upon *clu* knockdown, some feedback mechanism may be trying to compensate for

reduced levels of dopamine and potentially TH activity, providing us an entry point to study such

feedback loop which may also involve TH phosphorylation and other signaling mechanisms

(Dunkley et al. 2004; Tekin et al. 2014).

mask encodes a scaffolding molecule that has been studied in a number of different

biological contexts (DeAngelis et al. 2020; Zhu et al. 2017; Martinez et al. 2021; Fisher et al.

2018). In this study, *mask* knockdown showed reduction in both *TH* mRNA and protein levels,

suggesting that it may reduce dopamine by reducing TH transcriptionally. Interestingly, *mask* has

been implicated in multiple biological pathways that were highlighted from our screen, including

EGF signaling (e.g. *Egfr*, *Sev*, Raf), Hippo signaling (e.g. *hpo*, *wts*, *mats*), and mitochondrial

biology (Sidor et al. 2013, 2019; Sansores-Garcia et al. 2013; Smith et al. 2002; Zhu et al. 2015).

While Egf and Hippo signaling have well-known transcriptional effects (Lusk et al. 2017; Oh and

Irvine 2010), whether it may regulate TH transcription directly or indirectly in this context need

further investigations. It is also possible that because *mask* acts in multiple of these pathways,

which causes it to have a strong enough effect to show a significant change in dopamine in the

whole brain analysis while knockdown of individual pathway genes may have a more restricted

role (e.g. *Egfr*, *Sos*, *hpo*). Importantly, while reduction in TH mRNA could be the primary cause of

decreased TH protein levels upon *mask* knockdown, additional translational or post-translational

mechanisms may be involved considering the pleiotropic role of Mask. Interestingly,

haploinsufficiency of the human ortholog of *mask* (*ANKRD17)* was recently identified as the cause

of a complex neurodevelopmental disorder (Chopra et al. 2021). Further investigation of whether

ANKRD17 regulates dopamine levels in the mammalian brain and its mechanistic investigation

will likely provide a novel angle to study this understudied rare disease and related conditions.

In conclusion, while both of these genes caused a reduction in dopamine, they appear to

be acting through different mechanisms. Additional analysis on changes in enzymatic activity as

well as potential downstream targets would help identify exactly how these genes are acting.

Some regulators of dopamine levels may act locally

While we were able to identify two genes that significantly affect dopamine levels in the

brain, there were many genes that did not show a global change in the brain dopamine levels in

the head or brain. Also, due to the limitation in resources, we only measured dopamine in the

head of 35 prioritized candidates. There are still 97 conserved pigmentation genes that may affect

dopamine levels, some of which may function in the brain. Considering changes in head levels of

dopamine were not indicative of changes in brain dopamine, future experiments should be

prioritized based on the candidates' expression in DANs to increase the yield for novel dopamine

regulators.

In addition, when we examined the expression of some of our genes which we measured

dopamine levels from the head or brain (Supplemental Figure 5), we found that many of them do

not overlap with all DANs. Therefore, measuring global changes in dopamine may not be best for

determining effects on DA for many of these genes. They may require targeted dopamine

measurement, perhaps through a fixed or live reporter (Inagaki et al. 2012; Sun et al. 2020). It is

also possible to assess changes in dopamine using behavior, but this would require detailed

assessment for each gene since behaviors are often cluster specific (Kasture et al. 2018). For

example, the gene *rickets* (*rk*) is known to regulate TH activity in the cuticle, but there is no known

role for *rk* in adulthood. However, we found that *rk* was expressed in about ~33% of DANs in the

adult brain. Therefore, it may be difficult to see global changes in dopamine upon *rk* knockdown

considering the effect size and the degree of variability we observe in our biochemical assay, but

since its expression overlaps with all of the DANs in the PPL2ab cluster, we can potentially

measure alterations in behavior that this cluster is known to regulate, such as changes in food

preference (Liu et al. 2017). In conclusion, our screen of identified novel pigmentation genes, a

subset of which were identified as novel regulators of dopamine *in vivo*. Unbiased forward genetic

screens in model organisms are powerful ways to identify unanticipated links between distinct

biological pathways and provide new molecular handles to study the functional connections

between them. Application of such strategies to regulation of dopamine levels will likely continue

to identify novel factors, some of which will have impact on understanding the mechanism of

human neurological and neurodevelopmental diseases.

Materials and Methods

Fly Maintenance

Flies (*Drosophila melanogaster*) strains and crosses were reared at 25°C on a 12:12 Light/Dark (LD) program unless stated otherwise. Flies were fed a molasses-based food source. Crosses were set with 3-5 males and 3-5 virgin females and transferred every two days to increase the number of progenies. Different temperatures were used to achieve different levels of gene knockdown or overexpression since GAL4/UAS system is temperature sensitive. The flies were transferred 0-2 days post-eclosion (dpe) to separate them for age-appropriate analysis. The experiments were generally run on 3-7 dpe flies unless otherwise stated.

Fly Stocks and Crosses

Mutant and transgenic trains were obtained from Bloomington *Drosophila* Stock Center (BDSC, https://bdsc.indiana.edu/), Vienna Drosophila Research Center (VDRC, https://www.viennabiocenter.org/vbcf/vienna-drosophila-resource-center/), the Japanese National Institute of Genetics (https://shigen.nig.ac.jp/fly/nigfly/), or were gifts from scientists in the field or generated in house (Supplemental Table 1-5).

For the cuticle screen *pnr-GAL4* (Calleja et al. 2000) was crossed to the specified UAS-RNAi line (see Supplemental Table 1) at 29°C, 25°C, and/or 18°C and examined after pigmentation is completed (>1 dpe). For the behavior and HPLC analysis *TH-GAL4* females (also called *ple-GAL4*, BDSC stock #8848) were crossed to the specified RNAi line (see Supplemental Table). For all HPLC analysis, the controls were specific to the tested UAS-RNAi line. Thus, the TRiP collection of RNAi lines are compared to the *UAS-mCherry(mCh) RNAi* line (BDSC #35785) and the NIG collection of RNAi lines as well as the VDRC RNAi lines are compared to the *UAS-lacZ RNAi* line (Kennerdell and Carthew 2000). For the HPLC analysis performed on brains, we

recombined the *TH-GAL4* with another DAN driver *GMR58E02-GAL4* (BDSC #41347), which

expresses in a large group DANs in the PAM cluster (Riemensperger et al. 2013). The PAM

cluster of neurons are largely missed in the *TH-GAL4*, but the recombined stock captures the

majority of DANs (see Supplemental Figure 4). These flies are referred to as *TH,R58-GAL4*

throughout the article. The *TH,R58-GAL4* line was also used for qPCR and TH protein analysis

via immunohistochemistry.

For gene expression analysis *UAS-mCh::nls* (BDSC #38424) females were collected and crossed

with respective T2A-GAL4 lines (see Supplemental Table 5). Female and males were both

imaged at ~5-7dpe.

Notum Dissection and Imaging

For thorax analysis and dissection, males and females were both selected, though differences

were not generally observed, and females are shown here. Flies that were 3-5 dpe were collected

and placed in 70% EtOH. Notum dissection and imaging was performed as in (Yamamoto et al.,

2012). In summary, the thorax was dissected by removing the legs, abdomen, and head. Then

the thorax was cut such that a hole was on the ventral side to allow solution to pass into the

thorax. These dissected thoraces were placed in 10% KOH for 10 minutes at 90°C on a heat

block. Then, the KOH was removed and replaced with 70% EtOH solution. The thorax was

trimmed to only contain the notum (dorsal thorax), and any remaining tissue was removed. They

were mounted on a slide prepared with tape on two sides (to prevent the coverslip from crushing

the thorax) in mounting media (50% glycerol, 50% EtOH). The thoraces were imaged on a stereo

microscope (Leica MZ16) using OPTRONICS® MicroFIRE camera. The images are z-stack

brightfield images taken and collapsed using extended depth of field in Image-Pro Plus 7.0 and

In-Focus (Version 1.6).

Brain Expression Analysis

Brain dissections and imaging were performed as reported in (Marcogliese et al. 2022). In

summary, the brains were dissected at 3-7 dpe in ice cold PBS and then fixed in 4% PFA in 0.5%

PBST for 20 minutes. Then, they were washed with a quick wash in 0.5% PBST followed by three

10-20 minute washes in 0.5% PBST while on a rotator. The samples were placed in primary

antibody solution [anti-Tyrosine Hydroxylase 1:500, PelFreez Biologicals, rabbit, P40101 ; anti-

elav 1:100, Developmental Studies Hybridoma Bank, rat, 7E8A10 in solution (5% Normal Donkey

Serum, 0.1% NaN$_3$ in 0.5% PBST)] at 4°C for 3 days. Then, the samples were washed with a

quick wash in 0.5% PBST followed by three 10-20 minute washes in 0.5% PBST while on a

rotator. After the last wash, the samples were placed in a secondary antibody solution [anti-Rabbit

1:200 (Thermo Fisher Sci., Alexa-647, A-21208); anti-rat 1:200 (Thermo Fisher Sci., Alexa-488,

A-27040) in solution (5% Normal Donkey Serum, 0.1% NaN$_3$ in 0.5% PBST)] for two hours at

room temperature. The samples were washed again with one quick wash followed by three 10-

20 minutes washes in 0.5% PBST on a rotator. Then, the brains were mounted in Vectashield

mounting media and imaged on a confocal microscope (Zeiss LSM 710 or Zeiss LSM 880). All

images shown in this manuscript are Z-projection images that were generated using the ZEN

software (Zeiss).

RNAi-based Pigmentation Screen

We identified potential regulators of cuticle color from a primary screen performed in Jurgen

Knoblich's lab using the Vienna Drosophila Resource Center collection of RNAi lines by accessing

their public RNAi screening database (https://bristlescreen.imba.oeaw.ac.at/start.php). The

original screen was performed on 20,262 RNAi lines, encompassing 11,619 genes (~82% of

protein-coding genes) (Mummery-Widmer et al. 2009). We selected genes that showed a cuticle

pigmentation score for any RNAi line tested (selected gene scores ranged from two to ten). To

validate pigmentation defects, we selected genes that had RNAi lines within the Fly National

Institute of Genetics (NIG, 220 genes, 426 RNAi lines) or the Harvard Transgenic RNAi Project

(TRiP, 221 genes, 292 RNAi lines) collections. These RNAi lines were tested for a cuticle color

defects. Each RNAi line was tested at 29°C and 25°C. If there was no phenotype no further testing

was performed. If it was lethal, we tested it at 18°C.

Classification of Cuticle Phenotypes

Phenotypes were scored based on images taken for each genotype. The phenotype was scored

using a qualitative scoring system. This system used *UAS-TH RNAi* and *UAS-ebony RNAi*, some

of the strongest regulators of cuticle color, as a baseline for the "strong". Then, the other lines

were placed along the spectrum as mild, moderate, or strong. Phenotypes were scored by two

independent observers and differences were settled by a third independent observer. If there

was variability with one line or if there was variability amongst lines, this might appear as mild-

moderate, moderate-strong, or mild-strong. In these cases, the strongest phenotype observed

was documented as the recorded phenotype for a given gene.

Protein-protein interaction network and gene ontology analysis

The protein-protein interaction network was generated using STRING (search tool for recurring

instances of neighboring genes; https://string-db.org/). All of the 153 pigmentation gene hits were

input into multiple protein analysis for the species *Drosophila melanogaster*, and genes were

verified for the appropriate gene name. This database includes protein-protein interactions that

are physical interactions and functionally associations. For our data set we included these sources

of interactions: gene neighborhoods (genes that are found close together across species; (Snel

et al. 2000), curated databases (publicly available databases of protein interactions), and

experimental evidence (known complexes and pathways from curated sources) (Szklarczyk et al.

2021, 2023). Any genes that did not interact with other genes from our screen were not included

in the interaction network. This database was last accessed on 05/05/2023. The lines between

genes represents the confidence in the interactions. This is defined based on the strength of data

sources for the given interaction, which is based on complex algorithms that are dependent on

the source.

The gene ontology analysis was generated using GOrilla (Gene Ontology enRIchment anaLysis

and visuaLizAtion tool, https://cbl-gorilla.cs.technion.ac.il/). All of the 153 pigmentation genes hits

were included as target genes, and the background genes were the entire VDRC RNAi collection

included in the Mummery-Widmer et al. 2009 screen. Only the GO terms with a p-value greater

than 0.001 and a fold enrichment of greater than three were included in the analysis. Redundant

terms were removed using Revigo (http://revigo.irb.hr/) with a stringency of 0.7. Those that have

a fold enrichment greater than five were included in Supplementary Figure 1.

Human Neurological Disease Gene Classification

Homologs for each fly gene were identified using DIOPT (DRSC Integrative Ortholog Prediction

Tool, v8.0, https://www.flyrnai.org/diopt) (Hu et al. 2011). Genes were classified as orthologs if

they had a DIOPT score≥3. The human genes were scored as neurological disease-causing

genes if they had an OMIM (Online Mendelian Inheritance of Man, last accessed 01/01/2023,

https://www.omim.org/) disease association with any documented neurological phenotype

(Hamosh et al. 2021). Genes were also classified as human neurological disease genes if they

were included in the Simons Simplex Collection of Autism Spectrum Disorder gene list (last

accessed 06/01/2022, https://gene.sfari.org/) and had a score equal to or less than 3 (1, 2, or 3)

and/or syndromic (S) (Abrahams et al. 2013).

Behavior Analysis using the *Drosophila* Activity Monitor

For the behavior screen, the crosses were set in a 12:12 LD chamber at 25°C and transferred

every 2-3 days to increase the number of progenies. Upon eclosion, flies were transferred to 29°C

and kept for 2-3 more days before testing. Individual tubes (PPT5x65 Polycarbonate, Trikinetics

Inc, USA) appropriate for use with the *Drosophila* Activity Monitor (DAM, Model DAM2 for 5mm

tubes, TriKinetics Inc, USA) were loaded with approximately ½" worth of molasses-based food.

The end of the tube with food was sealed by placing the vial into Paraplast® (Sigma-Aldrich) wax

three times and allowing it to dry. Once they were dry, individual male flies were placed into each

vial, the vial was placed into the DAM recording chamber, and it was sealed using a 5mm tube

cap (CAP5, TriKinetics Inc, USA) with a hole stuffed with cotton. For each genotype, 16 individual

males were run, except in a few cases where we were unable to get that many flies. In which

case, as many living males as possible were run. The flies were loaded into a 12:12 LD chamber

at 29°C and monitored for 3 days. Then, the flies were removed and dead flies were eliminated

from analysis.

Locomotion and sleep analysis was run on a 24-hour period, which started at the first onset of

light after the flies were placed in the chamber (i.e. 18-24 hours after being placed in the chamber).

Locomotion was calculated based on the number of beam crossings over the full 24 hours, the

12 hours of light, and the 12 hours of dark. Sleep was classified as 5 minutes without any beam

crossings. Total Sleep was quantified as total time spent sleeping over the 12 hours of light and

the 12 hours of dark. Sleep latency was quantified as the amount of time after light onset or dark

onset before a 5 minute period of sleep. Sleep Bout Length was quantified as the average length

of sleep bouts (>5 minute period of sleep) during the 12 hours of light and 12 hours of dark.

"Outliers" were classified as lines that were more than two standard deviations from the mean of

all lines for any of the phenotypes scored (24 hour locomotion, 12 hour light locomotion, 12 hour

dark locomotion, 12 hour light total sleep, 12 hour dark total sleep, 12 hour light sleep bout length,

12 hour dark sleep bout length, sleep latency during the light, or sleep latency during the dark).

The total list of genes from all outliers was 50, which was on par with the other classifications. In

addition, there was not a great amount of overlap between the phenotypes scored (Figure 3) and

dopamine could cause an effect in potentially any of these phenotypes depending on what cells

the gene functions in. Thus, a gene was classified as a DAM "outlier" if it was an "outlier" in any

of the scored phenotypes. The data analysis of the DAM data was performed using Microsoft

Excel.

High Performance Liquid Chromatography (HPLC) Analysis of dopamine and serotonin

Sample Preparation

For the HPLC on fly heads and brain, we adapted and modified the protocol based on (Cichewicz

et al. 2017; Hardie and Hirsh 2006). Crosses were reared at 29°C and flies were transferred into

new tubes 0-2 days post eclosion (dpe). They remained at 29°C until they were 3-5 dpe. Female

heads were collected by anesthetizing flies with CO_2 and cutting their heads off with a razor blade.

Samples were collected between ZT04-ZT08. The heads were placed into 60 μL of 50 mM citrate

acetate (pH=4.5) and either used for analysis that day or frozen at -20°C. Five heads were used

per sample, and ~10 samples were run per genotype. The day of HPLC analysis, the samples

were thawed, grounded using a pestle for 30 seconds (Cordless Pestle Motor and Fisherbrand™

Disposable Pellet Pestle for 1.5mL tube). Then the samples were spun down at 13,000 rpm for

10 minutes. The supernatant was removed and placed into a new vial. 10 μL from each sample

was used for the Bradford Protein Analysis Assay. The rest of the sample solution (~40-50 μL)

was loaded into a vial (300 μL Polypropylene Sample Vials with 8mm Snap Caps) for HPLC

analysis.

For HPLC analysis on fly brains, flies were collected 3-7 dpe. Then, 10 fly brains (5 male, 5

female) were dissected in ice cold PBS. Right after dissection, the brains were transferred to 60uL

of ice cold 50mM citrate acetate (pH=4.5). Samples were frozen at -20°C and ran within 4 weeks

of collection. While not all brains were dissected for a given genotype on the same day, the

samples collected for each genotype compared was even for a given day (e.g. 1 control RNAi, 1

RNAi #1, 1 RNAi #2 OR 2 control RNAi, 2 RNAi #1, 2 RNAi #2). The day of HPLC analysis, the

samples were homogenized similarly to the fly heads, though for the Protein Assay, 20 μL of

sample was used. Then, the remaining solution was loaded for HPLC analysis.

Bradford Protein Analysis Assay

The Bio-Rad Bradford Assay was used for colorimetric scoring of total protein. See product details

for full description (Bio-Rad Protein Assay Kit I #5000001). Briefly, the dye reagent was diluted

1:4 in MilliQ H_2O. Then, the solution was filtered through a 0.22 µm SFCA Nalgene filter. A protein

standard was created (500 µg/mL, 250 µg/mL, 100 µg/mL, and 50 µg/mL) by diluting the protein

sample provided with the assay. 10 µL of each protein sample (standard or fly sample) was placed

into single wells of a 96-well plate. Then, 200 µL of diluted 1:4 dye reagent was added to each

well. The samples rested for 30-45 minutes and then absorbance was measured using BMG

Labtech FLUOstar OPTIMA microplate reader. Protein measurements for each sample were

calculated based on standards ran on the same plate.

HPLC Information

The HPLC used is a Antec® Scientific product with a LC110S pump, SYSTEC OEM MINI Vacuum

Degasser, AS110 autosampler, a SenCell flow cell with salt bridge reference electrode in the

Decade Lite. Data was collected and processed using DataApex Clarity™ chromatography

software. The column used is chosen to work well for neurotransmitters (Acquity UPLC BEH C18

Column, 130Å, 1.7 µm, 1 mm X 100 mm with Acquity In-Line 0.2 µm Filter). The mobile phase

was a 6% Acetonitrile mobile phase optimized for our samples (74.4 mg $NA_2EDTA \cdot 2H_2O$, 13.72

mL 85% w/v phosphoric acid, 42.04 g citric acid, 1.2 g OSA, 120 mL acetonitrile, H_2O up to 2L,

pH=6.0 using 50% NaOH solution). The mobile phase was degassed for 10 minutes using the

Bransonic® Ultrasonic Bath before being loaded into the HPLC machine.

The standards for HPLC were made by generating master stocks of 100 mM dopamine and 10

mM serotonin diluted in MilliQ H_2O, which were kept at 4°C. The day of HPLC analysis, the master

stocks were diluted to produce standards ranging from 5-100 nM for 5-HT and 5-1000 nM for

dopamine. Sample concentrations of dopamine (Sigma-Aldrich, Cat#H8502) and serotonin

(Sigma-Aldrich, Cat#H7752) were calculated based on standards run in the same batch.

Standards were compared to standards run on other days to assess for overall performance.

Standards were loaded into the 300 μL Polypropylene Sample Vials with 8mm Snap Caps for

HPLC analysis.

RNA expression analysis

Flies were raised at 29°C until 5-7 dpe, then heads were collected by cutting them off with a razor

blade. Twenty heads were collected per sample with a mixture of males and females and placed

directly on dry ice. Samples were kept on dry ice or at -80°C until RNA isolation. For RNA isolation

samples were homogenized in 200μL of TRIzol®. Then 800μL of TRIzol® was added and the

samples were mixed by pipetting. The samples incubated for 5 minutes at room temperature and

were then spun down at 12,000 rpm for 10 minutes at 4°C. The supernatant was removed and

200μL of chloroform was added. The samples were mixed by shaking and then incubated at room

temperature for 3 minutes. Then, they were spun at 10,000 rpm for 15 minutes at 4°C, and the

aqueous layer was collected and placed in a new tube. 500μL of isopropanol was added and the

samples were left to incubate for 10 minutes at room temperature. The samples were spun at

10,000 rpm for 10 minutes at 4°C and the supernatant was removed. The pellet was washed with

1mL of 75% EtOH and centrifuged at 5,000 rpm for 5 minutes. All EtOH was removed and the

pellet dried at room temperature for 10 minutes. Then, the pellet was dissolved in 100μL of H_2O.

RNA content and quality was measured using DeNovix DS-11 Fx spectrophotometer/fluorometer.

cDNA reverse transcription was performed according to the iScript™ reverse transcriptase Bio-

Rad kit and 2 μL of RNA sample was added in a 10 μL reaction for both the reverse and no

reverse transcriptase reaction. Samples were measured for content and quality and then they

were diluted to 100ng/μL. qPCR was performed using Bio-Rad iQ SYBR Green Supermix and

measured on Bio-Rad CFX96™ Real-Time System. The TH primers used Forward: 5'-

ATGTTCGCCATCAAGAAATCCT-3' and Reverse: 5'-GGGTCTCGAAACGGGCATC-3', and the

control primers were for Rpl32 were Forward: 5'-ATGCTAAGCTGTCGCACAAATG-3' and

Reverse: 5'-GTTCGATCCGTAACCGATGT-3'. For each sample, two technical replicates were

run for the reverse transcriptase reaction, and the final quantity was based on the average of the

two. If the no reverse transcriptase reaction produced a product that was a Cq within ten, the

sample was discarded.

DAN and TH Quantification

Samples were stained and imaged just like to the brain expression analysis, except the primarily

antibody solution included an Elav antibody [anti-Tyrosine Hydroxylase 1:500, PelFreez

Biologicals, rabbit, P40101; anti-Elav 1:100, Developmental Studies Hybridoma Bank, rat,

7E8A10 in solution (5% Normal Donkey Serum, 0.1% NaN_3 in 0.5% PBST)]. Confocal imaging

was performed as described above. Once the samples were imaged, anterior, posterior, and

whole brain projections were generated that included all fluorescent signals from the anterior (PAL

and PAM clusters) and posterior (PPL1, PPM2, PPM3, PPL2ab) neuron clusters of interest. The

number of neurons per cluster (excluding the PAM cluster) were counted for each sample. PPM2

and PPM3 clusters were counted together. For TH protein quantification samples were assessed

using ImageJ. A boundary was drawn around each individual cluster and quantified. Then, the

local background for that cluster was subtracted from the cluster to give the fluorescence for the

given sample. Each hemisphere was treated as a separate sample.

Statistical Analysis

Statistical analysis was performed using Graphpad Prism 10. Unless otherwise stated, all data

was subjected to an ROUT outlier test, where all outliers were removed, and then a one-way

ANOVA was performed. Each of the experimental samples were compared to the control for the

given samples. In the cases where there were only two samples (i.e. control and experimental),

a t-test was performed.

For the HPLC screen on heads, the samples were normalized to the controls run on that day. The

specific controls were specific to the given line such that the TRiP collection was normalized to

UAS-mCh RNAi control and the NIG collection was normalized to the *UAS-lacZ RNAi* control.

Figures and Figure Legends (Figures 1-6, Table 1)

Figure 1

Figure 1: Manipulation of most pigmentation-related enzymes do not have expected effects

on dopamine levels. (A) Pathway for dopamine metabolism and key cuticle components. Gene

names encoding pigmentation-related enzymes are listed in blue font and metabolites in black

font, respectively. (B-D) Thorax phenotypes observed upon knockdown of synthesis enzymes

Tyrosine hydroxylase (TH, also known as Pale) (C), Dopa decarboxylase (Ddc) (D) compared to

control (B). (E-E') Dopamine levels measured from the whole head (E) and whole brain (E') upon

knockdown of *TH* and *Ddc*. Pigmentation phenotypes observed from cuticle pigmentation-related

enzymes in classical mutant alleles, as observed in the thorax (F-L) and the along the body (F'-

L') (Arrow in I' indicates the darker region characteristic of *speck* mutants, which is consistent with

previous observations). *T2A-GAL4* alleles for *black*, *speck*, and *tan* show expression in the adult

brain that does not have major overlap with dopaminergic neurons (DANs) (M-P). (Q) Dopamine

measured from the head of classical pigmentation mutants leads to reduced levels for *black[1]*,

ebony[1], *straw[1]*, and *yellow[1]*. (R) There is no change in dopamine levels measured from dissected

brains of classical pigmentation mutants. Ordinary one-way ANOVA test with Dunlett multiple

comparisons test. $*=p<0.05$, $**=p<0.01$, $***=p<0.001$, $****=p<0.0001$. Error bars show standard

deviation (SD).

Figure 2

Figure 2: RNAi pigmentation screen reveals a broad spectrum of phenotypes and classes

of genes (A) Flow chart depicting screening process for identifying novel regulators of dopamine

in the fly brain. (B) Examples of the breath of phenotypes observed from the RNAi-based

pigmentation screen for flies with 'pale', 'dark', or 'mixed' phenotypes. (C) Protein-protein

interaction network for 153 genes identified as high confidence pigmentation gene hits from RNAi-

screen using the STRING network analysis. Strength of line indicates confidence or strength of

data support. Interactions include gene neighborhoods, curated databases, and experimental

evidence as defined in the methods. Genes that do not have any known interactions with other

genes are not displayed. (D-F) Pie charts showing the representation for the 153 pigmentation

genes hits for strength of phenotype (D), phenotype color (E), and conservation (F).

Table 1

Fly Gene	Human Homologs with Neurological Phenotypes (DIOPT Score, Max=18)	Notable Neurological Phenotypes
AP-1sigma	AP1S2 (12), AP1S1 (11), AP2S1 (3)	seizures, ataxia, developmental delay
ATP7	ATP7A (13), ATP7B (13)	seizures, neurodegeneration, dystonia, tremor
Atu	LEO1 (10)	ASD
byn	TBXT (11), TBX19 (10)	neural tube defects, seizures
CG16758	PNP (14)	spastic diplegia, ataxia, tremor, behavioral problems
CG4328	LMX1B (10)	spina bifida
CG43689	MYT1L (7)	DD, ID
CG4679	PTCD3 (14)	psychomotor delay, myoclonus, cerebral atrophy
CG5902	AMMECR1 (13)	early motor delay, early language delay
CG8728	PMPCA (15)	ataxia
ClC-c	CLCN3 (12), CLCN4 (11), CLCNKA (3)	ID, seizures, structural abnormalities, language delay
ctrip	TRIP12 (10)	psychomotor delay, ID, delayed speech
Cyp301a1	CYP24A1 (5), CYP27B1 (4), CYP27A1 (4)	motor delay, seizures, ataxia, psychiatric symptoms
CysRS-m	CARS2 (15)	psychomotor delay, epilepsy, cerebral atrophy
Ddc	DDC (14), HDC (4)	psychomotor delay, dystonia, sleep disturbance
Dgp-1	GTPBP2 (3)	psychomotor delay, ID, ataxia, dystonia, seizures
Dscam2	DSCAM (10)	ASD
Egfr	ERBB4 (10), ERBB2 (10), ERBB3 (9)	ALS, DD, axonal neuropathy, seizures
Gbeta13F	GNB1 (15), GNB4 (13), GNB2 (12), GNB5 (3)	ID, CMT, DD, language delay, ADHD
gig	TSC2 (13)	seizures, brain structural abnormalities
Hrb98DE	HNRNPA2B1 (12), HNRNPA1 (9)	dementia, motor neuropathy, ALS
Itpr	ITPR1 (14), ITPR3 (11)	motor delay, ataxia, cerebellar atrophy
Khc	KIF5C (12), KIF5A (12)	psychomotor delay, seizures, spastic paraplegia, ALS
Larp7	LARP7 (10)	ID, DD, unstable gait, anxiety
LeuRS-m	LARS2 (15)	cognitive impairment, DD, dystonia, tremor, seizures
mask	ANKRD17 (10)	developmental delay, ID, speech delay, motor delay
mbt	PAK1 (3), PAK3 (3), PAK2 (3)	ID, macrocephaly, seizures, speech delay
Membrin	GOSR2 (13)	DD, cerebral atrophy, seizures
Mp	COL18A1 (6)	ataxia, seizures
Mpcp2	SLC25A3 (14)	motor delay
ND-13B	NDUFA5 (13)	ASD

ND-42	NDUFA10 (14)	DD, structural abnormalities
Npc2b	NPC2 (7)	DD, dementia, dystonia, seizures, behavioral problems
per	PER3 (8), PER2 (7)	sleep abnormalities
PIG-A	PIGA (15)	seizures, structural abnormalities
ple	TH (15)	ataxia, dystonia, Parkinsonism, motor and speech delay
Prosap	SHANK2 (10), SHANK3 (8)	ASD, schizophrenia, DD, seizures
Raf	BRAF (14), RAF1 (11)	DD, seizures
Rbp9	ELAVL2 (12), ELAVL3 (10)	ASD
rdx	SPOP (10)	ID, DD, speech delay
robo1	ROBO3 (8)	structural abnormalities
SCOT	OXCT1 (15)	coma, psychomotor delay
sesB	SLC25A4 (12)	motor delay, hyperreflexia
slp1	FOXG1 (6), FOXI1 (3), FOXJ1 (3), FOXC1 (3)	dystonia, motor delay, sleep and structural abnormalities
Smurf	ITCH (4), NEDD4L (3)	ID, psychomotor delay, seizures
Sod3	SOD1 (4)	ALS, spastic tetraplegia
Sos	SOS1 (14), SOS2 (12)	cognitive impair, learning disabilities
Spn38F	SERPINC1 (4), SERPINI1 (4)	seizures, dementia, cerebral atrophy
stv	BAG3 (9)	abnormal gait, neuropathy, behavioral problems
tws	PPP2R2A (13)	ataxia
Usp2	USP8 (4)	behavioral problems
Usp7	USP7 (13)	psychomotor delay, ID, seizures, structural abnormalities

Table 1: Fly pigmentation genes identified from this screen with neurological disease

associated homologs in human. Human homologs were defined as fly genes with a Drosophila

Integrative Orthology Prediction Tool (DIOPT, version 8.0) score greater than or equal to three

for at least one human gene. Neurological disease associations were based on Online Mendelian

Inheritance in Man (OMIM)_clincal disease phenotypes and high-confident ASD genes listed in

Simons Foundation Autism Research Initiative (SFARI) gene database (class S, 1, 2 or 3).

ASD=Autism Spectrum Disorder, DD=Developmental Delay, ID=Intellectual Disability,

ALS=Amytrophic Lateral Sclerosis, CMT=Charcot-Marie-Tooth disease.

Figure 3

Figure 3: *Drosophila* Activity Monitor (DAM)-based screen identifies 60 lines

corresponding to 45 genes as behavioral "outliers". Activity and sleep data showing

representative quantification for the phenotypes measured in a neutral RNAi line (*mCherry RNAi*,

A) and an example of a behavioral "outlier" (*Raf RNAi*, B) knocked down in many dopaminergic

cells (*TH-GAL4*). (C) Bar graph showing range of all RNAi lines tested for 24hr locomotion

phenotype, where light pink represents lines ±1SD away from the mean, and dark pink are lines

±2SDs away from the mean (a.k.a. "outliers"). (D) Venn diagram for all "outliers" identified for the

four phenotypes examined (sleep latency, sleep bout length, locomotion, and total sleep). (E-H)

Bar graphs showing range of RNAi lines tested for locomotion (E), sleep bout length (F), total

sleep (G), and sleep latency (H).

Figure 4

Figure 4: Eleven genes with altered levels of head dopamine upon knockdown. (A) Venn

diagram showing three categories that were used to prioritze 39 genes (in yellow) tested for

altered levels of dopamine. (B) Representative chromatograms showing peaks for dopamine and

serotonin for upon knockdown or overexpression of TH in many dopaminergic cells (*TH-GAL4*).

(C) Quantification of dopamine (DA) and serotonin (5-HT) using HPLC upon knockdown (*TH>TH*

RNAi, blue) or overexpression of (*TH>TH cDNA*, green) of TH. (D-E) HPLC analysis on prioritized

gene hits for lines from the NIG (D) or the TRiP (E) collection of UAS-RNAi lines. Pie charts for

the prioritized genes hits (39 genes) showing the representation of cuticle color (F) and the trend

of change in dopamine (G). (H) XY plot showing pigmentation phenotype compared to dopamine

levels observed. Brown-Forsythe test showed different SDs for (C), so Brown-Forsythe ANOVA

test performed with Dunnett's T3 multiple comparisons test. For (D) and (E) ordinary One-way

ANOVA test with Dunnett's multiple comparisons test. **=$p < 0.01$, ***=$p < 0.001$, ****=$p < 0.0001$.

Error bars represent SD.

Figure 5

Figure 5: Analysis of 11 genes identifies *mask* and *clu* as novel regulators of dopamine

levels in the fly brain (A-D) Bsg^{T2A}-GAL4, ClC-c^{T2A}-GAL4, PIG-A^{T2A}-GAL4, and rk^{T2A}-GAL4

driving *UAS-mCh::nls* (nuclear localized mCherry) shows broad expression patterns in the brain.

(A'-D') These expression patterns overlaps with some to many dopaminergic neurons. Dopamine

measured from dissected adult brains upon knockdown of *Bsg* (E), *CG43689* (F), *ClC-c* (G), *Dpy-*

30L1 (H), *Membrin* (I), *PIG-A* (J), *rk* (K), *Spn38F* (L), *Sos* (M) do not show consistent changes in

dopamine using a novel recombined doaminergic GAL4 driver (*TH, R58-GAL4*). (N) Knockdown

of *clu* shows consistent reduction in total dopamine in the adult fly brain. (O) *clu::GFP* shows

broad expression in the brain and may be enriched in many DANs (P-R, P-R'). (S) Knockdown of

mask shows consistent reduction in total dopamine in the adult fly brain. (T) *mask::GFP* shows

broad expression in the brain and overlaps with many DANs (U-W, U'-W'). For samples with one

control to one RNAi, t-tests were performed (E, F, H, I, J, L). For samples with two samples for

one control, ordinary one-way ANOVA was done with Dunnett's multiple comparisons test (G, K,

M, R, W). Error bars represent SD. Scale bars are 50 μm.

Figure 6

Figure 6: *mask* and *clu* effects on Tyrosine Hydroxylase

Figure 6: *mask* and *clu* knockdown do not affect TH cell number and having different

affects TH levels (A) qRT-PCR of TH upon *mask* and *clu* knockdown shows reduced TH mRNA

upon *mask* knockdown and increased TH mRNA upon *clu* knockdown using the recombined

dopaminergic (*TH, R58-GAL4*) driver. (B-B') Image showing the primary clusters of DANs in the

anterior (B) and posterior (B') of the adult fly brain. (C) Quantification of total TH protein level

through immunofluorescence signals in given clusters of DANs upon knockdown of *mask* and *clu*.

(D) Quantification of the number of TH positive (TH+) neurons in given clusters of DANs upon

knockdown of *mask* and *clu*. Ordinary one-way ANOVAs performed with Dunnett's multiple

comparisons for those with significant differences. $*=p<0.05$, $**p<0.01$, $****p<0.0001$. Error bars

represent SD.

Acknowledgements

Acknowledgements

This work was supported by startup funds to SY from the Jan and Dan Duncan Neurological

Research Institute at Texas Children's Hospital and the Department of Molecular and Human

Genetics at Baylor College of Medicine. Confocal microscopy at BCM was supported in part by

the Intellectual and Developmental Disabilities Research Center (IDDRC, grant number

U54HD083092) from the Eunice Kennedy Shriver National Institute of Child Health & Human

Development. We thank the Bloomington *Drosophila* Stock Center (USA), the National Institutes

Genetics Fly Stock Center (Japan), *Vienna* Drosophila RNAi Center (Austria) and Dr. Hugo Bellen

for providing useful fly stocks and reagents for this project. We would like to thank Drs. Jonathan

Andrews, Hugo Bellen, Brigitte Dauwalder, Herman Dierick, Linsday Goodman, Oguz Kanca,

Kartik Venkatachalam, Michael Wangler, Sheng Zhang and Huda Zoghbi for useful suggestions,

discussions and advice on the overall project.

References

Abrahams, B.S., Arking, D.E., Campbell, D.B., Mefford, H.C., Morrow, E.M., Weiss, L.A., et al. (2013). SFARI Gene 2.0: A Community-Driven Knowledgebase for the Autism Spectrum Disorders (ASDs). Mol. Autism 4 (1). doi: 10.1186/2040-2392-4-36.

Armstrong, M.J., and Okun, M.S. (2020). Diagnosis and Treatment of Parkinson Disease: A Review. JAMA 323 (6): 548–60. doi: 10.1001/JAMA.2019.22360.

Bayersdorfer, F., Voigt, A., Schneuwly, S., and Botella, J.A. (2010). Dopamine-Dependent Neurodegeneration in Drosophila Models of Familial and Sporadic Parkinson's Disease. Neurobiol. Dis. 40 (1): 113–19. doi: 10.1016/J.NBD.2010.02.012.

Birman, S., Morgan, B., Anzivino, M., and Hirsh, J. (1994). A Novel and Major Isoform of Tyrosine Hydroxylase in Drosophila Is Generated by Alternative RNA Processing. J. Biol. Chem. 269 (42): 26559–67. doi: 10.1016/S0021-9258(18)47231-6.

Brand, A.H., and Perrimon, N. (1993). Targeted Gene Expression as a Means of Altering Cell Fates and Generating Dominant Phenotypes. Development 118 (2): 401–15. http://www.ncbi.nlm.nih.gov/pubmed/8223268.

Budnik, V., and White, K. (1987). Genetic Dissection of Dopamine and Serotonin Synthesis in the Nervous System of Drosophila Melanogaster. J. Neurogenet. 4 (6): 309–14. doi: 10.3109/01677068709167191.

Burbulla, L.F., Song, P., Mazzulli, J.R., Zampese, E., Wong, Y.C., Jeon, S., et al. (2017). Dopamine Oxidation Mediates Mitochondrial and Lysosomal Dysfunction in Parkinson's Disease. Science 357 (6357): 1255–61. doi: 10.1126/SCIENCE.AAM9080.

Buszczak, M., Paterno, S., Lighthouse, D., Bachman, J., Planck, J., Owen, S., et al. (2007). The Carnegie Protein Trap Library: A Versatile Tool for Drosophila Developmental Studies. Genetics 175 (3): 1505–31. doi: 10.1534/GENETICS.106.065961.

Calleja, M., Herranz, H., Estella, C., Casal, J., Lawrence, P., Simpson, P., et al. (2000).

Generation of Medial and Lateral Dorsal Body Domains by the Pannier Gene of

Drosophila. Development 127 (18): 3971–80. doi: 10.1242/DEV.127.18.3971.

Chaudhuri, A., Bowling, K., Funderburk, C., Lawal, H., Inamdar, A., Wang, Z., et al. (2007).

Interaction of Genetic and Environmental Factors in a Drosophila Parkinsonism Model. J.

Neurosci. 27 (10): 2457. doi: 10.1523/JNEUROSCI.4239-06.2007.

Chiu, J.C., Low, K.H., Pike, D.H., Yildirim, E., and Edery, I. (2010). Assaying Locomotor Activity

to Study Circadian Rhythms and Sleep Parameters in Drosophila. J. Vis. Exp., no. 43. doi:

10.3791/2157.

Chopra, M., McEntagart, M., Clayton-Smith, J., Platzer, K., Shukla, A., Girisha, K.M., et al.

(2021). Heterozygous ANKRD17 Loss-of-Function Variants Cause a Syndrome with

Intellectual Disability, Speech Delay, and Dysmorphism. Am. J. Hum. Genet. 108 (6):

1138–50. doi: 10.1016/J.AJHG.2021.04.007.

Church, F.C. (2021). Treatment Options for Motor and Non-Motor Symptoms of Parkinson's

Disease. Biomolecules 11 (4). doi: 10.3390/BIOM11040612.

Cichewicz, K., Garren, E.J., Adiele, C., Aso, Y., Wang, Z., Wu, M., et al. (2017). A New Brain

Dopamine-Deficient Drosophila and Its Pharmacological and Genetic Rescue. Genes.

Brain. Behav. 16 (3): 394–403. doi: 10.1111/GBB.12353.

Clark, I.E., Dodson, M.W., Jiang, C., Cao, J.H., Huh, J.R., Seol, J.H., et al. (2006). Drosophila

Pink1 Is Required for Mitochondrial Function and Interacts Genetically with Parkin. Nature

441 (7097): 1162–66. doi: 10.1038/nature04779.

Davis, M.M., O'Keefe, S.L., Primrose, D.A., and Hodgetts, R.B. (2007). A Neuropeptide

Hormone Cascade Controls the Precise Onset of Post-Eclosion Cuticular Tanning in

Drosophila Melanogaster. Development 134 (24): 4395–4404. doi: 10.1242/DEV.009902.

Davla, S., Artiushin, G., Li, Y., Chitsaz, D., Li, S., Sehgal, A., et al. (2020). AANAT1 Functions in

Astrocytes to Regulate Sleep Homeostasis. Elife 9 (September): 1–48. doi:

10.7554/ELIFE.53994.

DeAngelis, M.W., McGhie, E.W., Coolon, J.D., and Johnson, R.I. (2020). Mask, a Component of

the Hippo Pathway, Is Required for Drosophila Eye Morphogenesis. Dev. Biol. 464 (1): 53–

70. doi: 10.1016/J.YDBIO.2020.05.002.

Diao, F., Ironfield, H., Luan, H., Diao, F., Shropshire, W.C., Ewer, J., et al. (2015). Plug-and-

Play Genetic Access to Drosophila Cell Types Using Exchangeable Exon Cassettes. Cell

Rep. 10 (8): 1410–21. doi: 10.1016/j.celrep.2015.01.059.

Dietzl, G., Chen, D., Schnorrer, F., Su, K.C., Barinova, Y., Fellner, M., et al. (2007). A Genome-

Wide Transgenic RNAi Library for Conditional Gene Inactivation in Drosophila. Nature 448

(7150): 151–56. doi: 10.1038/NATURE05954.

Dunkley, P.R., Bobrovskaya, L., Graham, M.E., Nagy-Felsobuki, E.I. Von, and Dickson, P.W.

(2004). Tyrosine Hydroxylase Phosphorylation: Regulation and Consequences. J.

Neurochem. 91 (5): 1025–43. doi: 10.1111/J.1471-4159.2004.02797.X.

Fisher, K.H., Fragiadaki, M., Pugazhendhi, D., Bausek, N., Arredondo, M.A., Thomas, S.J., et

al. (2018). A Genome-Wide RNAi Screen Identifies MASK as a Positive Regulator of

Cytokine Receptor Stability. J. Cell Sci. 131 (13). doi: 10.1242/JCS.209551.

Freeman, A., Pranski, E., Miller, R.D., Radmard, S., Bernhard, D., Jinnah, H.A., et al. (2012).

Sleep Fragmentation and Motor Restlessness in a Drosophila Model of Restless Legs

Syndrome. Curr. Biol. 22 (12): 1142. doi: 10.1016/J.CUB.2012.04.027.

Frighetto, G., Zordan, M.A., Castiello, U., Megighian, A., and Martin, J.R. (2022). Dopamine

Modulation of Drosophila Ellipsoid Body Neurons, a Nod to the Mammalian Basal Ganglia.

Front. Physiol. 13 (April). doi: 10.3389/FPHYS.2022.849142.

Gervasi, N., Tchénio, P., and Preat, T. (2010). PKA Dynamics in a Drosophila Learning Center:

Coincidence Detection by Rutabaga Adenylyl Cyclase and Spatial Regulation by Dunce

Phosphodiesterase. Neuron 65 (4): 516–29. doi: 10.1016/J.NEURON.2010.01.014.

Gervasi, N.M., Scott, S.S., Aschrafi, A., Gale, J., Vohra, S.N., Macgibeny, M.A., et al. (2016).

The Local Expression and Trafficking of Tyrosine Hydroxylase MRNA in the Axons of

Sympathetic Neurons. RNA 22 (6): 883–95. doi: 10.1261/RNA.053272.115.

Goentoro, L.A., Yakoby, N., Goodhouse, J., Schüpbach, T., and Shvartsman, S.Y. (2006).

Quantitative Analysis of the GAL4/UAS System in Drosophila Oogenesis. Genesis 44 (2):

66–74. doi: 10.1002/GENE.20184.

Grace, A.A. (2016). Dysregulation of the Dopamine System in the Pathophysiology of

Schizophrenia and Depression. Nat. Rev. Neurosci. 17 (8): 524–32. doi:

10.1038/NRN.2016.57.

Hamosh, A., Amberger, J.S., Bocchini, C., Scott, A.F., and Rasmussen, S.A. (2021). Online

Mendelian Inheritance in Man (OMIM®): Victor McKusick's Magnum Opus. Am. J. Med.

Genet. A 185 (11): 3259–65. doi: 10.1002/AJMG.A.62407.

Hardie, S.L., and Hirsh, J. (2006). An Improved Method for the Separation and Detection of

Biogenic Amines in Adult Drosophila Brain Extracts by High Performance Liquid

Chromatography. J. Neurosci. Methods 153 (2): 243–49. doi:

10.1016/J.JNEUMETH.2005.11.001.

Heisenberg, M., Borst, A., Wagner, S., and Byers, D. (1985). Drosophila Mushroom Body

Mutants Are Deficient in Olfactory Learning. J. Neurogenet. 2 (1): 1–30. doi:

10.3109/01677068509100140.

Hovemann, B.T., Ryseck, R.P., Walldorf, U., Störtkuhl, K.F., Dietzel, I.D., and Dessen, E.

(1998). The Drosophila Ebony Gene Is Closely Related to Microbial Peptide Synthetases

and Shows Specific Cuticle and Nervous System Expression. Gene 221 (1): 1–9. doi:

10.1016/S0378-1119(98)00440-5.

Hu, Y., Flockhart, I., Vinayagam, A., Bergwitz, C., Berger, B., Perrimon, N., et al. (2011). An

Integrative Approach to Ortholog Prediction for Disease-Focused and Other Functional

Studies. BMC Bioinformatics 12 (August). doi: 10.1186/1471-2105-12-357.

Imperato, A., Puglisi-Allegra, S., Casolini, P., Zocchi, A., and Angelucci, L. (1989). Stress-

Induced Enhancement of Dopamine and Acetylcholine Release in Limbic Structures: Role

of Corticosterone. Eur. J. Pharmacol. 165 (2–3): 337–38. doi: 10.1016/0014-

2999(89)90735-8.

Inagaki, H.K., Ben-Tabou De-Leon, S., Wong, A.M., Jagadish, S., Ishimoto, H., Barnea, G., et

al. (2012). Visualizing Neuromodulation In Vivo: TANGO-Mapping of Dopamine Signaling

Reveals Appetite Control of Sugar Sensing. Cell 148 (3): 583. doi:

10.1016/J.CELL.2011.12.022.

Islam, R., Yang, L., Sah, M., Kannan, K., Anamani, D., Vijayan, C., et al. (2012). A

Neuroprotective Role of the Human Uncoupling Protein 2 (HUCP2) in a Drosophila

Parkinson's Disease Model. Neurobiol. Dis. 46 (1): 137–46. doi:

10.1016/J.NBD.2011.12.055.

Janssens, J., Aibar, S., Taskiran, I.I., Ismail, J.N., Gomez, A.E., Aughey, G., et al. (2022).

Decoding Gene Regulation in the Fly Brain. Nature 601 (7894): 630–36. doi:

10.1038/S41586-021-04262-Z.

Jenett, A., Rubin, G.M., Ngo, T.T.B., Shepherd, D., Murphy, C., Dionne, H., et al. (2012). A

GAL4-Driver Line Resource for Drosophila Neurobiology. Cell Rep. 2 (4): 991–1001. doi:

10.1016/J.CELREP.2012.09.011.

Kanca, O., Zirin, J., Garcia-Marques, J., Knight, S.M., Yang-Zhou, D., Amador, G., et al. (2019).

An Efficient CRISPR-Based Strategy to Insert Small and Large Fragments of DNA Using

Short Homology Arms. Elife 8 (November). doi: 10.7554/ELIFE.51539.

Kasture, A.S., Hummel, T., Sucic, S., and Freissmuth, M. (2018). Big Lessons from Tiny Flies:

Drosophila Melanogaster as a Model to Explore Dysfunction of Dopaminergic and

Serotonergic Neurotransmitter Systems. Int. J. Mol. Sci. 19 (6). doi:

10.3390/IJMS19061788.

Kawahata, I., Ohtaku, S., Tomioka, Y., Ichinose, H., and Yamakuni, T. (2015). Dopamine or

Biopterin Deficiency Potentiates Phosphorylation at (40)Ser and Ubiquitination of Tyrosine

Hydroxylase to Be Degraded by the Ubiquitin Proteasome System. Biochem. Biophys.

Res. Commun. 465 (1): 53–58. doi: 10.1016/J.BBRC.2015.07.125.

Kennerdell, J.R., and Carthew, R.W. (2000). Heritable Gene Silencing in Drosophila Using

Double-Stranded RNA. Nat. Biotechnol. 18 (8): 896–98. doi: 10.1038/78531.

Kitada, T., Asakawa, S., Hattori, N., Matsumine, H., Yamamura, Y., Minoshima, S., et al. (1998).

Mutations in the Parkin Gene Cause Autosomal Recessive Juvenile Parkinsonism. Nature

392 (6676): 605–8. doi: 10.1038/33416.

Knight, A.L., Yan, X., Hamamichi, S., Ajjuri, R.R., Mazzulli, J.R., Zhang, M.W., et al. (2014). The

Glycolytic Enzyme, GPI, Is a Functionally Conserved Modifier of Dopaminergic

Neurodegeneration in Parkinson's Models. Cell Metab. 20 (1): 145–57. doi:

10.1016/J.CMET.2014.04.017.

Kroemer, G., Mariño, G., and Levine, B. (2010). Autophagy and the Integrated Stress

Response. Mol. Cell 40 (2): 280. doi: 10.1016/J.MOLCEL.2010.09.023.

Lee, P.-T., Zirin, J., Kanca, O., Lin, W.-W., Schulze, K.L., Li-Kroeger, D., et al. (2018). A Gene-

Specific T2A-GAL4 Library for Drosophila. Elife 7 (March). doi: 10.7554/eLife.35574.

Letizia, A., Bario, R., and Campuzano, S. (2007). Antagonistic and Cooperative Actions of the

EGFR and Dpp Pathways on the Iroquois Genes Regulate Drosophila Mesothorax

Specification and Patterning. Development 134 (7): 1337–46. doi: 10.1242/DEV.02823.

Levitt, M., Spector, S., Sjoerdsma, A., and Udenfriend, S. (1965). ELUCIDATION OF THE

RATE-LIMITING STEP IN NOREPINEPHRINE BIOSYNTHESIS IN THE PERFUSED

GUINEA-PIG HEART. J. Pharmacol. Exp. Ther. 148 (1).

Li, H., Janssens, J., Waegeneer, M. de, Kolluru, S.S., Davie, K., Gardeux, V., et al. (2022). Fly

Cell Atlas: A Single-Nucleus Transcriptomic Atlas of the Adult Fruit Fly. Science (80-.). 375

(6584). doi:

10.1126/SCIENCE.ABK2432/SUPPL_FILE/SCIENCE.ABK2432_MDAR_REPRODUCIBILI

TY_CHECKLIST.PDF.

Lim, K.L., and Tan, J.M.M. (2007). Role of the Ubiquitin Proteasome System in Parkinson's

Disease. BMC Biochem. 8 (Suppl 1): S13. doi: 10.1186/1471-2091-8-S1-S13.

Liu, Q., Tabuchi, M., Liu, S., Kodama, L., Horiuchi, W., Daniels, J., et al. (2017). Branch-Specific Plasticity of a Bifunctional Dopamine Circuit Encodes Protein. Science 356 (6337): 534. doi: 10.1126/SCIENCE.AAL3245.

Livingstone, M.S., and Tempel, B.L. (1983). Genetic Dissection of Monoamine Neurotransmitter Synthesis in Drosophila. Nature 303 (5912): 67–70. doi: 10.1038/303067A0.

López-Cruz, L., San Miguel, N., Carratalá-Ros, C., Monferrer, L., Salamone, J.D., and Correa, M. (2018). Dopamine Depletion Shifts Behavior from Activity Based Reinforcers to More Sedentary Ones and Adenosine Receptor Antagonism Reverses That Shift: Relation to Ventral Striatum DARPP32 Phosphorylation Patterns. Neuropharmacology 138 (August): 349–59. doi: 10.1016/J.NEUROPHARM.2018.01.034.

Lu, J., Wang, Y., Wang, X., Wang, D., Pflugfelder, G.O., and Shen, J. (2022). The Tbx6 Transcription Factor Dorsocross Mediates Dpp Signaling to Regulate Drosophila Thorax Closure. Int. J. Mol. Sci. 23 (9). doi: 10.3390/IJMS23094543.

Lusk, J.B., Lam, V.Y.M., and Tolwinski, N.S. (2017). Epidermal Growth Factor Pathway Signaling in Drosophila Embryogenesis: Tools for Understanding Cancer. Cancers (Basel). 9 (2). doi: 10.3390/CANCERS9020016.

Mao, Z., and Davis, R.L. (2009). Eight Different Types of Dopaminergic Neurons Innervate the Drosophila Mushroom Body Neuropil: Anatomical and Physiological Heterogeneity. Front. Neural Circuits 3 (JUL): 5. doi: 10.3389/NEURO.04.005.2009/BIBTEX.

Marcogliese, P.C., Deal, S.L., Andrews, J., Harnish, J.M., Bhavana, V.H., Graves, H.K., et al. (2022). Drosophila Functional Screening of de Novo Variants in Autism Uncovers Damaging Variants and Facilitates Discovery of Rare Neurodevelopmental Diseases. Cell Rep. 38 (11). doi: 10.1016/J.CELREP.2022.110517.

Martinez, D., Zhu, M., Guidry, J.J., Majeste, N., Mao, H., Yanofsky, S.T., et al. (2021). Mask, the Drosophila Ankyrin Repeat and KH Domain-Containing Protein, Affects Microtubule

Stability. J. Cell Sci. 134 (20). doi: 10.1242/JCS.258512.

Massey, J.H., Akiyama, N., Bien, T., Dreisewerd, K., Wittkopp, P.J., Yew, J.Y., et al. (2019).

Pleiotropic Effects of Ebony and Tan on Pigmentation and Cuticular Hydrocarbon

Composition in Drosophila Melanogaster. Front. Physiol. 10 (MAY): 518. doi:

10.3389/FPHYS.2019.00518/BIBTEX.

McGary, K.L., Park, T.J., Woods, J.O., Cha, H.J., Wallingford, J.B., and Marcotte, E.M. (2010).

Systematic Discovery of Nonobvious Human Disease Models through Orthologous

Phenotypes. Proc. Natl. Acad. Sci. U. S. A. 107 (14): 6544–49. doi:

10.1073/pnas.0910200107.

McKinnon, C., Snoo, M.L. De, Gondard, E., Neudorfer, C., Chau, H., Ngana, S.G., et al. (2020).

Early-Onset Impairment of the Ubiquitin-Proteasome System in Dopaminergic Neurons

Caused by α-Synuclein. Acta Neuropathol. Commun. 8 (1). doi: 10.1186/S40478-020-

0894-0.

Meiser, J., Weindl, D., and Hiller, K. (2013). Complexity of Dopamine Metabolism. Cell

Commun. Signal. 11 (1). doi: 10.1186/1478-811X-11-34.

Morin, X., Daneman, R., Zavortink, M., and Chia, W. (2001). A Protein Trap Strategy to Detect

GFP-Tagged Proteins Expressed from Their Endogenous Loci in Drosophila. Proc. Natl.

Acad. Sci. U. S. A. 98 (26): 15050. doi: 10.1073/PNAS.261408198.

Mummery-Widmer, J.L., Yamazaki, M., Stoeger, T., Novatchkova, M., Bhalerao, S., Chen, D., et

al. (2009). Genome-Wide Analysis of Notch Signalling in Drosophila by Transgenic RNAi.

Nature 458 (7241): 987–92. doi: 10.1038/nature07936.

Nakashima, A., Ohnuma, S., Kodani, Y., Kaneko, Y.S., Nagasaki, H., Nagatsu, T., et al. (2016).

Inhibition of Deubiquitinating Activity of USP14 Decreases Tyrosine Hydroxylase

Phosphorylated at Ser19 in PC12D Cells. Biochem. Biophys. Res. Commun. 472 (4): 598–

602. doi: 10.1016/J.BBRC.2016.03.022.

Ni, J.Q., Liu, L.P., Binari, R., Hardy, R., Shim, H.S., Cavallaro, A., et al. (2009). A Drosophila

Resource of Transgenic RNAi Lines for Neurogenetics. Genetics 182 (4): 1089–1100. doi:

10.1534/GENETICS.109.103630.

Ni, J.Q., Markstein, M., Binari, R., Pfeiffer, B., Liu, L.P., Villalta, C., et al. (2007). Vector and

Parameters for Targeted Transgenic RNA Interference in Drosophila Melanogaster. Nat.

Methods 2008 51 5 (1): 49–51. doi: 10.1038/nmeth1146.

Ni, J.Q., Zhou, R., Czech, B., Liu, L.P., Holderbaum, L., Yang-Zhou, D., et al. (2011). A

Genome-Scale ShRNA Resource for Transgenic RNAi in Drosophila. Nat. Methods 8 (5):

405–7. doi: 10.1038/NMETH.1592.

Nicklas, W. m. do. J., Vyas, I., and Heikkila, R.E. (1985). Inhibition of NADH-Linked Oxidation in

Brain Mitochondria by 1-Methyl-4-Phenyl-Pyridine, a Metabolite of the Neurotoxin, 1-

Methyl-4-Phenyl-1,2,5,6-Tetrahydropyridine. Life Sci. 36 (26): 2503–8. doi: 10.1016/0024-

3205(85)90146-8.

Nutt, D.J., Lingford-Hughes, A., Erritzoe, D., and Stokes, P.R.A. (2015). The Dopamine Theory

of Addiction: 40 Years of Highs and Lows. Nat. Rev. Neurosci. 16 (5): 305–12. doi:

10.1038/NRN3939.

Oh, H., and Irvine, K.D. (2010). Yorkie: The Final Destination of Hippo Signaling. Trends Cell

Biol. 20 (7): 410. doi: 10.1016/J.TCB.2010.04.005.

Pan, Y., Li, W., Deng, Z., Sun, Y., Ma, X., Liang, R., et al. (2022). Myc Suppresses Male-Male

Courtship in Drosophila. EMBO J. 41 (7). doi: 10.15252/EMBJ.2021109905.

Pan, Y., and Monje, M. (2020). Activity Shapes Neural Circuit Form and Function: A Historical

Perspective. J. Neurosci. 40 (5): 944. doi: 10.1523/JNEUROSCI.0740-19.2019.

Pendleton, R.G., Rasheed, A., Sardina, T., Tully, T., and Hillman, R. (2002). Effects of Tyrosine

Hydroxylase Mutants on Locomotor Activity in Drosophila: A Study in Functional Genomics.

Behav. Genet. 32 (2): 89–94. doi: 10.1023/A:1015279221600.

Radwan, B., Liu, H., and Chaudhury, D. (2019). The Role of Dopamine in Mood Disorders and

the Associated Changes in Circadian Rhythms and Sleep-Wake Cycle. Brain Res. 1713

(June): 42–51. doi: 10.1016/J.BRAINRES.2018.11.031.

Rezatabar, S., Karimian, A., Rameshknia, V., Parsian, H., Majidinia, M., Kopi, T.A., et al. (2019).

1149   RAS/MAPK Signaling Functions in Oxidative Stress, DNA Damage Response and Cancer

1150   Progression. J. Cell. Physiol. 234 (9): 14951–65. doi: 10.1002/JCP.28334.

1151   Richardt, A., Rybak, J., Störtkuhl, K.F., Meinertzhagen, I.A., and Hovemann, B.T. (2002). Ebony

1152   Protein in the Drosophila Nervous System: Optic Neuropile Expression in Glial Cells. J.

1153   Comp. Neurol. 452 (1): 93–102. doi: 10.1002/CNE.10360.

1154   Riemensperger, T., Isabel, G., Coulom, H., Neuser, K., Seugnet, L., Kume, K., et al. (2011).

1155   Behavioral Consequences of Dopamine Deficiency in the Drosophila Central Nervous

1156   System. Proc. Natl. Acad. Sci. U. S. A. 108 (2): 834–39. doi: 10.1073/PNAS.1010930108.

1157   Riemensperger, T., Issa, A.R., Pech, U., Coulom, H., Nguyễn, M.V., Cassar, M., et al. (2013). A

1158   Single Dopamine Pathway Underlies Progressive Locomotor Deficits in a Drosophila Model

1159   of Parkinson Disease. Cell Rep. 5 (4): 952–60. doi: 10.1016/J.CELREP.2013.10.032.

1160   Rohwedder, A., Wenz, N.L., Stehle, B., Huser, A., Yamagata, N., Zlatic, M., et al. (2016). Four

1161   Individually Identified Paired Dopamine Neurons Signal Reward in Larval Drosophila. Curr.

1162   Biol. 26 (5): 661–69. doi: 10.1016/J.CUB.2016.01.012.

1163   Sansores-Garcia, L., Atkins, M., Moya, I.M., Shahmoradgoli, M., Tao, C., Mills, G.B., et al.

1164   (2013). Mask Is Required for the Activity of the Hippo Pathway Effector Yki/YAP. Curr. Biol.

1165   23 (3): 229–35. doi: 10.1016/J.CUB.2012.12.033.

1166   Sekine, Y., Hatanaka, R., Watanabe, T., Sono, N., Iemura, S. ichiro, Natsume, T., et al. (2012).

1167   The Kelch Repeat Protein KLHDC10 Regulates Oxidative Stress-Induced ASK1 Activation

1168   by Suppressing PP5. Mol. Cell 48 (5): 692–704. doi: 10.1016/J.MOLCEL.2012.09.018.

1169   Sekine, Y., Takagahara, S., Hatanaka, R., Watanabe, T., Oguchi, H., Noguchi, T., et al. (2011).

1170   P38 MAPKs Regulate the Expression of Genes in the Dopamine Synthesis Pathway

1171   through Phosphorylation of NR4A Nuclear Receptors. J. Cell Sci. 124 (Pt 17): 3006–16.

1172   doi: 10.1242/JCS.085902.

Sen, A., and Cox, R.T. (2016). Clueless Is a Conserved Ribonucleoprotein That Binds the

Ribosome at the Mitochondrial Outer Membrane. Biol. Open 5 (2): 195–203. doi:

10.1242/BIO.015313/-/DC1.

Sen, A., Damm, V.T., and Cox, R.T. (2013). Drosophila Clueless Is Highly Expressed in Larval

Neuroblasts, Affects Mitochondrial Localization and Suppresses Mitochondrial Oxidative

Damage. PLoS One 8 (1). doi: 10.1371/JOURNAL.PONE.0054283.

Sen, A., Kalvakuri, S., Bodmer, R., and Cox, R.T. (2015). Clueless, a Protein Required for

Mitochondrial Function, Interacts with the PINK1-Parkin Complex in Drosophila. Dis.

Model. Mech. 8 (6): 577–89. doi: 10.1242/DMM.019208.

Shafer, O.T., and Keene, A.C. (2021). The Regulation of Drosophila Sleep. Curr. Biol. 31 (1):

R38–49. doi: 10.1016/J.CUB.2020.10.082.

Shimizu, J., Kasai, T., Yoshida, H., Huynh, A.M., Nakao-Azuma, Y., Shinomoto, M., et al.

(2020). Novel Drosophila Model for Parkinsonism by Targeting Phosphoglycerate Kinase.

Neurochem. Int. 139 (October). doi: 10.1016/J.NEUINT.2020.104816.

Sidor, C., Borreguero-Munoz, N., Fletcher, G.C., Elbediwy, A., Guillermin, O., and Thompson,

B.J. (2019). Mask Family Proteins ANKHD1 and ANKRD17 Regulate YAP Nuclear Import

and Stability. Elife 8 (October). doi: 10.7554/ELIFE.48601.

Sidor, C.M., Brain, R., and Thompson, B.J. (2013). Mask Proteins Are Cofactors of Yorkie/YAP

in the Hippo Pathway. Curr. Biol. 23 (3): 223–28. doi: 10.1016/J.CUB.2012.11.061.

Smith, R.K., Carroll, P.M., Allard, J.D., and Simon, M.A. (2002). MASK, a Large Ankyrin Repeat

and KH Domain-Containing Protein Involved in Drosophila Receptor Tyrosine Kinase

Signaling. Development 129 (1): 71–82. doi: 10.1242/DEV.129.1.71.

Snel, B., Lehmann, G., Bork, P., and Huynen, M.A. (2000). STRING: A Web-Server to Retrieve

and Display the Repeatedly Occurring Neighbourhood of a Gene. Nucleic Acids Res. 28

(18): 3442. doi: 10.1093/NAR/28.18.3442.

Spana, E.P., Abrams, A.B., Ellis, K.T., Klein, J.C., Ruderman, B.T., Shi, A.H., et al. (2020).

Speck, First Identified in Drosophila Melanogaster in 1910, Is Encoded by the

Arylalkalamine N-Acetyltransferase (AANAT1) Gene. G3 (Bethesda). 10 (9): 3387–98. doi:

10.1534/G3.120.401470.

Spanos, M., Gras-Najjar, J., Letchworth, J.M., Sanford, A.L., Toups, J.V., and Sombers, L.A.

(2013). Quantitation of Hydrogen Peroxide Fluctuations and Their Modulation of Dopamine

Dynamics in the Rat Dorsal Striatum Using Fast-Scan Cyclic Voltammetry. ACS Chem.

Neurosci. 4 (5): 782–89. doi: 10.1021/CN4000499.

Steiner, H., and Kitai, S.T. (2001). Unilateral Striatal Dopamine Depletion: Time-Dependent

Effects on Cortical Function and Behavioural Correlates. Eur. J. Neurosci. 14 (8): 1390–

1404. doi: 10.1046/J.0953-816X.2001.01756.X.

Sugumaran, M., and Barek, H. (2016). Critical Analysis of the Melanogenic Pathway in Insects

and Higher Animals. Int. J. Mol. Sci. 17 (10). doi: 10.3390/IJMS17101753.

Sun, F., Zhou, J., Dai, B., Qian, T., Zeng, J., Li, X., et al. (2020). Next-Generation GRAB

Sensors for Monitoring Dopaminergic Activity in Vivo. Nat. Methods 17 (11): 1156. doi:

10.1038/S41592-020-00981-9.

Szklarczyk, D., Gable, A.L., Nastou, K.C., Lyon, D., Kirsch, R., Pyysalo, S., et al. (2021). The

STRING Database in 2021: Customizable Protein–Protein Networks, and Functional

Characterization of User-Uploaded Gene/Measurement Sets. Nucleic Acids Res. 49 (D1):

D605. doi: 10.1093/NAR/GKAA1074.

Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R., et al. (2023). The

STRING Database in 2023: Protein–Protein Association Networks and Functional

Enrichment Analyses for Any Sequenced Genome of Interest. Nucleic Acids Res. 51 (D1):

D638. doi: 10.1093/NAR/GKAC1000.

Takahashi, A. (2013). Pigmentation and Behavior: Potential Association through Pleiotropic

Genes in Drosophila. Genes Genet. Syst. 88 (3): 165–74. doi: 10.1266/GGS.88.165.

Takáts, S., Pircs, K., Nagy, P., Varga, Á., Kárpáti, M., Hegedus, K., et al. (2014). Interaction of

the HOPS Complex with Syntaxin 17 Mediates Autophagosome Clearance in Drosophila.

Mol. Biol. Cell 25 (8): 1338–54. doi: 10.1091/MBC.E13-08-0449.

Tank, A.W., Xu, L., Chen, X., Radcliffe, P., and Sterling, C.R. (2008). Post-Transcriptional

Regulation of Tyrosine Hydroxylase Expression in Adrenal Medulla and Brain. Ann. N. Y.

Acad. Sci. 1148: 238–48. doi: 10.1196/ANNALS.1410.054.

Tekin, I., Roskoski, R., Carkaci-Salli, N., and Vrana, K.E. (2014). Complex Molecular Regulation

of Tyrosine Hydroxylase. J. Neural Transm. 121 (12): 1451–81. doi: 10.1007/S00702-014-

1238-7.

Terhzaz, S., Cabrero, P., Chintapalli, V.R., Davies, S.A., and Dow, J.A.T. (2010). Mislocalization

of Mitochondria and Compromised Renal Function and Oxidative Stress Resistance in

Drosophila SesB Mutants. Physiol. Genomics 41 (1): 33–41. doi:

10.1152/PHYSIOLGENOMICS.00147.2009.

True, J.R., Edwards, K.A., Yamamoto, D., and Carroll, S.B. (1999). Drosophila Wing Melanin

Patterns Form by Vein-Dependent Elaboration of Enzymatic Prepatterns. Curr. Biol. 9 (23):

1382–91. doi: 10.1016/S0960-9822(00)80083-4.

Valente, E.M., Abou-Sleiman, P.M., Caputo, V., Muqit, M.M.K., Harvey, K., Gispert, S., et al.

(2004). Hereditary Early-Onset Parkinson's Disease Caused by Mutations in PINK1.

Science (80-.). 304 (5674): 1158–60. doi: 10.1126/science.1096284.

Vié, A., Cigna, M., Toci, R., and Birman, S. (1999). Differential Regulation of Drosophila

Tyrosine Hydroxylase Isoforms by Dopamine Binding and CAMP-Dependent

Phosphorylation. J. Biol. Chem. 274 (24): 16788–95. doi: 10.1074/JBC.274.24.16788.

Vissers, J.H.A., Manning, S.A., Kulkarni, A., and Harvey, K.F. (2016). A Drosophila RNAi Library

Modulates Hippo Pathway-Dependent Tissue Growth. Nat. Commun. 2016 71 7 (1): 1–6.

doi: 10.1038/ncomms10368.

Wang, J., Lou, H., Pedersen, C.J., Smith, A.D., and Perez, R.G. (2009). 14-3-3zeta Contributes

to Tyrosine Hydroxylase Activity in MN9D Cells: Localization of Dopamine Regulatory

Proteins to Mitochondria. J. Biol. Chem. 284 (21): 14011–19. doi:
10.1074/JBC.M901310200.

Wang, Z., Ferdousy, F., Lawal, H., Huang, Z., Daigle, J.G., Izevbaye, I., et al. (2011).
Catecholamines up Integrates Dopamine Synthesis and Synaptic Trafficking. J.
Neurochem. 119 (6): 1294–1305. doi: 10.1111/J.1471-4159.2011.07517.X.

Wang, Z.H., Clark, C., and Geisbrecht, E.R. (2016). Drosophila Clueless Is Involved in Parkin-
Dependent Mitophagy by Promoting VCP-Mediated Marf Degradation. Hum. Mol. Genet.
25 (10): 1946–64. doi: 10.1093/HMG/DDW067.

Wicker-Thomas, C., and Hamann, M. (2008). Interaction of Dopamine, Female Pheromones,
Locomotion and Sex Behavior in Drosophila Melanogaster. J. Insect Physiol. 54 (10–11):
1423–31. doi: 10.1016/J.JINSPHYS.2008.08.005.

William Langston, J., Ballard, P., Tetrud, J.W., and Irwin, I. (1983). Chronic Parkinsonism in
Humans Due to a Product of Meperidine-Analog Synthesis. Science 219 (4587): 979–80.
doi: 10.1126/SCIENCE.6823561.

Wittkopp, P.J., True, J.R., and Carroll, S.B. (2002). Reciprocal Functions of the Drosophila
Yellow and Ebony Proteins in the Development and Evolution of Pigment Patterns.
Development 129 (8): 1849–58. doi: 10.1242/DEV.129.8.1849.

Wright, T.R.F., Bewley, G.C., and Sherald, A.F. (1976). The Genetics of Dopa Decarboxylase in
Drosophila Melanogaster. II. Isolation and Characterization of Dopa-Decarboxylase-
Deficient Mutants and Their Relationship to the Alpha-Methyl-Dopa-Hypersensitive
Mutants. Genetics 84 (2): 287–310. doi: 10.1093/GENETICS/84.2.287.

Xie, T., Ho, M.C.W., Liu, Q., Horiuchi, W., Lin, C.C., Task, D., et al. (2018). A Genetic Toolkit for
Dissecting Dopamine Circuit Function in Drosophila. Cell Rep. 23 (2): 652–65. doi:
10.1016/J.CELREP.2018.03.068.

Xu, C., Luo, J., He, L., Montell, C., and Perrimon, N. (2017). Oxidative Stress Induces Stem Cell
Proliferation via TRPA1/RyR-Mediated Ca2+ Signaling in the Drosophila Midgut. Elife 6

(May). doi: 10.7554/ELIFE.22441.

Yamamoto-Hino, M., Yoshida, H., Ichimiya, T., Sakamura, S., Maeda, M., Kimura, Y., et al.

(2015). Phenotype-Based Clustering of Glycosylation-Related Genes by RNAi-Mediated

Gene Silencing. Genes Cells 20 (6): 521–42. doi: 10.1111/GTC.12246.

Yamamoto, S., Charng, W.-L., Rana, N.A., Kakuda, S., Jaiswal, M., Bayat, V., et al. (2012). A

Mutation in EGF Repeat-8 of Notch Discriminates Between Serrate/Jagged and Delta

Family Ligands. Science (80-.). 338 (6111): 1229–32. doi: 10.1126/science.1228745.

Yamamoto, S., Jaiswal, M., Charng, W.-L., Gambin, T., Karaca, E., Mirzaa, G., et al. (2014). A

Drosophila Genetic Resource of Mutants to Study Mechanisms Underlying Human Genetic

Diseases. Cell 159 (1): 200–214. doi: 10.1016/j.cell.2014.09.002.

Yamamoto, S., Kanca, O., Wangler, M.F., and Bellen, H.J. (2023). Integrating Non-Mammalian

Model Organisms in the Diagnosis of Rare Genetic Diseases in Humans. Nat. Rev. Genet.

In press.

Yamamoto, S., and Seto, E.S. (2014). Dopamine Dynamics and Signaling in Drosophila: An

Overview of Genes, Drugs and Behavioral Paradigms. Exp. Anim. 63 (2): 107–19. doi:

10.1538/EXPANIM.63.107.

Yang, H., Sibilla, C., Liu, R., Yun, J., Hay, B.A., Blackstone, C., et al. (2022). Clueless/CLUH

Regulates Mitochondrial Fission by Promoting Recruitment of Drp1 to Mitochondria. Nat.

Commun. 13 (1). doi: 10.1038/S41467-022-29071-4.

Yoon, S., and Baik, J.H. (2013). Dopamine D2 Receptor-Mediated Epidermal Growth Factor

Receptor Transactivation through a Disintegrin and Metalloprotease Regulates

Dopaminergic Neuron Development via Extracellular Signal-Related Kinase Activation. J.

Biol. Chem. 288 (40): 28435–46. doi: 10.1074/JBC.M113.461202.

Zeitlinger, J., and Bohmann, D. (1999). Thorax Closure in Drosophila: Involvement of Fos and

the JNK Pathway. Development 126 (17): 3947–56. doi: 10.1242/DEV.126.17.3947.

Zhang, S., Wang, R., and Wang, G. (2019). Impact of Dopamine Oxidation on Dopaminergic

Neurodegeneration. ACS Chem. Neurosci. 10 (2): 945–53. doi:

10.1021/ACSCHEMNEURO.8B00454/ASSET/IMAGES/MEDIUM/CN-2018-

004545_0005.GIF.

Zhang, Y.Q., Friedman, D.B., Wang, Z., Woodruff, E., Pan, L., O'Donnell, J., et al. (2005).

Protein Expression Profiling of the Drosophila Fragile X Mutant Brain Reveals Up-

Regulation of Monoamine Synthesis. Mol. Cell. Proteomics 4 (3): 278–90. doi:

10.1074/mcp.M400174-MCP200.

Zhu, M., Li, X., Tian, X., and Wu, C. (2015). Mask Loss-of-Function Rescues Mitochondrial

Impairment and Muscle Degeneration of Drosophila Pink1 and Parkin Mutants. Hum. Mol.

Genet. 24 (11): 3272–85. doi: 10.1093/HMG/DDV081.

Zhu, M., Zhang, S., Tian, X., and Wu, C. (2017). Mask Mitigates MAPT- and FUS-Induced

Degeneration by Enhancing Autophagy through Lysosomal Acidification. Autophagy 13

(11): 1924–38. doi: 10.1080/15548627.2017.1362524.

1316