Synthetic Eco-Evolutionary Dynamics in Simple Molecular Environment

Luca Casiraghi[a†], **Francesco Mambretti**[b†], **Anna Tovo**[b], **Elvezia Maria Paraboschi**[c,d], **Samir Suweis**[b], **Tommaso Bellini**[a]

[a]Dipartimento di Biotecnologie Mediche e Medicina Traslazionale, Università degli Studi di Milano, Via Fratelli Cervi, 93 - L.I.T.A., Segrate, 20054, Italy; [b]Dipartimento di Fisica e Astronomia, Università degli Studi di Padova, Via Marzolo 8, Padova, 35131, Italy; [c]Department of Biomedical Sciences, Humanitas University, Via Rita Levi Montalcini 4, Pieve Emanuele, 20072, Italy; [d]IRCCS, Humanitas Clinical and Research Center, Via Manzoni 56, Rozzano, 20089, Italy

*For correspondence:
elvezia_maria.paraboschi@
hunimed.eu (EMP);
samir.suweis@unipd.it (SS);
tommaso.bellini@unimi.it (TB)

[†]These authors contributed equally to this work.

Abstract   The understanding of eco-evolutionary dynamics, and in particular the mechanism of emergence of species, is still fragmentary and in need of test bench model systems. To this aim we developed a variant of SELEX in-vitro selection to study the evolution of a population of $\sim 10^{15}$ single-strand DNA oligonucleotide 'individuals'. We begin with a seed of random sequences which we select via affinity capture from $\sim 10^{12}$ DNA oligomers of fixed sequence ('resources') over which they compete. At each cycle ('generation'), the ecosystem is replenished via PCR amplification of survivors. Massive parallel sequencing indicates that across generations the variety of sequences ('species') drastically decreases, while some of them become populous and dominate the ecosystem. The simplicity of our approach, in which survival is granted by hybridization, enables a quantitative investigation of fitness through a statistical analysis of binding energies. We find that the strength of individual-resource binding dominates the selection in the first generations, while inter and intra-individual interactions becomes important in later stages, in parallel with the emergence of prototypical forms of mutualism and parasitism.

Introduction

A central effort in theoretical biology and ecology is to provide an effective description of the intimate but often subtle relationship between a given environment and the evolution of its ecosystem. In the case of simple environments without geographical isolation and physical barriers, as the one here considered, the emergence of species and phenotyping clustering *Davis* (*1943*); *Rundle and Nosil* (*2005*); *Keymer et al.* (*2012*); *Gupta et al.* (*2021*) is generally considered an outcome of the competition for the limited available resources *Dieckmann and Doebeli* (*1999*); *Pigolotti et al.* (*2007*); *de Aguiar et al.* (*2009*); *Anceschi et al.* (*2018*). The coexistence of stable species is ecologically understood in terms of "niches", indicating the unique role and position that a particular species occupies within an ecosystem. According to the "niche hypothesis" *Chase and Leibold* (*2009*); *Peterson* (*2011*); *Anceschi et al.* (*2018*), biodiversity is limited by the number of "niches" (or types of resources) that are present since no two species can occupy the same niche indefinitely, as one would eventually out-compete the other *Levin* (*1970*); *Gupta et al.* (*2021*).

Fitness, which quantifies species reproductive success and thus also their relationship with the environment *De Visser and Krug* (*2014*), cannot generally be defined in a predictive way even in the most idealized systems *Thurner et al.* (*2010*); *Wiser and Lenski* (*2015*). Indeed, despite the efforts

to identify simple case study systems, the evolution of populations formed by a variety of species remains difficult to model and to quantitatively characterize because of the inherent complexity of ecosystems and living beings, of the large number of potentially relevant variables of difficult access, and of the role of stochasticity *Catalán et al.* (*2017*).

In this scenario, introducing new tools to explore and test eco-evolutionary models, concepts and interpretations appear as the best strategy toward new understanding *Solé* (*2016*). Along this line, various synthetic biological platforms have been proposed in the last years *Ichihashi et al.* (*2013*); *Tizei et al.* (*2016*); *Parrilla-Gutierrez et al.* (*2017*); *Kauffman et al.* (*2018*); *Adamala and Szostak* (*2013*); *Katla et al.* (*2023*) that exploit different principles and mechanisms, and focused on in-vivo, in-vitro, ex-vivo, or in-silico approaches. For example, in *Ichihashi et al.* (*2013*) a "cell-like" model system is proposed, in which the evolution of one long genomic RNA (> 2,000 nt-long) was investigated in detail under the action of a selective pressure ultimately provided by its biological meaning. Despite the tremendous simplification provided by this approach, the constructed artificial cell is still a complex system that includes ribosomes, lipids, translation factors, accessory proteins, tRNAs and amino acids. While most of the synthetic biological platforms are based on the establishment of cell-like, compartmentalized systems involving complex biomolecular milieus and processes *Tizei et al.* (*2016*), a few are devoted to investigate, with distinct strategies, the competition between two coexisting vescicle-based species competing for resources (vescicle-forming molecules) in a non-biochemical context *Adamala and Szostak* (*2013*); *Katla et al.* (*2023*). These articles demonstrate that typical evolution concepts, such as competition or resources and niche exclusion principle, can be applied to "non-biological" systems.

We here propose a synthetic eco-evolutionary scheme, with no cell-like compartmentalization and no connection to the molecular biology of the cell: no coding sequences, no translation, no proteins involved. We consider an ecosystem formed by a large crowd ($\sim 10^{15}$) of distinct molecular individuals interacting with each other and competing for survival in an environment with fixed resources, and we focus on the process by which selection and competition drive the emergence of dominating species. Specifically, we study solutions of 50-base-long single strand DNA oligomers with random sequences, thus initially all distinct, whose mechanisms of survival and mutual interactions are based on DNA hybridization. By exploiting the wealth of tools and knowledge around DNA code selective pairing and DNA synthesis, amplification and sequencing, we create a condition in which a limited number of accessible and computable variables controls the destiny of an ecosystem formed by biological molecules that evolve in a non-biological way, i.e. with no reference to the biological meaning of their sequence.

The results of our work, demonstrating the intimate connection between fitness and ecological interactions, belong to the growing body of studies *Fussmann et al.* (*2007*); *Vetsigian* (*2017*); *Camacho Mateu et al.* (*2021*) showing that ecological and evolutionary processes are strictly connected in the emergence and maintenance of species.

Results

Affinity-based DNA Synthetic Evolution

We introduce here a variant of SELEX for in-vitro synthetic evolution of oligonucleotides to develop protein-binding aptamers *Ellington and Szostak* (*1990*); *Tuerk and Gold* (*1990*). In standard SELEX protocol, the evolving oligonucleotides are selected at each cycle by their interactions of with the target protein. In the experiment here reported, we implemented a selective mechanism based on the affinity capture provided by magnetic beads carrying single-stranded DNA (ssDNA) filaments of fixed length $L = 20$ and sequence, that act as targets (or resources, Figure 1a). Selection is thus primarily based on the sequence of the DNA individuals and its level of complementarity to the targets. This marks a significant difference with SELEX, in which the aptamer-protein interaction depends instead on higher order factors such as the secondary structure of the oligonucleotides and the variety of binding sites on the folded protein. Being these hard to model, predict and

Figure 1. Affinity-based DNA Synthetic Evolution (ADSE). a: structure of the DNA oligomers participating in ADSE as individuals (DNAi) and resources (Figure 1-figure supplement 1). b: steps in the ADSE (Figure 1-figure supplement 2). The process starts with a random-sequence DNAi population. The capture by magnetic bead-conjugated resources provides the selection: bead-bound DNAi are amplified to form the new generation, a small fraction of which is sequenced by MPS. The rest of the original solution is discarded. Red arrows mark the steps of each ADSE cycle. c: possible interactions involving DNAi.
The online version of this article includes the following figure supplement(s) for Figure 1:
Figure 1-figure supplement 1 DNA individuals structure and its interaction with the oligos.
Figure 1-figure supplement 2 Schematic representation of the Affinity-based DNA Synthetic Evolution protocol.

---

control, SELEX has never been considered, to the best of our knowledge, as a useful experimental
test bench to understand evolution.
In our Affinity-based DNA Synthetic Evolution (ADSE) protocol, evolution start from an initial
pool of DNA Individuals (DNAi), chosen to be of fixed length $L = 50$ and random-sequence. Since
the potential molecular variety is $4^{50} \sim 10^{30}$ while our experiments use about 10^{15} initial molecules,
each sequence is represented one time only. The following evolution process is sketched in Fig-
ure 1b, Figure 1-figure supplement 2, and is given by three steps.
(i) *Selection*: the seed population is mixed with a given amount of dispersed capture beads. After
a suitable incubation time, the beads are extracted from the solution and the bound oligomers
released and saved. The rest of the original solution is discarded;
(ii) *Amplification*: the pool of "survived" oligomers is PCR amplified about 1000 times to recover the
initial molarity;
(iii) *Sequencing*: a small portion of the amplified sample (1-3 10^6 molecules) is analyzed by massive
parallel sequencing.
These steps constitute a cycle - one generation of evolution - that we repeated up to 24 times in
two independent evolution histories.
In actuality, to enable amplification and sequencing, DNAi are built by flanking the 50mer with
two 25 base long fixed sequences that enable primer binding for a total length of 100 bases.
Such two terminal segments of the DNAi are made inactive during selection by hybridization with
oligomers of perfect complementarity, as sketched in Figure 1a, Figure 1-figure supplement 1.
A key feature of ADSE is that DNAi can interact not only with the resources, but also with itself
and with each other, and also potentially form complexes binding to the affinity beads as summa-
rized in Figure 1c. Indeed, the choice of a length of 50 for the DNAi interaction was thought to
enable - in principle - simultaneous resource and mutual binding.
PCR amplification can be of high fidelity or error-prone, the latter choice enabling genetic drift
and, potentially, speciation. Since our primary goal was to first investigate fitness in a pool of com-
peting species within a given niche - defined in our case by the 20 base long resources, we opted
for a high fidelity amplification, and left the investigation of high mutations regimes to a follow-up
experiment. Because of the many PCR cycles required in the ADSE scheme, the amplification in-
evitably leads to the formation of artifacts in the form of longer sequences *Tolle et al.* (*2014*), a
phenomenon that intrinsically sets a limit to the number of generations that can be explored (see
Appendix 1).

123 As detailed below, the ADSE protocol enables observing a non-trivial evolution of the DNAi
124 ecosystem across generations, the emergence of dominating species and nonmonotonic popula-
125 tion evolution. It also enables to appreciate the role of inter-specific interactions and their contri-
126 bution to fitness.

Evolution of the DNAi ecosystem

128 The main output of the synthetic evolution that we are proposing is the dataset $\{\text{DNAi}\}_j$ obtained
129 by sequencing DNAi at the various cycles ($1 \leq j \leq 24$ being the index of the cycle). We find that
130 the initial random ecosystem markedly changes with ADSE generations as shown in Figure 2a, in
131 which we have plotted the evolution of: (i) the fraction F_D of the total population formed by dis-
132 tinct nucleotide sequences (red dots). F_D drops from 1 to nearly 0, indicating that initially DNAi
133 are all different from each other, while after 24 generations the number of distinct sequences is
134 much smaller than the number of DNAi. This indicates that most of the initial sequences become
135 progressively "extinct"; (ii) The fraction F_{10} of the total population formed by the 10 most abundant
136 DNA sequences (black dots). F_{10} is initially close to 0 - being each sequence represented by a single
137 DNAi, and grows to about 25% at cycle 12 and 55% at cycle 24 - indicating that the offspring of 10
138 initial "mother" sequences becomes the majority of the system. This loss of diversity can also be
139 quantified by the zipped file size of the list of sequences *Benedetto et al.* (*2002*) or by computing
140 its Shannon entropy *Hill* (*1973*), both markedly decreasing during evolution (see Appendix 1).

141 Since in ADSE survival depends on bead capture, we expect the DNAi-resource binding strength
142 to be an essential ingredient of fitness. To this aim we computed the mean DNAi-resources bind-
143 ing free energy ΔG_{DR} across generations, plotted in Figure 2b (blue squares, right-hand side y-axis).
144 Specifically, ΔG_{DR} has been computed based on DNAi and resource sequences by using the stan-
145 dard "nearest-neighbor approximation" for the thermodynamics of DNA hybridization *SantaLu-*
146 *cia Jr* (*1998*); *Ghosh et al.* (*2019*); *Plata et al.* (*2021*), as implemented in the NUPACK tool in Python
147 *Zadeh et al.* (*2011*). Being this computation of some complexity, we could perform it only on
148 batches of 1000 DNAi randomly chosen within $\{DNA_i\}_j$, the error bars expressing the uncertainty
149 introduced by such downsampling (see Material and Methods section).

150 To perform a faster - and thus complete - analysis of the resource-binding energy in the evolv-
151 ing population, we introduced a simpler, albeit more approximate, quantifier: the length ω of the
152 longest consecutive number of basis within each DNAi that is complementary to the resource se-
153 quence (considering all possible relative positions of the two), in which we allow for single pairing
154 errors and 1-base bulges *Mambretti et al.* (*2022a*). While ω, missing key ingredients such as the
155 fraction of CG vs. AT base pairs and their sequence, is certainly inadequate to evaluate the binding
156 strength of a specific DNAi, we find that its average value $\langle \omega \rangle$ computed for the whole popula-
157 tion in each generation (Figure 2b, full red dots, left-hand side y-axis) grows almost identically to
158 $\langle \Delta G_{DR} \rangle$. The $\langle \omega \rangle$ axis has been scaled so to make its value computed on a pool of ideal randomly
159 sequence DNAi (dashed red line) match, on the plot, $\langle \Delta G_{DR} \rangle$ computed on the same population
160 (dashed blue line). A more detailed comparison between ω and ΔG_{DR} is in Figure 2c, showing box
161 plots with ΔG_{DR} on the y axis and ω expressed through color code, built on a random selection of
162 1000 distinct DNAi in the initial (left hand side box) and the final populations (central box), and on
163 top 10 most frequent sequences in the final generation (right hand side box). The result further
164 strengthens the validity, in our statistical context, of ω as a quantifier of interaction strength to the
165 resources *Mambretti et al.* (*2022a*).

166 Figure 2b reveals that, during ADSE, $\langle \omega \rangle$ tends to saturate at a value $\omega_{sat} \approx 10$, indicating that no
167 relevant selective advantage is gained when $\omega > 10$, in agreement with the notion that the residence
168 time of hybridized oligomers becomes larger than typical experimental times when $\omega > 12$ *Di Leo*
169 *et al.* (*2022*).

170 Figure 2d-f describes the evolution of the ecosystem $\{\text{DNAi}_j$ by showing $P(\omega)$, the fraction of
171 DNAi having overlap ω with the resource evaluated in the initial pool (panel d) and at generation 12
172 (e) and 24 (f). $P(\omega)$ clearly evolves, its small ω components being progressively lost, while individuals

with large ω grow in number, as they are more successful in being selected and amplified. It might be worth pointing out that the appearance of non-zero $p(\omega > 13)$ at generation 12, while $p(\omega > 13) = 0$ in the initial distribution, is a pure effect of the undersampling involved in the sequencing procedure.

The different destiny of DNAi with distinct ω values is shown in Figure 2g-i, where we show the temporal (i.e. across generation) evolution of the fraction of the total population whose ω is in the following ranges: $3 \leq \omega \leq 5$ (panel g), $6 \leq \omega \leq 10$ (panel h) and $11 \leq \omega$ (panel i). As expected, DNAi with weak affinity to the resources decrease while those with large affinity increase. Interestingly, DNAi with intermediate affinity exhibit a non-monotonic behavior, indicating that the conditions for survival change during the evolution, reflecting the evolution of the ecosystem.

Null Model and Eco-evolutionary Algorithm

In order to better understand the experimental outcomes, we first build a null random model without evolution and then an Individual-Based Eco-Evolutionary (IBEE) model enabling predictions for $P(\omega)$.

The null model describes the interaction between individuals in the initial pool $\{DNAi_{j=0}\}$ of random sequences and the resources within a purely combinatorial framework (see Methods and Appendix 3 for mathematical details). In this model, we attach a random string of length 50 to the resource string in a random position, and we compute the maximum consecutive overlap, accepting the binding only if it is at least formed by 3 complementary basis. For analytical tractability we do not allow the random string to shift over the resource, but we calculate the ω constrained to a specific binding position.

The resulting analytical $p_0(\omega)$, plotted in Figure 2d (orange crosses) closely matches the data obtained from sequencing, confirming the random-sequence nature of $\{DNAi_{j=0}\}$. The analytical $P_0(\omega)$ extends to a ω range where no data are available because of the limited size of the sequenced pool. The comparison between $P_0(\omega)$ and $P(\omega)$ in panels e and f shows that the exponential decay of $P_0(\omega)$ at large ω is maintained even in generation 12 and partly in generation 24, further supporting the notion that the selective advantage of ω saturates at large ω.

In the IBEE algorithm we consider N_p individuals. Each has a fitness $f_i(\omega)$ ($i = 1, 2, .., N_p$) that depends on its affinity ω with the resource. At time $t = 0$ we assign ω to each individual based on $P_0(\omega)$ from the null model. Then, for each evolutive cycle we model competition and selection so that out of N_p individuals, only N_r survives. This is attained as a combination of two processes: a fraction x of the N_r sequences results from sampling individuals from the N_p population, each with a survival probability $f(\omega) \in (0, 1)$; the remaining fraction $1 - x$ is formed by a straight random selection from the N_p individuals. The latter group is meant to mimic "neutral drift" (as it would be expressed in evolutionary language) provided by non-specific binding to the beads. The N_r survivers are amplified by identical copying back to N_p, the starting population of the next evolutionary cycle (the introduction of very rare mutations do not affect the results). We perform 24 evolutionary cycles.

As expected, the outcome of the eco-evolutionary dynamics strongly depends on the shape of the fitness function. We model it as:

$$f(\omega) = \left(\frac{\omega^*}{\omega_{max}} \right)^{\gamma}$$

where ω_{max} is the (cycle-dependent) largest value of ω within the actual population and, $\omega^* = \omega$ if $\omega < \omega_{sat}$, while $\omega^* = \omega_{sat}$ otherwise. ω_{sat} and γ are parameters to be tuned in the comparison with the observed $\langle \omega \rangle$. ω_{sat} expresses the loss of fitness gain for $\omega > \omega_{sat}$ yielding the saturation of $\langle \omega \rangle$. γ represents the strength with which $f(\omega)$ depends on ω, and thus the rate at which low ω individuals are discarded. γ may of course depend on time on account of the evolving ecosystem.

After grid search we find $x = 0.9$, indicating that the random drift contributes to about 10% of survival at each cycle, and $\omega_{th} = 10$, in agreement with the observations prompted by the saturating behavior of $\langle \omega \rangle$. We also find that the data cannot be approximated with a single value of γ, but they can be very well matched assuming $\gamma = 3$ for the first 5 cycles and $\gamma = 1$ for the remaining cycles, as

Figure 2. Evolution of the DNAi population. Time is expressed in ADSE cycles. a: fraction F_D of the total population formed by different sequences obtained from experiment (red dots) and computed with the IBEE model (green line); fraction F_{10} of the total population formed by the 10 most abundant sequences (experimental: black dots, IBEE model: grey line). c: boxplots and scatterplots of $\langle \Delta G_{DR} \rangle$ in ensembles of 1000 random sequences (left), 1000 sequences extracted from the experimental population at cycle 24 (middle) and top-10 most populous sequences at the same cycle (right). The color code is assigned to each point based on its ω value (color bar). d-f: probability distributions $P(\omega)$ for at cycles 0 (d), 12 (e) and 24 (f). The orange points and line are the distributions evaluated with the null model. g-i: evolution of the abundance (expressed as fraction of the total population F) of sequences whose ω is low ($3 \leq \omega \leq 5$- panel g), medium ($6 \leq \omega \leq 10$ - panel h) and high ($11 \leq \omega$ - panel i) as obtained from the experiments (dots) and with the IBEE model (green lines). Model results are averages over 20 simulations.

The online version of this article includes the following figure supplement(s) for Figure 2:

Figure 2-figure supplement 1. Time evolution of the fraction of PCR by-products and of their $\langle \omega \rangle$.
Figure 2-figure supplement 2. Probability distributions $P(\langle \omega \rangle)$ of PCR by-products at cycles 0 and 24.
Figure 2-figure supplement 3. Time evolution of $\langle \omega \rangle$ for the replica Oligo2.
Figure 2-figure supplement 4. Probability distributions $P(\langle \omega \rangle)$ at cycles 0 and 18 for the replica Oligo2.
Figure 2-figure supplement 5. Experimental vs IBEE time evolution of the population zip ratio and of the Shannon Entropy associated to the RSA distribution.
Figure 2-figure supplement 6. Experimental vs IBEE $\langle \omega \rangle$ time evolution, for two different IBEE hyperparameters choices.
Figure 2-figure supplement 7. Experimental vs IBEE $\langle \omega \rangle$ time evolution, for different IBEE starting population sizes and compositions.

.

²¹⁹ shown in Figure 2b, green line. With the same choice of parameters, the IBEE fitness based model
²²⁰ captures the decrease in the diversity of the DNAi ecosystem (Figure 2a, green and grey lines) and
²²¹ the temporal (i.e. across generations) evolution of the relative population abundance of DNAi in
²²² the three ω intervals in Figure 2, panels g, h and i.

²²³ The effect of the system size and initial conditions on the IBEE results, discussed in the Appendix
²²⁴ 2, do not qualitatively change the models results. Error-bars on simulations have been obtained
²²⁵ by averaging 20 independent runs, starting from the same initial conditions. As it can be observed,
²²⁶ the variability among simulations is negligible.

²²⁷ The change in γ, and thus in fitness, necessary to reproduce the observed $\langle\omega\rangle$ with the IBEE
²²⁸ model is remarkable because it contrasts with the generally simple (fixed resource and low muta-
²²⁹ tion rate) conditions of ADSE environment and indicates that survival is controlled by more than
²³⁰ affinity to the resources, as discussed in the analysis below.

Self and mutual DNAi interactions are evolutionary drivers

²³² While ω is certainly a key driver of the observed evolution, it is clearly not the only one. The fact
²³³ that the top 10 most represented sequences come to be, in the last cycle, about 55% of the total
²³⁴ population (Figure 2a) implies that, even among sequences with large ω, only a tiny minority come
²³⁵ to dominate, while the largest part of them eventually disappear. Moreover, the 10 most repre-
²³⁶ sented sequences do not stand out for their particularly large ΔG_{DR} or ω, as noticeable in Figure.
²³⁷ 2c. The same data are plotted in Figure 3a as a $P(\Delta G_{DR})$ distribution to enable comparing the free
²³⁸ energy distribution in the initial population (purple shading), in the final population (blue columns)
²³⁹ and in the top 10 (black columns). These elements support the notion that survival and dominance
²⁴⁰ must be also due to factors other than ω, which we thus explored.

²⁴¹ Figure 3b shows the value, across the 24 generations, of two other contributions to the total
²⁴² free energy, normalized to their value computed in random sequences, so to enable comparing
²⁴³ their relative variations: $\langle\Delta G_{\text{self}}\rangle$ (green symbols), the average unimolecular free energy, expressing
²⁴⁴ the average strength of the internal folding of each DNAi; $\langle\Delta G_{DD}\rangle$ (red dots), the total bimolecular
²⁴⁵ DNAi-DNAi free energy, comprising both self and mutual interactions. For comparison, we also
²⁴⁶ plot the normalized value of $\langle\Delta G_{DR}\rangle$ (blue dots). As for the case of $\langle\Delta G_{DR}\rangle$, $\langle\Delta G_{\text{self}}\rangle$ and $\langle\Delta G_{DD}\rangle$
²⁴⁷ are computed by randomly selecting 1000 individuals or pairs, respectively, from the population
²⁴⁸ at cycle j and using the NUPACK tool to compute the values.

²⁴⁹ Figure 3b reveals that $\langle\Delta G_{\text{self}}\rangle$ grows in time even more than $\langle\Delta G_{DR}\rangle$, but with a different pro-
²⁵⁰ gression: $\langle\Delta G_{DR}\rangle$ grows faster in the first cycles to later saturate, while the growth of $\langle\Delta G_{\text{self}}\rangle$ is
²⁵¹ more uniform.

²⁵² Since ΔG_{DR} is computed as the free energy of the whole DNAi-resource structure, it includes
²⁵³ contributions of self energy associated to hairpins in the DNAi. Thus the growth of $\langle\Delta G_{DR}\rangle$ could
²⁵⁴ actually depend on the growth in $\langle\Delta G_{\text{self}}\rangle$ (but not the contrary). To investigate this possibility, Fig-
²⁵⁵ ure 3c displays the scatter plot between ΔG_{DR} and ΔG_{self}, for a randomly chosen subset of DNAi
²⁵⁶ at $j = 24$. In evidence are the 10 points corresponding to the 10 most populous sequences. The
²⁵⁷ plot shows weak or no correlation, and a relevant shift with respect to $\Delta G_{\text{self}} = \Delta G_{DR}$ (black line),
²⁵⁸ indicating that there is no dependence of $\langle\Delta G_{DR}\rangle$ on $\langle\Delta G_{\text{self}}\rangle$ and indicating that these two quan-
²⁵⁹ tities reflect two independent driving forces in the selection mechanism of ADSE. Their behavior
²⁶⁰ suggests that in the first stages of ADSE the selection is mostly dominated by the affinity with the
²⁶¹ resources, while in the later stages the requirement of a stronger unimolecular folding becomes
²⁶² more important.

²⁶³ A similar scatter plot analysis for $\langle\Delta G_{DD}\rangle$ yields a different outcome. Figure 3d compares ΔG_{DD}
²⁶⁴ computed for two selected DNAi (individual "k" and "l") and the sum of ΔG_{self} of the same two DNAi.
²⁶⁵ The apparent correlation indicates that, *on average*, a large part of $\langle\Delta G_{DD}\rangle$ simply embodies the
²⁶⁶ growth of self-energy, although the difference $\Delta\Delta G \equiv \Delta G_{DD,kl} - \Delta G_{\text{self},k} - \Delta G_{\text{self},l}$ (orange arrow) is
²⁶⁷ non-negligible. Figure 3b shows the behavior of $\langle\Delta\Delta G\rangle$ across cycles (orange squares). Despite the
²⁶⁸ resulting mild growth of $\Delta\Delta G$ might appear not relevant, it actually indicated that ADSE, indepen-

dently of ω, select strings that have higher reciprocal affinity than a random DNAi set (see Appendix 2). Indeed, as we could have obtained a significant decrease of the same quantity. It should also be noted that mutual interactions might involve the unfolding of hairpins self-structures, in which case their strength would be much larger than $\Delta\Delta G$.

Self-interactions and mutual interactions can compete with the binding of DNAi to the resources either when they involve the same nucleobases or through steric hindrance. Therefore, we could expect that the increase of ΔG_{DR} would lead to a decrease of both ΔG_{self} and ΔG_{DD}. Their unforeseen growth in the ADSE process is hence an indication of the selective advantage they convey. We interpret this behavior as an indirect sign that mutual DNAi interactions, more than just a impediment, are a major deadly threats for their survival. To avoid them, DNAi need screening. Indeed, the strong growth of self interactions, and the mild increment in mutual interactions, could represent the emergence of "defensive" strategies, as discussed below.

In our search for evolution quantifiers other than resource binding strength we also found that selection favors binding to resource sequences close to their resource 3' terminal, away from the bead surface (see Figure 4 - figure supplement 2). While this is expected, being that terminal much less constrained and in a less crowded environment, it also provide useful clues to further analysis.

The evolution of DNAi species

While the trends in average free energies offer some insight into the nature of the ADSE ecosystem, it has also become clear that they do not enable full understanding, especially since the dominant species are not - as one could naively expect - at the extremes of the energies distributions. To gain further insight, we examined the change in numerosity of the groups of DNAi having equal sequence, which might be regarded as the "species" of ADSE. Accordingly, the 10 most populous sequences are the 10 most successful *species*. The evolution of the populousness, expressed as the fraction F of the total population that belong to a given species, is shown in Figure 4a for a limited set of (mainly) successful species (for a larger selection see SI). Species that are later becoming dominant "appear" in our analysis only at generation 5 in a single or a few copies. This is reasonable since we sequence a sample of $\sim 10^6$ DNAi over the much larger population of $\sim 10^{12}$ survived individuals. Species are named after their ω value, combined with an index (e.g. "12.0", "12.1", "12.2" …) expressing their ranking in populousness.

ADSE species populations do not have standard patterns of evolution. This is true even when limiting the observation to the successful species in Figure 4a whose population growths are varied and do not reflect in any simple way the energy quantifiers. Conversely, species with similar energy quantifiers may have opposite fate, either becoming dominant or rapidly extinguishing or else displaying a non-monotonic population evolution. It is worth pointing out that all species start equal, each represented in the original pool by a single molecule, which on this scale would appear as a $F \approx 10^{-12}$ (black dot in Figure 4a).

We thus decided to seek further understanding on the survival of the fittest in ADSE by inspecting the "natural history" of a few species (see Tables 2, 3).

13.0 and 11.0 (orange and gray dots in Figure. 4a) are, respectively, the most and second most populous species in the last generation. It might be relevant to mention that the replica of ADSE described in the SI leads to different dominant sequences, as expected on the basis of the distinct initial pool and on the key role of randomness and sampling in the first cycles. At generation 24, 13.0 and 11.0 have grown to more than 85% and 53% of DNAi with $\omega = 13$ and $\omega = 11$, respectively, a further indication that ω is only one of the ingredients of the fitness. 13.0, and 11.0 likewise, have a weak self-interaction (Figure 3c), corresponding to hairpins that provide a mild defense to mutual interactions, as schematically sketched in Figure 4, panel b1. The hairpins involve most of the bases that are complementary to the resource (green shading), leaving however a few unpaired bases that might act as toehold to initiate resource binding. The resource binding takes place at its 3' terminus (panel b2). A bit puzzled about the success of these two species we also explored their capacity of mutual interactions and found that both of them are capable of forming

Figure 3. Distribution and evolution of free energy quantifiers. a: probability distribution for the DNAi-resource binding free energy computed by NUPACK, $p(\langle \Delta G_{\mathrm{DR}} \rangle)$, for the initial population (grey shading), for the final population (random choice of 1000 DNAi - cyan columns, top 10 most populous sequences - black columns). b: time evolution, expressed in cycles, for various mean free energies $\langle \Delta G \rangle$, normalized to their value computed on pools of random sequences. All $\langle \Delta G \rangle$ are computed by NUPACK on sets of 1000 individuals: $\langle \Delta G_{\mathrm{DR}} \rangle$ (blue dots); unimolecular self interaction $\langle \Delta G_{\mathrm{self}} \rangle$ (green dots); bimolecular mutual DNAi interaction $\langle \Delta G_{\mathrm{DD}} \rangle$ (red dots); mutual, self-subtracted interaction $\langle \Delta\Delta G \rangle$ (yellow dots). c: scatter plot of ΔG_{self} vs. ΔG_{DR} computed for 1000 DNAi in the final population (blue squares). Red dots mark the point relative to the 10 most populous sequences, as identified by the labels. Note that the x-axis scale of panels a and c is the same, enabling identifying sequences. d: scatter plot computed on 10^4 DNAi pairs from the final population comparing $\Delta G_{\mathrm{self,j}} + \Delta G_{\mathrm{self,l}}$ and $\Delta G_{\mathrm{DD,kl}}$ (green squares). Red dots mark the pair formed by the 10 most populous sequences, some of which identified by labels. With respect to the condition $\Delta G_{\mathrm{DD,kl}} = \Delta G_{\mathrm{self,j}} + \Delta G_{\mathrm{self,l}}$ (black line), data are on average displaced by $\Delta\Delta G \sim 7.5$ kcal/mol (yellow arrow). The online version of this article includes the following figure supplement(s) for Figure 3:

Figure 3-figure supplement 1. Three examples of different relative positions for sequence-sequence attachment.

Figure 3-figure supplement 2. Scheme of target-consumer interaction, from a combinatorial standpoint.

Figure 3-figure supplement 3. Null model without threshold, analytical and simulated, with an even and uneven nucleotides distribution.

Figure 3-figure supplement 4. Null model with/without threshold, simulated and analytical.

.

homodimers, as shown in panel b3 for 13.0 (binding energy marked in Figure 3d). Such dimers can bind resources at both ends (green shading regions), giving 13.0 and 11.0 self-screening and divalent binding capacity. We argue this to be a crucial drive toward success of these two species, which also explains why their growth has such an increment in the latest generations, following the probability of dimer formation.

The third most populous sequence (species 8.0) is characterized by a weak resource interaction which includes a bulge (see Figure 3c). While its survival rationale appears clear - good defensive self-binding (panel c1) and interaction targeting the 3' end of the resource (panel c2) - it is hard to accept this could be the cause of the success. Again, inspecting the mutual interaction we found that 8.0 efficiently interacts with 13.0 (as marked Figure 3d), in a region (pink shading) that does not conflict with the capacity of 13.0 to bind to the resource (panel c3). We speculate that this parasitic capacity of 8.0 adds to the weak intrinsic binding to lead to a remarkable success of this species.

As a test to this concept, we focused on species 5.1, whose binding strength to the resource is the weakest in the top 10 species and one of the weakest among all the survivors of generation 24 (Figure 3c), a situation that cannot justify its success. However, by inspecting mutual interactions, we find its binding to 11.0 to be strong and stable (pink shading panel d1 and Figure 3d). An analogous behavior is found for species 5.0 and its interaction to 12.0 (panel d2), confirming that parasitism is an emerging successful survival solution in ADSE.

We then inspected 12.1 since it has largest ΔG_{DR} and ΔG_{self} among the top 10 species (Figure 3c). Species 12.1 has all the features to succeed in ADSE: it forms a weak hairpin with the bases involved in the binding with the resources and a strong hairpin that protects the rest of the ssDNA segment (panel e1); it strongly binds to the resource with the bond ending at its 3' terminal (panel e2); it forms homodimers capable of double resource binding, analogous to that of 13.0 (panel e3). Remarkably, 12.1 emerges only late in ADSE (Figure 4a, blue dots), a very unusual behavior when compared to the other many species we have considered. The combination of this late appearance and of the remarkable growth in F afterward suggests that 12.1 is the outcome of one of the rare *mutations* that can occur even with the high quality PCR that we adopted. By assuming the initial growth as with the dominant species (black arrows in panel a), we argue that such mutation occurred around generation 9, where we have found ancestors having sequence equal to 12.1 except for 4 bases.

F has a monotonic increase for most of the species that become dominant in the latest generations. This behavior is however not at all general. Non-monotonic F, growing in the first generation and decreasing to extinction afterward, is the general behavior for the majority of species that do not become extinct in the first cycle, as also shown - as mean behavior - in Figure 2h. An examples of these is species 7.X, which we plot in panel a of Figure 4 (empty dots).

More intriguing is when a non-monotonic behavior is observed for one of the dominating species. This is the case of 11.1, whose growth can be again attributed to an effective screening and a good binding to the resources, but whose decrease is harder to justify. To understand it we again investigated mutual interactions, and we found a strong 11.0-11.1 bond (sketched in panel f), which, differently from the DNAi-DNAi complex examined above, competes with resource binding. We thus speculate that the fall of 11.1 is a consequence of the rise of 11.0, which has become frequent enough to make the formation of the complex 11.0-11.1 probable, which lowers the survival probability of both species. This could also explain why species 11.0 has such an irregular growth pattern, with a slowing down when 11.1 becomes highly populated.

The insight gained by these case-studies enables appreciating how self and mutual interactions can affect the fate of species in ADSE beyond what we could discern through distributions and average quantifiers. Indeed, this analysis shows that the success of the dominant ASDE species can be achieved through different combinations of resource and mutual DNAi interactions, a fact that makes the evolution of population so varied and generation dependent and explains why, in the latest generations of ADSE, the population distribution $p(\Delta G)$ (Figure 3a) is so irregular.

Figure 4. Natural history of DNAi species. a: fraction F of {DNAi} that belong to a choice of specific species as a function of the ADSE cycles, in linear (left) and logarithmic (right) scale. Arrows connect the initial condition (one individual per species) to the earliest detection via sequencing, across the 6 order of magnitudes gap (grey shading) between the population of DNAi survived at each cycle and the amount of sequenced DNAi. The same growth is assumed for species 12.1, suggesting its appearance by mutation occurred at generation 9. b-f: self interactions (b1, c1, e1), resource interactions (b2, c2, e2, d1, d2) and mutual interactions (b3, c3, e3, d1, d2, f) of selected species, sketched as per the NUPACK output. Nucleobases color coded (G - black, C - blue, A - green, T - red). Paired bases are connected. Double and single stand regions are represented as straight and curved, respectively. As in Figure 1a, terminal blocks of DNAi are marked as graphic double helices colored according to the legend of panel a, and beads as sketched yellow spheres. Yellow shading: section of DNAi complementary to resources. Pink frames: regions of hybridization between DNAi. b: interactions involving species 13.0 including its homodimerization (b3). c: interactions involving species 8.0, including its binding to 13.0 (c3). e: interactions involving species 12.1, including its homodimerization (e3). d and f: DNAi heterodimers interactions suggesting parasitism (d1), and possibly mutualism (d2) and mutual damage (f). The online version of this article includes the following figure supplement(s) for Figure 4:

Figure 4-figure supplement 1. Evolution of the intra-species interaction strengths ($\langle \Delta G_{pp} \rangle$).
Figure 4-figure supplement 2. 2D probability for a consumer, shown at different experimental cycles, to have a given $\langle \omega \rangle$ jointly with a given attachment coordinate to a target strand.

Discussion and Conclusion

We have here introduced ADSE, a synthetic molecular evolution protocol that exploits sequencing technology and DNA interaction computability to provide a test-bed for key concepts in ecology and evolution. We performed experiments by adopting the most simple environment within this protocol, *i.e.* by holding fixed the capture sequence and minimizing mutations, to achieve a condition dominated by competition and selection, which could enable investigating the nature of fitness in this simplified evolution process.

Our experiments and their comparison with theoretical models indicate that fitness is not a simple function of the direct competitive advantage of strong binding to the resources, as it could have been naively expected. Resource binding is indeed the dominant factor in the first part of the evolution (cycles 1-5), producing a fast growth of $\langle \omega \rangle$ compatible with a strong dependence of the survival probability on ω ($f \propto \omega^3$). However, as $\langle \omega \rangle$ reaches a value corresponding to bonds of moderate stability **Woodside et al.** (**2006**) - approximately from generation 6 onward - the selective pressure related to resource binding decreases ($f \propto \omega$), a condition enabling to appreciate other factors at play. Which are these factors is suggested by the different growth patterns of ΔG_{DR} and ΔG_{self}, the former dominating the first cycles while the latter drives the later stages. However, the quantification of binding energies isn't sufficient to predict the fate of individual species, that also depends on interaction details (location of binding and hairpin spots, secondary structures) and possibly on kinetics.

In fact, by design, DNAi can interact with themselves, both internally - forming hairpin-like structures - and mutually - leading to the formation of complexes. These interactions can either coexist - when they involve non-overlapping sequences - or they can become competitive. Our experiments reveal that, among the wide variety of conditions made available by the initial random seed, ADSE generally promotes those that combine a good interaction with the resources and a structure capable of screening from mutual DNAi interactions conflicting with binding to the beads. By exploring hybridisation-dependent DNAi secondary structures - the only ones accessible with simple analytical tools, we found three basic motifs: hairpins due to self-interactions (as for species 12.1), formation of DNAi homoduplets (as for species 13.0 and 11.0), and formation of dimers of DNAi belonging to distinct species. The latter can bring to distinct prototypical behavior: it can be the basis of *parasitism*, as in the case of the pairs 8.0-13.0 and 5.1-11.0, it can lead to *extinction* with no benefit for either species (as in the case of 15.0-12.0), or it can provide - at least in principle - *mutualism*. While we do not have a solid proof, clues suggest that cooperation might be the case of the interaction between species 5.0 and 12.0, since 12.0 has a weak self-defensive system and since the populations of 12.0 and 5.0 evolve very similarly (see Table 2). Finally, we cannot exclude a possible role of the formation of multi-DNAi multivalent complexes. This is the case of species 12.0 and 15.0 whose pattern of multivalent mutual interactions can, in principle, allow higher order interactions (see Appendix 2).

Fitness in ADSE is the outcome of a complex interplay of all these elements. The history of single species in Figure 4 indicates that the nature of fitness is beyond what can be captured by analyzing the probability distributions of the relevant parameters describing structure and interactions. Moreover, the variety of population evolution enlightens the fundamental fact that despite the constancy of resources, the progressive modification of the ecosystem brings about a change in the competition patterns and thus in fitness. A forthcoming work focused on a limited number of species will be devoted to better disentangle the nature of fitness in ADSE.

The evolution of the ADSE ecosystem, which corresponds to a marked decrement of its entropy - with 2/3 of the initial species become extinct in the first 5 generations - and potentially terminating with the indication of a single winner species leads instead, in the last cycles, to a significant number of dominant species all still growing at the expenses of subservient ones, indicating that in their direct competition none of them is strongly prevailing despite - or maybe thanks to - their distinct survival strategies and their very different population share. Hence, while the niche hypothesis

could still be verified in the long run - beyond the experimental limitations due to PCR - it is clear that its drive is weak, suggesting that it could be overcome by environmental fluctuations, thus allowing for coexistence of different species even in a single niche environment.

Methods

Our experimental design takes advantage of a selective capture mechanism where magnetic beads carrying single-stranded DNA filaments of fixed length and sequence (resources) target DNA individuals (DNAi) (Figure 1 - figure supplement 1) present in a DNA library based on their level of complementarity. This process of selection is carried out through subsequent steps that are described in detail in the next paragraphs and represented in Figure 1-figure supplement 2.

Library design

The DNA library contains 100-nt-long sequences where a randomized central region of 50 nucleotides is flanked by 25-nt-long fixed sequences at both its 5' and 3' ends. The fixed regions provide an anchor point for primer annealing, required to perform PCR reactions during the amplification phase (see below). These terminal segments are made inactive by hybridization with oligomers of perfect complementarity (blockers), so that they are not involved in the selection phase. The blockers, as well as the fixed regions of the DNA library, have been designed to avoid any interactions with the resources. In addition, the blockers carry a phosphate group at their 3' end to prevent them from functioning as primers during the PCR amplification. Following the above described criteria, we designed two sets of sequences (Table 1), called Oligo1 (whose results are presented in the main text) and Oligo2 that was used as a replication experiment. All the oligonucleotides used in this work were purchased from Integrated DNA Technologies, Coralville, IA, USA (Table 1).

Beads preparation

The capture of DNAi within the DNA library was performed with carboxylic acid magnetic beads (M-270 Dynabeads, Invitrogen, Carlsbad, CA) coated with the resources. The resources are 20-nucleotide-long, 5'-amino-modified oligonucleotides. Their coupling to the beads surface was performed according to the manufacturer's instructions. Following the activation and coupling procedure, the beads were washed in Tris-HCl (50mM, pH 7.4), and stored in the same buffer in single-use aliquotes.

Sample preparation

The starting samples were prepared in 1X SSC buffer (0.15M sodium chloride, 15mM sodium citrate, pH 7.0). In detail, the Oligo1 (or Oligo2) library (0.75nmol) was mixed with the blockers (2.25nmol) in 1:3 molar ratio to saturate all the available interaction sites between the blockers and the fixed regions of the DNA library. The sample was then denatured at 95°C for 5 minutes, then slowly brought to room temperature using a thermal cycler (MasterCycler Nexus Gradient, Eppendorf, Hamburg, Germany).

Selection phase

The beads, coupled with the resources, are mixed with the sample, prepared as described above. The capture of the DNAi is carried out at 40°C for 2 hours in stirring (600rpm, ThermoMixer, Eppendorf). Once the selection phase is completed, the sample is incubated for 2 minutes on a magnet and the supernatant is removed. The beads, that are now bound to the captured DNAi, are washed three times in SSC 1X buffer to eliminate the aspecific sequences. Finally, the beads are resuspended in water and incubated for 5 minutes at 60°C to recover the DNAi from the resources. The sample is quickly placed on the magnet and after 2 minutes the supernatant containing the selected DNAi is collected.

Amplification phase

The captured DNAi are then amplified by PCR with Q5® Hot Start High-Fidelity DNA Polymerase (New England Biolabs, Ipswich, MA, USA). DNA samples were diluted 10 times and 3μl were used for each PCR reaction. The PCR was performed in a final volume of 25μL using 0.25μL of polymerase, and each primer had a final primer concentration of 2.5μM. To allow the regeneration of the single strand library, the reverse primer was designed to carry a 5' biotin modification (see next paragraph for details). The thermal protocol was the following: i) denaturation step at 98 ▪ C for 2 minutes, ii) 28 cycles characterized by three thermal steps: 10 seconds at 98▪C, 10 seconds at 68▪C for Oligo1 (69▪C for Oligo2) and 1 second at 72▪C, iii) 2 minutes at 72▪C. For each generation, 10 different PCR reactions were performed. The PCR products were then checked for size on a 2% TBE 1X agarose gel.

Regeneration phase

PCR products were purified by precipitation with 2.5 volumes of ethanol and 0.1 volumes of sodium acetate 2M, pH 5.2. After an overnight precipitation at -20▪C, the pellet was washed with ethanol 75%, resuspended in water and quantified using the Qubit fluorometer (Invitrogen, Waltham, MA, USA). The single-strand regeneration was performed with streptavidin-coated magnetic beads (M-270 Dynabeads, Invitrogen, Carlsbad, CA) according to manufacturer's instructions. Briefly, the PCR product was incubated for 15 minutes on a rotator with the magnetic beads. After 2 washes with binding and washing buffer (10mM Tris–HCl pH 7.5, 1mM EDTA, 0.2M NaCl) and a third one with SSC 1X buffer, an alkaline denaturation was performed by incubating the beads with 150mM NaOH for 10 minutes to induce the separation of the two DNA strands. This way it is possible to recover the unlabeled DNA strand in solution. The ssDNA is then collected and buffered with 1.25M acetic acid and TE 1X. The regenerated ssDNA is ready for the next round of selection and amplification.

Sequencing of the recovered products

To check the growth of DNAi species in our experimental model, some generations were sequenced through next-generation sequencing techniques. To prepare the sample for sequencing, the DNAi species captured as described in the "Selection phase" paragraph were first PCR-amplified using the same conditions discussed before. However, in this case, both forward and reverse primers were unlabelled. After PCR purification, performed as already described (see "regeneration phase" paragraph), the products were quantified and used to obtain libraries. About 300ng of DNA were used as starting material for the NEBNext® Ultra™ II DNA Library Prep Kit for Illumina® (New England Biolabs), and libraries were prepared following the manufacturer's instruction. Sequencing was performed with the NextSeq™ 550 sequencer (Illumina, San Diego, CA, USA) and a paired-end strategy to obtain 75-nt long reads.

Replicas

The experimental workflow was repeated on two different sets of DNA libraries: Oligo1, whose results are described in the main text, and Oligo2, used as a replicate. The experimental conditions and procedures were the same for both sets of oligonucleotides, the only differences being the sequences of the fixed parts of the DNA libraries (and hence of the blockers and primers) and of the resources (Table 1). As a consequence, also the PCR conditions had to be adjusted, in terms of the annealing temperature that is one degree higher for Oligo2.

Eco-Evolutionary algorithm

To support experimental observation with a simple abstract model, we developed an Evolutionary Algorithm where a population of N_p sequences evolves in presence of $N_r < N_p$ shorter sequences. In particular, N_r individuals in this population are selected at each cycle depending on their fitness, i.e. on their affinity to the resource, expressed via their ω with it. These survived sequences are

then amplified by a factor of (roughly) N_p/N_r. In this work, two types of fitness functions have been
explored: the first one is merely linearly proportional to the ω of each individual (i.e. $\frac{\omega}{\Sigma\omega}$, where
$\Sigma\omega$ is the sum of the ω values of the whole population), while the other one is a modification of
the previous fitness which sets it to be $\frac{\omega_{th}}{\Sigma\omega}$ beyond a threshold value ω_{th}. The code is written in
C++, exploiting MPI *Message Passing Interface Forum* (*2021*) and Armadillo *Sanderson and Curtin*
(*2016*, 2018) libraries for acceleration.

NUPACK calculations

We resorted to NUPACK for nucleotide sequences analysis, for the prediction of their free ener-
gies at equilibrium, either alone or when binding to one or two other oligomers. Schemes like
those shown in Figure 4 have been obtained via the NUPACK web application *Zadeh et al.* (*2011*),
while massive ΔG calculations have been performed with custom Python codes exploiting NU-
PACK Python package (v4.0.0.27) *Fornace et al.* (*2020*); *Dirks et al.* (*2007*). The model is dna04 and
ensemble='some-nupack3'; $T = 40$·C and $[Na^+] = 0.24$ M, as in the experiments. Concentration of
each species in the tube has been arbitrarily set to 10^{-6} M. Detection of hairpins and other sec-
ondary structures has been performed visually (thanks to the oxView software *Bohlin et al.* (*2022*);
Poppleton et al. (*2020*)).

Acknowledgments

528 T.B. acknowledges support from MIUR-PRIN (Grant No. 2017Z55KCW). F.M. and S.S. acknowledge
529 CloudVeneto (http://cloudveneto.it, *Andreetto et al.* (*2019*)) for the use of computing and storage
530 facilities, through the SEDES and HPC-Physics projects.

Conflict of interest

532 The authors declare no conflict of interest.

Availability of data and materials

534 The datasets generated and analysed during the current study are available from the correspond-
535 ing author on reasonable request.

Code availability

537 The Jupyter Notebooks used for the present analysis are available at https://github.com/francescomambretti/
538 stat_phys_synthetic_biodiversity.

Author Contributions

540 SS, EMP and TB conceived and designed the study; LC and EMP designed protocols and performed
541 the experiments; FM, AT and SS analyzed the data and developed the theoretical models; LC and
542 FM performed NUPACK analysis; LC, FM, SS and TB interpreted the data and wrote the manuscript.

References

Adamala K, Szostak JW. Competition between model protocells driven by an encapsulated catalyst. Nature
Chemistry. 2013 May; 5(6):495–501. https://doi.org/10.1038/nchem.1650, doi: 10.1038/nchem.1650.

de Aguiar MAM, Baranger M, Baptestini EM, Kaufman LS, Bar-Yam Y. Global patterns of speciation and diversity.
Nature. 2009; 460:384–387.

Anceschi N, Hidalgo J, Bellini T, Maritan A, Suweis S. How neutral and niche forces contribute to speciation
processes. arXiv: Populations and Evolution. 2018; .

Andreetto P, Chiarello F, Costa F, Crescente A, Fantinel S, Fanzago F, Konomi E, Mazzon P, Menguzzato M,
Segatta M, Sella G, Sgaravatto M, Traldi S, Verlato M, Zangrando L. Merging OpenStack-based private clouds:
the case of CloudVeneto.it. EPJ Web of Conferences. 2019; .

553 **Benedetto D**, Caglioti E, Loreto V. Language trees and zipping. Physical Review Letters. 2002; 88(4):048702.

554 **Bohlin J**, Matthies M, Poppleton E, Procyk J, Mallya A, Yan H, Šulc P. Design and simulation of DNA, RNA and
555 hybrid protein–nucleic acid nanostructures with oxView. Nature Protocols. 2022 Aug; 17(8):1762–1788. https:
556 //doi.org/10.1038/s41596-022-00688-5, doi: 10.1038/s41596-022-00688-5.

557 **Camacho Mateu J**, Sireci M, Muñoz MA. Phenotypic-dependent variability and the emergence of tolerance in
558 bacterial populations. PLoS computational biology. 2021; 17(9):e1009417.

559 **Catalán P**, Arias CF, Cuesta JA, Manrubia S. Adaptive multiscapes: an up-to-date metaphor to visualize molec-
560 ular adaptation. Biology Direct. 2017; 12:1–15.

561 **Charalambides CA**. Enumerative Combinatorics. SIGACT News. 2008 nov; 39(4):25–27. https://doi.org/10.
562 1145/1466390.1466395, doi: 10.1145/1466390.1466395.

563 **Chase JM**, Leibold MA. Ecological niches. University of Chicago Press; 2009.

564 **Davis WB**. Mayr, Ernst. Systematics and the Origin of Species. Columbia University Press, 334 pp., 29 figs.
565 Journal of Mammalogy. 1943 06; 24(2):273–274. https://doi.org/10.2307/1374810, doi: 10.2307/1374810.

566 **De Visser JAG**, Krug J. Empirical fitness landscapes and the predictability of evolution. Nature Reviews Genetics.
567 2014; 15(7):480–490.

568 **Di Leo S**, Marni S, Plata CA, Fraccia TP, Smith GP, Maritan A, Suweis S, Bellini T. Pairing statistics and melting of
569 random DNA oligomers: Finding your partner in superdiverse environments. PLoS computational biology.
570 2022; 18(4):e1010051.

571 **Dieckmann U**, Doebeli M. On the origin of species by sympatric speciation. Nature. 1999; 400:354–357.

572 **Dirks RM**, Bois JS, Schaeffer JM, Winfree E, Pierce NA. Thermodynamic Analysis of Interacting Nucleic Acid
573 Strands. SIAM Review. 2007; 49(1):65–88. https://doi.org/10.1137/060651100, doi: 10.1137/060651100.

574 **Ellington AD**, Szostak JW. In vitro selection of RNA molecules that bind specific ligands. Nature. 1990 Aug;
575 346(6287):818–822. https://doi.org/10.1038/346818a0, doi: 10.1038/346818a0.

576 **Fornace ME**, Porubsky NJ, Pierce NA. A Unified Dynamic Programming Framework for the Analysis of Interact-
577 ing Nucleic Acid Strands: Enhanced Models, Scalability, and Speed. ACS Synthetic Biology. 2020; 9(10):2665–
578 2678. https://doi.org/10.1021/acssynbio.9b00523, doi: 10.1021/acssynbio.9b00523, pMID: 32910644.

579 **Fussmann GF**, Loreau M, Abrams PA. Eco-evolutionary dynamics of communities and ecosystems. Functional
580 ecology. 2007; p. 465–477.

581 **Ghosh S**, Takahashi S, Endoh T, Tateishi-Karimata H, Hazra S, Sugimoto N. Validation of the nearest-neighbor
582 model for Watson–Crick self-complementary DNA duplexes in molecular crowding condition. Nucleic acids
583 research. 2019; 47(7):3284–3294.

584 **Gupta D**, Garlaschi S, Suweis S, Azaele S, Maritan A. Effective resource competition model for species coexis-
585 tence. Physical review letters. 2021; 127(20):208101.

586 **Hill MO**. Diversity and evenness: a unifying notation and its consequences. Ecology. 1973; 54(2):427–432.

587 **Ichihashi N**, Usui K, Kazuta Y, Sunami T, Matsuura T, Yomo T. Darwinian evolution in a translation-coupled
588 RNA replication system within a cell-like compartment. Nature Communications. 2013 Oct; 4(1). https://doi.
589 org/10.1038/ncomms3494, doi: 10.1038/ncomms3494.

590 **Katla SK**, Lin C, Pérez-Mercader J. Competitive exclusion principle among synthetic non-biochemical proto-
591 cells. Cell Reports Physical Science. 2023 Apr; 4(4):101359. https://doi.org/10.1016/j.xcrp.2023.101359, doi:
592 10.1016/j.xcrp.2023.101359.

593 **Kauffman WB**, Guha S, Wimley WC. Synthetic molecular evolution of hybrid cell penetrating peptides. Na-
594 ture Communications. 2018 Jul; 9(1). https://doi.org/10.1038/s41467-018-04874-6, doi: 10.1038/s41467-018-
595 04874-6.

596 **Keymer JE**, Fuentes MA, Marquet PA. Diversity emerging: from competitive exclusion to neutral coexistence
597 in ecosystems. Theoretical Ecology. 2012; 5:457–463.

598 **Levin SA**. Community equilibria and stability, and an extension of the competitive exclusion principle. The
599 American Naturalist. 1970; 104(939):413–423.

Mambretti F, Pedrani N, Casiraghi L, Paraboschi EM, Bellini T, Suweis S. OxDNA to Study Species Interactions. Entropy. 2022; 24(4). https://www.mdpi.com/1099-4300/24/4/458, doi: 10.3390/e24040458.

Mambretti F, Pedrani N, Casiraghi L, Paraboschi EM, Bellini T, Suweis S. OxDNA to Study Species Interactions. Entropy. 2022; 24(4). https://www.mdpi.com/1099-4300/24/4/458, doi: 10.3390/e24040458.

Message Passing Interface Forum. MPI: A Message-Passing Interface Standard Version 4.0; 2021, https://www.mpi-forum.org/docs/mpi-4.0/mpi40-report.pdf.

Metropolis N, Rosenbluth AW, Rosenbluth MN, Teller AH, Teller E. Equation of state calculations by fast computing machines. J Chem Phys. 1953 3; 6(21). doi: 10.2172/4390578.

Parrilla-Gutierrez JM, Tsuda S, Grizou J, Taylor J, Henson A, Cronin L. Adaptive artificial evolution of droplet protocells in a 3D-printed fluidic chemorobotic platform with configurable environments. Nature Communications. 2017 Oct; 8(1). https://doi.org/10.1038/s41467-017-01161-8, doi: 10.1038/s41467-017-01161-8.

Peterson AT. Ecological niche conservatism: A time-structured review of evidence. Journal of Biogeography. 2011; 38(5):817–827.

Pigolotti S, López C, Hernández-García E. Species clustering in competitive Lotka-Volterra models. Phys Rev Lett. 2007; doi: 10.1103/PhysRevLett.98.258101.

Plata CA, Marni S, Maritan A, Bellini T, Suweis S. Statistical physics of DNA hybridization. Physical Review E. 2021; 103(4):042503.

Poppleton E, Bohlin J, Matthies M, Sharma S, Zhang F, Šulc P. Design, optimization and analysis of large DNA and RNA nanostructures through interactive visualization, editing and molecular simulation. Nucleic Acids Res. 2020 Jul; 48(12):e72.

Rundle H, Nosil P. Ecological speciation - Rundle - 2005 - Ecology Letters - Wiley Online Library. Ecology letters. 2005 01; .

Sanderson C, Curtin R. Armadillo: a template-based C++ library for linear algebra. Journal of Open Source Software. 2016; 1(2):26. https://doi.org/10.21105/joss.00026, doi: 10.21105/joss.00026.

Sanderson C, Curtin R. A User-Friendly Hybrid Sparse Matrix Class in C++. In: Davenport JH, Kauers M, Labahn G, Urban J, editors. *Mathematical Software – ICMS 2018* Cham: Springer International Publishing; 2018. p. 422–430.

SantaLucia Jr J. A unified view of polymer, dumbbell, and oligonucleotide DNA nearest-neighbor thermodynamics. Proceedings of the National Academy of Sciences. 1998; 95(4):1460–1465.

Solé R, The major synthetic evolutionary transitions. The Royal Society; 2016.

Thurner S, Hanel R, Klimek P. Physics of evolution: Selection without fitness. Physica A: Statistical Mechanics and its Applications. 2010; 389(4):747–753.

Tizei PAG, Csibra E, Torres L, Pinheiro VB. Selection platforms for directed evolution in synthetic biology. Biochemical Society Transactions. 2016 Aug; 44(4):1165–1175. https://doi.org/10.1042/bst20160076, doi: 10.1042/bst20160076.

Tolle F, Wilke J, Wengel J, Mayer G. By-product formation in repetitive PCR amplification of DNA libraries during SELEX. PloS one. 2014; 9(12):e114693.

Tuerk C, Gold L. Systematic Evolution of Ligands by Exponential Enrichment: RNA Ligands to Bacteriophage T4 DNA Polymerase. Science. 1990; 249(4968):505–510. https://www.science.org/doi/abs/10.1126/science.2200121, doi: 10.1126/science.2200121.

Vetsigian K. Diverse modes of eco-evolutionary dynamics in communities of antibiotic-producing microorganisms. Nature Ecology & Evolution. 2017; 1(7):0189.

Wiser MJ, Lenski RE. A comparison of methods to measure fitness in Escherichia coli. PloS one. 2015; 10(5):e0126210.

Woodside MT, Behnke-Parks WM, Larizadeh K, Travers K, Herschlag D, Block SM. Nanomechanical measurements of the sequence-dependent folding landscapes of single nucleic acid hairpins. Proceedings of the National Academy of Sciences. 2006; 103(16):6190–6195.

Zadeh JN, Steenberg CD, Bois JS, Wolfe BR, Pierce MB, Khan AR, Dirks RM, Pierce NA. NUPACK: Analysis and design of nucleic acid systems. J Comput Chem. 2011 Jan; 32(1):170–173.

 Appendix 1

 ## Data analysis protocol

Since we performed a paired-end sequencing, for each experimental cycle two FASTQ files were
generated (R1 and R2).

The data shown in the paper all come from the cumulative analysis of R1 and R2 files. To
analyze such files, we have developed an ad-hoc package (in Python and C++), currently maintained
at https://github.com/francescomambretti/stat_phys_synthetic_biodiversity. Here follows a scheme of
the main steps of the analysis:

1. Read FASTQ file, extracting the list of single-stranded DNA (ssDNA) sequences. Reverse se-
quences are reverted and complemented, so to have all the sequences oriented as $5' \to 3'$.
Fixed sequences at the ends are removed in this way: if the full blocker is present, then the
50 bases after it are retained. If some of the last bases of the blocker are lost (it may happen),
then just remove the primer and keep the subsequent 50 bases. Slightly different choices for
this cut of the sequences turned out to be substantially irrelevant for all the results.

2. Each sequence can be either saved or discarded, according to some filtering procedure. Cri-
teria include the minimum quality of the read (≥ 10, measured according to the *Phred* score),
the minimum and maximum allowed length, the presence of invalid nucleotides (indicated
by 'N' in the FASTQ file) and other checks (see below in particular for PCR by-products).

3. The 'survived' sequences are processed, computing the related observables. Unique sequences
are identified and their degeneracy is computed. This allows to monitor the time evolution
of the abundance of each species (where the equivalence: DNA sequence-species holds).

4. For each of these unique ssDNAs, we compute the indicators quantifying the 'affinity' with
the target sequence: in particular, the maximum consecutive overlap (MCO) and eventually
the binding free energy with the resource and/or other oligomers.

5. Several post-processing options are available, such as: generating histograms of the ssDNA
population based on their affinity with the resource; tracking the time evolution of dominant
strands; determining the fraction of the total population covered by the n most abundant
strands. All these quantities can be plotted by suitable Python scripts, provided in the GitHub
repository.

By-Product Formation from PCR Amplification

During the amplification phase, described in our workflow, we observed the formation of PCR by-
products. This phenomenon is known in SELEX systems, especially when an oligonucleotide library
characterized by high heterogeneity is PCR-amplified *Tolle et al.* (*2014*). The generation of these by-
products is caused by unspecific hybridization of the primers and/or of oligonucleotides, leading
to PCR products abnormal in terms of length and distribution of random/fixed sequences. To limit
this phenomenon we adopted blocker oligos with a phosphate group at 3', so that they do not
act as primers, and a stringent PCR protocol, characterized by: i) a high annealing temperature, ii)
fewer PCR cycles, iii) a high primer concentration, iv) a very short elongation step (1s). Moreover,
during the data analysis step, we developed an algorithm to identify and remove these spurious
sequences.

PCR by-products are identified among the sequenced oligomers according to the following cri-
teria:

• if either the sequence (after the cut of the blocker as previously described) contains at least
n consecutive bases identical to n of the $5'$ blocker (i.e. an n-long MCO with a subsequence of
the Primer1-F)

• or it has at least n consecutive bases identical to n of the $3'$ blocker

then this sequence is considered as a by-product and excluded from the analysis.

Figure 2-figure supplement 1a shows how such unwanted sequences become $\approx 30\%$ of the total
sampled population at cycle 15 and then the striking majority ($> 95\%$) in the last cycle, setting *de*

698 *facto* an upper limit to the number of feasible experimental cycles. On the other hand, as we can
699 see from Figure 2-figure supplement 1b, the behaviour of $\langle \omega \rangle$ for the by-products is very similar to
700 the one observed considering only regular oligomers, therefore their presence does not affect the
701 main outcome of the experiment.
702 In Figure 2-figure supplement 2 the $P(\omega)$ for the first and last cycle of only the by-products of
703 Oligo 1 are displayed and compared with the null model prediction. The main evolutive trend is
704 also present in these sequences, which have however been discarded from all the analyses.

Further analyses of the Eco-Evolutionary Model

We here show some further results and sensitivity analyses of the Eco-Evolutionary stochastic in-
dividual based model. In Figure 2-figure supplement 5 we show how the eco-evolutionary model
can describe the emergent order and complexity, that is a loss of the initial huge diversity of the
random strings towards few dominating species, during evolution. This behaviour can be quanti-
fied 1) by the ratio CR between the zipped file size *Benedetto et al.* (*2002*) and the original file size
of all the sequences oligomers in the ecosystem or 2) by computing the Shannon entropy of the
Relative Species Abundance *Hill* (*1973*). We can see that that the model is able to reproduce, at
least qualitatively, the marked decrease of diversity observed during evolution.

In Figure 2-figure supplement 6a we show the outcome of the model if we do not include the
saturation at $\omega_{th} = 10$. As we can see, the effect is evident as after a few cycles the experimental
$\langle \omega \rangle$ lies significantly under the model (which promotes too much the strongest sequences). Simi-
larly, Figure 2-figure supplement 6b shows the evolution for the average MCO in the case we use
only $\gamma = 1$ in the simulation (here, $\omega_{sat} = 11$). It is very apparent how in this case the simulated
initial evolution is much slower with respect to the observed from the data, while the final trend
is captured. These results indicate that both saturation and the non-linear fitness time dependent
fitness function are fundamental ingredients to describe the empirical evolution of $\langle \omega \rangle$. Note that
in both cases of the above picture, there is a 10% random sampling drift.

Finally, Figure 2-figure supplement 7a and Figure 2-figure supplement 7b show the simulations
results for different total number of individuals ($N = 10^6$, 10^5 and 10^4) and resources ($R = N/100$,
keeping this ratio fixed) and for different initial conditions (i.e., different sequences populations),
respectively. We can see that the qualitative results concerning $\langle \omega \rangle$ as a function of time are robust,
beyond a given system's size (the silver data in Figure 2-figure supplement 7a are already of a
system's size large enough, while the blue ones correspond to a too small system). Considering
a different initial (random) population of sequences, keeping fixed all the other hyperparameters
(in particular $N = 10^6$ and $R = 1064$), as shown by Figure 2-figure supplement 7b preserves the
qualitative agreement with the experimental data, assessing the substantial independence on the
initial conditions. Overall, these further analyses show the robustness of the qualitative behavior
of results of the Eco-Evolutionary model, pointing out to two relevant phases of the evolution of
the oligomers, as highlighted in the main text.

Further results on the interactions between oligomers

In this section we highlight some further results on the interactions between species and resources
and among oligemers, respectively.

Figure 4-figure supplement 1 shows the average pair-wise interactions between 10 random
strings (as representative of the whole population) during the different evolutionary cycles ($t =
0, 9, 18, 24$), $\langle \Delta G_{pp} \rangle$. Such averages have been computed by giving to NUPACK a pool of 1000 se-
quences, randomly subdivided in 100 groups of 10 sequences each. We can see that, during evo-
lution, the selected oligomers increase their average interaction strength, a signature that the evo-
lution favors not only strings that have higher overlap with the resource, but also that interact in
some way with the other species.

Figure 4-figure supplement 2 is the 2D probability distribution $p(\omega, x_R)$, computed from experi-
mental data, to simultaneously have a given MCO$= \omega$ and x_R, the rightmost basis of the of the re-
source involved in the formation of such an MCO. Noticeably, we start from a distribution peaked
in $\omega = 4$ and $10 \leq x_R \leq 12$, essentially corresponding to purely random binding. Across cycles, the
probability distribution is deeply modified, with a shift towards the bottom right corner. Along the
horizontal axis this simply reflects the shift in the average $\langle \omega \rangle$ of the population, and also the fact
that winners have large overlaps. Interestingly, we observe a significant shift of x_R towards high
values, meaning the MCO are preferably formed far away from the bead surface, near the free end

₇₅₄ of target strands.

Results of the experimental replica

₇₅₆ In Figure 2-figure supplement 3 and Figure 2-figure supplement 4 show the main outcomes of the
₇₅₇ replica experiment Oligo2. The evolution of the average MCO $\langle \omega \rangle$ over cycles is qualitatively the
₇₅₈ same of that one found in Oligo1: we have a initial steep grow $\langle \omega \rangle$, compatible with a non linear
₇₅₉ fitness, and then we observe a slower increase for the last cycles, saturating for $\langle \omega \rangle \approx 7$. The
₇₆₀ histogram shows the whole $p(\omega)$ at cycles $t = 0$ and $t = 18$. As for what we found in Oligo1, at $t = 0$
₇₆₁ the behaviour of $p(\omega)$ is compatible with our null random model, while at $t = 18$ the oligomers with
₇₆₂ higher ω are selected, and the tail of $p(\omega)$ is fatter than the one expected by chance.
₇₆₃ These results show that indeed the main outcome of Oligo1 are replicated by Oligo2.

 Appendix 3

 A combinatorial null model with no evolution

As a first step to understand what happens in our experiments, we have built a null model, as-
suming purely random attachment sites, i.e. there is not any search for the relative positions of
binding filaments so to maximize the MCO. In this way we have a null expectation for the typical
MCO distribution in case of oligomers insensitive to the presence of target sequences.

Let us call a the number of consecutive bonded pairs between oligomers 1 (e.g. the species) and
2 (e.g. the resource), with $\max_{r_{1,2}} MCO = \omega$ and $R_{1,2}$ being the relative position of the left ends of
the two filaments, as also explained in *Mambretti et al.* (*2022b*) and in the main text. Consider the
initial population being uniformly random selected (in the high 4^L dimensional space of all possible
DNA strands). The first quantity we are interested to calculate is $\pi(a)$, the *probability density of a*
at time $t = 0$, i.e. before the evolution starts. This distribution is also the distribution of the sub-
sample (i.e, the $\approx 10^6$ strands which we are able to sequence at each cycle), if we assume purely
random attachment (no competition among the consumers, no optimization of the attachment
site).

The relative position of the two strands is important since the target can be either "inside" (Fig-
ure 3-figure supplement 1, panel a)) or surpassing one of the borders (Figure 3-figure supplement
1, panel b) and C)) of the longer oligomer after the attachment. In the picture, $R_{1,2} = 0$ when the
first nucleotide of the target overlaps with the first one of the consumer. From now on, we will
refer to $R_{1,2}$ simply as R.

Our goal consists in finding the MCO of L-long strand (the species) with a l-long target ssDNA
(the resource), where $l < L$. In the following calculations, we therefore assume that R is a random
uniform variable ($-l + 1 \leq r \leq L - 1$) and we will calculate $a|_R$ (for this R).

Such a problem is sketched in Figure 3-figure supplement 2 (where $R = 2$, but the following
holds for any R in the allowed range), where we indicate with d the number of mismatching bases
between the target (red) and the consumer (blue), with u the number of matching ones, being
$d + u := l$. Labeling as x_i the number of consecutive matching nucleotides comprised between
two consecutive mismatching bases, as shown in the picture, the problem can be reformulated in
terms of counting how many integer solutions can be found for the equation:

$$\sum_{i=0}^{d} x_i = u \tag{1}$$

that corresponds to determine the number of ways in which a set of x_i solving Eq. 1 can be
obtained. Note that x_i vanishes between two neighboring pairs of mismatching sites, as marked
by the arrows in the picture. In the case reported in the picture, e.g., $x_i = (1, 0, 1, 4, 0, 4, 0, 0, 2)$ and
$\sum_{i=0}^{8} x_i = 12$, $u = 12$ and $d = 8$.

Ref. *Charalambides* (*2008*) provides us with a formula to answer this question. We stress that
finding the number of solutions $A_{d,u}$ for the above equation is equivalent to obtaining the probabil-
ity distribution $\pi(a)$ for the relative occurrence of each a in a sample of DNA strands, with uniformly
random extracted relative positions of the two sequences.

Theorem 1. *Let us have the equation* $\sum_{i=1}^{n} x_i = u$. *Let us suppose that* $s_i \leq x_i \leq m_i, \forall i = 1, \ldots, n$ *for*
given integers $s_i, m_i, i = 1, \ldots, n$ *with* $s \leq u \leq m$, *being* $s := \sum_{i=1}^{n} s_i$, *and* $m = \sum_{i=1}^{n} m_i$. *Let us define*
$w_i = m_i - s_i \geq 0 \forall i$. *The number of integer solutions, having the shape* $(r_1, \ldots, r_n)$, *of this equation, with*
the restrictions aforementioned, is given by:

$$A_{n,u}(w_1, \ldots, w_n) = \binom{n + u - s - 1}{n - 1} +$$
$$\sum_{r=1}^{n} (-1)^r \sum_{\{i_1, i_2, \ldots, i_r\}} \binom{n + u - s - (\sum_{j=i_1}^{i_r} w_j) - r - 1}{n - 1} \tag{2}$$

where the inner sum is performed over all the r−combinations (having the shape $\{i_1, i_2, \ldots, i_r\}$)

of the n indices $1, \ldots, n$.

In our case the target string may not go beyond the consumer species boundaries (as the latter

is longer than the former). But we can modify the name of some variables to adapt the theorem

to our problem and make some constraints explicit:

- from $i = 1, \ldots, n$ it follows that $n = d + 1$ because we have $d + 1$ terms (from 0 to d).

- $s_i = 0 \, \forall i$, because each $x_i \geq 0$ by definition. Therefore, $s = 0$.

- at least one of the x_i must be equal to a

- in principle, m_i is equal to a for each i (which means that there are no consecutive overlaps

  larger than a).

- it must be that $a \leq u \leq \min((d + 1)a, l)$ (i.e. the total overlap u - which, at most is equal to $d + 1$

  times a - must not be larger than l)

In fact, by using that $s = 0$ and $d + u = l$, the theorem Eq. 2 can be rewritten as:

$$A_{d+1,u}(w_1, \ldots, w_n) = \binom{l}{d} + \sum_{r=1}^{d+1} (-1)^r \sum_{\{i_1, i_2, \ldots, i_r\}} \binom{l - (\sum_{j=i_0}^{i_r} w_j) - r}{d}$$

where the inner sum can be replaced by $\binom{d+1}{r}$ (how many ways are there to dispose the $d+1$ integer indexes to form r groups):

$$A_{d+1,u}(w_1, \ldots, w_n) = \binom{l}{d} + \sum_{r=1}^{d+1} (-1)^r \binom{d+1}{r} \binom{l - (\sum_{j=i_0}^{i_r} w_j) - r}{d}$$

First of all, $\binom{l}{d}$ can be placed inside the summation because it corresponds to the $r = 0$ case. Now, we have to apply the constraint that at least one of the $w_j = a$, which means having the first term where the w_j are always equal to a, minus the second term which includes all the cases where all the $w_j < a$:

$$A_{d+1,u}(w_1, \ldots, w_n) = \sum_{r=0}^{d+1} (-1)^r \binom{d+1}{r} \binom{l - ra - r}{d} -$$

$$\sum_{r=0}^{d+1} (-1)^r \binom{d+1}{r} \binom{l - r(a-1) - r}{d} =$$

$$\sum_{r=0}^{d+1} (-1)^r \binom{d+1}{r} \binom{l - r(a+1)}{d} - \sum_{r=0}^{d+1} (-1)^r \binom{d+1}{r} \binom{l - ra}{d}$$

Note that the previous expressions concern only the cases where $0 \leq R \leq L - l$. However, this

should not matter for practical purposes because R does not appear in the above expressions.

Similar formulas can be found when the target surpasses the left/right border:

if $-l + 1 \leq R \leq -1$

$$A_{d+1,u}(w_1, \ldots, w_n)^{(II)} = \sum_{r=0}^{d+1} (-1)^r \binom{d+1}{r} \binom{l + R - r(a+1)}{d} -$$

$$\sum_{r=0}^{d+1} (-1)^r \binom{d+1}{r} \binom{l + R - ra}{d}$$

while if $L - l \leq R \leq L - 1$

$$A_{d+1,u}(w_1, \ldots, w_n)^{(III)} = \sum_{r=0}^{d+1} (-1)^r \binom{d+1}{r} \binom{L - R - r(a+1)}{d} -$$

$$\sum_{r=0}^{d+1} (-1)^r \binom{d+1}{r} \binom{L - R - ra}{d}$$

Using all the previous results, the general equation for the number of ways which lead to a MCO a is given by:

$$n(a, R) = \sum_{k=1}^{3} n_k(a, R) \tag{3}$$

where the 3 options are originated by the 3 options for the relative position of the target and of the consumer sketched in Figure 3 - figure supplement 1.

Let us now derive $n_k, \forall k$. In particular, if the target "resource" is completely within the "consumer" species boundaries, then we have

$$n_1(a, R) = n_1(a) = (L - l + 1)4^{L-l} \sum_{d=0}^{l} 3^d n_{1,d}(a), \tag{4}$$

with $(L - l + 1)$ is the total number of possible positions R where the consumer can attach to the resource, 4^L is the number of ways to build a L-long string (i.e. all possible species), 3^d represents the number of arrangements for the non-matching bases (3 is because we exclude the only matching base) and $n_{1,d}(a) = A_{d+1,u}^{(I)}$ for $0 \leq R \leq L - l$.

Analogously,

$$n_2(a, R) = \sum_{R=-l+1}^{-1} 4^{L-l-R} \sum_{d=0}^{l+R} 3^d n_{2,d}(a, R), \tag{5}$$

where the summations run over the admitted integer R values as well as over the allowed d values. In this case, 4^{L-l-R} is the number of ways in which the nucleotides not involved in the attachment can be obtained. Here, $n_{2,d}(a, R) = A_{d+1,u}^{(II)}$. A similar equation holds for $n_3(a)$, where

$$\sum_{R=L-l+1}^{L-1} 4^R \sum_{d=0}^{L-R} 3^d n_{3,d}(a, R)$$

and $n_{3,d}(a, R) = A_{d+1,u}^{(III)}$.

To get the distribution of the overlap measure $\pi(a)$, it suffices to divide $n(a, R)$ by the total number of ways in which we can build a string, i.e. 4^L times the number of possible positions R where the attachment can happen, i.e. $L + l - 1$.

$$\pi(a) := \frac{n(a, R)}{\sum_{j=0}^{l} n(j, R)} = \frac{n(a, R)}{4^L (L + l - 1)} \tag{6}$$

$\pi(a)$ is exactly the probability density to find a given MCO= a between a l-long and a L-long sequence which found themselves attached in a random position R.

The underlying probability distribution $\pi(a)$ is naturally discretized and can be represented as an histogram, with the occupancy probability for each bin a (the affinity to the resource). $\pi(a)$ is represented in Figure 3-figure supplement 3a as orange bars; grey bars in the same panel represent the average of 50 independent simulations with $K = 10^6$ 50-mers uniformly generated, where their overlap with the target strand - once uniformly random extracted their relative position - is measured.

The analytical result for the null model and simulations yield almost exactly the same results, with the difference that for large MCO we have strong sampling effect in the simulations. In order to understand how possible biases in the initial frequency of each nucleotide in the starting population, we run $N = 50$ independent simulations with a non-uniform frequency for A,C,G,T in the sequences. For instance, the panel **b)** of Figure 3-figure supplement 3 compares the $\pi(a)$ obtained with A,C,G,T having a probability of 25% (grey data) with the corresponding distribution obtained having: A=23%, C=19%, G=29%, T=29%. The average bin occupancy does not change significantly, but only the large a tails are slightly affected by the bias and the error-bars are much larger for the majority of the a values.

We furthermore set a threshold T to model the physical constraint that if $a < T$, the two ssDNAs cannot bind. We thus obtain a **null model physically constrained** distribution $\pi'(a)$, where $\pi'(a) =$

0 if $a < T$, while is the same as before (after a proper normalization) for $a \geq T$. The new analytical distribution $\pi'(a)$ reads as

$$\pi'(a) = \frac{\pi(a) \times \frac{1}{2}[\text{sgn}(a - T) + 1]}{\sum_{j=T}^{l} \pi(j)},$$ (7)

where $\text{sgn}(0) := 0$. The corresponding pseudo-algorithm to generate random number from such distribution is

- sample a random number z from $\pi(a)$
- if $z < T$, sample another number from $\pi(a)$
- else, save it in the new distribution $\pi'(a)$

The comparison between analytical and numerical results are reported in Figure 3-figure supplement 4a and Figure 3-figure supplement 4b (linear and logarithmic scale on y axis, respectively). The cyan bars correspond to the analytical distribution for the null model with physical constraint ($\pi'(a)$), whereas the purple bars represented the average over $N = 200$ independent simulations of $\pi'(a)$ performed via Metropolis Monte Carlo accept-reject technique *Metropolis et al.* (*1953*). A custom Python script which simulates this process is available at the GitHub repository associated to this work. The null model that we present in the main text is the physical constraint one, i.e. $\pi'(a)$.