Distinguishing cophylogenetic signal from phylogenetic congruence clarifies the

interplay between evolutionary history and species interactions

Benoît Perez-Lamarque [1,2][*] and Hélène Morlon [1]

[1] Institut de biologie de l'École normale supérieure (IBENS), École normale supérieure, CNRS,

INSERM, Université PSL, 46 rue d'Ulm, 75 005 Paris, France

[2] Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum national d'histoire

naturelle, CNRS, Sorbonne Université, EPHE, UA, CP39, 57 rue Cuvier 75 005 Paris, France

[*] corresponding author: benoit.perez@ens.psl.eu; ORCID: 0000-0001-7112-7197

Abstract:

Interspecific interactions, including host-symbiont associations, can profoundly

affect the evolution of the interacting species. Given the phylogenies of host and

symbiont clades and knowledge of which host species interact with which symbiont,

two questions are often asked: "Do closely related hosts interact with closely related

symbionts?" and "Do host and symbiont phylogenies mirror one another?". These

questions are intertwined and can even collapse under specific situations, such that

they are often confused one with the other. However, in most situations, a positive

answer to the first question, hereafter referred to as "cophylogenetic signal", does not

imply a close match between the host and symbiont phylogenies. It suggests only that

past evolutionary history has contributed to shaping present-day interactions, which

can arise, for example, through present-day trait matching, or from a single ancient

vicariance event that increases the probability that closely related species overlap

geographically. A positive answer to the second, referred to as "phylogenetic

congruence", is more restrictive as it suggests a close match between the two

phylogenies, which may happen, for example, if symbiont diversification tracks host

diversification or if the diversifications of the two clades were subject to the same

succession of vicariance events.

Here we apply a set of methods (ParaFit, PACo, and eMPRess), which

significance is often interpreted as evidence for phylogenetic congruence, to

simulations under three biologically realistic scenarios of trait matching, a single

ancient vicariance event, and phylogenetic tracking. The latter is the only scenario that

generates phylogenetic congruence, whereas the first two generate a cophylogenetic

signal in the absence of phylogenetic congruence. We find that tests of global-fit

methods (ParaFit and PACo) are significant under the three scenarios, whereas tests

of event-based methods (eMPRess) are only significant under the scenario of

phylogenetic tracking. Therefore, significant results from global-fit methods should be

interpreted in terms of cophylogenetic signal and not phylogenetic congruence; such

significant results can arise under scenarios when hosts and symbionts had

independent evolutionary histories. Conversely, significant results from event-based

methods suggest a strong form of dependency between hosts and symbionts

evolutionary histories. Clarifying the patterns detected by different cophylogenetic

methods is key to understanding how interspecific interactions shape and are shaped

by evolution.

Keywords: cophylogeny, codiversification, phylogenetic methods, symbiosis,

parasitism, coevolution.

Introduction:

Antagonistic or mutualistic interactions, such as parasitism, herbivory, seed dispersal, or pollination, are key components of ecological communities (Bascompte and Jordano 2013; Mittelbach and McGill 2019). Patterns of interactions (i.e. who interacts with whom) are shaped by evolutionary history through a variety of processes. For example, interspecific interactions may be constrained to species with matching traits, as frequently observed in pollination or host-parasite interaction networks (Muchhala and Thomson 2009; Morand et al. 2015), in which case, species evolutionary history matters as soon as the traits involved in the interactions are evolutionarily conserved. Interspecific interactions may also be constrained by historical contingencies such as past geographic events: if lineages have been geographically isolated following a vicariance event, these lineages do not interact simply because they do not co-occur (Althoff et al. 2014; Perez-Lamarque et al. 2022a). Interspecific interactions can also be transmitted from generation to generation on evolutionary time scales, as in the case of symbionts that are vertically transmitted from parental to descendant hosts (Bright and Bulgheresi 2010). Reciprocally, interactions affect the evolution of the interacting species, for instance through disruptive selection, stabilizing selection, or coevolution. Over macroevolutionary scales, such effects can leave an imprint on the phylogenetic trees of the interacting clades (Harmon et al. 2019; Hembry and Weber 2020; Hayward et al. 2021). An extreme case corresponds to phylogenetic tracking, which can happen, for example, when host speciation events lead to the subsequent speciation of symbionts that are closely associated with them (Fahrenholz 1912). Analyzing interacting species through the lens of their past evolutionary history is therefore fundamental for understanding how interspecific interactions shape and are shaped by evolution.

Given host-symbiont cophylogenetic data, *i.e.* a phylogenetic tree for both the host and the symbiont clades and knowledge of which host species interact with which

symbiont, two patterns are often investigated (Box 1): (i) whether closely-related hosts interact with closely-related symbionts, hereafter referred to as "cophylogenetic signal", and (ii) whether host and symbiont phylogenies mirror one another, hereafter referred to as "phylogenetic congruence" (Blasco-Costa et al. 2021). By definition, phylogenetic congruence can occur only when the number of host and symbiont species is similar, with mostly 'one-to-one' interactions. In this case, patterns of cophylogenetic signal and phylogenetic congruence tend to collapse, such that they can be studied interchangeably. In this context, cophylogenetic methods have been developed to test for phylogenetic congruence using linear algebra technics, such as the fourth-corner statistics (ParaFit, Legendre et al. 2002) and procrustean superposition (PACo, Balbuena et al. 2013). Numerous cophylogenetic systems observed in nature, however, are characterized by many-to-many interactions (Bascompte and Jordano 2013; Perez-Lamarque et al. 2022b), and the same methods have been regularly applied to such systems, and interpreted as tests of phylogenetic congruence (see *e.g.* Fuzessy et al. 2022; Suetsugu et al. 2022; Suzuki et al. 2022 for recent examples). If, under these situations, these methods in fact do not measure phylogenetic congruence, this can lead to a misinterpretation of the results.

Patterns of cophylogenetic signal and phylogenetic congruence can reflect different processes (Fig. 1). The occurrence of cophylogenetic signal simply suggests that past evolutionary history has contributed to shaping present-day interactions. For instance, a cophylogenetic signal can arise if interactions are shaped by evolutionarily conserved traits (Fig. 1a "trait matching") or by past biogeographic events (Fig. 1b "vicariance"), even if the diversification of the host and symbiont clades was not influenced by the interactions between host and symbiont species. The occurrence of phylogenetic congruence, on the other hand, suggests a potential (co-)dependency between the host and symbiont diversifications, causing both phylogenetic trees to look alike. For instance, phylogenetic congruence can emerge if the diversification of vertically transmitted symbionts tracks host diversification (Fig. 1c – "phylogenetic

tracking"), if symbionts diversify by preferentially switching to closely-related host species ("pseudo-codiversification"), or if the diversifications of the two clades were subject to the same succession of vicariance events (de Vienne et al. 2013; Althoff et al. 2014). Phylogenetic congruence implies cophylogenetic signal, but the reverse is not true.

Here, in an effort to clarify the conclusions that can be drawn from various cophylogenetic methods, we analyze their outputs under various evolutionary scenarios, using simulations. Cophylogenetic methods can be divided into two main types of approaches (see de Vienne et al. (2013) and Dismukes et al. (2022) for reviews of these methods): the global-fit methods mentioned above, such as ParaFit and PACo, and event-based methods, such as TreeMap (Page 1994) or eMPRess (Santichaivekin et al. 2021). Event-based methods try to reconciliate the host and symbiont phylogenies by fitting a set of reconciliation events (e.g. cospeciation, host transfer, intra-host duplication, or symbiont loss) to the symbiont phylogeny. We have shown before, in a different context, that global-fit and event-based methods can sometimes output contrasting results (Perez-Lamarque and Morlon 2022), suggesting that they measure different patterns. We evaluate here their outputs under three biologically realistic scenarios of host-symbiont evolution by trait matching, vicariance, and phylogenetic tracking. The latter is the only scenario that generates phylogenetic congruence, whereas the first two generate a cophylogenetic signal in the absence of phylogenetic congruence. We find that global-fit methods provide a test of cophylogenetic signal rather than phylogenetic congruence, whereas event-based methods provide a test of phylogenetic congruence, and discuss the implications of these results for the study of cophylogenetic systems.

Simulations of three biologically realistic scenarios:

In the first simulated scenario, we assumed that present-day host-symbiont interactions are more likely between species having complementary traits following a trait-matching expression with a unidimensional continuous trait (Fig. 1a – "trait matching"; Supplementary Methods 1). We independently simulated two phylogenetic trees for the host and symbiont clades using a birth-death model and on each tree, we then independently simulated the evolution of traits modulating present-day interactions. By using a Brownian motion for trait evolution, closely-related hosts (resp. symbionts) tend to have similar trait values (*i.e.* phylogenetic signal in species traits). Finally, we assumed that the degree of specialization of the symbionts (*i.e.* the number of hosts that a given symbiont interacts with) follows a Poisson distribution and attributed the present-day host-symbiont interacting pairs following a trait-matching expression. As a result, closely related host species interact with closely related symbiont species (*i.e.* cophylogenetic signal), although the phylogenetic trees were simulated independently and are therefore not congruent. In the second simulated scenario, we assumed that host and symbiont species interact at random as long as they occupy the same biogeographic area (Fig. 1b – "vicariance"). At first, all hosts and symbionts simultaneously occupy a single area and diversify independently, until a vicariance event splits the area into three separate areas. Half of the host and symbiont species are then isolated in separate areas; the other half experiences allopatric speciation as their population is split into two. Following the vicariance event, each host and symbiont lineage diversifies independently in its area, resulting in a phylogenetic signal in biogeographic repartition (*i.e.* closely related species tend to occupy the same area). Although the host and symbiont diversifications are not independent (they both experienced a burst of speciation events at the time of vicariance), the host and symbiont phylogenetic trees are not congruent as they experienced different events of diversification before and after the vicariance (Fig. 1b). Finally, we assumed that the degree of specialization of the symbionts follows a

Poisson distribution and randomly attributed host-symbiont present-day interactions within each area. This scenario thus generates cophylogenetic signal but no phylogenetic congruence. In the third simulated scenario, we assumed that the symbiont diversification tracks the host diversification: symbionts species are vertically transmitted over long-time scales among hosts lineages and cospeciate at host speciation events resulting in codiversification (Fig. 1c – "phylogenetic tracking"). In addition, we assumed that symbiont lineages experience host transfers and intra-host duplication, and that host lineages can lose their symbionts. The latter processes dampen but maintain the phylogenetic congruence between host and symbiont phylogenies, rendering it more realistic. This third scenario generates both cophylogenetic signal and phylogenetic congruence. For each scenario, we performed 1,000 simulations with different host and symbiont species richness values (Supplementary Fig. S1). As expected, our simulations under phylogenetic tracking produce systems with a much higher proportion of one-to-one interactions than simulations under trait matching and vicariance (Supplementary Fig. S2).

Outputs of cophylogenetic methods:

For each simulated cophylogenetic data, we first applied the global-fit methods ParaFit and PACo (Ref; Supplementary Methods 2). We measured the strength of the cophylogenetic signal using the "ParaFit global statistic" in ParaFit, whereas, in PACo, we used $R^2 = 1 - m^2$, where m^2 is the sum of the squared residuals of the symmetric procrustean superposition (Blasco-Costa et al. 2021). R^2 is comprised between 0 and 1, and R^2 close to 0 (resp. 1) indicates low (resp. high) cophylogenetic signal. Following Legendre et al. (2002) and Hutchinson et al. (2017), the significance of the ParaFit and PACo tests was evaluated by independently shuffling which host species are associated with each symbiont species (hereafter referred to as "null model 1"). For ParaFit, we found significant tests in 39% of the trait-matching simulations, 52% of the vicariance simulations, and 99% of the phylogenetic tracking (Table 1). Significant tests

for PACo were even more frequent, reaching 50%, 68%, and 100% of the simulations

in the three scenarios, respectively (Table 1). Qualitatively similar results were

obtained when evaluating the significance by shuffling the host species labels

(hereafter referred to as "null model 2"; Supplementary Table S1). Global-fit methods

thus tend to measure cophylogenetic signal in general rather than specifically

phylogenetic congruence. Corroborating this result, we found that, under scenarios of

trait matching or vicariance, global-fit tests are more likely to be significant when there

are frequent many-to-many host-symbiont interactions (Supplementary Table S2;

Supplementary Figs. S3 & S4; generalized linear models (GLM): p-values<0.05), while

we would expect the contrary if these tests measured phylogenetic congruence. As

expected, global-fit tests were more often significant in simulations with more

interacting species (Supplementary Table S3, Supplementary Figs. S3 & S4).

Second, we applied the event-based method eMPRess to each simulation.

eMPRess reconciles the host and symbiont tree topologies by fitting, using maximum

parsimony, events of host transfers, duplications, and losses, each event being

associated with a given cost. We chose eMPRess over all the other existing event-based

methods, as it can be automatically and rapidly run using the command line. As with

most event-based methods, eMPRess does not handle symbiont species that interact

with multiple hosts (Dismukes et al. 2022). However, symbionts often interact with

several hosts in nature, and also in our simulated scenarios of trait matching and

vicariance. We thus tested two strategies for running eMPRess: (i) subsampling one

host at random per symbiont species (Su et al. 2022) or (ii) randomly generating

bifurcating sub-trees for symbionts with multiple hosts, such that each tip in the

symbiont tree is associated with a single host (Satler et al. 2019). A reconciliation is

considered significant if its total cost is lower than 95% of the costs of reconciliations

obtained after shuffling the host species labels ("null model 2"). Phylogenetic trees are

considered congruent if the reconciliation estimates less host transfer than

cospeciation events (Groussin et al. 2017; Perez-Lamarque and Morlon 2022). When

subsampling one host per symbiont species, the eMPRess reconciliation was significant in only 5% of the trait-matching simulations and 16% of the vicariance simulations, and congruent in none (Table 1). When generating random bifurcations in the symbiont tree, it was significant in 34% and 80% of the trait-matching and vicariance simulations, respectively, but congruent in none (Table 1). Indeed, eMPRess estimated on average 5 times more host transfers than cospeciations for both scenarios of trait matching and vicariance (Table 1; Supplementary Fig. S5). In contrast, when simulating phylogenetic tracking, eMPRess gave significant reconciliations in 100% of the simulations, 92% of which were congruent (Table 1). Results were qualitatively similar when choosing alternative event costs (Supplementary Tables S4, S5, & S6). Among all existing event-based methods, we only tried eMPRess for computational reasons. Yet, many of these other methods, e.g. TreeMap or Jane (Page 1994; Conow et al. 2010), are also cost-based and use maximum parsimony, so we suspect that they would behave as eMPRess. More sophisticated methods, such as the version of the amalgamated likelihood estimation (ALE) that considers branching orders in addition to tree topology (Szöllősi et al. 2013), may perform even better than eMPRess as it would provide only time-consistent reconciliations (Maestri et al. 2023). Overall, our findings suggest that event-based methods can specifically detect patterns of phylogenetic congruence and distinguish them from a "simple" cophylogenetic signal.

We investigated whether the effect size of the global-fit methods, *i.e.* values of the ParaFit global statistic and PACo's R^2, can be used as indicators of phylogenetic congruence (Blasco-Costa et al. 2021). The ParaFit global statistic varies with the total number of species, and is thus difficult to interpret in itself; it cannot be used to distinguish phylogenetic congruence from cophylogenetic signal (Fig. 2; Supplementary Fig. S3). In contrast, for PACo, we found that R^2 is generally higher than 0.25 when there is a pattern of phylogenetic congruence (in 99% of the scenario of phylogenetic tracking), whereas it is lower than 0.5 when there is a pattern of cophylogenetic signal alone (in 99% of the scenarios of trait-matching or vicariance;

Fig. 2). Hence, a significant test with $R^2 > 0.50$ is strong support for phylogenetic congruence (although not definite evidence, it may be possible to obtain higher R^2 values under scenarios without congruence with different simulation choices), whereas a significant test with $R^2 < 0.25$ suggests that the system presents a cophylogenetic signal without phylogenetic congruence. A significant test with R^2 values between 0.25 and 0.50 are harder to interpret, as R^2 tends to correlate positively with the ratio of one-to-one interactions and R^2 values above 0.25 are sometimes reported in simulations without phylogenetic congruence (Supplementary Fig. S4). PACo alone is therefore often not sufficient to identify a pattern of phylogenetic congruence.

Concluding remarks:

Using simulations, we have assessed the ability of different cophylogenetic methods to distinguish phylogenetic congruence from cophylogenetic signal. The distinction is important, as these patterns are indicative of different processes, phylogenetic congruence revealing more intricate evolutionary histories between hosts and symbionts than cophylogenetic signal (Fig. 3). We have shown that global-fit methods typically return significant results as soon as there is cophylogenetic signal in species interactions, including in the absence of phylogenetic congruence, meaning that they cannot distinguish the two patterns. In contrast, event-based methods can be used to specifically detect phylogenetic congruence.

Several recent studies have interpreted significant results of global-fit approaches as evidence for phylogenetic congruence and interpreted them as signs of codiversification (*e.g.* Fuzessy et al. 2022; Suetsugu et al. 2022; Suzuki et al. 2022). Our findings suggest that there is evidence for cophylogenetic signal in the studied systems, which is already insightful in itself, but that little can be said about phylogenetic congruence and codiversification before event-based methods are

applied. Applying event-based methods could drastically change the biological conclusions that have been drawn.

Our results suggest that the complementary use of PACo (fast and easy to run) and eMPRess (more computationally intensive but more informative) can be most useful for analyzing cophylogenetic data, with a careful interpretation of the results (Fig. 3). One could begin by using PACo; if the test is not significant, there is neither cophylogenetic signal nor phylogenetic congruence in the data, suggesting independent evolution. If the PACo test is significant but with a low R^2 ($R^2 < 0.25$), there is a cophylogenetic signal but no congruence, suggesting that evolutionary history played a role in shaping present-day interactions, but that there was not a strong co-dependency during the evolutionary history of the two clades. If the PACo test is significant with a higher R^2 ($R^2 > 0.25$), there is a cophylogenetic signal, and potentially also phylogenetic congruence, but the latter needs to be validated using eMPRess, as $R^2 > 0.25$ also frequently occur in systems without phylogenetic congruence. Before running eMPRess, if some symbiont species interact with several host species, we recommend randomly sampling one host species per symbiont (rather than, for example, generating random bifurcations in the symbiont tree, which seems to foster false positives). If the eMPRess test supports phylogenetic congruence (significant reconciliation with a larger number of cospeciation events compared with the number of host transfers), this suggests a strong co-dependency during the evolutionary history of the two clades, such as phylogenetic tracking, successive vicariance events, or pseudo-codiversification.

An advantage of global-fit methods is that their hypothesis testing is rather flexible. One can easily imagine designing more constrained randomization strategies, for example, to specifically test the influence of biogeography or trait matching on the observed cophylogenetic signal (Perez-Lamarque and Morlon 2022). In parallel, data augmentation and other machine-learning technics may allow overriding the

computational bottleneck that limits the implementation of more complex process-
based models. Altogether, such advancements will facilitate linking patterns to
processes in cophylogenetic systems.

To conclude, our results imply that using global-fit methods alone is not
sufficient to robustly assess a pattern of phylogenetic congruence. In a given
cophylogenetic system, finding both significant global-fit tests and significant
reconciliations using event-based approaches suggests that there is a pattern of
phylogenetic congruence that can be linked to various processes such as phylogenetic
tracking, successive vicariance events, or pseudo-codiversification. In contrast, finding
significant global-fit tests but no significant reconciliations with event-based
approaches suggests that phylogenetic congruence is unlikely; the cophylogenetic
signal in this system may rather emerge from processes such as trait matching or
vicariance, but not from phylogenetic tracking or pseudo-codiversification. Clearly
distinguishing patterns of cophylogenetic signal and phylogenetic congruence and
carefully interpreting outputs of cophylogenetic methods are key if we are to
understand the processes that shape present-day communities.

Acknowledgments:

This work was performed using HPC resources from GENCI-IDRIS (Grants 2021-
A0100312405 and 2022-AD010313735).

Code availability:

All the codes for generating the simulations and performing the analyses are available
through the following link:

https://github.com/BPerezLamarque/Scripts/tree/master/Cophylogenetic_signal.

Conflicts of interests:

The authors declare no conflict of interests.

Author contributions:

BPL and HM designed the study, BPL performed the analyses, and both authors wrote

the manuscript.

References:

Althoff D.M., Segraves K.A., Johnson M.T.J. 2014. Testing for coevolutionary

diversification: Linking pattern with process. Trends Ecol. Evol. 29:82–89.

Balbuena J.A., Míguez-Lozano R., Blasco-Costa I. 2013. PACo: A novel procrustes

application to cophylogenetic analysis. PLoS One. 8:e61048.

Bascompte J., Jordano P. 2013. Mutualistic networks. Princeton University Press.

Blasco-Costa I., Hayward A., Poulin R., Balbuena J.A. 2021. Next-generation

cophylogeny: unravelling eco-evolutionary processes. Trends Ecol. Evol. 36:907–

918.

Bright M., Bulgheresi S. 2010. A complex journey: transmission of microbial

symbionts. Nat Rev Microbiol. 8:218–230.

Conow C., Fielder D., Ovadia Y., Libeskind-Hadas R. 2010. Jane: A new tool for the

cophylogeny reconstruction problem. Algorithms Mol. Biol. 5:1–10.

Dismukes W., Braga M.P., Hembry D.H., Heath T.A., Landis M.J. 2022.

Cophylogenetic methods to untangle the evolutionary history of ecological

interactions. Annu. Rev. Ecol. Evol. Syst. 53:1–24.

Fahrenholz H. 1912. Ectoparasiten und abstammungslehre. Zool. Anz. 41:371–374.

Fuzessy L., Silveira F.A.O., Culot L., Jordano P., Verdú M. 2022. Phylogenetic

congruence between Neotropical primates and plants is driven by frugivory.

Ecol. Lett. 25:320–329.

Groussin M., Mazel F., Sanders J.G., Smillie C.S., Lavergne S., Thuiller W., Alm E.J.

2017. Unraveling the processes shaping mammalian gut microbiomes over

evolutionary time. Nat. Commun. 8:14319.

Harmon L.J., Andreazzi C.S., Débarre F., Drury J., Goldberg E.E., Martins A.B.,

Melián C.J., Narwani A., Nuismer S.L., Pennell M.W., Rudman S.M., Seehausen

O., Silvestro D., Weber M., Matthews B. 2019. Detecting the macroevolutionary

signal of species interactions. J. Evol. Biol. 32:769–782.

Hayward A., Poulin R., Nakagawa S. 2021. A broadscale analysis of host-symbiont

cophylogeny reveals the drivers of phylogenetic congruence. Ecol. Lett. 24:1681–

1696.

Hembry D.H., Weber M.G. 2020. Ecological interactions and macroevolution: A new

field with old roots. Annu. Rev. Ecol. Evol. Syst. 51:215–243.

Hutchinson M.C., Cagua E.F., Balbuena J.A., Stouffer D.B., Poisot T. 2017. paco:

implementing Procrustean Approach to Cophylogeny in R. Methods Ecol. Evol.

8:932–940.

Legendre P., Desdevises Y., Bazin E. 2002. A statistical test for host-parasite

coevolution. Syst. Biol. 51:217–234.

Maestri R., Perez-Lamarque B., Zhukova A., Morlon H. 2023. Recent evolutionary

origin and localized diversity hotspots of mammalian coronaviruses.

bioRxiv.:2023.03.09.531875.

Mittelbach G.G., McGill B.J. 2019. Community Ecology. Oxford University Press.

Morand S., Krasnov B.R., Littlewood D.T.J. 2015. Parasite diversity and

diversification. Cambridge University Press.

Muchhala N., Thomson J.D. 2009. Going to great lengths: Selection for long corolla

tubes in an extremely specialized bat-flower mutualism. Proc. R. Soc. B Biol. Sci.

276:2147–2152.

Page R.D.M. 1994. Parallel phylogenies: Reconstructing the history of host-parasite

assemblages. Cladistics. 10:155–173.

Perez-Lamarque B., Krehenwinkel H., Gillespie R.G., Morlon H. 2022a. Limited

evidence for microbial transmission in the phylosymbiosis between Hawaiian

spiders and their microbiota. mSystems. 7:e01104-21.

Perez-Lamarque B., Maliet O., Pichon B., Selosse M.-A., Martos F., Morlon H. 2022b.

Do closely related species interact with similar partners? Testing for

phylogenetic signal in bipartite interaction networks. Peer Community J. 2:e59.

Perez-Lamarque B., Morlon H. 2022. Comparing different computational approaches

for detecting long-term vertical transmission in host-associated microbiota. Mol.

Ecol.

Santichaivekin S., Yang Q., Liu J., Mawhorter R., Jiang J., Wesley T., Wu Y.C.,

Libeskind-Hadas R. 2021. EMPRess: A systematic cophylogeny reconciliation

tool. Bioinformatics. 37:2481–2482.

Satler J.D., Herre E.A., Jandér K.C., Eaton D.A.R., Machado C.A., Heath T.A., Nason

J.D. 2019. Inferring processes of coevolutionary diversification in a community

of Panamanian strangler figs and associated pollinating wasps. Evolution (N. Y).

73:2295–2311.

Su Z.-H., Sasaki A., Kusumi J., Chou P.-A., Tzeng H.-Y., Li H.-Q., Yu H. 2022.

Pollinator sharing, copollination, and speciation by host shifting among six

closely related dioecious fig species. Commun. Biol. 5:284.

Suetsugu K., Okada H., Hirota S.K., Suyama Y. 2022. Evolutionary history of

mycorrhizal associations between Japanese Oxygyne (Thismiaceae) species and

Glomeraceae fungi. New Phytol.

Suzuki T.A., Fitzstevens J.L., Schmidt V.T., Enav H., Huus K.E., Ngwese M.M.,

Grießhammer A., Pfleiderer A., Adegbite B.R., Zinsou J.F., Esen M., Velavan

T.P., Adegnika A.A., Song L.H., Spector T.D., Muehlbauer A.L., Marchi N., Kang

H., Maier L., Blekhman R., Ségurel L., Ko G.P., Youngblut N.D., Kremsner P.,

Ley R.E. 2022. Codiversification of gut microbiota with humans. Science (80-.).

377:1328–1332.

Szöllősi G.J., Rosikiewicz W., Boussau B., Tannier E., Daubin V., Szöllosi G.J.,

Rosikiewicz W., Boussau B., Tannier E., Daubin V. 2013. Efficient exploration of

the space of reconciled gene trees. Syst. Biol. 62:901–912.

de Vienne D.M., Refrégier G., López-Villavicencio M., Tellier A., Hood M.E., Giraud

T. 2013. Cospeciation vs host-shift speciation: Methods for testing, evidence from

natural associations and relation to coevolution. New Phytol. 198:347–385.

Box 1: Definitions

Host-symbiont bipartite interaction network: A network representing which symbiont species interact with which host, here encoded by a matrix with host species in rows and symbiont species in columns, and 1/0 representing the presence/absence of interaction, respectively.

Codiversification/Co-cladogenesis: Pattern of concomitant diversification events happening in both host and symbiont clades. Codiversification can occur due to processes of phylogenetic tracking or successive vicariance events affecting both clades.

Coevolution: Process of reciprocal evolutionary changes induced by reciprocal selective pressures in two (or more) interacting lineages.

Cophylogenetic signal: Pattern depicting the tendency of closely related species to interact with closely related partners.

Cophylogenetics: Study of the link between the host and symbiont evolutionary histories and extant interactions.

Cospeciation: Concomitant event of host and symbiont speciations.

Degree of specialization: Number of species interacting with a given focal species.

Event-based methods: Cophylogenetic methods reconciling the host and symbiont phylogenies by fitting reconciliation events (e.g. cospeciation, host transfer, duplication, or loss) on the symbiont phylogeny.

Global-fit methods: Cophylogenetic methods evaluating the presence of cophylogenetic signal in a bipartite network.

Intra-host duplications: Process of symbiont speciation within a host lineage.

Host transfer (syn. host switch, host shift): Process of transmission of a symbiont from a donor host to a receiver host.

Phylogenetic congruence: Pattern of high similarity of the phylogenetic trees of interacting host and symbiont clades in terms of topology and relative branch

---

lengths. If host and symbiont divergence times are matching, phylogenetic congruence can correspond to codiversification.

Phylogenetic signal: Pattern depicting the tendency of closely related species to have similar traits.

Phylogenetic tracking: Process of host (resp. symbiont) speciations generating subsequent symbiont (resp. host) speciations (e.g. because of vertical transmission that isolates symbiont populations between the two daughter host lineages). Alternatively, symbionts can experience events of host transfers, intra-host duplication, or loss. Phylogenetic tracking usually generates a pattern of codiversification.

Preferential host switching: Tendency of symbionts to experience host transfers toward closely related host species; when the transfer results in a speciation event in the symbiont lineage, this tends to generate phylogenetic congruence. This does not imply co-diversification though, as the divergence times of the symbionts may be much more recent than those of the hosts. We specifically refer to this resulting pattern as **pseudo-codiversification.**

Symbiont loss: Extinction of a symbiont in a given host lineage.

Tree topology: Branching pattern of the different nodes of a phylogenetic tree that does not consider branch lengths (*i.e.* evolutionary time).

Vertical transmission: Process of symbiont inheritance from host generation to host generation (at the level of individual or at the level of the whole lineage). Over long timescales, it can generate phylogenetic tracking.

Vicariance: Process of biogeographic changes leading to the isolation of populations and subsequent allopatric speciation. By simultaneously affecting hosts and symbionts, vicariance can lead to cophylogenetic signal, and potentially phylogenetic congruence if events of vicariance occur repetitively.

Table 1: ParaFit and PACo (global-fit methods) are often significant in all simulated scenarios, including scenarios of trait matching and vicariance that do not generate phylogenetic congruence, whereas eMPRess (an event-based method) supports phylogenetic congruence only under phylogenetic tracking which generates phylogenetic congruence:

The table indicates the percentages of simulations for which ParaFit, PACo, or eMPRess output a significant test, under three scenarios (Fig. 1): (a) present-day interactions dictated by trait matching, (b) present-day interactions at random following a single vicariance event, or (c) present-day interactions resulting from phylogenetic tracking. For eMPRess, we report the percentage of significant reconciliations based on permutations alone (P) or based on selecting among these the reconciliations that have more cospeciation than host transfer events (P+C). We consider eMPRess to support phylogenetic congruence when conditions P and C are met (in bold). Results by clade size are given in Supplementary Tables S1-S6. eMPRess results correspond to host-symbiont reconciliations ran with relative costs d=4, t=1, and l=1 for duplications, host transfers, and losses, respectively; other costs combinations provided qualitatively similar results – see Supplementary Tables S4-S6).

Methods		(a) Trait matching	(b) Vicariance	(c) Phylogenetic tracking
Global-fit methods	Percentage of significant tests using ParaFit	39%	52%	99%
	Percentage of significant tests using PACo	50%	68%	100%
Event-based methods	Percentage of significant tests using eMPRess with one host per symbiont	P: 5% **P+C: 0%**	P: 16% **P+C: 0%**	P: 100% **P+C: 92%**
	Percentage of significant tests using eMPRess with random bifurcations	P: 34% **P+C: 0%**	P: 80% **P+C: 0%**	

Figure 1: Three mock examples of host-symbiont systems (here represented as gophers and lice) that generate a cophylogenetic signal. Only the third example also generates phylogenetic congruence.

(a) Trait matching. Here host and symbiont phylogenetic trees evolve independently. Traits also evolve independently on each phylogeny following Brownian motion processes, which results in a phylogenetic signal in species traits (i.e. closely related species tend to have similar trait values, see the color gradient). Host-symbiont interactions at present are more likely between species having complementary traits, following a trait-matching expression. Because of trait matching, closely related host species interact with closely related symbiont species (*i.e.* cophylogenetic signal), although the phylogenetic trees are independent (*i.e.* no phylogenetic congruence).

(b) Vicariance: Here hosts and symbionts interact at random as long as they occupy the same biogeographic area. They occupy a single area until an event of vicariance splits this area into three separate areas, isolating populations and leading to speciation (a species occupying two areas at the time of vicariance immediately experiences allopatric speciation). Following the vicariance event, each host and symbiont lineage radiates independently on its area, without migration, which results in a phylogenetic signal in biogeographic repartition (i.e. closely related species tend live on the same area). Because interactions happen at random within each area, this scenario generates cophylogenetic signal without phylogenetic congruence.

(c) Phylogenetic tracking: Here symbiont species are vertically transmitted over long-time scales along host lineages and cospeciate at host speciation events resulting in codiversification. In addition, symbiont lineages can experience horizontal host transfers (with replacement of the previous symbiont lineage), intra-host duplication, and host lineages can lose their symbionts. This scenario generates both cophylogenetic signal and phylogenetic congruence.

(a) Trait matching

(c) Phylogenetic tracking

Figure 2: eMPRess correctly distinguishes patterns of phylogenetic congruence

(scenario of phylogenetic tracking) from cophylogenetic signal alone (scenarios of

trait matching and vicariance), whereas ParaFit cannot and PACo can only in some

cases:

Distribution of ParaFit global statistics (a), PACo's R^2 (b), and the ratio between the

number of host transfers and the number of cospeciations in eMPRess reconciliations

(c) as a function of the simulation scenario and the test significance. Here, we only

reported eMPRess reconciliations obtained when subsampling one host per symbiont

species, but similar results were observed when using random bifurcations

(Supplementary Fig. S5).

Figure 3: Patterns of cophylogenetic signal and phylogenetic congruence can be generated by various processes:

Event-based and global-fit methods differently measure these patterns: Event-based methods can robustly identify phylogenetic congruence, whereas global-fit methods measure cophylogenetic signal. Some of the statistics of global-fit methods (e.g. the R^2 of PACo) can inform whether the cophylogenetic signal may be due to phylogenetic congruence: a low R^2 ($R^2<0.25$) indicates that there is a cophylogenetic signal but no phylogenetic congruence, whereas a higher R^2 ($R^2>0.25$) suggests there is a cophylogenetic signal and potentially also phylogenetic congruence, but the latter needs to be validated using eMPRess, as $R^2>0.25$ also frequently occur in systems without phylogenetic congruence (Figure 2).