Exploring Liquid-Liquid Phase Separation in the organization of Golgi Matrix Proteins

Luis Felipe S. Mendes,[1,*,#] Carolina G. Oliveira,[2,#] Emanuel Kava,[2] Antonio J. Costa-Filho[2,*]

[1] Group of Biophysics and Structural Biology "Sergio Mascarenhas". São Carlos Institute of Physics, University of São Paulo, São Carlos-SP, Brazil

[2] Molecular Biophysics Laboratory, Department of Physics, Faculty of Philosophy, Sciences, and Letters at Ribeirão Preto, University of São Paulo, Ribeirão Preto-SP, Brazil

\# These authors contributed equally

To whom correspondence should be addressed: luis.felipe.mendes@usp.br; ajcosta@usp.br

Abstract

The Golgi apparatus is a critical organelle in protein sorting and lipid metabolism. Characterized by its stacked, flattened cisternal structure, the Golgi exhibits distinct polarity with its *cis-* and *trans-*faces orchestrating various protein maturation and transport processes. At the heart of its structural integrity and organization are the Golgi Matrix Proteins (GMPs), predominantly comprising Golgins and GRASPs. These proteins contribute to this organelle's unique stacked and polarized structure and ensure the precise localization of Golgi-resident enzymes, which is crucial for accurate protein processing. Despite over a century of research since its discovery, the Golgi architecture's intricate mechanisms still need to be fully understood. Here, we demonstrate that GMPs present a significant tendency to form biocondensates through Liquid-Liquid Phase Separation (LLPS) across different Eukaryotic lineages. Moreover, we validated experimentally that members of the GRASP family also exhibit a strong tendency for LLPS. Our findings offer a new perspective on the possible roles of protein disorder and LLPS of GMPs in the Golgi organization.

Keywords:

Liquid-Liquid Phase Separation, Golgi Complex, Golgins, GRASPs, Intrinsically Disordered Proteins

Introduction

Eukaryotic cells evolved to possess efficient secretory machinery capable of transporting a significant fraction of their proteome (1). Proteins entering the secretory pathway are efficiently sorted to a specific destination: the extracellular space, the plasma membrane, or the endomembrane system. The machinery also has efficient quality control, avoiding the transport of misfolded proteins or complexes (2). Proteins destined for this classical secretory pathway are synthesised in the endoplasmic reticulum (ER) interior, where they are first folded and possibly modified by the post-translation ER machinery (2). Subsequently, these secretory cargoes are transferred to the ER exit sites (ERES), where they are assembled into secretory vesicles coated with coatomers (COPII), routed for the ER-Golgi intermediate compartment (ERGIC), and later to the *cis*-Golgi face (3, 4).

The Golgi apparatus, a central organelle in eukaryotic cells, plays a pivotal role in trafficking, processing, and sorting proteins and lipids. This membrane-bound organelle exhibits a unique architecture consisting of a stack of flattened cisternae interconnected by tubular-vesicular networks. The cisternae can be further classified into three distinct regions: *cis*, *medial*, and *trans*, each responsible for specific functions in protein processing and glycosylation. The maintenance of this intricate structure is crucial for the Golgi's function and relies on a diverse group of proteins, collectively known as the Golgi matrix proteins (GMPs). In the organisation of the Golgi stacks and cisternae formation, it was observed that in cells treated with brefeldin A (an inhibitor of COPI-mediated transport from the Golgi to the ER), the Golgi membrane and its integral proteins relocated to the ER (5). On the other hand, the GMPs accumulated in the cytoplasm formed a ribbon-like reticulum in the correct Golgi location (5). Based on this observation, it seems that GMPs could generate the higher-order architecture of the Golgi by a mechanism independent of other Golgi proteins. Although an orchestrated action involving all the GMPs provides the total adhesive energy dictating Golgi cisternal stacking (6), how these proteins act in the general organisation/dynamics remains unknown.

The Golgins and the Golgi Reassembly and Stacking Proteins (GRASP) are the main protein families composing the GMPs. The Golgins are predominantly extended coiled-coil proteins predicted to reach an extension between 100-600 nm (7, 8). They attach to the Golgi cisternae through either their carboxy terminus, directly interacting with the membrane surface (7) or a membrane-associated partner (9). Golgins participate in vesicular traffic, Golgi maintenance and dynamics, and the positioning of the Golgi apparatus within cells (8, 10). On the other hand, GRASPs constitute a family of peripheral membrane proteins essential for general Golgi organisation into the ribbon-like structure (9) and Golgi integrity for the correct protein sorting and glycosylation along the secretory pathway (11). GRASPs are observed in all branches of the eukaryotic tree of life, although not equally spread in all supergroups (12).

GRASPs are collapsed intrinsically disordered proteins (IDPs) with high promiscuity for mediating protein-protein interactions with several different protein partners (13-15). Their unusual structural plasticity (16) might be the converging feature of GRASPs in multiple cell functional tasks, including their direct participation in Golgi dynamics during mitotic and apoptotic times (17-19) and their self-association during stress conditions triggering unconventional protein secretion pathways (20-23). GRASPs interact with the membrane via a lipidation of its Gly2 using a myristoyl-switch mechanism (24, 25). They also anchor some Golgin representatives, including GM130, Golgin45, and p115 in mammalian cells and the Golgin BUG1 in Yeast (9, 26). This anchoring is essential for the correct Golgi organisation (9, 27).

In 2019, James Rothman suggested that the Golgi, akin to the nucleolus, would fundamentally be liquid with a phase-separated internal organisation (28). This suggestion came after discovering that many human Golgins can efficiently populate a liquid-like droplet state once they are overexpressed *in vivo* and dispersed in the cytosol (29). In recent years, liquid-liquid phase separation (LLPS) was discovered to form and organize biomolecular condensates within cells. LLPS is a process by which a homogenous solution of macromolecules separates into distinct liquid phases, often leading to the assembly of membrane-less compartments with unique biophysical properties (30). This phenomenon has been implicated in cellular functions, such as signal transduction, RNA processing, and stress response (31-33).

Here, we build on the discussion of the role of LLPS in regulating cellular processes by expanding its potential to describe several characteristics of the Golgi apparatus, especially its dynamic nature and unusual flattened membrane organisation. It is important to note that the Golgi is a membrane-bound organelle, setting it apart from membrane-less organelles like the nucleolus. Furthermore, any LLPS at the Golgi involving Golgi-matrix proteins would need to occur on a membrane surface. The remodelling of cell membranes requires their bending. Membrane bending is an essential cellular process with implications in numerous aspects of cell biology, such as membrane traffic, cell motility, organelle creation, and cell division (34-36). Besides the traditional bending induced by well-shaped proteins, it has been recently shown that protein phase separation occurring on the surfaces of both synthetic and cell-derived membrane vesicles induces significant compressive stress within the membrane plane, which prompts the membrane to bend inward, forming protein-lined membrane tubules (35, 37). In this case, the proteins undergoing phase separation contain substantial regions of intrinsic disorder, which do not possess a stable three-dimensional structure (37, 38). Fascinatingly, disordered regions have recently been observed to form networks sustained by weak, multivalent contacts, facilitating the assembly of protein liquid phases on membrane surfaces (39, 40).

Since LLPS can affect membrane bending dramatically (35, 39, 41) and some Golgins
have been shown to undergo LLPS in cells (29), it would be reasonable to argue that there might
be a direct correlation between the LLPS-propensity of GMPs and the unusual Golgi cisternae
shape. However, it is unclear how the LLPS propensity spread among the GMPs and how this is
conserved in the remaining Eukaria tree. Here, we highlight the significant prevalence of
intrinsically disordered regions within GMPs across Eukarya and how these regions are linked
to a high propensity for those proteins to form liquid-like condensates via LLPS. We also
describe a detailed experimental study of LLPS of human GRASP55 and GRASP65 to confirm
some of our predictions.

Results and Discussions

Despite the commonly accepted presence of coiled-coil regions within Golgins, this still
lacks experimental validation. There has also been a limited investigation into the prevalence of
structural disorder in these proteins. Coiled-coil regions are frequently predicted as disordered
sequences at rates similar to genuine intrinsically disordered segments identified in DisProt (42,
43). Although coiled coils often transition from a disordered to an ordered state upon assembly,
they hardly ever evolve into highly compact structures. Their extended configuration and the
presence of extensive multivalent surfaces possibly underlie the observed propensity of specific
coiled-coil-containing proteins to undergo LLPS in cells (29, 44, 45). This behaviour resembles
the multivalency characteristic of disordered proteins, enabling many of them to engage in
numerous weak interactions concurrently. Such a feature significantly enhances their likelihood
of undergoing LLPS. Despite these insights, a thorough large-scale investigation remains
lacking, particularly one that evaluates GMP disorder and quantifies amino acid residues
conducive to more compact structures.

We started by exploring the structural properties of the metazoan GMPs. Identifying all
GMPs in metazoans is an active research area, and the comprehensive list has not been
definitively established. An extra difficulty is that different organisms can express different sets
of Golgins. We selected some well-known GMPs and associated proteins that are known to play
a role in maintaining Golgi structure and function. It is important to emphasise that we might
not have exhausted the full list of GMPs. Some GMPs commonly identified in metazoans are:
GM130 (GOLGA2), Golgin-45 (BLZF1), Golgin-160 (GOLGA3), AKAP9 (CG-NAP), Giantin
(GOLGB1), GCP60 (ACBD3), GCC88 (GCC1), Golgin-97, Golgin-245 (GOLGA4),
SCYL1BP1 (GORAB), GCC185 (GCC2), Bicaudal-D (BCD1 and BCD2), NEEC1
(JAKMIP2), GRASP65 (GORASP1) and GRASP55 (GORASP2). These families form the list
to be discussed now in the text. Other important GMPs and Golgi structure-associated proteins

like CASP (GOLGA8), Golgin-84 (GOLGA5), p115 (also known as USO1), Zinc Finger Protein-Like 1 (ZFPL1), and GMAP-210 (also known as TRIP11) are also found in metazoans but have a broader presence in the Eukary tree, and so will be discussed later in the text in a broader eukaryotic perspective. It is important to note that different proteins can have multiple names or synonyms, adding to the complexity of cataloguing them all.

Not all the proteins listed in the previous paragraph are Golgins. Golgins are typically predicted to have coiled-coil regions; many have a C-terminal GRIP domain or a similar structure. In the previous list, proteins like GRASP65 and GRASP55, while important to the function and structure of the Golgi, are not classified as Golgins. The same is true for the ZFPL1. The distinction between Golgins and other Golgi-associated proteins is not always clear, and there may be exceptions or debates over how specific proteins are classified. The pervasive presence of Golgins with their long rod-like structures establishes an extensive ribosome-free zone around the Golgi apparatus, a feature readily observable in microscopy images (46).

Our comprehensive analysis spanned >28,000 different protein sequences and employed metrics to shed light on the proteins' inherent tendency to adopt extended configurations (out of classical collapsed ones), consequently exposing their multivalent properties. Such multivalence is pivotal for LLPS and a central theme of our investigation. We started utilising the Hydrophobicity versus Net Charge analysis to address this issue. Initially proposed by Uversky, this method provides a binary classification of proteins (47). The method builds a clear boundary (red line in Figure S1) separating "natively disordered proteins" from "ordered proteins". Predominantly used to differentiate extended IDPs from globular proteins, this analytical approach unveiled a critical trait: most GMPs exhibited behaviours akin to extended IDPs (Figure 1). This finding underscores the conservation of sequence features within GMPs, including low net hydrophobicity and high net charge.

Notable exceptions to this trend are GRASPs and Golgin-45, which appear to reside more frequently on the "well-folded" side of the Uversky plot (Figure S1). GRASPs consist of two primary regions: the structured GRASP domain and a fully disordered C-terminus, the latter typically being the larger portion of those proteins (15). Previous molecular biophysics analyses indicated GRASPs as molten-globule proteins, suggesting an intermediate level of structural order (16, 21). Golgin-45 is the primary binding partner of mammalian GRASP55 (13, 48). Unlike GRASP, which is predicted to be present in the Last Eukaryotic Common Ancestor (LECA), Golgin-45 appears to be holozoan-specific (49). Prior research demonstrated that the progenitor of the paralogous proteins GRASP55 and GRASP65 could bind Golgin-45 with high affinity (13). In this context, GRASPs and Golgin-45 neither exhibit the mean net

charge/hydrophobicity ratio characteristic of extended IDPs, like other GMPs, nor fit in the
classic well-folded realm of protein structure.

To expand the previous observations, we explored the relative amount of IDRs in the
protein sequences to get more quantitative details of the disordered behaviour. Drawing from
the rich tapestry of protein families, we singled out two distinct families for comparison with
our database. On one end of the spectrum, we considered the mammalian Albumins—
representative of structurally well-ordered proteins—and on the opposite end, we included the
RNA binding protein FUS, a classic representative of the extended IDPs found in vertebrates.
The median of disorder fractions varied substantially from around 30% of the total protein
sequence for Golgin45 and NEEC to nearly 70% for SCYL1BP1 (Figure 1A). Our data showed
that most families were composed of a median ranging from 40-60%, explaining why these
proteins behave more similarly to extended IDPs in the Uversky plot analyses.

Figure 1: Analyses of Amino Acid Composition and LLPS-Propensity Across Metazoan GMPs. (A)
Representation of residue fractions exhibiting a score exceeding the 0.5 thresholds as per IUPred2 long
disorder predictions. (B) Enumeration of order-promoting residue fractions ('I', 'C', 'L', 'V', 'W', 'Y', 'F') as
defined by reference (50). (C) Application of the machine-learning-based PSPredictor (51) for LLPS
prediction. A protein sequence is categorised as phase separating if the PsPred Score exceeds 0.5. (D)
Assessment of LLPS propensity utilising the catGRANULE algorithm, where propensity scores are
evaluated in the context of proteome distribution, as conveyed through a cumulative distribution function.
A higher score corresponds to a higher propensity for a protein to phase separate. As dosage-sensitive

proteins are known to score approximately 0.5 (as per reference (52)), we designate this score as our
threshold in the analyses.

To deepen our understanding of the structural propensities of the GMPs, we analysed
their overall composition of order/disorder-promoting residues. In the realm of protein structure,
the concepts of "disorder-promoting residues" and "order-promoting residues" refer to the
propensities of specific amino acids to promote either a structured (ordered) or unstructured
(disordered) conformation of the protein chain. The results observed for the GMPs tested were
between the extreme situations observed for order-promoting residues in FUS (~15%) and
Albumin (~33%), (Figure S1), again suggesting a high degree of structural flexibility. GMPs
harboured fewer regions of order and exhibited higher flexibility (Figure 1B), suggesting that
their function is likely linked to these structural characteristics. For example, GMPs tend to
adopt extended structures, which give rise to the extensive ribosome-exclusion area surrounding
the Golgi in microscopy images (46, 53).

IDPs and IDRs are unique segments of proteins that lack a fixed or stable three-
dimensional structure under physiological conditions *in vitro*. They are associated with many
cellular functions, including molecular recognition, molecular chaperones, protein scaffolding,
flexible linkers, among others (54, 55). The flexibility and diverse binding potential of
IDPs/IDRs are often associated with an increased propensity for liquid-liquid phase separation
(LLPS) (40, 51). However, this does not necessarily imply a direct correlation. LLPS is like oil
droplets forming in water: specific proteins or protein complexes separate from the surrounding
solution to form concentrated, liquid-like droplets within the cell, and this phenomenon is now
recognised to be widely spread *in vivo* (56). Predicting the propensity of a protein or a protein
region to undergo LLPS is a vibrant area of investigation in current bioinformatics and
biophysical research. Several computational tools are available to predict LLPS propensity, with
many focusing on the presence of IDRs and sequence-based features like amino acid
composition, charge, and hydrophobicity (51, 52). These include tools like PSPred and
catGRANULE (51, 52).

PSPred is a machine-learning, sequence-based tool that queries and predicts potential
phase separation proteins. Its output is represented by the "PSP Score", which ranges from 0 to
1, with 1 indicating the highest propensity for LLPS (51). In contrast, the catGRANULE
algorithm delivers a protein-level score as a cumulative distribution (52). It is noteworthy that
catGRANULE does not prescribe a default cutoff, but based on how the model is trained, any
score greater than 0 already suggests that a protein is predisposed to phase separation (52).
Nevertheless, we have opted for a cutoff value of 0.5, which corresponds to the median in the
distribution of scores determined in the positive training set. We utilised these two analytical

tools to assess our GMP database, focusing on evaluating the propensity of LLPS and its

conservation within specific protein families. We only excluded the FUS family data from the

catGRANULE output analysis due to its distinct inclination towards a phase-separated state. Its

median value exceeds 4, skewing the graph and complicating the comparison with other protein

families (Figure S2).

Previously, metazoan-specific Golgins, such as GM130, Golgin-97, Golgin-245,

GCC88, and GCC185, were observed to undergo phase separation when overexpressed *in cell*

(29, 45). Interestingly, Golgin-97 and GCC88 families did not demonstrate a strong propensity

towards LLPS (Figures 1C and D). Similarly, Bicaudal-D, NEEC, and Golgin45 lacked such

strong propensity (Figures 1C and D). To delve deeper into these protein families with lower

propensities, we tested human protein representatives using the PhaSePred Algorithm. This tool

is a meta-predictor that aggregates the prediction scores of several LLPS-associated predictors

(57). Notably, the human Golgin-97 protein (Uniprot Q92805) returned a Feature PS-Part score

(a metric related to proteins whose phase separation behaviours are regulated by protein or

nucleic acid partner components) exceeding 75%. This score points to a strong, albeit partner-

dependent, propensity for LLPS. Similar patterns were detected for GCC88 (Uniprot Q96CN9),

Bicaudal-D (Uniprot Q96G01), and Golgin-45 (Uniprot Q9H2G9), which produced PS-Part

scores of approximately 64%, 77%, and 52% respectively. In contrast, NEEC (Uniprot

Q96AA8) generated a markedly lower score, at around 32%.

Therefore, all GM protein families tested (except for NEEC) are suggested to phase-

separate spontaneously or in a partner/context-dependent process. This prevalence was

particularly notable, considering the unique evolutionary paths of these proteins across Eukaria

and their low primary sequence conservation. The significant conservation of this

physicochemical property stood out, underscoring its potentially crucial role in those proteins'

functions and interactions. Since LLPS appears inherent to metazoan GMPs, it is tempting to

speculate that this property is important to the proteins' functionality. As previously discussed,

one of the principal roles of GMPs lies in organising and preserving the Golgi apparatus.

However, characteristics such as flattened cisternae and stacked Golgi are not unique to

metazoans.

Furthermore, considering the evolutionary pathways and conservation of certain GMPs,

there is a compelling argument that Golgi stacking may have originated in the common

eukaryotic ancestor. In a study grounded on phylogenetics, Barlow and colleagues identified

several proteins implicated in the Golgi structure, which were inferred to be present in the Last

Eukaryotic Common Ancestor (LECA). These protein families, which share roles in delineating

separate Golgi structures, include GMAP, GRASP, TMF, Golgin-84, CASP, p115, GRIP-
containing protein, and zinc finger protein-like 1 (ZFPL1) (49).

282   Therefore, we expanded our earlier analyses to encompass a database of these "ancient"
GMPs, where we collected a sum of different modern orthologs from all possible branches of
the Eukarya tree. This extension aimed to investigate whether the high LLPS tendency observed
in metazoans is also prevalent across Eukarya. Our analyses confirm a high prevalence of
protein regions with a high propensity to be intrinsically disordered, with medians ranging from
~20% (p115) to ~60% (TMF) of the entire protein sequence (Figure 2A). Interestingly, even
other properties like pI are highly conserved across the eukarya GMPs predicted to be present in
LECA (Figure 2B). In alignment with our previous findings, most GMPs also exhibited a high
propensity for spontaneous LLPS, including the families of GRASP, GMAP-21, TMF, Golgin-
84, and GRIP-CP (Figures 2C and 2D). The exceptions were ZFPL1, p115, and CASP, with
scores as low as those observed in our control dataset of Albumins.

Figure 2: Analyses of the LLPS propensity and IDR content in Golgi structure proteins inferred to be
present in the LECA according to (49). A) Representation of residue fractions exhibiting scores exceeding
the 0.5 thresholds, using IUPred2 long disorder predictions. B) pI distribution across the GMP database.
The pI values were calculated using an in-house Python script described in the Material and Methods
Section. C) Application of the machine-learning-based PSPredictor (51). for LLPS prediction. A protein
sequence is categorised as phase separating if the PsPred Score exceeds 0.5. D) Assessment of LLPS
propensity utilising the catGRANULE algorithm, where propensity scores are evaluated in the context of
proteome distribution, as conveyed through a cumulative distribution function. A higher score
corresponds to a higher propensity for a protein to phase separate. As dosage-sensitive proteins are known
to score approximately 0.5 (as per reference (52)), we designate this score as our threshold in the
analyses.

Building on the detailed examination of GMPs' theoretical propensity for LLPS, we
now turn to experimental evidence to support our findings. To explore this phenomenon, we
examined the physicochemical behaviour of human GRASPs in various solutions under
crowded situations (58), aiming to understand and characterise the potential LLPS phenomena
(Figure 3). Utilizing optical microscopy, we observed the formation of protein-rich droplets of
GRASP55 (Figure 3A), and this tendency was both protein, ionic strength and crowding agent
concentration-dependent (Figure S3). We initially focused on the impact of pH on the droplet
formation. Considering the intracellular environment, we selected a pH of 5.5 to mimic extreme
cellular conditions (59) and pH values of 6.0 and 6.3 to represent the values observed in *medial*
and *trans* regions of the Golgi apparatus, respectively (60). Lastly, we included a physiological
pH of 7.4 for comparative analysis. This approach allowed us to discern how varying pH
conditions influence the behaviour of GRASP., contributing to our understanding of their role
within the Golgi apparatus and potential implications for cellular function. Interestingly, despite
GRASP65 being a vertebrate paralogue of GRASP55 (61), our experiments did not reveal any
LLPS under the tested conditions (Figure 3B). The underlying factors contributing to this
absence of LLPS in GRASP65 are not immediately apparent and warrant further investigation,
which will be the subject of future investigation.

Our experiments revealed that GRASP55 underwent LLPS, although to a different
extent, in all examined pH environments (Figure 3A). Turbidity assays showed that the most
pronounced LLPS activity of GRASP55 occurred at pH levels corresponding to its native
positioning within the *medial* (pH 6.0) and *trans* (pH 6.3) cisternae of the Golgi apparatus
(Figure 3C) (62). This observation suggests a strong tendency for GRASP55 to engage in LLPS,
potentially contributing to the structural and functional self-organisation within the Golgi
complex.

Figure 3: pH-Dependent Propensity of GRASPs for LLPS. (A) Differential Interference Contrast (DIC) microscopy images of GRASP55 (5 μM), highlighting the physical structure and morphology of the protein. The second row features fluorescence microscopy images of the same GRASP55 protein stained with Alexa Fluor 555 dye. (B) Similar to item (A) but using GRASP65. (C) Turbidity Analysis of GRASP55 phase separation at various pH levels, demonstrating pH-sensitive behaviour. (D) GRASP55 droplets dissolved after addition of 10% (w/V) 1,6 Hexanediol, monitored by DIC microscopy. The images selected are representative ones. Scale bars of 5 μm for (A) and (D), and of 20 μm for (B).

We then established a pH of 6.3 for subsequent tests because it reflects the environment
in the *medial* part of the Golgi apparatus, where GRASP55 demonstrated the most effective
droplet formation. To further investigate the underlying forces driving LLPS in GRASP55, we
conducted experiments using NaCl and 1,6-Hexanediol across a gradient range of 15 – 550 mM
for NaCl and 0% - 15% for Hexanediol (Figure 3D and S3). The NaCl-containing samples
demonstrated increased droplet formation with higher salt concentrations (Figure S3). In
contrast, adding 10% Hexanediol led to the dissociation of the condensates after 10 minutes of
incubation (Figure 3D). This result implies that the weak hydrophobic interactions disrupted by
Hexanediol (63) are crucial in stabilising the GRASP55 droplets.

Besides examining the dependency of GRASP55 propensity to LLPS in a crowded
environment, we also explored additional properties of the formed droplets, particularly their
physical state – whether they were more liquid-like or gelatinous. We conducted permeability
experiments using SYPRO™ Orange Protein Gel Stain to assess this. This stain permeates
droplets if they are sufficiently liquid-like to allow for molecular exchange with their
surrounding environment. In the initial part of our experiment, we prepared GRASP55 samples
in LLPS-conditions and incubated them for 20 minutes to facilitate droplet formation.
Subsequently, we checked the samples under a microscope to confirm the presence of droplets.
Following this, we added SYPRO Orange to the solution at a concentration of 10% of the stock
solution and incubated it for an additional 10 minutes to allow the SYPRO Orange to permeate
the droplets. After 10 minutes of incubation, the samples were examined under a fluorescence
microscope. Our observations revealed that the droplets exhibited liquid characteristics, as
evidenced by the penetration of SYPRO Orange and subsequent fluorescence (Figure 4A). This
liquid characteristic suggests that even after forming the biomolecular condensate, where the
protein is at high concentrations, it can still exchange molecules with its surrounding
environment. Another liquid-like observation in GRASP55 droplets was the occurrence of
fusion or coalescence, as seen in Figure 4B, where individual protein droplets merged to form a
single, larger droplet.

Figure 4: Liquid-like properties of GRASP55 droplets. (A) DIC Microscopy of GRASP55 (concentration of 5 µM). The image captures the dynamic process of droplets merging over time, illustrating the fluid nature of these structures. (B) The GRASP55 droplets (concentration: 5 µM) are observed under DIC microscopy. The image highlights the interaction of these droplets with the surrounding medium, where SYPRO Orange dye is dissolved. The permeation of the SYPRO Orange into the droplets is evident, indicating the liquid-like properties of the droplets and their ability to exchange substances with their environment—scale bars: 5 μm.

Conclusion:

In conclusion, our study provides a novel perspective on the nature of GMPs and critical functional characteristics across Eukarya, focusing on metazoans. Our data distinctly highlight the high prevalence of intrinsic disorder within GMPs, a feature that appears to be conserved across diverse eukaryotic lineages, albeit with different levels of prevalence and extent. Interestingly, the disorder in GMPs seems to be central to the potential of these proteins for LLPS, a phenomenon widely observed in the intracellular environment and critical for various biological processes. The combined findings indicate that disorder and LLPS propensity, despite the low primary sequence conservation across GMP families, have been evolutionarily conserved, suggesting their critical roles in the proteins' functionality. To substantiate our

findings, we conducted experimental tests on the potential for LLPS in a previously unstudied
family of GMPs, namely GRASPs. These experiments confirmed their capacity for LLPS,
thereby reinforcing our hypothesis. This study highlights that the functional versatility and
adaptability of GMPs are likely derived from their structural disorder and/or elongated,
multivalent properties. This intrinsic characteristic enables them to engage in diverse
interactions and respond to various cellular signals, which is crucial for the organisation and
maintenance of the Golgi apparatus. Our findings open the way for future studies on the role of
LLPS in GMPs and their implication in various cellular processes. It will be particularly
interesting to delve further into the precise mechanisms by which these proteins influence the
architecture of the Golgi apparatus and the regulatory role of phase separation in these
processes. The evolutionary conservation of these traits among GMPs suggests their
fundamental importance in cell biology and invites further exploration.

Material and Methods

Database collection:

In order to collect a broadly representative collection of unique protein orthologue
sequences, we relied on the search for the protein family name in the hierarchical catalogue of
orthologs OrthoDB (64) and for protein families in the InterPro database for those who are not
catalogued in a single orthologue family (65). The protein sequences collected were as follows:
TMF (OrthoDB 2054925at2759, 3717503at2759 and 1221595at2759, N = 1629), CASP
(InterPro IPR012955, N = 2395), Golgin84 (InterPro IPR019177, N = 1500), Mammalian
Serum Albumin-Like (from InterPro IPR020858, N = 1500), GRASP (OrthoDB 36330at2759,
N = 2395), GM130 (OrthoDB 2880220at2759, N = 887), Golgin45 (OrthoDB 2879260at2759,
N = 830), Golgin160 (OrthoDB 5357451at2759, N = 560), AKAP9 (OrthoDB 158920at33208,
N = 1550), Giantin (OrthoDB 4227816at2759, N = 505), GCP60 (OrthoDB 38315at33208, N =
1384), GCC88 (OrthoDB 237371at33208, N = 831), Golgin97 (OrthoDB 42510at33208, N =
832), FUS RNA binding protein (OrthoDB 484351at7742, N = 982), Golgin245 (OrthoDB
2529099at33208, N = 871), protein bicaudal D homolog 2-like (OrthoDB 5350043at2759, N =
1886)), SCYL1BP1 (OrthoDB 2486431at33208, N = 708), p115 (OrthoDB 1774at2759, N =
1927), GCC185 (OrthoDB 814692at33208, N = 807), GRIP-Containing Proteins (N = 1500
randomly selected from InterPro IPR000237), GMAP (OrthoDB Group 5138860at2759, N =
886), ZFPL1 (OrthoDB Group 25738at2759, N = 903) and Janus kinase and microtubule
interacting protein 2 - NECC (OrthoDB 5403492at2759, N = 1461). We employed the online
resource found at https://fasta.bioch.virginia.edu/fasta_www2/clean_fasta.html (accessed in
2023) to process and clean our dataset's FASTA formatted protein names. For most larger

databases exceeding 1500 sequences, a custom Python script was utilised to randomly truncate the sequence data to the desired size of 1500 (script "fasta_random" in the SI). We have also generated a Python script to filter out unnatural amino acid residues within the protein sequences (script "fasta_filter" in the SI). This script reads the protein sequences from an input file, removes any character that is not a valid amino acid from each sequence, and writes the cleaned sequences to an output file, preserving the headers of the original sequences. This procedure was crucial given that most of the analysis tools we employed lacked the functionality to handle these unconventional letters appropriately. The full protein sequence database is available as a supplementary file.

Prediction of LLPS-Propensity:

The propensity for LLPS was evaluated using the PSPred and catGRANULE web-based platforms. Each algorithm accepts different sequence lengths: PSPred can simultaneously process segments of 100 amino acids, while catGRANULE can handle up to 500 amino acids per run. To streamline the upload process and accommodate these constraints, we developed Python scripts that partition our protein sequence database into appropriately sized segments for submission to each server. This script (script "fasta_split" in the SI) reads a FASTA file, splits the sequences into groups of a specified size, and writes each group to a separate FASTA file. The statistical analyses were performed using the software Origin 2022.

Prediction of IDRs and calculated fractions within the sequence

We quantified each amino acid's intrinsic disorder within a sequence using the IUPred2A tool, specifically designed for assessing long sequences (66). We engineered a Python script to interface with our comprehensive protein sequence databases to streamline the process (script "disorder_fractions" in the SI). This script seamlessly integrated with the IUPred2A software package we sourced directly from the official website. It then automated the computation of disorder scores, effectively isolating the fraction of residues with scores exceeding the threshold of 0.5. The statistical analyses were performed using the software Origin 2022.

Charge vs Hydrophobicity Plot:

We crafted a Python script implementing the Uversky methodology (script "fasta_CHA" in the SI) to systematically evaluate each protein sequence's mean hydrophobicity and net

charge in our database. Our user-friendly script allows customisation of the hydrophobicity
scale per user preference, exploring the potential described in (67). For this manuscript, we have
utilised the Kyte-Doolittle scale, originally employed by Uversky in (47). Our data presentation
shows a discrimination borderline, defined by a specific equation, categorising proteins as either
"natively disordered" or "natively ordered". The equation is $<R> = 2.785 <H> - 1.151$,
according to (47). In this work, $<H>$ and $<R>$ are the mean hydrophobicity and the mean
net charge of the protein, respectively.

Amino acid sequence general properties determination:

We engineered a Python script to efficiently perform multiple tasks to expedite the
protein sequence analysis process. This script reads protein sequences from a FASTA file,
conducts a comprehensive evaluation of various properties using the Biopython package, and
outputs these properties in an organised tabular format. The analysed properties include
fractions of residues promoting order/disorder, isoelectric point (pI), Grand Average of
Hydropathy (GRAVY), and the proportion of polar and charged residues. The statistical
analyses were performed using the software Origin 2022.

Protein expression and purification

GRASP55 and GRASP65 were purified as previously reported (68, 69). The purified
solutions were concentrated with an Amicon Ultra-15 Centrifugal Filter (NMWL of 10 kDa,
Merck Millipore, Burlington, USA). For protein concentration determination, the extinction
coefficient at 280 nm was calculated with the ProtParam web server (70).

For the fluorescence microscopy experiments, protein samples were labelled with Alexa
Fluor™ 555 NHS Ester (Succinimidyl Ester) (ThermoFisher). The dye was diluted in DMSO
(Dimethyl sulfoxide pure) stock solution at a concentration of 10 mM. Before adding the dye,
the protein solutions were exchanged with a 20 mM sodium phosphate buffer
(Na2HPO4/NaH2PO4) at pH 7.4 using a HiTrap® Desalting Column connected to an Äkta
purifier system (GE Healthcare). Subsequently, Alexa 555 dye was added at a concentration
double that of the protein and the mixture was then incubated for approximately one hour at 4°C
with gentle shaking. Post-incubation, excess unbound Alexa 555 was removed using a HiTrap®
Desalting Column connected to an Äkta purifier system (GE Healthcare). Finally, the proportion
of labelled to unlabelled protein was maintained at a 1% ratio for microscopy experiments.

Differential interference contrast (DIC) and fluorescence microscopies

The LLPS and droplet formation by the target proteins were visualised using an inverted

system microscope IX71 equipped with a PicoQuant MT 200 module (PicoQuant®, Germany)

under DIC and fluorescence mode with a 60x objective (NA = 1.2). The fluorescence excitation

intervals were selected using the mirror unities: U-MNU2 BP360/370; Blue: U-MWB2

BP460/490; Green: U-MWIG3 BP530-550. The long-pass filters HQ405LP, HQ460LP, BLP01-

488R, HQ550LP and HQ690/70m were used. The droplet formations were induced by a 10X

dilution method using a stock solution of 10 mg/mL unless otherwise stated. LLPS solutions

were composed of 20 mM Na_2HPO_4/NaH_2PO_4 + X% (W/V) polyethylene glycol (PEG) 8000

(ThermoFisher) adjusted to the desired pH and with X varied from 2-20%. All the samples were

incubated for 20 minutes in RT prior to measurements. The image acquisition was performed by

adding 5uL drop solution in the centre of coverslips 20 x 20 mm (Deckgläser). Liquid droplet

permeability was studied using 1X SYPRO™ Orange Protein Gel Stain (5,000X Concentrate in

DMSO - Thermo Fisher Scientific). Data collection and analysis were performed with the

SymPhoTime 64 software (PicoQuant).

Turbidity assays

The turbidity experiments were conducted by measuring the absorbance at a wavelength

of 350 nm in a Spectrophotometer Multiskan Go (ThermoFisher) equipment using the SkanIT

RE 6.1.1 program. The samples contained only protein without any dye and the corresponding

pH buffer in Na_2HPO_4/NaH_2PO_4 + X% (W/V) polyethylene glycol (PEG) 8000. The

experiments were performed in triplicate with fresh sample preparation.

Acknowledgements

514   The authors thank the *Fundação de Amparo à Pesquisa do Estado de São Paulo* (Grants

515   No. 2022/06006-0, 2021/10465-7 and 2020/15542-7) for the financial support. AJCF thanks

516   Conselho Nacional de Desenvolvimento Científico e Tecnlógico (CNPq) for the partial financial

517   support (Grant no. 306682/2018-4)

Supporting Information

In the supplementary material, we have included Figure S1-S3. Additionally, we have

provided the Python scripts employed for data analysis. Furthermore, an accompanying .zip file

contains the complete database referenced in our manuscript.

Conflict of interest

The authors declare that they have no conflict of interest.

References:

----

1. M. Uhlen *et al.*, Tissue-based map of the human proteome. *Science* **347** (2015).
2. R. Benyair, E. Ron, G. Z. Lederkremer, Protein quality control, retention and degradation
at the endoplasmic reticulum. *International Review of Cell and Molecular Biology, Vol*
292 **292**, 197-280 (2011).
3. J. Dancourt, C. Barlowe, Protein Sorting Receptors in the Early Secretory Pathway.
Annual Review of Biochemistry, Vol 79 **79**, 777-802 (2010).
4. G. Zanetti, K. B. Pahuja, S. Studer, S. Shim, R. Schekman, COPII and the regulation of
protein sorting in mammals. *Nature Cell Biology* **14**, 20-28 (2012).
5. J. Seemann, E. Jokitalo, M. Pypaert, G. Warren, Matrix proteins can generate the higher
order architecture of the Golgi apparatus. *Nature* **407**, 1022-1026 (2000).
6. I. Lee *et al.*, Membrane adhesion dictates Golgi stacking and cisternal morphology.
Proceedings of the National Academy of Sciences of the United States of America **111**,
1849-1854 (2014).
7. T. M. Witkos, M. Lowe, The Golgin Family of Coiled-Coil Tethering Proteins. *Frontiers in*
Cell and Developmental Biology **3** (2016).
8. A. K. Gillingham, S. Munro, Finding the Golgi: Golgin Coiled-Coil Proteins Show the
Way. *Trends in Cell Biology* **26**, 399-408 (2016).
9. Y. J. Zhang, J. Seemann, Rapid degradation of GRASP55 and GRASP65 reveals their
immediate impact on the Golgi structure. *Journal of Cell Biology* **220** (2020).
10. M. Wong, S. Munro, The specificity of vesicle traffic to the Golgi is encoded in the
golgin coiled-coil proteins. *Science* **346**, 601-+ (2014).
11. Y. Xiang *et al.*, Regulation of protein glycosylation and sorting by the Golgi matrix
proteins GRASP55/65. *Nature Communications* **4** (2013).
12. L. F. S. Mendes *et al.*, The GRASP domain in golgi reassembly and stacking proteins:
differences and similarities between lower and higher Eukaryotes. *Febs Journal* **286**,
3340-3358 (2019).
13. L. F. S. Mendes *et al.*, Resurrecting Golgi proteins to grasp Golgi ribbon formation and
self-association under stress. *bioRxiv*, 2021.2007.2027.453980 (2021).
14. L. F. S. Mendes *et al.*, Conformational flexibility of GRASPs and their constituent PDZ
subdomains reveals structural basis of their promiscuous interactome. *Febs Journal*
287, 3255-3272 (2020).
15. L. F. S. Mendes, N. A. Fontana, S. T. Reddy, V. N. Uversky, A. J. Costa, The exquisite
structural biophysics of the Golgi Reassembly and Stacking Proteins. *International*
Journal of Biological Macromolecules **164**, 3632-3644 (2020).
16. L. F. S. Mendes *et al.*, New structural insights into Golgi Reassembly and Stacking
Protein (GRASP) in solution. *Scientific Reports* **6** (2016).
17. I. Ayala, F. Mascanzoni, A. Colanzi, GRASP65-mediated control of the Golgi structure is
crucial for G2/M transition. *Molecular Biology of the Cell* **29** (2018).

18.   D. Tang, H. Yuan, Y. Wang, The Role of GRASP65 in Golgi Cisternal Stacking and Cell
Cycle Progression. *Traffic* **11**, 827-842 (2010).
19.   J. D. Lane *et al.*, Caspase-mediated cleavage of the stacking protein GRASP65 is
required for Golgi fragmentation during apoptosis. *Journal of Cell Biology* **156**, 495-509
(2002).
20.   N. A. Fontana, A. D. Rosse, A. Watts, P. S. R. Coelho, A. J. Costa-Filho, *In vivo* amyloid-
like fibrils produced under stress. *bioRxiv*, 2021.2002.2002.429251 (2021).
21.   N. A. Fontana, R. Fonseca-Maldonado, L. F. S. Mendes, L. P. Meleiro, A. J. Costa, The
yeast GRASP Grh1 displays a high polypeptide backbone mobility along with an
amyloidogenic behavior. *Scientific Reports* **8** (2018).
22.   A. J. Curwin *et al.*, ESCRT-III drives the final stages of CUPS maturation for
unconventional protein secretion. *Elife* **5** (2016).
23.   C. Bruns, J. M. McCaffery, A. J. Curwin, J. M. Duran, V. Malhotra, Biogenesis of a novel
compartment for autophagosome-mediated unconventional protein secretion. *Journal
of Cell Biology* **195**, 979-992 (2011).
24.   E. Kava, L. F. S. Mendes, M. R. B. Batista, A. J. Costa, Myristoylation and its effects on
the human Golgi Reassembly and Stacking Protein 55. *Biophysical Chemistry* **279**
(2021).
25.   E. Kava, C. V. Garbelotti, J. L. S. Lopes, A. J. Costa, Myristoylated GRASP55 dimerizes in
the presence of model membranes. *Journal of Biomolecular Structure & Dynamics*
(2024).
26.   R. Behnia, F. A. Barr, J. J. Flanagan, C. Barlowe, S. Munro, The yeast orthologue of
GRASP65 forms a complex with a coiled-coil protein that contributes to ER to Golgi
traffic. *Journal of Cell Biology* **176**, 255-261 (2007).
27.   R. Grond *et al.*, The function of GORASPs in Golgi apparatus organization in vivo. *J Cell
Biol* **219** (2020).
28.   J. E. Rothman, Jim's View: Is the Golgi stack a phase-separated liquid crystal? *Febs
Letters* **593**, 2701-2705 (2019).
29.   P. Ziltener, A. A. Rebane, M. Graham, A. M. Ernst, J. E. Rothman, The golgin family
exhibits a propensity to form condensates in living cells. *Febs Letters*.
30.   M. Fuxreiter, M. Vendruscolo, Generic nature of the condensed states of proteins.
Nature Cell Biology **23**, 587-594 (2021).
31.   S. Boeynaems *et al.*, Protein Phase Separation: A New Phase in Cell Biology. *Trends in
Cell Biology* **28**, 420-435 (2018).
32.   S. Alberti, D. Dormann, Liquid-Liquid Phase Separation in Disease. *Annual Review of
Genetics, Vol 53* **53**, 171-+ (2019).
33.   Y. Shin, C. P. Brangwynne, Liquid phase condensation in cell physiology and disease.
Science **357** (2017).
34.   H. T. McMahon, J. L. Gallop, Membrane curvature and mechanisms of dynamic cell
membrane remodelling. *Nature* **438**, 590-596 (2005).
35.   S. Mondal, T. Baumgart, Membrane reshaping by protein condensates. *Biochimica Et
Biophysica Acta-Biomembranes* **1865** (2023).
36.   H. T. McMahon, E. Boucrot, Membrane curvature at a glance. *Journal of Cell Science*
128, 1065-1070 (2015).
37.   F. Yuan *et al.*, Membrane bending by protein phase separation. *Proceedings of the
National Academy of Sciences of the United States of America* **118** (2021).
38.   V. N. Uversky, Intrinsically disordered proteins in overcrowded milieu: Membrane-less
organelles, phase separation, and intrinsic disorder. *Current Opinion in Structural
Biology* **44**, 18-30 (2017).
39.   C. Has, P. Sivadas, S. L. Das, Insights into Membrane Curvature Sensing and Membrane
Remodeling by Intrinsically Disordered Proteins and Protein Regions. *Journal of
Membrane Biology* **255**, 237-259 (2022).

40. K. K. Turoverov *et al.*, Stochasticity of Biological Soft Matter: Emerging Concepts in
Intrinsically Disordered Proteins and Biological Phase Separation. *Trends in Biochemical*
Sciences **44**, 716-728 (2019).

41. A. Mangiarotti, N. Chen, Z. Zhao, R. Lipowsky, R. Dimova, Wetting and complex
remodeling of membranes by biomolecular condensates. *Nature Communications* **14**,
2809 (2023).

42. B. Szappanos, D. Suveges, L. Nyitray, A. Perczel, Z. Gaspari, Folded-unfolded cross-
predictions and protein evolution: The case study of coiled-coils. *Febs Letters* **584**,
1623-1627 (2010).

43. Z. Gáspári, Is Five Percent Too Small? Analysis of the Overlaps between Disorder, Coiled
Coil and Collagen Predictions in Complete Proteomes. *Proteomes* **2**, 72-83 (2014).

44. D. A. Ramirez, L. E. Hough, M. R. Shirts, Coiled-coil domains are sufficient to drive
liquid-liquid phase separation of proteins in molecular models. *bioRxiv* (2023).

45. A. A. Rebane *et al.*, Liquid-liquid phase separation of the Golgi matrix protein GM130.
Febs Letters **594**, 1132-1144 (2020).

46. F. S. Sjostrand, V. Hanzon, ULTRASTRUCTURE OF GOLGI APPARATUS OF EXOCRINE CELLS
OF MOUSE PANCREAS. *Experimental Cell Research* **7**, 415-429 (1954).

47. V. N. Uversky, J. R. Gillespie, A. L. Fink, Why are "natively unfolded" proteins
unstructured under physiologic conditions? *Proteins-Structure Function and Genetics*
41, 415-427 (2000).

48. J. Zhao, B. Li, X. Huang, X. Morelli, N. Shi, Structural Basis for the Interaction between
Golgi Reassembly-stacking Protein GRASP55 and Golgin45. *Journal of Biological*
Chemistry **292**, 2956-2965 (2017).

49. L. D. Barlow, E. Nyvltova, M. Aguilar, J. Tachezy, J. B. Dacks, A sophisticated,
differentiated Golgi in the ancestor of eukaryotes. *Bmc Biology* **16** (2018).

50. L. F. S. Mendes *et al.*, New structural insights into Golgi Reassembly and Stacking
Protein (GRASP) in solution. *Scientific Reports* **6** (2016).

51. X. Q. Chu *et al.*, Prediction of liquid-liquid phase separating proteins using machine
learning. *Bmc Bioinformatics* **23** (2022).

52. B. Bolognesi *et al.*, A Concentration-Dependent Liquid Phase Separation Can Cause
Toxicity upon Increased Protein Expression. *Cell Reports* **16**, 222-231 (2016).

53. S. Mogelsvang, N. Gomez-Ospina, J. Soderholm, B. S. Glick, L. A. Staehelin,
Tomographic evidence for continuous turnover of Golgi cisternae in Pichia pastoris.
Molecular Biology of the Cell **14**, 2277-2291 (2003).

54. V. N. Uversky, The protein disorder cycle. *Biophysical Reviews* (2021).

55. Z. R. Liu, Y. Q. Huang, Advantages of proteins being disordered. *Protein Science* **23**, 539-
550 (2014).

56. F. C. Keber, T. Nguyen, C. P. Brangwynne, M. Wühr, Evidence for widespread
cytoplasmic structuring into mesoscopic condensates. *bioRxiv*, 2021.2012.2017.473234
(2021).

57. Z. M. Chen *et al.*, Screening membraneless organelle participants with machine-
learning models that integrate multimodal features. *Proceedings of the National*
Academy of Sciences of the United States of America **119** (2022).

58. S. Alberti, A. Gladfelter, T. Mittag, Considerations and Challenges in Studying Liquid-
Liquid Phase Separation and Biomolecular Condensates. *Cell* **176**, 419-434 (2019).

59. N. Demaurex, pH Homeostasis of cellular organelles. *News Physiol Sci* **17**, 1-5 (2002).

60. S. Kellokumpu, Golgi pH, Ion and Redox Homeostasis: How Much Do They Really
Matter? *Frontiers in Cell and Developmental Biology* **7** (2019).

61. L. F. S. Mendes *et al.*, Resurrecting Golgi proteins to grasp Golgi ribbon formation and
self-association under stress. *International Journal of Biological Macromolecules* **194**,
264-275 (2022).

62.   J. Shorter *et al.*, GRASP55, a second mammalian GRASP molecule involved in the
671   stacking of Golgi cisternae in a cell free system. *Molecular Biology of the Cell* **10**, 115A-
672   115A (1999).

673   63.   M. J. d. Amaral *et al.*, Copper drives prion protein phase separation and modulates
674   aggregation. *bioRxiv*, 2023.2002.2015.528739 (2023).

675   64.   D. Kuznetsov *et al.*, OrthoDB v11: annotation of orthologs in the widest sampling of
676   organismal diversity. *Nucleic Acids Research* (2022).

677   65.   T. Paysan-Lafosse *et al.*, InterPro in 2022. *Nucleic Acids Research* (2022).

678   66.   B. Meszaros, G. Erdos, Z. Dosztanyi, IUPred2A: context-dependent prediction of protein
679   disorder as a function of redox state and protein binding. *Nucleic Acids Research* **46**,
680   W329-W337 (2018).

681   67.   F. Huang *et al.*, Improving protein order-disorder classification using charge-hydropathy
682   plots. *Bmc Bioinformatics* **15** (2014).

683   68.   S. T. Reddy, L. F. S. Mendes, N. A. Fontana, A. J. Costa, Exploring structural aspects of
684   the human Golgi matrix protein GRASP55 in solution. *International Journal of Biological*
685   *Macromolecules* **135**, 481-489 (2019).

686   69.   S. T. Reddy, V. N. Uversky, A. J. Costa, Biophysical characterization of intrinsically
687   disordered human Golgi matrix protein GRASP65. *International Journal of Biological*
688   *Macromolecules* **162**, 1982-1993 (2020).

689   70.   M. R. Wilkins *et al.*, Protein identification and analysis tools in the ExPASy server. *2-D*
690   *Proteome Analysis Protocols* **112**, 531-552 (1999).

691