Cross-transmission of resistant gastrointestinal nematodes between wildlife and transhumant sheep

Camille Beaumelle[a,b,c,1,*], Carole Toïgo[d], Rodolphe Papet[e], Slimania Benabed[a,b], Mathieu Beurier[d], Léa Bordes[f], Anaïs Brignone[d], Nadine Curt-Grand-Gaudin[c], Mathieu Garel[d], Justine Ginot[b], Philippe Jacquiet[f], Christian Miquel[c], Marie-Thérèse Poirel[a,b], Anna Serafino[b], Eric Vannard[g], Gilles Bourgoin[a,b, †], Glenn Yannic[c, †]

[a] *Université de Lyon, Université Lyon 1, CNRS, Laboratoire de Biométrie et Biologie Evolutive UMR 5558, F-69100 Villeurbanne, France*

[b] *Université de Lyon, VetAgro Sup, Campus Vétérinaire de Lyon, F-69280 Marcy l'Etoile, France*

[c] *Université Grenoble Alpes, Université Savoie Mont Blanc, CNRS, LECA, 38000, Grenoble, France*

[d] Office Français de la Biodiversité, Unité Ongulés Sauvages, Gières, France

[e] Parc national des Écrins, Secteur du Champsaur-Valgaudemar, 05260 Saint Jean Saint Nicolas, France

[f] Université de Toulouse, UMT Pilotage de la Santé des Ruminants, Ecole Nationale Vétérinaire de Toulouse, France

[g] Parc national des Écrins, Secteur du Briançonnais, 05100 Briançon, France

*Corresponding author: beaumelle.camille@gmail.com

† Co-senior authors

ABSTRACT

Wild and domestic ungulates can be infected with the same species of gastrointestinal parasitic nematodes. These parasites have a free leaving stage in the environment that contributes to the ease of transmission among different host species. In addition, gastrointestinal nematodes have developed resistance to anthelmintics which is now considered a major problem for the livestock sector. In a context where wild and domestic ungulates share the same pastures, the maintenance and circulation of resistant gastrointestinal nematodes between species have rarely been explored.

In the European Alps, domestic sheep are driven in the high-altitude summer pastures leaving in sympatry with wild ungulates for several months. In this study we investigated the nemabiome of domestic sheep and Alpine ibex, *Capra ibex*, in three different areas of the French Alps to evaluate the parasites circulation between the two host species. The Alpine ibex is a protected mountain ungulate that is phylogenetically related to sheep and hosts nematode species common to sheep.

Using internal transcribed spacer 2 (ITS-2) nemabiome metabarcoding, we found sheep and ibex sharing similar gastrointestinal nematodes, except for a few species, such as *Marshallagia marshalli* and *Trichostrongylus axei*. This suggests that the long-term co-occurrence of sheep and ibex on mountain pastures has promoted the exchange of gastrointestinal nematodes between the two hosts. Based on the sequencing of the isotype 1 of the beta tubulin gene, associated with benzimidazole resistance, we found resistant nematodes in all sheep farms and in all Ibex populations. Our results demonstrated that Ibex can host and shed resistant strains before transhumant sheep arrive on pastures, and then can act as a reservoir or refugia for resistant gastrointestinal nematodes. The relative role of ibex to the nemabiome and in particular to the maintenance and circulation of resistant strains in sheep remain to be determined.

Keywords: ITS-2 rDNA, benzimidazole resistance; isotype-1 β-tubulin; livestock; nemabiome metabarcoding; wild ungulates, transhumant sheep, Alpine ibex

Introduction

Parasites represent a large proportion of animal diversity and are key components of food webs (Hudson et al., 2006) and are essential determinants of the health, fitness, population dynamics and community composition of their hosts (Tompkins et al., 2011). The parasites of the Nematodes class infect a wide range of species worldwide, including animals and plants (Blaxter and Koutsovoulos, 2015). In animals, the gastrointestinal nematode parasites are of major concern for livestock productivity and security as they can impact animal health implying reduced animal productions and economical losses (Charlier et al., 2020; Roeber et al., 2013).

To limit the parasite load and its impact on livestock health, the use of anthelmintics to treat livestock against gastrointestinal nematodes is a common and cost effective practice (Vercruysse et al., 2018). Nonetheless, the repeated use of anthelmintics has led to the selection of anthelmintic-resistant strains of gastrointestinal nematodes. Within the gastrointestinal nematodes, resistances to several families of anthelmintics (e.g., benzimidazole, macrocyclic lactones and levamisole) has been observed, and resistance to several drugs is increasing (Bordes et al., 2020; Kaplan and Vidyashankar, 2012; Rose et al., 2015; Rose Vineer et al., 2020). The first case of resistance to benzimidazole was reported in 1964 in lambs from central Kentucky (Drudge et al., 1964).

Today, resistance to benzimidazoles is widespread throughout the world (Kaplan and Vidyashankar, 2012), and is particularly common on sheep farms in Europe (Papadopoulos et al., 2012; Rose Vineer et al., 2020). The mechanisms of resistance to benzimidazole are well known and documented (Whittaker et al., 2017). The anthelmintic effect of benzimidazoles relies on the fixation of the molecule to the parasite β-tubulin isotype-1, resulting in the disruption of tubulin–microtubule equilibrium (Whittaker et al., 2017). In resistant nematodes, specific mutations of the β-tubulin isotype-1 gene have been correlated with the resistance to benzimidazole in several gastrointestinal nematode species (Charlier et al., 2022). These mutations were associated with alteration of the β-tubulin isotype-1 structure, decreasing the affinity of the protein with the benzimidazole and subsequently, activity of benzimidazole (Whittaker et al., 2017).

Some generalist gastrointestinal nematodes can infect several host species (Walker and Morgan, 2014), including both domestic and wild ungulates (e.g.,

Beaumelle et al., 2022; Cerutti et al., 2010). The transmission of gastrointestinal
nematodes among hosts, even if they do not simultaneously occupy the same
pastures, is possible thanks to their free-living infective larval stage that may rely
active until several months in the environment (Carlsson et al., 2013; Fiel et al., 2012;
Walker and Morgan, 2014). Transmitted parasites can also include gastrointestinal
nematodes resistant to anthelmintics. For instance, benzimidazole-resistant
nematodes have been detected in free-living populations of roe deer living in
sympatry with livestock (Chintoan-Uta et al., 2014; Nagy et al., 2017). To date, the
role of wild ungulates in the epidemiology of resistant nematodes remains to be
determined, but it has been suggested that wildlife may act as a reservoir of resistant
nematodes for livestock (Brown et al., 2022; Chintoan-Uta et al., 2014; Francis and
Šlapeta, 2023; Laca Megyesi et al., 2019; Walker and Morgan, 2014). Yet, we need
to investigate the presence of resistant nematodes between co-grazing wild and
domestic ungulates in different contexts, i.e., for different host species, in different
landscape, and under different climatic conditions, to accurately evaluate the
potential role of wildlife as reservoir for anthelmintic resistant gastrointestinal
nematodes.

Transhumant pastoralism is a common practice in the European Alps and
consists of the seasonal movement of grazing livestock from lowland areas to
mountain meadows in summer which provide fresh pasture for domestic ungulates,
i.e., mainly sheep, but also cows or goats (Biber, 2010). These mountainous areas
are inhabited year-round by wild ungulates, particularly those living at high altitude in
the European Alps, like Alpine ibex (*Capra ibex*), or Northern chamois (*Rupicapra*
rupicapra). While wild ungulates tend to avoid domestic herds spatially or temporarily
during the summer (Acevedo et al., 2008), certain factors may contribute to the use
of the same pastures by both wild and domestic ungulates.

Spatial segregation between wild and domestic ungulates is usually observed
once livestock arrive on pasture (Brivio et al., 2022; Ryser-Degiorgis et al., 2002).
Livestock are generally released onto the best grazing areas during the summer
season due to their high productivity and accessibility of water for animals, and salt
licks provides essential mineral nutrients in relatively nutrient-poor alpine ecosystems
(Chirichella et al., 2014; Richomme et al., 2006; Ryser-Degiorgis et al., 2002).
However, prior to the arrival of livestock, mountain ungulates have been observed
using preferentially the same grazing areas as those used by livestock on their arrival

(Brivio et al., 2022; Ryser-Degiorgis et al., 2002) and ibex, for example, have been observed to return to the sheep grazing area immediately after the sheep have left (Ryser-Degiorgis et al., 2002). The presence of wild and domestic ungulates in attracting zones such as salt licks, even if not simultaneous, offers good opportunities for parasites transmission, and these areas are therefore considered hotspots for parasites infection (Richomme et al., 2006; Ryser-Degiorgis et al., 2002; Utaaker et al., 2023).

Consequently, transhumant pastoralism represents a risk of pathogens transmission between wild and domestic ungulates in mountain areas (Rossi et al., 2019). Pathogen exchanges at the interface of wild and domestic ungulates have already been well documented. The Alpine ibex has been identified as the wildlife reservoir of brucellosis (*Brucella melitensis*) which was transmitted to cattle and human in the Bargy massif in northern French Alps (Marchand et al., 2017). In addition, sheep have been confirmed as the domestic reservoir of the Border disease, which induced a major virus outbreak in Southern chamois (*Rupicapra pyrenaica*) population in the Pyrenees (Luzzago et al., 2016). The transmission of gastrointestinal nematodes has already been described between wild ungulates and transhumant domestic ungulates in mountainous areas (Cerutti et al., 2010; Citterio et al., 2006; Khanyari et al., 2022; Zaffaroni et al., 2000). However, no study has yet investigated the transmission of anthelmintic-resistant nematode in a transhumant pastoral system.

In this study, we investigated the community of gastrointestinal nematodes infecting Alpine ibex and domestic sheep (*Ovis aries*) and the prevalence of resistance to benzimidazole, in three different regions of the French Alps. The Alpine ibex was close to extinction at the beginning of the 19[th] century but the reinforcement of its populations by several reintroductions in different part of the Alps has increased the species' overall abundance and range (Brambilla et al., 2022). Today,the Alpine ibex species is estimated at 52 000 individuals in Europe (Brambilla et al., 2020).

Given that the co-occurrence of sheep and ibex on the same pastures is limited to the summer period and since sheep, and that ibex usually host species-specific gastrointestinal nematodes (Walker and Morgan, 2014), we expect the nemabiome to be highly differentiated between the two species in the three mountain areas (H1). We expect sheep to host benzimidazole-resistant strains of gastrointestinal nematode, in line with the general pattern observed for sheep in France

(Papadopoulos et al., 2012; Rose Vineer et al., 2020). With the implementation of
reintroduction programs in the second half of the 20th century, ibex have colonized
pastures traditionally grazed by sheep. We therefore expect that ibex will also host
benzimidazole-resistant gastrointestinal nematodes but to a lesser extent, as
resistance do not represent a selective advantage for nematodes in the ibex
environment (Hahnel et al., 2018) (H2). Because there are very few documented ibex
dispersal events among the 3 ibex populations ((Brambilla, 2020), R. Papet, C. Toïgo
and E. Vannard, personal communication), we should observe genetic differences
among nematodes species/community or strains (ASV) among the populations of
ibex due to genetic drift (H3).
178
179 **Materials and Methods**
180
181 *Study area*
182
Samples of sheep and ibex feces were collected in the French Alps in 3
different mountain areas (Figure 1). The Belledonne mountain is located in the
western part of the Alps in southeast France. The Cerces and Champsaur mountains
are in the north and the south parts of the Ecrins National Park, respectively (Figure
1).

Figure 1 : Sampling locations in the French Alps were sheep and ibex feces were

collected. Pastoral unit: area where both ibex and sheep have been sampled.

Cerces : AL: Aiguillette de Lauzet, and MO: Montagne de l'Oule ; Belledonne : AB:

Ane Buyant, LP : La Pesée and SL : Sept Laux.

The 3 study areas are characterized by the presence of steep slopes, high

peaks (>2500m) and agropastoral activities. Climatic conditions are harsh in these

mountains with a mean temperature in winter (December-March) 2015-2019 of 0°C

in Belledonne (alt:1785m), 1°C in Champsaur (alt:1620m) and 1.5°C in Cerces

(alt:1553m). During summer (June-September), the mean temperature is 13°C in

Belledonne, 14°C in Champsaur and 16°C in the Cerces (réseau d'observation

météo du massif alpin). Champsaur is the southern study area and has a

Mediterranean influence. Consequently, rainfall is less important in this area

compared to Cerces and Belledonne. The vegetation is distributed along an elevation

gradient from coniferous woodland (*Abies alba* and *Picea abies* in Belledonne and

Larix decidua and *Pinus sylvestris* in Cerces and Champsaur) in the lower range of

ibex, to a landscape dominated by heathland with *Rhododendron ferrugineum*,

Vaccinium spp. and *Juniperus communis*, and grassland (*Carex* spp. *Festuca* spp.)
above the tree line (Ozenda, 1985).

The ibex populations were established in Belledonne, Cerces and Champsaur,
in 1983 with the introduction of 20 ibex, in 1959-1961 with the introduction of 6 ibex
and in 1994-1995 with the introduction of 30 ibex, respectively.

The distribution of ibex in Belledonne, Cerces and Champsaur range between
630m and 2860m on 200km², between 1410m and 3100m on 120km² and between
1320m and 3550m on 280km², respectively. Census populations sizes are estimated
to 800, 320 and 420 individuals in Belledonne, Cerces and Champsaur, respectively.

Traditional pastoral activity is practiced in all massifs, where sheep flocks
arrive early summer to graze mountain pastures coming from the plain on foot or by
truck. In Belledonne, the size of the herds are 750 ewes followed by their lambs in La
Pesée, 900 ewes in Sept Laux, and 1600 ewes in Ane Buyant. A dozen of rams are
also present within the La Pesée and Sept Laux herds, as well as some goats in La
Pesée. Each sheep herd from Belledonne belongs to one farmer while several
farmers grouped their sheep herds in Cerces and Champsaur. In Champsaur, the
herd included 4 breeding farms for a total of 1070 sheep and 5 goats. In Cerces, the
herd located in the West (Aiguillette du Lauzet) included 3 breeding farms for a total
of 800 sheep and the herd located in the East (Montagne de l'Oule) included 4
breeding farms for a total of 940 sheep.

Sample collection

We collected sheep feces up to 15 days after arrival on pasture to ensure that
we collected nematode species representative of sheep at the time of arrival on
pastures and to avoid detecting nematodes ingested secondarily on alpine pastures.
Similarly, ibex feces were collected prior to the arrival sheep and until 15 days after
arrival to ensure that the nematode community was not yet influenced by the arrival
of domestic livestock. Fresh ibex feces were mostly collected directly on the ground
and, in Belledonne, also during captures as part of the long-term monitoring program
conducted by the French Office for Biodiversity. Where possible, feces were collected
immediately after observation of ibex to avoid collecting of feces from the same
individual. Samples were stored in plastic bags, sealed after air removal, and

analyzed within 48h upon receipt maximum 15 days after field work (mean: 2.5 [min: 0 - max: 15] days) in the parasitology laboratory of the National Veterinary School of Lyon (ENVL, Marcy-l'Étoile, France). In total, we sampled 167 fecal samples from ibex and 90 fecal samples from 6 sheep herds, distributed over four study sites, i.e., Belledonne, Champsaur, Aiguillette du Lauzet, Cerces and Montagne de l'Oule, Cerces (Table 1, Figure S1).

Table 1 : Location, period and number of samples collected in the French alps. Coproscopic results of the gastro-intestinal (GI) nematodes (strongyles) eggs are also reported (med[min-max]). Cerces : AL: Aiguillette de Lauzet, and MO: Montagne de l'Oule ; Belledonne : AB: Ane Buyant, LP : La Pesée and SL : Sept Laux.

	Sampling date	N	GI nematodes epg
Cerces mountain			
Sheep	June 2019	AL: 15 MO: 15	AL: 7.5 [0-60] MO: 7.5 [0-225]
Ibex	AL: May-June 2019 MO: May 2018-2019	AL: 29 MO: 18	AL: 7.5 [0-30] MO: 7.5 [0-60]
Champsaur mountain			
Sheep	June 2019	15	7.5 [0-30]
Ibex	May 2019	40	7.5 [7.5-105]
Belledonne mountain			
Sheep	July 2019	AB: 15 LP: 15 SL: 15	AB: 30 [0-90] LP: 7.5 [0-165] SL: 0 [0-30]
Ibex	July 2018 and May-June 2019	80	15 [0-525]

Parasitological analyses

The number of gastro-intestinal nematodes eggs per gram of feces (epg) was counted following a modified McMaster protocol (Raynaud et al., 1970). In order that strongyles reach the L3 stage, coprocultures of feces were done at 24 ± 1 °C during 12-15 days with regular mixing and moistening. After the collection of L3 in tape water with a Baermann apparatus, we evaluated the success of the coproculture by counting the number of L3 in each sample. We extracted gastrointestinal nematodes DNA from samples for which there were at least 20 L3 and we limited the extraction to ~200 L3. DNA was extracted using extraction kit (Qiagen DNeasy® PowerSoil) following the manufacturer's instruction with an elution volume of 50 µl of water. We

extracted twice the DNA of 30 randomly chosen samples as internal extraction
controls. We quantified DNA concentration for all samples using Qubit 2.0
fluorometer (Life Technologies) and we homogenized DNA samples to a DNA
concentration of 1ng/µl (DNA samples were not diluted if the DNA concentration was
<1 ng/µl).

High throughput sequencing analyses

To determine the nemabiome of sheep and ibex we used a modified version of
the protocol developed by Avramenko et al., (2015). The ITS2 region of the nuclear
rDNA was amplified using the primer pair NC1 (Forward - 5'-
ACGTCTGGTTCAGGGTTGTT-3') and NC2 (Reverse- 5'-
TTAGTTTCTTTTCCTCCGCT-3') with the following PCR conditions: 10µl of Applied
Biosystems™ Master Mix AmpliTaq Gold™ 360, 5,84µl of molecular biology grade
water, 0,16µl of Bovine Serum Albumin, 2µL of 5 µM mixed F and R primers form, 2
µL of DNA lysate. The PCR was performed under the following conditions: 10 min
initial denaturation at 95°C; 35 cycles of denaturation (30 s at 95°C), annealing (30 s
at 54°C), and extension (1 min at 72°C); a final extension at 72°C for 7 min. The
thermocycling parameters were choose identically to Avramenko et al., (2015). To
detect the mutations responsible of the resistance of gastrointestinal nematode to
benzimidazole, we used a modified protocol of Avramenko et al., (2019). We
amplified the isotype-1 β-tubulin fragment comprising the codon at position 167, 198
and 200 with two pairs of primers in two independent PCR.

The PCR conditions were: 10µl of Applied Biosystems™ Master Mix
AmpliTaq Gold™ 360, 5.84 µl of molecular biology grade water, 0.16 µl of Bovine
Serum Albumin, 2 µL of 5 µM mixed forward and reverse primers mix, 2 µL of DNA
lysate. The PCR was performed under the following conditions: 10 min initial
denaturation at 95°C; 40 cycles of denaturation (30 s at 95°C), annealing (30 s at
65°C), and extension (30 s at 72°C); a final extension at 72°C for 7 min. We targeted
Teladorsagia circumcincta and *Trichostrongylus* spp. (Forward:5'-
CGCATTCWCTTGGAGGAGG-3' and Reverse: 5'-
GTGAGYTTCAAWGTGCGGAAG-3') and *Haemonchus contortus* (Forward:5'-
CGCATTCYTTGGGAGGAGG-3' and Reverse: 5'-GTGAGTTTYAAGGTGCGGAAG-

3') with the primers described by Avramenko et al., (2019). All forward and reverse

primers were tagged at 5' in order that each sample had a unique combination of

tagged primers.

In all PCRs, we added positive PCR controls (i.e., *Haemonchus contortus* and

Teladorsagia circumcincta DNA extracts), negative PCR controls (distilled H_2O) and

negative DNA extraction controls. All samples (including controls) were

independently amplified 4 times to ensure reliability of the sequencing, in 96-well

plates including 209 samples, 17 PCR positive controls, 13 PCR negative controls, 7

extraction negative controls, 30 DNA extraction controls, as well as 12 empty wells in

each 96-well plates to quantify tag jumping during PCR and sequencing steps (Figure

S2) (De Barba et al., 2014; Taberlet et al., 2018).

All PCR products of the ITS2 and the two β-tubulin isotype 1 sets were purified

using QIAquick® Spin Columns (QIAquick® PCR Purification KitQIAGEN) and

quantified using a Qubit 2.0 fluorometer (Life Technologies). Next, we pooled the 3

purified DNA pools (ITS2, two β-tubulin isotype 1) based on their initial concentration

and in proportion according to the following ratio: ITS2 50%, β-tubulin isotype 1 25%

for each. According to preliminary tests, we expect to achieve a sequencing depth of

20 000 reads per ITS2 DNA sample and 5 000 reads per β-tubulin isotype 1 DNA

sample. Sequencing was performed with pair-end sequencing technology on the

Illumina platform (2*250 bp Miseq) at Fasteris, Geneva, Switzerland.

Sequence analysis and taxon assignation

The sequence reads were first analyzed with the OBITOOLS package (Boyer et al.,

2016). Forward and reverse reads were assembled with the *alignpairedend* function,

and we kept only sequences with a good score of alignment (rnorm>0.8). Sequences

were attributed to their samples with the *ngsfilter* function with default parameters.

Subsequently, assigned sequences were analyzed with the dada2 package

(Callahan et al., 2016) following the pipeline available in www.nemabiome.ca. The

dada2 pipeline returns Amplicon Sequence Variant (ASV) which are sequence

variants differing by as little as one nucleotide (Callahan et al., 2017). Following

Beaumelle et al., (2021), the gastrointestinal nematodes were identified with three

different methods of assignation: BLASTn (Altschul et al., 1990) against the NCBI

database (November 2022), and AssignTaxonomy (Callahan et al., 2016; Wang et al., 2007) and IDTaxa (Murali et al., 2018) against the nematode ITS2 rDNA database 1.1.0 (Workentine et al., 2020) or the nematode β-tubulin isotype 1 DNA reference sequence supplied in Avramenko et al., (2019). We chose to attribute a confidence level to taxonomic identification at species level: high or moderate confidence if the three or two methods of assignation, respectively, were congruent. We also adjusted the sequence filtering based on an adapted procedure of Calderón‐Sanou et al., (2020). We kept only ASVs present in at least 2 replicates of the same samples and removed ASVs that were not assigned to genus level for the ITS2 and species level for β-tubulin isotype 1. We removed potential contaminants (reagent contaminants and cross-contaminations) following the procedure detailed in Calderón‐Sanou et al., (2020). For each sample, we sum the reads of the two replicates with the highest similarity and if this similarity is higher than the mean similarity among all replicates. At the end, we removed samples if they have <1000 reads of ITS2 and <500 reads of β-tubulin isotype 1 (Figure S3).

Identification of non-synonymous mutations in codons 167, 198 and 200

For each nematode species, all β-tubulin isotype 1 ASVs were aligned to one of the β-tubulin isotype 1 consensus sequences of the reference database (Avramenko et al., 2019) using the *AlignSeqs* function of the DECIPHER package (Wright, 2016). We examined each β-tubulin isotype 1 ASV at codon 167,198 and 200 positions to record whether the codon is associated with a non-synonymous mutation. However, we ignored the other polymorphism sites of the Exon.

Statistical analyses on measures of nemabiomes

To measure the differences of nemabiome among the two host species (sheep and ibex) and the 4 study sites (Aiguillette de Lauzet, Montagne de l'Oule, Champsaur and Belledonne), we considered two measures of diversity, i.e., the alpha diversity and the beta diversity. The alpha diversity was measured with the Shannon index that considers richness and evenness of communities and the beta diversity was measured with the weighted UniFrac index estimated using the R phyloseq package

(McMurdie and Holmes, 2013). The weighted UniFrac distance is a phylogenetic
distance between the set of ASV from each nemabiome weighted by the transformed
abundance of each ASV (Lozupone and Knight, 2005). The phylogenetic distances
were computed from a phylogenetic tree which was conducted using a maximum
likelihood tree with the GTR+G+I model according to the *ModelTest* function (Posada
and Crandall, 1998; Schliep, 2011). The exact count of ITS2 reads were transformed
with the Hellinger transformation (e.g., square root on relative frequencies) to account
for the high number of zero in community table and to decrease the influence of rare
ASVs in statistical analyses (Legendre and Legendre, 2012).

We tested the effects of host species and site, including their interaction, on alpha
diversity with linear models, and on beta diversity using perMANOVA (*adonis2*,
vegan R package (Oksanen et al., 2020)). All possible models including the null
model were computed. For perMANOVA models, we used a custom function to
compute Akaike's information criterion corrected for small sample size (AICc) based
on residual sums of squares (Dyson, 2018). In a model selection approach, for both
alpha and beta diversity, all possible models were ranked using the AICc and we
selected the model with the lowest AICc value. Models with ΔAICc ≤ 2 were
considered equivalent (Burnham and Anderson, 2002), and in this case, we
considered the most parsimonious one, i.e., the model with the lowest degree of
freedom.

All analyses were carried out using R 3.6 (R Core team, 2020. https://www.R-
project.org/).

Statistical analyses on measures of resistant nematode strains

To compare the importance of benzimidazole resistance of gastrointestinal
nematodes between ibex and sheep and among the 3 sites, we tested if the host
species, the site and the gastrointestinal nematode species influence the relative
abundance of ASVs with a resistant allele. For this purpose, we used a generalized
linear model with a binomial family and a model selection approach such as
described above.

We used *AlignSeqs* (Wright, 2016) to generate multi-sequence aligned β-tubulin
isotype 1 haplotype data. For each gastrointestinal species, we removed short ASVs,

e.g., ASVs with a sequence length <10% comparing to the median ASVs length.
PopART v1.7 (Leigh and Bryant, 2015) was used to draw the median joining
networks based on each gastrointestinal nematode haplotypes data.
397
398 **Results**
399
400 *Parasite material*
401
The median number of eggs of strongyles per gram of feces were lower in sheep
(7.5[0,148]$_{95\%IQR}$; n = 90) than in ibex (15[0,163]$_{95\%IQR}$; n = 167) feces (Mann–
Whitney U test; W = 9043.5, P = 0.006). As a result of the low level of infestation in
some samples, the number of L3 hatched from eggs were not sufficient (<20) for 48
ibex or sheep samples. These samples were not used for subsequent genetic
investigations. Specifically, all samples from the Sept Laux sheep herd (n=15,
Belledonne) were discarded. Therefore, the nemabiome was determined based on
the ITS2 for 196 (n= 55 sheep and n=141 ibex) out of 209 samples for which DNA
was extracted (Figure S1).
411
412 *Diversity of gastrointestinal nematodes in sheep and ibex*
413
In total, we detected 408 ASVs corresponding to 13 gastrointestinal nematode
species (Table 2, Figure S4). Eight ASVs were assigned to the genus level (i.e.,
Marshallagia spp., *Nematodirus* spp. and *Trichostrongylus* spp.) due to non-identical
assignation among taxonomic methods. An ASV corresponding to the lungworm
Cystocaulus ocreatus were discarded for the statistical analyses because we only
focus on gastrointestinal nematodes.
Teladorsagia circumcincta was the most prevalent nematode species and was
detected in 85% of samples (90%, n=127/141 ibex and 71%, n=39/56 sheep),
followed by *Trichostrongylus vitrinus,* 63% (73%, n=103/141 ibex and 36%, n=20/55
sheep) and *Haemonchus contortus,* 56% (70%, n=98/141 ibex and 22%, n=12/55
sheep) (Table 2, Figure S4). *Nematodirus* spp. and *Ostertagia leptospicularis* were
the rarest species and were detected in only 2 samples and 1 sample, respectively
and with a very small relative frequency (<0.1%).

The Nematodirus were not considered for the following results as our protocol is not
appropriate for this species (Hoberg et al., 2001)

Table 2 : Number of ASV, reads and samples for each nematode taxon, including the
results from the ITS2 (nemabiome) and the β-tubulin isotype 1 (resistance to
benzimidazole). N= number and percentage of samples in which the taxa was
detected. Host species and study sites are mixed here.

	ITS2			β-tubulin isotype 1		
	ASV	Reads	N (%)	ASV	Reads	N (%)
Bunostomum trigonocephalum	8	61 631	12 (6%)	-	-	-
Chabertia ovina	19	95 585	62 (32%)	-	-	-
Cooperia curticei	4	803	6 (3%)	-	-	-
Cooperia fuelleborni	2	358	5 (3%)	-	-	-
Haemonchus contortus	47	1 937 587	110 (56%)	38	392 110	96 (62%)
Marshallagia marshalli	23	338 064	57 (29%)	-	-	-
Marshallagia spp.	2	404	2 (1%)	-	-	-
Oesophagostomum venulosum	13	311 530	82 (42%)	-	-	-
Ostertagia leptospicularis	1	5	1 (1%)	-	-	-
Ostertagia ostertagi	2	269	2 (1%)	-	-	-
Teladorsagia circumcincta	235	3 311 094	166 (85%)	310	343 551	145 (94%)
Trichostrongylus axei	26	462 936	103 (53%)	33	68 414	104 (68%)
Trichostrongylus colubriformis	12	411 776	102 (52%)	8	144 643	112 (73%)
Trichostrongylus vitrinus	9	521 817	123 (63%)	44	92 834	107 (69%)
Trichostrongylus spp.	5	71 124	70 (36%)	-	-	-

According to the model selection approach, the host species was the only
factor that explained the alpha diversity (Table S1). The model indicates that sheep
had a lower alpha diversity compared to ibex (β = -0.42 ± 0.07, $P < 0.001$, R² of the
model=0.15) (Figure 2b). The site was not retained in the model selected to explain
the alpha diversity (Table S1).

Beta diversity is best explained by both factors, host species ($F_{1,188} = 27.69$, P
$= 0.001$) and site ($F_{3,188} = 16.39$, $P = 0.001$) and their interaction ($F_{3,188} = 25.61$, $P =$

0.001) according to our model selection (Table 3, Table S2). Some gastrointestinal

nematodes were mostly (*Trichostrongylus axei*: mean reads relative abundance

(RRA) of 19% [11%; 26%]$_{95CI}$ in sheep feces and 2% [1%;2%]$_{95CI}$ in ibex feces) or

only (*Bunostomum trigonocephalum, Cooperia* spp.) found in in sheep feces or only

in ibex feces (*Marshallagia* spp., *Ostertagia* spp.; Figure 2a). *Trichostrongylus*

colubriformis was more frequent in the Ecrins national park (mean RRA: 10% [6%;

13%]$_{95CI}$) than in Belledonne (mean RRA: 4% [2%; 6%]$_{95CI}$). Likewise, *Marshallagia*

spp. was more frequent in ibex feces in Cerces and Champsaur mountains (mean

RRA: 11% [6%;15%]$_{95CI}$) than in ibex feces in Belledonne (0.4% [-0.2%; 1%]$_{95CI}$). The

distribution of *Haemonchus contortus* in host species and sites had a particular

pattern. This parasite was more frequent in ibex feces compared to sheep feces in

Belledonne (mean RRA: 0.004% in sheep; 48% in ibex) and Champsaur (mean RRA:

0% in sheep; 41% in ibex), while the opposite was observed in the Cerces (Aiguillette

du Lauzet: mean RRA of 7% in sheep and 0.9% in ibex; Montagne de l'Oule: mean

RRA of 30% in sheep and 0% in ibex).

Table 3 : Parameters estimated from the best PerMANOVA model explaining the

beta diversity in ibex and sheep. The effect of host species (ibex or sheep) and site

(Aiguillette de Lauzet, Montagne de l'Oule, Champsaur or Belledonne) and the

interaction between the two factors are reported. Partial R² are reported with the

corresponding *F*-value and *p value* (P).

Diversity index	Best model selected	Variables	partial R²	F-value	P
Weighted UniFrac	β ~ Site x Host species	Residuals	0.55	-	-
		Site	0.14	16.39	0.001
		Host species	0.08	27.69	0.001
		Site: Host species	0.22	25.61	0.001

Figure 2 : (a) Mean relative frequencies of gastrointestinal nematodes species and
(b) Shannon diversity of ITS2 ASVs and (c) prevalence and mean relative
frequencies of β-tubulin isotype 1. For each host species (sheep or ibex) and each
site (Cerces: Montagne de l'Oule and Aiguillette de Lauzet; Champsaur and
Belledonne). The sample size for ITS2 ASVs (N_{ITS}; panel (a)) and for β-tubulin
isotype 1 ($N_β$; panels (b)) is given for each population. On the panel (c), the size of
the pie chart corresponds to the prevalence of the corresponding gastrointestinal
nematode species in the population, and the size of each sector to the mean
proportion of each allele.

Anthelmintic resistance

We found 433 different β-tubulin isotype 1 ASVs in 154 (n= 39 sheep and
n=115 ibex) out of 209 samples for which DNA was extracted. Among the 5
gastrointestinal nematode species targeted by primers, we detected, *H. contortus* in
96 samples, *Teladorsagia circumcincta* in 145 samples, *Trichostrongylus axei* in 104
samples, *Trichostrongylus vitrinus* in 107 samples, or *Trichostrongylus colubriformis*
in 112 samples (Table 2, Figure S5). No resistance mutation has been detected for *T.*
vitrinus. Therefore, *T. vitrinus* was not included in the model explaining the relative
abundance of resistant reads.

Resistance mutations were highly frequent (93.5%; n=144/154) with only 10
ibex feces (3 from Belledonne and 7 from Champsaur) in which no resistant mutation
has been detected.

Based on the best model for resistant RRA, the frequency of resistant nematodes
depended on gastrointestinal nematode species and the interaction between host
species and the study site (Table 4, Table S3). *Teladorsagia circumcincta* was the
species with the lowest resistant RRA and the nematode species with the higher
resistant RRA was *Trichostrongylus colubriformis* (Table 4). The mean observed
RRA of resistant nematodes differed between GIN species (*H. contortus:* 19%
$[13;25]_{95CI}$; *T. circumcincta:* 4% $[3;6]_{95CI}$; *T. axei:* 70% $[63;78]_{95CI}$; *T. colubriformis:*
96% $[93;99]_{95CI}$). The resistant RRA is generally lower in ibex compared to sheep (β
= -2.21 ± 0.66, *P* < 0.001). Resistant RRA were the lowest in the Aiguillette de Lauzet
but this is also the only site were ibex had significantly higher resistant RRA than
sheep (Table 4).

The most frequent resistance mutation is the F200Y present in the 4
gastrointestinal nematode species and 141 feces samples, follow by the E198A (46
samples, 2 nematodes species: *H. contortus* and *T. circumcincta*), the F167Y (17
samples, 2 nematodes species: *H. contortus* and *T. circumcincta*), the E198L (7
samples, *T. circumcincta*) and the E198G (4 samples, *H. contortus*) (Figure 2c).

Table 4 : Parameter estimates for the best generalized linear model explaining the
resistant reads relative abundance (RRA) in ibex and sheep. The effect of host
species (sheep as reference), study sites (Belledonne as reference), and their
interaction, in addition to the nematode species (*Teladorsagia circumcincta* as

reference) are reported. Parameter estimates with standard error (SE) are reported with the corresponding *z-value (z*-val*)* and *p value* (*P*). AL: Aiguillette de Lauzet; MO: Montagne de l'Oule; Ch: Champsaur; Hc : *Haemonchus contortus*; Ta: *Trichostrongylus axei*; Tcol: *Trichostrongylus colubriformis*.

Best model selected	Variables	Parameter estimate ± SE	z⬚val	*P*
Resistant RRA ~ host species x study sites + nematode species	Intercept	-2.02 ± 0.59	-3.45	5e-04
	Species	-2.21 ± 0.66	-3.35	8e-04
	Mountain(AL)	-2.89 ± 0.99	-2.93	0.003
	Mountain(MO)	-1.37 ± 0.89	1.54	0.123
	Mountain(Ch)	-0.10 ± 1.18	-0.08	0.935
	Nematode(Hc)	2.52 ± 0.62	4.05	5e-05
	Nematode(Ta)	4.68 ± 0.59	7.95	1e-15
	Nematode(Tcol)	7.38 ± 0.75	9.79	<2e-16
	Species:Site(AL)	2.90 ± 1.19	2.44	0.015
	Species:Site(MO)	0.04 ± 1.28	0.03	0.973
	Species:Site(Ch)	-0.75 ± 1.32	-0.56	0.574

Sheep and ibex shared 164 (38%) β-tubulin isotype 1 haplotypes and 238 (55%) β-tubulin isotype 1 haplotypes was only found in ibex samples (Figure 3). Most of the resistant haplotypes of *T. circumcincta* and *H. contortus*, e.g., containing a non-synonym mutation to the codon 167, 198 or 200, were genetic variants of a common sensitive haplotype shared by ibex and sheep (Figure 3). The resistant haplotypes of *T. axei* and *T. colubriformis* were more common than the sensitive haplotypes and the most similar sensitive haplotypes are found either in both sheep and ibex samples, or only in ibex samples. Both, *T. colubriformis* and *T. vitrinus* showed two distinct lineages, separated by ≥10 mutations. One of the lineages of *T. vitrinus* was only found in ibex from Belledonne and an ibex from Champsaur (Figure 3) while the ASVs of the second lineage are found both in ibex and sheep. One of the lineages of the *T. circumcincta* was more diverse (Figure 3).

528
Figure 3: Median joining network of β-tubulin isotype 1 haplotypes. Each point
represents a unique haplotype, and the colors correspond to the host species in
which the haplotype was detected. The size of the points is proportional to the
number of samples in which the haplotype was found. S: sensitive haplotype, R:

resistant haplotype. The tag above the points indicated the name of the mutation,
based on the codon position and the substitution of the amino acid.

Discussion

Because transhumant sheep and resident Alpine ibex use the same pastures
during the summer, we sought to assess the extent of nematode parasites sharing
between these two host species. Specifically, we investigated the presence of
anthelmintic-resistant nematode strains in sheep and ibex to determine the role of
transhumant sheep in contaminating alpine pastures, and whether ibex may play a
role in the maintenance and circulation of anthelmintic resistant nematodes. We used
a metabarcoding approach based the sequencing of ITS2 and the β-tubulin to
demonstrate that both sheep and ibex were infected by the same gastrointestinal
nematode species and shared anthelmintic-resistant strains, despite the absence of
sheep on alpine pastures for much of the year and therefore a narrow temporal
window of contamination.

In line with other studies investigating the gastrointestinal nematodes of sheep
and ibex (Burgess et al., 2012; Gruner et al., 2006; Redman et al., 2019; Zaffaroni et
al., 2000), the most prevalent and abundant species in both host species was
Teladorsagia circumcincta. Next, *Trichostrongylus vitrinus* was moderately prevalent,
but not abundant in sheep and ibex nemabiome. In accordance with the climatic
conditions of ibex environment, these two nematode species, as well as *Marshallagia*
spp., are better adapted to cold temperatures than the other nematode species
detected in this study (O'Connor et al., 2006; Zaffaroni et al., 2000). The sheep flocks
originate from French plains and/or the south of France and are driven into mountain
areas in early summer. Consequently, their nemabiome at the time of sampling is
representative of the gastrointestinal nematode communities present in sheep on the
farm, i.e., prior to transhumance. In a similar context, Gruner et al., (2006), observed
a high prevalence of *T. circumcincta* in two of three transhumant sheep flocks at the
beginning of the grazing season in the mountains south of the Alps. Furthermore,
transhumant sheep flocks appear to ingest mainly *T. circumcincta* when grazing in
the mountains, as this parasite remains the dominant species identified in feces and
tracer lambs during the summer (Gruner et al., 2006). Pastoral activity in

mountainous areas of France could therefore favor nematode species more adapted to cool and wet environmental conditions, such as *T. circumcincta* (O'Connor et al., 2006), compared with sheep grazing on the plains all year round. To confirm this hypothesis, a study of the variation in the nemabiome of transhumant sheep during the summer should be undertaken.

High relative frequency (>30%) of *H. contortus* was detected in ibex in Belledonne and Champsaur. In contrast, almost no *H. contortus* was observed in sheep flock droved in these mountains, raising the possibility that ibex may be contributing to the reinfection of sheep with *H. contortus*. To our knowledge, this is the first time that such relative abundance of *H. contortus* is reported in Alpine ibex (see previous studies based on morphological identification: Carcereri et al., 2021; Marreros et al., 2012; Zaffaroni et al., 2000). In addition, it should be noted that Alpine ibex were sampled before a potential contamination by domestic sheep could be detected, i.e., before the end of the pre-patent period: time between the infestation and the first eggs production, and at the end of spring – early summer, i.e., at the start of the epidemiological period at risk for *Haemonchus contortus* in a high-altitude mountain area. We can therefore expect higher levels of contaminations in late summer, when domestic sheep leave mountain pastures. In addition, we cannot exclude that some laboratory issues might have reduced the apparent prevalence and abundance of *H. contortus* as some samples from sheep have been kept in the fridge at 4°C during 2 to 3 days (including e.g., the sheep samples without *H. contortus* from Belledonne and Champsaur) which can reduce the proportion of *H. contortus* eggs hatching (McKenna, 1998).

The detection of *H. contortus* raises conservation issues for Alpine ibex as this nematode species is known to be highly pathogenic to sheep (Taylor et al., 2015). Infection of a phylogenetically related species, the Pyrenean ibex (*Capra pyrenaica pyrenaica*), with a few thousand of *H. contortus* resulted in severe clinical signs, including extremely low weight and hemorrhagic anemia (Lavín et al., 1997). In addition, *H. contortus* may have been involved, along with pneumonia, in the collapse of the Northern Chamois in the province of Lecco, Italy from November 2000 to March 2001 (Citterio et al., 2006). As gastrointestinal nematodes can have an impact on the demographic dynamics of the host population (Acerini et al., 2022; Albery et al., 2021; Albon et al., 2002), they are suspected of being involved in the low natality rates observed in the French Alpine ibex populations (Brambilla et al., 2020). While

Alpine ibex appears to be fairly resilient to parasite infections (Marreros et al., 2012), further investigations should be carried out to assess the consequences of gastrointestinal nematodes infections for ibex at both individual and population level.

We detected anthelmintic resistant alleles in 4 out of the 5 nematodes species, namely *H. contortus*, *T. circumcincta*, *Trichostrongylus axei*, *Trichostrongylus colubriformis*, but not *T. vitrinus*, for which the β-tubulin amplicons have been sequenced. Both sheep and ibex hosted resistant strains of the 4 nematode species and only 10 out of 116 ibex did not have resistant strains to any of the nematode species studied. The benzimidazole resistance was therefore very common in the studied sheep flocks, in agreement with the situation of sheep farms in Europe (Rose et al., 2015; Rose Vineer et al., 2020). The presence of anthelmintic resistant nematodes in ibex is most likely explained by the indirect transmission of resistant nematodes from sheep to ibex through the environment. The large number of shared β-tubulin ASVs between sheep and ibex and the high overlap between their nemabiome confirm this scenario (Figure 2c, Figure 3). This is in accordance with other studies investigating the share of nematode parasites at the interface of wild and domestic ungulates (Beaumelle et al., 2022; Cerutti et al., 2010; Laca Megyesi et al., 2019).

It is worth noting that feces of ibex were sampled before the arrival of sheep on pastures. This important result demonstrates that anthelmintic resistant nematodes can be maintained in mountainous areas from year to year in wild populations of ibex despite harsh winter environmental conditions, and in the absence of the main source of parasites during most of the year, i.e., the domestic sheep. The shedding of eggs from resistant nematodes by ibex prior to the arrival of domestic sheep suggests the potential role of ibex as a reservoir of anthelmintic resistant nematodes for other susceptible domestic and wild ungulates. In addition, the position of resistant mutant strains detected in ibex at the periphery of haplotypes networks (Figure 3) supported the relatively recent selection of benzimidazole resistance and the lack of benzimidazole resistant reversion since the resistant strains have been transmitted to ibex.

The 5 nematodes species for which we have studied the resistances seemed to have different selection dynamics which may reflect the life history traits of species (Redman et al., 2015). In fact, we detected no resistant allele in *T. vitrinus* and conversely, the proportion of benzimidazole resistant strains of *T. axei* and *T.*

colubriformis were high in sheep and in a lesser extent in ibex (Figure 2c). The
resistance proportions of *T. circumcincta* and *H. contortus* were lower compared to *T.*
axei and *T. colubriformis*, excepted for the *H. contortus* of the sheep flock of the
Montagne de l'Oule. In this study area, the proportion of resistant strains of *H.*
contortus was very high.

Consistent with our study, benzimidazole-resistant strains of *T. vitrinus* were rare
in other studies of sheep farms (in the UK, Avramenko et al., 2019, and in Canada,
Queiroz et al., 2020). In contrast, high frequencies of benzimidazole resistance in *T.*
axei and *T. colubriformis* (between 40% and 100%) were already reported in sheep;
in UK, *T. axei*: 26-27% and *T. colubriformis:* 53-62% (Avramenko et al., 2019); in
Austria, *T. colubriformis:* 77%-100%, (Hinney et al., 2020); in France, *T. axei*: 63%,
(Palcy et al., 2010). In contrast, Hinney et al., (2020), observed in transhumant sheep
flocks in Austria Alps, a higher mean frequency of the F200Y resistance allele in *T.*
circumcincta: 32.4 ± 6.8% (mean ± standard error of the mean) and *H. contortus:*
91.9 ± 3.7%, compared to the sheep flocks of this study (*T. circumcincta*, 6.6 ± 3.5%
and *H. contortus,* 69.9 ± 14.4%, all resistant alleles combined).

Several factors are suspected to contribute to interspecific differences in the
selection of resistance strains between nematode species, including specific
reproductive rate, seasonal dynamics, climatic conditions in the location of sheep
farms, anthelmintic strategies, e.g., treatment molecules, timing and rate of
anthelmintic treatments, grazing management and the cost of benzimidazole
resistance (Hodgkinson et al., 2019; Redman et al., 2015). However, the links
between traits of parasites and interspecific variation of resistant acquisition by
gastrointestinal nematodes have not been tested yet (Morgan et al., 2019). Our
results suggested a few clues in relation to the ecology of nematode species.

Firstly, nematode species have different abilities to practice hypobiosis, i.e., the
ability to halt embryonic development under environmental constraints (Gibbs, 1986).
H. contortus and *T. circumcincta* are known to arrest development more frequently
than *Trichostrongylus* spp. (Langrová et al., 2008), and hypobiotic larvae have been
shown to be less sensitive to drug (Sargison et al., 2007). Secondly, among the
Trichostrongylus spp., *T. vitrinus* may have a higher proportion of overwintering
larvae in pastures as this species is more resistant to cold temperature compared to
T. axei and *T. colubriformis* (O'Connor et al., 2006). As parasites on pasture are not

subject to selection pressure by anthelmintics, they are a source of susceptible
strains.

As the proportion of resistant strains is generally lower in ibex compared to sheep,
ibex may have contributed to a dilution effect of resistant strains, i.e., by hosting
susceptible nematodes. However, the role of ibex in the maintenance of a refugia
needs to be investigated by considering the relative number of susceptible strains
deposited by ibex on pasture compared with sheep. Furthermore, it seems that the
role of ibex in the maintenance of a refugia vary according to nematode species. For
example, ibex excrete a lower proportion of *T. axei* eggs than sheep (Figure 2a),
while resistant strains are also dominant in ibex (Figure 2c). In contrast, *T.*
circumcincta and *H. contortus* in ibex were more frequently susceptible and
genetically diverse (higher number of ITS2 and β-tubulin ASVs) compared with *T.*
axei and *T. colubriformis* ibex (Table 2, Figure 3). As *T. circumcincta* was dominant in
ibex, a refuge of susceptible *T. circumcincta* strains may have been maintained within
ibex and may contribute to limit the spread of resistance in sheep farms. The results
concerning *H. contortus* were variable according to the massifs since ibex from the
two Cerces sites had a low amount of *H. contortus* compared to the two other
populations, i.e., Belledonne and Champsaur.

The divergence between massifs can be observed at the community and genetic
level among sheep flocks and ibex populations. Indeed, it was expected that some
differences in nemabiome composition would be observed between the massifs and
sheep flocks, given that sheep and ibex from different areas never meet (R. Papet, C.
Toïgo, E. Vannard, Pers. communication). Furthermore, the sheep flocks come from
different locations and have been subjected to different anthelmintic strategies. For
the ibex, differences in the original population of translocated animals (Gauthier and
Villaret, 1990; Kessler et al., 2022) and potential founder effect – not all parasites
present in the source population were present in the newly established sink
population - may have had a long term impact on the composition of the nemabiome.
For example, the distinct population of *T. vitrinus*, found mainly in ibex in Belledonne,
may had been inherited from the founding ibex population. This highlights that
various reintroductions of ibex in the study area can also influence the composition of
the parasite community. This distinct population *of T. vitrinus* was absent in sheep
grazing in Belledonne, which hosted other strains of *T. vitrinus* and despite several
summers of co-occurrence of sheep and ibex in Belledonne. In is possible that sheep

never graze on pastures contaminated by these strains, but this observation may also be due to a sampling bias as the number of sheep sampled remains low. Within the same mountain area, few differences have, however, been observed among the nemabiome of ibex, e.g., between the ibex of the Aiguillette de Lauzet and those of the Montagne de l'Oule, whereas the sheep flocks host distinct nemabiome communities.

In conclusion, transmissions of gastrointestinal nematodes species, including resistant nematodes strains, have occurred between sheep and ibex even though the contact between the two species is limited to the summer period. In this study, we demonstrated more specifically that ibex can maintain and shed eggs of resistant gastrointestinal nematodes despite the absence of sheep on pastures for several months, suggesting a potential role of ibex as a reservoir for these nematodes. However, the extent to which each host species can influence the nematode community of the other during the transhumant period remains to be determined. To this end, it would be useful to analyze the nemabiome of sheep and ibex before and after the transhumant period. The lower proportion of resistance allele in ibex nemabiome compared with that of sheep underlines the possibility that ibex could contribute to the maintenance and circulation of susceptible strains in sheep. Based on our results, it seems that ibex have helped to limit the spread of anthelmintic resistance of *T. circumcincta* and *H. contortus* in sheep flocks, by maintaining parasite refugia not exposed to anthelmintic pressure. As with sheep and roe deer (Beaumelle et al., 2022), domestic sheep contributes to the modification of the nemabiome of ibex. This raises concerns about the conservation of the ibex, and the consequences of strongyles infection in ibex should be investigated. Indeed, ibex is characterized by low genetic diversity due to the strong demographic decline of this species, followed by multiple re-introduction (Grossen et al., 2018) and a high genetic structure of immunity-related loci among populations (Kessler et al., 2022) whereas both neutral and adaptive genetic diversity are known to have an influence parasite resistance in ungulates (Portanier et al., 2019).

Appendices

733 Supplementary data to this article can be found online in MendeleyData (DOI:
734 10.17632/cm97cg87d6.1).

Acknowledgements

The authors warmly thank all the professionals from the Office Français de la
Biodiversité and all the trainees for data collection, and J. S. Gilleard, E. Redman
from the University of Calgary and C. Lionnet from the Laboratoire d'Ecologie Alpine
(LECA) and other members of the lab to help us developing the deep sequencing
analyses for nematodes. The research benefited from the support of AnaBM (USMB)
and AEEM (UGA) laboratory facilities.

Funding

This project was founded by the Office Français de la Biodiversité, the Laboratoire
d'Ecologie Alpine (LECA) and VetAgro Sup - Pôle d'Expertise Vétérinaire et
Agronomique des Animaux Sauvages (EVAAS, France; http://evaas.vetagro￼sup.fr/;
DGAL—VetAgro Sup - INRAE funding). G. Bourgoin was supported by the
AgreenSkills+ fellowship program (European Union program; MarieCurie FP7
COFUND People Programme; grant agreement n_609398).

Conflict of interest disclosure

755 The authors declare that they have no conflict of interest.

Data, scripts, and supplementary information availability

759 The bioinformatic pipeline, the ASV analysed during the current study and the R
760 script of statistic analyses are available in MendeleyData (DOI:
761 10.17632/cm97cg87d6.1).

References

Acerini, C.I., Morris, S., Morris, A., Kenyon, F., McBean, D., Pemberton, J.M., Albery,
G.F., 2022. Helminth parasites are associated with reduced survival probability
in young red deer. Parasitology 149, 1702–1708.
https://doi.org/10.1017/S0031182022001111

Acevedo, P., Cassinello, J., Gortazar, C., 2008. The Iberian ibex is under an
expansion trend but displaced to suboptimal habitats by the presence of
extensive goat livestock in central Spain, in: Hawksworth, D.L., Bull, A.T.
(Eds.), Biodiversity and Conservation in Europe, Topics in Biodiversity and
Conservation. Springer Netherlands, Dordrecht, pp. 119–134.
https://doi.org/10.1007/978-1-4020-6865-2_9

Albery, G.F., Morris, A., Morris, S., Kenyon, F., Nussey, D.H., Pemberton, J.M., 2021.
Fitness costs of parasites explain multiple life-history trade-offs in a wild
mammal. Am. Nat. 197, 324–335. https://doi.org/10.1086/712633

Albon, S.D., Stien, A., Irvine, R.J., Langvatn, R., Ropstad, E., Halvorsen, O., 2002.
The role of parasites in the dynamics of a reindeer population. Proc. R. Soc.
Lond. B Biol. Sci. 269, 1625–1632. https://doi.org/10.1098/rspb.2002.2064

Avramenko, R.W., Redman, E.M., Lewis, R., Yazwinski, T.A., Wasmuth, J.D.,
Gilleard, J.S., 2015. Exploring the gastrointestinal "Nemabiome": deep
amplicon sequencing to quantify the species composition of parasitic
nematode communities. PLoS One 10(12), e0143559.
https://doi.org/10.1371/journal.pone.0143559

Avramenko, R.W., Redman, E.M., Melville, L., Bartley, Y., Wit, J., Queiroz, C.,
Bartley, D.J., Gilleard, J.S., 2019. Deep amplicon sequencing as a powerful
new tool to screen for sequence polymorphisms associated with anthelmintic
resistance in parasitic nematode populations. Int. J. Parasitol. 49, 13–26.
https://doi.org/10.1016/j.ijpara.2018.10.005

Beaumelle, C., Redman, E., Verheyden, H., Jacquiet, P., Bégoc, N., Veyssière, F.,
Benabed, S., Cargnelutti, B., Lourtet, B., Poirel, M.-T., de Rijke, J., Yannic, G.,
Gilleard, J.S., Bourgoin, G., 2022. Generalist nematodes dominate the
nemabiome of roe deer in sympatry with sheep at a regional level. Int. J.
Parasitol. https://doi.org/10.1016/j.ijpara.2022.07.005

Beaumelle, C., Redman, E.M., de Rijke, J., Wit, J., Benabed, S., Debias, F.,
Duhayer, J., Pardonnet, S., Poirel, M.-T., Capron, G., Chabot, S., Rey, B.,
Yannic, G., Gilleard, J.S., Bourgoin, G., 2021. Metabarcoding in two isolated

populations of wild roe deer (*Capreolus capreolus*) reveals variation in

gastrointestinal nematode community composition between regions and

among age classes. Parasit. Vectors 14, 594. https://doi.org/10.1186/s13071-

021-05087-5

Biber, J.-P., 2010. Transhumance in France. Pastoralism 1, 91–98.

https://doi.org/10.3362/2041-7136.2010.006

Blaxter, M., Koutsovoulos, G., 2015. The evolution of parasitism in Nematoda.

Parasitology 142, S26–S39. https://doi.org/10.1017/S0031182014000791

Bordes, L., Dumont, N., Lespine, A., Souil, E., Sutra, J.-F., Prévot, F., Grisez, C.,

Romanos, L., Dailledouze, A., Jacquiet, P., 2020. First report of multiple

resistance to eprinomectin and benzimidazole in *Haemonchus contortus* on a

dairy goat farm in France. Parasitol. Int. 76, 102063.

https://doi.org/10.1016/j.parint.2020.102063

Boyer, F., Mercier, C., Bonin, A., Le Bras, Y., Taberlet, P., Coissac, E., 2016.

obitools: a unix-inspired software package for DNA metabarcoding. Mol. Ecol.

Resour. 16, 176–182. https://doi.org/10.1111/1755-0998.12428

Brambilla, A., 2020. Caratterizzazione genetica delle popolazioni di stambecco delle

Alpi occidentali. Department of Evolutionary Biology and Environmental

Studies, Zurich.

Brambilla, A., Bassano, B., Biebach, I., Bollmann, K., Keller, L., Toïgo, C., von

Hardenberg, A., 2022. Alpine Ibex *Capra ibex* Linnaeus, 1758, in: Corlatti, L.,

Zachos, F.E. (Eds.), Terrestrial Cetartiodactyla, Handbook of the Mammals of

Europe. Springer International Publishing, Cham, pp. 383–408.

https://doi.org/10.1007/978-3-030-24475-0_32

Brambilla, A., Von Hardenberg, A., Nelli, L., Bassano, B., 2020. Distribution, status,

and recent population dynamics of Alpine ibex *Capra ibex* in Europe.

Mammal Rev. 50, 267–277. https://doi.org/10.1111/mam.12194

Brivio, F., Ciuti, S., Pipia, A., Grignolio, S., Apollonio, M., 2022. Livestock displace

European mouflon from optimal foraging sites. Eur. J. Wildl. Res. 68, 30.

https://doi.org/10.1007/s10344-022-01581-y

Brown, T.L., Airs, P.M., Porter, S., Caplat, P., Morgan, E.R., 2022. Understanding the

role of wild ruminants in anthelmintic resistance in livestock. Biol. Lett. 18,

20220057. https://doi.org/10.1098/rsbl.2022.0057

Burgess, C.G.S., Bartley, Y., Redman, E., Skuce, P.J., Nath, M., Whitelaw, F., Tait,

A., Gilleard, J.S., Jackson, F., 2012. A survey of the trichostrongylid nematode

species present on UK sheep farms and associated anthelmintic control

practices. Vet. Parasitol. 189, 299–307.

https://doi.org/10.1016/j.vetpar.2012.04.009

Burnham, K.P., Anderson, D.R., 2002. Model selection and multimodel inference: a

practical information-theoretic approach, 2nd ed. Springer, New York.

https://doi.org/10.1007/b97636

Calderón-Sanou, I., Münkemüller, T., Boyer, F., Zinger, L., Thuiller, W., 2020. From

environmental DNA sequences to ecological conclusions: how strong is the

influence of methodological choices? J. Biogeogr. 47, 193–206.

https://doi.org/10.1111/jbi.13681

Callahan, B.J., McMurdie, P.J., Holmes, S.P., 2017. Exact sequence variants should

replace operational taxonomic units in marker-gene data analysis. ISME J. 11,

2639–2643. https://doi.org/10.1038/ismej.2017.119

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes,

S.P., 2016. DADA2: high-resolution sample inference from Illumina amplicon

data. Nat. Methods 13, 581–583. https://doi.org/10.1038/nmeth.3869

Carcereri, A., Stancampiano, L., Marchiori, E., Sturaro, E., Ramanzin, M., Cassini,

R., 2021. Factors influencing gastrointestinal parasites in a colony of Alpine

ibex (*Capra ibex*) interacting with domestic ruminants. Hystrix Ital. J. Mammal.

32, 95–101. https://doi.org/10.4404/hystrix-00393-2020

Carlsson, A.M., Irvine, R.J., Wilson, K., Coulson, S.J., 2013. Adaptations to the

Arctic: low-temperature development and cold tolerance in the free-living

stages of a parasitic nematode from Svalbard. Polar Biol. 36, 997–1005.

https://doi.org/10.1007/s00300-013-1323-7

Cerutti, M.C., Citterio, C.V., Bazzocchi, C., Epis, S., D'Amelio, S., Ferrari, N.,

Lanfranchi, P., 2010. Genetic variability of *Haemonchus contortus* (Nematoda:

Trichostrongyloidea) in alpine ruminant host species. J. Helminthol. 84, 276–

283. https://doi.org/10.1017/S0022149X09990587

Charlier, J., Bartley, D.J., Sotiraki, S., Martinez-Valladares, M., Claerebout, E., von

Samson-Himmelstjerna, G., Thamsborg, S.M., Hoste, H., Morgan, E.R.,

Rinaldi, L., 2022. Chapter Three - Anthelmintic resistance in ruminants:

challenges and solutions, in: Rollinson, D., Stothard, R. (Eds.), Advances in

Parasitology. Academic Press, pp. 171–227.

https://doi.org/10.1016/bs.apar.2021.12.002

Charlier, J., Rinaldi, L., Musella, V., Ploeger, H.W., Chartier, C., Vineer, H.R., Hinney,

B., von Samson-Himmelstjerna, G., Băcescu, B., Mickiewicz, M., Mateus, T.L.,

Martinez-Valladares, M., Quealy, S., Azaizeh, H., Sekovska, B., Akkari, H.,

Petkevicius, S., Hektoen, L., Höglund, J., Morgan, E.R., Bartley, D.J.,

Claerebout, E., 2020. Initial assessment of the economic burden of major

parasitic helminth infections to the ruminant livestock industry in Europe. Prev.

Vet. Med. 182, 105103. https://doi.org/10.1016/j.prevetmed.2020.105103

Chintoan-Uta, C., Morgan, E.R., Skuce, P.J., Coles, G.C., 2014. Wild deer as

potential vectors of anthelmintic-resistant abomasal nematodes between cattle

and sheep farms. Proc. R. Soc. B Biol. Sci. 281, 20132985.

https://doi.org/10.1098/rspb.2013.2985

Chirichella, R., Apollonio, M., Putman, R., 2014. Competition between domestic and

wild Ungulates., in: Putman, R., Apollonio, M. (Eds.), Behaviour and

Management of European Ungulates. pp. 110–123.

Citterio, C.V., Caslini, C., Milani, F., Sala, M., Ferrari, N., Lanfranchi, P., 2006.

Abomasal Nematode Community in an Alpine Chamois (*Rupicapra r.*

rupicapra) Population before and after a Die-Off. J. Parasitol. 92, 918–927.

De Barba, M., Miquel, C., Boyer, F., Mercier, C., Rioux, D., Coissac, E., Taberlet, P.,

2014. DNA metabarcoding multiplexing and validation of data accuracy for diet

assessment: application to omnivorous diet. Mol. Ecol. Resour. 14, 306–323.

https://doi.org/10.1111/1755-0998.12188

Drudge, J.H., Szanto, J., Wyant, Z.N., Elam, G., 1964. Field studies on parasite

control in sheep: comparison of thia-bendazole, ruelene, and phenothiazine.

Am. J. Vet. Res. 25, 1512–1518.

Dyson, K., 2018. Custom community ecology helper R scripts.

https://github.com/kdyson/R_Scripts

Fiel, C.A., Fernández, A.S., Rodríguez, E.M., Fusé, L.A., Steffan, P.E., 2012.

Observations on the free-living stages of cattle gastrointestinal nematodes.

Vet. Parasitol. 187, 217–226. https://doi.org/10.1016/j.vetpar.2012.01.011

Francis, E.K., Šlapeta, J., 2023. Refugia or reservoir? Feral goats and their role in the

maintenance and circulation of benzimidazole-resistant gastrointestinal

nematodes on shared pastures. Parasitology 1–11.

https://doi.org/10.1017/S0031182023000380

Gauthier, D., Villaret, J., 1990. La réintroduction en France du bouquetin des Alpes.

Rev Eco Terre Vie 45, 97–120. https://doi.org/10.3406/revec.1990.6338

Gibbs, H.C., 1986. Hypobiosis in Parasitic Nematodes—An Update, in: Baker, J.R.,

Muller, R. (Eds.), Advances in Parasitology. Academic Press, pp. 129–174.

https://doi.org/10.1016/S0065-308X(08)60343-7

Grossen, C., Biebach, I., Angelone-Alasaad, S., Keller, L.F., Croll, D., 2018.

Population genomics analyses of European ibex species show lower diversity

and higher inbreeding in reintroduced populations. Evol. Appl. 11, 123–139.

https://doi.org/10.1111/eva.12490

Gruner, L., Sauvé, C., Boulard, C., Calamel, M., 2006. Analysis of the relationship

between land use and the parasitism of sheep during their transhumance.

Anim. Res. 55, 177–188. https://doi.org/10.1051/animres:2006009

Hahnel, S.R., Zdraljevic, S., Rodriguez, B.C., Zhao, Y., McGrath, P.T., Andersen,

E.C., 2018. Extreme allelic heterogeneity at a *Caenorhabditis elegans* beta-

tubulin locus explains natural resistance to benzimidazoles. PLOS Pathog. 14,

e1007226. https://doi.org/10.1371/journal.ppat.1007226

Hinney, B., Schoiswohl, J., Melville, L., Ameen, V.J., Wille-Piazzai, W., Bauer, K.,

Joachim, A., Krücken, J., Skuce, P.J., Krametter-Frötscher, R., 2020. High

frequency of benzimidazole resistance alleles in trichostrongyloids from

Austrian sheep flocks in an alpine transhumance management system. BMC

Vet. Res. 16, 132. https://doi.org/10.1186/s12917-020-02353-z

Hoberg, E.P., Kocan, A.A., Rickard, L.G., 2001. Gastrointestinal strongyles in wild

ruminants, in: William, S.M., Pybus, M.J., Kocan, A.A. (Eds.), Parasitic

Diseases of Wild Mammals. Iowa State University Press, Ames, Iowa. USA,

pp. 193–227.

Hodgkinson, J.E., Kaplan, R.M., Kenyon, F., Morgan, E.R., Park, A.W., Paterson, S.,

Babayan, S.A., Beesley, N.J., Britton, C., Chaudhry, U., Doyle, S.R., Ezenwa,

V.O., Fenton, A., Howell, S.B., Laing, R., Mable, B.K., Matthews, L., McIntyre,

J., Milne, C.E., Morrison, T.A., Prentice, J.C., Sargison, N.D., Williams, D.J.L.,

Wolstenholme, A.J., Devaney, E., 2019. Refugia and anthelmintic resistance:

Concepts and challenges. Int. J. Parasitol. Drugs Drug Resist. 10, 51–57.

https://doi.org/10.1016/j.ijpddr.2019.05.001

Hudson, P.J., Dobson, A.P., Lafferty, K.D., 2006. Is a healthy ecosystem one that is
rich in parasites? Trends Ecol. Evol., Twenty years of TREE - part 2 21, 381–
385. https://doi.org/10.1016/j.tree.2006.04.007

Kaplan, R.M., Vidyashankar, A.N., 2012. An inconvenient truth: Global worming and
anthelmintic resistance. Vet. Parasitol., Special issue: novel approaches to the
control of helminth parasites of livestock 186, 70–78.
https://doi.org/10.1016/j.vetpar.2011.11.048

Kessler, C., Brambilla, A., Waldvogel, D., Camenisch, G., Biebach, I., Leigh, D.M.,
Grossen, C., Croll, D., 2022. A robust sequencing assay of a thousand
amplicons for the high-throughput population monitoring of Alpine ibex
immunogenetics. Mol. Ecol. Resour. 22, 66–85. https://doi.org/10.1111/1755-
0998.13452

Khanyari, M., Robinson, S., Milner-Gulland, E.J., Morgan, E.R., Rana, R.S.,
Suryawanshi, K.R., 2022. Pastoralism in the high Himalayas: Understanding
changing practices and their implications for parasite transmission between
livestock and wildlife. Pastoralism 12, 44. https://doi.org/10.1186/s13570-022-
00257-1

Laca Megyesi, Š., Königová, A., Babják, M., Molnár, L., Rajský, M., Szestáková, E.,
Major, P., Soroka, J., Urda Dolinská, M., Komáromyová, M., Várady, M., 2019.
Wild ruminants as a potential risk factor for transmission of drug resistance in
the abomasal nematode *Haemonchus contortus*. Eur. J. Wildl. Res. 66, 9.
https://doi.org/10.1007/s10344-019-1351-x

Langrová, I., Makovcová, K., Vadlejch, J., Jankovská, I., Petrtýl, M., Fechtner, J.,
Keil, P., Lytvynets, A., Borkovcová, M., 2008. Arrested development of sheep
strongyles: onset and resumption under field conditions of Central Europe.
Parasitol. Res. 103, 387–392. https://doi.org/10.1007/s00436-008-0984-6

Lavín, S., Marco, I., Rossi, L., Meneguz, P.G., Viñas, L., 1997. Haemonchosis in
Spanish Ibex. J. Wildl. Dis. 33, 656–659. https://doi.org/10.7589/0090-3558-
33.3.656

Legendre, P., Legendre, L., 2012. Numerical Ecology. Elsevier.

Leigh, J.W., Bryant, D., 2015. popart: full-feature software for haplotype network
construction. Methods Ecol. Evol. 6, 1110–1116. https://doi.org/10.1111/2041-
210x.12410

Lozupone, C., Knight, R., 2005. UniFrac: a new phylogenetic method for comparing
microbial communities. Appl. Environ. Microbiol. 71, 8228–8235.
https://doi.org/10.1128/AEM.71.12.8228-8235.2005

Luzzago, C., Ebranati, E., Cabezón, O., Fernández-Sirera, L., Lavín, S., Rosell, R.,
Veo, C., Rossi, L., Cavallero, S., Lanfranchi, P., Marco, I., Zehender, G., 2016.
Spatial and Temporal Phylogeny of Border Disease Virus in Pyrenean
Chamois (*Rupicapra p. pyrenaica*). PLOS ONE 11, e0168232.
https://doi.org/10.1371/journal.pone.0168232

Marchand, P., Freycon, P., Herbaux, J.-P., Game, Y., Toïgo, C., Gilot-Fromont, E.,
Rossi, S., Hars, J., 2017. Sociospatial structure explains marked variation in
brucellosis seroprevalence in an Alpine ibex population. Sci. Rep. 7, 15592.
https://doi.org/10.1038/s41598-017-15803-w

Marreros, N., Frey, C.F., Willisch, C.S., Signer, C., Ryser-Degiorgis, M.-P., 2012.
Coprological analyses on apparently healthy Alpine ibex (*Capra ibex ibex*)
from two Swiss colonies. Vet. Parasitol. 186, 382–389.
https://doi.org/10.1016/j.vetpar.2011.11.009

McKenna, P.B., 1998. The effect of previous cold storage on the subsequent
recovery of infective third stage nematode larvae from sheep faeces. Vet.
Parasitol. 80, 167–172. https://doi.org/10.1016/S0304-4017(98)00203-9

McMurdie, P.J., Holmes, S., 2013. phyloseq: An R Package for Reproducible
Interactive Analysis and Graphics of Microbiome Census Data. PLOS ONE 8,
e61217. https://doi.org/10.1371/journal.pone.0061217

Morgan, E.R., Aziz, N.-A.A., Blanchard, A., Charlier, J., Charvet, C., Claerebout, E.,
Geldhof, P., Greer, A.W., Hertzberg, H., Hodgkinson, J., Höglund, J., Hoste,
H., Kaplan, R.M., Martínez-Valladares, M., Mitchell, S., Ploeger, H.W., Rinaldi,
L., von Samson-Himmelstjerna, G., Sotiraki, S., Schnyder, M., Skuce, P.,
Bartley, D., Kenyon, F., Thamsborg, S.M., Vineer, H.R., de Waal, T., Williams,
A.R., van Wyk, J.A., Vercruysse, J., 2019. 100 Questions in Livestock
Helminthology Research. Trends Parasitol. 35, 52–71.
https://doi.org/10.1016/j.pt.2018.10.006

Murali, A., Bhargava, A., Wright, E.S., 2018. IDTAXA: a novel approach for accurate
taxonomic classification of microbiome sequences. Microbiome 6, 140.
https://doi.org/10.1186/s40168-018-0521-5

Nagy, G., Csivincsik, Á., Sugár, L., Zsolnai, A., 2017. Benzimidazole resistance
within red deer, roe deer and sheep populations within a joint habitat in
Hungary. Small Rumin. Res. 149, 172–175.
https://doi.org/10.1016/j.smallrumres.2017.02.012

O'Connor, L.J., Walkden-Brown, S.W., Kahn, L.P., 2006. Ecology of the free-living
stages of major trichostrongylid parasites of sheep. Vet. Parasitol. 142, 1–15.
https://doi.org/10.1016/j.vetpar.2006.08.035

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D.,
Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H.,
Szoecs, E., Wagner, H., 2020. vegan□: community ecology package.

Ozenda, P., 1985. La végétation de la chaîne alpine dans l'espace montagnard
européen. Masson, Paris.

Palcy, C., Silvestre, A., Sauve, C., Cortet, J., Cabaret, J., 2010. Benzimidazole
resistance in *Trichostrongylus axei* in sheep: long-term monitoring of affected
sheep and genotypic evaluation of the parasite. Vet. J. 183, 68–74.
https://doi.org/10.1016/j.tvjl.2008.09.012

Papadopoulos, E., Gallidis, E., Ptochos, S., 2012. Anthelmintic resistance in sheep in
Europe: a selected review. Vet. Parasitol., Special issue: Update on Parasitic
Diseases of Sheep 189, 85–88. https://doi.org/10.1016/j.vetpar.2012.03.036

Portanier, E., Garel, M., Devillard, S., Maillard, D., Poissant, J., Galan, M., Benabed,
S., Poirel, M.-T., Duhayer, J., Itty, C., Bourgoin, G., 2019. Both candidate gene
and neutral genetic diversity correlate with parasite resistance in female
Mediterranean mouflon. BMC Ecol. 19, 12. https://doi.org/10.1186/s12898-
019-0228-x

Posada, D., Crandall, K.A., 1998. modeltest: testing the model of DNA substitution.
Bioinformatics 14, 817–818. https://doi.org/10.1093/bioinformatics/14.9.817

Queiroz, C., Levy, M., Avramenko, R., Redman, E., Kearns, K., Swain, L., Silas, H.,
Uehlinger, F., Gilleard, J.S., 2020. The use of ITS-2 rDNA nemabiome
metabarcoding to enhance anthelmintic resistance diagnosis and surveillance
of ovine gastrointestinal nematodes. Int. J. Parasitol. Drugs Drug Resist. 14,
105–117. https://doi.org/10.1016/j.ijpddr.2020.09.003

Raynaud, J.-P., William, G., Brunault, G., 1970. Etude de l'efficacité d'une technique
de coproscopie quantitative pour le diagnostic de routine et le contrôle des

infestations parasitaires des bovins, ovins, équins et porcins. Ann. Parasitol.

Hum. Comparée 45, 321–342. https://doi.org/10.1051/parasite/1970453321

Redman, E., Queiroz, C., Bartley, D.J., Levy, M., Avramenko, R.W., Gilleard, J.S.,

2019. Validation of ITS-2 rDNA nemabiome sequencing for ovine

gastrointestinal nematodes and its application to a large scale survey of UK

sheep farms. Vet. Parasitol. 275, 108933.

https://doi.org/10.1016/j.vetpar.2019.108933

Redman, E., Whitelaw, F., Tait, A., Burgess, C., Bartley, Y., Skuce, P.J., Jackson, F.,

Gilleard, J.S., 2015. The emergence of resistance to the benzimidazole

anthlemintics in parasitic nematodes of livestock is characterised by multiple

independent hard and soft selective sweeps. PLoS Negl. Trop. Dis. 9,

e0003494. https://doi.org/10.1371/journal.pntd.0003494

Richomme, C., Gauthier, D., Fromont, E., 2006. Contact rates and exposure to inter-

species disease transmission in mountain ungulates. Epidemiol. Infect. 134,

21–30. https://doi.org/10.1017/S0950268805004693

Roeber, F., Jex, A.R., Gasser, R.B., 2013. Impact of gastrointestinal parasitic

nematodes of sheep, and the role of advanced molecular tools for exploring

epidemiology and drug resistance - an Australian perspective. Parasit. Vectors

6, 153. https://doi.org/10.1186/1756-3305-6-153

Rose, H., Rinaldi, L., Bosco, A., Mavrot, F., Waal, T. de, Skuce, P., Charlier, J.,

Torgerson, P.R., Hertzberg, H., Hendrickx, G., Vercruysse, J., Morgan, E.R.,

2015. Widespread anthelmintic resistance in European farmed ruminants: a

systematic review. Vet. Rec. 176, 546–546. https://doi.org/10.1136/vr.102982

Rose Vineer, H., Morgan, E.R., Hertzberg, H., Bartley, D.J., Bosco, A., Charlier, J.,

Chartier, C., Claerebout, E., de Waal, T., Hendrickx, G., Hinney, B., Höglund,

J., Ježek, J., Kašný, M., Keane, O.M., Martínez-Valladares, M., Mateus, T.L.,

McIntyre, J., Mickiewicz, M., Munoz, A.M., Phythian, C.J., Ploeger, H.W.,

Rataj, A.V., Skuce, P.J., Simin, S., Sotiraki, S., Spinu, M., Stuen, S.,

Thamsborg, S.M., Vadlejch, J., Varady, M., von Samson-Himmelstjerna, G.,

Rinaldi, L., 2020. Increasing importance of anthelmintic resistance in

European livestock: creation and meta-analysis of an open database. Parasite

27, 69. https://doi.org/10.1051/parasite/2020062

Rossi, L., Tizzani, P., Rambozzi, L., Moroni, B., Meneguz, P.G., 2019. Sanitary emergencies at the wild/domestic caprines interface in Europe. Animals 9, 922. https://doi.org/10.3390/ani9110922

Ryser-Degiorgis, M.-P., Ingold, P., Tenhu, H., Less, A.M.T., Ryser, A., Giacometti, M., 2002. Encounters between Alpine ibex, Alpine chamois and domestic sheep in the Swiss Alps. Hystrix Ital. J. Mammal. 13. https://doi.org/10.4404/hystrix-13.1-2-4180

Sargison, N.D., Wilson, D.J., Bartley, D.J., Penny, C.D., Jackson, F., 2007. Haemonchosis and teladorsagiosis in a Scottish sheep flock putatively associated with the overwintering of hypobiotic fourth stage larvae. Vet. Parasitol. 147, 326–331. https://doi.org/10.1016/j.vetpar.2007.04.011

Schliep, K.P., 2011. phangorn: phylogenetic analysis in R. Bioinformatics 27, 592–593. https://doi.org/10.1093/bioinformatics/btq706

Taberlet, P., Bonin, A., Zinger, L., Coissac, E., 2018. Environmental DNA: for biodiversity research and monitoring. Oxford University Press, Oxford, UK.

Taylor, M.A., Coop, R.L., Wall, R.L., 2015. Veterinary Parasitology, 4th ed. ed. Wiley Blackwell, Chichester.

Tompkins, D.M., Dunn, A.M., Smith, M.J., Telfer, S., 2011. Wildlife diseases: from individuals to ecosystems. J. Anim. Ecol. 80, 19–38. https://doi.org/10.1111/j.1365-2656.2010.01742.x

Utaaker, K.S., Ytrehus, B., Davey, M.L., Fossøy, F., Davidson, R.K., Miller, A.L., Robertsen, P.-A., Strand, O., Rauset, G.R., 2023. Parasite spillover from domestic sheep to wild reindeer—The role of salt licks. Pathogens 12, 186. https://doi.org/10.3390/pathogens12020186

Vercruysse, J., Charlier, J., Dijk, J.V., Morgan, E.R., Geary, T., Samson-Himmelstjerna, G. von, Claerebout, E., 2018. Control of helminth ruminant infections by 2030. Parasitology 145, 1655–1664. https://doi.org/10.1017/S003118201700227X

Walker, J.G., Morgan, E.R., 2014. Generalists at the interface: nematode transmission between wild and domestic ungulates. Int. J. Parasitol. Parasites Wildl. 3, 242–250. https://doi.org/10.1016/j.ijppaw.2014.08.001

Wang, Q., Garrity, G.M., Tiedje, J.M., Cole, J.R., 2007. Naïve bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. Appl. Environ. Microbiol. 73, 5261–5267. https://doi.org/10.1128/AEM.00062-07

Whittaker, J.H., Carlson, S.A., Jones, D.E., Brewer, M.T., 2017. Molecular

mechanisms for anthelmintic resistance in strongyle nematode parasites of

veterinary importance. J. Vet. Pharmacol. Ther. 40, 105–115.

https://doi.org/10.1111/jvp.12330

Workentine, M.L., Chen, R., Zhu, S., Gavriliuc, S., Shaw, N., Rijke, J. de, Redman,

E.M., Avramenko, R.W., Wit, J., Poissant, J., Gilleard, J.S., 2020. A database

for ITS2 sequences from nematodes. BMC Genet. 21, 74.

https://doi.org/10.1186/s12863-020-00880-0

Wright, E.S., 2016. Using DECIPHER v2.0 to Analyze Big Biological Sequence Data

in R. R J. 8, 352–359.

Zaffaroni, E., Teresa Manfredi, M., Citterio, C., Sala, M., Piccolo, G., Lanfranchi, P.,

2000. Host specificity of abomasal nematodes in free ranging alpine

ruminants. Vet. Parasitol. 90, 221–230. https://doi.org/10.1016/S0304-

4017(00)00240-5

1112