Cross-transmission of resistant gastrointestinal nematodes between wildlife and transhumant sheep

Camille Beaumelle[a,b,c,1,*], Carole Toïgo[d], Rodolphe Papet[e], Slimania Benabed[a,b], Mathieu Beurier[d], Léa Bordes[f], Anaïs Brignone[d], Nadine Curt-Grand-Gaudin[c], Mathieu Garel[d], Justine Ginot[b], Philippe Jacquiet[f], Christian Miquel[c], Marie-Thérèse Poirel[a,b], Anna Serafino[b], Eric Vannard[g], Gilles Bourgoin[a,b, †], Glenn Yannic[c, †]

[a] *Université de Lyon, Université Lyon 1, CNRS, Laboratoire de Biométrie et Biologie Evolutive UMR 5558, F-69100 Villeurbanne, France*

[b] *Université de Lyon, VetAgro Sup, Campus Vétérinaire de Lyon, F-69280 Marcy l'Etoile, France*

[c] *Université Grenoble Alpes, Université Savoie Mont Blanc, CNRS, LECA, 38000, Grenoble, France*

[d] Office Français de la Biodiversité, Unité Ongulés Sauvages, Gières, France

[e] Parc national des Écrins, Secteur du Champsaur-Valgaudemar, 05260 Saint Jean Saint Nicolas, France

[f] Université de Toulouse, UMT Pilotage de la Santé des Ruminants, Ecole Nationale Vétérinaire de Toulouse, France

[g] Parc national des Écrins, Secteur du Briançonnais, 05100 Briançon, France

*Corresponding author: beaumelle.camille@gmail.com

† Co-senior authors

ABSTRACT

Wild and domestic ungulates can be infected with the same species of gastrointestinal parasitic nematodes. These parasites have free-living stages in the environment that contribute to the ease of transmission among different host species. In addition, gastrointestinal nematodes have developed resistance to anthelmintics which is now considered a major problem for the livestock sector. In a context where wild and domestic ungulates share the same pastures, the maintenance and circulation of resistant gastrointestinal nematodes between species have rarely been explored.

In the European Alps, domestic sheep are driven to high-altitude summer pastures and live in sympatry with wild ungulates for several months each year. In this study, we investigated the nemabiome of domestic sheep and Alpine ibex, *Capra ibex*, in three different areas of the French Alps to evaluate parasite circulation between the two host species. The Alpine ibex is a protected mountain ungulate that is phylogenetically related to sheep and hosts nematode species common to sheep.

Using internal transcribed spacer 2 (ITS-2) nemabiome metabarcoding, we found sheep and ibex share similar gastrointestinal nematodes, except for a few species such as *Marshallagia marshalli* and *Trichostrongylus axei*. This suggests that the long-term co-occurrence of sheep and ibex on mountain pastures has promoted the exchange of gastrointestinal nematodes between the two hosts. Based on the sequencing of the isotype 1 of the beta tubulin gene, associated with benzimidazole resistance, we found resistant nematodes in all sheep flocks and in all ibex populations. Our results demonstrated that ibex can host and shed resistant strains before transhumant sheep arrive on pastures, and thus could act as refuge or even contribute to maintain resistant gastrointestinal nematodes. The relative role of ibex in the maintenance and circulation of resistant strains in sheep remain to be determined.

Keywords: ITS-2 rDNA, benzimidazole resistance; β-tubulin isotype 1; livestock; nemabiome metabarcoding; wild ungulates, transhumant sheep, Alpine ibex

Introduction

Parasites represent a large proportion of animal diversity and are key components of food webs (Hudson et al., 2006). They are also essential determinants of the health, fitness, population dynamics and community composition of their hosts (Tompkins et al., 2011). Parasites of the Nematoda phylum infect a wide range of species worldwide, including animals and plants (Blaxter and Koutsovoulos, 2015). In animals, gastrointestinal nematode parasites are of major concern for livestock productivity and security as they can impact animal health and reduce animal production, with significant economic losses (Charlier et al., 2020; Roeber et al., 2013). Ungulates are usually infected by free-living larvae of gastrointestinal nematodes when they graze pasture. Infective larvae may survive several months in the environment depending on species and climatic conditions (O'Connor et al., 2006). Following ingestion, larvae finish their development to reach the adult stage in the digestive tract. Egg-laying occurs 2–4 weeks post infection. The duration of infection by nematodes varies depending on species but lasts for at least 2 months from the L3 ingestion (Deplazes et al., 2016). The larvae can arrest their development in the host (hypobiosis) during harsh climatic conditions, delaying egg-laying (Deplazes et al., 2016).

To limit parasite load and its impact on livestock health, the use of anthelmintics to treat livestock against gastrointestinal nematodes is a common and cost effective practice (Vercruysse et al., 2018). Nonetheless, the repeated use of anthelmintics has led to the selection of anthelmintic-resistant strains of gastrointestinal nematodes. Resistance to several families of anthelmintics (e.g., benzimidazole, macrocyclic lactones and levamisole) has been observed, and multi-resistance is increasing (Bordes et al., 2020; Kaplan and Vidyashankar, 2012; Rose et al., 2015; Rose Vineer et al., 2020).

In particular, resistance to benzimidazoles is widespread throughout the world (Kaplan and Vidyashankar, 2012), and is particularly common on sheep farms in Europe (Papadopoulos et al., 2012; Rose Vineer et al., 2020). Contrary to other anthelmintic families, the mechanisms of resistance to benzimidazoles are well known and documented (Whittaker et al., 2017); in resistant nematodes, specific mutations of the β-tubulin isotype-1 gene have been correlated with resistance (Charlier et al., 2022). Furthermore, large-scale screening based on molecular tools is now feasible for this resistance (Avramenko et al., 2019), whereas the recommended method in

livestock (i.e., the fecal egg count reduction test; Kaplan et al., 2023) for the diagnosis

of resistance to other anthelmintics requires techniques that are difficult to achieve in

wildlife leaving in remote locations.

Some generalist gastrointestinal nematodes can infect several host species

(Walker and Morgan, 2014), including both domestic and wild ungulates (e.g.,

Beaumelle et al., 2022; Cerutti et al., 2010). The transmission of gastrointestinal

nematodes among hosts, even if they do not simultaneously occupy the same

pastures, is possible thanks to their free-living infective larval stage that may remain

active several months in the environment (Carlsson et al., 2013; Fiel et al., 2012;

Walker and Morgan, 2014). Transmitted parasites can also include gastrointestinal

nematodes resistant to anthelmintics. For instance, benzimidazole-resistant

nematodes have been detected in free-living populations of roe deer, *Capreolus*

capreolus, living in sympatry with livestock (Chintoan-Uta et al., 2014; Nagy et al.,

2017). To date, the role of wild ungulates in the epidemiology of resistant nematodes

remains to be determined, but it has been suggested that wildlife may act as a reservoir

of resistant nematodes for livestock (Brown et al., 2022; Chintoan-Uta et al., 2014;

Francis and Šlapeta, 2023; Laca Megyesi et al., 2019; Walker and Morgan, 2014). To

accurately evaluate this hypothesis, we need to investigate the presence of resistant

nematodes in co-grazing wild and domestic ungulates in different contexts (i.e.,

different host species, different landscapes, and under different climatic conditions).

Transhumant pastoralism is a common practice in the European Alps and consists

in the seasonal movement of grazing livestock from lowland areas to mountain

meadows in summer which provide fresh pasture for domestic ungulates, i.e., mainly

sheep, but also cows or goats (Biber, 2010). These mountainous areas are inhabited

year-round by wild ungulates, particularly those living at high altitude in the European

Alps, like Alpine ibex (*Capra ibex*), or Northern chamois (*Rupicapra rupicapra*). While

wild ungulates tend to avoid domestic herds spatially or temporarily during the summer

(Acevedo et al., 2008), certain factors may contribute to the use of the same pastures

by both groups.

Spatial segregation between wild and domestic ungulates is usually observed once

livestock arrive on pasture (Brivio et al., 2022; Ryser-Degiorgis et al., 2002). Livestock

are generally released onto the best grazing areas during the summer season

(Chirichella et al., 2014; Richomme et al., 2006; Ryser-Degiorgis et al., 2002).

Mountain ungulates have been observed to preferentially use the same grazing areas

both before and after use by livestock (Brivio et al., 2022; Ryser-Degiorgis et al., 2002). The presence of wild and domestic ungulates in attracting zones such as salt licks, even if not simultaneous, offers good opportunities for parasite transmission, and these areas are therefore considered hotspots for parasite infection (Richomme et al., 2006; Ryser-Degiorgis et al., 2002; Utaaker et al., 2023).

Consequently, transhumant pastoralism represents a risk for pathogen transmission between wild and domestic ungulates in mountain areas (Rossi et al., 2019). Pathogen exchange at the interface of wild and domestic ungulates have already been well documented. The Alpine ibex has been identified as the wildlife reservoir of brucellosis (*Brucella melitensis*) which was transmitted to cattle and humans in the Bargy massif in northern French Alps (Marchand et al., 2017). In addition, sheep have been confirmed as the domestic reservoir of Border disease, which induced a major viral outbreak in Southern chamois (*Rupicapra pyrenaica*) populations in the Pyrenees (Luzzago et al., 2016). The transmission of gastrointestinal nematodes has already been described between wild ungulates and transhumant domestic ungulates in mountainous areas (Cerutti et al., 2010; Citterio et al., 2006; Khanyari et al., 2022; Zaffaroni et al., 2000). However, no study has yet investigated the transmission of anthelmintic-resistant nematodes in a transhumant pastoral system.

In this study, we investigated the community of gastrointestinal nematodes infecting Alpine ibex and domestic sheep (*Ovis aries*) and the prevalence of resistance to benzimidazole in three different regions of the French Alps. The Alpine ibex was close to extinction at the beginning of the 19th century but the reinforcement of its populations by several reintroductions in different parts of the Alps has increased the species' overall abundance and range (Brambilla et al., 2022). Today, the Alpine ibex population is estimated at 52 000 individuals in Europe (Brambilla et al., 2020).

Ibex usually host species-specific gastrointestinal nematodes (Walker and Morgan, 2014) but they may also be exposed to generalist nematodes deposited by other related ungulate (i.e., Northern chamois, *Rupicapra rupicapra*) or domestic species (e.g., sheep). Furthermore, anthelmintic treatments are frequently applied to livestock by farmers with the aim of reducing the parasite load and hence reducing the diversity of nematodes in sheep. Therefore, we expected the nemabiome to be highly differentiated between the two species in the three studied mountain areas, with higher nematode diversity in ibex compared to sheep (H1). We expected sheep to host

benzimidazole-resistant strains of gastrointestinal nematodes, in line with the general
pattern observed for sheep in France (Papadopoulos et al., 2012; Rose Vineer et al.,
2020). With the reintroduction programs in the second half of the 20^{th} century, ibex
colonized pastures traditionally grazed by sheep. We therefore expected that ibex
would also host benzimidazole-resistant gastrointestinal nematodes, but to a lesser
extent, as resistance does not represent a selective advantage for nematodes in the
ibex environment (Hahnel et al., 2018) (H2). Because there are very few documented
ibex dispersal events among the 3 ibex populations (Brambilla, 2020 R. Papet, C.
Toïgo and E. Vannard, personal communication), we predicted genetic differences
among nematode species/communities or strains (ASV: Amplicon sequence variant)
among the ibex populations due to genetic drift (H3).
178
Materials and Methods
180
Study area
182
183 Samples of sheep and ibex feces were collected in the French Alps in 3 different
184 mountain areas (Figure 1). The Belledonne mountain is located in the western part of
185 the Alps in southeast France. The Cerces and Champsaur mountains are in the north
186 and the south parts of the Ecrins National Park, respectively (Figure 1).

Figure 1 : Sampling locations in the French Alps where sheep and ibex feces were collected. Pastoral unit: area where both ibex and sheep have been sampled. Cerces : AL: Aiguillette de Lauzet, and MO: Montagne de l'Oule ; Belledonne : AB: Ane Buyant, LP : La Pesée and SL : Sept Laux.

The 3 study areas are characterized by the presence of steep slopes, high peaks (>2500m) and agropastoral activities. Climatic conditions are harsh in these mountains with a mean temperature in winter (December-March) 2015-2019 of 0°C in Belledonne (alt:1785m), 1°C in Champsaur (alt:1620m) and 1.5°C in Cerces (alt:1553m). During summer (June-September), the mean temperature is 13°C in Belledonne, 14°C in Champsaur and 16°C in the Cerces (Réseau d'Observation Météo du Massif Alpin; www.romma.fr). Champsaur, the most southern study area, has a Mediterranean influence. Consequently, rainfall is less important in this area compared to Cerces and Belledonne. Across areas, the vegetation is distributed along an elevation gradient from coniferous woodland (*Abies alba* and *Picea abies* in Belledonne and *Larix decidua* and *Pinus sylvestris* in Cerces and Champsaur) in the lower range of ibex, to a landscape dominated by heathland with *Rhododendron ferrugineum*, *Vaccinium* spp. and *Juniperus communis*, and grassland (*Carex* spp. *Festuca* spp.) above the tree line (Ozenda, 1985).

The ibex populations were established in Belledonne, Cerces and Champsaur, in 1983 with the introduction of 20 ibex, in 1959-1961 with the introduction of 6 ibex and in 1994-1995 with the introduction of 30 ibex, respectively. Traditional pastoral activity is practiced in all massifs, where sheep flocks arrive from the plain on foot or by truck in early summer to graze mountain pastures.

In the Belledonne mountain, the distribution of ibex range between 630m and 2860m over 200km². Population size is estimated at 800 individuals. The size of the sheep herds are 750 ewes followed by their lambs in La Pesée, 900 ewes in Sept Laux, and 1600 ewes in Ane Buyant. A dozen rams are also present within the La Pesée and Sept Laux herds, as well as some goats in La Pesée.

The ibex population located in the Cerces mountain is estimated at 320 individuals and occupies an area of 120km², between 1410m and 3100m. In Cerces, the sheep herd located in the West (Aiguillette du Lauzet) includes 3 breeding farms for a total of 800 sheep and the herd located in the East (Montagne de l'Oule) includes 4 breeding farms for a total of 940 sheep (Table 1).

In the Champsaur mountain, the distribution of ibex range between 1320m and 3550m over 280km². Population size is estimated at 420 individuals. In Champsaur, the sheep herd includes 4 breeding farms for a total of 1070 sheep and 5 goats (Table 1).

Each sheep herd belongs to one farmer while several farmers grouped their sheep herds in Cerces and Champsaur (Table 1).

Table 1 : Descriptive data on study sites.

	Area (km²)	Altitude (m)	Number of introduced ibex	Ibex Population size	Domestic flock size	Climate
Belledonne mountain	200	630-2860	20	800	AB: 1600 LP: 750 SL: 900	Mountain climate
Cerces mountain	120	1410-3100	6	320	AL: 800 MO: 940	Mountain climate
Champsaur mountain	280	1320-3550	30	420	1070	Mountain climate with Mediterranean influence

Sample collection

We collected sheep feces for 15 days after arrival on pasture to ensure that we collected nematode species representative of sheep at the time of arrival on pastures and not those ingested secondarily on alpine pastures. Similarly, ibex feces were collected prior to the arrival sheep and until 15 days after arrival to ensure that the nematode community was not influenced by the arrival of domestic livestock. Fresh ibex feces were mostly collected directly on the ground and, in Belledonne, also during captures as part of the long-term monitoring program conducted by the French Office for Biodiversity. Ibex feces were collected within each of the pastoral units in which we collected sheep feces. Where possible, feces were collected immediately after observation of ibex to avoid collecting of feces from the same individual. Feces from all age groups were collected. Samples were stored in plastic bags, sealed after air removal, and analyzed within 48h upon receipt in the parasitology laboratory of the National Veterinary School of Lyon (ENVL, Marcy-l'Étoile, France) or up to a maximum of 15 days after field collection (mean: 2.5 days). In total, we sampled 167 fecal samples from ibex and 90 fecal samples from 6 sheep herds, distributed over 6 pastoral units, i.e., Aiguillette du Lauzet, Montagne de l'Oule, Champsaur, Ane Buyant, La Pesée and Sept Laux (Table 2, Figure S1). In Belledonne, 21 samples were collected in 2018 following the sampling strategy of 2019. We controlled that year of sampling did not result in a drastic change in the nemabiome of ibex (Figure S6) and included those samples in analyses.

Table 2 : Location, period and number of samples of ibex and sheep collected in the French Alps. Coproscopic results of the gastro-intestinal (GI) nematodes (strongyles) in eggs per gram (epg) of feces are also reported (median[min-max]).

	Sampling date	N	GI nematodes epg
Cerces mountain			
Sheep	June 2019	AL: 15	AL: 7.5 [0-60]
		MO: 15	MO: 7.5 [0-225]
Ibex	AL: May-June 2019	AL: 29	AL: 7.5 [0-30]
	MO: May 2019	MO: 18	MO: 7.5 [0-60]
Champsaur mountain			
Sheep	June 2019	15	7.5 [0-30]
Ibex	May 2019	40	7.5 [7.5-105]
Belledonne mountain			
Sheep	July 2019	AB: 15	AB: 30 [0-90]
		LP: 15	LP: 7.5 [0-165]
		SL: 15	SL: 0 [0-30]
Ibex	July 2018 and May-June 2019	80	15 [0-525]

Cerces : AL: Aiguillette de Lauzet, and MO: Montagne de l'Oule ; Belledonne : AB: Ane Buyant, LP : La Pesée and SL : Sept Laux.

Parasitological analyses

The number of gastro-intestinal nematode eggs per gram of feces (epg) was counted following a modified McMaster protocol (Raynaud et al., 1970) with a solution of zinc sulphate ($ZnSO_4$, density = 1.36, 1/15 dilution). The eggs were counted on a McMaster slide with two chambers (theoretical sensitivity of 15 epg). We also checked for the presence of low abundant parasite propagules with a "control slide". We prepared this control slide with a 14mL tube filled with the remaining solution until a meniscus is formed. We covered the tube with a coverslip and centrifuge the tube with a basket centrifuge (5 min at 1200 rpm) to help propagules to rise and stick on the coverslip. After centrifugation, the coverslip was transferred on a microscope slide for microscopical observation. We attributed the value of 7.5 epg for parasites with no egg observed on the McMaster, but at least one egg observed on the control slide (for a similar procedure see Beaumelle et al, 2021).

In order for strongyles to reach the L3 stage, coprocultures of feces were done at 24 ± 1 °C during 12-15 days with regular mixing and moistening. We then collected

the L3 in tape water with a Baermann apparatus. We extracted gastrointestinal nematode DNA from samples for which there were at least 20 L3 and we limited the extraction to ~200 L3. DNA was extracted using an extraction kit (Qiagen DNeasy® PowerSoil) following the manufacturer's instruction with an elution volume of 50 µl of water. We extracted the DNA of 30 randomly chosen samples twice as internal extraction controls (Taberlet et al., 2018). We quantified the DNA concentration for all samples using Qubit 2.0 fluorometer (Life Technologies) and homogenized DNA samples to a DNA concentration of 1ng/µl (DNA samples were not diluted if the DNA concentration was <1 ng/µl).

High throughput sequencing analyses

To determine the nemabiome of sheep and ibex, we used a modified version of the protocol developed by Avramenko et al (2015). The ITS2 region of the nuclear rDNA was amplified using the primer pair NC1 (Forward - 5'-ACGTCTGGTTCAGGGTTGTT-3') and NC2 (Reverse- 5'-TTAGTTTCTTTTCCTCCGCT-3') with the following PCR conditions: 10µl of Applied Biosystems™ Master Mix AmpliTaq Gold™ 360, 5.84µl of molecular biology grade water, 0.16µl of Bovine Serum Albumin, 2µL of 5 µM mixed F and R primers form, 2 µL of DNA lysate. The PCR was performed under the following conditions: 10 min initial denaturation at 95°C; 35 cycles of denaturation (30 s at 95°C), annealing (30 s at 54°C), and extension (1 min at 72°C); a final extension at 72°C for 7 min.

To detect the mutations responsible for the resistance of gastrointestinal nematodes to benzimidazole, we used a modified protocol of Avramenko et al., (2019). Using the same mix as previously described, we amplified the β-tubulin isotype 1 fragment comprising the codons at position 167, 198 and 200 with two pairs of primers in two independent PCRs. The PCR was performed under the following conditions: 10 min initial denaturation at 95°C; 40 cycles of denaturation (30 s at 95°C), annealing (30 s at 65°C), and extension (30 s at 72°C); a final extension at 72°C for 7 min. We targeted *Teladorsagia circumcincta* and *Trichostrongylus* spp*.* (Forward:5'-CGCATTCWCTTGGAGGAGG-3' and Reverse: 5'-GTGAGYTTCAAWGTGCGGAAG-3') and *Haemonchus contortus* (Forward:5'-CGCATTCYTTGGGAGGAGG-3' and Reverse: 5'-GTGAGTTTYAAGGTGCGGAAG-

3') with the primers described by Avramenko et al (2019). All forward and reverse primers were tagged at 5' in order that each sample had a unique combination of tagged primers.

In all PCRs, we added positive PCR controls (i.e., *Haemonchus contortus* and *Teladorsagia circumcincta* DNA extracts), negative PCR controls (distilled H_2O) and negative DNA extraction controls. All samples (including controls) were tagged with unique barcode identifiers to allow pooling into a single amplicon library (Taberlet et al., 2018), and all samples were independently amplified 4 times to ensure sequencing reliability. Amplifications were carried out in 96-well plates, totaling 209 ibex and sheep samples, 17 PCR positive controls, 13 PCR negative controls, 7 extraction negative controls, 30 DNA extraction controls, as well as 12 empty wells in each plate to quantify tag jumping during PCR and sequencing steps (Figure S2; De Barba et al., 2014; Taberlet et al., 2018).

All PCR products of the ITS2 and the two β-tubulin isotype 1 sets were purified using QIAquick® Spin Columns (QIAquick® PCR Purification KitQIAGEN) and quantified using a Qubit 2.0 fluorometer (Life Technologies). Next, we pooled the 3 purified DNA pools (ITS2, two β-tubulin isotype 1) based on their initial concentration and in proportion according to the following ratio: ITS2 50%, β-tubulin isotype 1 25% for each. According to preliminary tests, we expected to achieve a sequencing depth of 20 000 reads per ITS2 DNA sample and 5 000 reads per β-tubulin isotype 1 DNA sample. Sequencing was performed with pair-end sequencing technology on the Illumina platform (2*250 bp Miseq) at Fasteris, Geneva, Switzerland.

Sequence analysis and taxon assignation

The sequence reads were first analyzed with the OBITOOLS package (Boyer et al., 2016). Forward and reverse reads were assembled with the *alignpairedend* function, and we kept only sequences with a good score of alignment (rnorm>0.8). Sequences were attributed to their samples with the *ngsfilter* function with default parameters. Subsequently, assigned sequences were analyzed with the dada2 package (Callahan et al., 2016) following the pipeline available in www.nemabiome.ca. The *dada2* pipeline returns Amplicon Sequence Variants (ASV) which are sequence variants differing by as little as one nucleotide (Callahan et al., 2017). Following Beaumelle et al. (2021),

gastrointestinal nematodes were identified with four different methods of assignation-databases: BLASTn (Altschul et al., 1990) based on (1) the NCBI database (Accessed: November 2022), and (2) AssignTaxonomy (Callahan et al., 2016; Wang et al., 2007) and (3) IDTaxa (Murali et al., 2018) based on the nematode ITS2 rDNA database 1.1.0 (Workentine et al., 2020). To identify the species associated with the β-tubulin sequences, we used IDTaxa against the nematode β-tubulin isotype 1 DNA reference sequences supplied in Avramenko et al., (2019). We chose to attribute a confidence level to taxonomic identifications at the species level: high or moderate confidence if three or two methods of assignation, respectively, were congruent. We also adjusted the sequence filtering based on an adapted procedure of Calderón-Sanou et al. (2020). We kept only ASVs present in at least 2 replicates of the same sample and removed ASVs that were not assigned to the genus level for the ITS2 and to the species level for β-tubulin isotype 1. We removed potential contaminants (reagent contaminants and cross-contaminations) following the procedure detailed in Calderón-Sanou et al., (2020). For each sample, we summed the reads of the two replicates with the highest similarity. If this similarity was lower than the mean similarity among all replicates, sample was discarded. Then, we verified that the 30 DNA extraction replicates had similar nemabiomes and kept one sample replicate out of two. At the end, we removed samples if they had <1000 reads of ITS2 and <500 reads of β-tubulin isotype 1 (Figure S3).

Identification of non-synonymous mutations in codons 167, 198 and 200

For each nematode species, all β-tubulin isotype 1 ASVs were aligned to one of the β-tubulin isotype 1 consensus sequences of the reference database (Avramenko et al., 2019) using the *AlignSeqs* function of the DECIPHER package (Wright, 2016). We examined each β-tubulin isotype 1 ASV at codons 167,198 and 200 to record whether the codon was associated with a non-synonymous mutation. However, we ignored the other polymorphic sites of the Exon.

Statistical analyses on measures of nemabiomes

We measured differences of nemabiomes among the two host species (sheep and ibex) and the 4 study sites (Aiguillette de Lauzet, Montagne de l'Oule, Champsaur and Belledonne). In the Cerces Massif, we considered independently the two sectors of the Aiguillette de Lauzet and the Montagne de l'Oule that correspond to distinct and distant pastures where sheep herds never met and where ibex feces were independently collected on these two pastoral units. To that end, we considered two measures of diversity, i.e., the alpha diversity and the beta diversity. The alpha diversity was measured with the Shannon index that considers richness and evenness of communities and the beta diversity was measured with the weighted UniFrac index estimated using the R phyloseq package (McMurdie and Holmes, 2013). The weighted UniFrac distance is a phylogenetic distance between the set of ASVs from each nemabiome weighted by the transformed abundance of each ASV (Lozupone and Knight, 2005). The phylogenetic distances were computed from a phylogenetic tree which was constructed using maximum likelihood with the GTR+G+I model according to the *ModelTest* function (Posada and Crandall, 1998; Schliep, 2011). The exact counts of ITS2 reads were transformed with the Hellinger transformation (e.g., square root of relative frequencies) to account for the high number of zeros in the community tables and to decrease the influence of rare ASVs in statistical analyses (Legendre and Legendre, 2012).

We tested the effects of host species and site, including their interaction, on alpha diversity with linear models, and on beta diversity using perMANOVA (*adonis2*, vegan R package (Oksanen et al., 2020)). All possible models including the null model were computed. For perMANOVA models, we used a custom function to compute Akaike's information criterion corrected for small sample size (AICc) based on residual sums of squares (Dyson, 2018). In a model selection approach, for both alpha and beta diversity, all possible models were ranked using the AICc and we selected the model with the lowest AICc value. Models with ΔAICc ≤ 2 were considered equivalent (Burnham and Anderson, 2002), and in this case, we considered the most parsimonious one, i.e., the model with the lowest degrees of freedom.

As sex and age were determined for some ibex, we also tested the effect of ibex classes (1: adult males, 2: females or kids/yearlings) on alpha and beta diversity

following the same model selection approach, including ibex classes and sites as explanatory variables.

All analyses were carried out using R 3.6 (R Core team, 2020. https://www.R-project.org/).

Statistical analyses on measures of resistant nematode strains

To compare the frequency of benzimidazole resistance in gastrointestinal nematodes between ibex and sheep and among the 3 sites, we tested if the host species, the site and the nematode species influenced the relative abundance of ASVs with a resistant allele. For this purpose, we used a generalized linear model with a binomial family and a model selection approach such as described above.

We used *AlignSeqs* (Wright, 2016) to generate multi-sequence aligned β-tubulin isotype 1 haplotype data. For each gastrointestinal species, we removed short ASVs, e.g., ASVs with a sequence length <10% compared to the median ASV length. PopART v1.7 (Leigh and Bryant, 2015) was used to draw median joining networks based on the haplotype data of each gastrointestinal nematode species.

Results

Parasite material

The median number of eggs of strongyles per gram of feces were lower in sheep ($7.5[0,148]_{95\%IQR}$; n = 90) than in ibex ($15[0,163]_{95\%IQR}$; n = 167) feces (Mann–Whitney U test; $W = 9043.5$, $P = 0.006$). As a result of the low level of infestation in some samples, the number of L3 hatched from eggs were not sufficient (n<20) for 48 samples. These samples were not used for subsequent genetic investigations. Specifically, all samples from the Sept Laux sheep herd (n=15, Belledonne) were discarded. Therefore, the nemabiome was determined based on the ITS2 for 196 (n= 55 sheep and n=141 ibex) out of 209 samples for which DNA was extracted (Figure S1).

Diversity of gastrointestinal nematodes in sheep and ibex

In total, we detected 408 ASVs corresponding to 13 gastrointestinal nematode species (Table 3, Figure S4). Eight ASVs were assigned to the genus level (i.e., *Marshallagia* spp., *Nematodirus* spp. and *Trichostrongylus* spp.) due to non-identical assignation among taxonomic methods. An ASV corresponding to the lungworm *Cystocaulus ocreatus* was discarded for the statistical analyses because we only focus on gastrointestinal nematodes.

Teladorsagia circumcincta was the most prevalent nematode species and was detected in 85% of samples (90%, n=127/141 ibex and 71%, n=39/55 sheep), followed by *Trichostrongylus vitrinus,* 63% (73%, n=103/141 ibex and 36%, n=20/55 sheep) and *Haemonchus contortus,* 56% (70%, n=98/141 ibex and 22%, n=12/55 sheep) (Table 3, Figure S4). *Nematodirus* spp. and *Ostertagia leptospicularis* were the rarest species and were detected in only 2 and 1 sample, respectively and with a very low relative frequency (<0.1%).

The parasite of the genus *Nematodirus* were not considered for the following results as our coproculture protocol is not the most appropriate for these parasites (i.e., some species need more than 2 weeks to reach the L3 stage and need to be exposed to cold temperatures before coproculture, which can be deleterious for other strongyle species such as *Haemonchus contortus*, van Wyk and Mayhew, 2013).

Table 3 : Number of ASVs, reads and samples for each nematode taxon, including the results from the ITS2 (nemabiome) and the β-tubulin isotype 1 (resistance to benzimidazole). N= number (and percentage) of samples in which a taxon was detected. Host species and study sites are mixed here.

	ITS2			β-tubulin isotype 1		
	ASVs	Reads	N (%)	ASVs	Reads	N (%)
Bunostomum trigonocephalum	8	61 631	12 (6%)	-	-	-
Chabertia ovina	19	95 585	62 (32%)	-	-	-
Cooperia curticei	4	803	6 (3%)	-	-	-
Cooperia fuelleborni	2	358	5 (3%)	-	-	-
Haemonchus contortus	47	1 937 587	110 (56%)	38	392 110	96 (62%)
Marshallagia marshalli	23	338 064	57 (29%)	-	-	-
Marshallagia spp.	2	404	2 (1%)	-	-	-
Oesophagostomum venulosum	13	311 530	82 (42%)	-	-	-
Ostertagia leptospicularis	1	5	1 (1%)	-	-	-
Ostertagia ostertagi	2	269	2 (1%)	-	-	-
Teladorsagia circumcincta	235	3 311 094	166 (85%)	310	343 551	145 (94%)
Trichostrongylus axei	26	462 936	103 (53%)	33	68 414	104 (68%)
Trichostrongylus colubriformis	12	411 776	102 (52%)	8	144 643	112 (73%)
Trichostrongylus vitrinus	9	521 817	123 (63%)	44	92 834	107 (69%)
Trichostrongylus spp.	5	71 124	70 (36%)	-	-	-

According to the model selection approach, host species was the only factor that explained the alpha diversity (Table S1). The model indicated that sheep had a lower alpha diversity compared to ibex (β = -0.42 ± 0.07, *P* < 0.001, R² of the model=0.15) (Figure 2b). Site was not retained in the selected model (Table S1).

Beta diversity was best explained by both factors, host species ($F_{1,188}$ = 27.69, *P* = 0.001) and site ($F_{3,188}$ = 16.39, *P* = 0.001) and their interaction ($F_{3,188}$ = 25.61, *P* = 0.001) according to model selection results (Table 4, Table S2). Some gastrointestinal nematodes were mostly found in sheep feces or only in ibex feces (Figure 2a). *Trichostrongylus colubriformis* was more frequent in the Ecrins national park (mean RRA: 10% [6%; 13%]$_{95CI}$) than in Belledonne (mean RRA: 4% [2%; 6%]$_{95CI}$). Likewise,

Marshallagia spp. was more frequent in ibex feces in Cerces and Champsaur mountains (mean RRA: 11% [6%;15%]$_{95CI}$) than in ibex feces in Belledonne (0.4% [-0.2%; 1%]$_{95CI}$). The distribution of *Haemonchus contortus* in host species and sites had a particular pattern. This parasite was more frequent in ibex feces compared to sheep feces in Belledonne (mean RRA: 0.004% in sheep; 48% in ibex) and Champsaur (mean RRA: 0% in sheep; 41% in ibex), while the opposite was observed in the Cerces (Aiguillette du Lauzet: mean RRA of 7% in sheep and 0.9% in ibex; Montagne de l'Oule: mean RRA of 30% in sheep and 0% in ibex).

The model selection approach retained the effects of site and class of individuals and the interaction between the two independent factors in the best model explaining alpha diversity (Table S3). We found significant differences among sites : Shannon index of alpha diversity was higher in Belledonne ($\beta = 0.62 \pm 0.16$, $P < 0.001$, R^2 of the model=0.27) and Champsaur ($\beta = 0.55 \pm 0.16$, $P < 0.001$) compared to the Aiguillette du Lauzet. The diversity of nematodes was higher also in males compared to females/yearlings ($\beta = 0.60 \pm 0.15$, $P < 0.001$), except in Champsaur where males present a lower alpha diversity than females with kids ($\beta = -0.52 \pm 0.20$, $P < 0.001$). For the beta diversity, the best model included only the site (($F_{2,62} = 15.93$, $P = 0.001$, Table S4).

Table 4 : Parameters estimated from the best PerMANOVA model explaining the beta diversity in ibex and sheep. The effect of host species (ibex or sheep) and site (Aiguillette de Lauzet, Montagne de l'Oule, Champsaur or Belledonne) and the interaction between the two factors are reported. Partial R^2 are reported with the corresponding *F*-value and *p value* (P).

Diversity index	Best model selected	Variables	partial R^2	F-value	P
Weighted UniFrac	$\beta \sim$ Site x Host species	Residuals	0.55	-	-
		Site	0.14	16.39	0.001
		Host species	0.08	27.69	0.001
		Site: Host species	0.22	25.61	0.001

Figure 2 : (a) Mean relative frequencies of gastrointestinal nematode species, (b) Shannon diversity of ITS2 ASVs and (c) prevalence and mean relative frequencies of β-tubulin isotype 1. Results and sample size (N_{ITS} and $N_β$) are presented for each host species (sheep or ibex) in each site (Cerces: Montagne de l'Oule and Aiguillette de Lauzet; Champsaur and Belledonne: Ane Buyant (AB) and La Pesée (LP)). On panel (c), the size of the pie chart corresponds to the prevalence of the corresponding gastrointestinal nematode species in the population, and the size of each slice to the mean proportion of each allele.

Anthelmintic resistance

We found 433 different β-tubulin isotype 1 ASVs in 154 (n= 39 sheep and n=115 ibex) out of 209 samples for which DNA was extracted. Among the 5 gastrointestinal nematode species targeted by specific primers, we detected, *Haemonchus contortus* in 96 samples, *Teladorsagia circumcincta* in 145 samples, *Trichostrongylus axei* in 104 samples, *Trichostrongylus vitrinus* in 107 samples and *Trichostrongylus colubriformis* in 112 samples (Table 3, Figure S5). No resistance mutation was detected for *Trichostrongylus vitrinus.* Therefore, *Trichostrongylus vitrinus* was not included in the model explaining the relative abundance of resistant reads.

Resistance mutations were highly frequent (93.5%; n=144/154) with only 10 ibex feces (3 from Belledonne and 7 from Champsaur) in which no resistant mutation was detected. Based on the best model for resistant RRA, the frequency of resistant nematodes depended on gastrointestinal nematode species and the interaction between host species and the study site (Table 5, Table S5). *Teladorsagia circumcincta* was the species with the lowest resistant RRA and *Trichostrongylus colubriformis* had the highest resistant RRA (Table 5). The mean observed RRA of resistant nematodes differed between gastrointestinal nematode species (*Haemonchus contortus:* 19% $[13;25]_{95CI}$; *Teladorsagia circumcincta:* 4% $[3;6]_{95CI}$; *Trichostrongylus axei:* 70% $[63;78]_{95CI}$; *Trichostrongylus colubriformis:* 96% $[93;99]_{95CI}$). Resistant RRA were generally lower in ibex compared to sheep (β = -2.21 ± 0.66, *P* < 0.001). Resistant RRA were the lowest in the Aiguillette de Lauzet but this was also the only site were ibex had a significantly higher resistant RRA compared to sheep (Table 5). We found no significative effect of ibex classes (males or females and kids/yearlings) on benzimidazole resistance frequencies ($F_{1,49}$=0.03, P = 0.863, ANOVA test).

The most frequent resistance mutation was the F200Y, present in the 4 gastrointestinal nematode species and 141 fecal samples, followed by the E198A (46 samples, 2 nematode species: *Haemonchus contortus* and *Teladorsagia circumcincta*), the F167Y (17 samples, 2 nematode species: *Haemonchus contortus* and *Teladorsagia circumcincta*), the E198L (7 samples, *Teladorsagia circumcincta*) and the E198G (4 samples, *Haemonchus contortus*) (Figure 2c).

Table 5 : Parameter estimates for the best generalized linear model explaining the resistant reads relative abundance (RRA) in ibex and sheep. The effect of host species (sheep as reference), study sites (Belledonne as reference), and their interaction, in addition to the nematode species (*Teladorsagia circumcincta* as reference) are reported. Parameter estimates with standard error (SE) are reported with the corresponding *z-value (z*-val*)* and *p value* (*P*). AL: Aiguillette de Lauzet; MO: Montagne de l'Oule; Ch: Champsaur; Hc : *Haemonchus contortus*; Ta: *Trichostrongylus axei*; Tcol: *Trichostrongylus colubriformis*.

Best model selected	Variables	Parameter estimate ± SE	z-val	*P*
Resistant RRA ~ host species x study sites + nematode species	Intercept	-2.02 ± 0.59	-3.45	**5e-04**
	Species	-2.21 ± 0.66	-3.35	**8e-04**
	Mountain(AL)	-2.89 ± 0.99	-2.93	**0.003**
	Mountain(MO)	-1.37 ± 0.89	1.54	0.123
	Mountain(Ch)	-0.10 ± 1.18	-0.08	0.935
	Nematode(Hc)	2.52 ± 0.62	4.05	**5e-05**
	Nematode(Ta)	4.68 ± 0.59	7.95	**1e-15**
	Nematode(Tcol)	7.38 ± 0.75	9.79	**<2e-16**
	Species:Site(AL)	2.90 ± 1.19	2.44	**0.015**
	Species:Site(MO)	0.04 ± 1.28	0.03	0.973
	Species:Site(Ch)	-0.75 ± 1.32	-0.56	0.574

Sheep and ibex shared 164 (38%) β-tubulin isotype 1 haplotypes and 238 (55%) β-tubulin isotype 1 haplotypes were only found in ibex samples (Figure 3). Most of the resistant haplotypes of *Teladorsagia circumcincta* and *Haemonchus contortus*, e.g., containing a non-synonymous mutation at the codon 167, 198 or 200, were genetic variants of a common sensitive haplotype shared by ibex and sheep (Figure 3). The resistant haplotypes of *Trichostrongylus axei* and *Trichostrongylus colubriformis* were more common than the sensitive haplotypes and the most similar sensitive haplotypes were found either in both sheep and ibex samples, or only in ibex samples. Both, *Trichostrongylus colubriformis* and *Trichostrongylus vitrinus* showed two distinct lineages, separated by ≥10 mutations. One of the lineages of *Trichostrongylus vitrinus* was only found in ibex from Belledonne and an ibex from Champsaur (Figure 3) while the ASVs of the second lineage were found both in ibex and sheep. One of the lineages of the *Teladorsagia circumcincta* was more diverse (Figure 3).

Figure 3 : Median joining network of β-tubulin isotype 1 haplotypes. Each point represents a unique haplotype, and the colors correspond to the host species in which the haplotype was detected. The size of the point is proportional to the number of samples in which the haplotype was found. S: sensitive haplotype, R: resistant

592   haplotype. The tag above the points indicates the name of the mutation, based on the
593   codon position and the substitution of the amino acid.

Discussion

Because resident Alpine ibex use pastures grazed by transhumant sheep during
summer, but not concurrently (unpublished data), we sought to assess the extent of
nematode sharing between these two host species. Specifically, we investigated the
presence of anthelmintic-resistant nematode strains in sheep and ibex to determine
the role of transhumant sheep in contaminating alpine pastures, and whether ibex may
play a role in the circulation and maintenance of anthelmintic resistant nematodes. We
used a metabarcoding approach based the sequencing of ITS2 and β-tubulin to
demonstrate that both sheep and ibex were infected by the same gastrointestinal
nematode species and shared anthelmintic-resistant strains, despite the absence of
sheep on alpine pastures for much of the year and therefore a narrow temporal window
for contamination.

In line with other studies investigating the gastrointestinal nematodes of sheep and
ibex (Burgess et al., 2012; Gruner et al., 2006; Redman et al., 2019; Zaffaroni et al.,
2000), the most prevalent and abundant species in both host species was
Teladorsagia circumcincta. Next, *Trichostrongylus vitrinus* was moderately prevalent,
but not abundant in sheep or ibex nemabiomes. These results are in accordance with
the year-round climatic conditions as these two nematode species, as well as
Marshallagia spp., are better adapted to cold temperatures than the other nematode
species detected in this study (O'Connor et al., 2006; Zaffaroni et al., 2000). The
studied sheep flocks originate from the French plains and/or the south of France and
are driven into mountain areas in early summer. Consequently, their nemabiome at the
time of sampling is representative of the gastrointestinal nematode communities
present in sheep on the farm, i.e., prior to transhumance. In a similar context, Gruner
et al., (2006) observed a high prevalence of *Teladorsagia circumcincta* in two of three
transhumant sheep flocks at the beginning of the grazing season in the southern Alps.
Furthermore, transhumant sheep flocks appear to ingest mainly *Teladorsagia*
circumcincta when grazing in the mountains, as this parasite remains the dominant
species identified in feces and tracer lambs during the summer (Gruner et al., 2006).
Pastoral activity in mountainous areas of France could therefore favor nematode

species more adapted to cool and wet environmental conditions, such as *Teladorsagia circumcincta* (O'Connor et al., 2006), compared with sheep grazing on the plains year round. To confirm this hypothesis, the nemabiome of transhumant sheep should be compared with the nemabiome of resident sheep that stay in farm all year around.

A high relative frequency (>30%) of *Haemonchus contortus* was detected in ibex in Belledonne and Champsaur. In contrast, almost no *Haemonchus contortus* was observed in sheep flocks driven to these mountains, raising the possibility that ibex may be contributing to the infection of sheep with *Haemonchus contortus*. To our knowledge, this is the first time that such a relative abundance of *Haemonchus contortus* is reported in Alpine ibex (see previous studies based on morphological identification: Carcereri et al., 2021; Marreros et al., 2012; Zaffaroni et al., 2000). In addition, it should be noted that Alpine ibex were sampled before a potential contamination by domestic sheep could be detected, i.e., before the end of the pre-patent period (time between infection and egg production) and at the end of spring – early summer, i.e., the start of the epidemiological period for *Haemonchus contortus* infection in high-altitude mountain areas. We can therefore expect higher levels of contamination in late summer, when domestic sheep leave mountain pastures. Nonetheless, we cannot exclude that some laboratory issues might have reduced the apparent prevalence and abundance of *Haemonchus contortus* as some samples from sheep were kept in the fridge at 4°C during 2 to 3 days (including e.g., the sheep samples without *Haemonchus contortus* from Belledonne and Champsaur) which could have reduced the proportion of *Haemonchus contortus* eggs hatching (McKenna, 1998).

The detection of Haemonchus *contortus* raises conservation issues for Alpine ibex as this nematode species is known to be highly pathogenic sheep (Taylor et al., 2015). Infection of a phylogenetically related species, the Pyrenean ibex (*Capra pyrenaica pyrenaica*), with a few thousand *Haemonchus contortus* resulted in severe clinical signs, including extremely low weight and hemorrhagic anemia (Lavín et al., 1997). In addition, *Haemonchus contortus* may have contributed to the collapse of the Northern Chamois, *Rupicapra rupicapra*, in the province of Lecco, Italy from November 2000 to March 2001 (Citterio et al., 2006). As gastrointestinal nematodes can have an impact on the demographic dynamics of the host population (Acerini et al., 2022; Albery et al., 2021; Albon et al., 2002), they are suspected of being behind the low natality rates observed in the French Alpine ibex populations (Brambilla et al., 2020). While Alpine

ibex appear to be fairly resilient to parasite infections (Marreros et al., 2012), further
investigations should be carried out to assess the consequences of gastrointestinal
nematode infections for ibex at both individual and population levels.

Several of the nematode species we found are common to other ungulate species
present in the study areas (Mediterranean mouflon, *Ovis gmelini musimon ✕ Ovis sp.*;
Northen chamois; domestic goat, *Capra hircus*; red deer*, Cervus elaphus*; and roe
deer, *Capreolus capreolus*) (Zaffaroni et al., 2000). In our study, only a few domestic
goats were present in Belledonne (n = 11 individuals) and in Champsaur (n =5
individuals) and represented less than 0.01% of the domestic flock in these areas. As
we collected feces directly on the ground, we cannot exclude that goat feces were
collected instead of sheep feces. In our opinion, domestic goats should not have a
significative influence on nemabiome of ibex in our study area considering the scarcity
of the species. In future analyses, we should consider the different domestic and wild
ungulate species living in the same study area. Especially because they have different
space use, different nemabiomes and could provide key information to better
understand the dynamic of nematode exchanges among domestic and wild ungulates.

We found a higher diversity of nematodes in adult males compared to females and
kids/yearlings. Sexual parasitism towards males is commonly observed in vertebrates
and ungulates in particular (Klein, 2000; Martínez-Guijosa et al., 2015; Oliver-Guimerá
et al., 2017, but see Beaumelle et al., 2021; Bourgoin et al., 2021). This parasitism
towards males is generally explained by both hormonal and behavioral differences
between the two sexes. Generally, males tend to allocate more energy to the
development of traits influenced by testosterone, such as secondary sexual
characteristics (e.g. the length of horns in ungulates) or courtship displays (e.g. male
aggression for mating opportunities). It is important to note that while testosterone is
necessary for the development of these secondary sexual characteristics in males,
high levels of the testosterone have also been linked to an altered immune system
(Klein, 2004), leading to increased parasitism. In addition, ibex segregate by sex
(Brambilla et al., 2022), providing less opportunities for intersexual transmission of
parasites. Contrary to females and kids, males feed on patches grazed by domestic
sheep before the grazing period (Margaillan, 2021), increasing the probability of
infection of males by over-wintering nematodes deposited by livestock during the
previous transhumance (O'Connor et al., 2006). However, we did not notice any
difference between classes concerning benzimidazole resistance frequencies.

However , more information regarding the spatial distribution of both sexes are required
if we want to investigate further the susceptibility of different sexes or age classes and
its importance in the spread and exchange parasites with domestic livestock (Bourgoin
et al., 2021).

We detected anthelmintic resistant alleles in 4 out of the 5 nematode species
tested, namely *Haemonchus contortus*, *Teladorsagia circumcincta*, *Trichostrongylus*
axei, *Trichostrongylus colubriformis*, but not in *Trichostrongylus vitrinus*. Both sheep
and ibex hosted resistant strains of the 4 nematode species and only 10 out of 116
ibex carried only susceptible strains. The benzimidazole resistance was therefore very
common in both host groups, in agreement with the situation of sheep farms in Europe
(Rose et al., 2015; Rose Vineer et al., 2020). The presence of anthelmintic resistant
nematodes in ibex is most likely explained by the indirect transmission of resistant
nematodes from sheep to ibex through the deposit of infected feces by sheep on
pastures. The large number of shared β-tubulin ASVs between sheep and ibex and the
high overlap between their nemabiomes tend to confirm this hypothesis (Figure 2c,
Figure 3). This is also in accordance with other studies investigating nematode
parasites exchange at the interface between wild and domestic ungulates (Beaumelle
et al., 2022; Cerutti et al., 2010; Laca Megyesi et al., 2019). Another pathway for the
contamination of ibex by resistant nematodes could be the sublethal exposure of the
free-living stages of nematodes to anthelmintic residues excreted in the environment
by treated sheep, which can select for anthelmintic resistance in situ (Dimunová et al.,
2022). Whereas sheep are generally treated just before their ascent to the mountain
pastures, excretion of anthelmintic drugs via sheep feces can occur during several
days after the administration and the molecule degradation last days, or even months
(Kolar et al., 2006). Unfortunately, the level of drugs in the environment, their
persistence and their spread in grazed mountainous area are totally unknown.
Environmental circulation of anthelmintic residues should be investigated in further
studies to understand its incidence on the presence of resistant nematodes in wildlife.

It is worth noting that feces of ibex were sampled before the arrival of sheep on
pastures. This demonstrates that anthelmintic resistant nematodes can be maintained
in mountainous areas from year to year in wild populations of ibex despite harsh winter
environmental conditions, and in the absence of the main source of parasites during
most of the year, i.e., the domestic sheep. The shedding of eggs from resistant
nematodes by ibex prior to the arrival of domestic sheep suggests the potential role of

ibex as a reservoir of anthelmintic resistant nematodes for other susceptible domestic
and wild ungulates. Once resistant strains have been selected, the absence of
selection pressure (i.e. absence of the use of anthelmintics) does not guarantee the
reversion of resistance (Hamilton et al., 2022; Leathwick et al., 2015). Consequently,
ibex may maintain benzimidazole-resistant strains for several years even in the
absence of selection pressure. In addition, the position of resistant mutant strains
detected at the periphery of haplotype networks (Figure 3) supports the lack of
benzimidazole resistant reversions and relatively recent selection of benzimidazole
resistance. The recent selection of resistance could result from the repeated use of
benzimidazole and also perhaps from the presence of anthelmintic drug residues in
the environment which maintain a selection pressure for gastro-intestinal nematodes
(Dimunová et al., 2022).

The 5 nematode species studied seemed to have different selection dynamics
which may reflect life history traits (Redman et al., 2015). In fact, we detected no
resistant allele in *Trichostrongylus vitrinus* and conversely, the proportion of
benzimidazole resistant strains of *Trichostrongylus axei* and *Trichostrongylus*
colubriformis were high in sheep, i.e., between 40% and 100% and slightly lower in
ibex, i.e., 33-100% (Figure 2c). These results are in line with observations in the
literature. Strains of Trichostrongylus vitrinus resistant to benzimidazoles were also
rare in other studies of sheep farms, e.g., in the UK (Avramenko et al., 2019), and in
Canada (Queiroz et al., 2020). Conversely, higher frequencies of benzimidazole
resistance in *Trichostrongylus axei* and *Trichostrongylus colubriformis* were already
reported in sheep, e.g., in UK (*Trichostrongylus axei*: 26-27% and *Trichostrongylus*
colubriformis: 53-62%; Avramenko et al., 2019), in Austria (*Trichostrongylus*
colubriformis: 77%-100%; Hinney et al., 2020), or in France, (*Trichostrongylus axei*:
63%; Palcy et al., 2010). Furthermore, we found a low proportion of resistance in
Teladorsagia circumcincta both in sheep and ibex (mean ± standard error of the mean:
6.6 ± 3.5%), in comparison for example with Trichostrongylus axei and
Trichostrongylus colubriformis, (Figure 2c). This contrast with previous results from the
literature, where a higher frequency of resistance allele was found for *Teladorsagia*
circumcincta: 32.4 ± 6.8% in Hinney et al. (2020) and ~66% in Avramenko et al. (2019).
Regarding *Haemonchus contortus*, we found high variability of resistance frequencies
among sheep flocks (69.9 ± 14.4%). Our results are similar to the observations of
Avramenko et al. (2019), but contrast with those of Hinney et al. (2020) who detected

high resistance frequencies for *Haemonchus contortus* (91.9 ± 3.7%) in transhumant sheep flocks in the Austrian Alps. In addition, *Haemonchus contortus* was the parasite species with the highest diversity in terms of non-synonymous mutations. Besides the commonly found polymorphisms at codons 167 and 200 (Avramenko et al., 2019; Redman et al., 2015), we detected several mutations at codon 198, occurring in particular with high frequency on the sheep farm in Montagne de l'Oule. In contrast to our study, previous research had found polymorphisms at codon 198, but they were rare (Francis and Šlapeta, 2023; Ramünke et al., 2016).

Several factors are suspected to contribute to interspecific differences in the selection of resistance strains among nematode species, including specific reproductive rates, seasonal dynamics, exposure to anthelmintic in the digestive tract, drug effective dose and fitness cost of benzimidazole resistance, and climatic conditions in the location of sheep farms, anthelmintic strategies, e.g., treatment molecules, timing and rate of anthelmintic treatments and grazing management (Hodgkinson et al., 2019; Redman et al., 2015). However, the links between parasite traits and interspecific variation of resistance acquisition by gastrointestinal nematodes has not been tested yet (Morgan et al., 2019). Our results suggest a few clues in relation to the ecology of the nematode species.

Firstly, nematode species have different abilities to practice hypobiosis, i.e., the ability to halt embryonic development under environmental constraints (Gibbs, 1986). *Haemonchus contortus* and *Teladorsagia circumcincta* are known to arrest development more frequently than *Trichostrongylus* spp. (Langrová et al., 2008), and hypobiotic larvae have been shown to be less sensitive to drugs (Sargison et al., 2007). Secondly, among the *Trichostrongylus* spp., *Trichostrongylus vitrinus* may have a higher proportion of overwintering larvae in pastures as this species is more resistant to cold temperature compared to *Trichostrongylus axei* and *Trichostrongylus colubriformis* (O'Connor et al., 2006). As parasites on pastures are less subject to selection pressure by anthelmintics, they are a source of susceptible strains.

As the proportion of resistant strains is generally lower in ibex compared to sheep, ibex may contribute to a dilution effect of resistant strains, i.e., by hosting susceptible nematodes. However, the role of ibex in the maintenance of a refugia needs to be investigated by considering the relative number of susceptible strains deposited by ibex on a pasture compared with sheep. Furthermore, it seems that the role of ibex in the maintenance of a refugia may vary according to nematode species. For example,

ibex excrete a lower proportion of *Trichostrongylus axei* eggs than sheep (Figure 2a),

but containing largely resistant strains (Figure 2c). In contrast, *Teladorsagia*

circumcincta and *Haemonchus contortus* in ibex were more frequently susceptible and

genetically diverse (higher number of ITS2 and β-tubulin ASVs) compared with

Trichostrongylus axei and *Trichostrongylus colubriformis* ibex (Table 3, Figure 3). As

Teladorsagia circumcincta was dominant in ibex, a refuge of susceptible *Teladorsagia*

circumcincta strains may be maintained within ibex and may contribute to limiting the

spread of resistance in sheep farms. Nematodes can negatively or positively interact

within the host gut, and interactions between species or between strains may have

important implications for the selection of resistance. However, the magnitude of

within-host interactions between nematode strains/species and its implication in the

management of resistance remains to be determined (Hellard et al., 2015; Lello et al.,

2004).

Differences in patterns among massifs were observed at the community and

genetic level among sheep flocks and ibex populations. Indeed, it was expected that

some differences in nemabiome composition would be observed between the massifs

and sheep flocks, given that sheep and ibex from different areas never meet (R. Papet,

C. Toïgo, E. Vannard, Pers. communication). Furthermore, the sheep flocks come from

different locations and have been subjected to different anthelmintic strategies. For the

ibex, differences in the original population of translocated animals (Gauthier and

Villaret, 1990; Kessler et al., 2022) and potential founder effects – not all parasites

present in the source population were present in the newly established population -

may have had a long term impact on the composition of the nemabiome. For example,

the distinct population of *Trichostrongylus vitrinus*, found mainly in ibex in Belledonne,

may have been inherited from the founding ibex population. This highlights that various

reintroductions of ibex in the study area can also influence the composition of the

parasite community. This distinct population *of Trichostrongylus vitrinus* was absent in

sheep grazing in Belledonne, which hosted other strains of *Trichostrongylus vitrinus*

despite summers of co-occurrence of sheep and ibex in Belledonne. It is possible that

this strain is adapted to ibex and incapable of developing in sheep or that this strain is

highly sensitive to antiparasitic treatments used in sheep farms and is systematically

eliminated when sheep are treated. but this observation may also be due to a sampling

bias as the number of sheep sampled remains low. Within the same mountain area,

few differences were observed among the nemabiomes of ibex, e.g., between the ibex

of the Aiguillette de Lauzet and those of the Montagne de l'Oule, whereas the sheep
flocks hosted distinct nemabiome communities.

In conclusion, transmission of gastrointestinal nematode species, including
resistant nematode strains, occur between sheep and ibex even though the contact
between the two species is limited to the summer period. In this study, we
demonstrated more specifically that ibex can maintain and shed eggs of resistant
gastrointestinal nematodes despite the absence of sheep on pastures for several
months, suggesting a potential role of ibex as a reservoir for these nematodes.
However, the extent to which each host species can influence the nematode
community of the other during the transhumant period remains to be determined. To
this end, a temporal sampling, before, during and after the different host species share
the same pasture should be considered. Analysis of parasite population structure using
appropriate genetic markers (i.e., microsatellites or SNPs) could also help to quantify
gene flow between ibex and sheep nematode populations (Cerutti et al., 2010). In
addition, intervention studies could be employed to infer the role of ibex in maintaining
nematode populations shared between the two host species (Viana et al., 2014).
Experimental infections of captive ibex or monitoring free-ranging ibex populations
after access to alpine pastures has been restricted to livestock should help us to refine
the ability of ibex to maintain nematodes from domestic ungulates, including resistant
nematodes. Finally, epidemiological models could be useful tools to better understand
the dynamics of resistant parasites at the livestock-wildlife interface (Brown et al.,
2022; Dickinson et al., 2024). The lower proportion of resistance alleles in ibex
compared to sheep underlines the possibility that ibex could contribute to the
maintenance and circulation of susceptible strains in sheep. Based on our results, it
seems that ibex have helped to limit the spread of anthelmintic resistance in
Teladorsagia circumcincta and *Haemonchus contortus* in sheep flocks, by maintaining
parasite refugia not exposed to anthelmintic pressure. As with roe deer (Beaumelle et
al., 2022), domestic sheep contribute to the modification of the nemabiome of ibex.
This raises concerns about ibex conservation, and the consequences of strongyle
infection in ibex should be investigated. Indeed, ibex are characterized by low genetic
diversity due to the strong demographic decline of this species followed by multiple re-
introductions (Grossen et al., 2018). The high genetic structure of immunity-related loci
among ibex populations (Kessler et al., 2022) raises additional concerns, as both

neutral and adaptive genetic diversity are known to have an influence on parasite resistance and tolerance in ungulates (Portanier et al., 2019).

Appendices

Supplementary data to this article can be found online in MendeleyData (DOI: 10.17632/cm97cg87d6.1).

Acknowledgements

The authors warmly thank all the professionals from the Office Français de la Biodiversité and all the trainees for data collection, J. S. Gilleard and E. Redman from the University of Calgary, C. Lionnet from the Laboratoire d'Ecologie Alpine (LECA) and other members of the lab to help us developing the deep sequencing analyses for nematodes. The research benefited from the support of AnaBM (USMB) and AEEM (UGA) laboratory facilities.

Funding

This project was founded by the Office Français de la Biodiversité, the Laboratoire d'Ecologie Alpine (LECA) and VetAgro Sup - Pôle d'Expertise Vétérinaire et Agronomique des Animaux Sauvages (EVAAS, France; http://evaas.vetagro-sup.fr/; DGAL—VetAgro Sup - INRAE funding). G. Bourgoin was supported by the AgreenSkills+ fellowship program (European Union program; MarieCurie FP7 COFUND People Programme; grant agreement n_609398).

Conflict of interest disclosure

The authors declare that they have no conflict of interest.

Data, scripts, and supplementary information availability

The bioinformatic pipeline, the ASV analysed during the current study and the R script of statistic analyses are available in MendeleyData (DOI: 10.17632/cm97cg87d6.1).

References

Acerini, C.I., Morris, S., Morris, A., Kenyon, F., McBean, D., Pemberton, J.M., Albery, G.F., 2022. Helminth parasites are associated with reduced survival probability in young red deer. Parasitology 149, 1702–1708. https://doi.org/10.1017/S0031182022001111

Acevedo, P., Cassinello, J., Gortazar, C., 2008. The Iberian ibex is under an expansion trend but displaced to suboptimal habitats by the presence of extensive goat livestock in central Spain, in: Hawksworth, D.L., Bull, A.T. (Eds.), Biodiversity and Conservation in Europe, Topics in Biodiversity and Conservation. Springer Netherlands, Dordrecht, pp. 119–134. https://doi.org/10.1007/978-1-4020-6865-2_9

Albery, G.F., Morris, A., Morris, S., Kenyon, F., Nussey, D.H., Pemberton, J.M., 2021. Fitness costs of parasites explain multiple life-history trade-offs in a wild mammal. Am. Nat. 197, 324–335. https://doi.org/10.1086/712633

Albon, S.D., Stien, A., Irvine, R.J., Langvatn, R., Ropstad, E., Halvorsen, O., 2002. The role of parasites in the dynamics of a reindeer population. Proc. R. Soc. Lond. B Biol. Sci. 269, 1625–1632. https://doi.org/10.1098/rspb.2002.2064

Avramenko, R.W., Redman, E.M., Lewis, R., Yazwinski, T.A., Wasmuth, J.D., Gilleard, J.S., 2015. Exploring the gastrointestinal "Nemabiome": deep amplicon sequencing to quantify the species composition of parasitic nematode communities. PLoS One 10(12), e0143559. https://doi.org/10.1371/journal.pone.0143559

Avramenko, R.W., Redman, E.M., Melville, L., Bartley, Y., Wit, J., Queiroz, C., Bartley, D.J., Gilleard, J.S., 2019. Deep amplicon sequencing as a powerful new tool to screen for sequence polymorphisms associated with anthelmintic resistance in parasitic nematode populations. Int. J. Parasitol. 49, 13–26. https://doi.org/10.1016/j.ijpara.2018.10.005

Beaumelle, C., Redman, E., Verheyden, H., Jacquiet, P., Bégoc, N., Veyssière, F., Benabed, S., Cargnelutti, B., Lourtet, B., Poirel, M.-T., de Rijke, J., Yannic, G.,

Gilleard, J.S., Bourgoin, G., 2022. Generalist nematodes dominate the nemabiome of roe deer in sympatry with sheep at a regional level. Int. J. Parasitol. https://doi.org/10.1016/j.ijpara.2022.07.005

Beaumelle, C., Redman, E.M., de Rijke, J., Wit, J., Benabed, S., Debias, F., Duhayer, J., Pardonnet, S., Poirel, M.-T., Capron, G., Chabot, S., Rey, B., Yannic, G., Gilleard, J.S., Bourgoin, G., 2021. Metabarcoding in two isolated populations of wild roe deer (*Capreolus capreolus*) reveals variation in gastrointestinal nematode community composition between regions and among age classes. Parasit. Vectors 14, 594. https://doi.org/10.1186/s13071-021-05087-5

Biber, J.-P., 2010. Transhumance in France. Pastoralism 1, 91–98. https://doi.org/10.3362/2041-7136.2010.006

Blaxter, M., Koutsovoulos, G., 2015. The evolution of parasitism in Nematoda. Parasitology 142, S26–S39. https://doi.org/10.1017/S0031182014000791

Bordes, L., Dumont, N., Lespine, A., Souil, E., Sutra, J.-F., Prévot, F., Grisez, C., Romanos, L., Dailledouze, A., Jacquiet, P., 2020. First report of multiple resistance to eprinomectin and benzimidazole in *Haemonchus contortus* on a dairy goat farm in France. Parasitol. Int. 76, 102063. https://doi.org/10.1016/j.parint.2020.102063

Bourgoin, G., Portanier, E., Poirel, M.-T., Itty, C., Duhayer, J., Benabed, S., Cockenpot, A., Callait-Cardinal, M.-P., Garel, M., 2021. Reproductive females and young mouflon (*Ovis gmelini musimon × Ovis* sp.) in poor body condition are the main spreaders of gastrointestinal parasites. Parasitology 148, 809–818. https://doi.org/10.1017/S0031182021000329

Boyer, F., Mercier, C., Bonin, A., Le Bras, Y., Taberlet, P., Coissac, E., 2016. obitools: a unix-inspired software package for DNA metabarcoding. Mol. Ecol. Resour. 16, 176–182. https://doi.org/10.1111/1755-0998.12428

Brambilla, A., 2020. Caratterizzazione genetica delle popolazioni di stambecco delle Alpi occidentali. Department of Evolutionary Biology and Environmental Studies, Zurich.

Brambilla, A., Bassano, B., Biebach, I., Bollmann, K., Keller, L., Toïgo, C., von Hardenberg, A., 2022. Alpine Ibex *Capra ibex* Linnaeus, 1758, in: Corlatti, L., Zachos, F.E. (Eds.), Terrestrial Cetartiodactyla, Handbook of the Mammals of Europe. Springer International Publishing, Cham, pp. 383–408. https://doi.org/10.1007/978-3-030-24475-0_32

Brambilla, A., Von Hardenberg, A., Nelli, L., Bassano, B., 2020. Distribution, status, and recent population dynamics of Alpine ibex *Capra ibex* in Europe. Mammal Rev. 50, 267–277. https://doi.org/10.1111/mam.12194

Brivio, F., Ciuti, S., Pipia, A., Grignolio, S., Apollonio, M., 2022. Livestock displace European mouflon from optimal foraging sites. Eur. J. Wildl. Res. 68, 30. https://doi.org/10.1007/s10344-022-01581-y

Brown, T.L., Airs, P.M., Porter, S., Caplat, P., Morgan, E.R., 2022. Understanding the role of wild ruminants in anthelmintic resistance in livestock. Biol. Lett. 18, 20220057. https://doi.org/10.1098/rsbl.2022.0057

Burgess, C.G.S., Bartley, Y., Redman, E., Skuce, P.J., Nath, M., Whitelaw, F., Tait, A., Gilleard, J.S., Jackson, F., 2012. A survey of the trichostrongylid nematode species present on UK sheep farms and associated anthelmintic control practices. Vet. Parasitol. 189, 299–307. https://doi.org/10.1016/j.vetpar.2012.04.009

Burnham, K.P., Anderson, D.R., 2002. Model selection and multimodel inference: a practical information-theoretic approach, 2nd ed. Springer, New York. https://doi.org/10.1007/b97636

Calderón-Sanou, I., Münkemüller, T., Boyer, F., Zinger, L., Thuiller, W., 2020. From environmental DNA sequences to ecological conclusions: how strong is the influence of methodological choices? J. Biogeogr. 47, 193–206. https://doi.org/10.1111/jbi.13681

Callahan, B.J., McMurdie, P.J., Holmes, S.P., 2017. Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. ISME J. 11, 2639–2643. https://doi.org/10.1038/ismej.2017.119

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016. DADA2: high-resolution sample inference from Illumina amplicon data. Nat. Methods 13, 581–583. https://doi.org/10.1038/nmeth.3869

Carcereri, A., Stancampiano, L., Marchiori, E., Sturaro, E., Ramanzin, M., Cassini, R., 2021. Factors influencing gastrointestinal parasites in a colony of Alpine ibex (*Capra ibex*) interacting with domestic ruminants. Hystrix Ital. J. Mammal. 32, 95–101. https://doi.org/10.4404/hystrix-00393-2020

Carlsson, A.M., Irvine, R.J., Wilson, K., Coulson, S.J., 2013. Adaptations to the Arctic: low-temperature development and cold tolerance in the free-living stages of a

parasitic nematode from Svalbard. Polar Biol. 36, 997–1005.
https://doi.org/10.1007/s00300-013-1323-7

Cerutti, M.C., Citterio, C.V., Bazzocchi, C., Epis, S., D'Amelio, S., Ferrari, N.,
Lanfranchi, P., 2010. Genetic variability of *Haemonchus contortus* (Nematoda:
Trichostrongyloidea) in alpine ruminant host species. J. Helminthol. 84, 276–
283. https://doi.org/10.1017/S0022149X09990587

Charlier, J., Bartley, D.J., Sotiraki, S., Martinez-Valladares, M., Claerebout, E., von
Samson-Himmelstjerna, G., Thamsborg, S.M., Hoste, H., Morgan, E.R., Rinaldi,
L., 2022. Chapter Three - Anthelmintic resistance in ruminants: challenges and
solutions, in: Rollinson, D., Stothard, R. (Eds.), Advances in Parasitology.
Academic Press, pp. 171–227. https://doi.org/10.1016/bs.apar.2021.12.002

Charlier, J., Rinaldi, L., Musella, V., Ploeger, H.W., Chartier, C., Vineer, H.R., Hinney,
B., von Samson-Himmelstjerna, G., Băcescu, B., Mickiewicz, M., Mateus, T.L.,
Martinez-Valladares, M., Quealy, S., Azaizeh, H., Sekovska, B., Akkari, H.,
Petkevicius, S., Hektoen, L., Höglund, J., Morgan, E.R., Bartley, D.J.,
Claerebout, E., 2020. Initial assessment of the economic burden of major
parasitic helminth infections to the ruminant livestock industry in Europe. Prev.
Vet. Med. 182, 105103. https://doi.org/10.1016/j.prevetmed.2020.105103

Chintoan-Uta, C., Morgan, E.R., Skuce, P.J., Coles, G.C., 2014. Wild deer as potential
vectors of anthelmintic-resistant abomasal nematodes between cattle and
sheep farms. Proc. R. Soc. B Biol. Sci. 281, 20132985.
https://doi.org/10.1098/rspb.2013.2985

Chirichella, R., Apollonio, M., Putman, R., 2014. Competition between domestic and
wild Ungulates., in: Putman, R., Apollonio, M. (Eds.), Behaviour and
Management of European Ungulates. pp. 110–123.

Citterio, C.V., Caslini, C., Milani, F., Sala, M., Ferrari, N., Lanfranchi, P., 2006.
Abomasal Nematode Community in an Alpine Chamois (*Rupicapra r. rupicapra*
) Population before and after a Die-Off. J. Parasitol. 92, 918–927.

De Barba, M., Miquel, C., Boyer, F., Mercier, C., Rioux, D., Coissac, E., Taberlet, P.,
2014. DNA metabarcoding multiplexing and validation of data accuracy for diet
assessment: application to omnivorous diet. Mol. Ecol. Resour. 14, 306–323.
https://doi.org/10.1111/1755-0998.12188

Deplazes, P., Eckert, J., Mathis, A., Samson-Himmelstjerna, G. von, Zahner, H., 2016.
Parasitology in veterinary medicine. Parasitology in veterinary medicine.

Dickinson, E.R., McFarland, C., Toïgo, C., Michael Scantlebury, D., Stephens, P.A.,
Marks, N.J., Morgan, E.R., 2024. Host movement dominates the predicted
effects of climate change on parasite transmission between wild and domestic
mountain ungulates. R. Soc. Open Sci. 11, 230469.
https://doi.org/10.1098/rsos.230469

Dimunová, D., Matoušková, P., Navrátilová, M., Nguyen, L.T., Ambrož, M., Vokřál, I.,
Szotáková, B., Skálová, L., 2022. Environmental circulation of the anthelmintic
drug albendazole affects expression and activity of resistance-related genes in
the parasitic nematode *Haemonchus contortus*. Sci. Total Environ. 822,
153527. https://doi.org/10.1016/j.scitotenv.2022.153527

Dyson, K., 2018. Custom community ecology helper R scripts. URL
https://github.com/kdyson/R_Scripts

Fiel, C.A., Fernández, A.S., Rodríguez, E.M., Fusé, L.A., Steffan, P.E., 2012.
Observations on the free-living stages of cattle gastrointestinal nematodes. Vet.
Parasitol. 187, 217–226. https://doi.org/10.1016/j.vetpar.2012.01.011

Francis, E.K., Šlapeta, J., 2023. Refugia or reservoir? Feral goats and their role in the
maintenance and circulation of benzimidazole-resistant gastrointestinal
nematodes on shared pastures. Parasitology 1–11.
https://doi.org/10.1017/S0031182023000380

Gauthier, D., Villaret, J., 1990. La réintroduction en France du bouquetin des Alpes.
Rev Eco Terre Vie 45, 97–120. https://doi.org/10.3406/revec.1990.6338

Gibbs, H.C., 1986. Hypobiosis in Parasitic Nematodes—An Update, in: Baker, J.R.,
Muller, R. (Eds.), Advances in Parasitology. Academic Press, pp. 129–174.
https://doi.org/10.1016/S0065-308X(08)60343-7

Grossen, C., Biebach, I., Angelone-Alasaad, S., Keller, L.F., Croll, D., 2018. Population
genomics analyses of European ibex species show lower diversity and higher
inbreeding in reintroduced populations. Evol. Appl. 11, 123–139.
https://doi.org/10.1111/eva.12490

Gruner, L., Sauvé, C., Boulard, C., Calamel, M., 2006. Analysis of the relationship
between land use and the parasitism of sheep during their transhumance. Anim.
Res. 55, 177–188. https://doi.org/10.1051/animres:2006009

Hahnel, S.R., Zdraljevic, S., Rodriguez, B.C., Zhao, Y., McGrath, P.T., Andersen, E.C.,
2018. Extreme allelic heterogeneity at a *Caenorhabditis elegans* beta-tubulin

locus explains natural resistance to benzimidazoles. PLOS Pathog. 14, e1007226. https://doi.org/10.1371/journal.ppat.1007226

Hamilton, K.M., Waghorn, T.S., de Waal, T., Keane, O.M., Green, P., Leathwick, D.M., 2022. *In vitro* evaluation of fitness parameters for isolates of *Teladorsagia circumcincta* resistant and susceptible to multiple anthelmintic classes. Vet. Parasitol. 310, 109791. https://doi.org/10.1016/j.vetpar.2022.109791

Hellard, E., Fouchet, D., Vavre, F., Pontier, D., 2015. Parasite–Parasite Interactions in the Wild: How To Detect Them? Trends Parasitol. 31, 640–652. https://doi.org/10.1016/j.pt.2015.07.005

Hinney, B., Schoiswohl, J., Melville, L., Ameen, V.J., Wille-Piazzai, W., Bauer, K., Joachim, A., Krücken, J., Skuce, P.J., Krametter-Frötscher, R., 2020. High frequency of benzimidazole resistance alleles in trichostrongyloids from Austrian sheep flocks in an alpine transhumance management system. BMC Vet. Res. 16, 132. https://doi.org/10.1186/s12917-020-02353-z

Hodgkinson, J.E., Kaplan, R.M., Kenyon, F., Morgan, E.R., Park, A.W., Paterson, S., Babayan, S.A., Beesley, N.J., Britton, C., Chaudhry, U., Doyle, S.R., Ezenwa, V.O., Fenton, A., Howell, S.B., Laing, R., Mable, B.K., Matthews, L., McIntyre, J., Milne, C.E., Morrison, T.A., Prentice, J.C., Sargison, N.D., Williams, D.J.L., Wolstenholme, A.J., Devaney, E., 2019. Refugia and anthelmintic resistance: Concepts and challenges. Int. J. Parasitol. Drugs Drug Resist. 10, 51–57. https://doi.org/10.1016/j.ijpddr.2019.05.001

Hudson, P.J., Dobson, A.P., Lafferty, K.D., 2006. Is a healthy ecosystem one that is rich in parasites? Trends Ecol. Evol., Twenty years of TREE - part 2 21, 381–385. https://doi.org/10.1016/j.tree.2006.04.007

Kaplan, R.M., Denwood, M.J., Nielsen, M.K., Thamsborg, S.M., Torgerson, P.R., Gilleard, J.S., Dobson, R.J., Vercruysse, J., Levecke, B., 2023. World Association for the Advancement of Veterinary Parasitology (W.A.A.V.P.) guideline for diagnosing anthelmintic resistance using the faecal egg count reduction test in ruminants, horses and swine. Vet. Parasitol. 318, 109936. https://doi.org/10.1016/j.vetpar.2023.109936

Kaplan, R.M., Vidyashankar, A.N., 2012. An inconvenient truth: Global worming and anthelmintic resistance. Vet. Parasitol., Special issue: novel approaches to the control of helminth parasites of livestock 186, 70–78. https://doi.org/10.1016/j.vetpar.2011.11.048

Kessler, C., Brambilla, A., Waldvogel, D., Camenisch, G., Biebach, I., Leigh, D.M., Grossen, C., Croll, D., 2022. A robust sequencing assay of a thousand amplicons for the high-throughput population monitoring of Alpine ibex immunogenetics. Mol. Ecol. Resour. 22, 66–85. https://doi.org/10.1111/1755-0998.13452

Khanyari, M., Robinson, S., Milner-Gulland, E.J., Morgan, E.R., Rana, R.S., Suryawanshi, K.R., 2022. Pastoralism in the high Himalayas: Understanding changing practices and their implications for parasite transmission between livestock and wildlife. Pastoralism 12, 44. https://doi.org/10.1186/s13570-022-00257-1

Klein, S.L., 2004. Hormonal and immunological mechanisms mediating sex differences in parasite infection. Parasite Immunol. 26, 247–264. https://doi.org/10.1111/j.0141-9838.2004.00710.x

Klein, S.L., 2000. The effects of hormones on sex differences in infection: from genes to behavior. Neurosci. Biobehav. Rev. 24, 627–638. https://doi.org/10.1016/S0149-7634(00)00027-0

Kolar, L., Flajs, V.C., Kužner, J., Marc, I., Pogačnik, M., Bidovec, A., van Gestel, C.A.M., Eržen, N.K., 2006. Time profile of abamectin and doramectin excretion and degradation in sheep faeces. Environ. Pollut., Soil and Sediment Remediation 144, 197–202. https://doi.org/10.1016/j.envpol.2005.12.019

Laca Megyesi, Š., Königová, A., Babják, M., Molnár, L., Rajský, M., Szestáková, E., Major, P., Soroka, J., Urda Dolinská, M., Komáromyová, M., Várady, M., 2019. Wild ruminants as a potential risk factor for transmission of drug resistance in the abomasal nematode *Haemonchus contortus*. Eur. J. Wildl. Res. 66, 9. https://doi.org/10.1007/s10344-019-1351-x

Langrová, I., Makovcová, K., Vadlejch, J., Jankovská, I., Petrtýl, M., Fechtner, J., Keil, P., Lytvynets, A., Borkovcová, M., 2008. Arrested development of sheep strongyles: onset and resumption under field conditions of Central Europe. Parasitol. Res. 103, 387–392. https://doi.org/10.1007/s00436-008-0984-6

Lavín, S., Marco, I., Rossi, L., Meneguz, P.G., Viñas, L., 1997. Haemonchosis in Spanish Ibex. J. Wildl. Dis. 33, 656–659. https://doi.org/10.7589/0090-3558-33.3.656

Leathwick, D.M., Ganesh, S., Waghorn, T.S., 2015. Evidence for reversion towards anthelmintic susceptibility in *Teladorsagia circumcincta* in response to

resistance management programmes. Int. J. Parasitol. Drugs Drug Resist. 5, 9–15. https://doi.org/10.1016/j.ijpddr.2015.01.001

Legendre, P., Legendre, L., 2012. Numerical Ecology. Elsevier.

Leigh, J.W., Bryant, D., 2015. popart: full-feature software for haplotype network construction. Methods Ecol. Evol. 6, 1110–1116. https://doi.org/10.1111/2041-210x.12410

Lello, J., Boag, B., Fenton, A., Stevenson, I.R., Hudson, P.J., 2004. Competition and mutualism among the gut helminths of a mammalian host. Nature 428, 840–844. https://doi.org/10.1038/nature02490

Lozupone, C., Knight, R., 2005. UniFrac: a new phylogenetic method for comparing microbial communities. Appl. Environ. Microbiol. 71, 8228–8235. https://doi.org/10.1128/AEM.71.12.8228-8235.2005

Luzzago, C., Ebranati, E., Cabezón, O., Fernández-Sirera, L., Lavín, S., Rosell, R., Veo, C., Rossi, L., Cavallero, S., Lanfranchi, P., Marco, I., Zehender, G., 2016. Spatial and Temporal Phylogeny of Border Disease Virus in Pyrenean Chamois (*Rupicapra p. pyrenaica*). PLOS ONE 11, e0168232. https://doi.org/10.1371/journal.pone.0168232

Marchand, P., Freycon, P., Herbaux, J.-P., Game, Y., Toïgo, C., Gilot-Fromont, E., Rossi, S., Hars, J., 2017. Sociospatial structure explains marked variation in brucellosis seroprevalence in an Alpine ibex population. Sci. Rep. 7, 15592. https://doi.org/10.1038/s41598-017-15803-w

Margaillan, L., 2021. Simultaneous GPS monitoring during summer reveals habitat selection in male Alpine ibex is shaped by resource and interference competition with sheep herds (Master thesis). Office Français de la Biodiversité.

Marreros, N., Frey, C.F., Willisch, C.S., Signer, C., Ryser-Degiorgis, M.-P., 2012. Coprological analyses on apparently healthy Alpine ibex (*Capra ibex ibex*) from two Swiss colonies. Vet. Parasitol. 186, 382–389. https://doi.org/10.1016/j.vetpar.2011.11.009

Martínez-Guijosa, J., Martínez-Carrasco, C., López-Olvera, J.R., Fernández-Aguilar, X., Colom-Cadena, A., Cabezón, O., Mentaberre, G., Ferrer, D., Velarde, R., Gassó, D., Garel, M., Rossi, L., Lavín, S., Serrano, E., 2015. Male-biased gastrointestinal parasitism in a nearly monomorphic mountain ungulate. Parasit. Vectors 8, 165. https://doi.org/10.1186/s13071-015-0774-9

McKenna, P.B., 1998. The effect of previous cold storage on the subsequent recovery
of infective third stage nematode larvae from sheep faeces. Vet. Parasitol. 80,
167–172. https://doi.org/10.1016/S0304-4017(98)00203-9

McMurdie, P.J., Holmes, S., 2013. phyloseq: An R Package for Reproducible
Interactive Analysis and Graphics of Microbiome Census Data. PLOS ONE 8,
e61217. https://doi.org/10.1371/journal.pone.0061217

Morgan, E.R., Aziz, N.-A.A., Blanchard, A., Charlier, J., Charvet, C., Claerebout, E.,
Geldhof, P., Greer, A.W., Hertzberg, H., Hodgkinson, J., Höglund, J., Hoste, H.,
Kaplan, R.M., Martínez-Valladares, M., Mitchell, S., Ploeger, H.W., Rinaldi, L.,
von Samson-Himmelstjerna, G., Sotiraki, S., Schnyder, M., Skuce, P., Bartley,
D., Kenyon, F., Thamsborg, S.M., Vineer, H.R., de Waal, T., Williams, A.R., van
Wyk, J.A., Vercruysse, J., 2019. 100 Questions in Livestock Helminthology
Research. Trends Parasitol. 35, 52–71. https://doi.org/10.1016/j.pt.2018.10.006

Murali, A., Bhargava, A., Wright, E.S., 2018. IDTAXA: a novel approach for accurate
taxonomic classification of microbiome sequences. Microbiome 6, 140.
https://doi.org/10.1186/s40168-018-0521-5

Nagy, G., Csivincsik, Á., Sugár, L., Zsolnai, A., 2017. Benzimidazole resistance within
red deer, roe deer and sheep populations within a joint habitat in Hungary. Small
Rumin. Res. 149, 172–175. https://doi.org/10.1016/j.smallrumres.2017.02.012

O'Connor, L.J., Walkden-Brown, S.W., Kahn, L.P., 2006. Ecology of the free-living
stages of major trichostrongylid parasites of sheep. Vet. Parasitol. 142, 1–15.
https://doi.org/10.1016/j.vetpar.2006.08.035

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D.,
Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H.,
Szoecs, E., Wagner, H., 2020. vegan : community ecology package.

Oliver-Guimerá, A., Martínez-Carrasco, C., Tvarijonaviciute, A., Ruiz de Ybáñez, M.R.,
Martínez-Guijosa, J., López-Olvera, J.R., Fernández-Aguilar, X., Colom-
Cadena, A., Mentaberre, G., Velarde, R., Gassó, D., Garel, M., Rossi, L., Lavín,
S., Serrano, E., 2017. The physiological cost of male-biased parasitism in a
nearly monomorphic mammal. Parasit. Vectors 10, 200.
https://doi.org/10.1186/s13071-017-2060-5

Ozenda, P., 1985. La végétation de la chaîne alpine dans l'espace montagnard
européen. Masson, Paris.

Palcy, C., Silvestre, A., Sauve, C., Cortet, J., Cabaret, J., 2010. Benzimidazole resistance in *Trichostrongylus axei* in sheep: long-term monitoring of affected sheep and genotypic evaluation of the parasite. Vet. J. 183, 68–74. https://doi.org/10.1016/j.tvjl.2008.09.012

Papadopoulos, E., Gallidis, E., Ptochos, S., 2012. Anthelmintic resistance in sheep in Europe: a selected review. Vet. Parasitol., Special issue: Update on Parasitic Diseases of Sheep 189, 85–88. https://doi.org/10.1016/j.vetpar.2012.03.036

Portanier, E., Garel, M., Devillard, S., Maillard, D., Poissant, J., Galan, M., Benabed, S., Poirel, M.-T., Duhayer, J., Itty, C., Bourgoin, G., 2019. Both candidate gene and neutral genetic diversity correlate with parasite resistance in female Mediterranean mouflon. BMC Ecol. 19, 12. https://doi.org/10.1186/s12898-019-0228-x

Posada, D., Crandall, K.A., 1998. modeltest: testing the model of DNA substitution. Bioinformatics 14, 817–818. https://doi.org/10.1093/bioinformatics/14.9.817

Queiroz, C., Levy, M., Avramenko, R., Redman, E., Kearns, K., Swain, L., Silas, H., Uehlinger, F., Gilleard, J.S., 2020. The use of ITS-2 rDNA nemabiome metabarcoding to enhance anthelmintic resistance diagnosis and surveillance of ovine gastrointestinal nematodes. Int. J. Parasitol. Drugs Drug Resist. 14, 105–117. https://doi.org/10.1016/j.ijpddr.2020.09.003

Ramünke, S., Melville, L., Rinaldi, L., Hertzberg, H., de Waal, T., von Samson-Himmelstjerna, G., Cringoli, G., Mavrot, F., Skuce, P., Krücken, J., Demeler, J., 2016. Benzimidazole resistance survey for *Haemonchus*, *Teladorsagia* and *Trichostrongylus* in three European countries using pyrosequencing including the development of new assays for *Trichostrongylus*. Int. J. Parasitol. Drugs Drug Resist. 6, 230–240. https://doi.org/10.1016/j.ijpddr.2016.10.002

Raynaud, J.-P., William, G., Brunault, G., 1970. Etude de l'efficacité d'une technique de coproscopie quantitative pour le diagnostic de routine et le contrôle des infestations parasitaires des bovins, ovins, équins et porcins. Ann. Parasitol. Hum. Comparée 45, 321–342. https://doi.org/10.1051/parasite/1970453321

Redman, E., Queiroz, C., Bartley, D.J., Levy, M., Avramenko, R.W., Gilleard, J.S., 2019. Validation of ITS-2 rDNA nemabiome sequencing for ovine gastrointestinal nematodes and its application to a large scale survey of UK sheep farms. Vet. Parasitol. 275, 108933. https://doi.org/10.1016/j.vetpar.2019.108933

Redman, E., Whitelaw, F., Tait, A., Burgess, C., Bartley, Y., Skuce, P.J., Jackson, F., Gilleard, J.S., 2015. The emergence of resistance to the benzimidazole anthlemintics in parasitic nematodes of livestock is characterised by multiple independent hard and soft selective sweeps. PLoS Negl. Trop. Dis. 9, e0003494. https://doi.org/10.1371/journal.pntd.0003494

Richomme, C., Gauthier, D., Fromont, E., 2006. Contact rates and exposure to inter-species disease transmission in mountain ungulates. Epidemiol. Infect. 134, 21–30. https://doi.org/10.1017/S0950268805004693

Roeber, F., Jex, A.R., Gasser, R.B., 2013. Impact of gastrointestinal parasitic nematodes of sheep, and the role of advanced molecular tools for exploring epidemiology and drug resistance - an Australian perspective. Parasit. Vectors 6, 153. https://doi.org/10.1186/1756-3305-6-153

Rose, H., Rinaldi, L., Bosco, A., Mavrot, F., Waal, T. de, Skuce, P., Charlier, J., Torgerson, P.R., Hertzberg, H., Hendrickx, G., Vercruysse, J., Morgan, E.R., 2015. Widespread anthelmintic resistance in European farmed ruminants: a systematic review. Vet. Rec. 176, 546–546. https://doi.org/10.1136/vr.102982

Rose Vineer, H., Morgan, E.R., Hertzberg, H., Bartley, D.J., Bosco, A., Charlier, J., Chartier, C., Claerebout, E., de Waal, T., Hendrickx, G., Hinney, B., Höglund, J., Ježek, J., Kašný, M., Keane, O.M., Martínez-Valladares, M., Mateus, T.L., McIntyre, J., Mickiewicz, M., Munoz, A.M., Phythian, C.J., Ploeger, H.W., Rataj, A.V., Skuce, P.J., Simin, S., Sotiraki, S., Spinu, M., Stuen, S., Thamsborg, S.M., Vadlejch, J., Varady, M., von Samson-Himmelstjerna, G., Rinaldi, L., 2020. Increasing importance of anthelmintic resistance in European livestock: creation and meta-analysis of an open database. Parasite 27, 69. https://doi.org/10.1051/parasite/2020062

Rossi, L., Tizzani, P., Rambozzi, L., Moroni, B., Meneguz, P.G., 2019. Sanitary emergencies at the wild/domestic caprines interface in Europe. Animals 9, 922. https://doi.org/10.3390/ani9110922

Ryser-Degiorgis, M.-P., Ingold, P., Tenhu, H., Less, A.M.T., Ryser, A., Giacometti, M., 2002. Encounters between Alpine ibex, Alpine chamois and domestic sheep in the Swiss Alps. Hystrix Ital. J. Mammal. 13. https://doi.org/10.4404/hystrix-13.1-2-4180

Sargison, N.D., Wilson, D.J., Bartley, D.J., Penny, C.D., Jackson, F., 2007. Haemonchosis and teladorsagiosis in a Scottish sheep flock putatively

associated with the overwintering of hypobiotic fourth stage larvae. Vet. Parasitol. 147, 326–331. https://doi.org/10.1016/j.vetpar.2007.04.011

Schliep, K.P., 2011. phangorn: phylogenetic analysis in R. Bioinformatics 27, 592–593. https://doi.org/10.1093/bioinformatics/btq706

Taberlet, P., Bonin, A., Zinger, L., Coissac, E., 2018. Environmental DNA: for biodiversity research and monitoring. Oxford University Press, Oxford, UK.

Taylor, M.A., Coop, R.L., Wall, R.L., 2015. Veterinary Parasitology, 4th ed. ed. Wiley Blackwell, Chichester.

Tompkins, D.M., Dunn, A.M., Smith, M.J., Telfer, S., 2011. Wildlife diseases: from individuals to ecosystems. J. Anim. Ecol. 80, 19–38. https://doi.org/10.1111/j.1365-2656.2010.01742.x

Utaaker, K.S., Ytrehus, B., Davey, M.L., Fossøy, F., Davidson, R.K., Miller, A.L., Robertsen, P.-A., Strand, O., Rauset, G.R., 2023. Parasite spillover from domestic sheep to wild reindeer—The role of salt licks. Pathogens 12, 186. https://doi.org/10.3390/pathogens12020186

van Wyk, J.A., Mayhew, E., 2013. Morphological identification of parasitic nematode infective larvae of small ruminants and cattle: a practical lab guide. Onderstepoort J. Vet. Res. 80, 00–00.

Vercruysse, J., Charlier, J., Dijk, J.V., Morgan, E.R., Geary, T., Samson-Himmelstjerna, G. von, Claerebout, E., 2018. Control of helminth ruminant infections by 2030. Parasitology 145, 1655–1664. https://doi.org/10.1017/S003118201700227X

Viana, M., Mancy, R., Biek, R., Cleaveland, S., Cross, P.C., Lloyd-Smith, J.O., Haydon, D.T., 2014. Assembling evidence for identifying reservoirs of infection. Trends Ecol. Evol. 29, 270–279. https://doi.org/10.1016/j.tree.2014.03.002

Walker, J.G., Morgan, E.R., 2014. Generalists at the interface: nematode transmission between wild and domestic ungulates. Int. J. Parasitol. Parasites Wildl. 3, 242–250. https://doi.org/10.1016/j.ijppaw.2014.08.001

Wang, Q., Garrity, G.M., Tiedje, J.M., Cole, J.R., 2007. Naïve bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. Appl. Environ. Microbiol. 73, 5261–5267. https://doi.org/10.1128/AEM.00062-07

Whittaker, J.H., Carlson, S.A., Jones, D.E., Brewer, M.T., 2017. Molecular mechanisms for anthelmintic resistance in strongyle nematode parasites of

veterinary importance. J. Vet. Pharmacol. Ther. 40, 105–115.
https://doi.org/10.1111/jvp.12330

Workentine, M.L., Chen, R., Zhu, S., Gavriliuc, S., Shaw, N., Rijke, J. de, Redman,
E.M., Avramenko, R.W., Wit, J., Poissant, J., Gilleard, J.S., 2020. A database
for ITS2 sequences from nematodes. BMC Genet. 21, 74.
https://doi.org/10.1186/s12863-020-00880-0

Wright, E.S., 2016. Using DECIPHER v2.0 to Analyze Big Biological Sequence Data
in R. R J. 8, 352–359.

Zaffaroni, E., Teresa Manfredi, M., Citterio, C., Sala, M., Piccolo, G., Lanfranchi, P.,
2000. Host specificity of abomasal nematodes in free ranging alpine ruminants.
Vet. Parasitol. 90, 221–230. https://doi.org/10.1016/S0304-4017(00)00240-5

1307