isGWAS: ultra-high-throughput, scalable and equitable inference of genetic associations with disease

Christopher N Foley[1,2*], Zhana Kuncheva[1,2], Riccardo E. Marioni[3], Heiko Runz[4], Benjamin B Sun[4*].

[1]Optima Partners, Edinburgh, UK.

[2]Bayes Centre, University of Edinburgh, Edinburgh, UK.

[3]Centre for Genomic and Experimental Medicine, Institute of Genetics and Cancer, University of Edinburgh, Edinburgh, UK.

[4]Translational Sciences, Biogen Inc., Cambridge, MA, US.

*Correspondence:

Dr Christopher N Foley

Email: chris.foley@optimapartners.co.uk or chris.neal.foley@gmail.com

Dr Benjamin B Sun

Email: benjamin.sun@biogen.com or bbsun92@outlook.com

Abstract

Genome-wide association studies (GWAS) have proven a powerful tool for human geneticists to generate biological insights or hypotheses for drug discovery. Nevertheless, a dependency on sensitive individual-level data together with ever-increasing cohort sample sizes, numbers of variants and phenotypes studied put a strain on existing algorithms, limiting the GWAS approach from maximising potential. Here we present in-silico GWAS (isGWAS), a uniquely scalable algorithm to infer regression parameters in case-control GWAS from cohort-level summary data. For any sample size, isGWAS computes a variant-disease association parameter in ~1 millisecond, or ~11m variants in UK-Biobank within ~4 minutes (~1500-fold faster than state-of-the-art). Extensive simulations and empirical tests demonstrate that isGWAS results are highly comparable to traditional regression-based approaches. We further introduce a heuristic re-sampling algorithm, leapfrog re-sampler (LRS), to extrapolate association results to semi-virtually enlarged cohorts. Owing to significant computational gains we anticipate a broad use of isGWAS and LRS which are customizable on a web interface.

Main

Genome wide association studies (GWAS) have been immensely successful in unravelling the genetic contribution to human disease. Cost-effective genotyping and large biobank cohorts now make it possible to routinely conduct GWAS for tens of millions of variants in hundreds of thousands of individuals across thousands of phenotypes[1]. With the advent of population-scale whole genome sequencing and expansion of GWAS to research participants of non-European ancestries, these numbers can be expected to increase by another magnitude over the next few years[2], [3].

Current GWAS approaches that compute variant-disease associations in a regression framework, such as PLINK[4], fastGWA[5], BOLT-LMM[6], SAIGE[7] and REGENIE[8], require access to and input from ever increasing individual-level data (ILD). The efforts of individual-level GWAS sample collection, genotyping and data analysis tend to grow as a polynomial function of sample size[7], [8]. Moreover, the exchange of ILD between researchers is non-trivial due partly to data size but increasingly to strict – but essential - data protection regulations, which can limit the scope of collaborative analyses and biological insights gained[9]–[12]. Finally, the substantial computational and financial burden of running massive-scale GWAS, especially for binary disease outcomes, is exacerbating inequity between researchers, typically favouring already well-equipped institutions. There is therefore a pressing need for innovative approaches that help attenuate the increasing resource and financial inequities for conducting contemporary GWAS and to help decide where limited resources should best be allocated.

Here we present in-silico GWAS (isGWAS), a biobank-scalable and computationally highly efficient algorithm to infer genetic regression parameters in case-control GWAS from just four broadly ascertained cohort-level summary parameters: the counts of cases and controls within a cohort, as well as case and control minor allele frequencies (MAFs). isGWAS is highly parallelisable, exceeding efficiencies of current GWAS analysis tools by several orders of magnitude. Furthermore, we demonstrate that isGWAS yield association summary statistics highly comparable to traditional ILD regression-based approaches through extensive simulations and empirical tests in UK Biobank[13], Biobank Japan[14] and the Psychiatric Genomics Consortium cohort[15]. Owing to the sizeable computational gains, we introduce a heuristic re-sampling algorithm, called the leapfrog re-sampler (LRS), which can confidently

49   extrapolate GWAS results to larger sample sizes, both at a locus or genome-wide scale. Our

50   underlying methodology also leads to several desirable high-utility properties. We release a

51   web tool available to the wider public to conduct customized isGWAS at www.optima-

52   isgwas.com.

Results

Genome-wide association testing from sufficient statistics

isGWAS assumes disease-variant associations can be evaluated via a logistic-link function and, similar to widely used methods[7], [8], uses a Firth adjusted maximum likelihood procedure and Newton-Raphson solver to estimate genetic effects, standard errors and association p-value [7], [8], [16]. isGWAS' notable advance is based on the insight that the Newton-Raphson procedure can be simplified so that: (a) elements of the Fisher information matrix and score function vector are collapsed by taking expectation over the empirical or *a priori* distribution of a genetic variant; and (b) sufficient statistics – a specific type of summary data - are used as input variables in the score function (see **Methods** for details). We provide several options to initialise the Newton-Raphson algorithm [17], [18] that improve computational performance and reduce analysis time (**Supplementary Information**). In brief, let y_i denote disease status for the i-th individual and $g_{ij,M}$ denote the j-th genetic variant under model M (e.g., additive, recessive or dominant). The sufficient statistic triple used by isGWAS is:

$$\left\{ T_{1j} = \sum_{i=1}^{N_j} y_i = N_j^*, \qquad T_{2j} = \sum_{i=1}^{N_j} y_i \, g_{ij,M}, \qquad T_{3j} = \sum_{i=1}^{N_j} g_{ij,M} \right\},$$

where T_{1j} is the total number of cases for variant j, T_{2j} is the covariance between the outcome y and genotype g for variant j under model M, and T_{3j} is the minor allele count for variant j under model M. For each variant, data can be provided as either: the sufficient statistic triple $\{T_{1j}, T_{2j}, T_{3j}\}$ plus sample size N_j (necessary for computing standard errors) or separately, on assuming Hardy-Weinberg equilibrium (HWE), as $\{N_j, N_j^*, MAF_{j,M}, MAF_{j,M}^*\}$. Default GWAS analyses assume HWE, making input data widely available [13]–[15], [19]–[21] for researchers to perform isGWAS, replicate or further expand on classical GWAS analyses (**Methods**). If MAFs for cases and controls are supplied, isGWAS will automatically convert to the pair

$\{MAF_{j,M}, MAF_{j,M}^*\}$ (**Methods**). After convergence, which is guaranteed for most scenarios by a

re-initialization approach (empirically all scenarios converged using isGWAS-Firth), the

estimated genetic effect parameter $\hat{\beta}_{j,M}$ and standard error $SE(\hat{\beta}_{j,M})$ are used to construct Wald

p-values (**Methods**). Additional options include a sample-level likelihood ratio-test or *p*-values

computed using sandwich-robust standard errors (**Supplementary Methods**). A simplified

illustration highlighting differences and computational advantages of isGWAS against ILD-

based genetic association analyses is summarised in **Figure 1**.

isGWAS reliably identifies genetic associations across cohorts and diseases

We benchmarked isGWAS in real-data settings and performed simulation studies to compare

isGWAS performance and results relative to several existing individual-level data (ILD)-based

approaches. Our assessments broadly fall into two categories: (1) methods which require ILD,

i.e., REGENIE[8] , logistic and Firth corrected regression[16], and (2) approaches which do not

require ILD directly, i.e., the logistic ad-hoc estimator[17] and Fisher's Exact Test (FET)[22].

We note FET was successfully leveraged for efficient large-scale GWAS analyses recently[23]

. Using data from UK Biobank (UKB), we first assessed isGWAS performance against the

popular ILD based regression approach REGENIE[8] by deploying both methods for analyses

of seven diseases some of which were previously used for establishing GWAS methodology[8].

Second, we evaluated isGWAS's ability to replicate 309 significantly associated variants from

the Biobank Japan (BBJ) meta-analysis of 30 diseases[24] using only the published sample-

summaries, i.e., without access to ILD, per variant and disease pair. Lastly, by considering

multiple iterations of nested schizophrenia meta-analysis GWAS[25]–[27] we assessed

isGWAS's ability to accurately predict genomic regions and significant novel associations. The

isGWAS sample size of the meta-analysis from 2014 was virtually expanded to match the

numbers from the larger 2022 GWAS whilst holding constant the allele frequency and

prevalence information in 2014 cohort.

Performance against ILD based regression GWAS in UK Biobank

We compared isGWAS results and computational performance against the state-of-the-art ILD-

based method REGENIE for seven diseases[8] in UKB: asthma (IC10:J45); atherosclerosis

(IC10:I25); colon cancer (IC10:C18); hypertension (IC10:I10); glaucoma (IC10:H40); stroke

(IC10:I63); and thyroid gland cancer (IC10:C73). Case-control ratios varied from 1:2 in

hypertension to 1:669 in thyroid gland cancer across the diseases (**Supplementary Table 1a**),

allowing for the review of performance in near balanced to highly imbalanced case-control

settings. To help attenuate the possible influence of confounders when deploying isGWAS,

particularly sample relatedness and population structure, we describe and apply additional data

quality control (QC) steps before computing the required sample-level sufficient statistics

(**Methods**). After additional QC, the total sample size analysed was ~335,000 individuals per

disease (**Supplementary Table 1**). This sample was used to perform and contrast analyses in

both isGWAS and REGENIE. We review approaches which leverage additional insight from

any removed samples in the **Discussion**. For each of the seven diseases, we applied isGWAS

(no covariates) and two-step REGENIE with Firth correction (including the covariates age,

gender and ethnicity principal components) to ~11 million autosomal variants. Results are

presented in **Table 1**, **Figures 2-3, Supplementary Figures 1-15, Supplementary Tables 1-9**

and Supplementary Files 1-2.

Across all seven traits tested we observed close to perfect consistency between REGENIE and

isGWAS association results, as illustrated in the mirrored Manhattan and *p-p* plots for asthma

(**Figure 2a**) and other diseases (**Supplementary Figures 1-6**). Concordance between

REGENIE and isGWAS is further validated by benchmarking accuracy (**Table 1**) and Pearson

correlations between estimated *p*-values $(cor(p_{isG}, p_R)) > 0.94$ at log_{10} scale

(**Supplementary Tables 2-3**). The results are consistent for varying prevalence levels

(**Supplementary Figures 1-6**) and are not affected by covariate adjustment (**Supplementary**

Tables 2-3). The consistency translated to the regional locus level. This is exemplified by a

locus zoom plot of the *FLG2* gene region for asthma (**Figure 2b**) where isGWAS not only

nominated the identical GWAS lead variants but also largely recapitulated the overall

association pattern identified through REGENIE. This observation is consistent across the lead

independent loci from the asthma GWAS (**Supplementary Figure 7**) and translates to all other

diseases studied (**Supplementary Figure 8**). For a comprehensive numerical comparison of

association results, we took REGENIE derived *p*-values as the ground truth, retaining all SNPs

with $p < 0.01$ and setting the true positive threshold as $p < 5 \times 10^{-8}$ (excluding stroke

(**Supplementary Figure 4**) which did not yield any significant associations). We computed the

accuracy, false positive (FPR), true positive (TPR) and false discovery rates (FDR) of isGWAS

(**Figure 3a** and **Table 1**). Accuracy of isGWAS was $\geq 99.98\%$ for each disease, highlighting

excellent overall correspondence between methods. The FPR was low, i.e., $FPR \lesssim 10^{-5}$, and

TPR was generally good at $> 88\%$ - excluding hypertension which had a $TPR = 0.63$. The

FDR was below $\leq 5\%$ for each disease, revealing that the positive predictive value of isGWAS

was greater than 95%.

Importantly, isGWAS and REGENIE results differed for two broad categories: (i) the

estimation of genetic effect sizes; and (ii) computational performance. When non-confounding

covariates are excluded (β_{no_cov}) or included (β_{cov}) in a model, previous and extensive

investigations of effect size estimates in logistic regression deduce that $|\beta_{no\_cov}| \leq |\beta_{cov}|$, i.e.,

regression estimates are smaller in magnitude when excluding covariates but the null-

hypothesis of no association is maintained [19-20]. Overall, we replicate these results in our

analyses. isGWAS computed effect sizes are smaller in absolute value, but largely concordant

with covariate-adjusted REGENIE. Moreover, we fail to reject the null hypothesis for the same

variants almost always between methods - suggesting that the isGWAS QC helps attenuate

issues of population confounding (**Figures 2d-e, Supplementary Figures 1d-e – 6d-e, 9, 10**

and 11). An investigation of the performance of isGWAS without removing related individuals

highlights potential expansion of isGWAS beyond the recommended QC (**Methods**,

Supplementary Information), but further investigation – possibly leveraging the re-sampling

potential of isGWAS - is required on the reliability of isGWAS in family-based cohorts and

ethnically diverse populations (**Supplementary Figures 12-13, Supplementary Tables 5-7)**.

In our full-QC analyses, all estimated effects between isGWAS and REGENIE were observed

to be in the same direction, and the correlation between estimates was on average

$cor(\beta_{isG}, \beta_{REGENIE}) \approx 0.7$ (**Supplementary Tables 1-3**). The relative drop in the correlation

between effect estimates (≈ 0.7) and p-values ($\gtrsim 0.94$) is anticipated[28] and can be explained

on noting that, across all diseases more precise effect estimates (i.e., those with smaller standard

errors) have stronger concordance between approaches (**Figure 2e and Supplementary**

Figures 1d-6d). Overall, we found that at least 98% of isGWAS and REGENIE confidence

intervals (CI) overlap, (**Supplementary Table 4**). When effect estimates are viewed as a

function of MAF, the absolute value of REGENIE-derived estimates seemingly increase (along

with standard errors) as MAF decreases across all scenarios. This contrasts with isGWAS where

the relationship between MAF and effect size is less clear: fewer variants with low MAF are

associated with relatively larger effect sizes. However, the correspondingly narrower standard

errors guarantee the same significance p-values as REGENIE. The isGWAS derived

distribution of effect sizes is consistent with the hypothesis of a flattened heritability

distribution under negative selection[29]. Genomic inflation computed from isGWAS results

across all analyses was on average ≈ 1.07 and ranged between (0.94, 1.26) which was similar

to REGENIE with average ≈ 1.1 and range (1.01,1.3) (**Supplementary Figure 14** and

Supplementary Table 8). We deploy isGWAS with genotype imputation in our primary

analyses and, as secondary sensitivity analyses, without imputation. Our investigation reveals

some surprising results. Imputation occasionally led to changes in MAF between cases and

controls such that estimated genetic effects switched sign (i.e., effect direction) relative to

results computed from non-imputed data (**Supplementary Figure 15, Supplementary Table**

3, Supplementary Files 2-3). The approach might be used to efficiently flag ambiguous

significant results in analyses that are the result of the missing values imputation strategy

(mean-imputed in the case of REGENIE).

Finally, the computational gains of isGWAS relative to REGENIE Step 2 are striking: a full

genome-wide association assessment for each disease took approximately 4 minutes using

isGWAS and, on average over different prevalence, this is around 1,300 times faster than a like-

for-like assessment using REGENIE Step-2 (**Figure 7, Supplementary Table 9, and**

Supplementary File 4).

Replicating significant associations in Biobank Japan analyses

Using only publicly available summary information from Biobank Japan (BBJ), i.e., without

access to ILD, we looked to compare and replicate BBJ GWAS results across 42 diseases [24].

We considered 309 variants that were identified in [24] as genome-wide significant ($p <$

5×10^{-8}) across 30 of the 42 diseases. Our results reveal very close alignment between

isGWAS computed associations and those of [24] - correlation between p-values at log_{10} scale

was $cor(p_{isG}, p_{BBJ}) = 0.98$ with 92.2% of isGWAS computed genetic effects within the 95%

CI of the original study (**Supplementary Figure 16).** Using the published study-level results

[24] as the ground truth, isGWAS demonstrated good sensitivity and specificity

(**Supplementary Figure 17**). We alternatively assessed performance when setting more

stringent significance thresholds - returning near identical conclusions when classifying

variants at $p < 9.58 \times 10^{-9}$ (used in the original publication). Results for X-chromosome

variants in males and females were similarly concordant (results not presented).

isGWAS model validation using simulations

We generated simulated datasets to assess performance of isGWAS - with and without Firth

correction - against a variety of classical methods which either: (a) do not require ILD, the

logistic ad-hoc estimator[17] and Fisher's Exact Test[22]; or (b) require ILD, logistic and Firth

corrected regression[7], [8], [16]. We perform two simulation studies (**Figure 4**,

Supplementary Figure 18, Supplementary Information). isGWAS-Firth outperformed all

other approaches in terms of either computational cost or robustness of results over the range

of scenarios considered. It is well documented that computational performance is reduced when

using Firth's bias correction in ILD regression analyses[7], [8], [16], we discover, however,

that no-ILD isGWAS-Firth regression has significantly improved performance relative to

uncorrected isGWAS (**Supplementary Table 21**). As anticipated[23], when disease prevalence

is rare (i.e., $\pi \leq 0.01$) parameter estimates computed using non-Firth corrected ILD regression

were unreliable. The MSE and distribution of parameters estimated via ILD logistic regression

were often orders of magnitude poorer than other methods (**Figure 4a-c**). **Figure 4f-h**

highlights the chronological evolution of no-ILD p-value estimates, from Sasieni's logistic ad-

hoc estimator (1997)[17], Fisher's Exact Test (1922)[22] to isGWAS-Firth, illustrating

improvements in estimation via successive approaches. See **Supplementary Information** for

detailed results review.

Leapfrog re-sampling: using isGWAS to extrapolate variant association results to larger sample sizes

When ILD are available, the computational benefits of isGWAS make it possible to deploy resampling approaches to estimate empirical effect sizes, *p*-values and corresponding confidence intervals, previously considered computationally daunting in GWAS[30], [31]. We extend the idea by introducing a heuristic leapfrog re-sampling (LRS) algorithm to help forecast future results in larger hypothetical GWAS sample sizes (**Methods**). The LRS is summarised in three key steps: (1) specify a target sample size along with the number and size of sub-samples to be generated; (2) (leapfrog-step) compute sufficient statistics in the sub-samples and re-scale the estimated number of cases and controls to match the larger target sample size; and (3) deploy isGWAS in each leapfrog sample to recover a distribution of association *p*-values over the collection of sub-samples. In our testing of the LRS, we use the median *p*-value as a generally robust estimate of a target p-value (weighted or distribution-based summaries can alternatively be considered). Thus, the LRS leverages variation in both genotype and disease status between individuals in the current sample to help predict updates of parameters after adding new samples. Despite perceived similarities, traditional GWA power calculators[32] and the isGWAS-LRS are different. isGWAS-LRS does not require input of case-control ratios, heritability (i.e., beta estimates) or type-I error rates. Instead, multiple regression analyses are combined to forecast and test parameter estimates in expanding sample sizes.

We run the leapfrog re-sampler in both simulation and real-data settings, informed by the seven studied diseases in UKB (**Methods and Supplementary Information**). We evaluate performance over a range of initialisations, starting from a 10% increase to a maximum of 100% (i.e., 2-fold) increase in GWAS sample size relative to the current actual UKB sample size. Results are presented in **Figure 5**, **Supplementary Tables 10-11**. As is standard, we assume a

true positive association of p<5×10−8 in the target sample. Our results in simulated scenarios

(**Figure 5f**) reveal that: when doubling sample size from $N_{current} = 276,204$ to a maximum

$N_{target} = 552,408$, the accuracy and TPR progressively dropped for subsequent increases in

the target sample size, but values for each measurement were typically ≥80% across the range.

Our real world LRS analyses of UK Biobank data replicate and further elucidate performance

across the six of the seven diseases (**Figure 5a-e**). Using a subsample size of $N_{current} \approx$

135,000 we increased target sample size up to $N_{target} \approx 270,000$, taken as the maximum

observed sample size we could benchmark against. For all choices of target sample size, and

across each disease, we observe high accuracy rates (≥95%). However, the TPR was sensitive

to disease prevalence, reducing monotonically as the target sample size increased. Broadly,

TPR remained reasonable (≳ 60%) up to a 2-fold increase in sample size, except for the very

rare (case-control ratio of 1:669) thyroid gland cancer. This is due to fewer significant variants

being included in the assessment as a result of lower percentage of heritability explained, which

can artificially reduce the TPR for each new locus with relatively high odds ratios. Naturally,

TPR reduces as a function of decreasing disease prevalence, as re-sampling from fewer cases

can increase the variability in MAFs and thus isGWAS forecasting. We note that our theoretical

sub-sampling approach had better predictive capabilities, owing to the prevalence preserving

sampling strategy taken (**Methods and Supplementary Figure 19**).

We also assessed isGWAS's ability to extrapolate results when ILD were not available, using

a highly constrained version of the leapfrog re-sampler (**Supplementary Information**). In this

scenario, MAF and disease prevalence per variant are fixed, computed from the maximum

current sample (i.e., without sub-sampling), and the number of cases and controls are

proportionately increased to match the target sample size. We did this for two GWAS of

schizophrenia: (a) 2014 analyses with up to $N = 77,096$ (cases $= 33,640$, controls $=$

43456) European ancestry individuals [25]; and (b) the larger (and future) 2022 analyses with

up to $N = 130,644$ (cases $= 53386$, controls $= 77258$) European ancestry individuals [27].

We treat the 2014 study as the current sample size and the 2022 sample size as the target future

state, which we benchmark predictive performance against. The studies were selected because,

for each variant j, necessary data to run isGWAS, i.e., $\{N_j, N_j^*, MAF_j, MAF_j^*\}$, were made

publicly available. Note these data are pooled estimates, computed across all European cohorts.

Despite not accessing ILD, our results reveal reasonable concordance between isGWAS 2014

extrapolated results and the published analyses of 2022 (**Figure 6**, **Supplementary Figures 20-**

21, Supplementary Tables 13-16). Like our Biobank Japan analyses, we also used a more

stringent significance threshold ($p < 10^{-10}$) to help attenuate false positives, observing

improved overall performance by recovering a good TPR $\geq 70\%$ (**Supplementary Table 13**).

We do not report FDR as these cannot be accurately computed when filtering results based on

a p-value inclusion/exclusion threshold. Of the overlapping 608 clumped variants considered,

isGWAS-LRS identified 136 associations that were not yet deemed GWAS significant (i.e,.

$p > 5e-8$) in the 2014 study but later identified as significant in the 2022 study. Moreover,

of the 436 significant associations predicted by isGWAS, 75% overlap with observed

significant associations in 2022. isGWAS predicted an additional 74 associations as significant

that were not significant in 2022 – of those 52 were near the significance threshold with $p <$

$9e-07$. There were 121 variants not correctly predicted by the 2014 cohort. This could be due

to increased ethnical and relatedness heterogeneity in the 2022 cohort that was not present in

the 2014 analysis.

Computational performance and convergence details

isGWAS is an iterative algorithm whose convergence (i.e., ability to estimate model

parameters) depends on several tuning parameters (**Methods**). Using default parameter settings,

isGWAS-Firth converged in all real-data and simulated scenarios tested (**Figure 4e and**

Supplementary Tables 17, 19-20). Convergence was achieved in around 0.001 seconds per

variant (**Supplementary Table 21**) on a 2.4 GHz 8-Core Intel Core i9 processor. The non-Firth

corrected isGWAS algorithm may require more iterations, particularly for diseases with lower

prevalence (e.g., case:control ratio of 1:94 and lower) which included scenarios where

convergence was not achieved (**Figure 4e, Supplementary Tables 17-20, Supplementary**

Information).

When distributed over 32 CPU cores on a high-performance cluster, Firth-corrected isGWAS

analysed a single disease from UK Biobank across ~11 million SNPs and for ~335,000

individuals in ~4 minutes (**Figure 7, Supplementary Tables 9 and 18**). This means that

isGWAS-Firth can perform around 1,500 disease GWAS for every one GWAS performed using

an alternative methodology. The same analysis with a small number of CPU cores was

completed in tens of minutes using isGWAS-Firth (**Figure 7**). Further computational gains at

larger sample sizes will likely be achieved as ILD methods can scale poorly with sample size,

whereas isGWAS has near fixed computational cost at any size. As isGWAS currently

computes associations for each variant independently, additional improvements such as

parallelisation are possible. Full details are available in **Supplementary File 4.**

Discussion

In this study, we developed isGWAS, an efficient, biobank-scalable method for genetic

association testing which can: (a) compute regression parameters and test for a variant-disease

association in real-time (i.e., approximately one millisecond) for any sample size; (b) bypass

the need to run large-scale GWAS using high-performance computing facilities owing to ultra-

low system resource demands (i.e., runtime and memory); and (c) infer GWAS results from

virtually enlarged sample sizes using a novel re-sampling procedure. The isGWAS algorithm

design allows analyses to be run without the need to hold or access individual-level data (ILD)

directly, thereby providing a single methodological framework to utilise a wide range of data

sources such as published summary-level data from biobanks and repositories.

isGWAS draws inspiration from classical methodologies to overcome significant

computational bottlenecks associated with massive-scale analyses. The practical simplicity and

quick runtime of classical approaches have seen them deployed in a recent large-scale

analysis[23]. Rather than using ILD, as contemporary GWAS regression analyses do, isGWAS

distils the required input data down to sufficient statistics – a low-dimensional summary of ILD

that captures all necessary information required to compute a genetic-disease association model

parameter. In combination with modifications to the Newton-Raphson procedure, used to

estimate model parameters in a logistic regression, our use of sufficient statistics dramatically

reduces the computational time for disease association testing relative to existing methods.

Achieving up to a 1,500-fold improvement in computational runtime, when benchmarked

against a state-of-the-art GWAS tool, isGWAS reduced time to genome-wide insight from

several days down to ~4 minutes. Thereby unlocking potential for massive scale exploration of

genetic-disease associations in real-time and making feasible the routine assessment of

thousands of disease endpoints and studies. Computational bottlenecks associated with existing

GWAS methodologies are fast approaching. Analyses of resources such as UK Biobank WGS

data, the emerging massive cohorts of the Global Biobank Initiative[1], and Our Future

Health[33] are expected to push current GWAS tools to their system resource limits with

significant associated time-to-insight penalties. Conversely, with no computational sensitivity

to sample size, as the number of variants assessed and sample sizes continue to increase, the

relative savings and benefits of isGWAS can be expected to grow.

To attenuate possible issues of confounding and population stratification, we propose that

additional QC-steps are performed before computing sufficient statistics for isGWAS. In our

analyses, these steps reduced our UK Biobank sample size from ~408k individuals (used in the

original testing of REGENIE[8]) down to a more homogeneous sample of ~335k individuals

used to generate and compare results from isGWAS and REGENIE. Our results reveal an often-

striking concordance between approaches genome-wide as well as at the regional locus level.

The reduction in sample size was compensated by the isGWAS leapfrog re-sampler (LRS),

which we demonstrate efficiently helped extrapolate GWA results onto larger sample sizes (up

to 2-times). While we note sensitivity of an LRS extrapolation to disease prevalence, across the

range considered the TPR and FPR were well calibrated to at least a 1.5-fold increase in sample

size. The LRS might therefore be leveraged to aid GWAS cohort design, for example to

quantify the potential benefit of sampling more participants with a disease of interest against

cost. In our analyses of the PGC Schizophrenia cohort, we deployed a highly restricted (i.e., no

ILD) version of the LRS: forecasting results from a smaller 2014 cohort[25] onto a sample size

that matched a future 2022 study[27]. Despite no guarantees of sufficiency, isGWAS LRS

identified 75% of significant variants that were later identified in the larger 2022 cohort (almost

double the size) while maintaining a low FDR. Unlike extrapolation via the ILD leapfrog re-

sampler, this naïve extrapolation does not account for differences in the MAF of cases and

controls between 2014 and 2022 data. Regardless, the above findings highlight potential for

isGWAS to furnish reasonable forecasts of future results without accessing ILD directly. Our

recent predictions from FinnGen consortium data [18], [34] provide confidence that the

isGWAS algorithm is applicable also to multi-ethnic GWAS through analyzing each ethnicity

separately and combining results in a meta-analysis, as it is common practice[35].

Beyond, isGWAS can be applied to help address routinely asked questions about future

scenarios and evaluate enrichment contribution of biobanks to disease-specific associations[34]

or to protein-specific variant associations[18], particularly in the rare spectrum. isGWAS-Firth

provides a timely, rapid regression-based analysis of common, rare and ultra-rare variants.

Unlike ILD-based analyses, where Firth's correction significantly increases computational

time[7], [8], there is no computational penalty when using Firth's correction in the isGWAS

framework - in fact, we observe improved computational performance. The advantage of

considerable improvements in computational runtime is that it allows for the introduction of

forecasting, re-sampling and other non-parametric techniques - the LRS being one example.

These might widen robust association testing strategies as well as provide new avenues to tackle

confounding or population sub-structure. For now, we envisage the possibility that the wider

human genetics community routinely compute and make available the sufficient statistics, i.e.

MAF in the cases and the cohort, and the corresponding sample sizes per variant, toward a

publicly available, privacy compliant, data asset. In addition to avoiding the need for expensive

high-performance computing facilities and memory intensive data storage, the data asset might

enhance meta-analyses and biological insight, improve equitable access, and enable faster

collaborations between teams and help bridge financial and resource gaps between institutions

and research groups internationally.

Methods

Disease SNP association model

Let $S_M \in \{1,2\}$ be the maximum number of copies of the effect allele for an individual under model $M \in \{A, D, R\}$, where A denotes an additive model, R a recessive model and D a dominant model, i.e.,

$$S_M = \begin{cases} 2, & M = A, \\ 1, & M = D \cap R. \end{cases}$$

Furthermore let,

$$MAF_{j,M} = MAF_j \mid M$$

$$MAF_{j,M}^* = MAF_j^* \mid M$$

where, for a given model M, $MAF_{j,M}$ is the minor allele frequency for variant j in the sample, ancestry, or population and $MAF_{j,M}^*$ the minor allele frequency in the cases. We let Y denote disease status and G_j the jth genotype in the sample. For convenience we write $G_{j,M} = G_j \mid M$. It is assumed that the outcome model for Y, conditional on G_j, is given by:

$$\mathbb{E}[Y \mid G_j, M] = h^{-1}(\alpha_{j,M} + \beta_{j,M} G_{j,M}), \qquad j = 1, 2, \dots, Q,$$

where, conditional on model M, the pair $\{\alpha_{j,M}\ \beta_{j,M}\}$ denote the intercept and genotype effect and h is a function linking the outcome to genotype $G_{j,M}$ for all $j = 1, 2, \dots, Q$ genetic variants considered. In deriving the isGWAS estimation procedure we assume that h is the logit function, i.e.,

$$\pi_{Y \mid G_{j,M}} = P(Y = 1 \mid G_j, M) = \frac{e^{(\alpha_{j,M} + \beta_{j,M} G_{j,M})}}{1 + e^{(\alpha_{j,M} + \beta_{j,M} G_{j,M})}}.$$

The isGWAS methodology can, however, be broadened to other link functions and outcome types. We take $\{\hat{\alpha}_{j,M}, \hat{\beta}_{j,M}, \hat{\sigma}_{\alpha_{j,M}}, \hat{\sigma}_{\beta_{j,M}}\}$ to be sample based estimates of the intercept $\alpha_{j,M}$ and coefficient $\beta_{j,M}$, and their associated standard errors $\{\hat{\sigma}_{\alpha_{j,M}}, \hat{\sigma}_{\beta_{j,M}}\}$. We allow the genetic effect

416 $\beta_{j,M}$ and genotype G_j (or an observation thereof g_j) to be analyzed on either (i) the standardized

417 scale or (ii) non standardized scale. To accommodate this, we introduce the variable s_m^*, so that:

418
$$\{\beta_{j,s_m^*}, g_{j,s_m^*}\} = \left\{(1 + (\sigma_{g_i} - 1)s_m^*)\beta_j, \left.g_j\middle/(1 + (\sigma_{g_i} - 1)s_m^*)\right.\right\}.$$

419 Hence, when $s_m^* = 0$ analyses are performed on the non-standardized scale and $s_m^* = 1$ on the

420 standardized scale. Default analyses assume the genetic effect is assessed on the non-

421 standardized scale, i.e., $s_m^* = 0$. Note that, while p-values are generally invariant to the choice

422 of effect scale s_m^*, betas and standard errors are dependent on the specification of s_m^*.

Sample-Level Newton-Raphson (SaLN-R) algorithm

425 Here we detail the isGWAS procedure for computing summary statistics

426 $\left\{\hat{\alpha}_{j,M}, \hat{\beta}_{j,M,s_m^*}, \hat{\sigma}_{\alpha_{j,M}}, \hat{\sigma}_{\beta_{j,M,s_m^*}}\right\}$ using only four data points,

427
$$\left\{N_j, \sum_{i=1}^{N_j} y_i = N_j^*, \sum_{i=1}^{N_j} y_i g_{ij,M}, \sum_{i=1}^{N_j} g_{ij,M} = n_{j1} + 2n_{j2}I(M = A)\right\}$$

428 or, as we show, the quadruple

429
$$\{N_j, N_j^*, MAF_{j,M}, MAF_{j,M}^*\},$$

where N_j denotes the study or population sample size and N_j^* the number of cases in the sample,

see ad-hoc estimator (Supplementary Information) for definitions of $n_{j.}$. Note that we have

allowed the sample size N and number of cases N^* to vary by genotype j, this is useful when

emulating results from GWAS. This is because the number of individuals analyzed in GWAS

can vary by genotype owing to (e.g.,) quality of imputation or available data per variant and

participant in a study. Ideally the sample size and number of cases would not vary by genotype

and when using isGWAS to forecast GWAS results, users do not need not vary N_j, i.e,.

437
$$N_j = N \quad \text{and} \quad N_j^* = N^*, \quad j = 1,2,\dots,Q.$$

438 Given a vector of observed data $\{\boldsymbol{y}, \boldsymbol{g}_{j,s_m^*}\}$, where $\boldsymbol{y} = \{y_1, y_2, \dots, y_{N_j}\}$ and $\boldsymbol{g}_{j,s_m^*} =$

439 $\{g_{1j,}, g_{2j}, \dots, g_{N_j}\}$, estimates of model parameters are typically derived by maximizing the log-

440 likelihood function

$$L\left(\boldsymbol{\beta}_{j,M,s_m^*}\right) = \sum_{i=1}^{N_j} \log P\left(y_i \,|\boldsymbol{\beta}_{j,M,s_m^*}, g_{ij,M,s_m^*}\right),$$

441 which is equivalent to identifying parameter values $\boldsymbol{\beta}_{j,M,s_m^*} = \{\alpha_{j,M}, \beta_{j,M,s_m^*}\}$ which satisfy:

$$\frac{\partial L\left(\boldsymbol{\beta}_{j,M,s_m^*}\right)}{\partial \boldsymbol{\beta}_{j,M,s_m^*}} = V\left(\boldsymbol{\beta}_{j,M,s_m^*}, I_F\right) = 0,$$

444 where $V\left(\boldsymbol{\beta}_{j,M,s_m^*}, I_F\right)$ denotes the logistic score function, i.e.,

$$V\left(\boldsymbol{\beta}_{j,M,s_m^*}, I_F\right) = \widetilde{\boldsymbol{g}}_{j,M,s_m^*}^T \left(\boldsymbol{y} - \pi_{y|\widetilde{g}_{j,M,s_m^*}}\right) + I_F \mathrm{K}\left(\boldsymbol{\beta}_{j,M,s_m^*}\right) = 0,$$

with I_F denoting an indicator function used to highlight that a Firth modified version of the

score function has been used. For ease of mathematical presentation initially, we detail the Firth

adjusted SaLN-R algorithm later, i.e., we set $I_F = 0$ in this section. Additionally, to improve

succinctness of notation, we drop the use of the parameter s_m^* - reintroducing where necessary

- and set $\boldsymbol{\beta}_{j,M} = \{\alpha_{j,M}, \beta_{j,M}\}$ and $\widetilde{\boldsymbol{g}}_{j,M} = (1, g_{j,M})$ above, so that $\boldsymbol{\beta}_{j,M}\widetilde{\boldsymbol{g}}_{j,M}^T = \alpha_{j,M} + \beta_{j,M}g_{j,M}$.

We compute candidate solutions to by expanding $V\left(\boldsymbol{\beta}_{j,M}\right)$ as a Taylor series about a value

$\boldsymbol{\beta}_{j0,M}$ and up to second order, i.e., using the Newton-Raphson (N-R) method:

$$V\left(\boldsymbol{\beta}_{j,M}\right) = V\left(\boldsymbol{\beta}_{j0,M}\right) + \frac{\partial V\left(\boldsymbol{\beta}_{j,M}\right)}{\partial \boldsymbol{\beta}_{j,M}}\Bigg|_{\boldsymbol{\beta}_{j0,M}} \left(\boldsymbol{\beta}_{j,M} - \boldsymbol{\beta}_{j0,M}\right) + \mathcal{O}\left(\left(\boldsymbol{\beta}_{j,M} - \boldsymbol{\beta}_{j0,M}\right)^2\right),$$

454 Which is re-written as

$$\boldsymbol{\beta}_{j,M} = \boldsymbol{\beta}_{j0,M} + \mathcal{I}^{-1}\left(\boldsymbol{\beta}_{j0,M}\right)V\left(\boldsymbol{\beta}_{j0,M}\right) + \mathcal{O}\left(\left(\boldsymbol{\beta}_{j,M} - \boldsymbol{\beta}_{j0,M}\right)^2\right)$$

456 and generalized into an N-R iterative algorithm:

457 $\boldsymbol{\beta}_{j(k+1),M,s_m^*} = \boldsymbol{\beta}_{jk,M,s_m^*} + \mathcal{I}^{-1}\left(\boldsymbol{\beta}_{jk,M,s_m^*}\right)V\left(\boldsymbol{\beta}_{jk,M,s_m^*}\right) + \mathcal{O}\left(\left(\boldsymbol{\beta}_{j(k+1),M,s_m^*} - \boldsymbol{\beta}_{jk,M,s_m^*}\right)^2\right), \quad k = 0,1,\dots,K,$

458   where we have re-introduced s_m^* to highlight that the algorithm is dependent on the choice of

459   effect scale. The variable $\mathcal{J}^{-1}$ denotes the inverse Fisher Information matrix, where

460
$$\mathcal{J}(\boldsymbol{\beta}_{jk,M}) = -\left.\frac{\partial V(\boldsymbol{\beta}_{j,M})}{\partial \boldsymbol{\beta}_{j,M}}\right|_{\boldsymbol{\beta}_{j,M}=\boldsymbol{\beta}_{jk,M}}$$

461
$$= \left.\tilde{\boldsymbol{g}}_{j,M}^T \mathrm{diag}\left(\pi_{y_i|g_{j,M}}\left(1-\pi_{y_i|g_{jk,M}}\right)\right)\tilde{\boldsymbol{g}}_{j,M}\right|_{\boldsymbol{\beta}_{j,M}=\boldsymbol{\beta}_{jk,M}}$$

462
$$= \left.\begin{pmatrix} \sum_{i=1}^{N_j}\pi_{y_i|g_{ij,M}}\left(1-\pi_{y_i|g_{ij,M}}\right) & \sum_{i=1}^{N_j}\pi_{y_i|g_{ij,M}}\left(1-\pi_{y_i|g_{ij,M}}\right)g_{ij,M} \\ \sum_{i=1}^{N_j}\pi_{y_i|g_{ij,M}}\left(1-\pi_{y_i|g_{ij,M}}\right)g_{ij,M} & \sum_{i=1}^{N_j}\pi_{y_i|g_{ij,M}}\left(1-\pi_{y_i|g_{ij,M}}\right)g_{ij,M}^2 \end{pmatrix}\right|_{\boldsymbol{\beta}_{j,M}=\boldsymbol{\beta}_{jk,M}}$$

463   and the score function is given by

464
$$V(\boldsymbol{\beta}_{jk,M}) = \left.\tilde{\boldsymbol{g}}_{j,M}^T\left(\boldsymbol{y}-\pi_{y|g_{j,M}}\right)\right|_{\boldsymbol{\beta}_{,Mj}=\boldsymbol{\beta}_{jk,M}}$$

465
$$= \left.\begin{pmatrix} \sum_{i=1}^{N_j}\left(y_i-\pi_{y_i|g_{ij,M}}\right) \\ \sum_{i=1}^{N_j}\left(y_i-\pi_{y_i|g_{ij,M}}\right)g_{ij,M} \end{pmatrix}\right|_{\boldsymbol{\beta}_{j,M}=\boldsymbol{\beta}_{jk,M}}$$

466   Both $\mathcal{J}(\boldsymbol{\beta}_{jk,M})$ and $V(\boldsymbol{\beta}_{jk,M})$ above require individual-level data to compute their values.

467   isGWAS aims to estimate values for these variables using sample-level information only,

468   thereby avoiding the immediate need for individual data. To achieve this, we approximate both

469   the Fisher Information matrix and the Score function via the pair $\{\mathcal{J}_{\text{e}}(\boldsymbol{\beta}_{jk,M}), V_{\text{e}}(\boldsymbol{\beta}_{jk,M})\}$,

470   where:

471
$$\mathcal{J}_{\text{e}}(\boldsymbol{\beta}_{jk,M}) = N_j\begin{pmatrix} \mathbb{E}_{g_{j,M}}\left[\pi_{y|g_{j,M}}\left(1-\pi_{y|g_{j,M}}\right); \boldsymbol{\beta}_{jk,M}\right] & \mathbb{E}_{g_{j,M}}\left[\pi_{y|g_{j,M}}\left(1-\pi_{y|g_{j,M}}\right)g_{j,M}; \boldsymbol{\beta}_{jk,M}\right] \\ \mathbb{E}_{g_{j,M}}\left[\pi_{y|g_{j,M}}\left(1-\pi_{y|g_{j,M}}\right)g_{j,M}; \boldsymbol{\beta}_{jk,M}\right] & \mathbb{E}_{g_{j,M}}\left[\pi_{y|g_{j,M}}\left(1-\pi_{y|g_{j,M}}\right)g_{j,M}^2; \boldsymbol{\beta}_{jk,M}\right] \end{pmatrix}$$

472   and

473
$$V_{\text{e}}(\boldsymbol{\beta}_{jk,M}) = \begin{pmatrix} \sum_{i=1}^{N_j}y_i - N_j\,\mathbb{E}_{g_{j,M}}\left[\pi_{y|g_{j,M}}; \boldsymbol{\beta}_{jk,M}\right] \\ \sum_{i=1}^{N_j}y_ig_{ij,M} - N_j\,\mathbb{E}_{g_{j,M}}\left[\pi_{y|g_{j,M}}g_{j,M}; \boldsymbol{\beta}_{jk,M}\right] \end{pmatrix}$$

474
$$\approx \begin{pmatrix} N_j^* - N_j \, \mathbb{E}_{g_{j,M}} \left[\pi_{y|g_{j,M}}; \, \boldsymbol{\beta}_{jk,M} \right] \\ N_j^* \mathbb{E}_{g_{j,M}} [g_{j,M} | \, y = 1] - N_j \mathbb{E}_{g_{j,M}} \left[\pi_{y|g_{j,M}} g_{j,M}; \, \boldsymbol{\beta}_{jk,M} \right] \end{pmatrix}, \quad N_j^* \gg 1.$$

with $\mathbb{E}_{g_{j,M}}[\,\cdot\,; \boldsymbol{\beta}_{jk,M}]$ denoting that expectation is taken with respect to $g_{j,M}$ and evaluated at

$\boldsymbol{\beta}_{j,M} = \boldsymbol{\beta}_{jk,M}$. Note that $\{\mathcal{I}_{\text{e}}(\boldsymbol{\beta}_{jk,M}), V_{\text{e}}(\boldsymbol{\beta}_{jk,M})\}$ are motivated by switching from empirical, i.e.,

sample-based, estimates in $\{\mathcal{I}(\boldsymbol{\beta}_{jk,M}), V(\boldsymbol{\beta}_{jk,M})\}$ to their expected value analogues, which

reverses the usual mode of estimation. Sample size N_j is presumed large and thus switching

from sample-based to expected values in the N-R algorithm is well motivated. However, when

the number of cases N_j^* is 'small', an approximation of $\sum_{i=1}^{N_j} y_i g_{j,M} \approx N_j^* \mathbb{E}_{g_{j,M}} [g_{j,M} | \, y = 1]$

becomes weaker and we recommend using the statistic $\sum_{i=1}^{N_j} y_i g_{j,M}$. Values for the elements in

$\{\mathcal{I}_{\text{e}}(\boldsymbol{\beta}_{jk,M}), V_{\text{e}}(\boldsymbol{\beta}_{jk,M})\}$ are computed via:

484
$$\mathbb{E}_{g_{j,M,s_m^*}} \left[\pi_{y|g_{j,M,s_m^*}} \left(1 - \pi_{y|g_{j,M,s_m^*}} \right) g_{j,M,s_m^*}^c; \, \boldsymbol{\beta}_{jk,M,s_m^*} \right] =$$

485
$$\sum_{l=0}^{s_M} \pi_{y|g_{j,M}=l/w_{s_m^*}; \boldsymbol{\beta}_{jk,M,s_m^*}} \left(1 - \pi_{y|g_{j,M}=l/w_{s_m^*}; \boldsymbol{\beta}_{jk,M,s_m^*}} \right) \left(\frac{l}{w_{s_m^*}} \right)^c p \left(g_{j,M} = l/w_{s_m^*} \right)$$

486
$$= e_{j,M}^{(c,k)}$$

487 and

488
$$\mathbb{E}_{g_{j,M,s_m^*}} \left[\pi_{y|g_{j,M,s_m^*}} g_{j,M,s_m^*}^c \mid \boldsymbol{\beta}_{jk,M,s_m^*} \right] =$$

489
$$\sum_{l=0}^{s_M} \pi_{y|g_{j,M}=l/w_{s_m^*}; \boldsymbol{\beta}_{jk,M,s_m^*}} \left(\frac{l}{w_{s_m^*}} \right)^c p \left(g_{j,M} = l/w_{s_m^*} \right)$$

490
$$= \tilde{e}_{j,M}^{(c,k)},$$

491 where $w_{s_m^*} = \left(1 + (\sigma_{g_i} - 1) s_m^* \right)$ and the superscript and subscript in $e_M^{(c,k)}$, $\tilde{e}_M^{(c,k)}$ are used to

492 highlight that expectation has been taken conditional on k-th iteration $\beta_{jk,M}$ and under

493 modelling assumption M (and implicitly effect scale s_m^*). Probability mass $p\left(g_{j,M} = {}^l/w_{s_m^*}\right)$

494 is either defined a-priori or can be approximated empirically, which we detail later. In

495 combination, therefore, it follows that:

496
$$\mathcal{I}_{\mathbb{e}}\left(\boldsymbol{\beta}_{jk,M,s_m^*}\right) = N_j \begin{pmatrix} e_{j,M}^{(0,k)} & e_{j,M}^{(1,k)} \\ e_{j,M}^{(1,k)} & e_{j,M}^{(2,k)} \end{pmatrix}$$

497 and

498
$$\mathcal{I}_{\mathbb{e}}^{-1}\left(\boldsymbol{\beta}_{jk,M,s_m^*}\right) = \frac{1}{N_j\left(e_{j,M}^{(0,k)} e_M^{(2,k)} - \left(e_{j,M}^{(1,k)}\right)^2\right)} \begin{pmatrix} e_{j,M}^{(2,k)} & -e_{j,M}^{(1,k)} \\ -e_{j,M}^{(1,k)} & e_{j,M}^{(0,k)} \end{pmatrix}.$$

499 Following the same process that led to the above, we re-write the sample-level Score function

500 $V_{\mathbb{e}}\left(\boldsymbol{\beta}_{jk,M,s_m^*}\right)$ as:

501
$$V_{\mathbb{e}}\left(\boldsymbol{\beta}_{jk,M,s_m^*}\right) = \begin{pmatrix} N_j^* - N_j\, \tilde{e}_{j,M}^{(0,0)} \\ \sum_{i\,:\,y_i=1} g_{ij,M,s_m^*} - N_j \tilde{e}_{j,M}^{(1,0)} \end{pmatrix}$$

502
$$\approx \begin{pmatrix} N_j^* - N_j\, \tilde{e}_{j,M}^{(0,0)} \\ \dfrac{S_M N_j^* MAF_{j,M}^*}{w_{s_m^*}} - N_j \tilde{e}_{j,M}^{(1,0)} \end{pmatrix}, \quad N_j^* \gg 1,$$

503 where we have used the following approximation:

504
$$\sum_{i\,:\,y_i=1} g_{ij,M,s_m^*} \approx N_j^* \mathbb{E}_{g_{j,M}}\left[g_{j,M,s_m^*} \mid y = 1\right] = \frac{S_M N_j^* MAF_{j,M}^*}{w_{s_m^*}}.$$

505 Hence, replacing the pair $\left\{\mathcal{I}\left(\boldsymbol{\beta}_{jk,M,s_m^*}\right), V\left(\boldsymbol{\beta}_{jk,M,s_m^*}\right)\right\}$ with the sample-level approximations

506 $\left\{\mathcal{I}_{\mathbb{e}}\left(\boldsymbol{\beta}_{jk,M,s_m^*}\right), V_{\mathbb{e}}\left(\boldsymbol{\beta}_{jk,M,s_m^*}\right)\right\}$ we furnish the SaLN-R algorithm:

507
$$\boldsymbol{\beta}_{j(k+1),M,s_m^*} = \boldsymbol{\beta}_{jk,M,s_m^*} + \mathcal{I}_{\mathbb{e}}^{-1}\left(\boldsymbol{\beta}_{jk,M,s_m^*}\right) V_{\mathbb{e}}\left(\boldsymbol{\beta}_{jk,M,s_m^*}\right)$$

508
$$= \boldsymbol{\beta}_{jk,M,s_m^*} + \frac{1}{N_j\left(e_{j,M}^{(0,k)} e_{j,M}^{(2,k)} - \left(e_{j,M}^{(1,k)}\right)^2\right)} \begin{pmatrix} e_{j,M}^{(2,k)} & -e_{j,M}^{(1,k)} \\ -e_{j,M}^{(1,k)} & e_{j,M}^{(0,k)} \end{pmatrix} \begin{pmatrix} N_j^* - N_j\, \tilde{e}_{j,M}^{(0,k)} \\ \dfrac{S_M N_j^* MAF_{j,M}^*}{w_{s_m^*}} - N_j \tilde{e}_{j,M}^{(1,k)} \end{pmatrix}.$$

509 The standard error of the updates, $\widehat{\boldsymbol{\sigma}}_{\boldsymbol{\beta}_{j(k+1),M,s_m^*}} = \left\{\hat{\sigma}_{\alpha_{j(k+1),M,s_m^*}}, \hat{\sigma}_{\beta_{j(k+1),M,s_m^*}}\right\}$, are given by the

510 diagonal of the inverse Fisher information matrix, i.e.,

$$511 \qquad \hat{\sigma}_{\alpha_{j(k+1),M,s_m^*}} = \sqrt{\frac{e_{j,M}^{(2,k)}}{N_j\left(e_{j,M}^{(0,k)}e_{j,M}^{(2,k)} - \left(e_{j,M}^{(1,k)}\right)^2\right)}},$$

$$512 \qquad \hat{\sigma}_{\beta_{j(k+1),M,s_m^*}} = \sqrt{\frac{e_{j,M}^{(0,k)}}{N_j\left(e_{j,M}^{(0,k)}e_{j,M}^{(2,k)} - \left(e_{j,M}^{(1,k)}\right)^2\right)}}.$$

513   It can be shown from the above that:

$$514 \qquad \boldsymbol{\beta}_{jk,M,s_m^*=1} = \sigma_{g_i}\boldsymbol{\beta}_{jk,M,s_m^*=0},$$

$$515 \qquad \hat{\sigma}_{\beta_{j(k+1),M,s_m^*=1}} = \sigma_{g_i}\hat{\sigma}_{\beta_{j(k+1),M,s_m^*=0}}.$$

516   We set $s_m^* = 0$ to compute values for the pair $\left\{\hat{\boldsymbol{\beta}}_{jk,M,0}, \hat{\sigma}_{\beta_{j(k+1),M,0}}\right\}$ and use the above identities

517   to return parameter estimates on the standardized scale $s_m^* = 1$. The data required to run the

518   SaLN-R algorithm are:

$$519 \qquad \left\{N_j, N_j^*, \sum_{i=1}^{N_j} y_i g_{ij,M}, p\big(g_{j,M} = 1\big)\right\} \bigcup \begin{cases} \emptyset, & S_M = 1, \\ p\big(g_{j,M} = 2\big), & S_M = 2. \end{cases}$$

520   We use the approximations

$$521 \qquad p\big(g_{j,M} = 1\big) \approx \frac{n_{j1}}{N_j} \xrightarrow[\substack{N_j \gg 1 \\ HWE}]{} \begin{cases} MAF_{j,M}, & S_M = 1, \\ 2MAF_{j,M}\big(1 - MAF_{j,M}\big), & S_M = 2, \end{cases}$$

$$522 \qquad p\big(g_{j,M} = 2\big) \approx \frac{n_{j2}}{N_j} \xrightarrow[\substack{N_j \gg 1 \\ HWE}]{} MAF_{j,M}^2,$$

523   where $\xrightarrow[HWE]{}$ is used to denote under Hardy-Weinberg equilibrium.

524   The SaLN-R algorithm is extended to include Firth's penalty function (see **Supplementary**

525   **Information for more details**):

$$526 \qquad \boldsymbol{\beta}_{j(k+1),M,s_m^*} = \boldsymbol{\beta}_{jk,M,s_m^*} + \frac{1}{\left(\left(e_M^{(0,k)} - \frac{\partial_{\alpha_{j,M}}K_{\oplus}^{(\alpha_{j,M})}}{N_j}\right)\left(e_M^{(2,k)} - \frac{\partial_{\beta_{j,M}}K_{\oplus}^{(\beta_{j,M})}}{N_j}\right) - \left(e_M^{(1,k)} - \frac{\partial_{\beta_{j,M}}K_{\oplus}^{(\alpha_{j,M})}}{N_j}\right)^2\right)}$$

$$\times \begin{pmatrix} e_M^{(2,k)} - \dfrac{\partial_{\alpha_{j,M}} K_\text{e}^{(\alpha_{j,M})}}{N_j} & -e_M^{(1,k)} + \dfrac{\partial_{\beta_{j,M}} K_\text{e}^{(\alpha_{j,M})}}{N_j} \\[3mm] -e_M^{(1,k)} + \dfrac{\partial_{\beta_{j,M}} K_\text{e}^{(\alpha_{j,M})}}{N_j} & e_M^{(0,k)} - \dfrac{\partial_{\beta_{j,M}} K_\text{e}^{(\beta_{j,M})}}{N_j} \end{pmatrix} \begin{pmatrix} \pi_j^* - \tilde{e}_M^{(0,k)} + K_\text{e}^{(\alpha_{j,M})} \big/ N_j \\[3mm] \dfrac{S_M \pi_j^* MAF_{j,M}^*}{w_{s_m^*}} - \tilde{e}_M^{(1,k)} + K_\text{e}^{(\beta_{j,M})} \big/ N_j \end{pmatrix},$$

529 where

530
$$K_\text{e}(\boldsymbol{\beta}_{j,M}) = \frac{1}{2\left(e_{j,M}^{(0,k)} e_{j,M}^{(2,k)} - \left(e_{j,M}^{(1,k)}\right)^2\right)} \begin{pmatrix} d_{\alpha,j,M}^{(0,k)} e_{j,M}^{(2,k)} + d_{\alpha,j,M}^{(2,k)} e_{j,M}^{(0,k)} - 2 d_{\alpha,j,M}^{(1,k)} e_{j,M}^{(1,k)} \\[2mm] d_{\beta,j,M}^{(0,k)} e_{j,M}^{(2,k)} + d_{\beta,j,M}^{(2,k)} e_{j,M}^{(0,k)} - 2 d_{\beta,j,M}^{(1,k)} e_{j,M}^{(1,k)} \end{pmatrix}$$

531 and

532
$$d_{\alpha,j,M}^{(c+1,k)} = d_{\beta,j,M}^{(c,k)} = \frac{\partial e_{j,M}^{(c,k)}}{\partial \beta_{j,M}}.$$

isGWAS is computed using sufficient statistics

535 Under Hardy-Weinberg equilibrium, the quadruple $\left\{N_j, N_j^*, MAF_{j,M}, MAF_{j,M}^*\right\}$ are combined to

536 form the global and local (under a wide radius of convergence) sufficient statistics from the

537 logistic model. Consequently, they hold all necessary information to compute regression

538 parameter estimates $\left\{\hat{\alpha}_{j,M}, \hat{\beta}_{j,M,s_m^*}, \hat{\sigma}_{\alpha_{j,M}}, \hat{\sigma}_{\beta_{j,M,s_m^*}}\right\}$ over a broad range of scenarios. Regardless

539 of Hardy-Weinberg being valid or not, we show that the triple $\left\{T_{1j}, T_{2j}, T_{3j}\right\}$,

540
$$\left\{T_{1j} = \sum_{i=1}^{N_j} y_i = N_j^*, \qquad T_{2j} = \sum_{i=1}^{N_j} y_i\, g_{ij,M,s_m^*}, \qquad T_{3j} = \sum_{i=1}^{N_j} g_{ij,M,s_m^*}\right\},$$

541 are the two global and one local sufficient statistics and these can alternatively be used as input

542 variables in isGWAS. To show this, we write:

543
$$L\left(\boldsymbol{\beta}_{j,M,s_m^*}\right) = \sum_{i=1}^{N_j} \log P\left(y_i \mid \boldsymbol{\beta}_{j,M,s_m^*}, g_{ij,M,s_m^*}\right)$$

$$544 \qquad = \alpha_{j,M,s_m^*} \sum_{i=1}^{N_j} y_i + \beta_{j,M,s_m^*} \sum_{i=1}^{N_j} y_i\, g_{ij,M,s_m^*}$$

$$545 \qquad + \sum_{i=1}^{N_j} \log\left(1 - P\left(y_i = 1 \,|\, \boldsymbol{\beta}_{j,M,s_m^*}, g_{ij,M,s_m^*}\right)\right)$$

$$546 \qquad = \alpha_{j,M,s_m^*} T_{1j} + \beta_{j,M,s_m^*} T_{2j} - \sum_{i=1}^{N_j} \log\left(1 + \exp\alpha_{j,M,s_m^*}\right)$$

$$547 \qquad - \sum_{i=1}^{N_j} \log\left(1 + \frac{\exp\alpha_{j,M,s_m^*}}{1 + \exp\alpha_{j,M,s_m^*}}\left(\left(\exp\beta_{j,M,s_m^*}\, g_{ij,M,s_m^*}\right) - 1\right)\right)$$

$$548 \qquad = \alpha_{j,M,s_m^*} T_{1j} + \beta_{j,M,s_m^*}\left(T_{2j} - \frac{\exp\alpha_{j,M,s_m^*}}{1 + \exp\alpha_{j,M,s_m^*}} T_{3j}\right) - N_j \log\left(1 - \exp\alpha_{j,M,s_m^*}\right)$$

$$549 \qquad + \mathcal{O}\left(\frac{\exp\alpha_{j,M,s_m^*}}{1 + \exp\alpha_{j,M,s_m^*}}\left(\beta_{j,M,s_m^*} g_{\cdot j,M,s_m^*}\right)^2\right)$$

$$550 \qquad = f\left(T_{1j}, T_{2j}, T_{3j}; \alpha_{j,M,s_m^*}, \beta_{j,M,s_m^*}\right) + \mathcal{O}\left(\frac{\exp\alpha_{j,M,s_m^*}}{1 + \exp\alpha_{j,M,s_m^*}}\left(\beta_{j,M,s_m^*} g_{\cdot j,M,s_m^*}\right)^2\right)$$

551 and valid when

$$552 \qquad \frac{\exp\alpha_{j,M,s_m^*}}{1 + \exp\alpha_{j,M,s_m^*}}\left|\left(\exp\beta_{j,M,s_m^*}\, g_{\cdot j,M,s_m^*}\right) - 1\right| < 1.$$

553 Hence, the global sufficient statistics are $\{T_{1j}, T_{2j}\}$ and (on assuming random g_{ij,M,s_m^*} as in the

554 SaLN-R algorithm) the locally sufficient statistic is $\{T_{3j}\}$, where:

$$555 \qquad T_{1j} = \sum_{i=1}^{N_j} y_i = N_j^*, \qquad T_{2j} = \sum_{i=1}^{N_j} y_i\, g_{ij,M,s_m^*}$$

556 and

$$557 \qquad T_{3j} = \sum_{i=1}^{N_j} g_{ij,M,s_m^*} = \begin{cases} n_{j1}, & S_M = 1, \\ n_{j1} + 2n_{j2}, & S_M = 2. \end{cases}$$

558 Under Hardy-Weinberg equilibrium, we can write

$$559 \qquad T_{2j} = s_m^* N_j^* MAF_j^* \quad \text{and} \quad T_{3j} = s_m^* N_j MAF_j.$$

Leapfrog re-sampler: forecasting results in target sample sizes

To estimate regression parameters $\{\alpha_{j,M,s_m^*}, \beta_{j,M,s_m^*}\}$ in larger target sample sizes, i.e., $\widehat{N}_j > N_j$, we propose the following strategy:

1. **Specify number K, sub-sample γ_1 and target sample γ_2 parameters**, where $K \geq 1$, $0 < \gamma_1 < 1$ and $\gamma_2 > 1$.

2. **Generate random sub-samples of individuals of size** $\widetilde{N}_j = \gamma_1 N_j < N_j$. For each of $k = 1, 2, \ldots, K$, generate a random sub-sample $D_{k,\gamma_1} \subset D$, where $\left| D_{k,\gamma_1} \right| = \widetilde{N}_j = \gamma_1 N_j$.

3. **(Leapfrog-step) Compute subsample quadruple and project to target sample size** $\widehat{N}_j$. For each subsample D_{k,γ_1}, compute values $\left\{ \widetilde{N}_{ij}^*, \widehat{MAF}_{kj,M}, \widehat{MAF}^*_{kj,M} \right\}$ and project these on to the target sample size, i.e., $d_{k,\gamma_{1,2}} = \left\{ \left(\frac{\gamma_2}{\gamma_1}\right) \widetilde{N}_{kj}, \left(\frac{\gamma_2}{\gamma_1}\right) \widetilde{N}_{ij}^*, \widehat{MAF}_{kj,M}, \widehat{MAF}^*_{kj,M} \right\}$ for sample D_{k,γ_1}

   - Note that $\left(\frac{\gamma_2}{\gamma_1}\right) \widetilde{N}_{kj} = \widehat{N}_j$, which is the target 'future' sample size.

4. **Deploy isGWAS across all K (projected) quadruples** $d_{k,\gamma_{1,2}}$ and record each estimate of the genetic effects, standard error and p-value $\left\{ \hat{\beta}_{k,j,M,s_m^*}, \hat{\sigma}_{\beta_{k,j,M,s_m^*}}, p_{k,j,M,s_m^*} \right\}_{k=1:K}$.

5. **Estimate p-value in target sample size** as a summary point estimate (e.g., median) or range across all K sub-samples, $$p_{target_{j,M,s_m^*}} = \text{median}\{p_{k,j,M,s_m^*}\}_{k=1:K}.$$

Data Quality Control: preparation of sufficient statistics for isGWAS

In order to deploy isGWAS successfully, the sufficient statistics are required to be prepared in a sample where only a single individual (preferably case) from pairs or n-tuples of 3^{rd}, 2^{nd} and 1^{st} degree relatives is retained. Additionally, ethnical outliers must also be removed. In summary, to deploy isGWAS successfully we require either: (a) access to the sufficient statistics computed after duplications of related n-tuples and ethnical outliers are removed; or (b) access to the individual level data, whereupon the sufficient statistics can be prepared as described in (a). We provide a detailed outline of recommended Quality Control for genetic Individual Level Data (ILD) to running successfully isGWAS in **Supplementary Information**.

Application to Biobank data

The GWAS results used in the assessment of isGWAS were taken from large-scale analyses of UK Biobank [13], Biobank Japan [14] and the Psychiatric Genomics Consortium [15].

The UK Biobank [13] is a large-scale biomedical database and research resource containing in-depth genetic and health information from half a million UK participants. From the full available UK Biobank cohort, we obtain phenotypes for seven different diseases with varying levels of prevalence. These are Hypertension (IC10:I10), Asthma (IC10:J45), Atherosclerosis (IC10:I25), Glaucoma (IC10:H40), Stroke (IC10:I63), Colon Cancer (IC10:C18) and Thyroid Gland Cancer (IC10:C73) patients. From a total cohort of 502,422 participants, we used the following inclusion criteria: white British (Field 22006), non-related (>3rd degree), no patients with difference in reported (Field 31) and genetic (Field 22001) sex, no patients with aneuploidy (Field 22019), no patients with unusual heterozygosity and high missing rates

(Field 22027). The ethnicity component is obtained from samples who self-identified as

'White British' according to Field 21000 and have very similar genetic ancestry based on a

principal components analysis of the genotypes. Retaining one related individual (where we

favour the retention of cases) we obtain a working sample size of ~335,000 individuals; the

approximate value is owing to small differences in the number of cases between disease

phenotypes (**Supplementary Information**). Comparative analysis for these varying

populations is reported in the main text. The prevalence ratios and exact number of cases and

controls are provided in **Supplementary Table 1**. The variant based statistics needed for

isGWAS were obtained from the imputed UK Biobank dataset. A quality info score>0.9 is

applied to the data, and the number of cases and controls per variant and the MAF for variant

in cases and controls is based on patients with non-missing genotypes for the variant using

software PLINK [4]. Sample-level MAF>0.001 is used as inclusion criteria for the variants to

analyse. For each disease, we run isGWAS analysis using default settings under the 'additive'

genetic model. In addition, we also perform GWAS analysis using two-step REGENIE [8]

applied to all variants with a MAF>0.001 and Genotype Score>0.99. Firth correction was

enabled and performed on variants with p-value<0.1. REGENIE was also adjusted for

covariate information (age, gender, ancestry). For each disease we provide the following

diagnostic plots: 1) mirrored Manhattan plot comparing directly p-values for isGWAS and

REGENIE, 2) p-value – p-value plots comparing REGENIE and isGWAS, 3) $\beta - \beta$ plots

comparing REGENIE and isGWAS where we have colored the values by a) MAF and b)

ratios of computed standard error (SE) between methods, i.e., $log_2(\frac{SE(isGWAS)}{SE(REGENIE)})$. Across all

diseases and variants considered, we compare performance of isGWAS and isGWAS-Firth to

REGENIE-Firth.

Schizophrenia data from the Psychiatric Genomics Consortium [15] was used to conduct two

different large-scale GWAS analysis. The first GWAS analysis was executed with data from

77,096 European individuals (33,640 cases, 43,456 controls) [25]. The second GWAS analysis

was executed with data from the larger 130,644 European individuals (53,386 cases, 77,258

controls) [27]. We used the 2014 dataset to infer the 2022 results. To do this, we refine the

significant results from both imputed 2014 and 2022 summary statistics [27] using clumping

with PLINK [4]. The European 1000 Genomes Project v3 [19] dataset was used as a reference

population for the clumping procedure. Twelve strategies for clumping were explored: three

were LD R^2-based only, the other nine were a combination of clumping by LD block

information and p-value thresholding. The refined variants are used to assess the inference

capabilities of isGWAS both within each of the two datasets and the enrichment capabilities of

isGWAS to infer p-values of the 2022 dataset using the 2014 dataset. For the 2014 dataset, 225

variants were remaining after the clumping. For the 2022 dataset, 451 variants were remaining

after the clumping. From those, 54 are overlapping and 608 is the unique set between the two

datasets.

The Biobank Japan data was used to conduct a large-scale GWAS with 212,453 Japanese

individuals across 42 different diseases [24]. We obtained the published significantly associated

loci (P < 5e-08) in autosomes from the GWAS findings which amounted to 309 variants across

30 different diseases. Similarly, we used the significantly associated X chromosome findings

for males and females that amounted to a total of nine significantly associated loci across five

diseases, although results are omitted from text. We applied isGWAS to the three different sets

of variants using default parameters to assess the performance of isGWAS. To aid association

interpretation, we use the following additional statistical tests to assess the accuracy and

sensitivity of the isGWAS calculator for the Biobank Japan data. First, a classical ROC curve

was produced where the true/false actual value was determined by various p-value thresholds

(benchmarked against published Biobank Japan results). The isGWAS calculator is an

inferential tool thus this usage of the ROC curve is unconventional, however, it provides us

with the opportunity to assess the sensitivity to the choice of thresholds used to correct for

multiple testing. These are 10^{-10}, 10^{-8}, 5×10^{-8}, 10^{-7}, where we have also used

9.58×10^{-9} for Biobank Japan as recommended by the authors [24]. AUC values were not

obtained as this is not a standard classification problem and they are not interpretable in this

context. Second, an adapted ROC curve was produced which accounts for two different

thresholds – one more stringent one to determine the true positive rate and one less stringent

one to determine the true negative rate. **Supplementary Figure 22** showcases this scenario and

highlights the importance of a threshold choice and its impact on a sensitivity analysis. The

main aim of isGWAS calculator is to be used as an inferential tool for truly significant or truly

non-significant genetic signals. Thus, using two thresholds – one for truly significant and one

for truly non-significant – provides us the assess the sensitivity of isGWAS to this scientific

question. Third, the obtained β values were compared to the true ones by obtaining the

percentage of 1) predicted β values in the 95% C.I.s of the true β values and 2) 95% C.I.s of

the predicted β values in the 95% C.I.s of the true β values.

Simulation scenarios

In the first scenario, for each individual i and iteration index k, we randomly generate disease

status via $y_{ik} \sim Ber(\pi_{ik}; \alpha_k, \beta)$ with probability of disease $\pi_{ik} = expit(\alpha_k + \beta g_{ik})$ and

$g_{ik} \sim Bin(2, MAF_k)$. Minor allele frequency is randomly selected from the set $MAF_k \in$

$\{10^{-4}, 5 \times 10^{-4}, 0.01\} \cup \{0.025, 0.05, ..., 0.5\}$ and the genetic effect on disease risk is fixed as

$\beta = 0.5$. In the second study, we allow the genetic effect to vary, i.e., $\beta \equiv \beta_k$, by fixing disease

status per individual and generating genotype data in controls $g_{ik}|y_{ik} = 0 \sim Bin(2, MAF_k)$ or

cases $g_{ik}|y_{ik} = 1 \sim Bin(2, MAF_k^*)$, where minor allele frequency in cases is taken as the outer

product with the sample minor allele frequency, with a random increase or decrease in

frequency (which controls the magnitude and direction of genetic effect), i.e., we introduce the

set $MAF_k^* \in MAF_k \otimes (1 \pm MAF_k)$. The parameter β_k is then estimated via each of the 5

estimators using the vector of simulated data $\{\mathbf{y}_k, \mathbf{g}_k\}$.

We compare isGWAS and isGWAS-Firth against classical logistic and Firth corrected

regression [16], [36], [37]. Details for the second scenario are provided alongside full

description of the simulation protocol in the **Supplementary Information**.

Leapfrog re-sampler: simulation and real-data analyses

The parameters $\{K, \gamma_1, \gamma_2\}$ in the leapfrog re-ampler are assessed over a variety of values. To

attenuate the computational burden of a 3-dimensional grid search, we considered scenarios in

which: $K = 100$, $\gamma_1 = {}^1/_{\gamma_2}$ and a γ_2-fold increase in sample size of $\gamma_2 \in$

$\{1.1, 1.25, 1.5, \ldots, 2.5\}$, i.e., a 10% to 150% increase in sample size. Furthermore, we take our

working sample size to be 276,204 individuals, which matches the number of all unrelated

individuals in our UKB sample (i.e., on not retaining any member of a related pair – which is

therefore fixed between diseases). We used our simulation protocol (**Supplementary**

Information) to generate synthetic samples and additionally assessed performance across all

7 disease datasets in UK Biobank. Variants for assessment were selected after pruning in

PLINK [4] was applied to the ~11 million variants with the following parameters: genotype

quality>0.99, MAF>0.01, HWE $p < 10e-15$, 1000 bp windows, 100 variant increments,

$R^2 > 0.9$. From the pruned variants, 5% were selected uniformly from variants with $p >$

10^{-6} and all variants with $p \leq 10^{-6}$ were retained. Final number of variants progressed for

LRS for the seven diseases are provided in **Supplementary Table 10**. For simulated data,

data for smaller sub-samples were simulated using full cohort and empirical distributions for

MAF and disease prevalence. In our tests of the LRS, we assess the predictive properties of

isGWAS on real-life data where the ground truth is either computed from the entire sample or

provided in the literature. Predictions from X-fold increases in sample size are compared

using standard accuracy, FDR, FPR and TPR measures based on a putative true significance

threshold of $5e - 08$.

Computational resources

Real-life analyses were performed using up to 48 virtual CPU cores of a 2.5 GHz Intel Xeon

Gold 6240R processor with 64 GB of memory. Simulation analyses were performed using up

to 8 virtual CPU cores of a 2.4 GHz Intel Core i9 processor.

Computational comparison protocol

We contrast the computational performance of isGWAS and REGENIE (Step-2 only). For

clarity, REGENIE Step-1 simplifies the outcome and model by projecting out covariate

information, before variant-disease association analyses are performed in Step-2. To directly

compare both methods, we performed individual GWA analyses of each of the seven diseases

considered in UK Biobank across ~11m variants for ~335,000 individuals. Owing to

computational cost of the ILD method, we summarise results from a single GWA analysis per

trait. Performance of isGWAS across repeated runs, for varying numbers of SNPs and

available CPUs, up to a maximum of 10m variants, is also performed.

Data availability

718   The genotype data, phenotype status and allele counts were extracted from UK Biobank [13] to

719   support the findings of this study. The genome-wide association summary data with available

720   allele frequencies and cohort counts that was used to support the findings of this study are

721   available from: Psychiatric Genomics Consortium [15] and Biobank Japan [14].

Code availability

723   The tool is available for use on the webportal www.optima-isgwas.com. The isGWAS

724   algorithm is also available on github (https://github.com/cnfoley/isgwas/).

References

[1]   W. Zhou *et al.*, 'Global Biobank Meta-analysis Initiative: Powering genetic discovery across human disease', *Cell genomics*, vol. 2, no. 10, Oct. 2022, doi: 10.1016/J.XGEN.2022.100192.

[2]   M. Claussnitzer *et al.*, 'A brief history of human disease genetics', *Nature*, vol. 577, no. 7789, pp. 179–189, Jan. 2020, doi: 10.1038/S41586-019-1879-7.

[3]   A. Abdellaoui, L. Yengo, K. J. H. Verweij, and P. M. Visscher, '15 years of GWAS discovery: Realizing the promise', *Am J Hum Genet*, vol. 110, no. 2, pp. 179–194, Feb. 2023, doi: 10.1016/J.AJHG.2022.12.011.

[4]   C. C. Chang, C. C. Chow, L. C. A. M. Tellier, S. Vattikuti, S. M. Purcell, and J. J. Lee, 'Second-generation PLINK: Rising to the challenge of larger and richer datasets', *Gigascience*, vol. 4, no. 1, Feb. 2015, doi: 10.1186/s13742-015-0047-8.

[5]   L. Jiang *et al.*, 'A resource-efficient tool for mixed model association analysis of large-scale data', *Nature Genetics 2019 51:12*, vol. 51, no. 12, pp. 1749–1755, Nov. 2019, doi: 10.1038/s41588-019-0530-8.

[6]   P. R. Loh *et al.*, 'Efficient Bayesian mixed-model analysis increases association power in large cohorts', *Nature Genetics 2015 47:3*, vol. 47, no. 3, pp. 284–290, Feb. 2015, doi: 10.1038/ng.3190.

[7]   W. Zhou *et al.*, 'Efficiently controlling for case-control imbalance and sample relatedness in large-scale genetic association studies', *Nature Genetics 2018 50:9*, vol. 50, no. 9, pp. 1335–1341, Aug. 2018, doi: 10.1038/s41588-018-0184-y.

748   [8]   J. Mbatchou *et al.*, 'Computationally efficient whole-genome regression for

749   quantitative and binary traits', *Nat Genet*, vol. 53, no. 7, pp. 1097–1103, Jul.

750   2021, doi: 10.1038/S41588-021-00870-7.

751   [9]   N. Homer *et al.*, 'Resolving Individuals Contributing Trace Amounts of DNA to

752   Highly Complex Mixtures Using High-Density SNP Genotyping Microarrays',

753   *PLoS Genet*, vol. 4, no. 8, p. e1000167, Aug. 2008, doi:

754   10.1371/JOURNAL.PGEN.1000167.

755   [10]   X. Shi and X. Wu, 'An overview of human genetic privacy', *Ann N Y Acad Sci*,

756   vol. 1387, no. 1, p. 61, Jan. 2017, doi: 10.1111/NYAS.13211.

757   [11]   E. Wright Clayton *et al.*, 'The law of genetic privacy: applications, implications,

758   and limitations', *J Law Biosci*, vol. 6, no. 1, pp. 1–36, Oct. 2019, doi:

759   10.1093/JLB/LSZ007.

760   [12]   Z. Wan, J. W. Hazel, E. W. Clayton, Y. Vorobeychik, M. Kantarcioglu, and B.

761   A. Malin, 'Sociotechnical safeguards for genomic data privacy', *Nature Reviews

762   Genetics 2022 23:7*, vol. 23, no. 7, pp. 429–445, Mar. 2022, doi:

763   10.1038/s41576-022-00455-y.

764   [13]   'UK Biobank - UK Biobank'. https://www.ukbiobank.ac.uk/ (accessed Aug. 17,

765   2022).

766   [14]   'BioBank Japan'. https://biobankjp.org/en/ (accessed Aug. 12, 2022).

767   [15]   'PGC – Psychiatric Genomics Consortium'. https://pgc.unc.edu/ (accessed Aug.

768   12, 2022).

769   [16]   D. Firth, 'Bias Reduction of Maximum Likelihood Estimates', *Source:

770   Biometrika*, vol. 80, no. 1, pp. 27–38, 1993, Accessed: Mar. 14, 2023. [Online].

771   Available:

http://www.jstor.orgURL:http://www.jstor.org/stable/2336755Accessed:22/05/2

00814:38

[17]   P. D. Sasieni, 'From Genotypes to Genes: Doubling the Sample Size',

Biometrics, vol. 53, no. 4, p. 1253, Dec. 1997, doi: 10.2307/2533494.

[18]   B. B. Sun *et al.*, 'Genetic associations of protein-coding variants in human

disease', *Nature*, vol. 603, no. 7899, pp. 95–102, Mar. 2022, doi:

10.1038/S41586-022-04394-W.

[19]   '1000 Genomes | A Deep Catalog of Human Genetic Variation'.

https://www.internationalgenome.org/ (accessed Aug. 16, 2022).

[20]   'Estonian Biobank'. https://genomics.ut.ee/en/content/estonian-biobank

(accessed Aug. 12, 2022).

[21]   'FinnGen research project is an expedition to the frontier of genomics and

medicine | FinnGen'. https://www.finngen.fi/en (accessed Aug. 12, 2022).

[22]   R. A. Fisher, 'Statistical methods for research workers.', *Statistical methods for*

research workers., no. 5th Ed, 1934.

[23]   K. Ishigaki *et al.*, 'Large-scale genome-wide association study in a Japanese

population identifies novel susceptibility loci across different diseases', *Nat*

Genet, vol. 52, no. 7, pp. 669–679, Jul. 2020, doi: 10.1038/s41588-020-0640-3.

[24]   S. Ripke *et al.*, 'Biological insights from 108 schizophrenia-associated genetic

loci', *Nature*, vol. 511, no. 7510, pp. 421–427, Jul. 2014, doi:

10.1038/nature13595.

[25]   M. Lam *et al.*, 'Comparative genetic architectures of schizophrenia in East Asian

and European populations', *Nat Genet*, vol. 51, no. 12, pp. 1670–1678, Dec.

2019, doi: 10.1038/s41588-019-0512-x.

[26]   V. Trubetskoy *et al.*, 'Mapping genomic loci implicates genes and synaptic

biology in schizophrenia', *Nature*, vol. 604, no. 7906, pp. 502–508, Apr. 2022,

doi: 10.1038/s41586-022-04434-5.

[27]   M. Pirinen, P. Donnelly, and C. C. A. Spencer, 'Including known covariates can

reduce power to detect genetic effects in case-control studies', *Nature Genetics*

2012 44:8, vol. 44, no. 8, pp. 848–851, Jul. 2012, doi: 10.1038/ng.2346.

[28]   L. J. O'Connor, A. P. Schoech, F. Hormozdiari, S. Gazal, N. Patterson, and A.

L. Price, 'Extreme Polygenicity of Complex Traits Is Explained by Negative

Selection', *Am J Hum Genet*, vol. 105, no. 3, p. 456, Sep. 2019, doi:

10.1016/J.AJHG.2019.07.003.

[29]   N. S. Raghavan et al., 'Whole-exome sequencing in 20,197 persons for rare

variants in Alzheimer's disease', Ann Clin Transl Neurol, vol. 5, no. 7, p. 832,

Jul. 2018, doi: 10.1002/ACN3.582.

[30]   K. Yu, F. Liang, J. Ciampa, and N. Chatterjee, 'Efficient p-value evaluation for

resampling-based tests', Biostatistics, vol. 12, no. 3, p. 582, Jul. 2011, doi:

10.1093/BIOSTATISTICS/KXQ078.

[31]   G. Kimmel and R. Shamir, 'A fast method for computing high-significance

disease association in large population-based studies', Am J Hum Genet, vol. 79,

no. 3, pp. 481–492, 2006, doi: 10.1086/507317.

[32]   P. M. Visscher et al., 'Statistical Power to Detect Genetic (Co)Variance of

Complex Traits Using SNP Data in Unrelated Samples', PLoS Genet, vol. 10,

no. 4, p. e1004269, 2014, doi: 10.1371/JOURNAL.PGEN.1004269.

[33]   'Our Future Health'. https://ourfuturehealth.org.uk/ (accessed Jul. 05, 2023).

[34]   M. I. Kurki et al., 'FinnGen provides genetic insights from a well-phenotyped

isolated population', Nature 2023 613:7944, vol. 613, no. 7944, pp. 508–518,

Jan. 2023, doi: 10.1038/s41586-022-05473-8.

[35]   J. K. Pritchard and P. Donnelly, 'Case-control studies of association in

structured or admixed populations'. 2001.

[36]   R. Puhr, G. Heinze, M. Nold, L. Lusa, and A. Geroldinger, 'Firth's logistic

regression with rare events: accurate effect estimates and predictions?', Stat

Med, vol. 36, no. 14, pp. 2302–2317, Jun. 2017, doi: 10.1002/SIM.7273.

[37]   G. Heinze, M. Ploner, D. Dunkler, H. Southworth, and L. Jiricka, 'Package

"logistf": Firth's Bias-Reduced Logistic Regression'. CRAN, Jan. 18, 2022.

Acknowledgements

834 The authors would like to thank Prof Peter Visscher for his insightful comments and suggestions

835 on the methodology. The authors are additionally grateful to Manju Dissanayake, Chris

836 Marshall, Michael Mulholland, Andrew Donald and the Optima Partners Data Science and

837 Engineering software development team.

Author contributions

CNF developed the mathematical and statistical methodologies, developed the statistical

software and conceptualized the design. ZK developed the statistical software and webtool,

designed the methodological analysis pipeline and conducted the real-life analyses. REM

contributed to the interpretation of results. HR conceptualized and supervised the study. BBS

conceptualized, designed the study, contributed to the method, application, and

contextualization of the study. All authors contributed to the writing of the manuscript.

Competing interests

846   BBS and HR are employed by Biogen. CNF and ZK are employed by Optima Partners. REM

847   is an advisor to the Epigenetic Clock Development Foundation and Optima Partners.

Tables

Table 1. Accuracy, true positive rate, false positive rate and false discovery rate of isGWAS using REGENIE results as gold-standard and a threshold of $p = 5e - 08$ as classification rule. Results are obtained on all 11,079,229 variants used for the analysis of seven diseases in UK Biobank without clumping/finemapping.

Disease (ICD code)	Case:control ratio	*TPR*	*FPR*	*Acc*	*FDR*
Hypertension (I10)	1:2	0.625	0.000026	0.99961	0.041
Asthma (J45)	1:6	0.982	0.000028	0.99994	0.016
Atherosclerosis (I25)	1:9	0.886	0.000011	0.99995	0.043
Glaucoma (H40)	1:26	0.891	0	0.99997	0.036
Stroke (I63)	1:56	NA	0	1	NA
Colon Cancer (C18)	1:94	0.944	0	0.99999	0.037
Thyroid Gland Cancer (C73)	1:669	1	0	0.99999	0.051

Figures

Figure 1. Diagram highlighting main differences between isGWAS and other GWAS approaches.

Figure 2. Comparative results for Asthma (IC10:J45) from UK Biobank. Subplot (a) is a mirror Manhattan plot comparing $-\log_{10} P$ values for isGWAS and REGENIE-Firth and subplot (b) is a locus zoom of the gene FLG2 region +/-250kbp on chromosome 7. Subplot (c) plots $-\log_{10} P$ values for isGWAS and REGNIE-Firth with the standard threshold P-value indicated colored by population-level MAF. Subplots (d) and (e) showcase $\beta - \beta$ effect size estimates for variants with p-values<0.05 and are coloured by population-level MAF and $\log_2(\frac{SE(isGWAS)}{SE(REGENIE)})$.

Figure 3. a) Accuracy/TPR/FPR comparing REGENIE-Firth and isGWAS results, where $p = 5e - 08$ threshold has been used as indicator for correct classification accuracy. Results are obtained on all 11,079,229 variants used for the analysis without clumping/finemapping. See Supplementary Table 1 for full results. Manhattan plots b), c), d), e) and f): Comparative results for five diseases from UK Biobank. Mirror Manhattan plots comparing $-\log_{10} P$ values for isGWAS and REGENIE-Firth for six different diseases obtained from UK Biobank. Stroke was excluded from the analysis due to no variants passing significance threshold.

Figure 4. Simulation I results. Clockwise from top: a) Mean square error; b) distribution of estimated beta values; c) distribution of associated standard errors and d) distribution of -log₁₀(p-value), for each model - logistic regression, firth regression, 'isGWAS', 'isGWAS_Firth' and, for p-values only, Fisher's Exact Test - and specification of disease prevalence. Panel e) presents the relationship between Firth regression derived -log₁₀(p-value), along the horizontal axis, and the corresponding 'isGWAS_Firth' and Fisher's Exact Test (FET) computed values, on the vertical axis. Results are presented in the range [0,15] as FET regularly failed to converge for very small p-values and coloured according to value of $\pi\delta_{MAF}$, where

π denotes prevalence and $\delta_{MAF} = \dfrac{(MAF^* - MAF)}{\left(MAF(1 - MAF)\right)}$. *In panels b)-d) a point denotes the median value and error-bars the first and ninth deciles of the range.*

Figure 5. Performance of Leapfrog re-sampler (LRS) benchmarked against results derived from a)-e) the sample of ~276,204 individuals from UK Biobank with five different diseases and b) a simulated sample of 552,408 individuals (i.e., double UK Biobank sample size). For each value of ξ, we subset the target sample size down to N_{target}/ξ individuals and deploy the LRS to compute predictions for the target sample N_{target}. As the maximum number of UK Biobank samples was 276,204, this was taken as the target. For example, when ξ = 2, we subset the full sample to 138,102 individuals and run the LRS to compute predictions of the larger 276,204 sample. We use results from the disease analysis, benchmarking LRS predictions against those computed on the pruned genome sampling uniformly across significance associations, resulting in ~3500 variants per studied disease. Colon cancer and Stroke are excluded from this figure as they don't have significant variants or a very low number of such after pruning. In the right panel we generated 1000 simulated datasets under the null of no genetic association or the alternative (see simulation protocol for details).

Figure 6. Prediction results for Schizophrenia (2022) using data from Schizophrenia (2014): population-level information for 608 significantly associated loci (P<1e-07) obtained from clumping with parameters ($R^2 = 0.2, p_1 = 1e - 7, p_2 = 1e - 7$) has been used to infer p-values. a) The figure compares reported GWAS Schizophrenia (2014) p-values and isGWAS predicted p-values using population-level information from Schizophrenia (2014) matching for the larger 2022 cohort size. b) The figure compares reported GWAS Schizophrenia (2022) p-values and isGWAS predicted p-values using population-level information from Schizophrenia (2014) matching for the larger 2022 cohort size. The dashed red line represents the threshold P=5e-08.

Figure 7. a) Computational CPU time (in seconds) for an increasing number of variants. The results compare performance of isGWAS with and without Firth distributed over different number of CPU cores. The data was obtained from UK Biobank ICD10:C73 disease with low disease prevalence (case-control ratio = 1:669). The x- and y-axis are on $\log_{10}$ scale. b) Computational CPU time (in seconds) for seven UK Biobank diseases on 11,079,229 variants for ~335,000 individuals. We compare the performance of isGWAS running on 16 and 30 CPUs vs the performance of REGENIE Step 2 running on 16 CPUs (with 16 threads) and 30 CPUs (with 30 threads). The computation is performed on the same machine. The y-axis is on $\log_{10}$ scale.