Optogenetic Stimulation of Single Ganglion Cells in the Living Primate Fovea

Peter J. Murphy[1,2*]**, Juliette E. McGregor**[2,3]**, Zhengyang Xu**[1,2]**, Qiang Yang**[2,3]**,
William Merigan**[1,3]**, David R. Williams**[1,2]

*For correspondence:
pmurp16@ur.rochester.edu (PM)

[1]The Institute of Optics, University of Rochester, Rochester, New York; [2]Center for
Visual Science, University of Rochester, Rochester, New York; [3]Flaum Eye Institute,
University of Rochester, Rochester, New York

Abstract   Though the responses of the rich variety of retinal ganglion cells (RGCs) reflect the
totality of visual processing in the retina and provide the sole conduit for those processed
responses to the brain, we have much to learn about how the brain uses these signals to guide
behavior. An impediment to developing a comprehensive understanding of the role of retinal
circuits in behavior is the paucity of causal studies in the intact primate visual system. Here we
demonstrate the ability to optogenetically activate individual RGCs with flashes of light focused
on single RGC somas *in vivo*, with no evidence for activation of neighboring cells. The ability to
selectively activate specific cells is the first step toward causal experiments that directly link
retinal circuits to visual experience and behavior.

Introduction

Though only a few classes of light-sensitive cells capture the retinal image in the primate eye, evo-
lution has produced many more classes of retinal ganglion cell (RGC), on the order of 20, each
morphologically and physiologically distinct and independently tiling the retina, to convey visual
information to the brain. This dramatic expansion of cell classes in the retinal output combined
with microelectrode recordings in excised tissue have made it increasingly clear that the diversity
of RGCs reflects instead a surprising amount of complex early processing in the mammalian retina,
processing that previously been had been thought to reside deeper in the brain (*Gollisch and Meis-
ter, 2010*). There is evidence, for example, for specialized circuits that convey directional selectivity
(DS) (*Barlow et al., 1964*), segregate object from background motion (*Lettvin et al., 1959*), and antic-
ipate moving stimuli (*Berry et al., 1999*). However, we remain uncertain about the direct functional
role of these RGC classes because, while ex vivo recordings can characterize the specific stimulus
requirements of RGC classes, they are often made in the excised retina that is detached from the
downstream visual pathways they serve. Such studies preclude direct manipulations of RGC ac-
tivity to observe the impact on visual experience or behavior in the awake primate. For example,
we cannot say with certainty whether the DS cells recently observed in the non-human primate
(NHP) are involved in eye movements, the perceptual experience of motion, or both. Indeed, there
is even lingering controversy about the perceptual roles of the most common primate RGCs, the
midget and parasol ganglion cells (*Freedland and Rieke, 2022*; *Patterson et al., 2019*).

    To address these controversies, one exciting possibility for the future is the development of
studies in which the effect of manipulating RGC responses in the awake behaving animal can be

measured with psychophysical or behavioral measures. There have been formidable challenges to deploying this causal paradigm to study retinal circuits because of their inaccessibility inside the moving eye of the awake behaving primate. Advances in high resolution retinal imaging with adaptive optics now make it possible to observe single RGCs in the living eye (*Gray et al., 2008*). AO has also enabled very accurate eye tracking, making it possible to not only image but also stimulate single photoreceptors repeatedly with focused flashes of light (*Schmidt et al., 2018a*,b; *Tuten et al., 2017*) in the moving human eye. In addition, viral vectors now allow expression of calcium indicators such as GCaMP6s (*Yin et al., 2013a*; *Godat et al., 2022*), enabling recording from single RGCs *in vivo*. Finally, optogenetic activation of RGCs has been demonstrated in both macaque (*McGregor et al., 2020*) and human (*Sahel et al., 2021*) retina *in vivo*. These recent technical advances taken together now offer the possibility of causal experiments in which single RGCs are optogenetically stimulated in the living primate eye while the behavioral consequences are measured. Here, we demonstrate the first step toward establishing this capability: showing successful stimulation of single RGCs in the living primate eye.

Results and Discussion

RGC activity was modulated via the optogenetic ChrimsonR and monitored via fluorescence imaging of the calcium indicator GCaMP6s. Isolated foveal RGCs were chosen for individual stimulation, being targeted with short pulses of light intended to elicit activity in that cell only. The response of both the selected cell and all nearby cells were monitored to determine activation status. Chosen cells and their fluorescence traces can be seen in Fig. 1. Targeted cells showed large responses to the optogenetic probe. The GCaMP responses rise to peak quickly, and then decay over several seconds. This time course is consistent with GCaMP responses to cone excitation of RGCs measured via in vivo AOSLO imaging (*Yin et al., 2013b*). Fig. 2 shows all cells' ΔF/F values plotted against their distance from the targeted cell. Non-targeted cells showed no significant response above the background noise in contrast to the targeted cells, the responses of which were between 3 and 4 times the background noise floor. While we cannot exclude the possibility that surrounding cells responded to the flash below the noise threshold of our measurements, clearly these conditions strongly favor the excitation of the target cell over that of the neighbors we were able to record from. We cannot rule out the possibility that, due to the random nature of the vector expression process, some of these non-targeted cells were inactive because they failed to express ChrimsonR. The degree of GCaMP6s and ChrimsonR expression varies from one RGC to another, introducing the possibility that some cells may only have significant expression of one or neither agent. This means that some optogenetically active cells may have had no or weak GCaMP6s expression, diminishing our ability to observe their responses in our recordings. Despite this uncertainty, we can be reasonably sure that most cells are strongly co-expressing both agents. This is achieved via fluorescence imaging of the tdTomato marking ChrimsonR expression, which shows substantial overlap with the observed GCaMP6s fluorescence.

Though this method has limitations, these may be overcome via further refinements of the technique. For this initial demonstration, we deliberately selected the most favorable conditions, especially by selecting cells to target that were spatially isolated from neighboring cells. Here, we show results for cells at the inner edge of the ring of RGCs serving the primate fovea. The ability to extrapolate the method to cells closely packed on all sides by neighboring cells, as occurs at slightly larger eccentricities, between 0.5 and 2.5 degrees, deeper in the foveal ring of labelled cells, has yet to be explored. In this region RGCs have little to no separation between cell somas. The transverse spatial resolution of an AOSLO is roughly 1.6 μm, which is more than adequate for targeting the much larger soma diameter. However, the axial resolution of an AOSLO is roughly 25 μm, roughly twice the soma diameter. This means that there would be some optogenetic excitation of the cells immediately above and below the cell of interest. But even in this less favorable case the balance of excitation could be shifted strongly to a subset of RGCs, even if perfect isolation were not possible. The ability to isolate single cells becomes far more favorable again in the periphery, where the RGC

Figure 1. Cells chosen for stimulation. The cell circled in red marks the targeted cell. The averaged fluorescence traces for each cell show the integrated intensity over the cell's soma. The red marks on the time axis represent the onset of the 800 ms optogenetic stimulus. The histograms show each cell's response in terms of z score. (a) Results from male macaque. (b) Results from female macaque. All targeted cells are at similar eccentricities.

Figure 2. ΔF/F for each cell soma plotted against that cell's distance from the targeted cell. Note that even the nearby (< 50 μm) cell somas do not show a significantly elevated response (p » 0.05, unpaired t-test) than other cells at more distant locations. The leftmost point on each plot, colored in red, corresponds to the targeted cell itself. The traces for the four closest neighbors are shown below each scatter plot.

somas are in a monolayer. However, in this retinal region the RGC somas and their cone receptive fields lie on top of each other, precluding the use of the spatial offset between cone receptive fields and their somas to separate cone and RGC optogenetic excitation. The use of two photon excitation coupled with the use of a more sensitive optogenetic, such as ChRmine may mitigate this concern by increasing the optogenetic excitation while reducing the efficacy of light falling on photoreceptors (*Marshel et al., 2019*). Other approaches to increasing the selective activation of targeted cells may include cortically-targeted retrograde injections or class-specific viral vectors.

The demonstration of *in vivo* selective activation of individual RGCs lays the foundation for future causal studies of RGC function. The causal experimental approach has been successful in understanding the roles of neuronal classes in cortex of NHPs for many years, such as the work of Newsome and colleagues who showed that electrical activation of DS cells in MT modifies the animal's choices in motion discrimination tasks (*Salzman et al., 1992*; *Cicmil and Krug, 2015*; *Parker and Newsome, 1998*). More recently, this paradigm has incorporated optogenetic activation to investigate the function role of localized neural circuitry in cortex (*Jazayeri et al., 2012*; *Soma et al., 2019*). Applied to RGCs, this approach would involve first, *in vivo* calcium imaging to classify each RGC, followed by psychophysical experiments combined with optogenetic stimulation designed to explore each cell's role in visual behavior. Such experiments would establish a direct link between activation of an individual RGC and visual behavior and could help determine whether a given cell class contributes to perceptual experience, subconscious visual behavior, or both. These experiments will face additional challenges such as deploying the technology demonstrated here in the awake behaving animal where eye motion will be larger than in the anesthetized animal. Nonetheless, the single cone tracking and stimulation experiments in the awake human inspire confidence that this will be possible (*Schmidt et al., 2018a*,b; *Tuten et al., 2017*). These experiments in which single cones are targeted with light have shown that visibility can be measured from photon absorption with flashes of light delivered through adaptive optics to single cones from each of the three foveal cone classes (*Williams et al., 1981*; *Schmidt et al., 2018b*; *Hofer et al., 2005*), encouraging the possibility that optogenetic stimulation of a single RGC might be sufficient to elicit a sensation in an awake behaving monkey a hypothesis that future experiments could test. Monkeys are remarkably good at psychophysical experiments, with performance rivalling that of

human observers (*Matsuno and Fujita, 2009*).

Interpretation of causal experiments through optogenetic stimulation of a single RGC is in-
evitably complicated by the fact that perceptual experience depends on cell populations and such
experiments would necessarily produce unnatural population activity. How the brain would inter-
pret such an unusual pattern of excitation is not entirely clear and it may be necessary to control
the stimulation of multiple nearby RGCs to develop a clear understanding of their function. Addi-
tionally, should it prove possible to realize this paradigm, it could provide valuable data to guide
optogenetics-based vision restoration therapies. This could eventually offer the opportunity to
simultaneously activate multiple classes of RGCs according to their individual response character-
istics, leading to a perceptual experience that more closely resembles that generated by the natural
environment in the sighted eye.

Methods

Animals and Animal Care

These experiments were performed in two anesthetized *macaca fascicularis*, one male (M1) and
one female (M2). Macaques were housed in an AAALAC accredited facility and were provided with
care by 4 full time veterinarians, 5 veterinary technicians, and an animal behaviorist from the De-
partment of Comparative Medicine. The animal care staff monitored each animal for signs of dis-
comfort at least twice daily. They were caged in pairs and had free access to lab chow and water.
Their diet was additionally supplemented with daily treats including dried fruits, fresh fruits, veg-
etables, and nuts. All animals have daily access to enrichment items such as puzzle feeders and
mirrors, and are provided with movies and/or music. More novel enrichment items such as treat
filled bags, snow, or forage boxes were provided weekly. The macaques had access to a large play
space with swings and perches on a rotating basis. This study was carried out in strict accordance
with the Association for Research in Vision and Ophthalmology (ARVO) Statement for the Use of
Animals and the recommendations in the Guide for the Care and Use of Laboratory Animals of
the National Institutes of Health. One animal had two lesions of the central fovea, which had re-
moved cone input to many of the labeled RGCs. The RGCs deafferented in this manner are easily
distinguishable from those with normal input, and were avoided for study in this experiment.

Injections for Expression of Optogenetic and Calcium Indicator.

Prior to injection, both animals received daily subcutaneous injections of cyclosporine to reduce
the possibility of immune reaction to the injection. Blood trough levels were monitored to titrate
the dose into the range of 150-200 ng/ml, and these levels were maintained throughout the testing
period. The ocular surface was disinfected before injection with 50% diluted Betadine. Coexpres-
sion of both an optogenetic actuator (ChrimsonR) and a calcium indicator (GCaMP6s) enabled both
stimulation of RGCs and recording of their responses. This was achieved with a single intravitreal
injection containing a mixture of two adeno-associated virus (AAV2) based vectors with a ubiq-
uitous CAG promoter, AAV2-CAG-tdTomato-ChrimsonR and AAV2-CAG-GCaMP6s. These vectors
were synthesized by the University of Pennsylvania Vector core. The injection was delivered via a
30 gauge needle with a tuberculin syringe into the middle of the vitreous roughly 2 mm behind
the limbus. The injection was made in the right eye of M1 and the left eye of M2, and had a total
volume of 75 µL. Post-injection, the eyes were monitored with conventional SLO imaging (Heidel-
berg Spectralis) for adverse events such as inflammation, which was not present in either animal
throughout data collection. As previously demonstrated (*Dalkara et al., 2013*), expression was
largely confined to an annulus spanning from 0.5° to 2°. The inner edge of this annular region cor-
responds to the most centrally located RGCs that receive input from the most central fovea, while
the outer edge corresponds to the most eccentric location where we presume the inner limiting
membrane remained thin enough for penetration of the viral vector. All RGCs that obtained ex-
pression are close enough to the foveal center to be displaced from their receptive fields, allowing

for direct optogenetic activation of cell somas without stimulation of the cones that drive them.

AO Imaging

Anesthesia and Animal Preparation

Anesthesia and animal preparation were performed by licensed veterinary technicians. Macaques
were fasted overnight prior to anesthesia in preparation for ventilation. The morning of an imaging
session, macaques were moved to the laboratory and anesthetized with isoflurane. They were
placed prone upon a stereotaxic cart, supported by cushions and covered with a warming blanket.
The animals were paralyzed for the duration of imaging via an injection of vecuronium bromide. For
the duration of paralysis, the animal's heart and respiratory rates, blood oxygenation levels, and
electrocardiogram were monitored by a veterinary technician. The animal's pupils were dilated
with drops of phenylephrine chloride (2.5%) and Tropicamide (1%), and the eye fitted with a rigid
gas permeable contact lens to maintain corneal hydration. The animal's head was stabilized with
ear bars and a chin rest, and the animal's whole body could be rotated via the stereotaxic cart to
position the animal for imaging. The macaque was monitored for several hours by a veterinary
technician after each session to ensure full and safe recovery from the anesthetic drugs. This
procedure was repeated no more than once per week on an individual macaque.

Imaging Procedure

High resolution imaging was performed with an adaptive optics scanning light ophthalmoscope
(AOSLO), a complete description of which can be found in (*Godat et al., 2022*). A Shack-Hartmann
wavefront sensor in conjunction with an 847 nm laser diode source (QPhotonics) was used for
wavefront sensing. The wavefront sensor data controlled a deformable mirror (ALPAO) in a closed-
loop configuration operating at 10 Hz to provide correction of ocular aberrations. Reflectance and
fluorescence signals were collected with photomultiplier tubes. The RGC layer was imaged using
the fluorescent light emitted by the calcium indicator excited with light from a 488 nm laser (Qiop-
tiq), and the optogenetic was excited using the 640 nm line of a multi-line laser (Toptica). The cone
mosaic was simultaneously imaged with light from a a 796 nm superluminescent diode (Super-
lum) to provide a higher SNR helpful for both real-time stimulus stabilization and post-recording
image registration. Both uses are based upon cross-correlation of small strips of the image (*Yang*
et al., 2014). For stimulus stabilization, only strips located near the stimulus' intended location
were used in the cross correlation, allowing the stimulus to be repeatedly presented to the same
retinal location. This stabilization is necessary because the heart beat and respiration of the ani-
mal cause lateral shifts in the retinal imaging throughout recording of approximately 50 μm. The
image registration application is similar in execution and is applied in turn to all sections of the
image to obtain a single stabilized video. The motion data from the strip registration of the cone
reflectance video was used to stabilize the calcium recording. These methods allowed us to obtain
cellular-scale recordings from the primate retina in vivo while correcting for the motion inherent
to the living eye.

Cell selection

For this first study, we selected RGCs lying on the innermost edge of the annulus of expression
often had large distances between their somas, making them easier to target with optogenetic
stimulation without exciting neighboring cells. This crosstalk could potentially enter our measure-
ments both when stimulating and recording. First, scatter from the 640 nm stimulus may illumi-
nate nearby cell somas or underlying fibers of passage. Additionally, these nearby cell somas, their
dendritic arbors, and fibers may contaminate the GCaMP recording if they overlap with the cell of
interest's soma. We chose cells two cells in each of the two animals that were > 20 μm from their
the nearest neighbor to mitigate the possibility of optical crosstalk. We have considered two pos-
sibilities by which light sources in the experiment other than the optogenetic probe might have
interfered with the RGC response, but calculations confirm that neither would be expected to ap-

Figure 3. Relevant absorption/emission spectra for our fluorophores and optogenetic actuator. The emission wavelengths of tdTomato and GCaMP6 are well suited to activating ChrimsonR, and the wavelength used to image GCaMP fluorescence stimulates both fluorophores well.

preciably influence the probe response. First, the 488 nm imaging light is expected to produce an optogenetic activation that is only about 2% of that of the optogenetic probe. Moreover, it illumi­nates all cells equally throughout the experiment and would be expected to produce a constant activation, unlike the probe which is confined in space and time. Second, the emission from GCaMP and tdTomato is at least 4 orders of magnitude lower than this 488 nm stimulus, making it unlikely that these flurophores could have caused a measurable cell response.

Cells within a 1°x1° field of view were imaged at 488 nm (25 µW) for 30 seconds to allow for adaptation to the maintained calcium imaging light. The RGCs showed some level of baseline ac­tivation in response to this. ChrimsonR was then activated by a circular stimulus (640 nm, 750 µW) with a diameter of 12.5 µm, approximately the size of a cell soma. This stimulus was formed by modulating the 640 nm laser source's output to only be on while scanning over the soma of interest at a rate of 25 Hz. The duration of excitation was 800 ms, throughout which the stimulus was stabilized on the cell's soma using the method describe in the imaging procedure section. This stimulus was repeated four times for each trial with a waiting period of 15 seconds between stimuli. We have seen no evidence that this modulated cell activity via the normal cone pathways.

Analysis

Analysis The GCaMP6s recordings were registered using strip registration and the frames averaged to obtain a high SNR image of the RGC mosaic. This registration is necessary because the heart beat and respiration of the animal cause lateral shifts in the retinal imaging throughout recording of approximately 50 µm. The resulting RGC image was manually segmented to create a binary mask that can be applied to each frame of the video. The mask is used to extract the mean brightness of each cell soma for each frame. These brightness values are then used to calculate ΔF/F values for each cell soma in response to each stimulus. This calculation was performed for all cells that remain within the field of view for the entirety of the recording. Cells near the edge of the field of view often move in and out of visibility, obscuring any signals that may be present. The mean and standard deviation of ΔF/F values are calculated for periods during which no stimulus was present using data taken from all labeled RGCs in the subject's retina. These are used to transform the cell responses to stimuli into z-scores.

Acknowledgments

The authors would like to thank Amber Walker for providing animal care and anesthesia and Sara Patterson for providing feedback on the manuscript and data analysis. We thank the vector core at the Perelman School of Medicine, University of Pennsylvania and the Genetically-Encoded Neuronal Indicator and Effector (GENIE) Project and the Janelia Research Campus of the Howard Hughes Medical Institute, specifically Vivek Jayaraman, Ph.D., Douglas S. Kim, Ph.D., Loren L. Looger, Ph.D., and Karel Svoboda, Ph.D.

Additional Information

Funding

Funding	Grant Reference Number	Author
National Eye Institute	R01EY031467	David Williams
National Eye Institute	P30EY001319	David Williams
Air Force Office of Scientific Research	FA9550-22-1-0167	David Williams

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Competing Interests

David R. Williams: Patents with the University of Rochester for adaptive optics retinal imaging: US patent 6,199,986 "Rapid, automatic measurement of the eye's wave aberration", US patent 6,264,328 "Wavefront sensor with off-axis illumination" and US patent 6,338,559 "Apparatus and method for improving vision and retinal imaging". Qiang Yang: Patents with the University of Rochester, Canon Inc. and the University of Montana, for image stabilization algorithms: US patent 9,226,656: "Real-time optical and digital image stabilization for adaptive optics scanning ophthalmoscopy", US patent 9,406,133: "System and method for real-time image registration", US patent 9,485,383, "Imaging based correction of distortion from a scanner" and US patent 9,454,084, "Light source modulation for a scanning microscope".

Author Contributions

Peter J. Murphy, Conceptualization, Data curation, Formal analysis, Investigation, Software, Visualization, Methodology, Writing - original draft, Writing - reviewing and editing; Zhengyang Xu, Investigation, Methodology; Juliette E. McGregor, Investigation, Methodology, Writing - reviewing and editing; Qiang Yang, Software, Methodology; William Merigan, Concepualization, Investigation, Methodology, Project administration, Supervision, Resources, Writing - reviewing and editing; David R. Williams, Conceptualization, Project administration, Supervision, Resources, Funding acquisition, Writing - original draft, Writing - reviewing and editing

Author ORCIDs

Peter J. Murphy https://orcid.org/0000-0002-9885-7779

Juliette E. McGregor https://orcid.org/0000-0002-8046-028X

William Merigan https://orcid.org/0000-0002-9359-5538

David R. Williams https://orcid.org/0000-0003-3227-8333

Data Availability

All fluorescence recordings, along with example analysis scripts and required supporting files, are available at https://osf.io/zgjsv/

References

Barlow HB, Hill RM, Levick WR. Retinal ganglion cells responding selectively to direction and speed of image motion in the rabbit. The Journal of Physiology. 1964; 173(3):377–407. https://physoc.onlinelibrary.wiley.com/doi/abs/10.1113/jphysiol.1964.sp007463, doi: https://doi.org/10.1113/jphysiol.1964.sp007463.

Berry MJ, Brivanlou IH, Jordan TA, Meister M. Anticipation of moving stimuli by the retina. Nature. 1999 Mar; 398(6725):334–338. https://doi.org/10.1038/18678, doi: 10.1038/18678.

Cicmil N, Krug K. Playing the electric light orchestra—how electrical stimulation of visual cortex elucidates the neural basis of perception. Philosophical Transactions of the Royal Society B: Biological Sciences. 2015; 370(1677):20140206. https://royalsocietypublishing.org/doi/abs/10.1098/rstb.2014.0206, doi: 10.1098/rstb.2014.0206.

Dalkara D, Byrne LC, Klimczak RR, Visel M, Yin L, Merigan WH, Flannery JG, Schaffer DV. In Vivo–Directed Evolution of a New Adeno-Associated Virus for Therapeutic Outer Retinal Gene Delivery from the Vitreous. Science Translational Medicine. 2013; 5(189):189ra76–189ra76. https://www.science.org/doi/abs/10.1126/scitranslmed.3005708, doi: 10.1126/scitranslmed.3005708.

Freedland J, Rieke F. Systematic reduction of the dimensionality of natural scenes allows accurate predictions of retinal ganglion cell spike outputs. Proceedings of the National Academy of Sciences. 2022; 119(46):e2121744119. https://www.pnas.org/doi/abs/10.1073/pnas.2121744119, doi: 10.1073/pnas.2121744119.

Godat T, Cottaris NP, Patterson S, Kohout K, Parkins K, Yang Q, Strazzeri JM, McGregor JE, Brainard DH, Merigan WH, Williams DR. In vivo chromatic and spatial tuning of foveolar retinal ganglion cells in Macaca fascicularis. PLoS One. 2022 Nov; 17(11):e0278261.

Gollisch T, Meister M. Eye Smarter than Scientists Believed: Neural Computations in Circuits of the Retina. Neuron. 2010; 65(2):150–164. https://www.sciencedirect.com/science/article/pii/S0896627309009994, doi: https://doi.org/10.1016/j.neuron.2009.12.009.

Gray DC, Wolfe R, Gee BP, Scoles D, Geng Y, Masella BD, Dubra A, Luque S, Williams DR, Merigan WH. In Vivo Imaging of the Fine Structure of Rhodamine-Labeled Macaque Retinal Ganglion Cells. Investigative Ophthalmology I& Visual Science. 2008 01; 49(1):467–473. https://doi.org/10.1167/iovs.07-0605, doi: 10.1167/iovs.07-0605.

Hofer H, Singer B, Williams DR. Different sensations from cones with the same photopigment. Journal of Vision. 2005 05; 5(5):5–5. https://doi.org/10.1167/5.5.5, doi: 10.1167/5.5.5.

Jazayeri M, Lindbloom-Brown Z, Horwitz GD. Saccadic eye movements evoked by optogenetic activation of primate V1. Nature Neuroscience. 2012 Oct; 15(10):1368–1370. https://doi.org/10.1038/nn.3210, doi: 10.1038/nn.3210.

Lettvin JY, Maturana HR, McCulloch WS, Pitts WH. What the Frog's Eye Tells the Frog's Brain. Proceedings of the IRE. 1959; 47(11):1940–1951. doi: 10.1109/JRPROC.1959.287207.

Marshel JH, Kim YS, Machado TA, Quirin S, Benson B, Kadmon J, Raja C, Chibukhchyan A, Ramakrishnan C, Inoue M, Shane JC, McKnight DJ, Yoshizawa S, Kato HE, Ganguli S, Deisseroth K. Cortical layer–specific critical dynamics triggering perception. Science. 2019; 365(6453):eaaw5202. https://www.science.org/doi/abs/10.1126/science.aaw5202, doi: 10.1126/science.aaw5202.

Matsuno T, Fujita K. A comparative psychophysical approach to visual perception in primates. Primates. 2009 Apr; 50(2):121–130. https://doi.org/10.1007/s10329-008-0128-8, doi: 10.1007/s10329-008-0128-8.

McGregor JE, Godat T, Dhakal KR, Parkins K, Strazzeri JM, Bateman BA, Fischer WS, Williams DR, Merigan WH. Optogenetic restoration of retinal ganglion cell activity in the living primate. Nature Communications. 2020 Apr; 11(1):1703. https://doi.org/10.1038/s41467-020-15317-6, doi: 10.1038/s41467-020-15317-6.

Parker AJ, Newsome WT. SENSE AND THE SINGLE NEURON: Probing the Physiology of Perception. Annual Review of Neuroscience. 1998; 21(1):227–277. https://doi.org/10.1146/annurev.neuro.21.1.227, doi: 10.1146/annurev.neuro.21.1.227, pMID: 9530497.

Patterson SS, Neitz M, Neitz J. Reconciling Color Vision Models With Midget Ganglion Cell Receptive Fields. Frontiers in Neuroscience. 2019; 13. https://www.frontiersin.org/articles/10.3389/fnins.2019.00865, doi: 10.3389/fnins.2019.00865.

331 **Sahel JA**, Boulanger-Scemama E, Pagot C, Arleo A, Galluppi F, Martel JN, Esposti SD, Delaux A, de Saint Aubert
332 JB, de Montleau C, Gutman E, Audo I, Duebel J, Picaud S, Dalkara D, Blouin L, Taiel M, Roska B. Partial recovery
333 of visual function in a blind patient after optogenetic therapy. Nature Medicine. 2021 Jul; 27(7):1223–1229.
334 https://doi.org/10.1038/s41591-021-01351-4, doi: 10.1038/s41591-021-01351-4.

335 **Salzman C**, Murasugi C, Britten K, Newsome W. Microstimulation in visual area MT: effects on direction discrim-
336 ination performance. Journal of Neuroscience. 1992; 12(6):2331–2355. https://www.jneurosci.org/content/
337 12/6/2331, doi: 10.1523/JNEUROSCI.12-06-02331.1992.

338 **Schmidt BP**, Boehm AE, Foote KG, Roorda A. The spectral identity of foveal cones is preserved in hue perception.
339 Journal of Vision. 2018 10; 18(11):19–19. https://doi.org/10.1167/18.11.19, doi: 10.1167/18.11.19.

340 **Schmidt BP**, Sabesan R, Tuten WS, Neitz J, Roorda A. Sensations from a single M-cone depend on the activity
341 of surrounding S-cones. Sci Rep. 2018 Jun; 8(1):8561.

342 **Soma S**, Yoshida J, Kato S, Takahashi Y, Nonomura S, Sugimura YK, Ríos A, Kawabata M, Kobayashi K, Kato
343 F, Sakai Y, Isomura Y. Ipsilateral-Dominant Control of Limb Movements in Rodent Posterior Parietal
344 Cortex. Journal of Neuroscience. 2019; 39(3):485–502. https://www.jneurosci.org/content/39/3/485, doi:
345 10.1523/JNEUROSCI.1584-18.2018.

346 **Tuten WS**, Harmening WM, Sabesan R, Roorda A, Sincich LC. Spatiochromatic Interactions between Individual
347 Cone Photoreceptors in the Human Retina. J Neurosci. 2017 Sep; 37(39):9498–9509.

348 **Williams DR**, MacLeod DI, Hayhoe MM. Punctate sensitivity of the blue-sensitive mechanism. Vision Res. 1981;
349 21(9):1357–1375.

350 **Yang Q**, Zhang J, Nozato K, Saito K, Williams DR, Roorda A, Rossi EA. Closed-loop optical stabilization and
351 digital image registration in adaptive optics scanning light ophthalmoscopy. Biomed Opt Express. 2014 Sep;
352 5(9):3174–3191. https://opg.optica.org/boe/abstract.cfm?URI=boe-5-9-3174, doi: 10.1364/BOE.5.003174.

353 **Yin L**, Geng Y, Osakada F, Sharma R, Cetin AH, Callaway EM, Williams DR, Merigan WH. Imaging light responses
354 of retinal ganglion cells in the living mouse eye. J Neurophysiol. 2013 Feb; 109(9):2415–2421.

355 **Yin L**, Geng Y, Osakada F, Sharma R, Cetin AH, Callaway EM, Williams DR, Merigan WH. Imaging light responses
356 of retinal ganglion cells in the living mouse eye. J Neurophysiol. 2013 Feb; 109(9):2415–2421.