**Title:** Developmental trajectories of EEG aperiodic and periodic components: Implications for understanding thalamocortical development during infancy.

**Authors:** Carol L. Wilkinson[1,2], Lisa Yankowitz[1], Jerry Y. Chao[3], Rodrigo Gutiérrez[4,9], Jeff L. Rhoades[5,6], Shlomo Shinnar[7,8], Patrick L. Purdon[9], Charles A. Nelson[1,2,10]

**Affiliations**

[1] Division of Developmental Medicine, Boston Children's Hospital, Boston, MA, United States.

[2] Harvard Medical School, Boston, MA, USA

[3] Department of Anesthesiology, Montefiore Medical Center, Children's Hospital at Montefiore, Albert Einstein College of Medicine, Bronx, NY, USA.

[4] Departamento de Anestesia y Medicina Perioperatoria, Hospital Clínico de la Universidad de Chile, Santiago, Chile.

[5] Department of Neurobiology, Harvard Medical School, Boston, MA, USA.

[6] Program in Neuroscience, Division of Medical Sciences, Graduate School of Arts and Sciences, Harvard University, Cambridge, MA, USA.

[7] The Saul R. Korey Department of Neurology, Montefiore Medical Center, Albert Einstein College of Medicine, Bronx, NY, USA.

[8] Department of Epidemiology and Population Health, Albert Einstein College of Medicine, Bronx, NY, USA.

[9] Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital, Boston, MA, United States.

[10] Harvard Graduate School of Education, Cambridge, MA, United States.

**Abstract**

The development of neural circuits has long-lasting effects on brain function, yet our understanding of early circuit development in humans remains limited. Here, periodic EEG power features and aperiodic components were examined from longitudinal EEGs collected from 592 healthy 2–44 month-old infants, revealing age-dependent nonlinear changes suggestive of distinct milestones in early brain maturation. Consistent with the transient developmental progression of thalamocortical circuitry, we observe the presence and then absence of periodic alpha and high beta peaks across the three-year period, as well as the emergence of a low beta peak (12-20Hz) after six months of age. We present preliminary evidence that the emergence of the low beta peak is associated with higher thalamocortical-dependent, anesthesia-induced alpha coherence. Together, these findings suggest that early age-dependent changes in alpha and beta periodic peaks may reflect the state of thalamocortical network development.

## Introduction

The infant brain undergoes dramatic structural and physiological change in the first year after birth. Rapid

increases in brain volume coincide with expansive synaptogenesis[1-3], as well as interneuron migration,

maturation and network integration[4]. In particular, during this early period thalamocortical connections

are established through an intricate sequence that plays a critical role in the development of sensory

cortical networks[5]. However, the detailed timing of interneuron and thalamocortical maturation in human

development is largely unknown. In rodent models, the development of thalamocortical circuitry is

notable for transient inhibitory connections that drive subsequent circuit formation and coincide with

critical periods of plasticity present during the first 2-3 postnatal weeks[6]. In humans, longitudinal resting-

state fMRI data suggest that while thalamic-sensorimotor connectivity networks are present at birth, other

networks (e.g., thalamus-medial-visual, thalamus-default-mode) do not emerge until 1 year of age[7].

However, MRI studies thus far have been limited to measuring annual changes in structural or functional

connectivity, preventing a detailed understanding of rapid developmental change during this period. In

contrast, electroencephalography (EEG) can provide frequent and non-invasive repeated measurement

of brain oscillations that directly result from transient developmental changes in inhibitory networks and

maturation of thalamocortical circuitry[5,8,9].

The EEG power spectrum is comprised of two physiologically distinct components reflecting underlying

neuronal activity: aperiodic and periodic activity. The aperiodic component defines the slope of the power

spectrum, following a 1/f power law distribution (Fig 1A) and reflects non-oscillatory neuronal spiking

activity[10-12]. In addition, recent evidence suggests that the aperiodic slope reflects the excitatory-

inhibitory (E/I) balance of the underlying neuronal network, where a flattened, reduced slope is

associated with increased excitation over inhibition, and a steeply more accelerated slope with increased

inhibition over excitation[11]. Longitudinal studies extending from childhood to adulthood have observed

decreases in aperiodic slope with age, suggestive of increases in E/I balance with age[13-16]. Changes in

the aperiodic component in early infancy are less well described, and we hypothesize they may be

substantially different from those in childhood, as the first year after birth includes rapid increases in

neuronal activity, synaptogenesis, and inhibitory neuron integration.

The periodic component of the power spectrum is defined as the portion of the absolute power spectrum

rising above the aperiodic slope (Fig 1B). Periodic power reflects oscillatory activity occurring in narrow

frequency bands that are highly correlated with various cognitive processes and behavioral states[17,18],

and provide the foundation for both local and long-range communication within the brain. The majority of

neural oscillations observed in the power spectrum are the direct result of inhibitory and thalamocortical

network responses to sensory input[19–23]. Thus, as a measure, the EEG power spectrum is well positioned

to shed light on the developmental timing of inhibitory and thalamocortical network maturation.

Thus far, developmental EEG studies have largely focused on theta/alpha oscillations which are

modulated by thalamocortical interactions and are associated with cognitive functions of attention and

memory[24–26]. Multiple studies of the first two years of life have observed a shift in peak frequency from a

5 Hz theta peak  at 5 months to 8 Hz alpha peak at 2 years, coinciding with increases in alpha power

across this period[27–31]. This dominant peak frequency continues to increase into the mature 10Hz

posterior alpha rhythm by adolescence[9,32]. It is hypothesized that the gradual shift in dominant peak

frequency is modulated by maturation of thalamocortical circuitry in concert with developmental gains in

cognitive functions[9]; however, the precise mechanisms remain unknown.

Unlike theta/alpha power, little is known about the early developmental changes in periodic beta power.

In adults, beta oscillations are strongly associated with sensorimotor processing in addition to higher-

order cognitive tasks such as working memory[33]. Similar to alpha oscillations, the generation of beta

oscillations relies on GABAergic interneuron networks and thalamocortical connectivity[34–36]. In adults,

low-dose GABA-modulating anesthetics induce a sedative state with 13-25Hz beta oscillations, whereas

higher doses used to maintain unconsciousness progressively slow these beta oscillations into coherent,

frontal specific, alpha oscillations[36–38]. However, GABA-dependent anesthesia-induced frontal alpha

coherence does not emerge in infants until *after* 10 months of age and is not consistently present until

15-20 months of age [39–41]. Anesthesia-induced alpha coherence is hypothesized to involve GABA-

dependent thalamocortical loops leading to hypersynchronization between thalamic and prefrontal

cortices [38,42,43]. Therefore, potential covariation of developing beta oscillations and anesthesia-induced

counterparts may lend insight into the role and time course of developing inhibition in human

thalamocortical circuit development.

Using longitudinal EEG data collected from 592 healthy infants (yielding a total of 1335 EEGs) from 2 to

44 months after birth, in this work we characterize early developmental trajectories of EEG aperiodic

components and periodic power from 2-50Hz. In doing so, this work identifies potential ages during

which sequential steps in inhibitory network and thalamocortical circuit development may occur.

Consistent with the transient and stepwise developmental progression of thalamocortical circuitry, we

observe transient periodic peaks in alpha power at 2-3 months and high beta power at 4-18 months. A

low beta peak (12-20Hz) also begins to emerge in infants starting as early as 6 months of age. We

hypothesized that emergence of this low beta peak reflects maturation of early connections between the

thalamus and cortex. To test this hypothesis, we leveraged a smaller dataset consisting of a cohort of

infants with EEG recordings before and during clinical anesthesia. Consistent with our hypothesis, we

find infants with an identifiable low beta peak have higher anesthesia-induced alpha coherence

compared to those that who do not, providing preliminary evidence that the emergence of this peak may

be associated with thalamocortical loop maturation.

**RESULTS**

Resting-state EEG were collected longitudinally from 592 healthy infants, aged 2-44 months, across 4

studies occurring in the same laboratory (Fig 1C, Table 1). Whole brain power spectra for each individual

were calculated using a multitaper spectral analysis[44] for each electrode, and then averaging across

electrodes (Supplemental Figure 1. Individual spectra shown in Supplemental Figure 2). For visualization

of developmental changes, spectra were then averaged across individuals within 8 age bins (Fig 1D).

Notable nonlinear changes in aperiodic and periodic power spectra were observed between age bins,

including transient peaks in the periodic spectrum across both alpha and beta frequency ranges (Fig 1E-

G). To further characterize these developmental changes in the spectra, we used generalized additive

mixed models (GAMMs) to model non-linear trajectories of power parameters. For each model an age-

by-sex interaction was tested for significance. If not significant, the interaction term was removed, and

the model was refit using sex as an additive covariate. All models also included random effects of study

and participant. Spaghetti plots of individual non-GAMMs modeled data are shown in Supplemental

Materials.

*Aperiodic activity increases most during first year of life*

First, we assessed age-dependent changes in the aperiodic component and observed the largest

developmental increases in aperiodic activity between 2 and 8 months after birth (Fig 1E). Aperiodic

offset and slope significantly increased with age (FDR-adjusted $q$ values <0.001), and age-by-sex

interactions were present for both aperiodic offset (F = 5.28, q = 0.002) and slope (F = 3.04, q = 0.03).

Modeled developmental trajectories of the aperiodic offset showed a qualitatively sharp linear increase

over the first year after birth for both males and females (Fig 1H). Modeled developmental trajectories of

the aperiodic slope showed a qualitatively gradual increase over the first year. These findings contrast

with consistent reports of *decreasing* offset and slope across child and adulthood[13–16], and likely reflect

the known increases in brain volume and synaptogenesis occurring across the first year of life.

Differences in developmental trajectories between 4 regions of interest (ROI) (frontal, central, temporal,

and posterior) were also assessed (Supplemental Figure 5 and 6). The posterior ROI had higher

aperiodic offset than all other ROIs, with the greatest increase in offset occurring in the first year (Frontal

F = -32.47, q <0.0001, Central F = -50.12, q < 0.0001, Temporal -39.29 , q < 0.0001; Supplemental Fig

3B).

**Figure 1**

**Figure 1. Developmental Trajectories of Aperiodic and Periodic Power Spectra**
**(A, B)** Example power spectra derived from 6-8 month-old participants from Study 2. Mean spectra with shaded errors describing 95% confidence intervals. **(A)** Aperiodic offset is defined as power at 2.5Hz. **(B)** Periodic power peaks defined as maxima within a defined frequency range. Periodic band power defined as the integral of the periodic power spectra between defined frequencies. **(C)** Longitudinal study enrollment. Each line is a participant with dots indicating when EEG was collected for that participant. **(D-F)** Absolute, Aperiodic, and Periodic power visualized across 8 age bins (see Table 1). Spectra from EEGs collected at within each age bin were averaged and shading describes 95% confidence intervals. **(G)** Heatmap showing age-related changes in periodic power binned every 2 months. **(H-I)** GAMMs modeled trajectories of aperiodic offset and slope for males (orange) and females (blue). Here, age is incorporated into the model using exact age in days, rather than age-bins. Relative inflection points are shown with circular markers. Below, heatmaps show the standardized change in offset or slope per day, defined as [change per day]/[standard deviation of measures across full age range]

*Transient 9.5Hz alpha peak observed in 2-4 month-old infants*

At the youngest age bin (2–4 months) *two* peaks with similar amplitude are observed across the

theta/alpha (4-12Hz) range in the majority of infants (69%; Fig 2A,C). A lower frequency peak is

observed in the theta (4-6Hz) range at 5.5 ± 0.3Hz, and higher frequency peak is observed in the alpha

(6-12Hz) range at 9.5 ± 0.45Hz. However, by 6-months only 15% of infants have two peaks in this range,

and for most infants it is the higher 9.5Hz peak that is no longer observed. At 6-months fewer than 40%

of infants exhibit a dominant peak in the "alpha" (6.5-12Hz) range (Fig 1D) and the average peak

frequency in the theta/alpha range is 6.3±1Hz. Age (in days) was negatively associated with the

probability of having two peaks across the 2-6 month age range in a generalized linear mixed effects

model (Odds Ratio = 0.98, p < 0.0001). This disappearance of the higher peak after 4-months of age

may reflect a transient step in thalamocortical circuit development. Previous research has observed a

gradual shift in peak frequency from 5 to 8Hz from infancy to early childhood, however these studies

started no earlier than 5 months of age[27–30]. In order to assess whether an increase in peak frequency

beginning at 5 months is present in our data set we modeled developmental trajectories of peak

amplitude and frequency between 4-12 Hz starting at 170 days, when the vast majority of EEGs

exhibited a single dominant peak. No age-by-sex interactions were observed in GAMMs modeled

trajectories, and consistent with previous studies peak frequency and peak amplitude significantly

increased with age (frequency: F = 138.3, q <0.0001 Fig 2E; amplitude: F = 416.62 q < 0.0001, Fig 2F).

Figure 2G-I show modeled trajectories for EEG power calculated over defined frequency bands

commonly used in infant EEG research: theta (4-6Hz), low alpha (6-9Hz), and high alpha (9-12Hz). An

age-by-sex interaction was present for theta power, although qualitatively the shapes of trajectories were

similar (Fig G; F = 5.5, q < 0.01).

# Figure 2

**Figure 2. Transient and nonlinear changes in periodic power between 4 and 12 Hz.**
**(A,B)** Individual periodic power spectra for 2-4 months, and 6-8 months old. Red markers show peaks between 4 and 12 Hz. **(C)** Proportion of infants with two peaks identified between 4 and 12 Hz at each age bin. **(D)** Proportion of infants with an identified peak between 6.5 and 12Hz at each age bin. **(E-I)** GAMMs modeled trajectories for males (orange) and females (blue). Relative inflection points are shown with circular markers. Below, heatmaps show the standardized change in offset or slope per day, defined as [change per day]/[standard deviation of measures across full age range]. Both male and female heatmaps shown for models with significant age x sex interaction.

*Transient beta peaks between 4-18 months*

Several age-dependent transient changes are observed in the low beta (13–20Hz) and high beta (20-

35Hz) range. First, the shape of the periodic power spectra in the low beta range is notable for the

absence of a low beta peak prior to 1 year of age (Fig 3A), with only 10% of infants (24/222) exhibiting a

peak between 6-8 months of age (Fig 3A-C). After 8 months, a low beta peak begins to emerge in some

of infants, with 48% (52/107) showing a peak at 18-20 months, and 70% (199/285) by 36 months (Fig

2C). As a low beta peak was not identified in many children across the age range, peak amplitude and

frequency was not modeled. In contrast, virtually all (99.5%) of the infants had an identifiable high beta

peak *prior* to 12 months of age (Fig 3A). However, notable nonlinear shifts in frequency and amplitude of

the high beta peak were observed (Fig 1F, 3D and E). During the first year after birth, the high beta peak

amplitude increases, peaking at 372 days (12.2 months), and then substantially decreases until 1021

days (2.8 years). High beta peak frequency trajectories are also nonlinear, with peak frequency at its

highest at 491 days (male 28.4Hz, female 28.8Hz), followed by a steady decline in frequency. Modeled

trajectories of periodic power for commonly used frequency bands are shown in Fig 3F-H: low beta (12-

20Hz), high beta (20-30Hz), and gamma (30-45Hz).

The observed nonlinear changes across the beta range are striking. While many EEG infant studies

group beta oscillations into a singular frequency range, the data presented here supports that low and

high beta have distinct developmental origins. Specifically, between 6-24 months we observe the gradual

emergence of a low beta peak, and simultaneously the rise and fall of a prominent high beta peak,

ultimately resolving into a broader beta peak by 36 months.

Traditionally, beta oscillations measured in children and adults are associated with sensory and motor

processing, where reductions in beta power are observed during the preparation or execution of motor

tasks[33]. However, beta activity has also been shown to be modulated during a wide range of nonmotor

cognitive tasks[33,45]. The developmental emergence of low beta oscillations may represent sensorimotor

skills (e.g., crawling, walking) gained during this period, but may also represent the developmental

maturation of neural circuitry. For example, GABAeric interneuron networks and thalamocortical

connectivity are highly associated with the generation of cortical beta oscillations, as well as anesthesia-

induced frontal alpha coherence, but neither are not fully established at birth[39].

## Figure 3

**Figure 3. Transient and nonlinear changes in periodic power between 12 and 35Hz.**
**(A,B)** Individual periodic power spectra 6-8 months, and 18-20 months old. Red markers show peaks between 12-20 Hz. Green markers show peaks between 20-35Hz. **(C)** Proportion of infants with peak identified between 12-20 Hz at each age bin. **(D-H)** GAMMS modeled trajectories for males (orange) and females (blue). Relative inflection points are shown with circular markers. Below, heatmaps show the standardized change in offset or slope per day, defined as [change per day]/[standard deviation of measures across full age range].

*Low beta peak associated with anesthesia-induced alpha coherence*

We wondered whether developmental changes in infant beta power measured in a resting state may represent concurrent maturation of GABAergic interneuron networks and thalamocortical connectivity. Multiple lines of evidence suggest that anesthesia-induced frontal alpha coherence is dependent on thalamocortical connectivity[38,42,43], and in infants robust levels of alpha coherence with anesthesia administration are not observed until 10 months of age[40]. We hypothesized that the emergence of low beta oscillations (as measured by the presence of a low beta peak) beginning after 7-months of age may reflect maturation of the thalamocortical loop also responsible for the developmental emergence of anesthesia-induced frontal alpha coherence around the same age[40].To explore this possibility, we analyzed data from infants participating in cross-sectional study where EEG data was collected before and during exposure to GABA-modulating sevoflurane anesthesia[46].  All infants were undergoing elective procedures (e.g., circumcision) and infants were excluded for prematurity, neurologic injury, epilepsy, or planned intracranial surgery. We then tested whether infants with a low beta peak during the awake, unanesthetized state had increased GABA-dependent anesthesia-induced frontal alpha coherence. EEG data from 36 infants across a broad age range (6-15 months old) were collected during the (1) awake pre-anesthetized and (2) anesthetized states. We first analyzed the baseline awake state (pre-anesthesia) EEG across all 36 infants to confirm that similar developmental changes were observed. Developmental changes in the periodic power spectra in this smaller dataset were qualitatively similar to those described above (Figure 4A), with a low beta peak beginning to emerge in some infants after 7 months and present in roughly half the infants between 7-12 months of age (11/23). To test our above hypothesis, we then limited our analysis to this 7-12 month age range (n=23), and compared alpha coherence *during anesthesia* in those with and without a low beta peak in the awake, unanesthetized state. As hypothesized, both ANCOVA and a generalized linear mixed effects (LME) model demonstrated a significant effect of low beta peak presence on median alpha coherence, with increased alpha coherence in those with a low beta peak compared to those without (ANOVA $F(1,20) = 5.25$, p =0.03; LME $\beta = 0.12$, SE = 0.051, p = 0.02; Figure 4B).

**Figure 4**

**Figure 4. Increased anesthesia-induced alpha coherence in infants with identified low beta peak in baseline EEG. (A)** Periodic power spectra of infants between 6 and 15 months old prior to receiving anesthesia **(B)** Mean alpha coherence during anesthesia in infants 7-12 months old, with (light blue) or without (white) an identified low beta peak. Both a generalized linear mixed effects model and Ancova showed significant association between low beta peak presence with anesthesia-induced alpha coherence (p < 0.05 for both tests). Dotted lines represent quartiles, with the long dash represented the median.

**Discussion**

We present, to our knowledge the largest-to-date longitudinal analysis of EEG data collected between 2

– 44 months of age. Findings provide insight into the developmental timing of inhibitory network and

thalamocortical circuit maturation during human infancy. Several age-dependent findings in our study

contrast with previous longitudinal studies of child and adulthood. First, we observe *increases* in both

aperiodic offset and slope, especially during the first year, whereas decreases in both measures are

observed starting as early as 4 years of age and continue to decrease with adulthood[13–16]. Second, while

expected shifts in the dominant peak from the theta to alpha range were observed between 5 to 44

months, in the 2-4 months age bin, a 9.5Hz peak was also transiently observed. Third, striking changes

within the beta (12-30Hz) range were observed, including the emergence of a low beta peak starting

after 6-months of age, and age-dependent shifts in high beta peak frequency and amplitude - first

increasing and then decreasing with age. Below we discuss how the above age-related changes may

represent sequential developmental maturation in the inhibitory system and thalamocortical network connections.

The aperiodic offset is hypothesized to represent broad band neuronal firing[10,47], and thus early increases in aperiodic offset are consistent with established increases in neuronal number, gray matter volume, and synaptic number during the first year[1–3,48,49]. Stabilization of aperiodic offset after 1-year of age is also consistent with MRI findings that gray matter volume doubles during the first postnatal year and then slows to 20% in its second year[48–50]. Regionally, we also observe differences between posterior and frontal aperiodic offset trajectories, which either plateau after 1 year (posterior), or have a slow continued increase (frontal) beyond 1 year of age (Supplemental Fig 3). Consistent with this pattern, synaptogenesis differs across cortical regions, with the posterior visual cortex exhibiting a burst in synapse formation between 3-4 months of age, whereas the prefrontal cortex shows peak synaptogenesis around 8 months of age and continued gains during the second year of life[1].

Our observed age-dependent *increases* in aperiodic slope in infancy also contrast with multiple studies covering child to adulthood, where decreases in slope have consistently been reported. Schaworonkov et. al.[51] also reported decreased slope with age in infants from 1 to 7-months-old, however the parameterization of the spectra in that study was limited to 1-10Hz due to excessive muscle noise in the data, and it is unclear how the shifts in 4-12Hz periodic power described below may affect modeling of the underlying aperiodic component in this range. We hypothesize that observed increases in aperiodic slope reflect changes in inhibitory networks that are unique to early development. Indeed, aperiodic slope from EEG recorded from sleeping newborns is observed to increase with age during the first 7 weeks after birth[52]. Growing evidence suggests that aperiodic slope is modulated by the balance between excitation and inhibition, with increased slope associated with a reduction in E/I ratio[11,14,53]. An age-dependent reduction in E/I ratio during the first postnatal year is consistent with the prolonged developmental timing of inhibitory network maturation in humans. Unlike excitatory neurons which are well established by birth, during the first postnatal year GABAergic inhibitory neurons continue to migrate from ventral subregions of the brain to the cortices where they ultimately mature and integrate into neuronal networks[54]. In addition, during this first year GABAergic responses switch from being excitatory to inhibitory due to changes in the concentration of chloride channels on cell membranes[55–57]. Overall, inhibitory neuron network integration and the excitatory-to-inhibitory GABA switch are unique to this developmental period and likely lead to increased inhibitory tone during the first year.

Observed changes in the periodic spectra may reflect sequential steps in inhibitory network and thalamocortical circuit development. Transient neural circuits are common in postnatal development and play critical steps in normal development of thalamocortical circuitry[6]. For example, transient circuits

between sublate neurons (SPN) and thalamo-recipient layer 4 spiny stellate neurons help establish thalamocortical connections prior to the maturation of primary sensory cortices[58]. Studies of postmortem fetal monkey and human brains suggests that the SPN in primates and humans slowly begins to disappear in the 3rd trimester but may persist until 6 months, with an overlapping period in which the thalamus makes connections with both the SPN and cortical layer IV neurons[6,59,60]. We hypothesize that the 9.5Hz peak observed at 2-4 months, but not at 6 months, reflects this transient period when mature excitatory subplate neurons are still receiving and relaying thalamic input to cortices, resulting in higher frequency alpha oscillations. Additionally, newly established connections between the thalamus and layer IV produce lower frequency theta rhythms that will later become the dominant alpha rhythm.

The thalamus is thought to play a central role in the generation of the mature posterior alpha rhythm[61–63]. A shift in dominant oscillatory frequency in the theta/alpha range (4-12Hz) across early childhood has been observed now in many studies[28,32,64]. Here, we both confirm and extend those findings over the first 3 years after birth, with peak frequency increasing most between 4 and 18 months. What factors are potential contributors to this shift in peak frequency? The dynamic circuit motif model (DCM) proposes that cortical network rhythms result from a combination of the intrinsic resonant frequency of a neuronal population and the time course properties of the inhibitory inputs on the neuronal population[65]. Under the DCM model, prior to the maturation of both local inhibitory circuitry and thalamocortical feedback loops, peak frequency oscillations as measured by scalp EEG are more likely to represent the intrinsic properties of cortical networks, with thalamic inputs beginning to play a larger role with age. For example, lower frequency 4-7Hz oscillations are intrinsically generated by *isolated* layer 5 cortical neurons, and the range of oscillations increases to 5-12Hz when connections to other cortical layers remain intact[66]. Thalamic neurons in the lateral geniculate nucleus also fire across the theta and alpha range. In vitro slice experiments from cats suggest that cortical input to thalamus modulates whether theta versus alpha oscillations are dominant[22,23]. Thus, the developmental shift in peak frequency from the canonical theta to alpha range over the first three years after birth, may represent the integration and maturation of cortical inhibitory neurons, as well as the establishment and maturation of thalamocortical connections.

Finally, our study identified early age-dependent changes in periodic beta power that we hypothesize are associated with thalamocortical loop maturation. We observe the emergence of a low beta peak in infants older than 6-months of age and find that the presence of a low beta peak is associated with higher anesthesia-induced frontal alpha coherence. Biophysical models demonstrate that this frontal anesthesia-induced alpha coherence requires inputs from both the thalamus and cortex[38]. Together these findings suggest that low beta oscillations may directly reflect thalamocortical loop maturation. Beta rhythms are thought to be generated both locally in the cortex through pyramidal-interneuron loops, as well as through thalamus to cortical connections that also rely on inhibitory inputs[33]. The emergence of

the low beta peak in awake infants may reflect the combination of newly established network connections between thalamic nuclei and cortical layers, as well as the maturation of interneurons within the thalamocortical pathways.

It is also possible that developmental changes in beta power are related to infant movement. During EEG acquisition, infants are held in their parent's lap and behavioral supports are in place to keep the infants calm. However, it is not possible to control the infants' movement, and movements both small (hand movements) and large (head turns, leg and arm movements) ubiquitously occur across recordings - likely increasing over the first year as infants become more mobile. Our preprocessing artifact removal pipelines (see Methods) includes several steps for removing high frequency noise from muscle artifact. However this would not remove EEG signal in response to sensorimotor processing. Infant jaw and upper limb movements have been shown to increase power between 9–20Hz along frontal and occipital sites, while hand and lower limb movements do not have significant effects[67]. In our dataset, increases in low beta power were most prominent in central (not frontal or occipital) ROIs (Supplemental Fig 3K). In addition, age-dependent shifts of both low and high beta are observed visually across individual power spectra plots (Supplemental Figure 2) and similar age-dependent shifts are observed in infants who were noted to have either limited or substantial movement during baseline EEG acquisition (Supplemental Figure 9).

Transient high beta peaks were also observed across this early period of development. Specifically, we observed early increases in high beta peak amplitude, which reached a maximum at 12.2 months, followed by decreases in both high beta peak amplitude and frequency, such that by 36 months the low beta peak is the dominant peak across the 12-30Hz range. The neurobiological mechanism of this high beta peak is unclear. As discussed above, administration of GABA agonists induces beta activity. In addition, several neurodevelopmental disorders associated with GABA receptor dysfunction show prominent beta peaks on EEG; individuals with Duplication 15q have a prominent beta peak at 23Hz[68,69], and we have observed that children with Fragile X Syndrome (FXS), aged 3-7 years, have a prominent 30Hz peak[70]. This 30hz peak observed in FXS children is qualitatively similar to the 30Hz peak observed in the present dataset at a much younger age. Further analysis of data previously published from FXS children shows that the observed high beta peak decreases with age (Figure 5), suggesting delayed brain development. Such observations highlight the value of the longitudinal EEG trajectories presented in this paper in placing findings from neurodevelopmental disorders in the broader context of developmental brain maturation.

**Figure 5. FXS children have increased high beta peak that decreases with age. (A)** Periodic power spectra of 35-48 month-old children with and without FXS. **(B)** Individual periodic power spectra of FXS children from 32-84 months old, with line hue corresponding to age in months.

1
2
There are several limitations that may affect the interpretation or generalizability of findings. First, there
are significant non-neural anatomical developmental changes that occur during infancy, including
increases in skull thickness, gradual closure of fontanels and sutures, and changes in cerebrospinal fluid
volume. Such changes can alter conductive properties of the skull, and in turn impact EEG signals,
including measures such as aperiodic offset and slope[71–73]. Second, more work is needed to firmly
establish the association between changes in periodic power and development of thalamocortical
circuitry. While we provide some preliminary evidence that the emergence of low beta is associated with
maturation of thalamocortical connectivity, the dataset is small and still an indirect measure. Future
research that combines EEG and MRI in infants from 4-6 months and again at 10-15 months, could
provide more direct evidence linking alpha and beta periodic EEG findings with the development of
thalamocortical connectivity. Third, impedance thresholds were kept below 100kΩ, rather than 50 kΩ, in
order to reduce the time needed to optimize impendences and in turn prevent infants and toddlers from
becoming fussy. While all data were collected in electrically-shielded rooms, it is possible that recordings
with impedances above 50 kΩ have reduced signal quality. However the developmental changes
described are robust and observed across individuals as observed in Supplemental Figure 2. Fourth, the
findings presented here utilize a specific artifact removal and processing pipeline (BEAPP and
HAPPE1.0). An automated pipeline was used to improve reproducibility of our findings, and the code is
available on Open Science Framework (https://osf.io/u3gp4). We do note that similar developmental

changes in the low and high beta from 9 and 12 months are shown in Figure 1 of Rayson et al. 2023 which used a combination of MADE and NEAR pipelines. Interestingly, recently Rico-Pico et al. 2023 also described developmental changes in the periodic power spectra using MADE processing pipelines but limited their analysis to 1-20Hz "due to a power bump" in the gamma range, assumed to be related to muscle artifact, but likely the same high beta peak we observe most prominent at 7-8 months of age. Power spectra from 6, 9, and 16 months shown in Supplemental Figure 8 of Rico-Pico et al. 2023 visually show a similar pattern in beta band change across development. We hope that our comprehensive characterization of developmental changes using a single pipeline with frequent age sampling across the first 3 years of development spur other infant EEG researchers to evaluate the patterns we describe in their own data.

In summary, our work highlights the dynamic developmental changes in neural activity occurring during the first three years after birth and provides insights in potential ways these age-dependent and sometimes transient changes may coincide with sequences in thalamocortical and inhibitory network maturation. Our findings help to ground cross-sectional work occurring at these early ages and provide a foundation to compare developmental trajectories of various neurodevelopmental disorders including autism, ADHD, and rare genetic disorders. Future studies examining early trajectories of functional connectivity and phase amplitude coupling across this age range will provide additional insight into the timing of critical periods in brain maturation.

**METHODS**

**Studies and Participants:** Lab based EEGs for this paper were collected as part of four different studies occurring over 15 years conducted at our lab at Boston Children's Hospital (Figure 1A). Sample numbers for each age are shown in Table 1. Study 1, the Healthy Baby Study (IRB-P00019083), was a longitudinal study, enrolling infants starting at 2 months of age, from the Boston Children's Hospital Primary Care Center, which predominantly services families from low-income backgrounds. EEG was collected and developmental assessment using the Mullen Scales of Early Learning (MSEL) was performed at 2, 6, 9, 12, 24, and 36 months.

Study 2, The Infant Sibling Study (IRB-X06-08-0374), and Study 4, the Infant Screening Project (IRB-P00018377), were both prospective, longitudinal studies, enrolling infants with and without first degree family history of ASD starting as early as 3-months of age. For this analysis only infants *without* family history of ASD were included. Study 4 also included a group of infants with elevated social communication concerns at 12 months of age, and they were also excluded from this analysis. EEGs

and MSEL were performed at 3, 12, 18, 24, and 36 months for both studies, as well as 6 and 9 months for Study 2. Infants were specifically assessed for ASD using the Autism Diagnostic Observation Schedule (ADOS) in conjunction with clinical best estimate at 24– and 36–month visits.

Study 3, the Emotion Project (IRB-P00002876), was a cohort/longitudinal study. Infants were enrolled at *either* 5, 7, or 12 months, and then followed through 7 years of age. In addition to the first time point, EEG data was again collected at 3 years of age. There were no developmental assessments performed for this study, however parent questionnaires regarding child development, diagnoses (e.g., ASD), and therapies were collected.

All infants had a minimal gestational age of 36 weeks, no history of prenatal or postnatal medical or neurological problems, and no known genetic disorders. Infants who were later diagnosed with ASD (either by assessment during the study, or by community diagnosis disclosed by parents prior to age of 5) were not included in this study.

Sample characteristics across and within studies are shown in Table 1. The analysis included 1335 EEGs collected from 592 participants. While all studies took place in the same laboratory, participant demographics vary between studies as was expected given differences in recruitment and research aims of each study. In addition, studies differed in the age of enrollment and when subsequent visits were completed (Figure 1C). Combined, the sample remained predominantly white (74%).

**Table 1: Sample Characteristics**

| | Combined Studies N =592 | Study 1 N = 49 | Study 2 N = 72 | Study 3 N = 363 | Study 4 N =108 |
|---|---|---|---|---|---|
| **Sex,** *% Female (n)* | 46.3 (274) | 51.0 (25) | 45.8 (33) | 46.3 (168) | 44.4 (48) |
| **Ethnicity,** *% (n)* | | | | | |
| Hispanic | 8.6 (51) | 16.3 (8) | 1.4 (1) | 10.5 (38) | 3.7 (4) |
| Non-Hispanic | 90.2 (534) | 81.6 (40) | 97.2 (70) | 88.4 (321) | 95.4 (103) |
| Not Answered | 1.2 (7) | 2.0 (1) | 1.4 (1) | 1.1 (4) | 0.9 (1) |
| **Race, % (n)** | | | | | |
| White | 74.0 (438) | 10.2 (5) | 86.1 (62) | 79.1 (287) | 77.8 (84) |
| Black or African-American | 6.4 (38) | 53.1 (26) | 1.4 (1) | 2.2 (8) | 2.8 (3) |
| Asian | 3.9 (23) | 4.1 (2) | 2.8 (2) | 4.1 (15) | 3.7 (4) |
| Mixed Race | 12.8 (76) | 14.3 (7) | 8.3 (6) | 12.9 (47) | 14.8 (16) |
| Other | 1.4 (8) | 12.2 (6) | 0 | 0.6 (2) | 0 |
| Not Answered | 1.5 (9) | 6.1 (3) | 1.4 (1) | 1.1 (4) | 0.9 (1) |
| **Family Income, % (n)*** | | | | | |
| <$35,000 | 5.4 (32) | 32.7 (16) | 4.2 (3) | 2.5 (9) | 3.7 (4) |
| $35,000 - $75,000 | 11.1 (66) | 18.4 (9) | 9.7 (7) | 13.2 (48) | 1.9 (2) |
| >$75,000 | 73.8 (437) | 18.4 (9) | 70.8 (51) | 76.3 (277) | 92.6 (100) |
| Not Answered or Don't Know | 9.6 (57) | 30.6 (15) | 15.3 (11) | 8.0 (29) | 1.9 (2) |
| **Participant EEG Data Included in Analysis,** *n (% with longitudinal data)* | | | | | |
| 2-4 m | 97 (91) | 46 (93) | 10 (100) | - | 41 (85) |
| 4-6 m | 119 (46) | - | 2 (100) | 113 (45) | 4 (50) |
| 6-8 m | 223 (68) | 43 (100) | 50 (100) | 130 (45) | - |
| 8-11 m | 95 (98) | 37 (100) | 58 (98) | - | |
| 11-15 m | 291 (79) | 40 (100) | 62 (100) | 102 (45) | 87 (95) |
| 18-20m | 107 (97) | - | 40 (100) | - | 67 (95) |
| 23-30m | 118 (99) | 22 (100) | 44 (100) | - | 52 (98) |
| 35-44m | 285 (92) | 18 (100) | 48 (95) | 174 (90) | 45 (95) |
| By participant, mean # EEGs ± SD | 3.2 ± 1.7 | 4.6 ± 1.1 | 4.9 ± 1.3 | 1.6 ± 0.5 | 3.2 ± 0.9 |

*2 infants reported to have income of $30-39,000 were placed in <35,000 category. 1 infant with reported income of $70-79,000 placed in >75,000 category.

**Lab-based EEG data collection:** Baseline, non-task-related EEG data was collected using similar

methods and rooms for all four studies. The infant was held by their seated caregiver in a dimly lit, sound

attenuated room with a low-electrical-signal background. For Study 2, a research assistant ensured that

the infant remained calm by blowing bubbles and/or showing toys. For Studies 1 and 3, a video of infant

toys was shown for 2–5 minutes and 2 minutes, respectively. For Study 4, a video of an abstract moving

objects was shown for 2–5 minutes. Continuous scalp EEG for Studies 1, 3, and 4 was recorded using a

128-channel Hydrocel Geodesic Sensor Nets (Electrical Geodesics, Inc., Eugene, OR) connected to a

NetAmps 300 amplifier (Electrical Geodesic Inc.) and sampled at 500Hz. Study 2 included recordings

using 64-channel Geodesic Sensor (<10% of data) or a 128-channel Hydrocel Geodesic Sensor Nets

(Electrical Geodesics, Inc., Eugene, OR), connected to either a NetAmps 200 or 300 amplifier (Electrical

Geodesic Inc.) and sampled at either 250 or 500Hz. Additional statistical analysis related to differences

in net and amps is described below in EEG Power analysis. For all studies, data was referenced online

to a single vertex electrode (Cz) and impedances were kept below 100kΩ in accordance with the

impedance capabilities of the high-impedance amplifiers inside the electrically shielded room.

Electrooculographic electrodes were removed to improve the child's comfort.

**EEG pre-processing:** Raw Netstation (Electrical Geodesics, Inc) files were exported to MATLAB

(version R2017a) for preprocessing and absolute power calculations using the Batch Automated

Processing Platform (BEAPP[74]) with integrated Harvard Automated Preprocessing Pipeline for EEG

(HAPPE[75]). For each EEG, a 1Hz high-pass and 100Hz low-pass filter were applied, data sampled at

500Hz were resampled to 250Hz, and then run through the HAPPE module consisting of 60Hz line noise

removal, bad channel rejection, and artifact removal using combined wavelet-enhanced independent

component analysis (ICA) and Multiple Artifact Rejection Algorithm (MARA[5,6]). The following channels, in

addition to the 10-20 electrodes, were used for MARA: 64-channel net – 16, 9, 8, 3, 58, 57, 21, 25, 18,

30, 43, 50, 53, 32, 33, 38, 41, 45; and 128-channel net - 28, 19, 4, 117, 13, 112, 41, 47, 37, 55, 87, 103,

98, 65, 67, 77, 90, 75. These electrodes were chosen as they evenly cover all brain regions of interest

(Supplemental Figure 1). After artifact removal, channels removed during bad channel rejection were

then interpolated, data were referenced to the average reference, detrended to the signal mean, and

segmented into 2-second segments. Any segments with retained artifact were rejected using HAPPE's

amplitude and joint probability criteria.

**EEG rejection criteria:** EEG recordings were rejected using the following HAPPE data quality

measures: Fewer than 20 segments (40 seconds of total EEG), percent good channels < 80%, percent

independent components rejected >80%, mean artifact probability of components kept > 0.3, and

percent variance retained < 25%. Expected differences between studies in number of segments

remaining post-segment rejection were observed, with Study 3 with the shortest resting state recording

period having fewer segments. All other quality metrics were similar across studies (Table 2).

**Table 2: EEG data quality metrics**

| | Combined Studies N = 1335 | Study 1 N = 206 | Study 2 N = 314 | Study 3 N = 519 | Study 4 N = 296 |
|---|---|---|---|---|---|
| **EEG quality metrics, Mean±SD** | | | | | |
| Number of Segments Retained after processing | 81±39.5 | 126±27.5 | 85.8±39.9 | 48.8±7.1 | 105.2±30.3 |
| Percent Good Channels | 92.2±4.5 | 92.2±4.8 | 92.6±4.4 | 92.0±4.6 | 93.5±4.3 |
| Percent ICs Rejected | 35.7±10.3 | 35.7±10.6 | 35.4±10.1 | 34.1±14.0 | 35.0±10.5 |
| Percent Variance Kept of Post Waveleted Data | 67.2±17.0 | 63.1±16.1 | 66.5±15.7 | 68.8±18.3 | 68.1±16.1 |
| Mean Artifact Probability of Kept ICs. | 0.12±0.05 | 0.12±0.04 | 0.11±0.04 | 0.12±0.05 | 0.12±0.05 |

**EEG Power analysis:** The power spectral density at each electrode, for each 2-second segment, was calculated in the BEAPP Power Spectral Density (PSD) module using a multitaper spectral analysis[44] and three orthogonal tapers. For each electrode, the PSD was averaged across segments, and then further averaged across all available electrodes, or frontal, temporal, central, and posterior regions of interest (Supplemental Figure 1C,D). The PSD was then further analyzed using a modified version of SpecParam v1.0.0[8](https://github.com/fooof-tools/fooof; in Python v3.6.8) in order to model periodic and aperiodic components of the power spectra. SpecParam required modification for use in this age range, as power spectrum models for 2-7 month ages showed poor model fit (increased mean squared error) for frequencies between 10-20Hz. Specifically, the SpecParam modeled curves did not accurately capture the "trough" in the power spectra visually observed in this frequency range at younger ages (Supplemental Figure 7 and 8) when alpha and high beta peaks are prominent, but a low beta peak is not present. In the original SpecParam, the power spectrum is first fit with an estimated aperiodic component which is subtracted from the raw signal. Any negative data, signal falling below 0, is converted to 0, and peaks are identified through iterative gaussian fits. Once all peaks are identified, these peaks are removed from the raw spectra and a final aperiodic component is fit. Between 2-7 months, signal between 10-20Hz often falls below the first estimated aperiodic fit and is thus converted to 0, impacting both the periodic peaks identified and the final fit of the aperiodic component. To improve model fit, the robust_ap_fit function, which initially defines the aperiodic component, was modified so that the initial estimate of the flattened power spectra (flatspec) has a baseline elevated such that the lowest point is ≥ 0, to avoid omitting data important across the 2-7 month age range. In the original and modified scripts, this initial fit is combined with thresholding to render a more robust second round of aperiodic parameters. After these second aperiodic parameters have been defined, the fit function re-estimates the flattened spectra (spectrum_flat). At this point, prior to fitting spectra peaks, the modified code sets

negative data in the flattened spectra equal to 0, similar to the approach of the original code during the initial aperiodic fit. In both versions, aperiodic parameters are fit a third and final time to the spectra with peaks removed (spectrum_peak_rm). The SpecParam model was used in the fixed mode (no spectral knee) with *peak_width_limits* set to [0.5, 18.0], *max_n_peaks* = 7, and *peak_threshold* = 2. Code is available (osf.io/u3gp4) which runs both the original and modified versions of SpecParam (with edits marked in code). Changes are also shown in Supplemental Materials. The code also graphs the RMSE across frequencies for both versions, separated by age. Comparisons model fits for each age for are presented in Supplemental Figure 4, as well as RMSE across frequencies for infants 5, 6, or 7 months old. Further analyses were subsequently restricted to 2.5-50Hz given elevated error between 2-2.5, and 50-55Hz. Mean $R^2$ for the full sample using this modified version of SpecParam was 0.997 (STD 0.008; range 0.890-0.999). Mean $R^2$ for each age bin ranged from 0.991-0.999. Mean estimated error for the sample was 0.01 (STD 0.01, range 0.002-0.09).

SpecParam provides two parameters to describe the aperiodic 1/f signal: offset and slope. As the SpecParam-determined offset is extrapolated to the estimated aperiodic power at 0Hz, where there are high amounts of error, we instead calculated the aperiodic offset based on aperiodic power at 2.5hz (Figure 1A). The periodic power spectrum (Figure 1B, F) was determined by subtracting the SpecParam estimated aperiodic spectrum (Figure 1E) from the absolute power spectrum (Figure 1D). To further characterize peaks within the power spectra across development, the periodic spectrum was then smoothed using a savgol filter (scipy.signal.savgoal_filter, window length = 101, polyorder = 8). Individual periodic power spectrum plots before and after savgol filter are shown in Supplemental Figure 2. We decided to use this method instead of using the SpecParam estimated peak_fit as the high beta peak appeared to have a non-gaussian shape at some ages, and thus peak_fit estimates did not accurately identify the high beta peak frequency. Using the smoothed periodic spectra, maxima were identified within the following frequency ranges: 4-6.5 (theta), 4-12Hz (theta/alpha), 12-20Hz (low beta) and 20-35Hz (high beta). Aperiodic and periodic power across the following canonical frequency bands was calculated taking the integral of each parametrized spectra between the following frequency ranges: theta (4-6Hz), low alpha (6-9Hz), high alpha (9-12Hz), low beta (12-20Hz), high beta (20-30Hz), and gamma (30-45Hz).

To visualize differences aperiodic and periodic EEG measures across different regions, topoplots for 12 measures across eight age bins are shown in Supplemental Figure 6. As SpecParam estimates were based on power spectra across averaged electrodes, topoplots were similarly created using averaged measures across electrodes from 6 regions of interested (left frontal, right frontal, central, left temporal, right temporal, and posterior).

As Study 2 collected data with 2 net types and 2 amplifiers, data from 6-, 9-, and 12-month age bins were assessed for spectra differences in total (2-50hz) aperiodic and periodic power as well as aperiodic slope and intercept measure from central electrodes between either 64 and 128 channel nets, or NetAmps 200 or 300 amplifiers. Of the 24 analyses performed, 3 showed significant differences. Net-type differences were observed for 9- and 12-month central aperiodic slope (p=0.04 for both) and an amplifier difference was observed for 12-month central offset (p=0.04). None of these were significant after correcting for multiple comparisons.

**Anesthesia cohort:** EEG data were collected from infants undergoing anesthesia as part of a prospective observational study approved by the Institutional Review Board at Montefiore Medical Center, Albert Einstein College of Medicine[46,76]. Infants scheduled for elective surgical procedures (e.g., circumcision, hernia repair) were recruited. Infants were excluded for prematurity, known neurologic injury, epilepsy, or planned intracranial surgery. Infants less than 6 months of age, or those documented to be asleep or crying during baseline (pre-anesthesia) recordings were excluded from further analysis. All subjects received general anesthesia with sevoflurane.

EEG recordings were obtained using a Food and Drug Administration (FDA) approved 26 channel device recording from scalp locations designated by the International 10-20 System, reference midline occipital channel (Oz) (microEEG System, Biosignal, Acton, MA)[77]. Data was collected from 21 electrodes (FpZ, Fp1, Fp2, Fz, F3, F4, F7, F8, Cz, C3, C4, T3, T4, Pz, P3, P4, T5, T6, Oz, O1, O2) both prior to (baseline) and during sevoflurane induction and maintenance. End tidal sevoflurane concentration was recorded, locked in time with EEG recording. Data were sampled at a frequency of 250 Hz.

Baseline EEGs (n = 45) were visually inspected and approximately 2 minutes of continuous EEG with minimal artifact was segmented and then processed using BEAPP/HAPPE[74,75]. 9 EEGs were excluded due to excessive artifact during visual inspection. No additional EEGs were excluded based on HAPPE data quality criteria. The final sample (n = 36) had an average age of 9.1 months (range 6-15 months), and was predominantly male (n=25). Central periodic power using Pz and Fz electrodes was calculated using multitaper spectral analysis using three orthogonal tapers. PSD was then further analyzed using a modified version of SpecParam v1.0.0 as described above, in order to model periodic and aperiodic components of the power spectra. Low beta peak was identified as described above from the baseline EEG.

**Alpha coherence analysis**: Preprocessing of Anesthesia EEG data: We used a bipolar montage to analyze the data (F7-Fp1 and F8-Fp2). We developed an automatized method to exclude epochs with high-amplitude noise based on the standard deviation of the time series signal. Three members of the

team (CW, JC, and RG) visually inspected the remaining epochs to select all available 30-second artifact-free segments. We then selected all epochs with a stable sevoflurane concentration defined as two consecutive minutes of end-tidal sevoflurane levels within 0.2% preceding the selection of an epoch of data. EEG data were band-pass filtered from 0.1 to 30 Hz.

For each subject, corresponding EEG data collected during anesthesia were inspected to identify all 30 second segments where both the sevoflurane concentration was stable, and segments were "artifact free" (eg. no motion or electrocautery artifacts). Epochs with high-amplitude noise in frontal electrodes (F7-Fp1; F8-Fp2) based on standard deviation of the time series were automatically excluded. Blinded visual inspection by members of the team (CW, JC, and RG) identified remaining epochs with stable sevoflurane concentration without other anesthesia (e.g. propofol bolus) interference. Stable sevoflurane concentration was defined as two consecutive minutes of end-tidal sevoflurane levels within 0.2% preceding the selection of an epoch of data.

Coherence Analysis: EEG data were band-pass filtered from 0.1 to 30 Hz. Custom EEG analysis scripts were written using MATLAB (version R2021a, MathWorks, Natick, MA), employing functions in the Chronux toolbox[78]. Coherence analysis was calculated between F7 and F8 using the multitaper method with the following parameters: window length T = 6s with no overlap, time- bandwidth product TW = 3, number of tapers K = 5, and spectral resolution of 2W = 1 Hz. We calculated coherence by quantifying the degree of correlation between both signals across a frequency range as previously described[79]:

$$C_{xy}(f) = \frac{\left|S_{xy}(f)\right|}{\sqrt{S_{xx}(f)S_{yy}(f)}}$$

Where $S_{xy}(f)$ is the cross-spectrum between the signals x(t) and y(t) (i.e., F7 and F8 electrodes), $S_{xx}(f)$ is the power spectrum of the signal x(t) and $S_{yy}(f)$ is the power spectrum of the signal y(t). Then, the median coherence within the alpha band was used for analyses.

**Statistical Analyses:**

GAMMs: To assess developmental trajectories of power spectral measures, we used generalized additive mixed models (GAMM). GAMMs are similar to generalized linear mixed models, with the advantage that predicts can be modeled linearly and nonlinearly. In GAMMs, smooth linear or nonlinear functions of the relationship between predictors and the outcome are simultaneously estimated, and then summed. GAMM is therefore an advantageous framework for exploring the relationship between power

spectral measures and age, for which the underlying form of the relationship is not yet known. Models

were fit using *mgcv* package[80] (version 1.8-38) and R (version 4.1.2).

A separate GAMM was fit to predict each power spectral measure, for each region of interest (e.g., whole

brain theta power, frontal aperiodic offset). First, to determine whether to include an age-by-sex

interaction, two models were fit with the following forms:

(1) Power Measure ~ oSex + s(age_days, k = 4, fx = T)  + s(New_ID, bs = 're') + s(Study, bs

9            = 're')"

10       (2) Power Measure ~ oSex + s(age_days, k = 4, fx = T) + s(age_days, by = oSex, k = 4, fx =

11           T) +  s(New_ID, bs = 're') + s(Study, bs = 're')

oSex represents sex stored as an ordered factor; coding sex as an ordered factor is necessary for the

GAMMs model to produce a single significance value for the age-by-sex interaction. This is more

interpretable than the alternative of including sex as a categorical factor where two separate smooth

effects of age are modeled for each sex, but no direct comparison is provided. s(age_days, k = 4, fx = T)

is a smoothed age term. Study and New_ID are each included as random effects to account for repeated

observations and clustering of observations within studies. These models were compared by ANOVA,

and model 2 (including the interaction term) was chosen if the difference was significant ($p<0.05$). To

correct for multiple comparisons, the false discovery rate (FDR) was controlled using the Benjamini and

Hochberg method[81], which was applied for each model term within each region of interest across  the 16

measure types (e.g., theta power, beta power, aperiodic offset) to produce q-values (FDR-corrected p-

values).

To further understand the nonlinear trajectories of change, inflection points were calculated using the

argrelextrema function from scipy in python with order = 100.  A standardized rate of change per day was

calculated to visualize developmental changes within features. The modeled value of a feature at a given

age (in days) was subtracted from the modeled value from the subsequent day, and this was divided by

the standard deviation of the modeled values of that feature across the age range.

To assess the differences between regions of interest, GAMM models were fit with the following form:

(3) Power Measure ~ s(age_days) + oSex + ROI+ s(New_ID, bs = 're') + s(Study, bs = 're')

where the terms have the same meanings as above, and ROI is a factor representing the four regions

(frontal, central, temporal, posterior). Because prior literature and preliminary visual inspection of the

data indicated that the posterior ROI is most unique in the time course of development, the posterior ROI

was set as the reference factor. Thus, the effect and significance associated with each of the other ROIs is a measure of the difference between that ROI and the posterior ROI.

To assess the relationship between age and probability of having one or two peaks within the broad alpha range at 2-6 months, a generalized linear mixed effects model was evaluated using the lme4 package in R[82].

Anesthesia Statistical Analysis: ANCOVA, with sevoflurane levels as a covariate, was used to determine effects of presence of low beta peak on anesthesia induced alpha coherence.

Figures were created using Python v3.6.8 and python data visualization libraries (*matplotlib*(60) and *Seaborn (https://seaborn.pydata.org/index.html*) or in R (version 4.1.2).

**Data availability:** Consents obtained from human participants prohibit sharing of de-identified individual data without data use agreement in place. Please contact the corresponding author with reasonable data requests.

**Code availability:** Code used for EEG processing and analyses used in this paper can be found on the Open Science Framework (https://osf.io/u3gp4).

**Funding Support:** This research was supported by the National Institutes of Health (R01-DC010290 and MH078829 to CAN, K23DC07983 and T32MH112510 to CLW, and UL1TR002556, KL2TR002558, and K23DA057499 to JYC. Research was also supported by the JPB Research Network on Toxic Stress.

**Acknowledgements:** We thank all the children and families who generously participated in this research. We thank all the research staff involved in participant recruitment, data collection, and database administration.

**Ethics Declarations:** Dr. Patrick Purdon is an inventor on pending patents on brain monitoring using the electroencephalogram. One of these patents is under non-exclusive license by Massachusetts General Hospital to Masimo Corporation. Dr. Purdon receives institutionally-distributed licensing royalties for this license. Dr. Purdon is also a co-founder of PASCALL Systems, Inc., a start-up company developing closed-loop physiological control for anesthesiology.

**Author Contributions:**

**Carol Wilkinson:** Conceptualization, Methodology, Software, Formal analysis, Data Curation, Writing – Original Draft, Visualization. **Lisa Yankowitz:** Methdology, Software, Formal analysis, Writing – Review & Editing, Visualization. **Jerry Y. Chao:** Data collection (anesthesia cohort), Methodology, Formal analysis, Investigation, Data Curation, Resources, Writing – Review & Editing. **Rodrigo Gutiérrez:** Methodology, Software, Formal analysis, Data Curation, Writing – Review & Editing. **Jeff L. Rhoades**: Software, Writing – Review & Editing. **Shlomo Shinnar:** Project administration, Supervision, Writing – Review & editing. **Patrick L. Purdon:** Methodology, Supervision, Writing – Review & Editing. **Charles A. Nelson:** Investigation, Resources, Writing – Review & Editing, Supervision, Project administration, Funding acquisition.

**REFEREENCES**

1.  Huttenlocher, P. R. & Dabholkar, A. S. Regional differences in synaptogenesis in human cerebral cortex. *J Comp Neurol* **387**, 167–178 (1997).

2.  Glantz, L. A., Gilmore, J. H., Hamer, R. M., Lieberman, J. A. & Jarskog, L. F. Synaptophysin and postsynaptic density protein 95 in the human prefrontal cortex from mid-gestation into early adulthood. *Neuroscience* **149**, 582–591 (2007).

3.  Tau, G. Z. & Peterson, B. S. Normal Development of Brain Circuits. *Neuropsychopharmacology* **35**, 147–168 (2010).

4.  Kim, J. Y. & Paredes, M. F. Implications of extended inhibitory neuron development. *International Journal of Molecular Sciences* **22**, 1–16 (2021).

5.  Antón-Bolaños, N., Espinosa, A. & López-Bendito, G. Developmental interactions between thalamus and cortex: a true love reciprocal story. *Curr Opin Neurobiol* **52**, 33–41 (2018).

6.  Molnár, Z., Luhmann, H. J. & Kanold, P. O. Transient cortical circuits match spontaneous and sensory-driven activity during development. *Science* **370**, eabb2153 (2020).

7.  Alcauter, S. *et al.* Development of Thalamocortical Connectivity during Infancy and Its Cognitive Correlations. *The Journal of Neuroscience* **34**, 9067 (2014).

8.  Doesburg, S. M., Ward, L. M. & Ribary, U. The alpha-theta-gamma ( ATG ) switch : Toward unified principles of cortical processing. *Current Trends in Neurology* **9**, 1–12 (2015).

9.  Tröndle, M., Popov, T., Dziemian, S. & Langer, N. Decomposing the role of alpha oscillations during brain maturation. *eLife* **11**, e77571 (2022).

10. Manning, J. R., Jacobs, J., Fried, I. & Kahana, M. J. Broadband shifts in local field potential power spectra are correlated with single-neuron spiking in humans. *Journal of Neuroscience* **29**, 13613–13620 (2009).

11. Gao, R., Peterson, E. J. & Voytek, B. Inferring synaptic excitation/inhibition balance from field potentials. *Neuroimage* **158**, 70–78 (2017).

12. Miller, K. J., Sorensen, L. B., Ojemann, J. G. & Nijs, M. den. Power-Law Scaling in the Brain Surface Electric Potential. *PLOS Computational Biology* **5**, e1000609 (2009).

13. Merkin, A. *et al.* Do age-related differences in aperiodic neural activity explain differences in resting EEG alpha? *Neurobiol Aging* **121**, 78–87 (2023).

14. Voytek, B. *et al.* Age-related changes in 1/f neural electrophysiological noise. *Journal of Neuroscience* **35**, 13257–13265 (2015).

15. Hill, A. T., Clark, G. M., Bigelow, F. J., Lum, J. A. G. & Enticott, P. G. Periodic and aperiodic neural activity displays age-dependent changes across early-to-middle childhood. *Developmental Cognitive Neuroscience* **54**, 101076 (2022).

16. McSweeney, M. *et al.* Age-related trends in aperiodic EEG activity and alpha oscillations during early-to middle-childhood. *Neuroimage* **269**, 119925 (2023).

17. Buzsáki, G., Logothetis, N. & Singer, W. Scaling brain size, keeping timing: evolutionary preservation of brain rhythms. *Neuron* **80**, 751–764 (2013).

18. Donoghue, T. *et al.* Parameterizing neural power spectra into periodic and aperiodic components. *Nature Neuroscience* **23**, 1655–1665 (2020).

19. Buzsáki, G. & Chrobak, J. J. Temporal structure in spatially organized neuronal ensembles: a role for interneuronal networks. *Curr Opin Neurobiol* **5**, 504–510 (1995).

20. Buzsáki, G., Lai-Wo S., L. & Vanderwolf, C. H. Cellular bases of hippocampal EEG in the behaving rat. *Brain Research Reviews* **6**, 139–171 (1983).

21. Lopes da Silva, F. Neural mechanisms underlying brain waves: from neural membranes to networks. *Electroencephalography and Clinical Neurophysiology* **79**, 81–93 (1991).

22. Hughes, S. W. & Crunelli, V. Thalamic mechanisms of EEG alpha rhythms and their pathological implications. *The Neuroscientist : a review journal bringing neurobiology, neurology and psychiatry* **11**, 357–372 (2005).

23. Hughes, S. W. & Crunelli, V. Just a phase they're going through: The complex interaction of intrinsic high-threshold bursting and gap junctions in the generation of thalamic α and θ rhythms. *International Journal of Psychophysiology* **64**, 3–17 (2007).

24. Klimesch, W. Alpha-band oscillations, attention, and controlled access to stored information. *Trends in Cognitive Sciences* **16**, 606–617 (2012).

25. Ball, G. *et al.* Thalamocortical Connectivity Predicts Cognition in Children Born Preterm. *Cerebral Cortex* **25**, 4310–4318 (2015).

26. Foxe, J. & Snyder, A. The Role of Alpha-Band Brain Oscillations as a Sensory Suppression Mechanism during Selective Attention. *Frontiers in Psychology* **2**, (2011).

27. Orekhova, E. V., Stroganova, T. A. & Posikera, I. N. Theta synchronization during sustained anticipatory attention in infants over the second half of the first year of life. *International Journal of Psychophysiology* **32**, 151172 (1999).

28. Marshall, P. J., Bar-Haim, Y. & Fox, N. A. Development of the EEG from 5 months to 4 years of age. *Clinical Neurophysiology* **113**, 1199–1208 (2002).

29. Lindsley, D. B. A Longitudinal Study of the Occipital Alpha Rhythm in Normal Children: Frequency and Amplitude Standards. *The Pedagogical Seminary and Journal of Genetic Psychology* **55**, 197–213 (1939).

30. Smith, J. R. The Frequency Growth of the Human Alpha Rhythms During Normal Infancy and Childhood. *The Journal of Psychology* **11**, 177–198 (1941).

31. Rico-Picó, J. *et al.* Early development of electrophysiological activity: Contribution of periodic and aperiodic components of the EEG signal. *Psychophysiology* **60**, e14360 (2023).

32. Cellier, D., Riddle, J., Petersen, I. & Hwang, K. The development of theta and alpha neural oscillations from ages 3 to 24 years. *Developmental Cognitive Neuroscience* **50**, 100969 (2021).

33. Spitzer, B. & Haegens, S. Beyond the Status Quo: A Role for Beta Oscillations in Endogenous Content (Re)Activation. *eNeuro* **4**, (2017).

34. Kuki, T. *et al.* Contribution of parvalbumin and somatostatin-expressing GABAergic neurons to slow oscillations and the balance in beta-gamma oscillations across cortical layers. *Frontiers in Neural Circuits* **9**, (2015).

35. Chen, G. *et al.* Distinct Inhibitory Circuits Orchestrate Cortical beta and gamma Band Oscillations. *Neuron* **96**, 1403-1418.e6 (2017).

36. McCarthy, M. M., Brown, E. N. & Kopell, N. Potential network mechanisms mediating electroencephalographic beta rhythm changes during propofol-induced paradoxical excitation. *J Neurosci* **28**, 13488–13504 (2008).

37. Purdon, P. L., Sampson, A., Pavone, K. J. & Brown, E. N. Clinical Electroencephalography for Anesthesiologists: Part I: Background and Basic Signatures. *Anesthesiology* **123**, 937–960 (2015).

38. Ching, S., Cimenser, A., Purdon, P. L., Brown, E. N. & Kopell, N. J. Thalamocortical model for a

propofol-induced α-rhythm associated with loss of consciousness. *Proceedings of the National*

*Academy of Sciences* **107**, 22665–22670 (2010).

39. Cornelissen, L., Kim, S.-E., Purdon, P. L., Brown, E. N. & Berde, C. B. Age-dependent

electroencephalogram (EEG) patterns during sevoflurane general anesthesia in infants. *Elife* **4**,

e06513 (2015).

40. Cornelissen, L. *et al.* Electroencephalographic markers of brain development during sevoflurane

anaesthesia in children up to 3 years old. *British Journal of Anaesthesia* **120**, 1274–1286 (2018).

41. Akeju, O. *et al.* Age-dependency of sevoflurane-induced electroencephalogram dynamics in children.

*Br J Anaesth* **115 Suppl 1**, i66–i76 (2015).

42. Flores, F. J. *et al.* Thalamocortical synchronization during induction and emergence from propofol-

induced unconsciousness. *Proc Natl Acad Sci U S A* **114**, E6660–E6668 (2017).

43. Bastos, A. M. *et al.* Neural effects of propofol-induced unconsciousness and its reversal using

thalamic stimulation. *eLife* **10**, e60824 (2021).

44. Babadi, B. & Brown, E. N. A review of multitaper spectral analysis. *IEEE Transactions on Biomedical*

*Engineering* **61**, 1555–1564 (2014).

45. Engel, A. K. & Fries, P. Beta-band oscillations--signalling the status quo? *Curr Opin Neurobiol* **20**, 156–

165 (2010).

46. Chao, J. Y. *et al.* Electroencephalographic Findings and Clinical Behavior During Induction of

Anesthesia With Sevoflurane in Human Infants: A Prospective Observational Study. *Anesthesia &*

*Analgesia* **130**, e161 (2020).

47. Miller, K. J. Broadband spectral change: Evidence for a macroscale correlate of population firing

rate? *Journal of Neuroscience* **30**, 6477–6479 (2010).

48. Knickmeyer, R. C. *et al.* A structural MRI study of human brain development from birth to 2 years. *Journal of Neuroscience* **28**, 12176–12182 (2008).

49. Gilmore, J. H. *et al.* Longitudinal development of cortical and subcortical gray matter from birth to 2 years. *Cereb Cortex* **22**, 2478–2485 (2012).

50. Gilmore, J. H., Knickmeyer, R. C. & Gao, W. Imaging structural and functional brain development in early childhood. *Nat Rev Neurosci* **19**, 123–137 (2018).

51. Schaworonkow, N. & Voytek, B. Longitudinal changes in aperiodic and periodic activity in electrophysiological recordings in the first seven months of life. *Dev Cogn Neurosci* **47**, 100895 (2021).

52. Chini, M., Pfeffer, T. & Hanganu-Opatz, I. An increase of inhibition drives the developmental decorrelation of neural activity. *eLife* **11**, e78811 (2022).

53. Voytek, B. & Knight, R. T. Dynamic Network Communication as a Unifying Neural Basis for Cognition, Development, Aging, and Disease. *Biological Psychiatry* **77**, 1089–1097 (2015).

54. Paredes, M. F. *et al.* Extensive migration of young neurons into the infant human frontal lobe. *Science* **354**, aaf7073 (2016).

55. Mueller, A. L., Taube, J. S. & Schwartzkroin, P. A. Development of hyperpolarizing inhibitory postsynaptic potentials and hyperpolarizing response to gamma-aminobutyric acid in rabbit hippocampus studied in vitro. *J Neurosci* **4**, 860–867 (1984).

56. Cherubini, E., Gaiarsa, J. L. & Ben-Ari, Y. GABA: an excitatory transmitter in early postnatal life. *Trends Neurosci* **14**, 515–519 (1991).

57. Xu, G. *et al.* Late Development of the GABAergic System in the Human Cerebral Cortex and White Matter. *J Neuropathol Exp Neurol* **70**, 841–858 (2011).

58. Kanold, P. O. & Shatz, C. J. Subplate neurons regulate maturation of cortical inhibition and outcome of ocular dominance plasticity. *Neuron* **51**, 627–638 (2006).

59. Kostovic, I. & Rakic, P. Developmental history of the transient subplate zone in the visual and somatosensory cortex of the macaque monkey and human brain. *Journal of Comparative Neurology* **297**, 441–470 (1990).

60. McQuillen, P. S., Sheldon, R. A., Shatz, C. J. & Ferriero, D. M. Selective Vulnerability of Subplate Neurons after Early Neonatal Hypoxia-Ischemia. *J Neurosci* **23**, 3308–3315 (2003).

61. Hughes, S. W. *et al.* Thalamic Gap Junctions Control Local Neuronal Synchrony and Influence Macroscopic Oscillation Amplitude during EEG Alpha Rhythms. *Front Psychol* **2**, 193 (2011).

62. Lorincz, M. L., Kékesi, K. A., Juhász, G., Crunelli, V. & Hughes, S. W. Temporal framing of thalamic relay-mode firing by phasic inhibition during the alpha rhythm. *Neuron* **63**, 683–696 (2009).

63. Saalmann, Y. B., Pinsk, M. A., Wang, L., Li, X. & Kastner, S. The pulvinar regulates information transmission between cortical areas based on attention demands. *Science* **337**, 753–756 (2012).

64. Orekhova, E. V., Stroganova, T. A., Posikera, I. N. & Elam, M. EEG theta rhythm in infants and preschool children. *Clinical Neurophysiology* **117**, 1047–1062 (2006).

65. Womelsdorf, T., Valiante, T. A., Sahin, N. T., Miller, K. J. & Tiesinga, P. Dynamic circuit motifs underlying rhythmic gain control, gating and integration. *Nat Neurosci* **17**, 1031–1039 (2014).

66. Silva, L. R., Amitai, Y. & Connors, B. W. Intrinsic Oscillations of Neocortex Generated by Layer 5 Pyramidal Neurons. *Science* **251**, 432 (1991).

67. Georgieva, S. *et al.* Toward the Understanding of Topographical and Spectral Signatures of Infant Movement Artifacts in Naturalistic EEG. *Frontiers in Neuroscience* **14**, (2020).

68. Frohlich, J. *et al.* A Quantitative Electrophysiological Biomarker of Duplication 15q11.2-q13.1 Syndrome. (2016) doi:10.1371/journal.pone.0167179.

69. Frohlich, J. *et al.* Mechanisms underlying the EEG biomarker in Dup15q syndrome. *Molecular Autism* **10**, 29 (2019).

70. Wilkinson, C. L. & Nelson, C. A. Increased aperiodic gamma power in young boys with Fragile X Syndrome is associated with better language ability. *Molecular Autism* 1–15 (2021) doi:10.1186/s13229-021-00425-x.

71. Gajawelli, N. *et al.* Neurocranium thickness mapping in early childhood. *Sci Rep* **10**, 16651 (2020).

72. Lew, S. *et al.* Effects of sutures and fontanels on MEG and EEG source analysis in a realistic infant head model. *Neuroimage* **76**, 282–293 (2013).

73. Antonakakis, M. *et al.* Inter-Subject Variability of Skull Conductivity and Thickness in Calibrated Realistic Head Models. *NeuroImage* **223**, 117353 (2020).

74. Levin, A. R., Méndez Leal, A. S., Gabard-Durnam, L. J. & O'Leary, H. M. BEAPP: The Batch Electroencephalography Automated Processing Platform. *Frontiers in Neuroscience* **12**, 513 (2018).

75. Gabard-Durnam, L. J., Mendez Leal, A. S., Wilkinson, C. L. & Levin, A. R. The Harvard Automated Processing Pipeline for Electroencephalography (HAPPE): standardized processing software for developmental and high-artifact data. *Frontiers in Neuroscience* **12**, 97 (2018).

76. Chao, J. Y. *et al.* Decreased Electroencephalographic Alpha Power During Anesthesia Induction Is Associated With EEG Discontinuity in Human Infants. *Anesthesia & Analgesia* **135**, 1207–1216 (2022).

77. Grant, A. C. *et al.* Diagnostic accuracy of microEEG: a miniature, wireless EEG device. *Epilepsy Behav* **34**, 81–85 (2014).

78. Bokil, H., Andrews, P., Kulkarni, J. E., Mehta, S. & Mitra, P. P. Chronux: a platform for analyzing neural signals. *J Neurosci Methods* **192**, 146–151 (2010).

79. Akeju, O. *et al.* Effects of sevoflurane and propofol on frontal electroencephalogram power and coherence. *Anesthesiology* **121**, 990–998 (2014).

80. Wood, S. N. *Generalized Additive Models: An Introduction with R, Second Edition*. (CRC Press, 2017).

81. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B (Methodological)* **57**, 289–300 (1995).

82. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using lme4. *J. Stat. Soft.* **67**, 1–48 (2015).

**Supplemental Figure 1 - Electrode layout:** (A) 128-channel Hydrocel Geodesic Sensor Net. (B) 64-channel Geodesic Sensor Net. Pink circles denote 10-20 electrodes, and blue circles denote the additional electrodes included in ICA and MARA steps of pre-processing. (C), (D) Electrodes averaged for frontal (yellow), central (blue), temporal (orange), and posterior (green) regions of interest.

**Supplemental Figure 2:** Individual plots of the periodic spectrum averaged across the whole ROI for each age bin.

**Supplemental Figure 3:** Spaghetti plots of aperiodic and periodic power measures

**Supplemental Figure 4:** Spaghetti plots of aperiodic and periodic power measures limited to participants with more than 4 data points and graphed by Age bins.

**Supplemental Figure 5:** GAMMs modeled trajectories of 12 aperiodic and periodic power measures, with ROI, study, smoothed age, and sex as predictor terms.

**Supplemental Figure 6**

## Supplemental Figure 6 cont.

**Supplemental Figure 6:** Modeled estimates of the 12 aperiodic and periodic power measures from the for the four ROIs shown in Supplemental Figure 1 are shown topographically.

*Note, at 2-4 months two peaks are present in the 4-12hz frequency range. Peak amplitude and frequency shown here is based on which ever peak had higher amplitude.

**Supplemental Figure 7:** For each age bin the average SpecParam modeled spectra (green) across participants in each age bin is graphed along with the averaged original power spectrum (red), and averaged SpecParam estimated aperiodic spectrum (blue). (A) Unedited SpecParam estimates. (B) Modified SpecParam estimates. (C) Comparison of squared error across frequencies of unedited (orange) and modified (blue) SpecParam.

**A. Unedited SpecParam model estimates**   - Original Spectrum   - SpecParam modeled Spectrum   - Aperiodic Fit

**B. Modified SpecParam model estimates**   - Original Spectrum   - SpecParam modeled Spectrum   - Aperiodic Fit

**Supplemental Figure 8:** Comparison of squared error across frequencies based of unedited (orange) and modified (blue) SpecParam estimates of whole brain power.

**Supplemental Figure 9:** Periodic power graphed by behavioral scores provided during EEG acquisition of Study 4. No other studies included behavioral ratings. At the time of acquisition a behavioral score 1-5 was provided by the research assistant based on the infant's behavioral regulation during baseline EEG acquisition, with higher scores indicating more behavioral compliance.

### Number of participants

| Behavior Score | | 2m | 12m | 18m | 24m | 36m |
|---|---|---|---|---|---|---|
| 2 | Frequently upset, only short periods of calm between agitation | 0 | 44 | 28 | 22 | 14 |
| 3 | Sometimes upset (30-50% of session) with long periods of calm between protests | 13 | 69 | 42 | 25 | 29 |
| 4 | Rarely upset with agitation/upset 10-30% of session | 17 | 54 | 39 | 33 | 24 |
| 5 | Calm for majority (90%) of session | 10 | 22 | 27 | 18 | 23 |

**Description of code edits to SpecParam. Please see code on Open Science Framework: osf.io/u3gp4**

In the original robust_ap_fit function (https://github.com/fooof-tools/fooof/blob/5e655d73c9d7a47d0411b5177657aaed67c69d6d/specparam/objs/fit.py#L964-L1024), the first estimated flatspec was calculated by subtracting the initial ap fit from the power spectra, AND any value below 0 was converted to 0 (line 993). This leads to the increased error between the original spectrum and the foood estimated spectrum in the 10-20Hz region for 2-7 month olds, since many values fell below 0. Our modified code 'new_robust_ap_fit' elevates the first estimated flat spec such that the lowest point in the flatspec is ≥ 0.

```
24      # Flatten power_spectrum based on initial aperiodic fit
25      flatspec = power_spectrum - initial_fit
26
27      # OLD: Flatten outliers, defined as any points that drop below 0
28      # flatspec[flatspec < 0] = 0   #ORIGINAL
29      # NEW: Increase baseline to prevent fitting negative values
30      if min(flatspec) < 0:
31          flatspec -= min(flatspec)
32
```

Following this step, a second more robust aperiodic fit is estimated, and then the fit function re-estimates the flattened spectra (spectrum_flat). The original fit function found here: https://github.com/fooof-tools/fooof/blob/5e655d73c9d7a47d0411b5177657aaed67c69d6d/specparam/objs/fit.py#L427-L525. Our modified code then sets any negative data in the flattened spectra equal to 0 (similar to the approach used in the original code during the initial aperiodic fit).

```
41      # In rare cases, the model fails to fit, and so uses try / except
42      try:
43
44          # Fit the aperiodic component
45  #         self.aperiodic_params_ = self._robust_ap_fit(self.freqs, self.power_spectrum)  #ORIGINAL
46          self.aperiodic_params_ = self._new_robust_ap_fit(self.freqs, self.power_spectrum)  #NEW
47          self._ap_fit = fooof.sim.gen.gen_aperiodic(self.freqs, self.aperiodic_params_)  #edited to make standalo
48
49          # Flatten the power spectrum using fit aperiodic fit
50          self._spectrum_flat = self.power_spectrum - self._ap_fit
51
52          self._spectrum_flat[self._spectrum_flat < 0] = 0   #NEW
53
```

Supplemental Table 1: GAMMs statistics for models of aperiodic offset and slope, and periodic power features

| | Aperiodic Offset | Aperiodic Slope | 4-12Hz Peak Amplitude | 4-12Hz Peak Frequency | High Beta Peak Amplitude | High Beta Peak Frequency |
|---|---|---|---|---|---|---|
| Age F (q value) | 93.42 (0) | 8.43 (0) | 138.3 (0) | 416.62 (0) | 59.31 (0) | 48.59 (0) |
| Sex F (q value) | -0.03 (0.98) | 0.7 (0.60) | -0.54 (0.6708) | 2.23 (0.14) | 1.31 (0.41) | 2.3 (0.14) |
| AgexFemale F (qvalue) | 5.28 (0.0019) | 3.04 (0.0282) | NA (NA) | NA (NA) | NA (NA) | NA (NA) |

Supplemental Table 2: GAMMs statistics for models of periodic power across 6 frequency bands.

| | Periodic Power | | | | | |
|---|---|---|---|---|---|---|
| | Theta | Low Alpha | High Alpha | Low Beta | High Beta | Gamma |
| Age F (q value) | 115.12 (0) | 243.83 (0) | 440.63 (0) | 133.04 (0) | 28.19 (0) | 220.69 (0) |
| Sex F (q value) | --1.86 (0.25) | 0.76 (0.60) | 1.27 (0.41) | 0.92 (0.57) | 1.01 (0.56) | 2.9 (0.06) |
| AgexFemale F (q value) | 5.5 (0.0019) | NA (NA) | NA (NA) | NA (NA) | NA (NA) | NA (NA) |

Supplemental Table 3: GAMM model statistics assessing differences between regions of interest.

| | Periodic Power | | | | | |
|---|---|---|---|---|---|---|
| | Theta | Low Alpha | High Alpha | Low Beta | High Beta | Gamma |
| Frontal F (q value) | 2.25 (0.025) | -10.61 (0) | -18.45 (0) | 6.88 (0) | 29.76 (0) | 20.1 (0) |
| Central F (q value) | 4.32 (0) | 16.31 (0) | 10.64 (0) | 26.78 (0) | 40.34 (0) | 28.66 (0) |
| Temporal F (q value) | 8.96 (0) | -6.43 (0) | -15.98 (0) | 6.73 (0) | 30.32 (0) | 17.85 (0) |