Growth directions and stiffness across cell layers determine whether tissues stay smooth or buckle

Avilash Singh Yadav[2]†, Lilan Hong[1]†, Patrick M. Klees[2], Annamaria Kiss[3], Manuel Petit[3], Xi He[1], Iselle M. Barrios[2], Michelle Heeney[2], Anabella Maria D. Galang[2], Richard S. Smith[4], Arezki Boudaoud[5], and Adrienne H.K. Roeder[2]*

[1]Institute of Nuclear Agricultural Sciences, Key Laboratory of Nuclear Agricultural Sciences of Ministry of Agriculture and Zhejiang Province, College of Agriculture and Biotechnology, Zhejiang University, Hangzhou 310058, China
[2]Weill Institute for Cell and Molecular Biology and Section of Plant Biology, School of Integrative Plant Sciences, Cornell University, Ithaca, NY 14853, USA
[3]Laboratoire Reproduction et Développement des Plantes, Univ Lyon, ENS de Lyon, UCB Lyon, CNRS, INRAE, INRIA, F-69342 Lyon, France
[4]The John Innes Centre, Norwich NR4 7UH, UK
[5]LadHyX, Ecole Polytechnique, CNRS, IP Paris, 91128 Palaiseau Cedex, France
†These authors contributed equally to this work.
*Corresponding author. ahr75@cornell.edu

Abstract

From smooth shapes to buckles, nature exhibits organs of various shapes and forms. How cells grow to produce smooth shaped leaves and sepals remain unclear. Here, we show that growth along the longitudinal axis during early developmental stages and comparable stiffness across both epidermal layers of Arabidopsis sepals are essential for smoothness, as seen in the wild type. We identified a mutant (*as2-7D*) with ectopic expression of *ASYMMETRIC LEAVES 2* (*AS2*) on the outer epidermis. Our analysis reveals that ectopic *AS2* expression causes the outer epidermis of *as2-7D* sepals to buckle during early stages of sepal development. We show that buckling of the outer epidermis occurs due to conflicting cell growth directions and unequal tissue stiffness across the epidermal layers. Overexpression of cyclin-dependent kinase (CDK) inhibitor Kip-related protein 1 (KRP1) in *as2-7D* restores sepal smoothness by aligning the growth directions of the outer epidermal cells along the longitudinal axis, while also increasing the overall stiffness of the outer epidermis. Furthermore, buckling is associated with the convergence of auxin efflux transporter protein PIN-FORMED 1 (PIN1) to generate outgrowth in the sepals at later stages, suggesting that buckling can initiate outgrowths. Our findings suggest that in addition to molecular cues influencing tissue mechanics, tissue mechanics can also modulate molecular signals, giving rise to well-defined shapes.

One-Sentence Summary

The *asymmetric leaves 2-7D* mutant sepals buckle due to discoordination of growth between the two epidermal layers.

INTRODUCTION

During development, many organs such as the brain, intestine, and orchid flower petals form complex shapes with undulations, folds, and loops [1-6]; conversely, other organs such as Arabidopsis leaf blades and fly wings develop smooth forms [7-10]. These undulations, folds, and loops generally form through mechanical buckling, which is the bending of tissue that occurs when its growth patterns generate compressive forces [11-14]. How cells within tissues regulate their growth and division to generate buckling in certain instances and maintain smoothness in others is not well understood.

Buckling plays a significant role in organ morphogenesis and development across organisms. In animals for instance, numerous studies indicate the role of buckling in organ morphogenesis [2-4,15-17]. Differential growth of connected tissue layers gives rise to buckling. Buckling creates complex organ shapes starting from simple geometries, which is important for functional specialization. These include gyrification of the cerebral cortex to increase neuron content [1], formation of gut villi and looping to increase contact with food and lengthen the gastrointestinal tract [2-4], and initiation of new branches in lung airway epithelium to enhance gas exchange and oxygen transfer [16]. Buckling also serves key roles in plant development. For instance, it generates helical twists in roots to enhance penetrative forces at the growing tip [18,19], triggers lobe deformation in Venus flytraps for rapid energy storage and release to capture prey [20,21], and induces nanoscale cuticular wrinkles on petals to produce iridescence [5,6,22]. Buckling has also been proposed to drive formation and positioning of primordia on the shoot apical meristem [23-25].

Buckling, however, is not always beneficial. Weight of the inflorescence and external forces such as wind or rain can compress and buckle the stems of cereal crops, which causes damage and reduced yield [26]. Similarly, slender tree stems are also vulnerable to buckling [27]. To counter this, plants need to allocate resources to increase stem thickness and stiffness [28,29]. While buckling generates diverse leaf shapes [30-32], the majority of the leaves suppress buckling, generating smooth leaf blades with a relatively flat structure that enhance light absorption and gas exchange for efficient photosynthesis [33,34]. Similarly, sepals, which are the outermost leaf-like floral organs, also exhibit smooth surfaces, which is important for all four sepals to properly enclose and protect the developing flower bud [35]. However, what growth mechanisms and tissue properties underlie the development of smooth organ morphology remain relatively unknown.

Using *Arabidopsis thaliana* (henceforth Arabidopsis) sepals as the model organ, we identified and characterized a gain-of-function mutant of the transcription factor *ASYMMETRIC LEAVES 2* (*AS2*), *as2-7D*, which exhibits folds and outgrowths on the outer epidermis of the sepal. We used

78 *as2-7D* as a tool to investigate what cell growth patterns and mechanical properties of the tissue
79 facilitate the formation of smooth sepals in wild type.
80
81 **RESULTS**
82
83 **Ectopic expression of *ASYMMETRIC LEAVES 2* (*AS2*) disrupts sepal smoothness**
84
To determine how the sepals achieve smooth morphology, we performed a forward genetic
screen [36] and identified an Arabidopsis mutant that disrupts sepal smoothness. This mutant
exhibits an irregular morphology with folds and outgrowths (Fig. 1, A to F). The folds comprise
ridges and invaginations, whereas the outgrowths, which are pointed epidermal projections, are
typically located at the end of the folds (Fig. 1F and fig. S1, A and B). These folds and
outgrowths are visible only on the outer (abaxial) epidermis, while the inner (adaxial) epidermis
is smooth like the wild type (Fig. 1, E to H, fig. S1, A and B and movie S1). Positional cloning
identified a G-to-A point mutation 1,484 bps upstream of the translational start site of
ASYMMETRIC LEAVES 2 (*AS2*) (Fig. 1I). *AS2* encodes a LATERAL ORGAN BOUNDARIES
(LOB) domain transcription factor that plays a critical role in leaf development, particularly in
specifying the upper (adaxial) layer of the leaf [37-40]. An identical mutation in the same locus was
identified previously [41]. However, the fold and outgrowth phenotype in the sepals was not
reported [41], since the phenotype is very subtle in the Columbia (Col-0) background compared to
what we observe in the Landsberg *erecta* (L*er*-0) background (hereafter referred to as wild type)
(Fig. 1, A to D and fig. S1, C to F). Previously, Wu et. al. [41] demonstrated that this mutation in
the *AS2* promoter disrupts the binding site of the repressor KANADI, a transcription factor
involved in specifying cell fate on the lower (abaxial) side of the leaf [42,43]. They further showed
that this mutation leads to ectopic *AS2* expression on the abaxial side, while normally in the wild
type, the *AS2* expression domain is restricted to the adaxial side (Fig.1I). To verify that this
mutation also leads to ectopic *AS2* expression in sepals of the L*er*-0 background, we expressed
GFP under the wild-type *AS2* promoter (*pAS2^{WT}::GUS-GFP*) and mutant *as2-7D AS2* promoter
(*pAS2^{as2-7D}::GUS-GFP*), in the wild-type (L*er*-0) background. While the *AS2* promoter from
wild type drives GFP specifically on the inner sepal epidermal layer (Fig. 1, J and K), the mutant
AS2 promoter drives GFP on both the inner and outer epidermal layers (Fig. 1L). The mutant
AS2 promoter drove GFP expression evenly throughout the outer epidermis, and no instance of
patchy *AS2* expression patterns were observed on the outer epidermis. In addition, *AS2* is still
expressed in its regular expression domain on the inner epidermis in *as2-7D* mutants (Fig. 1L).
This analysis confirms that the mutation causes a gain-of-function through ectopic *AS2*
expression on the outer epidermis, consistent with the dominant nature of this allele (see
Materials and methods). Therefore, following the convention of previous reports [41,44], we named
the mutant allele *as2-7D*.
116

To test whether ectopic *AS2* expression on the outer epidermal layer is sufficient to generate the
folds and outgrowths, we drove *AS2* expression under epidermal specific *MERISTEM LAYER 1*
(*ML1*) promoter (*pML1::AS2*), which ectopically expresses *AS2* on both inner and outer
epidermis of the wild type. All the transgenic plants form lumps and outgrowths on the outer
epidermis of the sepals, similar to *as2-7D*, which suggests that ectopic outer epidermal
expression of *AS2* is sufficient to replicate the *as2-7D* phenotype (Fig. 1, M to P). Additionally,
we do observe folds and outgrowths form on the epidermis of the leaves and fruits (fig. S1, G to
N), consistent with previous reports [39], indicating a potential stronger effect of constitutive
epidermal *AS2* expression in the transgenics than in the *as2-7D* gain-of-function mutant. In
summary, the mutation in *as2-7D* induces ectopic *AS2* expression on the outer epidermal layer,
which is sufficient to disrupt smooth sepal morphology by generating folds and outgrowths on
the outer epidermal layer.
129
130   **The *as2-7D* outer epidermis appears to buckle first and then form outgrowths**
131
To understand how ectopic *AS2* expression on the outer epidermis disrupts smooth sepal
morphology, we live-imaged wild type and *as2-7D* flowers from early sepal developmental
stages (Fig. 2, A and B, fig. S2, A to D, and movie S2). Initially, the outer epidermis of both the
wild type and *as2-7D* sepals are smooth (Fig. 2, A and B; 0-24h). While the outer epidermis of
wild-type sepals remains smooth throughout live imaging, *as2-7D* starts to develop a fold in the
middle of the sepal blade at 48 hours (Fig. 2, B and C). As the flower develops, the folds
intensify further, forming several inward invaginations and outward ridges throughout the outer
epidermis of *as2-7D* mutant (Fig. 2, B to D and movie S2). As the folds intensify over time, we
begin to see the development of phenotypically distinct pointed outgrowths. The outgrowths
become apparent only in the later stages as the folds intensify, which suggests that formation of
the folds precedes the emergence of the outgrowths (Fig. 2, B to D and fig. S2, C and D).
143
We considered two hypotheses for how the folds form: (1) due to localized growth at each ridge
or (2) due to mechanical buckling caused by a surface-wide growth disruption. To test the first
hypothesis, we compared the cell growth rates and division rates between wild type and *as2-7D*
mutant on the outer sepal epidermis at 48-hour interval over a 96-hour growth period (Fig. 2, E
to H and fig. S2, E to N). We do not see any localized fast growth or cell division in the region
where the first fold is formed in *as2-7D* at 48 hours (Fig. 2, C to F). In the initial stages (0-48
hours), the wild type exhibits the stereotypical basipetal growth pattern described earlier [45],
exhibiting fast cell growth at the tip, moderate growth in the center, and slow growth towards the
base. In contrast, the cells grow at relatively homogeneous rates throughout the outer epidermis
of *as2-7D* (Fig. 2, E and F; 0-48 hours). Although the pattern of growth is different, we observe
no difference in the average cell growth rates between the wild type and *as2-7D* (fig. S2M). This
pattern is also reflected in cell division rates, where we observe a basipetal pattern in wild type
and a more uniform pattern throughout the sepal epidermis in *as2-7D* (Fig. 2, G and H; 0-48

hours). Our results suggest that the folds form due to surface-wide disruption of cell growth
patterns, and not because of localized fast cell growth and division at the location of the ridge. In
the later time point (48-96 hrs) the growth is still relatively uniform along the places where the
as2-7D folds form. In contrast to folds, we do observe localized growth and division later at sites
where outgrowths form (Fig. 2F and fig. S2, E to L; 48-96 hrs). We therefore considered the
hypothesis that the folds on *as2-7D* outer epidermis may form through mechanical buckling
caused by surface-wide growth defects.
164
165 **Modeling predicts two criteria required for buckling**
166
Based on our observations of *as2-7D* development, we formulated a simple two-dimensional
(2D) analytical model to determine the basic conditions that could give rise to buckling (see
Materials and methods). Our model indicates that buckling can occur if the following two
conditions are simultaneously fulfilled. Firstly, if the average specified growth rate (the cell-
autonomous growth rate achieved if the layers were isolated) of the outer and inner epidermis is
higher than that of the mesophyll, then the epidermal layers will be under compression, which
will favor buckling in the epidermal layers. Secondly, considering that the mesophyll is known to
be softer than the epidermis [46], a significantly lower stiffness of the outer epidermis compared to
the inner epidermis can cause the outer epidermis specifically to buckle, without causing any
buckling of the inner epidermis.
177
We further tested the validity of the analytical model with simulations on a simple geometry: a
cross-sectional representation of a growing elastic system, comprising of the outer epidermis, the
mesophyll, the inner epidermis, and the edges (Fig. 2, I to K, fig. S3 and S4). In these
simulations, only the outer epidermis was allowed to grow in width, which is equivalent to the
outer epidermis having a higher specified growth rate than the other two layers. Under these
conditions, when the outer and inner epidermis exhibit similar stiffness, the tissue curves while
maintaining a smooth morphology for both the outer and inner epidermis (Fig. 2I). However,
when the outer epidermis is softer than the inner epidermis, the outer epidermis buckles, while
the inner epidermis shows no change in morphology (Fig. 2J). In contrast, increasing the
stiffness of the outer epidermis relative to the inner epidermis restores the smooth morphology of
the outer epidermis, while the inner epidermal morphology remains unaffected (Fig. 2K). In
summary, although our simple analytical model and simulations do not account for all the
complex dynamics of tissue growth in 3D. However, our model does help guide our
investigations towards testing whether differences in growth and tissue stiffness properties
between the outer and inner epidermal layers contribute to buckling in *as2-7D*.
193
194 **Aligned of cell growth orientation between epidermal layers is essential for the formation**
195 **of smooth sepals**
196

To test the first of our model predictions, we live-imaged the wild type and *as2-7D* sepals
through the stages prior to buckling (Fig. 3, A and B), and extracted the inner and outer
epidermal layers of the same growing sepals by image processing (Fig. 3, C and D). We found
that cells throughout the wild-type inner epidermis grow significantly faster than the outer
epidermal cells (Fig. 3E and fig. S5, A, B, and E). We also observe fast cell divisions on the
inner epidermis (Fig. 3G and fig. S5, F, G, and J). The fast growth of the inner epidermis was
surprising, since for the sepals to eventually grow into flat shape with smooth morphology, one
would intuitively expect the cells on both epidermal layers to grow at similar rates. However,
during the initial developmental stages, the sepals are highly curved along the longitudinal
direction, due to the smaller surface area of the inner compared to the outer epidermis. Hence, to
transition from an initially curved shape to a subsequently flat structure, the inner epidermis of
the sepal grows faster compared to the outer epidermis. (Fig. 3C and D, Fig. 4, and movie S2).
Like wild type, the inner epidermal cells of *as2-7D* mutant also grow faster than outer epidermal
cells; however, the inner epidermal cells of *as2-7D* do not grow as fast as wild type (Fig. 3, E
and F and fig. S5, A to E). In concert with the growth rate, the cell division rate is also lower on
the inner epidermis of *as2-7D* compared to wild type (Fig. 3, G and H and fig. S5, F to J).
213
Since buckling is a surface-wide phenomenon in our model, we analyzed the differences in
whole inner versus outer epidermal growth rates. On an epidermis-wide scale, we found that the
outer epidermis of wild type and *as2-7D* grow at the same rate (Fig. 4, A, B and E). This
observation is consistent with the growth rate data of individual cell lineages, where the average
outer epidermal cell growth rates of wild type and *as2-7D* show no significant difference (Fig. 4,
A, B and E and fig. S5E). However, the inner epidermis of *as2-7D* grows significantly slower
than wild type, which also aligns with the growth rate data of individual cell lineages on the
inner epidermis (Fig. 4, C, D and E and fig. S5E). Our findings indicate that the rate at which the
inner epidermis of *as2-7D* grows is insufficient to catch up with the increasing outer epidermal
surface area (Fig. 4F).
224
We then asked whether outer epidermal buckling in *as2-7D* could potentially be attributed to
local hypergrowth of the mesophyll cells underneath the outer epidermis. To test this, we
processed the sepal images using Convoluted Neural Network (see Materials and methods) [47] to
enhance the conspicuity of mesophyll cell membranes (fig. S6, A to H). We analyzed the growth
rates of the outermost layer of mesophyll cells which contains fewer airspaces in early stages,
thus allowing us to lineage track the cells over time. Our findings revealed that the spatial growth
patterns of the mesophyll cells of both wild type and *as2-7D* mimic their respective outer
epidermal cell growth patterns (fig. S6, I and J). Specifically, the wild-type mesophyll cells
exhibit the same basipetal growth gradient as that of the wild-type outer epidermal cells (Fig. 3E
and fig. S6I). The mesophyll cells of *as2-7D* exhibit homogeneous cell growth rates, similar to
the *as2-7D* outer epidermis (Fig. 3F and fig. S6J). Comparable to the outer epidermal cell growth
patterns, we did not find any significant difference in the average growth rate of the mesophyll

cells between wild type and *as2-7D* (fig. S6M). Notably, there was no evidence of local
hypergrowth in the central region of the sepal blade, where the initial buckling occurs (fig. S6J).
To test our model predictions, we then compared the magnitude of growth of the outer and inner
epidermal cells relative to the mesophyll (fig. S6N). We found that the average growth rates of
the inner and outer epidermal cells significantly exceed that of the mesophyll cells for both wild
type and *as2-7D* (fig. S6N), which aligns with the predictions of our model, suggesting that the
epidermal layers are under compression.
244
Since cell growth directions are critical for determining organ shapes [48-50], we assessed the
principal direction of maximal cell growth (PDG_{max}) in the wild type and *as2-7D* mutant. The
wild-type cells on both epidermal layers grow in a highly aligned manner along the sepal
longitudinal axis (Fig. 3I). In *as2-7D*, although inner epidermal cells grow parallel to the sepal
longitudinal axis, the outer epidermal cells exhibit highly misaligned growth, with the majority
of PDG_{max} tensors oriented in the transverse direction (Fig. 3J). Similar to the growth directions
of the outer epidermal cells, the mesophyll cells of wild type grow longitudinally, while the
mesophyll cells of *as2-7D* exhibit misaligned transverse growth (fig. S6, K and L). We
quantified the growth directions of the epidermal cells by calculating the deviation angle of
PDG_{max} tensors relative to the sepal's longitudinal axis (Fig. 3K). The heatmaps indicate minimal
deviation in cell growth directions of the wild-type epidermal layers and the inner epidermal
layer of *as2-7D* (Fig. 3, L and M and fig. S5, K to O). In contrast, the outer layer of *as2-7D*
displayed a high degree of deviation in growth directions (Fig. 3M and fig. S5, K to O). Since
microtubules are known to align perpendicular to the maximal cell growth direction [51,52], we then
tested the microtubule orientation pattern in comparable regions of the wild type and *as2-7D*
outer epidermis (Fig. 3, N and O). For the wild type, the microtubules show the typical
transverse alignment pattern, which is perpendicular to the sepal longitudinal growth direction
(Fig. 3N). Conversely, the microtubules are largely misoriented in the *as2-7D,* and in certain
instances, mirror the transverse cell growth pattern (yellow arrowheads, Fig. 3O).
264
Taken together, overgrowth of the *as2-7D* outer epidermis in width compared to the inner
epidermis contributes to the buckling of the outer epidermis. Our observations align with the
prediction of our model simulations, which indicates that the transverse expansion of the outer
epidermis, constrained along the edges by the inner epidermis would favor buckling. Overall, our
data suggests that anisotropic growth in an aligned manner on both the epidermal layers is
important for maintaining smooth sepal morphology.
271
Since we hypothesized that the transverse growth orientation of the outer epidermal cells causes
the outer epidermis to buckle, we tested whether forcing the cells to grow along the longitudinal
direction can rescue the *as2-7D* phenotype. Overexpressing cyclin-dependent kinase (CDK)
inhibitor Kip-related protein 1 (KRP1) throughout the epidermis (*pML1::KRP1*) at moderate
levels trigger the epidermal cells to skip division and undergo endoreduplication early during

development, resulting in an increase in giant cell formation [53,54]. Giant cells, which are
exclusive to the outer epidermis, exhibit the unique property of growing in a highly anisotropic
manner [53] along the longitudinal sepal axis (Fig. 5, A, C, E and G, fig. S7, A and C, and movie
S3). Therefore, we wondered whether increasing giant cell formation can restore directional
growth in *as2-7D* and consequently reduce buckling. Indeed, overexpressing *KRP1* in *as2-7D*
(*pML1::KRP1 as2-7D*) largely suppresses buckling of the outer sepal epidermis (Fig. 5, A to H,
fig. S7, A to D and movie S3). As expected, overexpressing KRP1 increases the number and size
of the giant cells, which grow in a highly aligned manner along the longitudinal sepal direction
(Fig. 5, E to P and movie S3). The PDG$_{max}$ tensors of *pML1::KRP1 as2-7D* outer epidermal cells
are similar to those of *pML1::KRP1*, wherein the transverse growth orientation exhibited by *as2-*
7D is largely re-oriented longitudinally (Fig. 5, M to P and fig. S7, E to L). In addition to
reduced buckling, the characteristic outgrowths of the *as2-7D* outer epidermis that are formed
downstream of the buckles, are also nearly absent in *pML1::KRP1 as2-7D* (Fig. 5, E to H),
further suggesting that buckling is required for the formation of the outgrowths. While there is no
obvious association between the location of giant cells and small cells and the buckles in the *as2-*
7D mutant, it is possible that the other properties caused by overexpression of *KRP1*, such as
changes in stiffness (verified below), increased outer epidermal thickness, and reduction of cell
division in *pML1::KRP1 as2-7D*, in addition to the growth orientation contributes to the
suppression of buckling. Overall, consistent with our hypothesis, our findings suggest that
transverse growth of the outer epidermal cells may be a key driver of buckling in *as2-7D*.
297
298   **Reduced stiffness of the outer epidermis compared to the inner epidermis is associated with**
299   **buckling of the outer epidermis in *as2-7D* mutants**
300
The second condition of our 2D analytical model predicts that buckling can be restricted to the
outer epidermis if it is softer than the inner epidermis. To test the relative stiffness of the outer
and the inner epidermal layers, we performed osmotic treatments [55,56] and compared how the
epidermal layers of *as2-7D* shrinks relative to wild type. Briefly, when exposed to a
hyperosmotic solution, cells undergo plasmolysis and no longer exert turgor pressure on the cell
wall, causing the cell wall to contract (Fig. 5, Q and R). The extent of cell wall shrinkage gives a
proxy measure for cell wall stiffness, with low shrinkage indicating the wall is stiff (high
Young's modulus) and high shrinkage indicating the wall is soft (low Young's modulus) [55]. We
exposed early sepals to salt solution and measured the epidermal shrinkage (Fig. 5, S and T, fig.
S7, M to P, and see Materials and methods). We found that the wild-type epidermal inner and
outer layers exhibit similar shrinkage, which indicates similar stiffness across epidermal layers
(Fig. 5, U and Y). In contrast, the outer epidermis of *as2-7D* shrank more than the inner
epidermis, indicating that *as2-7D* outer epidermis is softer (Fig. 5, V and Y). This observation is
also consistent with the second condition of our analytical model, which proposes that reduced
stiffness of the outer epidermis is an essential factor for buckling (Fig. 2, I and J). We then
wondered whether increasing giant cell formation affects the stiffness of the outer epidermis. To

test this, we performed osmotic treatments on *ML1::KRP1* and *ML1::KRP1 as2-7D*. Indeed, we found increased stiffness of the outer epidermis in both *ML1::KRP1* and *ML1::KRP1 as2-7D* (Fig. 5, W to Y), which aligns with our simulation predictions that increasing outer epidermal stiffness may restore smooth epidermal morphology (Fig. 2K).

Taken together, our results suggest that aligning growth along the longitudinal direction, decreasing cell division, and increasing outer epidermal stiffness suppresses buckling and subsequent outgrowth formation in *as2-7D*. However, we cannot also decouple the possibility that accelerated differentiation resulting from ectopic *KRP1* expression could also potentially contribute as an additional factor to the rescue. Our findings collectively suggest that relative stiffness and growth orientations of the epidermal layers play a critical role in shaping sepal morphology. Specifically, growth along the same direction and comparable stiffness across both epidermal layers are essential criteria for smooth sepal formation, as observed in the wild type.

Buckles in early *as2-7D* sepals are not caused by an apparent disruption of abaxial-adaxial patterning

We have focused on characterizing how aberrations in growth and mechanical properties lead to buckling on the outer epidermis of *as2-7D* sepals. As described previously, the buckles appear as folds, and these folds are phenotypically very distinct from the outgrowths (which appear as pointed projections later during sepal development). While formation of the folds and its characterization as buckles was previously unreported, the genetic basis of how the outgrowths form on aerial tissues is well established. Waites and Hudson showed nearly 30 years ago how distinct outer (abaxial/lower) and inner (adaxial/upper) tissue identities are established, and more importantly, how the juxtaposition of the outer and inner identities is essential for the lateral growth of the leaf lamina [57]. In the leaves of *phantastica* (*phan*) mutants of *Antirrhinum majus* (snapdragon), patches of inner tissue on the outer surface generate ectopic ridges that resemble lamina margins, indicating that juxtaposition of outer and inner identities can trigger the initiation of ectopic laminar outgrowths [57]. Since the time Waites-Hudson proposed how outgrowths are formed, several studies have independently shown that disruption in the gene regulatory network which determines outer-inner identities can generate ectopic outgrowths [58].

Broadly, outer identity is specified by *KANADI*, *YABBY*, and *AUXIN RESPONSE FACTOR* (*ARF*) gene families, while inner identity is established by the HD-ZIP class III genes, *ASSYMETRIC LEAVES 1* (*AS1*) and *AS2* (Fukushima and Hasebe 2014). While *kan1* and *kan2* single mutants show mild phenotypic defects, the *kan1 kan2* double mutants exhibit radialized organs with ectopic outgrowths on the outer side [42,43]. In the *kan1 kan2 kan4* triple mutants, the ectopic outgrowths are observed even in the hypocotyl and outer surface of the cotyledons [59]. Loss of function double mutants of YABBY genes *FILAMENTOUS FLOWER* (*FIL*) and *YABBY3* (*YAB3*), *fil yab3*, show only mild polarity defects compared to *kan1 kan2*, suggesting

partial redundancy or distinct functions [42]. In line with this, the characteristic folds and

outgrowths seen in *kan1 kan2* are largely absent in the quadruple mutants *kan1 kan2 fil yab3*,

although the leaves are radialized and severely reduced in size [60]. The outgrowths in *kan1 kan2*

double mutants also exhibit elevated levels of *FIL* expression, indicating that YABBYs are

essential for progression of the outgrowths [60].

Initiation of leaf primordium occurs at the boundary between the concentric, non-overlapping

expression domains of *REVOLUTA* (*REV*) (a member of the HD-ZIP class III gene family) and

KAN1 [61]. The *WUSCHEL-RELATED HOMEOBOX* (*WOX*) genes *PRESSED FLOWER*

(*PRS*)/*WOX3* and *WOX1*, which are expressed in the interface between inner and outer domains,

are also required for outgrowth formation[62]. Ectopic *WOX1* expression is sufficient to induce

outgrowth formation. While *prs wox1* double mutants exhibit narrow, upward-curved leaves, the

ectopic outgrowths are highly suppressed in the *prs wox1 kan1 kan2* quadruple mutants, although

the leaves still exhibit a folded morphology [62]. Similarly, *prs wox1 as2* triple mutants fail to

produce outgrowths and instead develop radial filamentous leaves. On the contrary, *ARF2*, *ARF3*

and *ARF4*, which restrict the expression domain of *WOX1/PRS*, repress the formation of ectopic

outgrowths [63]. In particular, the *arf2 arf3 arf4* triple mutants show outgrowths on the outer

surface of the leaves due to expanded *WOX1/PRS* expression [63]. Thus, WOX activity is required

for the formation of ectopic outer outgrowths.

As stated previously, *AS2* is a key transcription factor that is essential for establishing inner

identity[37-40]. Fundamentally, transcription factors cannot drive shape formation without

regulating tissue mechanics. Hence, *AS2* must exert its effects on tissue growth and mechanics

by regulating downstream genes. It is well known that overexpression of *AS2* cause outgrowth

formation on the outer surface of the leaves, and this is accompanied by repression of outer

identity genes (*KAN1*, *KAN2*, *FIL*, *YAB3*) [39]. We have shown that in addition to the outgrowths,

as2-7D also exhibit folds on the outer sepal epidermis. While there exists substantial evidence

for the genetic basis of outgrowth formation, the existence of the epidermal folds, as well as how

these folds are formed in the first place was not well characterized before. Our evidence strongly

supports our hypothesis that the folds observed in the outer epidermis of *as2-7D* in the initial

stages are in fact buckles, formed due to disrupted growth patterns and mechanical properties.

However, since *AS2* is known to promote inner epidermal identity [37-39], we tested the alternative

hypothesis that the initial folds observed on the outer epidermis of *as2-7D* are caused by ectopic

juxtaposition of inner and outer identity. Therefore, we looked at the expression pattern of *REV*

as an inner identity transcription factor [64], and *FIL* as an outer identity transcription factor [65]. We

observe no difference in *REV* and *FIL* expression patterns between wild type and *as2-7D*, and

particularly, no ectopic juxtaposition between inner and outer gene expression was formed in the

as2-7D sepal surface (Fig. 6, A to H). Since REV promotes adaxial identity in conjunction with

AS2, we tested whether REV is necessary or sufficient for the *as2-7D* phenotype. Ectopic *AS2*

expression in the *rev-6* mutant (*pML1::AS2 rev-6*) leads to folds and outgrowth formation similar

to *pML1::AS2* in wild type (Fig. 1, M to P, fig. S1, G to N, Fig. 6, I to L and fig. S8, A to I),
indicating REV is not necessary for either the buckles or the outgrowths. Conversely, ectopic
expression of *REV* in the wild-type sepals is insufficient to generate buckles or outgrowths (fig.
S8, J and K). Further, ectopic *REV* expression in *as2-7D* also does not enhance the phenotype
(fig. S8, L and M).
402
Since abaxial (outer) identity promoting KANADI transcription factors directly repress *AS2*, and
the disruption of KANADI binding site leads to ectopic *AS2* expression in *as2-7D* [41], we
assessed the sepal phenotypes of *kanadi* loss of function mutants (*kan1-2 kan2-1*). Consistent
with previous reports [38,42,43], the *kan1-2 kan2-1* double mutants display severe floral
development and phyllotaxy defects, characterized by irregular initiation and arrangement of
sepals, petals and stamens, and conspicuously enlarged carpels (fig. S9, A and B, E to L). In
particular, the *kan1-2 kan2-1* sepals from early developmental stages are strongly radialized with
little sepal lamina, and they fail to enclose the floral bud (fig. S9, F to L). These phenotypes are
distinct from those observed in *as2-7D* mutants and *pML1::AS2* in wild-type transgenics, which
show no such defects (Fig. 1 B, M to P and Fig. 2, A and B). While the mature *kan1-2 kan2-1*
sepals are curved downwards, indicative of strong adaxialization (predominantly inner identity),
we do not observe the strong folds and associated outgrowths on the outer epidermis of *kan1-2*
kan2-1 sepals unlike *as2-7D* (Fig. 1 E and F, fig. S9, C and D). However, as reported previously
[37,38,42,43,66], we do observe the folds and outgrowths on the outer epidermis of the *kan1-2 kan2-*1
leaves, similar to *AS2* constitutive overexpression phenotypes in the leaves (fig. S9, M to O). The
positions of outgrowths typically overlap folds (fig. S9O), which suggests a potential link
between buckling and ectopic outgrowth in the *kan1-2 kan2-1* mutant as well.
420
421   Overall, our investigations indicate that the outer epidermal buckling exhibited by *as2-7D* sepals
422   may not be associated with an obvious ectopic inner-outer polarity boundary, unlike outgrowths.
423   However, our investigations do not rule out the possibility that other polarity-associated genes
424   are disrupted in *as2-7D*.
425
426   **Buckling in *as2-7D* generates outgrowths via ectopic PIN convergence driving auxin**
427   **maxima formation**
428
429   In addition to the buckles, we observe the emergence of pointed outgrowths at the ends of the
430   buckles in the later stages in *as2-7D* sepals (Fig. 1F). While constitutive *AS2* overexpression has
431   previously been reported to cause outgrowths [39], what local signal triggers the formation of these
432   outgrowths at specific sites remains elusive.
433
The polar localization of PIN-FORMED 1 (PIN1) protein, an auxin efflux transporter, in the
plasma membrane on a particular side of the cell determines the directionality of auxin flow.
When PIN1 polarity from multiple neighboring cells converges to the same location, auxin

accumulates at that site, creating an auxin maximum [67]. It was shown that PIN1 convergence
sites form at the boundary between the *REV* and *KAN1* expression domains, which coincides
with the site of leaf initiation[61]. In the context of the outgrowths, Izhaki and colleagues first
showed that the site of outgrowth formation in *kan1 kan2 kan4* triple mutants coincide with the
site of PIN1 convergence, indicating that outgrowth formation involves PIN1 [59]. This was also
later confirmed by Abley and colleagues, who showed that PIN1 convergence sites form before
the outgrowth initiates in *kan1 kan2* mutant leaves [68]. Localized auxin import is essential for
outgrowth formation since Auxin importer genes *AUXIN RESISTANT 1* (*AUX1*) and *LIKE-AUX1*
(*LAX1*) are expressed at the tip of the outgrowths in *kan1 kan2* mutants, while the outgrowths
were severely reduced in *kan1 kan2 aux1 lax1 lax2 lax3* hextuple mutants [68]. Auxin biosynthesis
genes *YUCCA1* and *YUCCA4* were also expressed in the region surrounding the outgrowths.
Moreover, *CUP-SHAPED COTYLEDON2* (*CUC2*) guides the formation of PIN1 convergence
sites and localize auxin at the site of outgrowth formation in *kan1 kan2* leaves, indicating the
requirement of boundary genes in driving outgrowth formation[68].
451
To test whether outgrowths formed on the outer epidermis of *as2-7D* sepals also involve PIN1
convergence, we analyzed PIN1-GFP localization patterns in the wild type and *as2-7D* mutant.
In the inflorescence and early-stage flowers, the PIN1-GFP accumulation pattern in *as2-7D* is
similar to that of wild type (Fig. 7, A and B). However, in the later stages we observe PIN1-GFP
convergence specifically at the sites where the outgrowths form (Fig. 7, C and D, fig. S10, A and
B). In contrast, no PIN convergence sites are observed in the middle of wild-type sepals (Fig. 7,
C and D, fig. S10, A and B). We also observe that the auxin-responsive promoter DR5 is
specifically expressed where outgrowths are formed in *as2-7D* (fig. S10, C and D), which is
consistent with the fact that *PIN1* convergence creates local auxin peaks [69].
461
The polarity of PIN1 is mechano-sensitive in nature and responds to mechanical stress [70-72]. In
the later stages when *as2-7D* mutants exhibit both buckles and outgrowths, we observe PIN1
convergence specifically in the outgrowths and not the buckles (Fig. 7, C and D, fig. S10, A and
B). Since buckling inherently alters mechanical stress distributions [73,74], and buckling precedes
outgrowth formation, we hypothesized that buckling induced mechanical stress causes PIN1
convergence subsequently leading to outgrowth formation in *as2-7D*. To test our hypothesis, we
live imaged growing *pPIN1::PIN1-GFP as2-7D* sepals from initial stages prior to buckling,
through the stages where the outer epidermis buckles and outgrowths are evident (Fig. 8).
Initially, when the sepal outer epidermis uniformly curves outwards, no ectopic PIN1
convergence is observed on the sepal blade (Fig. 8, A to D). As the sepal grows, while buckle
formation occurs in the middle of the sepal blade, we do not observe PIN1 expression in the
region of the buckle (Fig. 8, E and F). Although outgrowth formation is not evident at this stage,
we do observe PIN1 convergence at the tip of the buckle, which marks the site of outgrowth
initiation (Fig. 8, E to H). In the following time-point, an outgrowth is evident in the region
where PIN1 convergence site was formed in the preceding stage (Fig. 8, I to L). As the
outgrowth progresses, the site of PIN1 convergence shifts towards the growing tip of the

outgrowth, while PIN1 expression continues to be absent in the region where the sepal buckles
(Fig. 8, M to P). Collectively, our observations suggest that the tissue buckles first and then PIN1
convergence occurs at the end of a buckle, which precedes the formation of the outgrowths. Our
data is consistent with the hypothesis that buckling induced mechanical stress may cause PIN1 to
converge and initiate an outgrowth.
483
We next tested whether PIN-mediated polar auxin transport is required specifically for the
formation of outgrowths and not buckles. We treated growing *as2-7D* inflorescences with polar
auxin transport inhibitor N-1-naphthylphthalamic acid (NPA) and compared the phenotype to
mock-treated samples. While *as2-7D* live-imaged in the presence of NPA exhibited
characteristic buckles, the formation of outgrowths was largely attenuated (Fig. 7, E and F, fig.
S10, E to H, and movie S4). Buckling continues to occur since the characteristic growth patterns
and misdirected sideways PDG_{max} tensors leading to buckling remain unaltered with NPA
treatment (Fig. 7, G to J). However, the localized fast growth leading to outgrowth formation
was largely reduced in the NPA-treated samples (Fig. 7, K and L). Our results suggest that polar
auxin transport is critical for the formation of outgrowth characterized by localized fast growth.
Overall, our results are consistent with our hypothesis that mechanical buckling causes ectopic
PIN1 convergence, generating local auxin maxima and fast growth, leading to outgrowth
formation.

DISCUSSION
498
While buckling is critical for generating complex organ shapes required for specialized
functional roles [1-6,16,18], buckling is not always advantageous, and needs to be circumvented to
generate smooth organs like sepals [75]. Using Arabidopsis sepals as the model, our study suggests
that unidirectional growth of both the outer and inner epidermal cells along the sepal longitudinal
axis, and similar tissue stiffness across the epidermal layers are key drivers of smooth sepal
formation in wild type. Conversely, the outer epidermis of *as2-7D* mutant buckles due to the
imbalance of growth and tissue stiffness across the outer and inner epidermal layers.
506
Overexpression of *AS2* leading to the formation of ectopic outgrowths has been documented by
previous studies [37,39,76]. However, we identify buckling mediated fold formation as an upstream
morphological deformity that occurs before the outgrowths are even evident, in the sepals. PIN1
convergence sites form specifically at the tip of buckles, marking the site of outgrowth initiation.
Since the Waites Hudson model in 1995 [57], the initiation of ectopic outgrowths has been
attributed to juxtaposition of abaxial-adaxial identities. We add a second mechanism through
which outgrowths can initiate via buckling, which generates auxin maxima via PIN1
convergence. We expect that auxin maxima trigger the gene regulatory networks involved in
outgrowth formation, that other studies have extensively characterized.
516

Although one would intuitively expect *AS2* ectopic expression to affect only the outer epidermis,
we find the growth rate and stiffness of the inner epidermis, where *AS2* is normally expressed, is
affected as well in *as2-7D*. Changes in inner epidermal growth properties in *as2-7D* could be
attributed to the fact that the outer and inner epidermis are connected via the mesophyll and at
the sepal margins. Altering growth patterns in the outer epidermis creates mechanical constraints
on the inner epidermis, thereby affecting its growth indirectly. Simply put, the outer epidermis of
as2-7D buckles as the sideways growth of the outer epidermis is limited by the inner epidermis.
Conceptually, this resembles trying to fit a cloth inside a box. If the cloth fits perfectly along the
edges, it remains smooth; otherwise, the cloth must wrinkle (buckle) to fit inside the box. Our
results show how the growth of an epidermal cell layer (or speculatively an animal epithelium [77])
relative to the tissue to which it is attached, drives organ shape.

While how molecular cues influence tissue mechanics has been relatively well studied [78-81], how
tissue mechanics inform molecular signals remain elusive. Our study suggests that not only does
AS2 influence tissue mechanics generating buckles, but also that buckles feed back into
molecular signals. We show that buckling is associated with convergence of the PIN1-auxin
efflux transporter. Our findings support the previous observations from multiple independent
studies that mechanical stress affect PIN1 polarity [70,72,82]. We find an association between the
tips of buckles and the establishment of PIN1 polarity convergence sites, causing the formation
of auxin maxima to subsequently trigger the initiation of epidermal outgrowths. There is a long-
standing debate as to whether primordial initiation in the SAM is also influenced by buckling, a
theory originally proposed by Paul Green [23]. However, PIN1 polarizes before primordia bumps
are visible, indicating that PIN1 can converge without buckling [83]. Our collective evidence
suggests that in certain contexts, buckling might be a causative factor, converging PIN1 and
generating outgrowths.

Limitations of the study

First, we used the *as2-7D* mutant as a tool to help us identify how aberrations in cell growth and
tissue properties generate buckles, as opposed to smooth sepal surfaces observed in the wild
type. However, *as2-7D* being a gain-of-function mutant does not inform much about the native
function of *AS2*. We have thus carefully refrained from extrapolating the native role of *AS2*
based on our findings.

Second, our numerical simulations are based on a cross-sectional 2D representation of a sepal,
since simulating the exact dynamics of a multilayered growing tissue in 3D is extremely
difficult. Hence, the Young's moduli values chosen to achieve outer epidermal buckling in the
numerical simulations are not comparable to the differences we observe experimentally. We used
the analytical model and numerical simulations as a guide for determining our experimental
trajectory, rather than a means to validate our findings.

557
Third, we used osmotic treatments to assess stiffness of the outer and inner epidermal layers,
mainly because the inner epidermis at early stages of sepal development is practically
inaccessible via dissection, without inevitably damaging the tissue[84]. As such, alternative
methods like AFM or indentation techniques [85] which require the tissue to be accessible, are
impractical for testing stiffness of the inner epidermis of young sepals. Such methods also
typically measure stiffness in highly localized regions, which again does not align with our
experimental purpose. Considering these factors, and the fact that osmotic treatments have
successfully been used to measure stiffness in multiple studies [36,55,56,86,87], we deemed osmotic
treatments as a suitable approach.
567
Fourth, there are currently very limited genetic tools available to orient growth and change tissue
mechanical parameters. Although a crude tool, we have chosen to over-expression *KRP1*, since
KRP1 is known to cause anisotropic growth [53], and is also unrelated to the core polarity
associated genetic network in which *AS2* is involved. However, genetic manipulations are not
entirely free from pleotropic phenotypic outcomes. While KRP1 inhibits cell division, the
evidence suggests it is not the inhibition of cell division that is suppressing buckling indirectly;
the *ML::KRP1 as2-7D* sepals still have dividing cells that could generate buckles and
outgrowths, but we do not observe these. Likewise, overexpression of *KRP1* could potentially
suppress buckling in *as2-7D* by accelerating differentiation; however, the live imaging shows
that the stomatal differentiation is not accelerated in *ML1::KRP1 as2-7D*. Overall, while the data
is consistent with our hypothesis, it may not be an absolute proof.
579
580 Finally, we acknowledge that we essentially provide negative data where we find no evidence
581 that the buckling is caused due to juxtaposed outer-inner identity. However, as a limitation, there
582 are of course a plethora of genes that *AS2* influences, that we could not test in this study.
583

ACKNOWLEDGEMENTS

We thank Isabella Burda, Frances Clark, Shuyao Kong, Byron Rusnak, Maura Zimmermann,
Erich Schwarz, Phoebe Roeder, Scott D. Emr and Stéphane Verger for their valuable comments
and suggestions on the manuscript. We thank Olivier Hamant for the *pPDF1::mCitrine-MBD*
microtubule marker plasmid. We thank John Paul Alvarez and John Bowman for sharing *kan1-2*
kan2-1/+ seeds. We thank Soren Strauss for his guidance with image processing. We thank
Mingyuan Zhu for his input on osmotic treatment experiments. We thank Trinh Duy Chi for his
input on imaging adult sepals. Finally, we thank Arabidopsis Biological Resource Center for
providing seed stocks for this work.

FUNDING

Research reported in this publication was supported by the National Institute of General Medical
Sciences of the National Institutes of Health under Award Number R01GM134037 to AHKR.
French National Research Agency (ANR, grant ANR-21-CE30-0039-01 GrowFlat, to AB). 2023
Sam and Nancy Fleming Research Fellowship to ASY. National Natural Science Foundation of
China grant no. 32270867 and Hundred-Talent Program of Zhejiang University to LH.
Cornell China Center and Zhejiang University for a joint seed fund grant to AHKR and LH.
IMB was supported by NSF REU Award (1850796).
602
The authors acknowledge the use of facilities and instrumentation supported by NSF through the
Cornell University Materials Research Science and Engineering Center DMR-1719875. The
content is solely the responsibility of the authors and does not necessarily represent the official
views of the National Institutes of Health or other funders.

AUTHOR CONTRIBUTIONS

Conceptualization: AHKR, ASY, LH. Experimentation: ASY, LH, PMK, XH, MH. Imaging:
ASY, LH, MH, PMK. Image processing: ASY, LH, PMK, IMB, AMG, RSS. Data analysis:
ASY, LH, AK. Analytical modeling: AK, AB. Simulations: AK, MP, AB. Funding acquisition:
AHKR, AB, LH. Project administration: AHKR, AB. Supervision: AHKR, AB, LH. Writing –
original draft: ASY, LH, AHKR. Writing – review & editing: ASY, LH, PMK, AK, XH, IMB,
MH, AMG, RSS, AB and AHKR.

COMPETING INTERESTS

Authors declare that they have no competing interests.

RESOURCE AVAILABILITY
The raw datasets used for making the graphs are provided in Supplementary Data S1. Primers
used for this study are available in Table S1 in Supplementary materials. The transgenic lines for
live imaging purposes generated in this study have been deposited with the Arabidopsis
Biological Resource Center (ABRC), associated with the following accession numbers: CS73559
(pLH13 L*er*-0), CS73560 (pLH13 *as2-7D*), CS73561 (pLH13 *pML1::KRP1* L*er*-0) and
CS73562 (pLH13 *pML1::KRP1 as2-7D*). The meshes and images are publicly available at osf.io
(https://osf.io/p5q39/). The R codes used for generating the plots are available at Github
(https://github.com/Avilash1993/R_codes_for_plots_as2-7D_sepal_buckling) under GNU
General Public License v3.0. Numerical model implementation and the codes for exploration of
the parameter space are available under LGPL license at
https://gitlab.inria.fr/mosaic/publications/growth_buckling.

FIGURE TITLES AND LEGENDS

634
635

Fig. 1. The ectopic expression of *ASYMMETRIC LEAVES 2* (*AS2*) on the outer epidermal layer disrupts sepal smoothness in *as2-7D* mutant.

(**A** and **B**) Whole inflorescences of (**A**) Wild type (WT, L*er*-0) and **B**) *as2-7D* mutant (in L*er*-0 background). Yellow arrowheads indicate outgrowth. Scalebar, 1 cm; representative images of *n* = 3 inflorescences for both genotypes.

(**C** and **D**) Mature flowers (stage 14 [88]) of (**C**) WT and (**D**) *as2-7D* mutant. WT sepals (green leaf-like organs) are marked with the letter "S" in white (C). Flowers are enclosed within four sepals. Note the outgrowths (yellow arrowheads) on the outer epidermis of *as2-7D* sepals (D). Scale bar, 1 cm; representative images of *n* = 3 flowers.

(**E** and **F**) Scanning electron microscopy (SEM) images of the outer epidermis of (**E**) WT and
(**F**) *as2-7D* mutant sepals. Note the folds (*) and outgrowths (yellow arrowheads) on the *as2-7D*
outer epidermis. Scale bar, 100 µm; representative images of *n* = 5 for WT and *n* = 10 for *as2-*
7D.
(**G** and **H**) Confocal microscopy images of the inner epidermis of (**G**) WT and (**H**) *as2-7D*
mutant. The images are representative of T2s harboring pLH13 (*35S::mCitrine-RCI2A*), a yellow
fluorescent protein-based cell membrane marker [89]. Scale bar, 100 µm; representative images of
n = 3 for both genotypes.
(**I**) Schematic diagram representing the genetic mutation in *as2-7D*. Gene body of *AS2* showing
the G-to-A point mutation (red arrow) at position 1484 bps upstream of ATG in *as2-7D* (Top
half). The *AS2* expression (red dots) domain in the WT and *as2-7D* sepals is shown (Bottom
half).
(**J**) Anatomical diagram representing (Top) an early stage 6 flower and (Bottom) an optical
longitudinal section through the flower. The letter "S" in black represents sepals. The outer and
inner epidermal layers of the outer sepal are marked.
(**K** and **L**) Longitudinal section of stage 6 flowers of WT. Cell walls are stained with propidium-
iodide (PI, greyscale). Images show the GFP expression pattern (green) driven by (**K**) WT *AS2*
promoter (*pAS2^{WT}*) and (**L**) *AS2* promoter from *as2-7D* (*pAS2^{as2-7D}*). Arrows indicate the outer
and inner epidermal layers of the outer sepal. Scale bar = 20 µm; *n* = 3.
(**M**) Whole inflorescence of a representative transgenic (T1) harboring *pML1::AS2* (*AS2*
expression driven by *ATML1* epidermal specific promoter) transgene in WT. Yellow arrows
mark the outgrowths on the sepal epidermis. Independent T1s assessed = 28, all of which show
similar levels of outgrowth formation. Representative image of *n* = 3 inflorescences.
(**N** to **P**) Confocal images of PI-stained (greyscale) inflorescences of (**N**) WT, (**O**) representative
transgenic (T1) harboring *pML1::AS2* transgene in WT, and (**P**) *as2-7D* mutant. WT outer
epidermis is smooth, whereas *pML1::AS2* exhibits outgrowths (yellow arrowheads) like *as2-7D*.
Scale bar, 50 µm; *n* = 3 inflorescences per genotype.

Fig. 2. The *as2-7D* sepal outer epidermal morphology transitions from smooth to buckles
and subsequently outgrowths.
(**A** and **B**) Representative live-imaging series of (**A**) a wild-type (WT) flower and (**B**) an *as2-7D*
mutant flower starting from developmental stage 4 over a course of 96 hours, imaged at 24-hour
intervals ($n = 3$). Images capture morphological changes of the outer sepal. Both genotypes
harbor the pLH13 (*35S::mCitrine-RCI2A*) cell membrane marker (greyscale). Trichomes were
removed from the outer sepal epidermis by image processing to better visualization the sepal
morphology. Scale bar, 50 µm.
(**C**) Tilted view of *as2-7D* mutant flower at the 48-hour time point. Note the appearance of the
out-of-plane protrusion (ridge) in the middle of the sepal, circled in white. Scale bar, 50 µm.
(**D**) The *as2-7D* mutant flower at the 96-hour time point. Note the dramatic increase in the
number and extent of ridges (red circles) and invaginations (white circles), characteristic of
buckles, on the outer sepal epidermis. Yellow arrowheads indicate outgrowths. Scale bar, 50 µm.
(**E** and **F**) Cell area growth rate over a 48-hour period for the (**E**) WT and (**F**) *as2-7D* mutant
sepals. Growth rate is calculated as the ratio of cell lineage area at the later time point divided by
that of the earlier time point. The heatmap is projected onto the later time point. Red to blue
gradient shows fast to slow growth. The growth rate over 0-48 hours (left) and 48-96 hours
(right) is shown. Arrowheads point to the outgrowths. Scale bar, 50 µm.
(**G** and **H**) Cell division rate over a 48-hour period for the (**G**) WT and (**H**) *as2-7D*. Cell division
rate is calculated based on the number of daughter cells per lineage, represented as a heatmap
projected onto the later time point. Red to blue gradient shows more to less division. The cell
lineages were tracked using MorphoGraphX software [90]. Segmented cells are outlined in black.
Representative heatmaps for one of the three replicates ($n = 3$) for WT and *as2-7D* are shown.
Scale bar, 50 µm.
(**I** to **K**) Simulations based on the analytical model (see Materials and methods), providing visual
illustrations of the conditions that may favor buckling of the outer epidermis. The simulations
illustrate change in tissue morphology along the cross-section when the outer epidermis was
allowed to grow in width over two time points. The elastic modulus (E) of the outer epidermis
(E_o), inner epidermis (E_i), mesophyll (E_m) and edges (E_{end}) are shown. Initial growth tensor along
the width for all the four components were set to 1, with the outer epidermal growth rate set to
0.1, while the growth rates of the other components was set to 0. (**I**) Simulation outcomes when
the outer and inner epidermis have similar stiffness (E_o and $E_i = 10^2$), causing the tissue to curve.
(**J**) When the stiffness of the outer epidermis ($E_o = 10^2$) is less than that of the inner epidermis
($E_o = 10^6$), the outer epidermis buckles, while the inner epidermis remains unaffected. (**K**)
Simulation outcomes wherein the stiffness of the outer epidermis ($E_o = 10^6$) is more than that of
the inner epidermis ($E_o = 10^2$), leading to a slight increase in tissue curvature over time.

709
710

Fig. 3. Unidirectional growth on both epidermal layers is essential for smooth sepal development.

(**A** and **B**) Representative confocal images of the (**A**) Wild type (WT) and (**B**) *as2-7D* mutant live-imaged with a high laser intensity (~16-20%) every 24 hours. Both the genotypes harbor pLH13 (*35S::mCitrine-RCI2A*) cell membrane marker (green). Scale bar, 20 µm. One of the three representative replicates (*n* = 3) is shown.

(**C** and **D**) Segmented cells in grey superimposed onto the plasma membrane signal (also in grey) projected on mesh surfaces generated using MorphoGraphX [90]. Mesh surfaces for 0 and 48-hour time points of the outer (left) and inner (right) epidermis of the (**C**) WT and (**D**) *as2-7D* mutant are shown. Note the strikingly reduced number of cells on the inner versus outer surface at the 0-hour time point. Scale bar, 20 µm; *n* = 3.

(**E** and **F**) Cell growth rates over a 48-hour period for the (**E**) WT and (**F**) *as2-7D* mutants are shown. The heatmap has been projected onto the 48-hour time-point surface. In each panel, the growth rate of cells on the outer surface (left) and inner surface (right) is shown. Red to blue heatmap gradient shows fast to slow growth. Scale bar, 20 µm; *n* = 3.

(**G** and **H**) Cell division rates over a 48-hour period for the (**G**) WT and (**H**) *as2-7D* mutant on the outer surface (left) and inner surface (right) are shown. Segmented cells are outlined in black. Red to blue gradient shows more to less/no cell division respectively. Scale bar, 20 µm; *n* = 3.

(**I** and **J**) Principal directions of maximal growth (PDG$_{max}$, yellow) of the cells on the outer (left) and inner (right) epidermis of (**K**) WT and (**L**) *as2-7D*. Segmented cells in grey are superimposed on the plasma membrane signal (also in grey). Length of any given PDG$_{max}$ tensor is directly proportional to the cell growth rate in that direction. A blue arrow indicates the longitudinal sepal axis. Note the misoriented PDG$_{max}$ tensors on the *as2-7D* outer epidermis. Scale bar, 20 µm; *n* = 3.

(**K**) A schematic diagram depicting the calculation of the angle of deviation between PDG$_{max}$ tensor and the longitudinal axis to obtain a quantitative measure of the deviation.

(**L** and **M**) Heatmaps show the angle of deviation of the PDG$_{max}$ tensor from the sepal longitudinal axis for the outer (left) and inner (right) epidermal cells of the (**L**) WT and (**M**) *as2-7D* mutant. Scale for the angle of deviation ranges from 90º (highly deviated, red) to 0º (no deviation, blue). Scale bar, 20 µm; *n* = 3.

(**N** and **O**) Cortical microtubule orientation (greyscale) from comparable regions of stage 6 flower outer epidermis for the (**I**) WT and (**J**) *as2-7D* mutant is shown. These plants harbor *pPDF1::mCitrine-MBD*, a YFP based microtubule marker [91]. . Cells segmented using meshes generated from PI-stained images are outlined in gray. White arrows point towards the longitudinal axis of the sepals. Microtubules in the WT are oriented perpendicular to the sepal longitudinal axis, whereas, in *as2-7D*, the microtubules are misoriented. Yellow arrowheads point towards the misoriented microtubules. Scale bar, 10 µm; *n* = 3.

Fig. 4. Slower growth on the inner epidermis leads to overgrowth of the outer epidermis in *as2-7D*.

(**A** to **D**) Heatmaps represent growth rates of (**A**) Wild-type (WT) outer, (**B**) *as2-7D* outer, (**C**) WT inner, and (**D**) *as2-7D* inner epidermal surface at every 24-hour interval, over a 48-hour live imaging period. Growth rate is projected onto the later time point. Growth rate for the region demarcated in cyan at 0-hour time point was tracked over time. Heatmap: red, fast growth, and blue, slow growth. Representative heatmaps for $n = 3$ biological replicates are shown. Scalebar, 20 µm.

(**E**) Graph shows epidermal surface growth rates every 24 hours for the WT (black) and *as2-7D* (red) is shown ($n = 3$). Lowercase letters on top of the data points represent significant differences based on one-way ANOVA with Tukey's HSD post-hoc test ($p < 0.05$, similar letters represent no significant difference).

763
Fig. 5. Aligning growth along the longitudinal direction and increasing outer epidermal stiffness is sufficient to rescue buckling in *as2-7D*.

(**A** to **D**) Representative stage 14 flowers of (**A**) Wild type (WT), (**B**) *as2-7D* mutant, (**C**) *AtML1::KRP1,* and (**D**) *AtML1::KRP1 as2-7D*. The WT sepals are marked with the letter "S" in

white (**A**). Yellow arrowheads point to outgrowths in *as2-7D* (**B**). Note the smooth sepal of

AtML1::KRP1 as2-7D compared to *as2-7D*. Scale bar, 1 cm; $n = 3$ flowers for all genotypes.

(**E** to **H**) Confocal images of the outer epidermis of (**E**) WT, (**F**) *as2-7D* mutant, (**G**)

AtML1::KRP1, and (**H**) *AtML1::KRP1 as2-7D* stage 14 sepals. The sepals were stained with PI

(grey, cell walls). Note the increased number of highly elongated giant cells in *AtML1::KRP1*

and *AtML1::KRP1 as2-7D* sepal, and the smooth outer epidermis of *AtML1::KRP1 as2-7D* sepal

compared to the highly rugose morphology of *as2-7D*. Scale bar, 200 µm; $n = 3$.

(**I** to **L**) Magnified insets of the (**I**) WT, (**J**) *as2-7D* mutant, (**K**) *AtML1::KRP1,* and (**L**)

AtML1::KRP1 as2-7D sepal for the regions marked by white boxes in panels E-H respectively.

Asterisks denote a typical giant cell. Scale bar, 200 µm; $n = 3$.

(**M** to **P**) The principal direction of growth (PDG$_{max}$) for the outer epidermal cells of the (**M**)

WT, (**N**) *as2-7D,* (**O**) *AtML1::KRP1,* and (**P**) *AtML1::KRP1 as2-7D* sepals. Yellow lines mark

the PDG$_{max}$ tensors calculated over a 48-hour period, projected onto the later time point. For WT

and *as2-7D*, the flower images used are the same as in Figure 2. Segmented cells in grey are

superimposed onto the plasma membrane signal (also in grey). Scale bar, 20 µm; $n = 3$.

(**Q** and **R**) Schematic diagrams demonstrating the concept of cell wall stiffness assessment by

osmotic treatment assays [55]. (**Q**) A typical plant epidermal cell before osmotic treatment. The cell

wall and cell membrane are shown in black and green respectively. (**R**) A cell that has undergone

plasmolysis in a hyperosmotic solution (0.35 M NaCl).

(**S** and **T**) Anatomical diagrams demonstrate the use of osmotic treatments to estimate epidermal

stiffness. Longitudinal sections of an early stage 6 flower (**S**) before and (**T**) after osmotic

treatment are shown. The outer (green) and inner (blue) sepal epidermis are marked. Arrowheads

(red) demonstrate aerial shrinkage.

(**U** to **X**) Heatmaps show the aerial shrinkage (in percentage) of a longitudinal sepal cross-section

for the inner and outer epidermis of (**U**) WT, (**V**) *as2-7D* mutant, (**W**) *AtML1::KRP1* and (**X**)

AtML1::KRP1 as2-7D, post osmotic treatment in 0.35M NaCl solution. Shrinkage was computed

as the decrease in area of a comparable cross-sectional region from an early stage 6 flower before

and after osmotic treatment. Heatmaps are projected on the post-treatment surface. More

shrinkage is represented in red and less in blue. Scale bar, 20 µm. Heatmaps for each genotype

are representative of 5 independent replicates ($n = 5$).

(**Y**) Graph showing the percentage aerial shrinkage post osmotic treatment. Data across 5

independent replicates each for the WT (black dots), *as2-7D* (red dots), *AtML1::KRP1* (grey

dots), and *AtML1::KRP1 as2-7D* (pink dots) for both the outer and inner epidermis are shown.

The letters above the plots represent significant differences based on one-way analysis of

variance (ANOVA) with Tukey's honest significant difference (HSD) post hoc correction ($p <$

0.05). Similar letters indicate no significant difference.

Fig. 6. Buckling in *as2-7D* is not caused due to disruption of outer (abaxial) – inner
(adaxial) polarity.
(**A** to **D**) REVOLUTA (REV) protein reporter (*proREV::REV-mGFP*) shows nuclear localization
of REV, restricted exclusively to the inner epidermis of the wild type (WT) and *as2-7D* sepals.
(**A** and **B**) Confocal images of inflorescence meristems show adaxial specific REV expression
(yellow arrowheads) in the young sepals of (**A**) WT and (**B**) *as2-7D* mutant. (**C** and **D**)
Representative middle stage (Stage 8-9) flowers of (**C**) WT and (**D**) *as2-7D* show no detectable
REV expression on the outer epidermis of the sepals in the folds or outgrowths. Images of
representative T3s (*n* = 3) show mGFP in green and chlorophyll in greyscale. Scalebar, 50 µm.
(**E** to **H**) *FILAMENTOUS FLOWER* (*FIL*) promoter reporter (*proFIL::mCherry-RCI2A*)
expression pattern is restricted to the outer sepal epidermis and shows no difference between the

WT and *as2-7D* mutant. (**E** and **F**) Confocal images of inflorescence meristems show *FIL*
expression pattern in the early floral developmental stages of (**E**) WT and (**F**) *as2-7D* mutant.
The membrane localized mCherry is shown in green and chlorophyll in greyscale. (**G** and **H**)
Representative middle stage (Stage 7-8) flowers of (**G**) WT and (**H**) *as2-7D* mutant showing the
FIL promoter reporter signal (mCherry, green). Images are representative of three T3s ($n = 3$).
Scalebar, 50 µm.
(**I** to **L**) Whole inflorescence images of (**I**) WT, (**J**) *rev-6* mutant, (**K**) a representative transgenic
(T1) harboring *pML1::AS2* transgene in *rev-6* and (**L**) *as2-7D* mutant. Note the outgrowths
(yellow arrows) on the sepal outer epidermis of *pML1::AS2 rev-6* are similar to those of *as2-7D*
and *pML1::AS2* in WT (Fig. 1M). Multiple independent T1s for *pML1::AS2* in *rev-6* were
assayed ($n = 19$), all of which show similar outgrowth formation on the sepal outer epidermis in
the *rev-6* background.
833

Fig. 7. Mechanical buckling drives ectopic PIN convergence creating auxin maxima, causing fast local growth and subsequent outgrowth formation.

(**A** to **D**) Confocal stack images showing the expression pattern of auxin efflux carrier protein PIN-FORMED 1 (PIN1) in the wild type (WT) and *as2-7D*. (**A** and **B**) Expression pattern of PIN1 protein reporter (*pPIN1::PIN1-GFP* [92]) in the inflorescence meristem of (**A**) the WT, (**B**) *as2-7D* mutant. The *PIN1* expression pattern in *as2-7D* is identical to WT in the young floral development stages, wherein PIN1 polarity is directed towards the insipient primordia and growing sepal edges, as described previously [92]. Scalebar, 50 µm. (**C** and **D**) Expression pattern of *PIN1* in a representative middle stage (stage 8) flower of the (**C**) WT and (**D**) *as2-7D* mutant. Note the expression of *PIN1* at the growing tip of the outgrowths (yellow arrows) in *as2-7D*, whereas no *PIN1* expression is detected in the blade region where the epidermis buckles. In panels A to D, the images are representative of three *pPIN1::PIN1-GFP* and *pPIN1::PIN1-GFP as2-7D* plants (*n* = 3). PI is shown in greyscale and GFP in green. Trichomes were removed from the outer sepal epidermis by image processing. Scale bar, 50 µm.

(**E** to **F**) Representative live-imaging series of (**E**) Mock treated *as2-7D* and (**F**) 10µM NPA treated *as2-7D* developing flowers over 96 hours, imaged at every 24-hour interval (representative of *n* = 3 flowers). Note the development of outgrowths from 72-hour time-point onwards (yellow arrowheads) in the mock-treated *as2-7D*, which are absent in NPA treated *as2-7D* flowers. Plants harbor pLH13 (*35S::mCitrine-RCI2A*) cell membrane marker (greyscale). Scale bar, 50 µm.

(**G** to **H**) The PDG_{max} tensors (yellow) of the cells on the outer epidermis of (**G**) mock-treated and (**H**) 10µM NPA treated *as2-7D* over the first 48-hour growth period for the representative samples shown in panels E to F. Characteristic misoriented PDG_{max} tensors (relative to the longitudinal sepal axis) on the outer epidermis of *as2-7D* remain unaffected in the NPA-treated samples, thus driving buckling in later stages. Scale bar, 50 µm.

(**I** to **L**) Outer epidermal cell growth rates over (**I** and **J**) 0–48-hour growth period and (**K** and **L**) 48–96-hour growth period of the mock and 10µM NPA treated *as2-7D* flowers respectively. Heatmaps are projected onto the later time point. Fast (red) to slow (blue) growth rates are shown. In panel K, note the fast cell growth driving outgrowth formation (yellow arrowheads) which is absent in NPA treated sample (panel L). Scale bar, 50 µm.

Fig. 8. PIN1-GFP converges at the end of a buckle where an outgrowth subsequently forms in *as2-7D* mutants.

Confocal live imaging series of a *pPIN1::PIN1-GFP as2-7D* flower, imaged every 12 hours starting from stage 3 (representative of *n = 3*). PIN1-GFP (green) and chlorophyll autofluorescence (gray) are shown. We analyzed the curvature (minimal curvature), in which warm colors denote out-of-plane (bulging) and cool colors represent inward (saddling) deformations. Scalebars, 20 µm.

(**A** to **D**) Confocal image of *pPIN1::PIN1-GFP as2-7D* stage 4 flower at 24-hour time point (**A**),
and the analysis of its minimal curvature (**B**) is shown. At this stage, the sepal exhibits uniform
outward curvature. The inset (**C**) and the cross-section through the inset (**D**) for the region
highlighted in panel A (yellow box) depicts the site where an outgrowth will form in the later
stages. Note the absence of *PIN1-GFP* expression at this site (white arrow).
(**E** to **H**) The *pPIN1::PIN1-GFP as2-7D* flower at the 36-hour time point as it transitions into
stages 6 (**E**), and its minimal curvature (**F**) is shown. At this stage, the initial buckle (marked by
a white circle) appears as an out-of-plane fold in the center of the sepal blade (deep red on the
curvature). No *PIN1* expression is observed at one end of the buckle. The inset (**G**) and the
cross-section through the inset (**H**) for the region highlighted in panel **E** (yellow box) is shown.
As the outer layer buckles, *PIN1* expression begins to converge in the highlighted region (white
arrow), marking the site of the later outgrowth.
(**I** to **L**) The *pPIN1::PIN1-GFP as2-7D* flower at the 48-hour time point (**I**), and its minimal
curvature (**J**) is shown. At this stage, the outgrowth begins to form (yellow arrow pointing
towards the red bulge) at the end of the buckle. The inset (**K**) and its corresponding cross-section
(**L**) for the region highlighted in panel E (yellow box) shows the *PIN1* expression associated
with the insipient outgrowth (white arrow).
(**M** to **P**) The *pPIN1::PIN1-GFP as2-7D* flower at the 84-hour time point as it transitions into
stages 9 (**M**), and its minimal curvature (**N**) is shown. The inset (**O**) and the cross-section
through the inset (**P**) for the region highlighted in panel **M** (yellow box) where the outgrowth has
already formed, is shown. By this stage, *PIN1* expression (white arrow) converges towards the
growing tip of the outgrowth.

MATERIALS AND METHODS

Plant Materials

Arabidopsis thaliana genotypes used in this study are in Landsberg *erecta* (L*er*) background,

which is referred to as the wild type, unless otherwise mentioned. The mutants *rev-6* [64] and *as2-*

5D mutant (in Col-0 background) [41] have been previously described, and were obtained from

Arabidopsis Biological Resource Center (ABRC). The *kan1-2 kan2-1* mutant previously

described [93] was a kind gift from John Alvarez (Monash University). The plants are homozygous

for *kan1-2* while *kan2-1* is segregating; the phenotypes were assayed for plants that are

homozygous for both *kan1-2* and *kan2-1*. The *as2-7D* mutant in the L*er*-0 background was

identified from ethyl methanesulfonate (EMS) based forward genetic screen. Briefly, M2 seeds

(second generation) obtained from self-fertilized EMS mutagenized L*er* plants were purchased

from Lehle Seeds [94]. M2 plants were screened under a stereo microscope for sepal defects [53,95].

The *as2-7D* mutant was isolated based on the lumpy sepal phenotype. A mapping population

was generated by crossing the *as2-7D* mutant with a Columbia (Col-0) accession plant. Standard

map-based cloning techniques were used to clone the underlying mutation [96]. The *as2-7D*

mutation is a G to A change at base 1,484 upstream of the ATG translational start site of

AT1G65620/*AS2*. To reduce background mutations, the *as2-7D* mutant was backcrossed twice to

wild-type L*er* plants twice before further analysis. In the backcrosses, we found that *as2-7D*

mutant behaves as a dominant allele as the F1s exhibit the *as2-7D* phenotype, although it is not

as strong. The *as2-7D* mutation can be PCR-genotyped by amplifying with primer pairs oLH108

and oLH324 at 55°C annealing temperature, followed by digesting the product with EcoRI to

produce an 82-bp wild-type product or a 111-bp mutant product. Primers are listed in

Supplementary Table S1.

Plant Growth

Plants were grown in Percival (Percival-Scientific) growth chambers programmed for long day

conditions (16-hour light/8-hour dark cycles). Growth temperature (22ºC), relative humidity

(65%) and fluorescence intensity (~100 µmol m^{-1}s^{-1}) were maintained.

For soil-based experiments, plants were grown on soil premixed with Vermaculite (LM111). For

stratification, the trays were placed in a 4 ºC cold room for three days prior to moving them to

the growth chambers. For BASTA (glufosinate ammonium) selection, the soil was autoclaved

twice (Program 3- 15 min sterilization, 30 min dry, Primus ST-1 Autoclave (Model PSS5-B-

MSSD)), and sterilized seeds were planted at a density of 50-100 seeds per pot. For live-imaging

and adult-stage phenotyping experiments, one plant was grown per pot (6.5×8.5 inches) to

facilitate healthy vegetative growth.

For plate-based experiments, seeds were washed twice with a sterilization solution containing
70% ethanol + 0.1% Triton X-100, replacing the solution after each wash. Two final rounds of
washes were provided with 100% ethanol. Seeds were spread on filter paper (Whatman 1,
diameter 90 mm) and allowed to dry inside a sterile safety cabinet (Class II, Type A2, Labgard).
Dried seeds were evenly distributed on freshly prepared MS plates (1/2 MS basal salt mixture,
0.05% MES w/v, 1% sucrose w/v, pH ~ 5.8 (adjusted using KOH) containing 0.8% phytoagar.
The plates were sealed with 3M micropore tape, wrapped in aluminum foil, and stratified for
three days at 4 ºC before being moved to the growth chamber.
964
965 **Phenotyping whole inflorescences, flowers, and plants**
966
967 Whole inflorescence and stage 14 flower images were taken either using Canon PowerShot A640
968 10MP Digital camera attached to Zeiss Stemi 2000-C stereo microscope or Accu-Scope Excelis
969 4K UHD camera attached to Zeiss Stemi 508 stereo microscope. For inflorescence images, and
970 Stage 14 flowers an optical zoom of 1.0 and 1.6 were used respectively. Whole plant images
971 were captured using iPhone XR with a 26 mm focal length and aperture of f/1.8.
972
973 **Schematic diagrams**
974
975 The schematic diagrams shown in Fig. 1I and Fig. 4 Q to T were hand-drawn using the
976 Notability application in Apple iPad (model A1701).
977
978 **Genetic crosses**
979
To generate *pATML1::KRP1 as2-7D*, the *pATML1::KRP1* in L*er* background [53,54] were crossed
into *as2-7D* mutants. Segregation analysis and confocal microscopy was used to find
pATML1::KRP1 as2-7D plants. The *pPIN1::PIN1-GFP DR5::VENUS* [92] plants and
pPIN1::PIN1-GFP pin1-4 [92] plants were crossed into *as2-7D* mutants. The *pPIN1::PIN1-GFP*
pin1-4 plants (control) exhibit a mild phyllotaxy defect. Genetic segregation analysis and
confocal microscopy was used to find *DR5::VENUS as2-7D* and *pPIN1::PIN1-GFP as2-7D*
plants.
987
988 **Cloning overexpression constructs and reporters**
989 The pLH13 (*p35S::mCitrine-RCI2A,* hygromycin resistance) and pAR169 (*pATML1::RCI2A-*
990 *mCtrine*, basta resistance) plasma membrane markers have been described previously [53,97]. The
991 *pPDF1::mCitrine-MBD* microtubule marker [91] was a kind gift from Olivier Hamant (RDP
992 institute, ENS de Lyon, France).
993
994 For cloning *pML1::AS2*, the AT1G65620/*AS2* coding sequence (600 bps) was amplified from
995 Col-0 cDNA using primers oLH111 and oLH112 by Phusion PCR (NEB) and cloned into

996　pENTR™/D-TOPO™ (Invitrogen™). The *AS2* coding sequence was cloned into the *pATML1*
997　containing gateway vector pAR176 (Basta resistance [53]) by LR reaction.
998
999　For *pML1::REV*, the AT5G60690/*REV* full length coding sequence spanning 2,529 bp was
1000　amplified from Col-0 inflorescence cDNA using primer pairs oAS_005 and oAS_006. The
1001　resulting PCR product was subcloned into pCR™8TOPO™ vector (Invitrogen™) by TA
1002　cloning. LR reaction was used to clone the *REV* coding sequence into pAR176.
1003
To clone *pAS2 WT::GFP-GUS*, the 4.1 kb upstream promoter region and part of the first exon of
AS2 was PCR amplified from L*er*-0 genomic DNA using primer pairs oLH347 and oLH370 and
cloned into pENTR™/D-TOPO™ vector (Invitrogen™). The resultant vector was LR
recombined into the gateway vector pBGWFS7 [98]. For cloning *pAS2^as2-7D^::GFP-GUS,* the 4.1 kb
upstream promoter region and part of the first exon of *AS2* was PCR amplified from *as2-7D*
genomic DNA, and the subsequent cloning steps were similar to cloning *pAS2::GFP-GUS.*
1010
1011　 For *proREV::REV-GFP*, the 7.75 kb *proREV::REV* fragment (amplified from L*er*-0) and 1.08
1012　kb mGFP-T35S fragment amplified from *pML1::H2B-mGFP* (pAR180 [53]) were assembled into
1013　the PmeI site of pMOA33 (Kanamycin resistance).
1014
For cloning *proFIL::mCherry-RCI2A*, a 3.92 kb *FIL* promoter fragment was amplified from L*er*-
0. This fragment was assembled with a 2.19 kb *omega-mCherry-RCI2A-T35S* fragment
amplified from *p35S::mCherry-RCI2A* (pMZ11 [99]) into the SacI and SpeI site of pMOA34
destination vector (hygromycin plant resistance, [100]). All the intermediate and final constructs
were verified for sequence and directionality by sequencing. Primers are listed in Supplementary
Table S1.
1021
1022　**Plant transformation and selection**
1023
The binary vector constructs were transformed into *Agrobacterium tumefaciens* GV3101 strain
(Rifampicin and Gentamycin resistant) by triparental mating method [101]. Briefly, two *E.coli*
strains, one containing the binary vector and the other containing the helper (conjugative)
plasmid, and *Agrobacterium* strain GV3101 were mated together. The GV3101 clones were
streaked onto Rifampicin + Gentamycin + binary vector specific antibiotic plates for selection,
and positive clones were isolated by colony PCR. The T-DNAs were transformed into plants by
Agrobacterium-mediated floral dipping. Transformants were selected either on ½ MS media
plates (for Hygromycin and Kanamycin selection) or on soil (for Basta selection). For selecting
pLH13 in wild type and *as2-7D*, *proFIL::mCherry-RCI2A* in wild type and *as2-7D* T1s, a final
concentration of 25 -30 µg/mL Hygromycin was used. For selecting *proREV::REV-mGFP* in
wild type and *as2-7D* T1s, a final concentration of 50 µg/mL Kanamycin was used. Healthy,
non-chlorotic plants were transplanted to the soil post two weeks of growth. For selecting

pML1::AS2 in wild type and in *rev-6*, *pML1::REV* in wild type and in *as2-7D*, *pAS2^WT^::GFP-*
GUS and *pAS2^as2-7D^::GFP-GUS* in wild type, *pPDF1::mCitrine-MBD* in wild type and in *as2-7D*
T1s, BASTA selection was performed on soil grown plants. Seedlings aged 1 to 2 weeks were
selected with 100 µg/mL BASTA solution by spraying, and non-chlorotic plants were
transplanted to fresh pots after a week. At least twenty independent T1 plants were characterized
for each construct. For fluorescent reporter lines, T1 plant was chosen for each line based on
fluorescence. Confocal microscopy was used to assess the stability of fluorescence in every
generation.
1044
1045 **Flower staging**
1046
1047 Floral staging was performed based on Smyth et al [88]. Briefly, sepal primordia barely arise in
1048 Stage 3 and overlie the floral meristem in Stage 4. Sepals enclose the floral bud completely by
1049 late stage 6. Flowers beyond stage 6 were staged based on size and morphology [35].
1050
1051 **Sepal selected for microscopy**
1052
For flower imaging across experiments (live imaging and single time point reporter imaging), we
focused primarily on the outer sepal of the flower, unless specified. Briefly, flowers are enclosed
within four sepals, wherein the foremost sepal, which faces away from the meristem is denoted
as the outer sepal, whereas the sepal that is diametrically opposite to the outer sepal and facing
towards the meristem is denoted as the inner sepal. The remaining two sepals positioned on
either side are the lateral sepals. For a detailed description, please refer to Roeder et. al [35].
1059
1060 **Scanning electron microscopy (SEM)**
1061
1062 Scanning electron microscopy was performed as described previously [53]. Briefly, Stage 14
1063 flowers were fixed in an FAA solution (50% ethanol, 5% acetic acid, and 3.7% formaldehyde)
1064 for 4 hours, dehydrated using an ethanol series, and critical point dried. Sepals were dissected,
1065 sputter-coated with platinum palladium and then imaged using a LEICA 440 scanning electron
1066 microscope.
1067
1068 **Confocal Microscopy**
1069
Live imaging
For live imaging sepal development, primary inflorescences from six- to eight-week-old plants
containing the *p35S::mCitrine-RCI2A* membrane marker (pLH13 in wild type, *as2-7D*,
pML1::KRP1, and *pML1::KRP1as2-7D* T2s) were excised. Dissection was performed as
described previously [99], with modifications. Briefly, siliques were removed using surgical
scissors (Artman Instruments). Flowers older than stage 7 were dissected using Dumont tweezers

(Electron Microscopy Sciences, Style 5) under a stereo microscope (ZEISS Stemi 2000).
Dissected inflorescences were inserted upright into small petri-plates containing live imaging
media (1/2 MS media supplemented with 1X Gamborg Vitamin mix and 1000-fold diluted plant
preservative mixture) with 1.2% w/v phytoagar. The plates were filled with autoclaved water and
allowed to stand for two-three hours for the dissected inflorescence to rehydrate. Residual buds
were dissected down to Stage 3-4 in water using a needle (BD Micro-Fine[TM] IV Insulin
Syringes). Inflorescences were moved to fresh media plates and transferred to the growth
chamber for at least 24 hours for recovery from dissection, prior to imaging. Stage 4 flowers
were selected as the 0- hour time point across all live imaging series.
1085
Post recovery, dissected tissues for live imaging were positioned at an angle to ensure the outer
sepal of the chosen stage 4 flower directly faces the objective of the confocal microscope. The
flowers were submerged in MillQ water containing 0.01% v/v SILWET L-77 surfactant and
imaged using Zeiss 710 upright confocal microscope using 20X water dipping objective (Plan-
Apochromat 20x/1.0 DIC D=0.17 M27 75mm). The following settings were used: mCitrine
(yellow fluorescent protein), excitation laser 514 nm (intensity- 2.4-6.0%), emission collection
range 519-596 nm and a z-step of 0.415-0.420 µm. For high laser intensity based live imaging of
pLH13 in wild type and in *as2-7D* to visualize the adaxial sepal epidermis, imaging settings for
mCitrine used: excitation laser 514 nm (intensity- 16-18%), emission collection range 519-596
nm and a z-step of 0.244 µm. Post imaging, the plates were returned to the growth chamber, and
the process was repeated for each time-point. Inflorescence tissues were transferred to fresh
media every 48 hours to prevent fungal contamination.
1098
1099 **NPA treatment assay**
1100
Primary inflorescences of six- to eight-week-old pLH13 in *as2-7D* T2s were dissected down to
stage 3 and were divided into two groups of 5-6 samples. One group of samples were placed in ½
MS live imaging media plates containing 10 µM NPA. The other group, serving as the control,
was placed in media plates containing an equivalent amount of DMSO (20 µL DMSO per 100
ml media). The tissues were dissected 48 hours prior to imaging. Post dissection, the tissues were
treated every 12 hours with a 100 µL liquid solution, applied directly above the inflorescence
tissue by pipetting. For the 10 µM NPA treatment samples, the liquid solution comprised of 1 µL
of 1M NPA+19 µL DMSO+10 µL Silwet in 100 ml water, whereas the control (mock) samples
were treated with 20 µL DMSO+10 µL Silwet in 100 ml water. The aim of administering liquid
solution was to enhance the effectiveness of the treatment, which was performed every 12 hours
throughout the experimental duration. Stage 4 flowers, 48 hours post dissection were selected as
the 0- hour time point for imaging mock and 10 µM NPA treated samples. Live imaging was
performed as described above, with the samples being moved to fresh mock or 10 µM NPA
plates every 48 hours. Imaging settings used: mCitrine (yellow fluorescent protein), excitation

laser 514 nm (intensity -2.6-2.8%), emission collection range 519-596 nm, and a z-step of 0.415

µm.

1117

1118   **Propidium Iodide (PI) cell wall staining and imaging**

1119

A 5 mg/ml stock solution was prepared for PI staining, with 5 mgs of PI (Sigma) solubilized in 1

mL of MilliQ water. The stock solution was stored in a foil-covered 1.7 ml microcentrifuge tube

and refrigerated at 4ºC. For staining, stock solution was five-fold diluted with water containing

0.01% v/v Silwet surfactant (PI working solution). Whole inflorescences and flowers below

Stage 10 were stained for 6-8 minutes, stage 14 sepals were stained for 12 minutes, and leaves

were stained for 25 minutes. Care was taken to ensure that the tissue was fully submerged in the

staining solution, and no air bubbles surrounded the tissue. Following staining, the tissue was

washed in a fresh solution containing just 0.01% v/v Silwet, prior to imaging. Fresh PI working

solution was prepared before every experiment. For imaging PI-stained inflorescences of wild

type, *as2-7D* and the transgenics (T1s) of *pML1::AS2* in wild type, *pML1::REV* in wild type,

pML1::REV in *as2-7D* and the inflorescences, flowers of *kan1-2 kan 2-1* mutants, the following

settings were used: excitation 514 nm (laser intensity -18-22%), emission 566-650 nm. For

imaging sections of PI-stained *kan1-2 kan 2-1* mutant leaves, samples were excited with 514 nm

(laser intensity - 70%) and emission was collected between 566-643 nm range.

1134

1135   **Imaging reporters**

1136

For single time point imaging of the reporters, inflorescences were dissected down to stage 3-4.

The recovery period was extended to 48 hours to allow ample time for stabilization of the

reporter signal. The inflorescences were placed upright in the live imaging media plates with the

meristem facing the objective. Prior to taking images of late-stage flowers, the inflorescences

were allowed to grow for a few days. For the reporters imaged together with PI, fresh

inflorescences were dissected for imaging late-stage flowers. Images were taken with Zeiss 710

upright confocal microscope using 20X water dipping objective.

1144

The following settings specific for each of the reporters were used. For *pAS2 ^{WT}::GFP-GUS* and

pAS2 $^{as2-7D}$::GFP-GUS images, PI - excitation laser 594 nm (intensity- 2%), emission range 599-

650 nm and GFP - excitation laser 488 nm (intensity-20%), emission range 493-550 nm, with z-

step of 0.240 µm. For *proREV::REV-GFP*, Chlorophyll and GFP were excited with 488 nm laser

(intensity- 40%), emission range was collected between 647-722 nm for Cholorophyll and 493-

550 nm for GFP. Z-step, 0.240 µm (inflorescences), 0.4 µm (flowers). For *proFIL::mCherry-

RCI2A*, Chlorophyll and mCherry were excited with 561 nm laser (intensity- 12-15%), emission

range was collected between 655-722 nm for Chlorophyll and 578-650 nm for mCherry. Z-step,

0.240 µm. For *pPIN1::PIN1-GFP* single time-point images, PI - excitation laser 594 nm

(intensity- 2%), emission collection range 599-650 nm and GFP - excitation laser 488 nm

(intensity- 25-30%), emission collection range 493-550 nm, with z-steps of 0.240-0.5 µm. For
pPIN1::PIN1-GFP as2-7D live-imaging, GFP – excitation laser 488 nm (intensity – 20-25%),
emission collection range 493-538 nm and Chlorophyll – excitation laser 488 nm (intensity – 20-
25%), emission collection range 647-722 nm, with z-steps of 0.5 µm. For *pDR5::VENUS*,
samples were excited with an argon laser (514 nm), and data were collected in the YFP (520–565
nm) channel. For *pPDF1::mCitrine-MBD,* PI and mCitrine were excited with 514 nm (intensity-
30-32%), and emission range was collected between 603-643 nm for PI and 519-567 nm for
mCitrine. Z-step, 0.2 µm.
1163
1164 **Imaging adult sepals**
1165
For imaging fully mature sepals, all flowers were dissected off the inflorescence except for a
single stage 14 flowers. All the internal floral organs from the stage 14 flower were carefully
dissected off, leaving just the outer sepal [35] attached to the pedicel. The sepal was wetted in
0.01% v/v Silwet water and mounted between two coverslips filled with 0.01% v/v Silwet water.
To prevent the sepal from drying out during imaging, the coverslips were sealed with vacuum
grease. Sepals were imaged using 10X objective (C-Apochromat 10X/0.45 W M27). Three-
horizontally tiled images were taken with an overlap of 5% and were auto-stitched with a
threshold of 0.70 in the ZEISS ZEN software. For imaging fully mature sepals of pLH13 in wild
type and in *as2-7D*, mCitrine was excited with 514 nm laser (intensity- 20-22%) and emission
was collected at 519-621 nm. For imaging PI-stained sepals of wild types, *as2-7D*, *pML1::KRP1*
and *pML1::KRP1 as2-7D*, PI was excited at 514 nm (intensity- 27-30%) and emission was
collected at 566-648 nm. For epidermal structure observation, mature flowers harboring pAR169
(*pATML1::RCI2A-mCtrine*) were transverse sectioned using a razor blade. Both mCitrine and
chlorophyll were excited using 514 nm laser, and emission was collected at 520-565 nm for
mCitrine and 651-700 nm for chlorophyll.
1181
1182 **Osmotic treatments for measuring epidermal stiffness**
1183
Osmotic treatments were performed as described previously [36,55], with modifications. In brief,
primarily inflorescences from pLH13 in wild type, pLH13 in *as2-7D*, pLH13 in *pML1::KRP1*
and pLH13 in *pML1::KRP1 as2-7D* six- to eight-week-old plants were dissected down to stage 3
to 4 and placed in ½ MS live imaging media plate. Dissected tissues were allowed to recover for
24 hours in the growth chamber. Early stage 6 buds were precisely chosen for osmotic treatment,
since this is the stage right before *as2-7D* outer epidermis starts buckling.
1190
Each bud (corresponding to one replicate) was subjected to osmotic treatment individually and
was imaged pre- and post-treatment. Pre-treatment imaging involved staining the bud with a PI
working solution for precisely 6 minutes. After washing off the PI stain, the tissue was
transferred to a fresh media plate and was oriented to focus on the outer sepal. The tissue was

submerged in 0.01% v/v Silwet water, and the pre-treatment image was captured. Imaging
settings used for mCitrine – excitation 514 nm (intensity- 20-25%), emission 519-567 nm and PI
- excitation 514 nm (intensity- 20-25%), emission 603-643 nm.
1198
For osmotic treatment, the bud was immersed in 0.35M NaCl solution containing 0.01% v/v
Silwet for exactly 20 minutes. Post treatment, the bud was re-stained in a PI working solution
prepared in 0.35M NaCl (1 part 5 mg/mL PI (solubilized in 0.35M NaCl) and 4 parts 0.35M
NaCl solution containing 0.01% v/v Silwet)) for 10 minutes, leading to a cumulative osmotic
treatment duration of 30 minutes. This timeframe was found to be optimum for inducing
plasmolysis in epidermal cells of early stage 6 flowers, wherein the plasma membrane gets
visibly detached from the cell wall at the corners. Re-staining the buds with PI is necessary since
NaCl solution erodes off the PI stain. The bud was finally re-positioned in a fresh media plate
and submerged in 0.35M NaCl solution. Before image acquisition, the tissue was allowed to
stabilize for a few minutes due to its increased fragility post osmotic treatment. The post-
treatment image acquisition setting was as follows: mCitrine – excitation 514 nm (intensity- 25-
35%), emission 519-567 nm and PI - excitation 514 nm (intensity- 25-35%), emission 603-643
nm. This procedure was iterated for all samples individually.
1212
1213 **Image processing**
1214
Converting raw images to tiffs, taking screenshots and videos
The raw images, generated either in lsm or czi formats from ZEN imaging software, were
converted to tiff files in ImageJ distribution Fiji. For multichannel images, images were split
based on channels, which were saved individually. The MorphoGraphX (MGX) [90] software was
used for image processing and taking the screenshots using the built-in screenshot option in
MGX. For taking screenshots of the reporter lines, the reporter image was loaded onto Stack 1,
and either chlorophyll or PI was loaded onto Stack 2. Colormaps for Stack 1 and Stack 2 were
set to green and gray respectively. Flower images and meristems were rotated in the desired
orientation. The voxels not associated with the flower or meristem of interest, were removed
using the Voxel Edit tool in MGX. In addition, trichomes from the outer sepal epidermis were
removed in a similar manner, for better visualization of the sepal surface. Following this,
screenshots were captured. From flower images, trichomes on the outer sepal were removed by
image processing for better visualization of the sepal epidermis. The brightness and contrast
were uniformly adjusted across samples for any given experiment. Screenshots of MGX
generated heatmaps were taken using the default brightness and contrast settings, without any
adjustments.
1231
Videos were captured as a series of two-dimensional tiffs files using the built-in "Record movie"
option in MGX. The 3D stack images were rotated in the MGX window for the desired duration
and time points. For capturing videos involving two genotypes/treatments, the images were

loaded onto Stack 1 and Stack 2 and rotated independently. The stack of tiffs generated was
loaded onto Fiji and saved as a .avi file without compression. The .avi files were assembled and
labeled in Adobe Premiere Pro 2023.
1238
1239 **Generating surfaces for analyzing outer epidermal growth**
1240
Making the surfaces and analysis of growth data was performed in MGX as previously described
in Zhu et. al. and Hong et. al. [36,102], with modifications. The following process was used to
generate the surfaces for analyzing cell growth parameters on the outer epidermis of wild type,
as2-7D, *pML1::KRP1*, *pML1::KRP1 as2-7D*, *as2-7D* mock and 10 µM NPA treatment samples.
Briefly, the tiffs were imported onto Stack 1 Main Store in the software. Trichomes on the outer
epidermis were removed using the voxel edit tool to get a better surface. Since the outer
epidermal morphology of *as2-7D* becomes increasingly irregular in later stages, Level Sets [103]
was used to extract the structure representing the general flower shape. For this, Level Sets was
initialized with an up/down threshold value of 3 to 4. The structure was allowed to evolve (Level
Set Evolve) with default parameters, except for "view steps", which was set to 1 for visualization
at more frequent intervals. For performing Edge Detect Angle, the mask was oriented such that
the outer sepal directly faces the viewer. The Edge Detect Angle process was then run with
default parameters to get a finer structure that closely resembles the image. The holes generated
in the structure (which corresponds to places in the image with low membrane marker signal)
were closed using Stack/Morphology/Closing - X/Y/Z radius of 3 to 8. A 2.5D surface fitted
with a triangular mesh was created from the structure using Marching Cube Surface with a cube
size of 3 to 5 µm, followed by smoothing once (Smooth Mesh, passes = 10) and subdividing
twice. The membrane signal was projected onto the 2.5D surface with a min. and max. distance
range of 1 to 8 µm from the image.
1260
1261 **Generating surfaces for analyzing outer vs inner epidermal growth in wild type and *as2-7D***
1262
To obtain the inner epidermal surface, clipping plane 1 (View Tab/Clip 1), set to a width of 10 to
15 µm, was used to clip the flower image along the longitudinal direction. The voxel editing tool
was used to carefully remove the voxels below the inner epidermis of the sepal within the
clipping plane region. The clipping plane was then moved to the adjacent region and the same
process was repeated, while tilting the plane to ensure voxels from the inner epidermis itself are
not removed. Once the voxels below the inner epidermis were removed throughout, the edited
tiff was copied back to the main stack and saved. To generate the surface for the inner epidermis,
the Z axis of the stack was reversed (Stack/Canvas/Reverse Axes). The stack was blurred using
Gaussian blur (X/Y/Z sigma (µm) = 2 to 2.5), and structure was extracted using Edge Detect
with a threshold of 2000 to 3000. A triangular mesh surface was created using a cube size of 3
µm, followed by smoothing once (passes = 10) and subdividing twice. Plasma membrane signal

from the inner surface was projected onto the mesh surface with a min. and max. distance range

of 3 to 8 µm.

1276

1277   To obtain the outer epidermal surface, the processed tiff, prior to reversing the axes, was used.

1278   To process the stack, Gaussian blur (X/Y/Z sigma (µm) = 1 to 1.5) followed by Edge detect

1279   (threshold of 8000 to 10,000) were used sequentially. Mesh surface was created as described

1280   above, and the membrane signal was projected within a 2 to 4 µm distance range from the image.

1281

1282   **Generating surfaces for analyzing growth of outermost mesophyll layer in wild type and**

1283   ***as2-7D***

1284

1285   To enhance the membrane marker signal for the mesophyll cells, we utilized a deep learning

1286   approach based on a Convolutional Neural Network (CNN). We used the inbuilt CNN Unet 3D

1287   prediction algorithm VijayanUnet.pt [47] in MorphoGraphX to precisely predict the cell outlines.

1288   The algorithm then uses these predictions for enhancing and de-noising the membrane marker

1289   signal and generates a new image based on the predictions.

1290

For both wild type and *as2-7D* mutants harboring pLH13 membrane marker, the CNN algorithm

was run on the pre-processed confocal images of the outer sepal, where the parts of the flower

not associated with the outer sepal were already removed by voxel editing (see 'Generating

surfaces for analyzing outer vs inner epidermal growth in wild type and *as2-7D*' section in

Materials and Methods). Using the pre-processed sepal images instead of whole flowers for

running the CNN algorithm largely reduced computational processing times. Briefly, for wild

tpye and *as2-7D* mutants, CNN-generated images of the sepals for each time point were obtained

by running Stack/CNN/UNet3D Prediction/VijayanUnet.pt using default parameters. The CNN

Unet3D prediction process usually also generates voxels unrelated to the sepals, which were

removed using the 'Voxel edit' tool. These edited CNN-generated images were then used to run

Stack/Mesh Interaction/Annihilate function, to remove the voxels associated with the outer

epidermis and the inner layers and obtain the outermost mesophyll layer (layer closest to the

outer epidermis). For sepals at 0-hour and 24-hour time points, the outermost mesophyll layer

was extracted by setting the minimum distance values for Annihilate function to 5/6 and the

maximum distance values to 15. For the 48-hour time points, the minimum and maximum

distance value for the Annihilate function was set to 6 and 20 respectively. The resulting images

generated post running Annihilate function were used to generate the 2.5D surface for the

outermost mesophyll layer. Briefly, for each time-point and genotype, the annihilated image

stack was loaded onto Stack 1 and blurred using Gaussian Blur, with X/Y/Z sigma (µm) value of

3. The surface was extracted using Edge Detect threshold of 2000 to 3000. The mesh surface was

created using a cube size of 3 µm, followed by smoothing once (passes = 10) and subdividing

twice. The membrane signal was projected onto the mesh surface with a minimum and maximum

distance range of 7 to 8 µm.

1314
1315
Lineage tracking
For meshes across all time points, individual cells were manually seeded. Cells were segmented
using the watershed algorithm, and unlabeled vertices were fixed by running Cell Mesh/Fix
Corner Triangles. Cell lineages were parent tracked to the earlier time-point to analyze cell
growth. Parent (lineage) tracking involves transferring the parent labels from cells in the earlier
time point onto the daughter cells at the later time point. For this, the meshes corresponding to
the earlier (Mesh 1) and later (Mesh 2) time points were opened side by side on the same MGX
window. A few daughter cells in Mesh 2 were manually parent labeled, and using these as
landmarks, the remaining daughter cells were auto parent labeled (by running
Deformation/Morphing/Set Correct Parents followed by Deformation/Mesh 2D/Auto Parent
Labelling 2D). Errors in segmentation or parent tracking were manually corrected. Parent
tracking was checked for accuracy using Cell Axis/PDG/Check Correspondence, which checks
whether the cell junctions are identical across time points. The final parent labels were saved as
.csv files. A custom R script (combinelineages.r) was run to automatically correlate the lineage
information through multiple time points, followed by manual correction.
1331
1332 **Analyzing cell growth parameters**
1333
Meshes corresponding to the earlier (Mesh 1) and later time points (Mesh 2) were loaded side by
side in the same MGX window, with parent labels loaded onto Mesh 2. Cell growth rate was
calculated based on the total area of the daughter cells in the second time point divided by the
area of the parent cell in the earlier time point. Growth rate heatmaps were generated using Heat
Map/Heat Map Classic/Heat Map with the Type set to Area. The cell division rate was calculated
based on the total number of daughters formed in the second time point for each cell in the
earlier time point. Cell division heatmaps were generated using Lineage Tracking/Heat Map
Proliferation. The values for growth rate and cell division rate were exported to .csv files.
1342
The Principal Direction of Growth (PDG) uses information of corresponding junctions between
the earlier and later time points, to compute the direction of maximum and minimum growth of
each individual cell in the earlier time point [90]. In the figures, the PDGs represent the maximum
growth direction (PDG_{max}) tensor (yellow lines), the length of which is directly proportional to
the magnitude along the direction of maximum growth. The PDGs were calculated for the cells
in the earlier time point (using the information of parent labels from the later time point), the
PDG_{max} tensors were projected onto the later time point for visualization. For displaying PDG_{max},
the values for line width and line scale were set to 5 and 3 respectively.
1351
The Angle of Deviation of the PDG_{max} tensor was calculated relative to the longitudinal axis of
the sepal. The sepal longitudinal axis was set using a Bezier line on the earlier time point, since

PDGs and deviation angles are calculated on the earlier time point. Briefly, a new Bezier grid
(Misc/Measure/New Bezier) was created, which was then collapsed into a Bezier line
(Misc/Bezier/Collapse Bezier Points). The supporting points on the Bezier line were placed on
the mesh in a manner such that the Bezier line is parallel to the sepal longitudinal direction, and
the line shape fits the curvature of the sepal. The custom axis for each cell was set using this
Bezier line (Cell Axis/Custom/Create Bezier Line Directions) followed by smoothing (Smooth
Custom Directions). The PDGs were loaded onto the mesh, and the angle of deviation of the
PDG_{max} from custom X direction (which represents the longitudinal axis) was calculated using
Cell Axis/Custom/Custom Direction Angle. The values of the angle of deviation heatmap were
exported as .csv files.
1364
1365 **Analyzing osmotic treatment induced epidermal shrinkage**
1366
Our osmotic treatment conditions were optimized such that the cell walls shrank, but the cell
membrane barely plasmolyzed. Post osmotic treatment, the membrane just visibly pulled away
from the cell wall corners, thereby rendering the wall and membrane area almost comparable.
Since the PI signal (corresponding to the cell wall) is very weak post-treatment, especially for the
inner layers, the PI and mCitrine (membrane marker) images were merged as a single image
MGX (Stack/Multistack/Merge Stacks).
1373
For analyzing epidermal stiffness, we assume that the surface tension across genotypes is
comparable. The shrinkage in area of the epidermal layers along a comparable longitudinal
section was analyzed to infer tissue stiffness. For this, the pre- and post-treatment merged stacks
for any given sample were opened side by side in the same MGX window. Using clipping plane
1 (Clip 1), longitudinal cross-section corresponding to the same regions were extracted from the
flower images. The tiffs were processed using Levels Sets to extract the global shape. Mesh
surfaces were made with a cube size of 3 µm, which was smoothed once (10 passes) and
subdivided twice. Signal was projected with a 1.5 to 3 µm distance range from the image. From
the cross-section surface, the outer epidermis and inner epidermis were seeded and segmented as
single "cells". Care was taken to ensure that the epidermal region being seeded was comparable
between the pre- and post-treatment surfaces. For measuring shrinkage, parent labels from the
pre-treatment mesh were transferred onto the post-treatment mesh, following which the
shrinkage ratios were quantified (Mesh/Heatmap/Heatmap Classic). The areal shrinkage values
were stored as a .csv file, which was converted to percentages for generating heatmaps shown in
the figures.
1389
1390 **Computing minimum curvature**
1391
In MorphographX, minimal curvature analysis quantifies the least degree of curvature at any
given location on the surface. The output is displayed in the form of a heatmap where warm

colors represent regions of higher curvature or outward bulging, whereas cool colors represent
regions of lower curvature, such as inward deformations (saddles). For calculating the minimal
curvature for the sepals from the live imaging series of *pPIN1::PIN1-GFP as2-7D* flowers, we
used the images obtained from the chlorophyll autofluorescence channel to construct the mesh
surfaces. Briefly, the chlorophyll image corresponding to a given flower (for which the minimal
curvature is to be analyzed) was loaded onto Stack 1. The structure representing the general
shape of the flower was extracted using Level Sets (Stack/Lyon/Init Level Set), with an up and
down threshold values set to 4. The structure was allowed to evolve (Stack/Lyon/Level Set
Evolve) with default parameters. The triangular 2.5D mesh was created using
Mesh/Creation/Marching Cube Surface with 5 µm cube size, followed by one round of
smoothing (Mesh/Structure/Smooth Mesh, passes = 5) and subdividing
(Mesh/Structure/subdivide). Minimal curvature was then calculated using Mesh/Signal/Project
Mesh Curvature, selecting Type = Minimal and setting the neighborhood radius value to 50 µm.
For visualizing the resulting heatmap, the transfer function was set to "Jet".
1408
1409

1410   **Analytical model for a growing trilayer**

1411

System and assumptions. We consider a growing elastic system made of three layers,
corresponding to the inner epidermis (denoted i), mesophyll (m), and outer epidermis (o) of a
lateral organ. We consider that the three layers may grow at different (specified or target) rates
and have different elastic moduli. To simplify the formulae, we assume that the epidermal layers
have similar thicknesses. We also assume that the Poisson ratios of all layers are equal (value v).
Finally, we neglect the bending modulus of each layer considering their small thickness. Note,
these three assumptions do not affect the qualitative behavior of the system.

We assumed E_i, E_m, and E_o be the values of elastic modulus, h, h_m, and h the values of
thickness, and g_i, g_m, and g_o as the values of strain associated with target (specified) growth rates
for the inner epidermis, the mesophyll, and the outer epidermis, respectively.

1422

The differences in growth rates between the layers may lead to changes in system geometry. As
the three layers are bound to each other, they have approximately the same curvature c
(corresponding to the radius of curvature measured at the middle of the mesophyll), while the
strain in the inner and outer layers can be deduced from the strain in the middle layer ε using
curvatures and assuming that the reference state is flat. From kinematics of the system, the
elastic (residual) strains in the three layers are given by

1429
$$\varepsilon_i = \varepsilon - \frac{1}{2}(h_m + h)c - g_i$$
1430
$$\varepsilon_m = \varepsilon - g_m$$

1431
$$\varepsilon_o = \varepsilon + \frac{1}{2}(h_m + h)c - g_o$$

1432 The elastic energy of the system takes the form

1433
$$E = \frac{1}{2(1-v^2)}[E_i h \varepsilon_i^2 + E_m h_m \varepsilon_m^2 + E_o h \varepsilon_o^2]$$

1434

1435 Minimizing the elastic energy with respect to strain and curvature yields their values at
1436 equilibrium. In particular, using the notation

1437
$$A = \frac{1}{(4E_i E_o h + E_m (E_i + E_o) h_m)} > 0$$

1438 we obtain the equilibrium values for the strain

1439
$$\varepsilon = A(2E_i E_o h(g_i + g_o) + E_m (E_i + E_o) h_m g_m)$$

1440 and the curvature

1441
$$c = \frac{2A}{h+h_m}\left(2E_i E_o h(g_o - g_i) + E_m E_o h_m(g_o - g_m) - E_m E_i h_m(g_i - g_m)\right).$$

1442 This curvature is in general non-zero, the system curves with the abaxial side on the convex side
1443 as soon as the abaxial growth rate is sufficiently larger than the adaxial growth rate.

1444 To determine whether buckling of the abaxial epidermis may occur, we compute the elastic
1445 (residual) strain in this layer at equilibrium:

1446
$$\varepsilon_o = AE_m E_i h_m(2g_m - g_i - g_o).$$

1447 ε_o can become compressive (<0) only if $2g_m - g_i - g_o < 0$. When it falls beyond a threshold
1448 [104], this leads to buckling of the abaxial side, with patterns as described [105], depending on the
1449 anisotropy of the residual strain.

1450 Interestingly, the elastic strain of the adaxial epidermis,

1451
$$\varepsilon_i = AE_m E_o h_m(2g_m - g_i - g_o)$$

always has the same sign as that of the abaxial epidermis. This means that either the two
epidermal layers are in compression, or the two epidermal layers are in tension. A situation
where one epidermis would be in tension and the other would be in compression does not
correspond to mechanical equilibrium. Nevertheless, the amount of compression may differ
between the two epidermal layers: the ratio of the elastic deformations is related to the ratio of
Young's moduli

1458
$$\frac{\varepsilon_o}{\varepsilon_i} = \frac{E_i}{E_o}.$$

1459 Accordingly, it is possible that only one of the two epidermal layers reaches the buckling
1460 threshold. For instance, a much smaller Young's modulus for the outer epidermis compared to
1461 the inner epidermis would lead to buckling of the outer epidermis exclusively.

1462

1463 Finally, we mention that the elastic strain in the mesophyll has an opposite sign

1464
$$\varepsilon_m = -2AE_i E_o h(2g_m - g_i - g_o).$$

1465 This means that as soon as the epidermis is under compression, the mesophyll is under tension
1466 and vice versa.

1467
1468 **Numerical model for a growing layered sepal**
1469

We again considered a growing elastic system comprising of three layers (inner epidermis (i),
mesophyll (m), and outer epidermis (o)). To better represent the geometry of the transversal
section of a sepal, we also considered two lateral semi-circles (the edges) to connect the inner
and outer epidermis (Fig. 2, I to K). These semi-circles are denoted "end", while the region
enclosed by the epidermal layers and the edges denote the mesophyll of the trilayer. This
geometry is characterized by three parameters: the epidermal layer thickness h, the mesophyll
thickness h_m and the width of the trilayer part L.

We adopt the framework of morpho-elasticity [106], assuming that the residual stresses are solely
created by a local growth deformation tensor F_g, and the observed deformation gradient can be
decomposed multiplicatively on the elastic and the growth deformation tensors. In this
framework we consider a quadratic energy density in the Green-Lagrange strain tensor E_e,

1481
$$W = \frac{1}{2}E_e : H : E_e,$$

1482 where H, the elasticity tensor of rank four, represents the mechanical properties of the material.
1483 The proposed model assumes all regions of the geometry to be isotropic materials. Therefore, the
1484 elasticity tensor field H, is characterized by independent parameters: the Young's moduli of the
1485 four regions (E_i, E_m, E_o and E_{end}), and the common Poisson ratio v. An equation for the time
1486 evolution of the growth tensor F_g completes the model,

1487
$$\begin{aligned} \frac{\partial F_g}{\partial t} &= GF_g, \quad t > 0 \\ F_g &= 1, \quad\quad t = 0 \end{aligned},$$

1488 with G the growth rate tensor. We consider here that the growth rate has both an imposed target
1489 (or specified) component G_T, as well as a component that stands for an influence of the elastic
1490 strain on growth,

1491
$$G = G_T + CE_e,$$

with C being the extensibility (assumed to be scalar). We assume that the target growth is
unidirectional along the trilayer, so that the target growth rates γ_i, γ_m, γ_o and γ_{end} for the inner
epidermis, the mesophyll, the outer epidermis, and the edge respectively are the only
independent parameters encoded in the target growth rate tensor G_T. As for the extensibility C, we
have four additional parameters C_i, C_m, C_o and C_{end} corresponding to the four regions.

In comparison with the analytical model presented above, all geometrical and mechanical
parameters are the same, while the growth parameters g_i, g_m, and g_o can be seen as the relative
growth cumulated until time t, which can be approximated as when $C = 0$.

The model of the growing layered sepal was solved using the Finite Element Method and
implemented with the FEniCS computing platform [107] with its Python interface, with a special
use of the BVPy Python library [108].

1503

We simulated the deformation of the defined geometry when the outer epidermis is growing (
$\gamma_o \neq 0$), while the other regions do not ($\gamma_i = \gamma_m = \gamma_{end} = 0$). This is one of the simplest situations,
when the mean growth rate of the two epidermal layers is higher than the growth rate of the
mesophyll, and according to the predictions of the analytical model the epidermal layers are
under compression. This first condition for buckling being satisfied, we studied whether buckling
indeed occurs depending on the moduli of the mesophyll and epidermal layers (fig. S3). To
validate the approach, we also compared the equilibrium curvature predicted by the analytical
model to the mean midline curvature of the deformed sepal in the numerical model (see fig. S4).

1512

For all our simulations we considered fixed values for the length $L = 10$, and thickness $2h + h_m =$
5 of the sepal, the target growth rate of the outer epidermis $\gamma_o = 0.1$, and the Poisson ratio $v =$
0.49. The value of the extensibility is chosen low for all regions but relatively higher for the
mesophyll region, $C_o = C_i = C_{end} = 0.01$ and $C_m = 0.1$. The Young's modulus of the mesophyll is
chosen as the lowest among all regions, $E_m = 1$, while the Young's modulus of the edge
epidermis is a high value, $E_{end} = 10^6$. As for the Young moduli of the outer and inner epidermis,
we performed an exploration of the parameter space defined by these two extreme values,
$(E_o, E_i) \in [1, 10^6] \times [1, 10^6]$. We present the results of this analysis for different values of the
epidermis thickness, $h \in \{0.1, 0.2, 0.3, 0.4\}$ (fig. S3 and S4).

1522

1523
1524 **Graph plotting**
1525
1526 Graphs were generated in R (version 4.2.2 (2022-10-31 ucrt)) using ggplot, raw data for which is
1527 provided in Data S1.

1528   **SUPPLEMENTARY FIGURE TITLES AND LEGENDS**

1529

Fig. S1. *AS2* **overexpression causes characteristic ridges, invaginations, and outgrowths on the outer epidermis, associated with Main Fig. 1.**

(**A** and **B**) Representative optical cross-sections of Stage 11 flowers of (**A**) the wild type (WT) and (**B**) *as2-7D* mutant. Images are representative of T2s harboring pAR169 (*pATML1::RCI2A-mCitrine*), an epidermal layer specific plasma membrane marker. Green and grey represent epidermal cells and chlorophyll containing mesophyll cells respectively. WT sepals (A) have a smooth shape, while the *as2-7D* sepal (B) outer epidermis exhibit outward protrusions (ridges), inward indents (invaginations) and pointed epidermal extensions (outgrowths). Note that the outgrowths lack mesophyll cells (grey) underneath whereas mesophylls occupy the space beneath ridges and invaginations.

(**C** and **D**) Whole inflorescence images of (**C**) Col-0 (control) and (**D**) *as2-5D* (in Col-0, isolated by Wu et. al [41]). Images are representative of $n = 3$ inflorescences. Scalebar, 1 cm.

(**E** and **F**) Stage 14 flowers of (**E**) Col-0 and (**F**) *as2-5D* mutant, representative of $n = 3$ flowers. Yellow arrowhead points toward an outgrowth. Note the mild nature of outgrowth formation in *as2-5D* mutant as compared to *as2-7D* (Fig. 1). Scalebar, 1 cm.

(**G** to **J**) Representative six-week-old adult plants of (**G**) WT, (**H**) *as2-7D* and (**I-J**) *pML1::AS2* in WT T1s. The transgenics show strong (17/28 plants, panel I) to extreme (9/28 plants, panel J) developmental defects, categorized qualitatively by the extent of leaf deformity. The transgenics exhibit outgrowths on the lower (abaxial) surface of the leaves as well, in addition to severely stunted growth. Scalebar, 1 cm.

(**K** to **N**) Fully formed siliques of (**K**) WT, (**L**) *as2-7D*, (**M-N**) *pML1::AS2* in WT T1s. Note the outgrowths on the silique epidermis. In panels M and N, siliques are representative of strong and extreme categories, described for panels I and J respectively. Three siliques on each panel correspond to three independent plants. Total number of independent transgenics (T1s) assessed $n = 28$.

Fig. S2. The outer epidermis of *as2-7D* exhibits loss of stereotypical growth patterns,
associated with Main Fig. 2.

(**A** to **D**) Confocal stack images of (**A**) Wild type (WT) replicate 2, (**B**) WT replicate 3, (**C**) *as2-*
7D replicate 2 and (**D**) *as2-7D* replicate 3 developing flowers, transformed with the pLH13
(*35S::mCitrine-RCI2A*) membrane marker. Images correspond to the 48-hour (left) and 96-hour
(right) time points from WT and *as2-7D* live imaging experiment ($n = 3$). Yellow arrowheads
show outgrowths. Trichomes were removed from the outer sepal epidermis by image processing.
Scalebar, 50 µm.
(**E** to **H**) Cell area growth rates of (**E**) WT replicate 2, (**F**) WT replicate 3, (**G**) *as2-7D* replicate 2
and (**H**) *as2-7D* replicate 3. The heatmap is projected onto the later time point. Red to blue
gradient shows fast to slow growth. Yellow arrowheads point towards the outgrowth which
exhibit localized fast growth. Scalebar, 50 µm.
(**I** to **L**) The cell division rates of (**I**) WT replicate 2, (**J**) WT replicate 3, (**K**) *as2-7D* replicate 2
and (**L**) *as2-7D* replicate 3. Red to blue gradient shows more to less division. Cell division over
0-48 hours is shown on the left and over 48-96 hours on the right. Scalebar, 50 µm.
(**M** and **N**) Graphical representation of (**M**) cell growth rates and (**N**) cell division rates across all
three independent biological replicates of the WT (black dots) and *as2-7D* mutant (red dots). Dot
represents the growth rate (M) and division rate (N) of a given cell lineage. Dot plots are
superimposed with box plots, where the central line indicates the median. Left and right panels
display data for 0-48 and 48-96 hours, respectively. Note the reduced spatial variability in the
growth rates of *as2-7D* (panel M). Lines above data points indicate statistical comparisons. The
p-values for Bonferroni corrected pairwise t-test (two-tailed) are shown. *p-values*: ***<0.0001,
**<0.001, *<0.05, ns = non-significant. Exact *p-values* for growth rates: WT vs *as2-7D* for 0-48
hours, 0.9884; WT vs *as2-7D* for 48-96 hours, 1.565172×10^{-93}. Exact *p-values* for cell division
rates: WT vs *as2-7D* for 0-48 hours, 7.800489×10^{-05}; WT vs *as2-7D* for 48 - 96 hours, 9.47882
$\times 10^{-07}$.

1582
1583
Fig. S3. Numerical simulations for a sepal cross-section with the outer epidermis growing
in width, showing that buckling is determined by the relative Young's moduli of the outer
and inner epidermis, associated with Main Fig. 2.

Parameter space exploration for the numerical model of the growing sepal with three layers,
corresponding to the outer epidermis (blue), mesophyll (yellow) and the inner epidermis (orange)
as shown in Fig. 2, I to K. As a proxy for detecting whether outer epidermis buckles or not, we
monitor the number of curvature maxima on the surface of the outer epidermis. A value for
maxima higher than one indicates the occurrence of a buckling event. Block rows in the grid
correspond to different epidermal thickness values $h \in \{0.1, 0.2, 0.3, 0.4\}$, while block columns
correspond to different time points during growth. In each block the Young's moduli of the outer
and inner epidermis are variable, $(E_o, E_i) \in [1, 10^6] \times [1, 10^6]$. All other parameters are kept
constant: $L = 10, 2h + h_m = 5, \gamma_o = 0.1, \gamma_i = \gamma_m = \gamma_{end} = 0, \nu = 0.49, C_o = C_i = C_{end} =$
$0.01, C_m = 0.1, E_m = 1, E_{end} = 10^6$. Holes in the grid are simulations for which the numerical
solver diverged for the given parameter set and time point.

1598

1599

Fig. S4. Cross validation of the analytical and numerical models for the growing layered
sepal, associated with Main Fig. 2.
(**A**) Equilibrium curvature value predicted by the analytical tri-layer model. (**B**) Mean midline
curvature of the deformed sepal in the numerical model. (**A** and **B**) Parameter values and the
organization of the grids are the same as described in fig. S3. The curvature values in the
numerical model deviate from the analytical model prediction primarily when the inner
epidermis is much softer than the outer epidermis and the edges. We attribute this deviation to
the difference in geometry between the two models, namely the presence of a rigid edge in the
numerical model, that connects the inner and outer epidermal layers.

1609
1610
Fig. S5. Slower growth on the inner epidermis together with misdirected outer epidermal
growth trigger buckling in *as2-7D*, associated with Main Fig. 3.

(**A** to **D**) Growth rate heatmaps for (**A**) Wild type (WT) replicate 2, (**B**) WT replicate 3, (**C**) *as2-*
7D replicate 2 and (**D**) *as2-7D* replicate 3 developing sepals. Red to blue heatmap gradient
shows fast to slow growth. Outer (left) and inner (right) epidermal growth rates over 48 hours is
projected as a heatmap onto the later time point. Scalebar, 20 µm.
(**E**) Graph shows cell growth rates over a 48-hour period for the WT (black) and *as2-7D* (red)
mutant ($n = 3$ biological replicates). Each dot represents the growth rate for a given cell lineage.
Dot plots are overlayed with boxplots, where the central line represents the median. The *p-values*
for Bonferroni corrected pairwise t-test (two-tailed) are shown. *p-values*: ***<0.0001, **<0.001,
*<0.05, ns = non-significant. Exact *p*-values for epidermal cell growth rate: WT outer vs inner,
5.751359×10^{-35}; *as2-7D* outer vs inner, 3.039502×10^{-25}; WT vs *as2-7D* outer, 0.1011; WT vs
as2-7D inner, 8.496122×10^{-16}.
(**F** to **I**) Cell division heatmaps for (**F**) WT replicate 2, (**G**) WT replicate 3, (**H**) *as2-7D* replicate
2 and (**I**) *as2-7D* replicate 3. Red to blue color gradient shows more to less cell division. The
data is presented in the same format as described for panels A-D. Scalebar, 20 µm.
(**J**) Graph shows cell division rates for the WT and *as2-7D* mutant ($n = 3$ biological replicates)
Dots represent the number of cell divisions for each cell lineage. Data presentation and statistical
comparisons are same as described above for panel E. Exact *p*-values for epidermal cell division
rates: WT outer vs inner, 4.08502×10^{-28}; *as2-7D* outer vs inner, 0.004; WT vs *as2-7D* outer,
3.677026×10^{-23}; WT vs *as2-7D* inner, 1.865456×10^{-06}.
(**K** to **N**) Heatmap representing the angle of deviation of PDG$_{max}$ tensors from the sepal
longitudinal axis for (**K**) WT replicate 2, (**L**) WT replicate 3, (**M**) *as2-7D* replicate 2 and (**N**)
as2-7D replicate 3. Scalebar represents a deviation of 90º in red and 0º (no deviation) in blue.
Scalebar, 20 µm.
(**O**) Graph showing the angle of deviation for the WT and *as2-7D* mutant ($n = 3$ biological
replicates). Each dot represents the angle of deviation for a given cell lineage. Data
representation and statistical comparisons are shown in the same format as described for panels E
and J. Exact *p-values* for epidermal angle of deviation: WT outer vs inner, 3.793354×10^{-05}; *as2-*
7D outer vs inner, 1.572509×10^{-20}; WT vs *as2-7D* outer, 8.052118×10^{-10}; WT vs *as2-7D*
inner, 0.349.
1642
1643

Sepal image   Longitudinal section   Longitudinal section (from CNN generated image)   Outer mesophyll surface

WT

as2-7D

Mesophyll cell growth rate

Principal Directions of Maximum Growth

1644

Fig. S6. Growth patterns of the mesophyll cells mimic that of the outer epidermis for both
wild type (WT) and *as2-7D*, associated with Main Fig. 3.
(**A** to **B**) Processed confocal images of (**A**) the wild-type (WT) and (**B**) *as2-7D* mutant stage 7
sepals, harboring the pLH13 membrane marker, at the 48-hour time point (representative of *n =*
3). Scalebars, 20 µm.
(**C** to **D**) View of the longitudinal cross section through the center of the stage 7 sepal blade for
(**C**) the WT and (**D**) *as2-7D* mutant (representative of *n = 3*). Scalebars, 20 µm.
(**E** to **F**) Longitudinal sections of stage 7 sepal images of (**E**) the WT and (**F**) *as2-7D* mutant
processed with CNN (Unet display/Vijayan.Unet algorithm, see Materials and Methods), to
enhance the cell outlines (representative of *n = 3*). The gray areas mark the mesophyll layer
closest to the outer epidermis (outer mesophyll), generated after running (Mesh
Interaction/Annihilate) process in MorphographX to remove the outer epidermis. The annihilated
image (gray) was overlayed on the CNN generated image (green) for comparison. The outer and
inner epidermal layers are shown (yellow arrows). Scalebars, 20 µm.
(**G** to **H**) Plasma membrane signal from the outer mesophyll layer of (**G**) to WT and (**H**) the *as2-*
7D mutant projected onto the mesh surface generated using MorphographX. Scalebars, 20 µm.
(**I** to **J**) Growth rates of the mesophyll cells over a 48-hour period for (**I**) the WT and (**J**) *as2-7D*
mutant are shown (representative of *n = 3*). Red to blue heatmap gradient shows fast to slow
growth. Scalebars, 20 µm.
(**K** to **L**) Principal directions of maximal growth (PDGmax, yellow) of the mesophyll cells of
(**K**) WT and (**L**) *as2-7D* mutant are shown. Segmented cells in grey are superimposed on the
plasma membrane signal (also in grey). Note that both the growth rates (panels I to J) and growth
directions of the mesophyll cells are similar to those of the outer epidermis, as shown in Main
Figure 3. Scalebars, 20 µm.
(**M**) Graph shows the cell growth rates of the mesophylls cells over a 48-hour period for the WT
(black) and *as2-7D* (red) mutant (*n = 3* biological replicates). Each dot represents the growth rate
for a given cell lineage. The dot plots are overlayed with box plots, where the central line
represents the median. The p-value from Tukey's test (two-tailed) is shown. *p-values*: ns = non-
significant (*p> 0.05*).
(**N**) Graph shows the average cell growth rates of the outer epidermis (Outer), inner epidermis
(inner), average of the outer and inner epidermis and the mesophyll cells of the WT (black) and
as2-7D (red) mutant. Each dot represents the average growth rate across all cell lineages in each
replicate (*n = 3* biological replicates). The lines represent the mean growth rate across three
biological replicates. The p-value from Tukey's test (two-tailed) is shown. *p-values*: *<0.05, ns =
non-significant. Exact p-values for WT average (Outer & Inner) vs Mesophyll, 0.0036; WT
average (Outer & Inner) vs Mesophyll, 0.0036.
1681

1682

Fig. S7. Analysis of cell growth direction and stiffness due to *KRP1* overexpression in wild
type (WT) and *as2-7D*, associated with Main Fig. 5.
(**A** to **D**) Confocal images of the inner epidermis of (**A**) Wild type (WT), (**B**) *as2-7D* mutant, (**C**)
AtML1::KRP1 and (**D**) *AtML1::KRP1 as2-7D* stage 14 sepals. The sepals were stained with PI.
Note the smooth morphology of the inner epidermis across all genotypes. Scalebar, 200 µm,
images representative of $n = 3$ adult sepals.
(**E** to **L**) Principal direction of growth (PDG$_{max}$) for the outer epidermal cells of (**E**) WT
Replicate 2, (**F**) *as2-7D* mutant Replicate 2, (**G**) *AtML1::KRP1* Replicate 2, (**H**) *AtML1::KRP1*
as2-7D Replicate 2, (**I**) WT Replicate 3, (**J**) *as2-7D* mutant Replicate 3, (**K**) *AtML1::KRP1*
Replicate 3 and (**L**) *AtML1::KRP1 as2-7D* Replicate 3 sepals. PDG$_{max}$ tensors (yellow)
calculated over a 48-hour live imaging period starting from stage 4, projected on a later time
point surface. Segmented cells in grey are superimposed onto the plasma membrane signal (also
in grey). Scalebar, 20 µm, $n = 3$.
(**M** and **N**) Representative confocal images of WT ($n = 5$) early stage 6 flower (**M**) before and
(**N**) after osmotic treatment in 0.35M NaCl. The PI (cell wall stain) and membrane marker
(pLH13, *35S::mCitrine-RCI2A*) signals have been merged together and shown. The highlighted
region of the outer sepal (white rectangle) indicates the comparable longitudinal section
extracted for analyzing epidermal shrinkage.
(**O** and **P**) MorphoGraphX generated surface of the longitudinal section (**O**) before treatment and
(**P**) after osmotic treatment, extracted from flower images shown in panels M and N respectively.
Comparable regions of the outer and inner sepal epidermis were segmented (outlined in cyan) to
calculate the percentage areal shrinkage.

Fig. S8. Ectopic *AS2* expression mediated buckles and subsequent outgrowth formation occur independent of *REVOLUTA*, associated with Main Fig. 6.

(**A**) Representation of the previously characterized meristem initiation defect phenotype of *rev-6*
mutant. Images show an excised lateral inflorescence meristem of the WT (left) and *rev-6* mutant
(right). Majority of the cauline leaves axils (white arrows) in the *rev-6* mutant are bare, as they
fail to produce axillary meristem, as described by Otsuga et. al. [64]. Scalebar, 1cm.
(**B** to **D**) Fully developed siliques of (**B**) *rev-6* and (**C** and **D**) *pML1::AS2* in rev-6 T1s. No of
independent T1s assessed $n = 19$. The three siliques correspond to three independent plants. The
numbers represent the total number of plants showing strong (12/19 plants, panel C) to extreme
(7/19 plants, panel D) phenotypes, categorized based on the extent of leaf deformity in the adult
vegetative stage.
(**E** to **I**) Representative six-week-old adult plants of (**E**) WT, (**F**) *rev-6*, (**G**) *as2-7D* and (**H** and
I) *pML1::AS2* in *rev-6* T1s. Total number of T1s assessed $n = 19$. The transgenics show strong
(12/19 plants, panel H) to extreme (7/19 plants, panel I) defects in leaf development,
characterized by outgrowth formation on the abaxial epidermis and stunted growth. Scalebar, 1
cm.
(**J** to **M**) Confocal images of PI-stained (greyscale) inflorescences of (**J**) WT, (**K**) representative
pML1::REV in WT T1, (**L**) *as2-7D* mutant, and (**M**) representative *pML1::REV* in *as2-7D* T1.
Note that the transgenics ectopically expressing *REV* looks like their respective backgrounds in
either the WT (panels J and K) or *as2-7D* (panels L and M). Yellow arrowheads point to the
outgrowths. For each *pML1::REV* transgenics in WT and in *as2-7D* backgrounds, number of
independent T1s assessed, $n = 28$. Scalebar, 50 µm.

Fig. S9. KANADI loss-of-function mutant (*kan1-2 kan2-1*) exhibits floral development defects which are distinct from *as2-7D*, associated with Main Fig. 6.

(**A**) Whole inflorescence images of *kan1-2 kan2-1*. The image is representative of *n* = 3 inflorescences. Scalebar, 1 mm.

(**B**) Stage 14 flower of *kan1-2 kan2-1*. The image is representative of $n = 3$ inflorescences.

Scalebar, 1 mm.

(**C**) Confocal image of PI-stained (greyscale) *kan1-2 kan2-1* mature sepal showing the outer

(abaxial) epidermis. The image is representative of $n = 3$ sepals. Scalebar, 100 µm. Note the

twisted and curled morphology of the sepal.

(**D**) Inset of *kan1-2 kan2-1* mature sepal outlined in panel (C). Note the folds on the abaxial

epidermis of *kan1-2 kan2-1* sepal, which are less severe as compared to *as2-7D*. Scalebar, 50

µm.

(**E**) Confocal image of PI-stained (greyscale) *kan1-2 kan2-1* inflorescence. The image is

representative of $n = 3$ inflorescences.

(**F** to **L**) Confocal images of PI-stained (greyscale) *kan1-2 kan2-1* flowers of different

developmental stages (staged based on floral organ development in accordance with Smyth et al.,

1990 [88]. The sepals are shown by white arrows. All the floral organs are morphologically

deformed. Note that the sepals fail to enclose the floral meristem from initial stage onwards and

exhibit a narrow blade development in the later stages. Parts of the inflorescence tissue not

associated with the flowers were removed by image processing. Scalebar, 20 µm.

(**M**) Representative five-week-old adult *kan1-2 kan2-1* plant. The image is representative of $n =$

3 plants. Scalebar, 1 cm.

(**N**) Representative rosette leaf of *kan1-2 kan2-1*. Yellow arrowheads show the outgrowths. The

image is representative of $n = 3$ leaves. Scalebar, 1 cm.

(**O**) Confocal image showing a region of the abaxial epidermis of kan*1-2 kan2-1* leaf (PI-stained,

grayscale). Yellow arrows show the outgrowth, while the white arrows denote the folds on the

leaf. Note that the outgrowth forms in the region where the folds form. Scalebar, 200 µm.

1769

Fig. S10. Outgrowths on *as2-7D* outer epidermis are formed due to buckling mediated PIN1 convergence and subsequent auxin maxima formation, associated with Main Fig. 7.

(**A** and **B**) Confocal images show the expression pattern of PIN1 protein reporter (*pPIN1::PIN1-GFP*) in (**A**) wild-type (WT) and (**B**) *as2-7D* inflorescence meristems. Note the expression of *PIN1* directed towards the emerging outgrowths on the outer epidermis of *as2-7D* older flowers. Scalebar, 50 µm, *n* = 3.

(**C** and **D**) Expression pattern of auxin response reporter (*pDR5::VENUS*) in stage 11 flower of *as2-7D* mutant. Shown is (**C**) VENUS signal by itself (green) and (**D**) VENUS signal merged with PI (grey). Note the presence of DR5 signal specifically in the outgrowths (yellow arrowheads). Scalebar, 100 µm, *n* = 3.

(**E** to **H**) Representative live-imaging series showing the second replicate of (**E**) Mock treated, (**F**) 10µM NPA treated developing flowers, and third replicate of (**G**) Mock treated, (**H**) 10µM NPA treated developing flowers of *as2-7D* mutant. Images taken at every 24-hour interval over 96-hour period. Note the development of outgrowths from 72-hour time-point onwards (yellow arrowheads) in the mock treated *as2-7D* samples (panels E and G), which are absent in NPA treated *as2-7D* flowers (panels F and H). The *as2-7D* flowers harbor pLH13 (*35S::mCitrine-RCI2A*) cell membrane marker (greyscale). Trichomes were removed from the outer sepal epidermis by image processing. Scalebar, 50 µm.

Supplementary Table S1: List of primers used in this study.

Supplementary Data S1: Raw data for making the plots

Supplementary Movie S1: Confocal images of the outer and inner epidermis of adult sepals of the wild type (WT, left) and *as2-7D* mutant (right). Genotypes harbor pLH13 (*p35S::mCitrine-RCI2A*) plasma membrane marker. Scalebar, 200 µm.

Supplementary Movie S2: Confocal images of developing wild-type (WT, left) and *as2-7D* mutant (right) flowers imaged every 24 hours, over 96-hour time period. Genotypes harbor pLH13 (*p35S::mCitrine-RCI2A*) plasma membrane marker. Trichomes were removed from the outer sepal epidermis by image processing for better visualization of sepal morphology. Scalebar, 50 µm.

Supplementary Movie S3: Confocal images show the outer epidermis of adult sepals of wild type (WT, top left), *as2-7D* (top right), *pML1::KRP1* (bottom left) and *pML1::KRP1 as2-7D* (bottom right). Note the largely rescued sepal smoothness in *pML1::KRP1 as2-7D* compared to *as2-7D*, due to an increase in the longitudinally aligned giant cells. Sepals were stained with Propidium-iodide, which stains the cell walls. Scalebar, 200 µm.

Supplementary Movie S4: Confocal images of developing *as2-7D* mutant flowers grown in the presence of mock (left) or 10µM NPA (right). Images were captured every 24 hours over the 96-hour time-period. The *as2-7D* mutant harbors pLH13 (*p35S::mCitrine-RCI2A*) plasma membrane marker. Trichomes were removed from the outer sepal epidermis by image processing. Note the reduction in outgrowth formation in 10µM NPA-treated samples despite buckling. Scalebar, 50 µm.

REFERENCES AND NOTES

1.   Richman, D.P., Stewart, R.M., Hutchinson, J.W., and Caviness, V.S., Jr. (1975). Mechanical model of brain convolutional development. Science *189*, 18-21. 10.1126/science.1135626.
2.   Hannezo, E., Prost, J., and Joanny, J.F. (2011). Instabilities of monolayered epithelia: shape and structure of villi and crypts. Phys Rev Lett *107*, 078104. 10.1103/PhysRevLett.107.078104.
3.   Savin, T., Kurpios, N.A., Shyer, A.E., Florescu, P., Liang, H., Mahadevan, L., and Tabin, C.J. (2011). On the growth and form of the gut. Nature *476*, 57-62. 10.1038/nature10277.

4.   Shyer, A.E., Tallinen, T., Nerurkar, N.L., Wei, Z., Gil, E.S., Kaplan, D.L., Tabin, C.J.,
and Mahadevan, L. (2013). Villification: how the gut gets its villi. Science *342*, 212-218.
10.1126/science.1238842.
5.   Antoniou Kourounioti, R.L., Band, L.R., Fozard, J.A., Hampstead, A., Lovrics, A.,
Moyroud, E., Vignolini, S., King, J.R., Jensen, O.E., and Glover, B.J. (2013). Buckling as
an origin of ordered cuticular patterns in flower petals. Journal of The Royal Society
Interface *10*, 20120847. doi:10.1098/rsif.2012.0847.
6.   Moyroud, E., Airoldi, C.A., Ferria, J., Giorio, C., Steimer, S.S., Rudall, P.J., Prychid,
C.J., Halliwell, S., Walker, J.F., Robinson, S., et al. (2022). Cuticle chemistry drives the
development of diffraction gratings on the surface of Hibiscus trionum petals. Current
Biology *32*, 5323-5334.e5326. https://doi.org/10.1016/j.cub.2022.10.065.
7.   Waddington, C.H. (1939). Preliminary Notes on the Development of the Wings in
Normal and Mutant Strains of Drosophila. Proc Natl Acad Sci U S A *25*, 299-307.
10.1073/pnas.25.7.299.
8.   Thomson, J.A. (1917). On Growth and Form. Nature *100*, 21-22. 10.1038/100021a0.
9.   Tsukaya, H. (2005). Leaf shape: genetic controls and environmental factors. Int J Dev
Biol *49*, 547-555. 10.1387/ijdb.041921ht.
10.   Tsukaya, H. (2006). Mechanism of leaf-shape determination. Annu Rev Plant Biol *57*,
477-496. 10.1146/annurev.arplant.57.032905.105320.
11.   Boudaoud, A. (2010). An introduction to the mechanics of morphogenesis for plant
biologists. Trends Plant Sci *15*, 353-360. 10.1016/j.tplants.2010.04.002.
12.   Li, B., Cao, Y.-P., Feng, X.-Q., and Gao, H. (2012). Mechanics of morphological
instabilities and surface wrinkling in soft materials: a review. Soft Matter *8*, 5728-5745.
10.1039/C2SM00011C.
13.   Nelson, C.M. (2016). On Buckling Morphogenesis. J Biomech Eng *138*, 021005.
10.1115/1.4032128.
14.   Borowska-Wykręt, D., and Kwiatkowska, D. (2018). Folding, Wrinkling, and Buckling
in Plant Cell Walls. In Plant Biomechanics: From Structure to Function at Multiple
Scales, A. Geitmann, and J. Gril, eds. (Springer International Publishing), pp. 209-233.
10.1007/978-3-319-79099-2_10.
15.   Shyer, A.E., Rodrigues, A.R., Schroeder, G.G., Kassianidou, E., Kumar, S., and Harland,
R.M. (2017). Emergent cellular self-organization and mechanosensation initiate follicle
pattern in the avian skin. Science *357*, 811-815. 10.1126/science.aai7868.
16.   Varner, V.D., Gleghorn, J.P., Miller, E., Radisky, D.C., and Nelson, C.M. (2015).
Mechanically patterning the embryonic airway epithelium. Proceedings of the National
Academy of Sciences *112*, 9230-9235. doi:10.1073/pnas.1504102112.
17.   Maroudas-Sacks, Y., and Keren, K. (2021). Mechanical Patterning in Animal
Morphogenesis. Annual Review of Cell and Developmental Biology *37*, 469-493.
https://doi.org/10.1146/annurev-cellbio-120319-030931.

18.   Silverberg, J.L., Noar, R.D., Packer, M.S., Harrison, M.J., Henley, C.L., Cohen, I., and
1877   Gerbode, S.J. (2012). 3D imaging and mechanical modeling of helical buckling in
1878   Medicago truncatula plant roots. Proc Natl Acad Sci U S A *109*, 16794-16799.
1879   10.1073/pnas.1209287109.
1880   19.   Yan, J., Wang, B., Zhou, Y., and Hao, S. (2018). Resistance from agar medium impacts
1881   the helical growth of Arabidopsis primary roots. Journal of the Mechanical Behavior of
1882   Biomedical Materials *85*, 43-50. https://doi.org/10.1016/j.jmbbm.2018.05.018.
1883   20.   Sachse, R., Westermeier, A., Mylo, M., Nadasdi, J., Bischoff, M., Speck, T., and
1884   Poppinga, S. (2020). Snapping mechanics of the Venus flytrap (Dionaea muscipula). Proc
1885   Natl Acad Sci U S A *117*, 16035-16042. 10.1073/pnas.2002707117.
1886   21.   Forterre, Y., Skotheim, J.M., Dumais, J., and Mahadevan, L. (2005). How the Venus
1887   flytrap snaps. Nature *433*, 421-425. 10.1038/nature03185.
1888   22.   Airoldi, C.A., Lugo, C.A., Wightman, R., Glover, B.J., and Robinson, S. (2021).
1889   Mechanical buckling can pattern the light-diffracting cuticle of Hibiscus trionum. Cell
1890   Reports *36*, 109715. https://doi.org/10.1016/j.celrep.2021.109715.
1891   23.   Green, P.B. (1992). Pattern Formation in Shoots: A Likely Role for Minimal Energy
1892   Configurations of the Tunica. International Journal of Plant Sciences *153*, S59-S75.
1893   10.1086/297064.
1894   24.   Dumais, J., and Steele, C.R. (2000). New Evidence for the Role of Mechanical Forces in
1895   the Shoot Apical Meristem. Journal of Plant Growth Regulation *19*, 7-18.
1896   10.1007/s003440000003.
1897   25.   Rambaud-Lavigne, L., Chatterjee, A., Bovio, S., Battu, V., Lavigne, Q., Gundiah, N.,
1898   Boudaoud, A., and Das, P. (2024). Heterogeneous identity, stiffness and growth
1899   characterise the shoot apex of Arabidopsis stem cell mutants. Development *151*.
1900   10.1242/dev.202810.
1901   26.   Żebrowski, J. (1999). Dynamic behaviour of inflorescence-bearing Triticale and Triticum
1902   stems. Planta *207*, 410-417.
1903   27.   Fournier, M., Dlouhá, J., Jaouen, G., and Almeras, T. (2013). Integrative biomechanics
1904   for tree ecology: beyond wood density and strength. Journal of Experimental Botany *64*,
1905   4793-4815. 10.1093/jxb/ert279.
1906   28.   Chapotin, S.M., Razanameharizaka, J.H., and Holbrook, N.M. (2006). A biomechanical
1907   perspective on the role of large stem volume and high water content in baobab trees
1908   (Adansonia spp.; Bombacaceae). Am J Bot *93*, 1251-1264. 10.3732/ajb.93.9.1251.
1909   29.   Méndez-Alonzo, R., Moctezuma, C., Ordoñez, V.R., Angeles, G., Martínez, A.J., and
1910   López-Portillo, J. (2015). Root biomechanics in Rhizophora mangle: anatomy,
1911   morphology and ecology of mangrove's flying buttresses. Annals of Botany *115*, 833-
1912   840. 10.1093/aob/mcv002.
1913   30.   Huang, C., Wang, Z., Quinn, D., Suresh, S., and Hsia, K.J. (2018). Differential growth
1914   and shape formation in plant organs. Proceedings of the National Academy of Sciences
1915   *115*, 12359-12364. doi:10.1073/pnas.1811296115.

31. Derr, J., Bastien, R., Couturier, É., and Douady, S. (2018). Fluttering of growing leaves as a way to reach flatness: experimental evidence on *Persea americana*. Journal of The Royal Society Interface *15*, 20170595. doi:10.1098/rsif.2017.0595.

32. Wang, T., Fu, C., Potier-Ferry, M., and Xu, F. (2024). Morphomechanics of growing curled petals and leaves. Journal of the Mechanics and Physics of Solids *184*, 105534. https://doi.org/10.1016/j.jmps.2023.105534.

33. Whitewoods, C.D. (2021). Riddled with holes: Understanding air space formation in plant leaves. PLoS Biol *19*, e3001475. 10.1371/journal.pbio.3001475.

34. Evans, J.R. (2013). Improving photosynthesis. Plant Physiol *162*, 1780-1793. 10.1104/pp.113.219006.

35. Roeder, A.H.K. (2021). Arabidopsis sepals: A model system for the emergent process of morphogenesis. Quantitative Plant Biology *2*, e14, e14. 10.1017/qpb.2021.12.

36. Hong, L., Dumond, M., Tsugawa, S., Sapala, A., Routier-Kierzkowska, A.L., Zhou, Y., Chen, C., Kiss, A., Zhu, M., Hamant, O., et al. (2016). Variable Cell Growth Yields Reproducible OrganDevelopment through Spatiotemporal Averaging. Dev Cell *38*, 15-32. 10.1016/j.devcel.2016.06.016.

37. Iwakawa, H., Ueno, Y., Semiarti, E., Onouchi, H., Kojima, S., Tsukaya, H., Hasebe, M., Soma, T., Ikezaki, M., Machida, C., and Machida, Y. (2002). The ASYMMETRIC LEAVES2 gene of Arabidopsis thaliana, required for formation of a symmetric flat leaf lamina, encodes a member of a novel family of proteins characterized by cysteine repeats and a leucine zipper. Plant Cell Physiol *43*, 467-478. 10.1093/pcp/pcf077.

38. Bowman, J.L., Eshed, Y., and Baum, S.F. (2002). Establishment of polarity in angiosperm lateral organs. Trends Genet *18*, 134-141. 10.1016/s0168-9525(01)02601-4.

39. Lin, W.C., Shuai, B., and Springer, P.S. (2003). The Arabidopsis LATERAL ORGAN BOUNDARIES-domain gene ASYMMETRIC LEAVES2 functions in the repression of KNOX gene expression and in adaxial-abaxial patterning. Plant Cell *15*, 2241-2252. 10.1105/tpc.014969.

40. Scacchi, E., Paszkiewicz, G., Thi Nguyen, K., Meda, S., Burian, A., de Back, W., and Timmermans, M.C.P. (2024). A diffusible small-RNA-based Turing system dynamically coordinates organ polarity. Nature Plants *10*, 412-422. 10.1038/s41477-024-01634-x.

41. Wu, G., Lin, W.C., Huang, T., Poethig, R.S., Springer, P.S., and Kerstetter, R.A. (2008). KANADI1 regulates adaxial-abaxial polarity in Arabidopsis by directly repressing the transcription of ASYMMETRIC LEAVES2. Proc Natl Acad Sci U S A *105*, 16392-16397. 10.1073/pnas.0803997105.

42. Eshed, Y., Baum, S.F., Perea, J.V., and Bowman, J.L. (2001). Establishment of polarity in lateral organs of plants. Curr Biol *11*, 1251-1260. 10.1016/s0960-9822(01)00392-x.

43. Kerstetter, R.A., Bollman, K., Taylor, R.A., Bomblies, K., and Poethig, R.S. (2001). KANADI regulates organ polarity in Arabidopsis. Nature *411*, 706-709. 10.1038/35079629.

44. Lin, X., Gu, D., Zhao, H., Peng, Y., Zhang, G., Yuan, T., Li, M., Wang, Z., Wang, X.,
and Cui, S. (2018). LFR is functionally associated with AS2 to mediate leaf development
in Arabidopsis. Plant J. 10.1111/tpj.13973.
45. Hervieux, N., Dumond, M., Sapala, A., Routier-Kierzkowska, A.L., Kierzkowski, D.,
Roeder, A.H., Smith, R.S., Boudaoud, A., and Hamant, O. (2016). A Mechanical
Feedback Restricts Sepal Growth and Shape in Arabidopsis. Curr Biol.
10.1016/j.cub.2016.03.004.
46. Onoda, Y., Schieving, F., and Anten, N.P. (2015). A novel method of measuring leaf
epidermis and mesophyll stiffness shows the ubiquitous nature of the sandwich structure
of leaf laminas in broad-leaved angiosperm species. J Exp Bot *66*, 2487-2499.
10.1093/jxb/erv024.
47. Vijayan, A., Tofanelli, R., Strauss, S., Cerrone, L., Wolny, A., Strohmeier, J., Kreshuk,
A., Hamprecht, F.A., Smith, R.S., and Schneitz, K. (2021). A digital 3D reference atlas
reveals cellular growth patterns shaping the Arabidopsis ovule. eLife *10*, e63262.
10.7554/eLife.63262.
48. Mollier, C., Skrzydeł, J., Borowska-Wykręt, D., Majda, M., Bayle, V., Battu, V.,
Totozafy, J.C., Dulski, M., Fruleux, A., Wrzalik, R., et al. (2023). Spatial consistency of
cell growth direction during organ morphogenesis requires CELLULOSE SYNTHASE
INTERACTIVE1. Cell Rep *42*, 112689. 10.1016/j.celrep.2023.112689.
49. Harmansa, S., Erlich, A., Eloy, C., Zurlo, G., and Lecuit, T. (2023). Growth anisotropy of
the extracellular matrix shapes a developing organ. Nature Communications *14*, 1220.
10.1038/s41467-023-36739-y.
50. Hong, L., Dumond, M., Zhu, M., Tsugawa, S., Li, C.-B., Boudaoud, A., Hamant, O., and
Roeder, A.H.K. (2018). Heterogeneity and Robustness in Plant Morphogenesis: From
Cells to Organs. Annual Review of Plant Biology *69*, 469-495. 10.1146/annurev-arplant-
042817-040517.
51. Sylvester, A.W., Williams, M.H., and Green, P.B. (1989). Orientation of cortical
microtubules correlates with cell shape and division direction. Protoplasma *153*, 91-103.
10.1007/BF01322469.
52. Hamant, O., Heisler, M.G., Jönsson, H., Krupinski, P., Uyttewaal, M., Bokov, P., Corson,
F., Sahlin, P., Boudaoud, A., Meyerowitz, E.M., et al. (2008). Developmental patterning
by mechanical signals in Arabidopsis. Science *322*, 1650-1655.
10.1126/science.1165594.
53. Roeder, A.H., Chickarmane, V., Cunha, A., Obara, B., Manjunath, B.S., and Meyerowitz,
E.M. (2010). Variability in the control of cell division underlies sepal epidermal
patterning in Arabidopsis thaliana. PLoS Biol *8*, e1000367.
10.1371/journal.pbio.1000367.
54. Bemis, S.M., and Torii, K.U. (2007). Autonomy of cell proliferation and developmental
programs during Arabidopsis aboveground organ morphogenesis. Dev Biol *304*, 367-381.
10.1016/j.ydbio.2006.12.049.

55.   Sapala, A., and Smith, R.S. (2020). Osmotic Treatment for Quantifying Cell Wall
1996   Elasticity in the Sepal of Arabidopsis thaliana. Methods Mol Biol *2094*, 101-112.
1997   10.1007/978-1-0716-0183-9_11.
1998   56.   Kierzkowski, D., Nakayama, N., Routier-Kierzkowska, A.L., Weber, A., Bayer, E.,
1999   Schorderet, M., Reinhardt, D., Kuhlemeier, C., and Smith, R.S. (2012). Elastic domains
2000   regulate growth and organogenesis in the plant shoot apical meristem. Science *335*, 1096-
2001   1099. 10.1126/science.1213100.
2002   57.   Waites, R., and Hudson, A. (1995). phantastica: a gene required for dorsoventrality of
2003   leaves in Antirrhinum majus. Development *121*, 2143-2154. 10.1242/dev.121.7.2143.
2004   58.   Fukushima, K., and Hasebe, M. (2014). Adaxial-abaxial polarity: the developmental basis
2005   of leaf shape diversity. Genesis *52*, 1-18. 10.1002/dvg.22728.
2006   59.   Izhaki, A., and Bowman, J.L. (2007). KANADI and Class III HD-Zip Gene Families
2007   Regulate Embryo Patterning and Modulate Auxin Flow during Embryogenesis in
2008   Arabidopsis. The Plant Cell *19*, 495-508. 10.1105/tpc.106.047472.
2009   60.   Eshed, Y., Izhaki, A., Baum, S.F., Floyd, S.K., and Bowman, J.L. (2004). Asymmetric
2010   leaf development and blade expansion in Arabidopsis are mediated by KANADI and
2011   YABBY activities. Development *131*, 2997-3006. 10.1242/dev.01186.
2012   61.   Caggiano, M.P., Yu, X., Bhatia, N., Larsson, A., Ram, H., Ohno, C.K., Sappl, P.,
2013   Meyerowitz, E.M., Jönsson, H., and Heisler, M.G. (2017). Cell type boundaries organize
2014   plant development. eLife *6*, e27421. 10.7554/eLife.27421.
2015   62.   Nakata, M., and Okada, K. (2012). The three-domain model: a new model for the early
2016   development of leaves in Arabidopsis thaliana. Plant Signal Behav *7*, 1423-1427.
2017   10.4161/psb.21959.
2018   63.   Guan, C., Wu, B., Yu, T., Wang, Q., Krogan, N.T., Liu, X., and Jiao, Y. (2017). Spatial
2019   Auxin Signaling Controls Leaf Flattening in Arabidopsis. Current Biology *27*, 2940-
2020   2950.e2944. https://doi.org/10.1016/j.cub.2017.08.042.
2021   64.   Otsuga, D., DeGuzman, B., Prigge, M.J., Drews, G.N., and Clark, S.E. (2001).
2022   REVOLUTA regulates meristem initiation at lateral positions. Plant J *25*, 223-236.
2023   10.1046/j.1365-313x.2001.00959.x.
2024   65.   Watanabe, K., and Okada, K. (2003). Two discrete cis elements control the Abaxial side-
2025   specific expression of the FILAMENTOUS FLOWER gene in Arabidopsis. Plant Cell
2026   *15*, 2592-2602. 10.1105/tpc.015214.
2027   66.   Semiarti, E., Ueno, Y., Tsukaya, H., Iwakawa, H., Machida, C., and Machida, Y. (2001).
2028   The ASYMMETRIC LEAVES2 gene of Arabidopsis thaliana regulates formation of a
2029   symmetric lamina, establishment of venation and repression of meristem-related
2030   homeobox genes in leaves. Development *128*, 1771-1783. 10.1242/dev.128.10.1771.
2031   67.   Gälweiler, L., Guan, C., Müller, A., Wisman, E., Mendgen, K., Yephremov, A., and
2032   Palme, K. (1998). Regulation of polar auxin transport by AtPIN1 in Arabidopsis vascular
2033   tissue. Science *282*, 2226-2230. 10.1126/science.282.5397.2226.

68. Abley, K., Sauret-Güeto, S., Marée, A.F., and Coen, E. (2016). Formation of polarity
convergences underlying shoot outgrowths. Elife *5*. 10.7554/eLife.18165.
69. Reinhardt, D., Pesce, E.-R., Stieger, P., Mandel, T., Baltensperger, K., Bennett, M.,
Traas, J., Friml, J., and Kuhlemeier, C. (2003). Regulation of phyllotaxis by polar auxin
transport. Nature *426*, 255-260. 10.1038/nature02081.
70. Heisler, M.G., Hamant, O., Krupinski, P., Uyttewaal, M., Ohno, C., Jönsson, H., Traas,
J., and Meyerowitz, E.M. (2010). Alignment between PIN1 polarity and microtubule
orientation in the shoot apical meristem reveals a tight coupling between morphogenesis
and auxin transport. PLoS Biol *8*, e1000516. 10.1371/journal.pbio.1000516.
71. Hamant, O., Meyerowitz, E.M., and Traas, J. (2011). Is cell polarity under mechanical
control in plants? Plant Signal Behav *6*, 137-139. 10.4161/psb.6.1.14269.
72. Nakayama, N., Smith, Richard S., Mandel, T., Robinson, S., Kimura, S., Boudaoud, A.,
and Kuhlemeier, C. (2012). Mechanical Regulation of Auxin-Mediated Growth. Current
Biology *22*, 1468-1476. https://doi.org/10.1016/j.cub.2012.06.050.
73. Kabir, A.M.R., Inoue, D., Afrin, T., Mayama, H., Sada, K., and Kakugo, A. (2015).
Buckling of Microtubules on a 2D Elastic Medium. Scientific Reports *5*, 17222.
10.1038/srep17222.
74. Sanyal, A., and Han, H.C. (2015). Artery buckling affects the mechanical stress in
atherosclerotic plaques. Biomed Eng Online *14 Suppl 1*, S4. 10.1186/1475-925x-14-s1-
s4.
75. Roeder, A.H.K. (2021). Arabidopsis sepals: A model system for the emergent process of
morphogenesis. Quant Plant Biol *2*. 10.1017/qpb.2021.12.
76. Xu, L., Xu, Y., Dong, A., Sun, Y., Pi, L., Xu, Y., and Huang, H. (2003). Novel as1 and
as2 defects in leaf adaxial-abaxial polarity reveal the requirement for ASYMMETRIC
LEAVES1 and 2 and ERECTA functions in specifying leaf adaxial identity.
Development *130*, 4097-4107. 10.1242/dev.00622.
77. Friesen, S., and Hariharan, I.K. (2023). Coordinated growth of linked epithelia is
mediated by the Hippo pathway. bioRxiv, 2023.2002.2026.530099.
10.1101/2023.02.26.530099.
78. Kelly-Bellow, R., Lee, K., Kennaway, R., Barclay, J.E., Whibley, A., Bushell, C.,
Spooner, J., Yu, M., Brett, P., Kular, B., et al. (2023). Brassinosteroid coordinates cell
layer interactions in plants via cell wall and tissue mechanics. Science *380*, 1275-1281.
doi:10.1126/science.adf0752.
79. Du, J., Kirui, A., Huang, S., Wang, L., Barnes, W.J., Kiemle, S.N., Zheng, Y., Rui, Y.,
Ruan, M., Qi, S., et al. (2020). Mutations in the Pectin Methyltransferase
QUASIMODO2 Influence Cellulose Biosynthesis and Wall Integrity in Arabidopsis. The
Plant Cell *32*, 3576-3597. 10.1105/tpc.20.00252.
80. Qi, J., Wu, B., Feng, S., Lü, S., Guan, C., Zhang, X., Qiu, D., Hu, Y., Zhou, Y., Li, C., et
al. (2017). Mechanical regulation of organ asymmetry in leaves. Nat Plants *3*, 724-733.
10.1038/s41477-017-0008-6.

81. Enugutti, B., Kirchhelle, C., and Schneitz, K. (2013). On the genetic control of planar
growth during tissue morphogenesis in plants. Protoplasma *250*, 651-661.
10.1007/s00709-012-0452-0.
82. Braybrook, S.A., and Peaucelle, A. (2013). Mechano-chemical aspects of organ
formation in Arabidopsis thaliana: the relationship between auxin and pectin. PLoS One
8, e57813. 10.1371/journal.pone.0057813.
83. Kuhlemeier, C. (2007). Phyllotaxis. Trends Plant Sci *12*, 143-150.
10.1016/j.tplants.2007.03.004.
84. Yadav, A.S., and Roeder, A.H.K. (2024). An optimized live imaging and growth analysis
approach for Arabidopsis Sepals. bioRxiv, 2024.2001.2022.576735.
10.1101/2024.01.22.576735.
85. Routier-Kierzkowska, A.L., and Smith, R.S. (2013). Measuring the mechanics of
morphogenesis. Curr Opin Plant Biol *16*, 25-32. 10.1016/j.pbi.2012.11.002.
86. Liu, S., Strauss, S., Adibi, M., Mosca, G., Yoshida, S., Dello Ioio, R., Runions, A.,
Andersen, T.G., Grossmann, G., Huijser, P., et al. (2022). Cytokinin promotes growth
cessation in the Arabidopsis root. Current Biology *32*, 1974-1985.e1973.
https://doi.org/10.1016/j.cub.2022.03.019.
87. Natonik-Białoń, S., Borowska-Wykręt, D., Mosca, G., Grelowski, M., Wrzalik, R.,
Smith, R.S., and Kwiatkowska, D. (2020). Deformation of a cell monolayer due to
osmotic treatment: a case study of onion scale epidermis. Botany *98*, 21-36. 10.1139/cjb-
2019-0027.
88. Smyth, D.R., Bowman, J.L., and Meyerowitz, E.M. (1990). Early flower development in
Arabidopsis. The Plant Cell *2*, 755-767. 10.1105/tpc.2.8.755.
89. Robinson, D.O., Coate, J.E., Singh, A., Hong, L., Bush, M., Doyle, J.J., and Roeder,
A.H.K. (2018). Ploidy and Size at Multiple Scales in the Arabidopsis Sepal. The Plant
Cell *30*, 2308-2329. 10.1105/tpc.18.00344.
90. Strauss, S., Runions, A., Lane, B., Eschweiler, D., Bajpai, N., Trozzi, N., Routier-
Kierzkowska, A.-L., Yoshida, S., Rodrigues da Silveira, S., Vijayan, A., et al. (2022).
Using positional information to provide context for biological image analysis with
MorphoGraphX 2.0. eLife *11*, e72601. 10.7554/eLife.72601.
91. Armezzani, A., Abad, U., Ali, O., Andres Robin, A., Vachez, L., Larrieu, A.,
Mellerowicz, E.J., Taconnat, L., Battu, V., Stanislas, T., et al. (2018). Transcriptional
induction of cell wall remodelling genes is coupled to microtubule-driven growth
isotropy at the shoot apex in Arabidopsis. Development *145*. 10.1242/dev.162255.
92. Heisler, M.G., Ohno, C., Das, P., Sieber, P., Reddy, G.V., Long, J.A., and Meyerowitz,
E.M. (2005). Patterns of Auxin Transport and Gene Expression during Primordium
Development Revealed by Live Imaging of the Arabidopsis Inflorescence Meristem.
Current Biology *15*, 1899-1911. https://doi.org/10.1016/j.cub.2005.09.052.

93. Eshed, Y., Baum, S.F., Perea, J.V., and Bowman, J.L. (2001). Establishment of polarity
in lateral organs of plants. Current Biology *11*, 1251-1260. 10.1016/S0960-
9822(01)00392-X.
94. Hong, L., Brown, J., Segerson, N.A., Rose, J.K., and Roeder, A.H. (2017). CUTIN
SYNTHASE 2 Maintains Progressively Developing Cuticular Ridges in Arabidopsis
Sepals. Mol Plant *10*, 560-574. 10.1016/j.molp.2017.01.002.
95. Roeder, A.H., Cunha, A., Ohno, C.K., and Meyerowitz, E.M. (2012). Cell cycle regulates
cell type in the Arabidopsis sepal. Development *139*, 4416-4427. 10.1242/dev.082925.
96. Lukowitz, W., Gillmor, C.S., and Scheible, W.-R.d. (2000). Positional Cloning in
Arabidopsis. Why It Feels Good to Have a Genome Initiative Working for You1. Plant
Physiology *123*, 795-806. 10.1104/pp.123.3.795.
97. Robinson, D.O., Coate, J.E., Singh, A., Hong, L., Bush, M., Doyle, J.J., and Roeder,
A.H.K. (2018). Ploidy and Size at Multiple Scales in the Arabidopsis Sepal. Plant Cell
30, 2308-2329. 10.1105/tpc.18.00344.
98. Karimi, M., Inze, D., and Depicker, A. (2002). GATEWAY vectors for Agrobacterium-
mediated plant transformation. Trends Plant Sci *7*, 193-195.
99. Zhu, M., Chen, W., Mirabet, V., Hong, L., Bovio, S., Strauss, S., Schwarz, E.M.,
Tsugawa, S., Wang, Z., Smith, R.S., et al. (2020). Robust organ size requires robust
timing of initiation orchestrated by focused auxin and cytokinin signalling. Nat Plants *6*,
686-698. 10.1038/s41477-020-0666-7.
100. Barrell, P.J., and Conner, A.J. (2006). Minimal T-DNA vectors suitable for agricultural
deployment of transgenic plants. BioTechniques *41*, 708-710. 10.2144/000112306.
101. Wise, A.A., Liu, Z., and Binns, A.N. (2006). Three methods for the introduction of
foreign DNA into Agrobacterium. Methods Mol Biol *343*, 43-53. 10.1385/1-59745-130-
4:43.
102. Zhu, M., and Roeder, A.H. (2020). Plants are better engineers: the complexity of plant
organ morphogenesis. Curr Opin Genet Dev *63*, 16-23. 10.1016/j.gde.2020.02.008.
103. Kiss, A., Moreau, T., Mirabet, V., Calugaru, C.I., Boudaoud, A., and Das, P. (2017).
Segmentation of 3D images of plant tissues at multiple scales using the level set method.
Plant Methods *13*, 114. 10.1186/s13007-017-0264-5.
104. Huang, Z.Y., Hong, W., and Suo, Z. (2005). Nonlinear analyses of wrinkles in a film
bonded to a compliant substrate. Journal of the Mechanics and Physics of Solids *53*,
2101-2118. https://doi.org/10.1016/j.jmps.2005.03.007.
105. Audoly, B., and Boudaoud, A. (2008). Buckling of a stiff film bound to a compliant
substrate—Part I:: Formulation, linear stability of cylindrical patterns, secondary
bifurcations. Journal of the Mechanics and Physics of Solids *56*, 2401-2421.
https://doi.org/10.1016/j.jmps.2008.03.003.
106. Goriely, A. (2016). Morpho-elasticity: The mechanics and mathematics of biological
growth (Springer New York).

107.   Alnæs, M., Blechta, J., Hake, J., Johansson, A., Kehlet, B., Logg, A., Richardson, C.,
2152   Ring, J., Rognes, M., and Wells, G. (2015). The FEniCS project version 1.5. *3*.
2153   10.11588/ans.2015.100.20553.
2154   108.   Gacon, F., Godin, C., and Ali, O. (2021). BVPy: A FEniCS-based Python package to
2155   ease the expression and study of boundary value problems in Biology. Journal of Open
2156   Source Software *6*, 1-6. 10.21105/joss.02831.
2157