1 **Growth directions and stiffness across cell layers determine whether tissues**
2 **stay smooth or buckle**
3

4 Avilash S. Yadav[2]†, Lilan Hong[1]†, Patrick M. Klees[2], Annamaria Kiss[3], Xi He[1], Iselle M.
5 Barrios[2], Michelle Heeney[2], Anabella Maria D. Galang[2], Richard S. Smith[4], Arezki Boudaoud[5],
6 and Adrienne H.K. Roeder[2]*
7

[1]Institute of Nuclear Agricultural Sciences, Key Laboratory of Nuclear Agricultural Sciences of
Ministry of Agriculture and Zhejiang Province, College of Agriculture and Biotechnology,
Zhejiang University, Hangzhou 310058, China
[2]Weill Institute for Cell and Molecular Biology and Section of Plant Biology, School of
Integrative Plant Sciences, Cornell University, Ithaca, NY 14853, USA
[3]Laboratoire de Reproduction et Développement des Plantes, Université de Lyon, ENS de Lyon,
UCBL, INRA, CNRS, 46 Allée d'Italie, 69364 Lyon Cedex 07, France
[4]The John Innes Centre, Norwich NR4 7UH, UK
[5]LadHyX, Ecole Polytechnique, CNRS, IP Paris, 91128 Palaiseau Cedex, France
†These authors contributed equally to this work.
*Corresponding author. ahr75@cornell.edu

Abstract

From smooth to buckled, nature exhibits organs of various shapes and forms. How cellular
growth patterns produce smooth organ shapes like those of leaves and sepals remains unclear.
Here we show that unidirectional growth and comparable stiffness across both epidermal layers
of Arabidopsis sepals are essential for smoothness. We identified a mutant with ectopic
ASYMMETRIC LEAVES 2 (*AS2*) expression that exhibits buckles on the outer epidermis due to
conflicting growth directions and unequal epidermal stiffnesses. Aligning growth direction and
increasing stiffness restores smoothness. Furthermore, buckling generates outgrowth by
influencing auxin efflux transporter protein PIN-FORMED 1 polarity, indicating buckling can
initiate organogenesis. Our findings suggest that in addition to molecular cues influencing tissue
mechanics, tissue mechanics can also modulate molecular signals, giving rise to well-defined
shapes.

One-Sentence Summary
The keys to unlocking sepal smoothness are unidirectional growth and even stiffness; misplace
them, it's time to buckle up!
35
36
37

INTRODUCTION

During development, many organs such as the brain, intestine, and orchid flower petals form complex shapes with undulations, folds, and loops (*1-5*); conversely, other organs such as Arabidopsis leaves and fly wings develop smooth forms (*6-9*). These undulations, folds, and loops generally form through mechanical buckling (*10-12*), which is the bending of tissue when its growth patterns generate compressive forces (*13-16*). How cells within tissues regulate their growth and division to generate buckling in certain instances and maintain smoothness in others is not well understood.

Buckling plays a significant role in organ morphogenesis and development across organisms. In animals for instance, numerous studies demonstrate the necessity of buckling in morphogenesis (*2-4, 17-21*). Differential growth of connected tissue layers gives rise to buckling. Buckling creates complex organ shapes starting from simple geometries, which is important for functional specialization. These include gyrification of the cerebral cortex to increase neuron content (*2*), formation of gut villi and looping to increase contact with digested food and length of the gastrointestinal tract (*3, 4, 19, 20*), and initiation of new branches in lung airway epithelium to enhance gas exchange and oxygen transfer (*18*). Buckling also serves key roles in plant development. For instance, it generates helical twists in roots to enhance penetrative forces at the growing tip (*22-25*), triggers lobe deformation in Venus flytraps for rapid energy storage and release to capture prey (*26, 27*), and induces nanoscale cuticular wrinkles on petals to produce iridescence (*28-30*). Buckling has also been proposed to drive formation and positioning of primordia on the shoot apical meristem (*31, 32*).

Buckling, however, is not always beneficial. Inflorescence weight and external forces such as wind or rain can compress and buckle the stems of cereal crops, thereby causing damage and reduced yield (*33*). Similarly, slender tree stems are also vulnerable to buckling (*34, 35*). To counter this, plants need to allocate resources to increase stem thickness and stiffness (*36-38*). While buckling generates diverse leaf shapes (*5, 39*), majority of the leaves suppress buckling, generating smooth shapes that enhance light absorption and gas exchange for efficient photosynthesis (*40, 41*). Similarly, sepals, which are the outermost leaf-like floral organs, grow to smooth shapes to properly enclose and protect the developing flower bud (*42*). Therefore, plants need to modulate growth and tissue properties of leaves and sepals to determine whether they remain smooth or buckle during development. In this study, we used a gain-of-function mutant of the transcription factor *ASYMMETRIC LEAVES 2* (*AS2*), *as2-7D,* as a tool to determine how cell growth patterns are regulated to prevent buckling, and thus produce smooth sepals in the wild-type (WT).

78   **RESULTS**
79
80   **Ectopic expression of *ASYMMETRIC LEAVES 2* (*AS2*) disrupts sepal smoothness**
81
To determine how the sepals achieve smooth morphology, we carried out a forward genetic
screen (*43*) and identified an *Arabidopsis thaliana* mutant that disrupts sepal smoothness. This
mutant exhibits an irregular morphology with folds and outgrowths (Fig. 1, A to F). The folds
comprise ridges and invaginations, whereas the outgrowths, which are pointed epidermal
projections, are typically located at the end of the folds (Fig. 1F and fig. S1, A and B). These
folds and outgrowths are visible only on the outer (abaxial) epidermis, while the inner (adaxial)
epidermis is smooth like the WT (Fig. 1, E to H, fig. S1, A and B and movie S1). Positional
cloning identified a G-to-A point mutation 1,484 bps upstream of the translational start site of
ASYMMETRIC LEAVES 2 (*AS2*) (Fig. 1I). *AS2* encodes a LATERAL ORGAN BOUNDARIES
(LOB) domain transcription factor that plays a critical role in leaf development, particularly in
specifying the upper (adaxial) layer of the leaf (*44-49*). An identical mutation in the same locus
was identified previously (*48*). However, the fold and outgrowth phenotype in the sepals was not
reported (*48*), since the phenotype is very subtle in the Columbia (Col-0) background compared
to what we observe in the Landsberg *erecta* (L*er*-0) background (hereafter referred to as WT)
(Fig. 1, A to D and fig. S1, C to F). Previously, Wu et. al. (*48*) demonstrated that the mutation in
the *AS2* promoter disrupts the binding site of the repressor KANADI, a transcription factor
involved in specifying cell fate on the lower (abaxial) side of the leaf (*50-53*). They further
showed that this mutation leads to ectopic *AS2* expression on the abaxial side, while normally in
the WT, the *AS2* expression domain is restricted to the adaxial side (Fig.1I). To verify that this
mutation also leads to ectopic *AS2* expression in sepals of the L*er*-0 background, we drove GFP
expression under the WT and mutant *AS2* promoters (*pAS2::GUS-GFP*) in WT. While the *AS2*
promoter from the WT drives GFP specifically on the inner sepal epidermal layer (Fig. 1, J and
K), the mutant *AS2* promoter drives GFP on both the inner and outer epidermal layers (Fig. 1L).
This confirms that the mutation causes a gain-of-function through ectopic *AS2* expression,
consistent with the dominant nature of this allele (see Materials and methods). Therefore,
following the convention of Wu et al. (*48, 54*), we named the mutant allele *as2-7D*.
108
To test whether ectopic *AS2* expression is sufficient to generate the folds and outgrowths, we
ectopically expressed *AS2* on both inner and outer epidermis of the WT (*pML1::AS2*). All the
transgenic plants form lumps and outgrowths on the sepal outer epidermis, similar to *as2-7D*
(Fig. 1, M to P). Additionally, we observe folds and outgrowths form on the epidermis of the
leaves and fruits (fig. S1, G to N), consistent with previous reports (*46*), indicating a potential
stronger effect of constitutive epidermal *AS2* expression than in the *as2-7D* gain-of-function
mutant. In summary, the mutation in *as2-7D* induces ectopic *AS2* expression on the outer
epidermal layer, which leads to disruption of smooth sepal morphology by generating folds and
outgrowths on the outer epidermal layer.

The *as2-7D* outer epidermis appears to buckle

To understand how ectopic *AS2* expression on the outer epidermis disrupts smooth sepal morphology, we live-imaged WT and *as2-7D* flowers from early sepal developmental stages (Fig. 2, A and B, fig. S2, A to D, and movie S2). Initially, both WT and *as2-7D* sepals are smooth (Fig. 2, A and B; 0-24h). While WT sepals remain smooth throughout, *as2-7D* starts to develop a fold in the middle of the sepal at 48 hours (Fig. 2, B and C). As the flower develops, folding intensifies further, forming several inward (invaginations) and outward (ridges) folds throughout the outer epidermis of *as2-7D* mutant (Fig. 2, B to D and movie S2). In addition to these folds, we also observe the development of phenotypically distinct pointed outgrowths. The outgrowths are not observed until the later stages when the folds intensify, suggesting they form downstream of folding (Fig. 2, B to D and fig. S2, C and D).

We considered two hypotheses for how the folds form: (1) due to localized growth underlying each ridge or (2) due to mechanical buckling caused by a surface-wide growth disruption. To test the first hypothesis, we compared the cell growth rates and division rates between WT and *as2-7D* mutant on the outer sepal epidermis every 48 hours over 96 hours (Fig. 2, E to H and fig. S2, E to N). We do not see any localized fast growth or cell division in the region where the first fold is formed in *as2-7D* at 48 hours (Fig. 2, C to F). In *as2-7D,* the cells grow at moderate rates throughout the epidermis (Fig. 2, E to F; 0-48 hrs). This pattern is also reflected in cell division rates, where we observe fast cell division throughout the sepal epidermis (Fig. 2, G and H; 0-48 hrs). This suggests that fold formation is a result of surface-wide disruption of cell growth patterns, and not because of localized cell growth and division. In the later time point (48-96 hrs) the growth is still relatively uniform along the places where the *as2-7D* buckles form; however, we do observe localized growth and division later at sites where outgrowths form (Fig. 2F and fig. S2, G and H; 48-96 hrs). We therefore considered the hypothesis that the folds on *as2-7D* outer epidermis form through mechanical buckling caused by surface-wide growth defects.

Analytical modeling predicts two criteria required for buckling

Based on our observations of *as2-7D* development, we built a simple two-dimensional (2D) analytical model to determine the conditions that can cause buckling (see Materials and methods). We consider a growing elastic system comprising three layers, corresponding to the inner epidermis, mesophyll, and outer epidermis of a sepal (Fig 2I). In the model, if the growth rate of the outer epidermis exceeds that of the inner epidermis, the tissue will curve (Fig. 2J). If the growth rate of the epidermal layers is sufficiently large compared to the mesophyll, then these two layers will be under compression, potentially causing buckling. If, in addition, the stiffness of the outer epidermis is less than the inner epidermis, the outer epidermis can, in theory, buckle, without any buckling of the inner epidermis (Fig. 2K). From the model and experimental observations, we predicted that (1) the growth of the outer epidermis of the *as2-7D*

158 mutant should exceed the inner epidermis and (2) the outer epidermis should be softer than the
159 inner epidermis.
160
161 **Synchrony of cell growth rate and directionality between epidermal layers is essential for**
162 **the formation of smooth sepals**
163
To test our model predictions, we live imaged WT and *as2-7D* sepals through pre-buckling
stages (Fig. 3, A and B), and extracted the inner and outer epidermal layers of the same growing
sepals by image processing (Fig. 3, C and D). We found that cells throughout the WT inner
epidermis grow significantly faster than the outer epidermal cells (Fig. 3E and fig. S3, A, B, and
E), overlapping with the cell division patterns (Fig. 3G and fig. S3, F, G, and J). This was
surprising, as intuitively for flat sepal growth, we expected the cells on both epidermal layers to
grow at similar rates. However, during the initial stages of sepal growth the sepal is highly
curved in the longitudinal direction, due to the smaller surface area of the inner epidermis. This
necessitates fast inner epidermal growth for achieving flatness (Fig. 3C and D, fig. S4F and
movie S2). Likewise, in *as2-7D*, inner epidermal cells grow faster than outer epidermal cells;
however, the inner epidermal cells of *as2-7D* do not grow as fast as the WT (Fig. 3, E and F and
fig. S3, A to E). In concert with the growth rate, the cell division rate is also lower on the inner
epidermis of *as2-7D* compared to the WT (Fig. 3, G and H and fig. S3, F to J). The fact that fast
growth of the inner epidermis is disrupted in *as2-7D* suggests that this fast inner epidermal
growth is an essential component of sepal smoothness.
179
Since buckling is a surface-wide phenomenon in our model, we tested the differences in whole
inner versus outer epidermal growth rates. We found that the outer epidermal layers of the WT
and *as2-7D* grow at the same rate, aligning with our cell growth rate data where the median
epidermal cell growth rates show no significant difference (fig. S3E and fig. S4, A, B and E).
However, the inner epidermis of *as2-7D* grows significantly slower than the WT (fig. S3E and
fig. S4, C, D and E). Our findings, therefore, indicate that the rate at which the inner epidermis of
as2-7D grows is insufficient to catch up with the increasing outer epidermal surface area (fig.
S4F).
188
Since cell growth directions have been shown to be critical for determining organ shapes (*55-*
57), we assessed the principal direction of maximal cell growth (PDG$_{max}$) in the WT and *as2-7D*
mutant. We found that the WT cells on both epidermal layers show highly aligned growth along
the sepal longitudinal axis (Fig. 3I). In *as2-7D*, although inner epidermal cells grow parallel to
the sepal longitudinal axis, the outer epidermal cells exhibit highly misaligned growth relative to
the longitudinal axis, with PDG$_{max}$ tensors oriented transversely (Fig. 3J). We quantified the cell
growth direction by the deviation angle of PDG$_{max}$ tensors relative to the sepal's longitudinal axis
(Fig. 3K). The heatmaps reveal minimal deviation in cell growth directions of the WT epidermal
layers and the inner epidermal layer of *as2-7D* (Fig. 3, L and M and fig. S3, K to O). In contrast,

the outer layer of *as2-7D* displayed a high degree of deviation in growth directions (Fig. 3M and
fig. S3, K to O). Since microtubules are known to align perpendicular to the maximal cell growth
direction (*58*), we then tested the microtubule orientation pattern in comparable regions of the
WT and *as2-7D* outer epidermis (Fig. 3, N and O). Microtubules show the typical transverse
alignment pattern in WT, which is perpendicular to sepal longitudinal growth direction (Fig. 3N).
Conversely, the microtubules are largely misoriented in the *as2-7D,* and in certain instances,
mirror the transverse cell growth pattern (yellow arrowheads, Fig. 3O). Taken together, our data
illustrates the complexity of biology beyond our simple 2D analytical model. Thus, we
hypothesize that in the *as2-7D* mutant, buckling occurs on the outer epidermis because it
overgrows in width while being constrained by the inner epidermis growing longitudinally. In
theory, transverse growth in the outer epidermis favors longitudinal buckles (*59*), consistent with
our observations. Overall, our data indicate that smooth sepal formation requires unidirectional
growth on both epidermal layers, along with fast growth of the inner layer to achieve flatness.
211

212 **Limiting transverse growth orientation and increasing stiffness of the outer epidermis**
213 **rescues *as2-7D* phenotype, by preventing buckling**
214

Having shown that *as2-7D* outer epidermal cells exhibit misaligned growth directions relative to
the sepal longitudinal axis, we hypothesized that the transversal growth orientation of the outer
epidermal cells causes buckling. Therefore, we tested whether prompting the cells to grow along
the longitudinal direction can rescue the *as2-7D* phenotype. Overexpressing cyclin-dependent
kinase (CDK) inhibitor Kip-related protein 1 (KRP1) throughout the epidermis (*pML1::KRP1*) at
moderate levels trigger epidermal cells to skip division and undergo endoreduplication early
during development, resulting in increased giant cell formation (*60, 61*). Giant cells, exclusive to
the outer epidermis, exhibit the unique property of growing in a highly directional manner (*60*)
along the longitudinal sepal axis (Fig. 4, A, C, E and G, fig. S5, A and C, and movie S3).
Therefore, we wondered whether increasing giant cell formation can restore directional growth
in *as2-7D* and consequently reduce buckling. Indeed, overexpressing *KRP1* in *as2-7D*
(*pML1::KRP1 as2-7D*) results in a striking smoothing of the outer sepal epidermis (Fig. 4, A to
H, fig. S5, A to D and movie S3). As expected, overexpressing KRP1 increases the number and
size of the giant cells, which grow in a highly aligned manner along the longitudinal sepal
direction (Fig. 4, E to P and movie S3). The PDG$_{max}$ tensors of *pML1::KRP1 as2-7D* outer
epidermal cells are similar to those of *pML1::KRP1*, wherein the transverse growth orientation
exhibited by *as2-7D* is largely re-oriented longitudinally (Fig. 4, M to P and fig. S5, E to L). In
addition to reduced buckling, the characteristic outgrowths of the *as2-7D* outer epidermis are
also nearly absent in *pML1::KRP1 as2-7D* (Fig. 4, E to H), suggesting that outgrowths form as a
result of buckling. Overall, consistent with our hypothesis, our findings show that misdirected
growth of the outer epidermal cells is a key driver of buckling in *as2-7D*.
236

The second condition of our 2D analytical model states that buckling can be restricted to the
outer epidermis if it is softer than the inner epidermis. To test the relative stiffness of the outer
and the inner epidermal layers, we performed osmotic treatments (*62*) and compared the
shrinkage of the WT and *as2-7D* epidermal layers. When exposed to a salt solution, cells
undergo plasmolysis while the cell wall contracts (Fig. 4, Q and R). The extent of cell wall
shrinkage gives a proxy measure for cell wall stiffness, with stiffness being inversely
proportional to the wall shrinkage (*63*). We exposed early sepals to salt solution and measured
the epidermal shrinkage as a proxy for epidermal stiffness (Fig. 4, S and T, fig. S5, M to P, and
see Materials and methods). We found that the WT epidermal layers exhibit similar shrinkage,
indicating comparable stiffness (Fig. 4, U and Y). In contrast, the outer epidermis of *as2-7D*
shrank more than the inner epidermis, indicating that *as2-7D* outer epidermis is softer, which
aligns with our model's prediction (Fig. 4, V and Y). We then wondered whether increasing
giant cell formation affects the stiffness of the outer epidermis. To test this, we performed
osmotic treatments on *ML1::KRP1* and *ML1::KRP1 as2-7D*. Indeed, we found increased
stiffness of the outer epidermis in both *ML1::KRP1* and *ML1::KRP1 as2-7D* (Fig. 4, W to Y),
suggesting that aligning growth along the longitudinal direction and increasing outer epidermal
stiffness rescues the *as2-7D* phenotype. The osmotic treatment results are consistent with our
initial hypothesis that reduced stiffness in the outer epidermis is another key factor in buckling.
Taken together, our findings indicate that unidirectional growth and comparable stiffness across
both epidermal layers are essential criteria for smooth sepal formation.
257
258   **Buckling in early *as2-7D* sepal stages is primarily due to growth defects rather than**
259   **abaxial-adaxial patterning disruption**
260
It has long been established that the juxtaposition of inner (adaxial) and outer (abaxial) identity
leads to the growth of the leaf blade (*50, 64, 65*). Ectopic boundary formation between these
identities can result in abnormal growths on the leaf blade that disrupt the smooth leaf surface
(*65-68*). An ectopic boundary can emerge when the inner identity-promoting transcription factors
are mis-expressed on the outer side, creating a juxtaposition between inner and outer identity
(*69*). Our evidence strongly supports our hypothesis that ectopic *AS2* expression disrupts growth
patterns and tissue properties, leading to buckling. However, since *AS2* is known to promote
inner epidermal identity (*44-46, 48, 70*), we tested the alternative hypothesis that *as2-7D*
phenotype is caused by ectopic juxtaposition of inner and outer identity. We therefore looked at
the expression pattern of *REVOLUTA* (*REV*) as an inner identity transcription factor (*53, 71, 72*),
and *FILAMENTOUS FLOWER* (*FIL*) as an outer identity transcription factor (*73-75*). We
observe no difference in *REV* and *FIL* expression patterns between WT and *as2-7D*, and
particularly, no ectopic juxtaposition between inner and outer gene expression was formed in the
as2-7D sepal surface (fig. S6, A to H). Since REV promotes adaxial identity in conjunction with
AS2, we tested whether REV is necessary or sufficient for the *as2-7D* phenotype. Ectopic *AS2*
expression in the *rev-6* mutant (*pML1::AS2* in *rev-6*) leads to folds and outgrowth formation

similar to *pML1::AS2* in WT (Fig. 1, M to P, fig. S1, G to N, S6, I to L and S7, A to I),
indicating REV is not necessary for the phenotype. Conversely, ectopic *REV* expression in WT
sepals is insufficient to generate buckles or outgrowths (fig. S7, J and K). Further, ectopic *REV*
expression in *as2-7D* also does not enhance the phenotype (fig. S7, L and M). Overall, our
investigations indicate no obvious inner-outer polarity disruption in *as2-7D*, which is consistent
with our hypothesis that buckling in *as2-7D* is indeed attributable to growth defects. However,
we cannot completely rule out the possibility that other polarity-associated genes are disrupted in
as2-7D.
285
286 **Buckling in *as2-7D* generates outgrowths via ectopic PIN convergence driving auxin**
287 **maxima formation**
288
289 In addition to the buckles, we observe the emergence of pointed outgrowths at the ends of the
290 buckles in the later stages in *as2-7D* (Fig. 1F). While constitutive *AS2* overexpression has
291 previously been reported to cause outgrowths (*46*), what local signal triggers the formation of
292 these outgrowths at specific sites remains elusive. In previously characterized outgrowths,
293 expression of auxin biosynthesis genes has been reported (*76*), suggesting auxin plays a role.
294
The polar localization of PIN-FORMED 1 (PIN1) protein, an auxin transporter, to the plasma
membrane on a particular side of the cell determines the directionality of auxin flow. When PIN1
polarity from multiple neighboring cells converges to the same location, auxin accumulates at
that site, creating an auxin maximum (*77*). In the shoot apical meristem (SAM), PIN1 converges
to form auxin maxima causing the outgrowth of primordia (*77*). PIN1 polarity also responds to
mechanical stress (*78*). Since buckling alters patterns of mechanical stress (*79, 80*), we
hypothesized that bucking causes PIN1 convergence which subsequently leads to auxin
accumulation and outgrowth formation in *as2-7D*. In the inflorescence and early-stage flowers,
the *PIN1* expression pattern in *as2-7D* is similar to that of WT (Fig. 5, A and B). However, in
stages where *as2-7D* exhibits both buckles and outgrowths, we observe PIN1 convergence
specifically in the sites where the outgrowths are formed (Fig. 5, C and D, fig. S8, A and B). In
contrast, no PIN convergence sites are observed in the middle of the WT sepals (Fig. 5, C and D,
fig. S8, A and B). We also observe the expression of auxin-responsive promoter DR5
specifically where outgrowths are formed in *as2-7D* (fig. S8, C and D), which is consistent with
the fact that *PIN1* convergence is essential for creating local auxin peaks (*81*).
310
We next tested whether PIN-mediated polar auxin transport is required for the formation of
outgrowths. We treated growing *as2-7D* inflorescences with polar auxin transport inhibitor N-1-
naphthylphthalamic acid (NPA) and compared the phenotype to mock-treated samples. While
as2-7D live-imaged in the presence of NPA exhibited characteristic buckles, the formation of
outgrowths was largely attenuated (Fig. 5, E and F, fig. S8, E to H, and movie S4). Buckling
continues to occur since the characteristic growth patterns and misdirected sideways PDG_{max}

tensors leading to buckling are unaltered with NPA treatment (Fig. 5, G to J). However, the
localized fast growth leading to outgrowth formation was largely reduced in the NPA-treated
samples (Fig. 5, K and L). This indicates that polar auxin transport is critical for the formation of
outgrowths, which are characterized by localized fast growth. Overall, our results are consistent
with our hypothesis that mechanical buckling causes ectopic PIN1 convergence, generating local
auxin maxima formation and fast growth, leading to outgrowth formation.

DISCUSSION
324
While buckling is critical for generating complex organ shapes required for specialized
functional roles (*1-5, 10, 11*), buckling is not always advantageous, and needs to be
circumvented to generate smooth organs like sepals (*82*). How cell growth patterns influence
tissue growth, and thereby organ shape remains one of the major open questions in biology.
Using Arabidopsis sepals as the model, our study demonstrates that unidirectional growth of both
the outer and inner epidermis along the sepal longitudinal axis, and similar tissue stiffness
between epidermal layers are key drivers of smooth sepal formation. Ectopic *AS2* expression on
the outer epidermis of *as2-7D* creates an imbalance of growth and tissue stiffness across outer
and inner epidermal layers, leading to outer epidermal buckling.
334
While our simple 2D analytical model directed us to the importance of analyzing growth and
stiffness across epidermal layers, our experiments show that buckling of growing biological
tissues is a complex three-dimensional phenomenon involving unequal growth directions and
tissue stiffness. Interestingly, we find the growth rate and stiffness of the inner epidermis in *as2-*
7D to be affected. Intuitively, one might expect that ectopic *AS2* expression on the outer
epidermis would alter cell growth properties only on the outer epidermis of *as2-7D*. However,
changes in inner epidermal growth properties in *as2-7D* could be due to the fact that the outer
and inner epidermis are connected at the sepal margins. Altering growth patterns in the outer
epidermis creates mechanical constraints on the inner epidermis, thereby affecting its growth
indirectly. Broadly, our results show how the growth of an epidermal cell layer (or speculatively
an animal epithelium (*83*)) relative to the tissue to which it is attached, drives organ shape.
346
AS2 has been studied primarily in the context of outer-inner polarity determination (*44-46, 48,*
70, 84). However, we demonstrate that the role of *AS2* extends beyond polarity specification.
Given that ectopic *AS2* expression disrupts the coordination of growth patterns across epidermal
layers, we propose that AS2 plays a central role as a growth coordinator. Growth coordination is
a phenomenon that is barely understood across plants and animals (*83, 85*). We hereby define
growth coordination as the exchange of positional, growth rate, and growth directional
information among cells across tissue layers for developing robust, functional organs. The
precise mechanical and molecular mechanisms through which AS2 relays information across cell
layers to coordinate growth across the organ, remain to be explored in future studies.

While how molecular cues influence tissue mechanics has been relatively well studied, how
tissue mechanics inform molecular signals remain elusive. Our study shows that, not only does
AS2 influence tissue mechanics generating buckles, but also that buckles feed back into
molecular signals to influence organogenesis. We show that buckling influences the polarity of
the PIN1-auxin efflux transporter. Our findings support the previous observations that
mechanical perturbations affect PIN1 polarity (*78, 86*). We further show that buckling is
sufficient to converge PIN1 polarity causing the formation of an auxin maxima, and subsequent
epidermal outgrowth. Thus, we find that buckling can initiate organogenesis. There is a long-
standing debate as to whether primordial initiation in the shoot apical meristem (SAM) is
influenced by buckling, a theory originally proposed by Paul Green (*31, 32, 87*). Contradicting
evidence shows that PIN1 polarizes before primordia bumps are visible (*88*). While our evidence
shows that buckling is sufficient to converge PIN1 and generate outgrowths, whether the same
phenomenon occurs in the shoot apical meristem to generate primordia, requires further
investigation.
370

ACKNOWLEDGEMENTS

We thank Isabella Burda, Frances Clark, Shuyao Kong, Byron Rusnak, Maura Zimmermann,
Erich Schwarz and Phoebe Roeder for their valuable comments and suggestions on the
manuscript. We thank Olivier Hamant for the *pPDF1::mCitrine-MBD* microtubule marker
plasmid. We thank Soren Strauss for his guidance with image processing. We thank Mingyuan
Zhu for his input on osmotic treatment experiments. We thank Trinh Duy Chi for his input on
imaging adult sepals. We thank Arabidopsis Biological Resource Center for providing seed
stocks for this work.
379

FUNDING
Research reported in this publication was supported by the National Institute of General Medical
Sciences of the National Institutes of Health under Award Number R01GM134037 to AHKR.
French National Research Agency (ANR, grant ANR-21-CE30-0039-01 GrowFlat, to AB).
National Natural Science Foundation of China grant no. 32270867 and Hundred-Talent Program
of Zhejiang University to LH.
Cornell China Center and Zhejiang University for a joint seed fund grant to AHKR and LH.
IMB was supported by NSF REU Award (1850796).
The authors acknowledge the use of facilities and instrumentation supported by NSF through the
Cornell University Materials Research Science and Engineering Center DMR-1719875. The
content is solely the responsibility of the authors and does not necessarily represent the official
views of the National Institutes of Health or other funders.

AUTHOR CONTRIBUTIONS

Conceptualization: AHKR, ASY, LH. Experimentation: ASY, LH, PMK, XH, MH. Imaging: ASY, LH, MH, PMK. Image processing: ASY, LH, PMK, IMB, AMG, RSS. Data analysis: ASY, LH, AK. Analytical modeling: AK, AB. Funding acquisition: AHKR, AB, LH. Project administration: AHKR, AB. Supervision: AHKR, AB, LH. Writing – original draft: ASY, LH, AHKR. Writing – review & editing: ASY, LH, PMK, AK, XH, IMB, MH, AMG, RSS, AB and AHKR.

COMPETING INTERESTS

Authors declare that they have no competing interests.

DATA AND MATERIALS AVAILABILITY

The raw datasets used for making the graphs are provided in Supplementary Data S1. Primers used for this study are available in Table S1 in Supplementary materials. The meshes and images will be uploaded in Zenodo. The supplementary movies can be downloaded from the link https://drive.google.com/drive/folders/1mI-y4dlJO6jXn8jJCxFTqsNS2jeNQtgK?usp=sharing. The R codes used for generating the plots can be accessed from GitHub: Avilash1993/R-codes-for-the-paper-Yadav-et.-al.-2023- as2-7D (github.com).

FIGURE TITLES AND LEGENDS

Fig. 1. Ectopic expression of *ASYMMETRIC LEAVES 2* (*AS2*) on the outer epidermal layer disrupts sepal smoothness in *as2-7D* mutant.
(**A** and **B**) Whole inflorescences of (**A**) WT (L*er*-0) and **B**) *as2-7D* mutant (in L*er*-0 background). Yellow arrowheads indicate outgrowths. Scalebar, 1 cm; representative images of *n* = 3 inflorescences for both genotypes.
(**C** and **D**) Mature flowers (stage 14 (*89*)) of (**C**) WT and (**D**) *as2-7D* mutant. WT sepals (green leaf-like organs) are marked with the letter "S" in white (C). Flowers are enclosed within four sepals. Note the outgrowths (yellow arrowheads) on the outer epidermis of *as2-7D* sepals (D). Scale bar, 1 cm; representative images of *n* = 3 flowers.
(**E** and **F**) Scanning electron microscopy (SEM) images of the outer epidermis of (**E**) WT and (**F**) *as2-7D* mutant sepals. Note the folds (*) and outgrowths (yellow arrowheads) on the *as2-7D* outer epidermis. Scale bar, 100 µm; representative images of *n* = 5 for WT and *n* = 10 for *as2-7D*.
(**G** and **H**) Confocal microscopy images of the inner epidermis of (**G**) WT and (**H**) *as2-7D* mutant. The images are representative of T2s harboring pLH13 (*35S::mCitrine-RCI2A*), a yellow fluorescent protein-based cell membrane marker (*90*). Scale bar, 100 µm; representative images of *n* = 3 for both genotypes.
(**I**) Schematic diagram representing the genetic mutation in *as2-7D*. Gene body of *AS2* showing the G-to-A point mutation (red arrow) at position 1484 bps upstream of ATG in *as2-7D* (Top half). The *AS2* expression (red dots) domain in the WT and *as2-7D* sepals is shown (Bottom half).
(**J**) Anatomical diagram representing (Top) an early stage 6 flower and (Bottom) an optical longitudinal section through the flower. The letter "S" in black represents sepals. The outer and inner epidermal layers of the outer sepal are marked.
(**K** and **L**) Longitudinal section of stage 6 flowers of WT. Cell walls are stained with propidium-iodide (PI, greyscale). Images show the GFP expression pattern (green) driven by (**K**) WT *AS2* promoter (*pAS2^{WT}*) and (**L**) *AS2* promoter from *as2-7D* (*pAS2^{as2-7D}*). Arrows indicate the outer and inner epidermal layers of the outer sepal. Scale bar = 20 µm; *n* = 3.
(**M**) Whole inflorescence of a representative transgenic (T1) harboring *pML1::AS2* (*AS2* expression driven by *ATML1* epidermal specific promoter) transgene in WT. Yellow arrows mark the outgrowths on the sepal epidermis. Independent T1s assessed = 28, all of which show similar levels of outgrowth formation. Representative image of *n* = 3 inflorescences.
(**N** to **P**) Confocal images of PI-stained (greyscale) inflorescences of (**N**) WT, (**O**) representative transgenic (T1) harboring *pML1::AS2* transgene in WT, and (**P**) *as2-7D* mutant. WT outer epidermis is smooth, whereas *pML1::AS2* exhibits outgrowths (yellow arrowheads) like *as2-7D*. Scale bar, 50 µm; *n* = 3 inflorescences per genotype.

Fig. 2. The *as2-7D* sepal outer epidermal morphology transitions from smooth to buckles
and subsequently outgrowths.
(**A** and **B**) Representative live-imaging series of (**A**) a WT flower and (**B**) an *as2-7D* mutant
flower starting from developmental stage 4 over a course of 96 hours, imaged at 24-hour
intervals (*n* = 3). Images capture morphological changes of the outer sepal. Both genotypes
harbor the pLH13 (*35S::mCitrine-RCI2A*) cell membrane marker (greyscale). Trichomes were
removed from the outer sepal epidermis by image processing to better visualization the sepal
morphology. Scale bar, 50 µm.
(**C**) Tilted view of *as2-7D* mutant flower at the 48-hour time point. Note the appearance of the
out-of-plane protrusion (ridge) in the middle of the sepal, circled in white. Scale bar, 50 µm.
(**D**) The *as2-7D* mutant flower at the 96-hour time point. Note the dramatic increase in the
number and extent of ridges (red circles) and invaginations (white circles), characteristic of
buckles, on the outer sepal epidermis. Yellow arrowheads indicate outgrowths. Scale bar, 50 µm.
(**E** and **F**) Cell area growth rate over a 48-hour period for the (**E**) WT and (**F**) *as2-7D* mutant
sepals. Growth rate is calculated as the ratio of cell lineage area at the later time point divided by
that of the earlier time point. The heatmap is projected onto the later time point. Red to blue
gradient shows fast to slow growth. The growth rate over 0-48 hours (left) and 48-96 hours
(right) is shown. Arrowheads point to the outgrowths. Scale bar, 50 µm.
(**G** and **H**) Cell division rate over a 48-hour period for the (**G**) WT and (**H**) *as2-7D*. Cell division
rate is calculated based on the number of daughter cells per lineage, represented as a heatmap
projected onto the later time point. Red to blue gradient shows more to less division. The cell
lineages were tracked using MorphoGraphX software (*91*). Segmented cells are outlined in
black. Representative heatmaps for one of the three replicates (*n* = 3) for WT and *as2-7D* are
shown. Scale bar, 50 µm.
(**I** to **K**) Visual representation of the analytical model (see Materials and methods) used to
determine buckling criteria. (**I**) A growing elastic system composed of three layers,
corresponding to the outer epidermis (denoted o), mesophyll (m), and inner epidermis (i) of a
sepal. Letters E, g and h represent Elastic modulus, growth rate, and thickness respectively.
Diagrams represent growth conditions causing (**J**) the tissue to curve and (**K**) buckling of the
outer epidermis.

Fig. 3. Unidirectional growth on both epidermal layers is essential for smooth sepal
development.
(**A** and **B**) Representative confocal images of the (**A**) WT and (**B**) *as2-7D* mutant live-imaged
with a high laser intensity (~16-20%) every 24 hours. Both the genotypes harbor pLH13
(*35S::mCitrine-RCI2A*) cell membrane marker (green). Scale bar, 20 µm. One of the three
representative replicates ($n = 3$) is shown.
(**C** and **D**) Segmented cells in grey superimposed onto the plasma membrane signal (also in grey)
projected on mesh surfaces generated using MorphoGraphX (*91*). Mesh surfaces for 0 and 48-
hour time points of the outer (left) and inner (right) epidermis of the (**C**) WT and (**D**) *as2-7D*
mutant are shown. Note the strikingly reduced number of cells on the inner versus outer surface
at the 0-hour time point. Scale bar, 20 µm; $n = 3$.
(**E** and **F**) Cell growth rates over a 48-hour period for the (**E**) WT and (**F**) *as2-7D* mutants are
shown. The heatmap has been projected onto the 48-hour time-point surface. In each panel, the
growth rate of cells on the outer surface (left) and inner surface (right) is shown. Red to blue
heatmap gradient shows fast to slow growth. Scale bar, 20 µm; $n = 3$.
(**G** and **H**) Cell division rates over a 48-hour period for the (**G**) WT and (**H**) *as2-7D* mutant on
the outer surface (left) and inner surface (right) are shown. Segmented cells are outlined in black.
Red to blue gradient shows more to less/no cell division respectively. Scale bar, 20 µm; $n = 3$.
(**I** and **J**) Principal directions of maximal growth (PDG$_{max}$, yellow) of the cells on the outer (left)
and inner (right) epidermis of (**K**) WT and (**L**) *as2-7D*. Segmented cells in grey are
superimposed on the plasma membrane signal (also in grey). Length of any given PDG$_{max}$ tensor
is directly proportional to the cell growth rate in that direction. A blue arrow indicates the
longitudinal sepal axis. Note the misoriented PDG$_{max}$ tensors on the *as2-7D* outer epidermis.
Scale bar, 20 µm; $n = 3$.
(**K**) A schematic diagram depicting the calculation of the angle of deviation between PDG$_{max}$
tensor and the longitudinal axis to obtain a quantitative measure of the deviation.
(**L** and **M**) Heatmaps show the angle of deviation of the PDG$_{max}$ tensor from the sepal
longitudinal axis for the outer (left) and inner (right) epidermal cells of the (**L**) WT and (**M**) *as2-*
7D mutant. Scale for the angle of deviation ranges from 90º (highly deviated, red) to 0º (no
deviation, blue). Scale bar, 20 µm; $n = 3$.
(**N** and **O**) Cortical microtubule orientation (greyscale) from comparable regions of stage 6
flower outer epidermis for the (**I**) WT and (**J**) *as2-7D* mutant is shown. These plants harbor
pPDF1::mCitrine-MBD, a YFP based microtubule marker (*92*). White arrows point towards the
longitudinal axis of the sepals. Microtubules in the WT are oriented perpendicular to the sepal
longitudinal axis, whereas, in *as2-7D*, the microtubules are misoriented. Yellow arrowheads
point towards the misoriented microtubules. Scale bar, 10 µm; $n = 3$.

Fig. 4. Aligning growth along the longitudinal direction and increasing outer epidermal stiffness is sufficient to rescue buckling in *as2-7D*.

(**A** to **D**) Representative stage 14 flowers of (**A**) WT, (**B**) *as2-7D* mutant, (**C**) *AtML1::KRP1,* and (**D**) *AtML1::KRP1 as2-7D*. The WT sepals are marked with the letter "S" in white (A). Yellow arrowheads point to outgrowths in *as2-7D* (B). Note the smooth sepal of *AtML1::KRP1 as2-7D* compared to *as2-7D*. Scale bar, 1 cm; *n = 3* flowers for all genotypes.

(**E** to **H**) Confocal images of the outer epidermis of (**E**) WT, (**F**) *as2-7D* mutant, (**G**) *AtML1::KRP1*, and (**H**) *AtML1::KRP1 as2-7D* stage 14 sepals. The sepals were stained with PI (grey, cell walls). Note the increased number of highly elongated giant cells in *AtML1::KRP1* and *AtML1::KRP1 as2-7D* sepal, and the smooth outer epidermis of *AtML1::KRP1 as2-7D* sepal compared to the highly rugose morphology of *as2-7D*. Scale bar, 200 µm; *n = 3*.

(**I** to **L**) Magnified insets of the (**I**) WT, (**J**) *as2-7D* mutant, (**K**) *AtML1::KRP1,* and (**L**) *AtML1::KRP1 as2-7D* sepal for the regions marked by white boxes in panels E-H respectively. Asterisks denote a typical giant cell. Scale bar, 200 µm; *n = 3*.

(**M** to **P**) The principal direction of growth (PDG$_{max}$) for the outer epidermal cells of the (**M**) WT, (**N**) *as2-7D,* (**O**) *AtML1::KRP1,* and (**P**) *AtML1::KRP1 as2-7D* sepals. Yellow lines mark the PDG$_{max}$ tensors calculated over a 48-hour period, projected onto the later time point. For WT and *as2-7D*, the flower images used are the same as in Figure 2. Segmented cells in grey are superimposed onto the plasma membrane signal (also in grey). Scale bar, 20 µm; *n = 3*.

(**Q** and **R**) Schematic diagrams demonstrating the concept of cell wall stiffness assessment by osmotic treatment assays (*62*). (**Q**) A typical plant epidermal cell before osmotic treatment. The cell wall and cell membrane are shown in black and green respectively. (**R**) A cell that has undergone plasmolysis in a hyperosmotic solution (0.35 M NaCl).

(**S** and **T**) Anatomical diagrams demonstrate using osmotic treatments to estimate epidermal stiffness. Longitudinal sections of an early stage 6 flower (**S**) before and (**T**) after osmotic treatment are shown. The outer (green) and inner (blue) sepal epidermis are marked. Arrowheads (red) demonstrate aerial shrinkage.

(**U** to **X**) Heatmaps show the aerial shrinkage (in percentage) of a longitudinal sepal cross-section for the inner and outer epidermis of (**U**) WT, (**V**) *as2-7D* mutant, (**W**) *AtML1::KRP1* and (**X**) *AtML1::KRP1 as2-7D*, post osmotic treatment in 0.35M NaCl solution. Shrinkage was computed as the decrease in area of a comparable cross-sectional region from an early stage 6 flower before and after osmotic treatment. Heatmaps are projected on the post-treatment surface. More shrinkage is represented in red and less in blue. Scale bar, 20 µm. Heatmaps for each genotype are representative of 5 independent replicates (*n = 5*).

(**Y**) Graph showing the percentage aerial shrinkage post osmotic treatment. Data across 5 independent replicates each for the WT (black dots), *as2-7D* (red dots), *AtML1::KRP1* (grey dots), and *AtML1::KRP1 as2-7D* (pink dots) for both the outer and inner epidermis are shown. The letters above the plots represent significant differences based on one-way analysis of variance (ANOVA) with Tukey's honest significant difference (HSD) post hoc correction (*p* < 0.05). Similar letters indicate no significant difference.

Fig. 5. Mechanical buckling drives ectopic PIN convergence creating auxin maxima, causing fast local growth and subsequent outgrowth formation.

(**A** to **D**) Confocal stack images showing the expression pattern of auxin efflux carrier protein PIN-FORMED 1 (PIN1) in the WT and *as2-7D*. (**A** and **B**) Expression pattern of PIN1 protein reporter (*pPIN1::PIN1-GFP* (*93*)) in the inflorescence meristem of (**A**) the WT, (**B**) *as2-7D* mutant. The *PIN1* expression pattern in *as2-7D* is identical to WT in the young floral development stages, wherein PIN1 polarity is directed towards the insipient primordia and growing sepal edges, as described previously (*93*). Scalebar, 50 µm. (**C** and **D**) Expression pattern of *PIN1* in a representative middle stage (stage 8) flower of the (**C**) WT and (**D**) *as2-7D* mutant. Note the expression of *PIN1* at the growing tip of the outgrowths (yellow arrows) in *as2-7D*, whereas no *PIN1* expression is detected in the blade region where the epidermis buckles. In panels A to D, the images are representative of three *pPIN1::PIN1-GFP* and *pPIN1::PIN1-GFP as2-7D* plants (*n* = 3). PI is shown in greyscale and GFP in green. Trichomes were removed from the outer sepal epidermis by image processing. Scale bar, 50 µm.

(**E** to **F**) Representative live-imaging series of (**E**) Mock treated *as2-7D* and (**F**) 10µM NPA treated *as2-7D* developing flowers over 96 hours, imaged at every 24-hour interval (representative of *n* = 3 flowers). Note the development of outgrowths from 72-hour time-point onwards (yellow arrowheads) in the mock-treated *as2-7D*, which are absent in NPA treated *as2-7D* flowers. Plants harbor pLH13 (*35S::mCitrine-RCI2A*) cell membrane marker (greyscale). Scale bar, 50 µm.

(**G** to **H**) The PDG_{max} tensors (yellow) of the cells on the outer epidermis of (**G**) mock-treated and (**H**) 10µM NPA treated *as2-7D* over the first 48-hour growth period for the representative samples shown in panels E to F. Characteristic misoriented PDG_{max} tensors (relative to the longitudinal sepal axis) on the outer epidermis of *as2-7D* remain unaffected in the NPA-treated samples, thus driving buckling in later stages. Scale bar, 50 µm.

(**I** to **L**) Outer epidermal cell growth rates over (**I** and **J**) 0–48-hour growth period and (**K** and **L**) 48–96-hour growth period of the mock and 10µM NPA treated *as2-7D* flowers respectively. Heatmaps are projected onto the later time point. Fast (red) to slow (blue) growth rates are shown. In panel K, note the fast cell growth driving outgrowth formation (yellow arrowheads) which is absent in NPA treated sample (panel L). Scale bar, 50 µm.

MATERIALS AND METHODS

Plant Materials

Arabidopsis thaliana genotypes used in this study are in Landsberg *erecta* (L*er*) background, which is referred to as the wild type (WT), unless otherwise mentioned. The mutants *rev-6* (*71*) and *as2-5D* mutant (in Col-0 background) (*48*) have been previously described, and were obtained from Arabidopsis Biological Resource Center (ABRC). The *as2-7D* mutant in the L*er*-0 background was identified from ethyl methanesulfonate (EMS) based forward genetic screen. Briefly, M2 seeds (second generation) obtained from self-fertilized EMS mutagenized L*er* plants were purchased from Lehle Seeds (*94*). M2 plants were screened under a stereo microscope for sepal defects (*60, 95*). The *as2-7D* mutant was isolated based on the lumpy sepal phenotype. A mapping population was generated by crossing the *as2-7D* mutant with a Columbia (Col-0) accession plant. Standard map-based cloning techniques were used to clone the underlying mutation (*96*). The *as2-7D* mutation is a G to A change at base 1,484 upstream of the ATG translational start site of AT1G65620/*AS2*. To reduce background mutations, the *as2-7D* mutant was backcrossed twice to wild-type L*er* plants twice before further analysis. In the backcrosses, we found that *as2-7D* mutant behaves as a dominant allele as the F1s exhibit the *as2-7D* phenotype, although it is not as strong. The *as2-7D* mutation can be PCR-genotyped by amplifying with primer pairs oLH108 and oLH324 at 55°C annealing temperature, followed by digesting the product with EcoRI to produce an 82-bp WT product or a 111-bp mutant product. Primers are listed in Supplementary Table S1.

Plant Growth

Plants were grown in Percival (Percival-Scientific) growth chambers programmed for long day conditions (16-hour light/8-hour dark cycles). Growth temperature (22ºC), relative humidity (65%) and fluorescence intensity (~100 µmol m^{-1}s^{-1}) were maintained.

For soil-based experiments, plants were grown on soil premixed with Vermaculite (LM111). For stratification, the trays were placed in a 4 ºC cold room for three days prior to moving them to the growth chambers. For BASTA (glufosinate ammonium) selection, the soil was autoclaved twice (Program 3- 15 min sterilization, 30 min dry, Primus ST-1 Autoclave (Model PSS5-B-MSSD)), and sterilized seeds were planted at a density of 50-100 seeds per pot. For live-imaging and adult-stage phenotyping experiments, one plant was grown per pot (6.5×8.5 inches) to facilitate healthy vegetative growth.

For plate-based experiments, seeds were washed twice with a sterilization solution containing 70% ethanol + 0.1% Triton X-100, replacing the solution after each wash. Two final rounds of washes were provided with 100% ethanol. Seeds were spread on filter paper (Whatman 1,

626   diameter 90 mm) and allowed to dry inside a sterile safety cabinet (Class II, Type A2, Labgard).
627   Dried seeds were evenly distributed on freshly prepared MS plates (1/2 MS basal salt mixture,
628   0.05% MES w/v, 1% sucrose w/v, pH ~ 5.8 (adjusted using KOH) containing 0.8% phytoagar.
629   The plates were sealed with 3M micropore tape, wrapped in aluminum foil, and stratified for
630   three days at 4 ºC before being moved to the growth chamber.
631

Phenotyping whole inflorescences, flowers, and plants

Whole inflorescence and stage 14 flower images were taken either using Canon PowerShot A640
10MP Digital camera attached to Zeiss Stemi 2000-C stereo microscope or Accu-Scope Excelis
4K UHD camera attached to Zeiss Stemi 508 stereo microscope. For inflorescence images, and
Stage 14 flowers an optical zoom of 1.0 and 1.6 were used respectively. Whole plant images
were captured using iPhone XR with a 26 mm focal length and aperture of f/1.8.

Schematic diagrams

The schematic diagrams shown in Fig. 1I and Fig. 4 Q to T were hand-drawn using the
Notability application in Apple iPad (model A1701).

Genetic crosses

Crosses were performed as described previously (*97*). The *pATML1::KRP1* in L*er* background
(*60, 61*) were crossed into *as2-7D* mutants. Segregation analysis and confocal microscopy was
used to find *pATML1::KRP1 as2-7D* plants. The *pPIN1::PIN1-GFP DR5::VENUS* (*93*) plants
and *pPIN1::PIN1-GFP pin1-4* (*93*) plants were crossed into *as2-7D* mutants. The *pPIN1::PIN1-*
GFP pin1-4 plants (control) exhibit a mild phyllotaxy defect. Genetic segregation analysis and
confocal microscopy was used to find *DR5::VENUS as2-7D* and *pPIN1::PIN1-GFP as2-7D*
plants.

Cloning overexpression constructs and reporters
The pLH13 (*p35S::mCitrine-RCI2A,* hygromycin resistance) and pAR169 (*pATML1::RCI2A-*
mCtrine, basta resistance) plasma membrane markers have been described previously (*60, 98*).
The *pPDF1::mCitrine-MBD* microtubule marker (*92*) was a kind gift from Olivier Hamant (RDP
institute, ENS de Lyon, France).

For cloning *pML1::AS2*, the AT1G65620/*AS2* coding sequence (600 bps) was amplified from
Col-0 cDNA using primers oLH111 and oLH112 by Phusion PCR (NEB), and cloned into
pENTR™/D-TOPO™ (Invitrogen™). The *AS2* coding sequence was cloned into the *pATML1*
containing gateway vector pAR176 (Basta resistance (*60*)) by LR reaction.

For *pML1::REV*, the AT5G60690/*REV* full length coding sequence spanning 2,529 bp was amplified from Col-0 inflorescence cDNA using primer pairs oAS_005 and oAS_006. The resulting PCR product was subcloned into pCR™8TOPO™ vector (Invitrogen™) by TA cloning (*99*). LR reaction was used to clone the *REV* coding sequence into pAR176.

To clone *pAS2 ^{WT}::GFP-GUS*, the 4.1 kb upstream promoter region and part of the first exon of *AS2* was PCR amplified from L*er*-0 genomic DNA using primer pairs oLH347 and oLH370, and cloned into pENTR™/D-TOPO™ vector (Invitrogen™). The resultant vector was LR recombined into the gateway vector pBGWFS7 (*100*). For cloning *pAS2 $^{as2-7D}$::GFP-GUS,* the 4.1 kb upstream promoter region and part of the first exon of *AS2* was PCR amplified from *as2-7D* genomic DNA, and the subsequent cloning steps were similar to cloning *pAS2::GFP-GUS*.

 For *proREV::REV-GFP*, the 7.75 kb *proREV::REV* fragment (amplified from L*er*-0) and 1.08 kb mGFP-T35S fragment amplified from *pML1::H2B-mGFP* (pAR180 (*60*)) were assembled into the PmeI site of pMOA33 (Kanamycin resistance).

For cloning *proFIL::mCherry-RCI2A*, a 3.92 kb *FIL* promoter fragment was amplified from L*er*-0. This fragment was assembled with a 2.19 kb *omega-mCherry-RCI2A-T35S* fragment amplified from *p35S::mCherry-RCI2A* (pMZ11 (*101*)) into the SacI and SpeI site of pMOA34 destination vector (hygromycin plant resistance, (*102*)). All the intermediate and final constructs were verified for sequence and directionality by sequencing. Primers are listed in Supplementary Table S1.

Plant transformation and selection

The binary vector constructs were transformed into *Agrobacterium tumefaciens* GV3101 strain (Rifampicin and Gentamycin resistant) by triparental mating method (*103*). Briefly, two *E.coli* strains, one containing the binary vector and the other containing the helper (conjugative) plasmid, and *Agrobacterium* strain GV3101 were mated together. The GV3101 clones were streaked onto Rifampicin + Gentamycin + binary vector specific antibiotic plates for selection, and positive clones were isolated by colony PCR. The T-DNAs were transformed into plants by Agrobacterium-mediated floral dipping (*104*). Transformants were selected either on ½ MS media plates (for Hygromycin and Kanamycin selection) or on soil (for Basta selection). For selecting pLH13 in WT and *as2-7D*, *proFIL::mCherry-RCI2A* in WT and *as2-7D* T1s, a final concentration of 25 -30 µg/mL Hygromycin was used. For selecting *proREV::REV-mGFP* in WT and *as2-7D* T1s, a final concentration of 50 µg/mL Kanamycin was used. Healthy, non-chlorotic plants were transplanted to the soil post two weeks of growth. For selecting *pML1::AS2* in WT and in *rev-6*, *pML1::REV* in WT and in *as2-7D*, *pAS2 ^{WT}::GFP-GUS* and *pAS2 $^{as2-7D}$::GFP-GUS* in WT, *pPDF1::mCitrine-MBD* in WT and in *as2-7D* T1s, BASTA selection was performed on soil grown plants. Seedlings aged 1 to 2 weeks were selected with 100 µg/mL

706   BASTA solution by spraying, and non-chlorotic plants were transplanted to fresh pots after a
707   week. At least twenty independent T1 plants were characterized for each construct. For
708   fluorescent reporter lines, T1 plant was chosen for each line based on fluorescence. Confocal
709   microscopy was used to assess the stability of fluorescence in every generation.
710
711   **Flower staging**
712
713   Floral staging was performed based on Smyth et al (*89*). Briefly, sepal primordia barely arise in
714   Stage 3, and overlie the floral meristem in Stage 4. Sepals enclose the floral bud completely by
715   late stage 6. Flowers beyond stage 6 were staged based on size and morphology (*42*).
716
717   **Sepal selected for microscopy**
718
For flower imaging across experiments (live imaging and single time point reporter imaging), we
focused primarily on the outer sepal of the flower, unless specified. Briefly, flowers are enclosed
within four sepals, wherein the foremost sepal, which faces away from the meristem is denoted
as the outer sepal, whereas the sepal that is diametrically opposite to the outer sepal and facing
towards the meristem is denoted as the inner sepal. The remaining two sepals positioned on
either side are the lateral sepals. For a detailed description, please refer to Roeder et. al (*42*).
725
726   **Scanning electron microscopy (SEM)**
727
728   Scanning electron microscopy was performed as described previously (*60*). Briefly, Stage 14
729   flowers were fixed in an FAA solution (50% ethanol, 5% acetic acid, and 3.7% formaldehyde)
730   for 4 hours, dehydrated using an ethanol series, and critical point dried. Sepals were dissected,
731   sputter-coated with platinum palladium and then imaged using a LEICA 440 scanning electron
732   microscope.
733
734   **Confocal Microscopy**
735
Live imaging
For live imaging sepal development, primary inflorescences from six- to eight-week-old plants
containing the *p35S::mCitrine-RCI2A* membrane marker (pLH13 in WT, *as2-7D*, *pML1::KRP1,*
and *pML1::KRP1as2-7D* T2s) were excised. Dissection was performed as described previously
(*101, 105*), with modifications. Briefly, siliques were removed using surgical scissors (Artman
Instruments). Flowers older than stage 7 were dissected using Dumont tweezers (Electron
Microscopy Sciences, Style 5) under a stereo microscope (ZEISS Stemi 2000). Dissected
inflorescences were inserted upright into small petri-plates containing live imaging media (1/2
MS media supplemented with 1X Gamborg Vitamin mix and 1000-fold diluted plant
preservative mixture) with 1.2% w/v phytoagar. The plates were filled with autoclaved water and

allowed to stand for two-three hours for the dissected inflorescence to rehydrate. Residual buds
were dissected down to Stage 3-4 in water using a needle (BD Micro-Fine^{TM} IV Insulin
Syringes). Inflorescences were moved to fresh media plates and transferred to the growth
chamber for at least 24 hours for recovery from dissection, prior to imaging. Stage 4 flowers
were selected as the 0- hour time point across all live imaging series.
751
Post recovery, dissected tissues for live imaging were positioned at an angle to ensure the outer
sepal of the chosen stage 4 flower directly faces the objective of the confocal microscope. The
flowers were submerged in MillQ water containing 0.01% v/v SILWET L-77 surfactant and
imaged using Zeiss 710 upright confocal microscope using 20X water dipping objective (Plan-
Apochromat 20x/1.0 DIC D=0.17 M27 75mm). The following settings were used: mCitrine
(yellow fluorescent protein), excitation laser 514 nm (intensity- 2.4-6.0%), emission collection
range 519-596 nm and a z-step of 0.415-0.420 µm. For high laser intensity based live imaging of
pLH13 in WT and in *as2-7D* to visualize the adaxial sepal epidermis, imaging settings for
mCitrine used: excitation laser 514 nm (intensity- 16-18%), emission collection range 519-596
nm and a z-step of 0.244 µm. Post imaging, the plates were returned to the growth chamber and
the process was repeated for each time-point. Inflorescence tissues were transferred to fresh
media every 48 hours to prevent fungal contamination.
764
765 **NPA treatment assay**
766
Primary inflorescences of six- to eight-week-old pLH13 in *as2-7D* T2s were dissected down to
stage 3 and were divided into two groups of 5-6 samples. One group of samples were placed in ½
MS live imaging media plates containing 10 µM NPA. The other group, serving as the control,
was placed in media plates containing an equivalent amount of DMSO (20 µL DMSO per 100
ml media). The tissues were dissected 48 hours prior to imaging. Post dissection, the tissues were
treated every 12 hours with a 100 µL liquid solution, applied directly above the inflorescence
tissue by pipetting. For the 10 µM NPA treatment samples, the liquid solution comprised of 1 µL
of 1M NPA+19 µL DMSO+10 µL Silwet in 100 ml water, whereas the control (mock) samples
were treated with 20 µL DMSO+10 µL Silwet in 100 ml water. The aim of administering liquid
solution was to enhance the effectiveness of the treatment, which was performed every 12 hours
throughout the experimental duration. Stage 4 flowers, 48 hours post dissection were selected as
the 0- hour time point for imaging mock and 10 µM NPA treated samples. Live imaging was
performed as described above, with the samples being moved to fresh mock or 10 µM NPA
plates every 48 hours. Imaging settings used: mCitrine (yellow fluorescent protein), excitation
laser 514 nm (intensity -2.6-2.8%), emission collection range 519-596 nm, and a z-step of 0.415
µm.
783
784 **Propidium Iodide (PI) cell wall staining and imaging**
785

A 5 mg/ml stock solution was prepared for PI staining, with 5 mgs of PI (Sigma) solubilized in 1

mL of MilliQ water. The stock solution was stored in a foil-covered 1.7 ml microcentrifuge tube

and refrigerated at 4ºC. For staining, stock solution was five-fold diluted with water containing

0.01% v/v Silwet surfactant (PI working solution). Whole inflorescences and flowers below

Stage 10 were stained for 6-8 minutes, and stage 14 sepals were stained for 12 minutes. Care was

taken to ensure that the tissue was fully submerged in the staining solution, and no air bubbles

surrounded the tissue. Following staining, the tissue was washed in a fresh solution containing

just 0.01% v/v Silwet, prior to imaging. Fresh PI working solution was prepared before every

experiment. For imaging PI-stained inflorescences of WT, *as2-7D* and the transgenics (T1s) of

pML1::AS2 in WT, *pML1::REV* in WT and *pML1::REV* in *as2-7D*, the following settings were

used: excitation 514 nm (laser intensity -18-22%), emission 566-650 nm.

Imaging reporters

For single time point imaging of the reporters, inflorescences were dissected down to stage 3-4.

The recovery period was extended to 48 hours to allow ample time for stabilization of the

reporter signal. The inflorescences were placed upright in the live imaging media plates with the

meristem facing the objective. Prior to taking images of late-stage flowers, the inflorescences

were allowed to grow for a few days. For the reporters imaged together with PI, fresh

inflorescences were dissected for imaging late-stage flowers. Images were taken with Zeiss 710

upright confocal microscope using 20X water dipping objective.

The following settings specific for each of the reporters were used. For *pAS2 ^{WT}::GFP-GUS* and

pAS2 $^{as2-7D}$::GFP-GUS images, PI - excitation laser 594 nm (intensity- 2%), emission range 599-

650 nm and GFP - excitation laser 488 nm (intensity-20%), emission range 493-550 nm, with z-

step of 0.240 µm. For *proREV::REV-GFP*, Chlorophyll and GFP were excited with 488 nm laser

(intensity- 40%), emission range was collected between 647-722 nm for Cholorophyll and 493-

550 nm for GFP. Z-step, 0.240 µm (inflorescences), 0.4 µm (flowers). For *proFIL::mCherry-*

RCI2A, Chlorophyll and mCherry were excited with 561 nm laser (intensity- 12-15%), emission

range was collected between 655-722 nm for Chlorophyll and 578-650 nm for mCherry. Z-step,

0.240 µm. For *pPIN1::PIN1-GFP*, PI - excitation laser 594 nm (intensity- 2%), emission

collection range 599-650 nm and GFP - excitation laser 488 nm (intensity- 25-30%), emission

collection range 493-550 nm, with z-steps of 0.240-0.5 µm. For *pDR5::VENUS*, samples were

excited with an argon laser (514 nm), and data were collected in the YFP (520–565 nm)

channelFor *pPDF1::mCitrine-MBD,* PI and mCitrine were excited with 514 nm (intensity- 30-

32%), and emission range was collected between 603-643 nm for PI and 519-567 nm for

mCitrine. Z-step, 0.2 µm.

Imaging adult sepals

For imaging fully mature sepals, all flowers were dissected off the inflorescence except for a
single stage 14 flowers. All the internal floral organs from the stage 14 flower were carefully
dissected off, leaving just the outer sepal (*42*) attached to the pedicel. The sepal was wetted in
0.01% v/v Silwet water and mounted between two coverslips filled with 0.01% v/v Silwet water.
To prevent the sepal from drying out during imaging, the coverslips were sealed with vacuum
grease. Sepals were imaged using 10X objective (C-Apochromat 10X/0.45 W M27). Three-
horizontally tiled images were taken with an overlap of 5% and were auto-stitched with a
threshold of 0.70 in the ZEISS ZEN software. For imaging fully mature sepals of pLH13 in WT
and in *as2-7D*, mCitrine was excited with 514 nm laser (intensity- 20-22%) and emission was
collected at 519-621 nm. For imaging PI stained sepals of WT, *as2-7D*, *pML1::KRP1* and
pML1::KRP1 as2-7D, PI was excited at 514 nm (intensity- 27-30%) and emission was collected
at 566-648 nm. For epidermal structure observation, mature flowers harboring pAR169
(*pATML1::RCI2A-mCtrine*) were transverse sectioned using a razor blade. Both mCitrine and
chlorophyll were excited using 514 nm laser, and emission was collected at 520-565 nm for
mCitrine and 651-700 nm for chlorophyll.
841
842   **Osmotic treatments for measuring epidermal stiffness**
843
Osmotic treatments were performed as described previously (*43, 62*), with modifications. In
brief, primarily inflorescences from pLH13 in WT, pLH13 in *as2-7D*, pLH13 in *pML1::KRP1*
and pLH13 in *pML1::KRP1 as2-7D* six to eight week old plants were dissected down to stage 3
to 4, and placed in ½ MS live imaging media plate. Dissected tissues were allowed to recover for
24 hours in the growth chamber. Early stage 6 buds were precisely chosen for osmotic treatment,
since this is the stage right before *as2-7D* outer epidermis starts buckling.
850
Each bud (corresponding to one replicate) was subjected to osmotic treatment individually and
was imaged pre- and post-treatment. Pre-treatment imaging involved staining the bud with a PI
working solution for precisely 6 minutes. After washing off the PI stain, the tissue was
transferred to a fresh media plate and was oriented to focus on the outer sepal. The tissue was
submerged in 0.01% v/v Silwet water, and the pre-treatment image was captured. Imaging
settings used for mCitrine – excitation 514 nm (intensity- 20-25%), emission 519-567 nm and PI
- excitation 514 nm (intensity- 20-25%), emission 603-643 nm.
858
For osmotic treatment, the bud was immersed in 0.35M NaCl solution containing 0.01% v/v
Silwet for exactly 20 minutes. Post treatment, the bud was re-stained in a PI working solution
prepared in 0.35M NaCl (1 part 5 mg/mL PI (solubilized in 0.35M NaCl) and 4 parts 0.35M
NaCl solution containing 0.01% v/v Silwet)) for 10 minutes, leading to a cumulative osmotic
treatment duration of 30 minutes. This timeframe was found to be optimum for inducing
plasmolysis in epidermal cells of early stage 6 flowers, wherein the plasma membrane gets
visibly detached from the cell wall at the corners. Re-staining the buds with PI is necessary since

NaCl solution erodes off the PI stain. The bud was finally re-positioned in a fresh media plate
and submerged in 0.35M NaCl solution. Before image acquisition, the tissue was allowed to
stabilize for a few minutes due to its increased fragility post osmotic treatment. The post-
treatment image acquisition setting was as follows: mCitrine – excitation 514 nm (intensity- 25-
35%), emission 519-567 nm and PI - excitation 514 nm (intensity- 25-35%), emission 603-643
nm. This procedure was iterated for all samples individually.
872
873   **Image processing**
874
Converting raw images to tiffs, taking screenshots and videos
The raw images, generated either in .lsm or .czi formats from ZEN imaging software, were
converted to tiff files in ImageJ distribution Fiji (*106*). For multichannel images, images were
split based on channels, which were saved individually. The MorphoGraphX (MGX) (*91*)
software was used for image processing and taking the screenshots using the built-in screenshot
option in MGX. For taking screenshots of the reporter lines, the reporter image was loaded onto
Stack 1, and either chlorophyll or PI was loaded onto Stack 2. Colormaps for Stack 1 and Stack 2
were set to green and gray respectively. Flower images and meristems were rotated in the desired
orientation. The voxels not associated with the flower or meristem of interest, were removed
using the Voxel Edit tool in MGX. In addition, trichomes from the outer sepal epidermis were
removed in a similar manner, for better visualization of the sepal surface. Following this,
screenshots were captured. From flower images, trichomes on the outer sepal were removed by
image processing for better visualization of the sepal epidermis. The brightness and contrast
were uniformly adjusted across samples for any given experiment. Screenshots of MGX
generated heatmaps were taken using the default brightness and contrast settings, without any
adjustments.
891
Videos were captured as a series of two-dimensional tiffs files using the built-in "Record movie"
option in MGX. The 3D stack images were rotated in the MGX window for the desired duration
and time points. For capturing videos involving two genotypes/treatments, the images were
loaded onto Stack 1 and Stack 2 and rotated independently. The stack of tiffs generated was
loaded onto Fiji and saved as a .avi file without compression. The .avi files were assembled and
labeled in Adobe Premiere Pro 2023.
898
899   **Generating surfaces for analyzing outer epidermal growth**
900
Making the surfaces and analysis of growth data was performed in MGX as previously described
in Zhu et. al. and Hong et. al. (*43, 107*), with modifications. The following process was used to
generate the surfaces for analyzing cell growth parameters on the outer epidermis of WT, *as2-*
7D, *pML1::KRP1*, *pML1::KRP1 as2-7D*, *as2-7D* mock and 10 µM NPA treatment samples.
Briefly, the tiffs were imported onto Stack 1 Main Store in the software. Trichomes on the outer

epidermis were removed using the voxel edit tool to get a better surface. Since the outer
epidermal morphology of *as2-7D* becomes increasingly irregular in later stages, Level Sets (*108*)
was used to extract the structure representing the general flower shape. For this, Level Sets was
initialized with an up/down threshold value of 3 to 4. The structure was allowed to evolve (Level
Set Evolve) with default parameters, except for "view steps", which was set to 1 for visualization
at more frequent intervals. For performing Edge Detect Angle, the mask was oriented such that
the outer sepal directly faces the viewer. The Edge Detect Angle process was then run with
default parameters to get a finer structure that closely resembles the image. The holes generated
in the structure (which corresponds to places in the image with low membrane marker signal)
were closed using Stack/Morphology/Closing - X/Y/Z radius of 3 to 8. A 2.5D surface fitted
with a triangular mesh was created from the structure using Marching Cube Surface with a cube
size of 3 to 5 µm, followed by smoothing once (Smooth Mesh, passes = 10) and subdividing
twice. The membrane signal was projected onto the 2.5D surface with a min. and max. distance
range of 1 to 8 µm from the image.
920
921 **Generating surfaces for analyzing outer vs inner epidermal growth in WT and *as2-7D***
922
To obtain the inner epidermal surface, clipping plane 1 (View Tab/Clip 1), set to a width of 10 to
15 µm, was used to clip the flower image along the longitudinal direction. The voxel editing tool
was used to carefully remove the voxels below the inner epidermis of the sepal within the
clipping plane region. The clipping plane was then moved to the adjacent region and the same
process was repeated, while tilting the plane to ensure voxels from the inner epidermis itself are
not removed. Once the voxels below the inner epidermis were removed throughout, the edited
tiff was copied back to the main stack and saved. To generate the surface for the inner epidermis,
the Z axis of the stack was reversed (Stack/Canvas/Reverse Axes). The stack was blurred using
Gaussian blur (X/Y/Z sigma (µm) = 2 to 2.5), and structure was extracted using Edge Detect
with a threshold of 2000 to 3000. A triangular mesh surface was created using a cube size of 3
µm, followed by smoothing once (passes = 10) and subdividing twice. Plasma membrane signal
from the inner surface was projected onto the mesh surface with a min. and max. distance range
of 3 to 8 µm.
936
937 To obtain the outer epidermal surface, the processed tiff, prior to reversing the axes, was used.
938 To process the stack, Gaussian blur (X/Y/Z sigma (µm) = 1 to 1.5) followed by Edge detect
939 (threshold of 8000 to 10,000) were used sequentially. Mesh surface was created as described
940 above, and the membrane signal was projected within a 2 to 4 µm distance range from the image.
941
942 **Lineage tracking**
943
For meshes across all time points, individual cells were manually seeded. Cells were segmented
using the watershed algorithm, and unlabeled vertices were fixed by running Cell Mesh/Fix

Corner Triangles. Cell lineages were parent tracked to the earlier time point to analyze cell
growth. Parent (lineage) tracking involves transferring the parent labels from cells in the earlier
time point onto the daughter cells at the later time point. For this, the meshes corresponding to
the earlier (Mesh 1) and later (Mesh 2) time points were opened side by side on the same MGX
window. A few daughter cells in Mesh 2 were manually parent labeled, and using these as
landmarks, the remaining daughter cells were auto parent labeled (by running
Deformation/Morphing/Set Correct Parents followed by Deformation/Mesh 2D/Auto Parent
Labelling 2D). Errors in segmentation or parent tracking were manually corrected. Parent
tracking was checked for accuracy using Cell Axis/PDG/Check Correspondence, which checks
whether the cell junctions are identical across time points. The final parent labels were saved as
.csv files. A custom R script (combinelineages.r) was run to automatically correlate the lineage
information through multiple time points, followed by manual correction.
958
959   **Analyzing cell growth parameters**
960
Meshes corresponding to the earlier (Mesh 1) and later time points (Mesh 2) were loaded side by
side in the same MGX window, with parent labels loaded onto Mesh 2. Cell growth rate was
calculated based on the total area of the daughter cells in the second time point divided by the
area of the parent cell in the earlier time point. Growth rate heatmaps were generated using Heat
Map/Heat Map Classic/Heat Map with the Type set to Area. The cell division rate was calculated
based on the total number of daughters formed in the second time point for each cell in the
earlier time point. Cell division heatmaps were generated using Lineage Tracking/Heat Map
Proliferation. The values for growth rate and cell division rate were exported to .csv files.
969
The Principal Direction of Growth (PDG) uses information of corresponding junctions between
the earlier and later time points, to compute the direction of maximum and minimum growth of
each individual cell in the earlier time point (*109*). In the figures, the PDGs represent the
maximum growth direction (PDG$_{max}$) tensor (yellow lines), the length of which is directly
proportional to the magnitude along the direction of maximum growth. The PDGs were
calculated for the cells in the earlier time point (using the information of parent labels from the
later time point), the PDG$_{max}$ tensors were projected onto the later time point for visualization.
For displaying PDG$_{max}$, the values for line width and line scale were set to 5 and 3 respectively.
978
The Angle of Deviation of the PDG$_{max}$ tensor was calculated relative to the longitudinal axis of
the sepal. The sepal longitudinal axis was set using a Bezier line on the earlier time point, since
PDGs and deviation angles are calculated on the earlier time point. Briefly, a new Bezier grid
(Misc/Measure/New Bezier) was created, which was then collapsed into a Bezier line
(Misc/Bezier/Collapse Bezier Points). The supporting points on the Bezier line were placed on
the mesh in a manner such that the Bezier line is parallel to the sepal longitudinal direction, and
the line shape fits the curvature of the sepal. The custom axis for each cell was set using this

Bezier line (Cell Axis/Custom/Create Bezier Line Directions) followed by smoothing (Smooth
Custom Directions). The PDGs were loaded onto the mesh, and the angle of deviation of the
PDG_{max} from custom X direction (which represents the longitudinal axis) was calculated using
Cell Axis/Custom/Custom Direction Angle. The values of the angle of deviation heatmap were
exported as .csv files.
991
992 **Analyzing osmotic treatment induced epidermal shrinkage**
993
Our osmotic treatment conditions were optimized such that the cell walls shrank, but the cell
membrane barely plasmolyzed. Post osmotic treatment, the membrane just visibly pulled away
from the cell wall corners, thereby rendering the wall and membrane area almost comparable.
Since the PI signal (corresponding to the cell wall) is very weak post-treatment, especially for the
inner layers, the PI and mCitrine (membrane marker) images were merged as a single image
MGX (Stack/Multistack/Merge Stacks).
1000
For analyzing epidermal stiffness, we assumed that the surface tension across genotypes is
comparable. The shrinkage in area of the epidermal layers along a comparable longitudinal
section was analyzed to infer tissue stiffness. For this, the pre- and post-treatment merged stacks
for any given sample were opened side by side in the same MGX window. Using clipping plane
1 (Clip 1), longitudinal cross-section corresponding to the same regions were extracted from the
flower images. The tiffs were processed using Levels Sets to extract the global shape. Mesh
surfaces were made with a cube size of 3 µm, which was smoothed once (10 passes) and
subdivided twice. Signal was projected with a 1.5 to 3 µm distance range from the image. From
the cross-section surface, the outer epidermis and inner epidermis were seeded and segmented as
single "cells". Care was taken to ensure that the epidermal region being seeded was comparable
between the pre- and post-treatment surfaces. For measuring shrinkage, parent labels from the
pre-treatment mesh were transferred onto the post-treatment mesh, following which the
shrinkage ratios were quantified (Mesh/Heatmap/Heatmap Classic). The areal shrinkage values
were stored as a .csv file, which was converted to percentages for generating heatmaps shown in
the figures.
1016
1017 **Analytical model for buckling**
1018
System and assumptions. We consider a growing elastic system made of three layers,
corresponding to the inner epidermis (denoted i), mesophyll (m), and outer epidermis (o) of a
lateral organ. We consider that the three layers may grow at different (specified) rates and have
different elastic moduli. To simplify the formulae obtained here, we assume that the epidermal
layers have similar thicknesses. We assume that the Poisson ratios of all layers are equal (value
v). Finally, we neglect the bending modulus of each layer in view of their small thickness. These
three assumptions do not affect the qualitative behavior of the system.

1026

1027 *Formulation.* Let E_i, E_m, and E_o be the values of elastic modulus, h,h_m, and h the values of
1028 thickness. g_i,g_m, and g_o are the values of strain associated with target growth rates, for the inner
1029 epidermis, the mesophyll, and the outer epidermis, respectively.

1030

The differences in growth rates between the layers may lead to changes in system geometry. As
the three layers are bound to each other, they have approximately the same curvature c
(corresponding to the radius of curvature measured at the middle of the mesophyll), while the
strain in the inner and outer layers can be deduced from the strain in the middle layer ε using
curvatures and assuming that the reference state is flat. From kinematics of the system, the
elastic (residual) strains in the three layers are given by

1037
$$\varepsilon_i = \varepsilon - \frac{1}{2}(h_m + h)c - g_i$$

1038
$$\varepsilon_m = \varepsilon - g_m$$

1039
$$\varepsilon_o = \varepsilon + \frac{1}{2}(h_m + h)c - g_o$$

1040 The elastic energy of the system takes the form

1041
$$E = \frac{1}{2(1 - v^2)}[E_i h \varepsilon_i^2 + E_m h_m \varepsilon_m^2 + E_o h \varepsilon_o^2]$$

1042

1043 *Results.* Minimizing the elastic energy with respect to strain and curvature yields their
1044 values at equilibrium. In particular, using the notation

1045
$$A = \frac{1}{(4E_i E_o h + E_m(E_i + E_o)h_m)} > 0$$

1046 we obtain the equilibrium values for the strain

1047
$$\varepsilon = A(2E_i E_o h(g_i + g_o) + E_m(E_i + E_o)h_m g_m)$$

1048 and the curvature

1049
$$c = \frac{2A}{h+h_m}\left(2E_i E_o h(g_o - g_i) + E_m E_o h_m(g_o - g_m) - E_m E_i h_m(g_i - g_m)\right).$$

1050 This curvature is in general non-zero, the system curves with the abaxial side on the convex side
1051 as soon as the abaxial growth rate is sufficiently larger than the adaxial growth rate.

1052 To determine whether buckling of the abaxial epidermis may occur, we compute the elastic
1053 (residual) strain in this layer at equilibrium:

1054
$$\varepsilon_o = AE_m E_i h_m(2g_m - g_i - g_o).$$

1055 ε_o can become compressive (<0) only if $2g_m - g_i - g_o < 0$. When it falls beyond a threshold
1056 (*110*), this leads to buckling of the abaxial side, with patterns as described (*59*), depending on the
1057 anisotropy of the residual strain.

1058 Interestingly, the elastic strain of the adaxial epidermis,

1059
$$\varepsilon_i = A E_m E_o h_m (2g_m - g_i - g_o)$$

always has the same sign as that of the abaxial epidermis. This means that either the two
epidermal layers are in compression, or the two epidermal layers are in tension. A situation
where one epidermis would be in tension and the other would be in compression does not
correspond to mechanical equilibrium. Nevertheless, the amount of compression may differ
between the two epidermal layers: the ratio of the elastic deformations is related to the ratio of
Young moduli

$\frac{\varepsilon_o}{\varepsilon_i} = \frac{E_i}{E_o}.$

Accordingly, it is possible that only one of the two epidermal layers reaches the buckling
threshold. For instance, a much smaller Young modulus for the outer epidermis compared to the
inner epidermis would lead to buckling of the outer epidermis exclusively.

1070

1071 Finally, we mention that the elastic strain in the mesophyll has an opposite sign

1072
$$\varepsilon_m = -2 A E_i E_o h (2g_m - g_i - g_o).$$

1073 This means that as soon as the epidermis is under compression, the mesophyll is under tension
1074 and vice versa.

1075
1076 **Graph plotting**
1077
1078 Graphs were generated in R (version 4.2.2 (2022-10-31 ucrt)) using ggplot, raw data for which is
1079 provided in Data S1.

 SUPPLEMENTARY FIGURE TITLES AND LEGENDS
1081

Fig. S1. *AS2* overexpression causes characteristic ridges, invaginations, and outgrowths on
the outer epidermis, associated with Main Fig. 1.
(**A** and **B**) Representative optical cross-sections of Stage 11 flowers of (**A**) the WT and (**B**) *as2-*
7D mutant. Images are representative of T2s harboring pAR169 (*pATML1::RCI2A-mCitrine*), an
epidermal layer specific plasma membrane marker. Green and grey represent epidermal cells and
chlorophyll containing mesophyll cells respectively. WT sepals (A) have a smooth shape, while
the *as2-7D* sepal (B) outer epidermis exhibit outward protrusions (ridges), inward indents
(invaginations) and pointed epidermal extensions (outgrowths). Note that the outgrowths lack
mesophyll cells (grey) underneath whereas mesophylls occupy the space beneath ridges and
invaginations.
(**C** and **D**) Whole inflorescence images of (**C**) Col-0 (control) and (**D**) *as2-5D* (in Col-0, isolated
by Wu et. al (*48*)). Images are representative of *n = 3* inflorescences. Scalebar, 1 cm.
(**E** and **F**) Stage 14 flowers of (**E**) Col-0 and (**F**) *as2-5D* mutant, representative of *n = 3* flowers.
Yellow arrowhead points toward an outgrowth. Note the mild nature of outgrowth formation in
as2-5D mutant as compared to *as2-7D* (Fig. 1). Scalebar, 1 cm.
(**G** to **J**) Representative six-week-old adult plants of (**G**) WT, (**H**) *as2-7D* and (**I-J**) *pML1::AS2*
in WT T1s. The transgenics show strong (17/28 plants, panel I) to extreme (9/28 plants, panel J)
developmental defects, categorized qualitatively by the extent of leaf deformity. The transgenics
exhibit outgrowths on the lower (abaxial) surface of the leaves as well, in addition to severely
stunted growth. Scalebar, 1 cm.
(**K** to **N**) Fully formed siliques of (**K**) WT, (**L**) *as2-7D*, (**M-N**) *pML1::AS2* in WT T1s. Note the
outgrowths on the silique epidermis. In panels M and N, siliques are representative of strong and
extreme categories, described for panels I and J respectively. Three siliques on each panel
correspond to three independent plants. Total number of independent transgenics (T1s) assessed
n = 28.

Fig. S2. The outer epidermis of *as2-7D* exhibits loss of stereotypical growth patterns,
associated with Main Fig. 2.
(**A** to **D**) Confocal stack images of (**A**) WT replicate 2, (**B**) WT replicate 3, (**C**) *as2-7D* replicate
2 and (**D**) *as2-7D* replicate 3 developing flowers, transformed with the pLH13 (*35S::mCitrine-*
RCI2A) membrane marker. Images correspond to the 48-hour (left) and 96-hour (right) time
points from WT and *as2-7D* live imaging experiment ($n = 3$). Yellow arrowheads show
outgrowths. Trichomes were removed from the outer sepal epidermis by image processing.
Scalebar, 50 µm.
(**E** to **H**) Cell area growth rates of (**E**) WT replicate 2, (**F**) WT replicate 3, (**G**) *as2-7D* replicate 2
and (**H**) *as2-7D* replicate 3. The heatmap is projected onto the later time point. Red to blue
gradient shows fast to slow growth. Yellow arrowheads point towards the outgrowth which
exhibit localized fast growth. Scalebar, 50 µm.
(**I** to **L**) The cell division rates of (**I**) WT replicate 2, (**J**) WT replicate 3, (**K**) *as2-7D* replicate 2
and (**L**) *as2-7D* replicate 3. Red to blue gradient shows more to less division. Cell division over
0-48 hours is shown on the left and over 48-96 hours on the right. Scalebar, 50 µm.
(**M** and **N**) Graphical representation of (**M**) cell growth rates and (**N**) cell division rates across all
three independent biological replicates of the WT (black dots) and *as2-7D* mutant (red dots). Dot
represents the growth rate (M) and division rate (N) of a given cell lineage. Dot plots are
superimposed with box plots, where the central line indicates the median. Left and right panels
display data for 0-48 and 48-96 hours, respectively. Note the reduced spatial variability in the
growth rates of *as2-7D* (panel M). Lines above data points indicate statistical comparisons. The
p-values for Bonferroni corrected pairwise t-test (two-tailed) are shown. *p-values*: ***<0.0001,
**<0.001, *<0.05, ns = non-significant. Exact *p-values* for growth rates: WT vs *as2-7D* for 0-48
hours, 0.9884; WT vs *as2-7D* for 48-96 hours, 1.565172×10^{-93}. Exact *p-values* for cell division
rates: WT vs *as2-7D* for 0-48 hours, 7.800489×10^{-05}; WT vs *as2-7D* for 48 - 96 hours, 9.47882
$\times 10^{-07}$.

Fig. S3. Slower growth on the inner epidermis together with misdirected outer epidermal
growth trigger buckling in *as2-7D*, associated with Main Fig. 3.
(**A** to **D**) Growth rate heatmaps for (**A**) WT replicate 2, (**B**) WT replicate 3, (**C**) *as2-7D* replicate
2 and (**D**) *as2-7D* replicate 3 developing sepals. Red to blue heatmap gradient shows fast to slow
growth. Outer (left) and inner (right) epidermal growth rates over 48 hours is projected as a
heatmap onto the later time point. Scalebar, 20 µm.
(**E**) Graph shows cell growth rates over a 48-hour period for the WT (black) and *as2-7D* (red)
mutant ($n = 3$ biological replicates). Each dot represents the growth rate for a given cell lineage.
Dot plots are overlayed with boxplots, where the central line represents the median. The *p-values*
for Bonferroni corrected pairwise t-test (two-tailed) are shown. *p-values*: ***<0.0001, **<0.001,
*<0.05, ns = non-significant. Exact *p*-values for epidermal cell growth rate: WT outer vs inner,
5.751359×10^{-35}; *as2-7D* outer vs inner, 3.039502×10^{-25}; WT vs *as2-7D* outer, 0.1011; WT vs
as2-7D inner, 8.496122×10^{-16}.
(**F** to **I**) Cell division heatmaps for (**F**) WT replicate 2, (**G**) WT replicate 3, (**H**) *as2-7D* replicate
2 and (**I**) *as2-7D* replicate 3. Red to blue color gradient shows more to less cell division. The
data is presented in the same format as described for panels A-D. Scalebar, 20 µm.
(**J**) Graph shows cell division rates for the WT and *as2-7D* mutant ($n = 3$ biological replicates)
Dots represent the number of cell divisions for each cell lineage. Data presentation and statistical
comparisons are same as described above for panel E. Exact *p*-values for epidermal cell division
rates: WT outer vs inner, 4.08502×10^{-28}; *as2-7D* outer vs inner, 0.004; WT vs *as2-7D* outer,
3.677026×10^{-23}; WT vs *as2-7D* inner, 1.865456×10^{-06}.
(**K** to **N**) Heatmap representing the angle of deviation of PDG_{max} tensors from the sepal
longitudinal axis for (**K**) WT replicate 2, (**L**) WT replicate 3, (**M**) *as2-7D* replicate 2 and (**N**)
as2-7D replicate 3. Scalebar represents a deviation of 90º in red and 0º (no deviation) in blue.
Scalebar, 20 µm.
(**O**) Graph showing the angle of deviation for the WT and *as2-7D* mutant ($n = 3$ biological
replicates). Each dot represents the angle of deviation for a given cell lineage. Data
representation and statistical comparisons are shown in the same format as described for panels E
and J. Exact *p-values* for epidermal angle of deviation: WT outer vs inner, 3.793354×10^{-05}; *as2-*
7D outer vs inner, 1.572509×10^{-20}; WT vs *as2-7D* outer, 8.052118×10^{-10}; WT vs *as2-7D*
inner, 0.349.

Fig. S4. Slower growth on the inner epidermis leads to overgrowth of the outer epidermis
in *as2-7D*, associated with Main Fig. 3.
(**A** to **D**) Heatmaps represent growth rates of (**A**) WT outer, (**B**) *as2-7D* outer, (**C**) WT inner, and
(**D**) *as2-7D* inner epidermal surface at every 24-hour interval, over a 48-hour live imaging
period. Growth rate is projected onto the later time point. Growth rate for the region demarcated
in cyan at 0-hour time point was tracked over time. Heatmap: red, fast growth, and blue, slow
growth. Representative heatmaps for $n = 3$ biological replicates are shown. Scalebar, 20 µm.
(**E**) Graph shows epidermal surface growth rates every 24 hours for the WT (black) and *as2-7D*
(red) is shown ($n = 3$). Lowercase letters on top of the data points represent significant
differences based on one-way ANOVA with Tukey's HSD post-hoc test ($p < 0.05$, similar letters
represent no significant difference).
(**F**) Graph shows the ratio of the outer vs inner epidermal surface area of WT (black) and *as2-7D*
(red) for each time point ($n = 3$). Note that the outer vs inner ratio of the WT decreases sharply
over time compared to *as2-7D*. Error bars indicate standard error.

Fig. S5. Analysis of cell growth direction and stiffness due to *KRP1* overexpression in WT
and *as2-7D*, associated with Main Fig. 4.
(**A** to **D**) Confocal images of the inner epidermis of (**A**) WT, (**B**) *as2-7D* mutant, (**C**)
AtML1::KRP1 and (**D**) *AtML1::KRP1 as2-7D* stage 14 sepals. The sepals were stained with PI.
Note the smooth morphology of the inner epidermis across all genotypes. Scalebar, 200 µm,
images representative of $n = 3$ adult sepals.
(**E** to **L**) Principal direction of growth (PDG$_{max}$) for the outer epidermal cells of (**E**) WT
Replicate 2, (**F**) *as2-7D* mutant Replicate 2, (**G**) *AtML1::KRP1* Replicate 2, (**H**) *AtML1::KRP1*
as2-7D Replicate 2, (**I**) WT Replicate 3, (**J**) *as2-7D* mutant Replicate 3, (**K**) *AtML1::KRP1*
Replicate 3 and (**L**) *AtML1::KRP1 as2-7D* Replicate 3 sepals. PDG$_{max}$ tensors (yellow)
calculated over a 48-hour live imaging period starting from stage 4, projected on a later time
point surface. Segmented cells in grey are superimposed onto the plasma membrane signal (also
in grey). Scalebar, 20 µm, $n = 3$.
(**M** and **N**) Representative confocal images of WT ($n = 5$) early stage 6 flower (**M**) before and
(**N**) after osmotic treatment in 0.35M NaCl. The PI (cell wall stain) and membrane marker
(pLH13, *35S::mCitrine-RCI2A*) signals have been merged together and shown. The highlighted
region of the outer sepal (white rectangle) indicates the comparable longitudinal section
extracted for analyzing epidermal shrinkage.
(**O** and **P**) MorphoGraphX generated surface of the longitudinal section (**O**) before treatment and
(**P**) after osmotic treatment, extracted from flower images shown in panels M and N respectively.
Comparable regions of the outer and inner sepal epidermis were segmented (outlined in cyan) to
calculate the percentage areal shrinkage.

Fig. S6. Buckling in *as2-7D* is not caused due to disruption of outer (abaxial) – inner (adaxial) polarity.

(**A** to **D**) REVOLUTA (REV) protein reporter (*proREV::REV-mGFP*) shows nuclear localization of REV, restricted exclusively to the inner epidermis of WT and *as2-7D* sepals. (**A** and **B**) Confocal images of inflorescence meristems show adaxial specific REV expression (yellow arrowheads) in the young sepals of (**A**) WT and (**B**) *as2-7D* mutant. (**C** and **D**) Representative middle stage (Stage 8-9) flowers of (**C**) WT and (**D**) *as2-7D* show no detectable REV expression on the outer epidermis of the sepals in the folds or outgrowths. Images of representative T3s (*n* = 3) show mGFP in green and chlorophyll in greyscale. Scalebar, 50 μm.

(**E** to **H**) *FILAMENTOUS FLOWER* (*FIL*) promoter reporter (*proFIL::mCherry-RCI2A*) expression pattern is restricted to the outer sepal epidermis and shows no difference between the WT and *as2-7D* mutant. (**E** and **F**) Confocal images of inflorescence meristems show *FIL* expression pattern in the early floral developmental stages of (**E**) WT and (**F**) *as2-7D* mutant. The membrane localized mCherry is shown in green and chlorophyll in greyscale. (**G** and **H**) Representative middle stage (Stage 7-8) flowers of (**G**) WT and (**H**) *as2-7D* mutant showing the *FIL* promoter reporter signal (mCherry, green). Images are representative of three T3s (*n* = 3). Scalebar, 50 μm.

(**I** to **L**) Whole inflorescence images of (**I**) WT, (**J**) *rev-6* mutant, (**K**) a representative transgenic (T1) harboring *pML1::AS2* transgene in *rev-6* and (**L**) *as2-7D* mutant. Note the outgrowths (yellow arrows) on the sepal outer epidermis of *pML1::AS2 rev-6* are similar to those of *as2-7D* and *pML1::AS2* in WT (Fig. 1M). Multiple independent T1s for *pML1::AS2* in *rev-6* were assayed (*n* = 19), all of which show similar outgrowth formation on the sepal outer epidermis in the *rev-6* background.

Fig. S7. Ectopic *AS2* expression mediated buckles and subsequent outgrowth formation occur independent of *REVOLUTA*.

(**A**) Representation of the previously characterized meristem initiation defect phenotype of *rev-6* mutant. Images show an excised lateral inflorescence meristem of the WT (left) and *rev-6* mutant (right). Majority of the cauline leaves axils (white arrows) in the *rev-6* mutant are bare, as they fail to produce axillary meristem, as described by Otsuga et. al. (*71*). Scalebar, 1cm.

(**B** to **D**) Fully developed siliques of (**B**) *rev-6* and (**C** and **D**) *pML1::AS2* in rev-6 T1s. No of independent T1s assessed *n* = 19. The three siliques correspond to three independent plants. The numbers represent the total number of plants showing strong (12/19 plants, panel C) to extreme (7/19 plants, panel D) phenotypes, categorized based on the extent of leaf deformity in the adult vegetative stage.

(**E** to **I**) Representative six-week-old adult plants of (**E**) WT, (**F**) *rev-6*, (**G**) *as2-7D* and (**H** and **I**) *pML1::AS2* in *rev-6* T1s. Total number of T1s assessed *n* = 19. The transgenics show strong (12/19 plants, panel H) to extreme (7/19 plants, panel I) defects in leaf development, characterized by outgrowth formation on the abaxial epidermis and stunted growth. Scalebar, 1 cm.

(**J** to **M**) Confocal images of PI-stained (greyscale) inflorescences of (**J**) WT, (**K**) representative *pML1::REV* in WT T1, (**L**) *as2-7D* mutant, and (**M**) representative *pML1::REV* in *as2-7D* T1. Note that the transgenics ectopically expressing *REV* looks like their respective backgrounds in either the WT (panels J and K) or *as2-7D* (panels L and M). Yellow arrowheads point to the outgrowths. For each *pML1::REV* transgenics in WT and in *as2-7D* backgrounds, number of independent T1s assessed, *n* = 28. Scalebar, 50 µm.

Fig. S8. Outgrowths on *as2-7D* outer epidermis are formed due to buckling mediated PIN1 convergence and subsequent auxin maxima formation, associated with Main Fig. 5.

(**A** and **B**) Confocal images show the expression pattern of PIN1 protein reporter (*pPIN1::PIN1-GFP*) in (**A**) WT and (**B**) *as2-7D* inflorescence meristems. Note the expression of *PIN1* directed towards the emerging outgrowths on the outer epidermis of *as2-7D* older flowers. Scalebar, 50 µm, $n = 3$.

(**C** and **D**) Expression pattern of auxin response reporter (*pDR5::VENUS*) in stage 11 flower of *as2-7D* mutant. Shown is (**C**) VENUS signal by itself (green) and (**D**) VENUS signal merged with PI (grey). Note the presence of DR5 signal specifically in the outgrowths (yellow arrowheads). Scalebar, 100 µm, $n = 3$.

(**E** to **H**) Representative live-imaging series showing the second replicate of (**E**) Mock treated, (**F**) 10µM NPA treated developing flowers, and third replicate of (**G**) Mock treated, (**H**) 10µM NPA treated developing flowers of *as2-7D* mutant. Images taken at every 24-hour interval over 96-hour period. Note the development of outgrowths from 72-hour time-point onwards (yellow arrowheads) in the mock treated *as2-7D* samples (panels E and G), which are absent in NPA treated *as2-7D* flowers (panels F and H). The *as2-7D* flowers harbor pLH13 (*35S::mCitrine-RCI2A*) cell membrane marker (greyscale). Trichomes were removed from the outer sepal epidermis by image processing. Scalebar, 50 µm.

1263

Supplementary Table S1: List of primers used in this study.

1265

Primer Name	Sequence	Purpose
oLH108	5'-TTAATGCGAATTACTAGTAGAGAGT-3'	F_Primer for PCR genotyping *as2-6D* mutation
oLH324	5'-TGAAGAATTTTGCTCTTTTTATTCGAATT-3'	R_Primer for PCR genotyping *as2-6D* mutation
oLH347	5'-CACCTGGTAGCTAGCGTTGTTGACA-3'	F_Primer for primer for cloning *AS2* (At1G65620) promoter from WT and *as2-7D* into pENTR
oLH370	5'-GAACGTTTGCGAATTTTTGTGG-3'	R_primer for cloning *AS2* (At1G65620) promoter from WT and *as2-7D* into pENTR
oLH111	5'-CACCATGGCATCTTCTTCAACAAACTCAC-3'	F_Primer for Cloning *AS2* CDS into pENTR
oLH112	5'-TCAAGACGGATCAACAGTACG-3'	R_Primer for Cloning *AS2* CDS into pENTR
oXH172	5'-ctctagacacggggtgtttGTGTGAGCCAAAAACGAG-3'	F_Primer for amplifying *proREV::REV* fragment from Ler-0
oXH176	5'-cggcagcagccgcagcagaaacCACAAAAGACCAGTTTACAAAG-3'	R_Primer for amplifying *proREV::REV* fragment from Ler-0
oXH177	5'-aaactggtctttgtggtttctgctgcggctgctgccg-3'	F_Primer for amplifying mGFP for cloning *proREV::REV-GFP*
oXH173	5'-gtggtaccgcggccgcgtttAGGTCACTGGATTTTGGTTTTAG-3'	R_Primer for amplifying mGFP for cloning *proREV::REV-GFP*
oXH174	5'-atatatcctgtcaaagagctTCTTTGTTCTGCGCAACAG -3'	F_Primer for amplifying promoter of *FIL*
oXH178	5'-ggtaattgttgtaaaaatacgcccAGTGAGAAGGGAAAGGGG-3'	R_Primer for amplifying promoter of *FIL*
oXH179	5'-CCCCTTTCCCTTCTCACTgggcgtattttacaacaattacc-3'	F_Primer for amplifying mCherry for cloning *proFIL::mcherry-RCI2A*
oXH175	5'-taaacgcggccgcggtaccaAGGTCACTGGATTTTGGTTTTAG -3'	R_Primer for amplifying mCherry for cloning *proFIL::mcherry-RCI2A*
oAS_005	5'-aaaATGGAGATGGCGGTGGC-3'	F_Primer amplying full length *REVOLUTA* CDS
oAS_006	5'-TCACACAAAAGACCAGTTTACAAAGG-3'	R_Primer amplying full length *REVOLUTA* CDS
oAS_007	5'-TCCGTGAATGTTCCATTTTG-3'	F_Primer *REVOLUTA* qRT-PCR, product size - 129 bp
oAS_008	5'-CAGAGAGCTTCCGGTTTACG-3'	R_Primer *REVOLUTA* qRT-PCR, product size - 129 bp
oAS_017	5'-GTAAAACGACGGCCAG-3'	M13 forward primer
oAS_031	5'-CCACATCCTCTTCCTCACCT-3'	F_Primer on *ATML1* for colony PCR & sequencing

1267

Supplementary Data S1: Raw data for making the plots

1269

Supplementary Movie S1: Confocal images of the outer and inner epidermis of adult sepals of the WT (left) and *as2-7D* mutant (right). Genotypes harbor pLH13 (*p35S::mCitrine-RCI2A*) plasma membrane marker. Scalebar, 200 µm.

Supplementary Movie S2: Confocal images of developing WT (left) and *as2-7D* mutant (right) flowers imaged every 24 hours, over 96-hour time period. Genotypes harbor pLH13 (*p35S::mCitrine-RCI2A*) plasma membrane marker. Trichomes were removed from the outer sepal epidermis by image processing for better visualization of sepal morphology. Scalebar, 50 µm.

Supplementary Movie S3: Confocal images show the outer epidermis of adult sepals of WT (top left), *as2-7D* (top right), *pML1::KRP1* (bottom left) and *pML1::KRP1 as2-7D* (bottom right). Note the largely rescued sepal smoothness in *pML1::KRP1 as2-7D* compared to *as2-7D*, due to an increase in the longitudinally aligned giant cells. Sepals were stained with Propidium-iodide, which stains the cell walls. Scalebar, 200 µm.

Supplementary Movie S4: Confocal images of developing *as2-7D* mutant flowers grown in the presence of mock (left) or 10µM NPA (right). Images were captured every 24 hours over the 96 hour time period. The *as2-7D* mutant harbors pLH13 (*p35S::mCitrine-RCI2A*) plasma membrane marker. Trichomes were removed from the outer sepal epidermis by image processing. Note the reduction in outgrowth formation in 10µM NPA-treated samples despite buckling. Scalebar, 50 µm.

1292

1293

REFERENCES AND NOTES

1. D. P. Richman, R. M. Stewart, J. W. Hutchinson, V. S. Caviness, Jr., Mechanical model of brain convolutional development. *Science* **189**, 18-21 (1975).
2. K. E. Garcia, C. D. Kroenke, P. V. Bayly, Mechanics of cortical folding: stress, growth and stability. *Philos Trans R Soc Lond B Biol Sci* **373**, (2018).
3. E. Hannezo, J. Prost, J. F. Joanny, Instabilities of monolayered epithelia: shape and structure of villi and crypts. *Phys Rev Lett* **107**, 078104 (2011).
4. A. E. Shyer *et al.*, Villification: how the gut gets its villi. *Science* **342**, 212-218 (2013).
5. C. Huang, Z. Wang, D. Quinn, S. Suresh, K. J. Hsia, Differential growth and shape formation in plant organs. *Proceedings of the National Academy of Sciences* **115**, 12359-12364 (2018).
6. C. H. Waddington, Preliminary Notes on the Development of the Wings in Normal and Mutant Strains of Drosophila. *Proc Natl Acad Sci U S A* **25**, 299-307 (1939).
7. J. A. Thomson, On Growth and Form. *Nature* **100**, 21-22 (1917).
8. H. Tsukaya, Leaf shape: genetic controls and environmental factors. *Int J Dev Biol* **49**, 547-555 (2005).
9. H. Tsukaya, Mechanism of leaf-shape determination. *Annu Rev Plant Biol* **57**, 477-496 (2006).
10. C. M. Nelson, On Buckling Morphogenesis. *J Biomech Eng* **138**, 021005 (2016).
11. E. Cerda, L. Mahadevan, Geometry and physics of wrinkling. *Phys Rev Lett* **90**, 074302 (2003).
12. E. Sharon, E. Efrati, The mechanics of non-Euclidean plates. *Soft Matter* **6**, 5693-5704 (2010).
13. A. Boudaoud, An introduction to the mechanics of morphogenesis for plant biologists. *Trends Plant Sci* **15**, 353-360 (2010).
14. B. Li, Y.-P. Cao, X.-Q. Feng, H. Gao, Mechanics of morphological instabilities and surface wrinkling in soft materials: a review. *Soft Matter* **8**, 5728-5745 (2012).
15. S. Emam, W. Lacarbonara, A review on buckling and postbuckling of thin elastic beams. *European Journal of Mechanics - A/Solids* **92**, 104449 (2022).
16. Y. Wang, Z. Chi, J. Liu, On buckling behaviors of a typical bending-dominated periodic lattice. *Composite Structures* **258**, 113204 (2021).
17. J. W. Osborn, A model of growth restraints to explain the development and evolution of tooth shapes in mammals. *J Theor Biol* **255**, 338-343 (2008).
18. V. D. Varner, J. P. Gleghorn, E. Miller, D. C. Radisky, C. M. Nelson, Mechanically patterning the embryonic airway epithelium. *Proceedings of the National Academy of Sciences* **112**, 9230-9235 (2015).
19. T. Savin *et al.*, On the growth and form of the gut. *Nature* **476**, 57-62 (2011).

20. N. L. Nerurkar, L. Mahadevan, C. J. Tabin, BMP signaling controls buckling forces to
modulate looping morphogenesis of the gut. *Proceedings of the National Academy of*
Sciences **114**, 2277-2282 (2017).
21. A. E. Shyer *et al.*, Emergent cellular self-organization and mechanosensation initiate
follicle pattern in the avian skin. *Science* **357**, 811-815 (2017).
22. J. L. Silverberg *et al.*, 3D imaging and mechanical modeling of helical buckling in
Medicago truncatula plant roots. *Proc Natl Acad Sci U S A* **109**, 16794-16799 (2012).
23. G. M. Whiteley, J. S. Hewitt, A. R. Dexter, The buckling of plant roots. *Physiologia*
Plantarum **54**, 333-342 (1982).
24. F. Bizet, A. G. Bengough, I. Hummel, M.-B. Bogeat-Triboulot, L. X. Dupuy, 3D
deformation field in growing plant roots reveals both mechanical and biological
responses to axial mechanical forces. *Journal of Experimental Botany* **67**, 5605-5614
(2016).
25. J. Yan, B. Wang, Y. Zhou, S. Hao, Resistance from agar medium impacts the helical
growth of Arabidopsis primary roots. *Journal of the Mechanical Behavior of Biomedical*
Materials **85**, 43-50 (2018).
26. R. Sachse *et al.*, Snapping mechanics of the Venus flytrap (Dionaea muscipula). *Proc*
Natl Acad Sci U S A **117**, 16035-16042 (2020).
27. Y. Forterre, J. M. Skotheim, J. Dumais, L. Mahadevan, How the Venus flytrap snaps.
Nature **433**, 421-425 (2005).
28. C. A. Airoldi, C. A. Lugo, R. Wightman, B. J. Glover, S. Robinson, Mechanical buckling
can pattern the light-diffracting cuticle of Hibiscus trionum. *Cell Reports* **36**, 109715
(2021).
29. E. Moyroud *et al.*, Cuticle chemistry drives the development of diffraction gratings on the
surface of Hibiscus trionum petals. *Current Biology* **32**, 5323-5334.e5326 (2022).
30. R. L. Antoniou Kourounioti *et al.*, Buckling as an origin of ordered cuticular patterns in
flower petals. *Journal of The Royal Society Interface* **10**, 20120847 (2013).
31. P. B. Green, Pattern Formation in Shoots: A Likely Role for Minimal Energy
Configurations of the Tunica. *International Journal of Plant Sciences* **153**, S59-S75
(1992).
32. J. Dumais, C. R. Steele, New Evidence for the Role of Mechanical Forces in the Shoot
Apical Meristem. *Journal of Plant Growth Regulation* **19**, 7-18 (2000).
33. J. Żebrowski, Dynamic behaviour of inflorescence-bearing Triticale and Triticum stems.
Planta **207**, 410-417 (1999).
34. M. Fournier, J. Dlouhá, G. Jaouen, T. Almeras, Integrative biomechanics for tree
ecology: beyond wood density and strength. *Journal of Experimental Botany* **64**, 4793-
4815 (2013).
35. K. J. Niklas, *Plant biomechanics. An engineering approach to plant form and function*.
(University of Chicago Press, Chicago, 1992), pp. xiii + 607 pp.

36. S. M. Chapotin, J. H. Razanameharizaka, N. M. Holbrook, A biomechanical perspective
on the role of large stem volume and high water content in baobab trees (Adansonia spp.;
Bombacaceae). *Am J Bot* **93**, 1251-1264 (2006).

37. K. J. Niklas, Plant Height and the Properties of Some Herbaceous Stems. *Annals of*
Botany **75**, 133-142 (1995).

38. R. Méndez-Alonzo *et al.*, Root biomechanics in Rhizophora mangle: anatomy,
morphology and ecology of mangrove's flying buttresses. *Annals of Botany* **115**, 833-840
(2015).

39. J. Derr, R. Bastien, É. Couturier, S. Douady, Fluttering of growing leaves as a way to
reach flatness: experimental evidence on <i>Persea americana</i>. *Journal of The Royal*
Society Interface **15**, 20170595 (2018).

40. C. D. Whitewoods, Riddled with holes: Understanding air space formation in plant
leaves. *PLoS Biol* **19**, e3001475 (2021).

41. J. R. Evans, Improving photosynthesis. *Plant Physiol* **162**, 1780-1793 (2013).

42. A. H. K. Roeder, Arabidopsis sepals: A model system for the emergent process of
morphogenesis. *Quantitative Plant Biology* **2**, e14 (2021).

43. L. Hong *et al.*, Variable Cell Growth Yields Reproducible OrganDevelopment through
Spatiotemporal Averaging. *Dev Cell* **38**, 15-32 (2016).

44. E. Semiarti *et al.*, The ASYMMETRIC LEAVES2 gene of Arabidopsis thaliana regulates
formation of a symmetric lamina, establishment of venation and repression of meristem-
related homeobox genes in leaves. *Development* **128**, 1771-1783 (2001).

45. H. Iwakawa *et al.*, The ASYMMETRIC LEAVES2 gene of Arabidopsis thaliana,
required for formation of a symmetric flat leaf lamina, encodes a member of a novel
family of proteins characterized by cysteine repeats and a leucine zipper. *Plant Cell*
Physiol **43**, 467-478 (2002).

46. W. C. Lin, B. Shuai, P. S. Springer, The Arabidopsis LATERAL ORGAN
BOUNDARIES-domain gene ASYMMETRIC LEAVES2 functions in the repression of
KNOX gene expression and in adaxial-abaxial patterning. *Plant Cell* **15**, 2241-2252
(2003).

47. A. Chalfun-Junior *et al.*, ASYMMETRIC LEAVES2-LIKE1 gene, a member of the
AS2/LOB family, controls proximal-distal patterning in Arabidopsis petals. *Plant Mol*
Biol **57**, 559-575 (2005).

48. G. Wu *et al.*, KANADI1 regulates adaxial-abaxial polarity in Arabidopsis by directly
repressing the transcription of ASYMMETRIC LEAVES2. *Proc Natl Acad Sci U S A*
105, 16392-16397 (2008).

49. Y. Matsumura, H. Iwakawa, Y. Machida, C. Machida, Characterization of genes in the
ASYMMETRIC LEAVES2/LATERAL ORGAN BOUNDARIES (AS2/LOB) family in
Arabidopsis thaliana, and functional and molecular comparisons between AS2 and other
family members. *Plant J* **58**, 525-537 (2009).

50. Y. Eshed, S. F. Baum, J. V. Perea, J. L. Bowman, Establishment of polarity in lateral
organs of plants. *Curr Biol* **11**, 1251-1260 (2001).
51. R. A. Kerstetter, K. Bollman, R. A. Taylor, K. Bomblies, R. S. Poethig, KANADI
regulates organ polarity in Arabidopsis. *Nature* **411**, 706-709 (2001).
52. J. L. Bowman, Y. Eshed, S. F. Baum, Establishment of polarity in angiosperm lateral
organs. *Trends Genet* **18**, 134-141 (2002).
53. J. F. Emery *et al.*, Radial patterning of Arabidopsis shoots by class III HD-ZIP and
KANADI genes. *Curr Biol* **13**, 1768-1774 (2003).
54. X. Lin *et al.*, LFR is functionally associated with AS2 to mediate leaf development in
Arabidopsis. *Plant J*, (2018).
55. C. Mollier *et al.*, Spatial consistency of cell growth direction during organ morphogenesis
requires CELLULOSE SYNTHASE INTERACTIVE1. *Cell Rep* **42**, 112689 (2023).
56. S. Harmansa, A. Erlich, C. Eloy, G. Zurlo, T. Lecuit, Growth anisotropy of the
extracellular matrix shapes a developing organ. *Nature Communications* **14**, 1220 (2023).
57. L. Hong *et al.*, Heterogeneity and Robustness in Plant Morphogenesis: From Cells to
Organs. *Annual Review of Plant Biology* **69**, 469-495 (2018).
58. A. W. Sylvester, M. H. Williams, P. B. Green, Orientation of cortical microtubules
correlates with cell shape and division direction. *Protoplasma* **153**, 91-103 (1989).
59. B. Audoly, A. Boudaoud, Buckling of a stiff film bound to a compliant substrate—Part I::
Formulation, linear stability of cylindrical patterns, secondary bifurcations. *Journal of the
Mechanics and Physics of Solids* **56**, 2401-2421 (2008).
60. A. H. Roeder *et al.*, Variability in the control of cell division underlies sepal epidermal
patterning in Arabidopsis thaliana. *PLoS Biol* **8**, e1000367 (2010).
61. S. M. Bemis, K. U. Torii, Autonomy of cell proliferation and developmental programs
during Arabidopsis aboveground organ morphogenesis. *Dev Biol* **304**, 367-381 (2007).
62. A. Sapala, R. S. Smith, Osmotic Treatment for Quantifying Cell Wall Elasticity in the
Sepal of Arabidopsis thaliana. *Methods Mol Biol* **2094**, 101-112 (2020).
63. A. Sapala, R. S. Smith, in *Plant Stem Cells: Methods and Protocols,* M. Naseem, T.
Dandekar, Eds. (Springer US, New York, NY, 2020), pp. 101-112.
64. J. R. McConnell, M. K. Barton, Leaf polarity and meristem formation in Arabidopsis.
Development **125**, 2935-2942 (1998).
65. T. Yamaguchi, A. Nukazuka, H. Tsukaya, Leaf adaxial-abaxial polarity specification and
lamina outgrowth: evolution and development. *Plant Cell Physiol* **53**, 1180-1194 (2012).
66. K. Fukushima, M. Hasebe, Adaxial–abaxial polarity: The developmental basis of leaf
shape diversity. *genesis* **52**, 1-18 (2014).
67. A. Y. Husbands, D. H. Chitwood, Y. Plavskin, M. C. Timmermans, Signals and
prepatterns: new insights into organ polarity in plants. *Genes Dev* **23**, 1986-1997 (2009).
68. R. Waites, A. Hudson, phantastica: a gene required for dorsoventrality of leaves in
Antirrhinum majus. *Development* **121**, 2143-2154 (1995).

69. M. P. Caggiano *et al.*, Cell type boundaries organize plant development. *eLife* **6**, e27421
(2017).
70. S. B. Sun, J. P. Song, L. S. Meng, ASYMMETRIC LEAVES2 gene, a member of
LOB/AS2 family of Arabidopsis thaliana, causes an abaxializing leaves in transgenic
cockscomb. *Mol Biol Rep* **39**, 4927-4935 (2012).
71. D. Otsuga, B. DeGuzman, M. J. Prigge, G. N. Drews, S. E. Clark, REVOLUTA regulates
meristem initiation at lateral positions. *Plant J* **25**, 223-236 (2001).
72. P. Merelo, E. B. Paredes, M. G. Heisler, S. Wenkel, The shady side of leaf development:
the role of the REVOLUTA/KANADI1 module in leaf patterning and auxin-mediated
growth promotion. *Curr Opin Plant Biol* **35**, 111-116 (2017).
73. Q. Chen *et al.*, The Arabidopsis FILAMENTOUS FLOWER gene is required for flower
formation. *Development* **126**, 2715-2726 (1999).
74. S. Sawa, T. Ito, Y. Shimura, K. Okada, FILAMENTOUS FLOWER controls the
formation and development of arabidopsis inflorescences and floral meristems. *Plant Cell*
11, 69-86 (1999).
75. K. Watanabe, K. Okada, Two discrete cis elements control the Abaxial side-specific
expression of the FILAMENTOUS FLOWER gene in Arabidopsis. *Plant Cell* **15**, 2592-
2602 (2003).
76. W. Wang *et al.*, YUCCA genes are expressed in response to leaf adaxial-abaxial
juxtaposition and are required for leaf margin development. *Plant Physiol* **157**, 1805-
1819 (2011).
77. L. Gälweiler *et al.*, Regulation of polar auxin transport by AtPIN1 in Arabidopsis
vascular tissue. *Science* **282**, 2226-2230 (1998).
78. M. G. Heisler *et al.*, Alignment between PIN1 polarity and microtubule orientation in the
shoot apical meristem reveals a tight coupling between morphogenesis and auxin
transport. *PLoS Biol* **8**, e1000516 (2010).
79. A. M. R. Kabir *et al.*, Buckling of Microtubules on a 2D Elastic Medium. *Scientific*
Reports **5**, 17222 (2015).
80. A. Sanyal, H. C. Han, Artery buckling affects the mechanical stress in atherosclerotic
plaques. *Biomed Eng Online* **14 Suppl 1**, S4 (2015).
81. D. Reinhardt *et al.*, Regulation of phyllotaxis by polar auxin transport. *Nature* **426**, 255-
260 (2003).
82. A. H. K. Roeder, Arabidopsis sepals: A model system for the emergent process of
morphogenesis. *Quant Plant Biol* **2**, (2021).
83. S. Friesen, I. K. Hariharan, Coordinated growth of linked epithelia is mediated by the
Hippo pathway. *bioRxiv*, 2023.2002.2026.530099 (2023).
84. A. Y. Husbands, A. H. Benkovics, F. T. Nogueira, M. Lodha, M. C. Timmermans, The
ASYMMETRIC LEAVES Complex Employs Multiple Modes of Regulation to Affect
Adaxial-Abaxial Patterning and Leaf Complexity. *Plant Cell* **27**, 3321-3335 (2015).

85. R. Kelly-Bellow *et al.*, Brassinosteroid coordinates cell layer interactions in plants via
cell wall and tissue mechanics. *Science* **380**, 1275-1281 (2023).
86. N. Nakayama *et al.*, Mechanical Regulation of Auxin-Mediated Growth. *Current Biology*
22, 1468-1476 (2012).
87. P. B. Green, Expression of pattern in plants: combining molecular and calculus-based
biophysical paradigms. *Am J Bot* **86**, 1059-1076 (1999).
88. C. Kuhlemeier, Phyllotaxis. *Trends Plant Sci* **12**, 143-150 (2007).
89. D. R. Smyth, J. L. Bowman, E. M. Meyerowitz, Early flower development in
Arabidopsis. *The Plant Cell* **2**, 755-767 (1990).
90. D. O. Robinson *et al.*, Ploidy and Size at Multiple Scales in the Arabidopsis Sepal. *The*
Plant Cell **30**, 2308-2329 (2018).
91. S. Strauss *et al.*, Using positional information to provide context for biological image
analysis with MorphoGraphX 2.0. *eLife* **11**, e72601 (2022).
92. A. Armezzani *et al.*, Transcriptional induction of cell wall remodelling genes is coupled
to microtubule-driven growth isotropy at the shoot apex in Arabidopsis. *Development*
145, (2018).
93. M. G. Heisler *et al.*, Patterns of Auxin Transport and Gene Expression during
Primordium Development Revealed by Live Imaging of the Arabidopsis Inflorescence
Meristem. *Current Biology* **15**, 1899-1911 (2005).
94. L. Hong, J. Brown, N. A. Segerson, J. K. Rose, A. H. Roeder, CUTIN SYNTHASE 2
Maintains Progressively Developing Cuticular Ridges in Arabidopsis Sepals. *Mol Plant*
10, 560-574 (2017).
95. A. H. Roeder, A. Cunha, C. K. Ohno, E. M. Meyerowitz, Cell cycle regulates cell type in
the Arabidopsis sepal. *Development* **139**, 4416-4427 (2012).
96. W. Lukowitz, C. S. Gillmor, W.-R. d. Scheible, Positional Cloning in Arabidopsis. Why
It Feels Good to Have a Genome Initiative Working for You1. *Plant Physiology* **123**,
795-806 (2000).
97. D. Weigel, J. Glazebrook, Setting up Arabidopsis crosses. *CSH Protoc* **2006**, (2006).
98. D. O. Robinson *et al.*, Ploidy and Size at Multiple Scales in the Arabidopsis Sepal. *Plant*
Cell **30**, 2308-2329 (2018).
99. M. Y. Zhou, C. E. Gomez-Sanchez, Universal TA cloning. *Curr Issues Mol Biol* **2**, 1-7
(2000).
100. M. Karimi, D. Inze, A. Depicker, GATEWAY vectors for Agrobacterium-mediated plant
transformation. *Trends Plant Sci* **7**, 193-195 (2002).
101. M. Zhu *et al.*, Robust organ size requires robust timing of initiation orchestrated by
focused auxin and cytokinin signalling. *Nat Plants* **6**, 686-698 (2020).
102. P. J. Barrell, A. J. Conner, Minimal T-DNA vectors suitable for agricultural deployment
of transgenic plants. *BioTechniques* **41**, 708-710 (2006).
103. A. A. Wise, Z. Liu, A. N. Binns, Three methods for the introduction of foreign DNA into
Agrobacterium. *Methods Mol Biol* **343**, 43-53 (2006).

104.   S. J. Clough, A. F. Bent, Floral dip: a simplified method for Agrobacterium-mediated
1529   transformation of Arabidopsis thaliana. *Plant J* **16**, 735-743 (1998).
1530   105.   G. V. Reddy, M. G. Heisler, D. W. Ehrhardt, E. M. Meyerowitz, Real-time lineage
1531   analysis reveals oriented cell divisions associated with morphogenesis at the shoot apex
1532   of Arabidopsis thaliana. *Development* **131**, 4225-4237 (2004).
1533   106.   C. T. Rueden *et al.*, ImageJ2: ImageJ for the next generation of scientific image data.
1534   *BMC Bioinformatics* **18**, 529 (2017).
1535   107.   M. Zhu, A. H. Roeder, Plants are better engineers: the complexity of plant organ
1536   morphogenesis. *Curr Opin Genet Dev* **63**, 16-23 (2020).
1537   108.   A. Kiss *et al.*, Segmentation of 3D images of plant tissues at multiple scales using the
1538   level set method. *Plant Methods* **13**, 114 (2017).
1539   109.   P. Barbier de Reuille *et al.*, MorphoGraphX: A platform for quantifying morphogenesis
1540   in 4D. *eLife* **4**, e05864 (2015).
1541   110.   Z. Y. Huang, W. Hong, Z. Suo, Nonlinear analyses of wrinkles in a film bonded to a
1542   compliant substrate. *Journal of the Mechanics and Physics of Solids* **53**, 2101-2118
1543   (2005).
1544

Inflorescence **Stage 14 flower** **Outer epidermis** **Inner epidermis**

WT

A

C

s s s

WT

E

F

as2-7D

WT

G

H

as2-7D

as2-7D

B

D

I

-1484 bps

aagaataagaataaa

ATG

AS2 - ASSYMETRIC LEAVES 2

•AS2

Outer
Mesophyll
Inner

•AS2

Outer epidermis

WT **as2-7D**

J

Stage 6 flower

Front view

s
s
s s

Optical
section

Rotated 90°

Longitudinal section

Outer
epidermis

K *pAS2^WT::GUS-GFP* in WT

Outer

Inner

PI

GFP

L *pAS2^as2-7D::GUS-GFP* in WT

Outer

Inner

M *pML1::AS2* in WT T1

N WT

O *pML1::AS2* in WT T1

P *as2-7D*

A WT

0 hr 24 hr 48 hr 72 hr 96 hr

p35S::mCitrine-RCI2A

B *as2-7D*

0 hr 24 hr 48 hr 72 hr 96 hr

p35S::mCitrine-RCI2A

C *as2-7D* 48 hr timepoint
(sideview)

D *as2-7D* 96 hr timepoint

Invaginations
Ridges
} Buckles

Growth Rate

E WT
0 - 48 hrs   48 - 96 hrs

Fast
12
Growth rate heatmap scale
2
Slow

F *as2-7D*
0 - 48 hrs   48 - 96 hrs

Cell division rate

G WT
0 - 48 hrs   48 - 96 hrs

More
10
9
Cell division heatmap scale
1
Less

H *as2-7D*
0 - 48 hrs   48 - 96 hrs

I
Outer E_o g_o
Mesophyll E_m g_m
Inner E_i g_i

J

K

A 0 hr   24 hr   48 hr

WT

B

as2-7D

p35S::mCitrine-RCI2A

C Abaxial/Outer Surface   Adaxial/Inner surface
0 hr   48 hr   0 hr   48 hr

WT

D

as2-7D

Cell growth Rate

E   Outer   Inner

WT

Fast
Growth rate scale
Slow

F   Outer   Inner

as2-7D

More
Cell division scale
Less

Cell division Rate

G   Outer   Inner

WT

H   Outer   Inner

as2-7D

N MBD-mCitrine in WT

O MBD-mCitrine in *as2-7D*

Principal Directions of Maximum Growth

I   Outer   Inner

WT

Longitudinal sepal axis

J   Outer   Inner

as2-7D

Longitudinal sepal axis

K

More
Angle of deviation from longitudinal axis
Less

90°
80°
70°
60°
50°
40°
30°
20°
10°
0°

Longitudinal axis

Angle of deviation

PDG

Cell

Transverse axis

Angle of deviation from longitudinal axis

L   Outer   Inner

WT

M   Outer   Inner

as2-7D

Stage 14 flower

WT

as2-7D

AtML1::KRP1

AtML1::KRP1 as2-7D

Outer epidermis

WT   as2-7D   AtML1::KRP1   AtML1::KRP1 as2-7D

WT   as2-7D

AtML1::KRP1   AtML1::KRP1 as2-7D

Q Before treatment

Epidermal Cell

Cytosol   Membrane   Cell Wall   Shrinkage

R After treatment (0.35M NaCl)

Water   Stiffness

(Cell wall shrinks)

S Cross-section of outer sepal

Outer epidermis   Inner epidermis

T (Shrunk sepal)

More → Less
% Shrinkage in area

WT

as2-7D

AtML1::KRP1

AtML1::KRP1 as2-7D

Y

% Shrinkage in area

a   a   b   c   c   ab   c   a

Outer | Inner   Outer | Inner   Outer | Inner   Outer | Inner

WT   as2-7D   AtML1::KRP1   AtML1::KRP1 as2-7D

pPIN1::PIN1-GFP

A Inflorescence — WT

B *as2-7D*

GFP
PI

C Stage 8 Flower — WT

D Stage 8 Flower — *as2-7D*

GFP
PI

E *as2-7D* (Mock) | 0 hr | 24 hr | 48 hr | 72 hr | 96 hr

p35S::mCitrine-RCI2A

F *as2-7D* (NPA treatment) | 0 hr | 24 hr | 48 hr | 72 hr | 96 hr

p35S::mCitrine-RCI2A

G PDGs (0 - 48 hrs) — *as2-7D* (Mock)

H *as2-7D* (NPA)

I Cell growth Rate (0 - 48 hrs) — *as2-7D* (Mock)

J *as2-7D* (NPA)

K Cell growth Rate (48 - 96 hrs) — *as2-7D* (Mock)

L *as2-7D* (NPA)

Growth rate heatmap scale
Fast 10 — 8 — 6 — 4 — 2 — Slow