Growth directions and stiffness across cell layers determine whether tissues stay smooth or buckle

Avilash S. Yadav[2]†, Lilan Hong[1]†, Patrick M. Klees[2], Annamaria Kiss[3], Manuel Petit[3], Xi He[1], Iselle M. Barrios[2], Michelle Heeney[2], Anabella Maria D. Galang[2], Richard S. Smith[4], Arezki Boudaoud[5], and Adrienne H.K. Roeder[2]*

[1]Institute of Nuclear Agricultural Sciences, Key Laboratory of Nuclear Agricultural Sciences of Ministry of Agriculture and Zhejiang Province, College of Agriculture and Biotechnology, Zhejiang University, Hangzhou 310058, China
[2]Weill Institute for Cell and Molecular Biology and Section of Plant Biology, School of Integrative Plant Sciences, Cornell University, Ithaca, NY 14853, USA
[3]Laboratoire Reproduction et Développement des Plantes, Univ Lyon, ENS de Lyon, UCB Lyon1, CNRS, INRAE, INRIA, F-69342 Lyon, France
[4]The John Innes Centre, Norwich NR4 7UH, UK
[5]LadHyX, Ecole Polytechnique, CNRS, IP Paris, 91128 Palaiseau Cedex, France
†These authors contributed equally to this work.
*Corresponding author. ahr75@cornell.edu

Abstract

From smooth to buckled, nature exhibits organs of various shapes and forms. How cellular growth patterns produce smooth organ shapes such as leaves and sepals remains unclear. Here we show that unidirectional growth and comparable stiffness across both epidermal layers of Arabidopsis sepals are essential for smoothness. We identified a mutant with ectopic *ASYMMETRIC LEAVES 2* (*AS2*) expression on the outer epidermis. Our analysis reveals that ectopic *AS2* expression causes outer epidermal buckling at early stages of sepal development, due to conflicting growth directions and unequal epidermal stiffnesses. Aligning growth direction and increasing stiffness of the outer epidermis restores smoothness. Furthermore, buckling influences auxin efflux transporter protein PIN-FORMED 1 polarity to generate outgrowth in the later stages, suggesting that buckling is sufficient to initiate outgrowths. Our findings suggest that in addition to molecular cues influencing tissue mechanics, tissue mechanics can also modulate molecular signals, giving rise to well-defined shapes.

One-Sentence Summary

The *asymmetric leaves 2-7D* mutant sepals buckle due to discoordination of growth between the two epidermal layers.

INTRODUCTION

During development, many organs such as the brain, intestine, and orchid flower petals form complex shapes with undulations, folds, and loops (*1-6*); conversely, other organs such as Arabidopsis leaves and fly wings develop smooth forms (*7-10*). These undulations, folds, and loops generally form through mechanical buckling, which is the bending of tissue when its growth patterns generate compressive forces (*11-13*). How cells within tissues regulate their growth and division to generate buckling in certain instances and maintain smoothness in others is not well understood.

Buckling plays a significant role in organ morphogenesis and development across organisms. In animals for instance, numerous studies demonstrate the importance of buckling in organ morphogenesis (*2-4, 14, 15*). Differential growth of connected tissue layers gives rise to buckling. Buckling creates complex organ shapes starting from simple geometries, which is important for functional specialization. These include gyrification of the cerebral cortex to increase neuron content (*1*), formation of gut villi and looping to increase contact with digested food and lengthen the gastrointestinal tract (*2-4*), and initiation of new branches in lung airway epithelium to enhance gas exchange and oxygen transfer (*15*). Buckling also serves key roles in plant development. For instance, it generates helical twists in roots to enhance penetrative forces at the growing tip (*16, 17*), triggers lobe deformation in Venus flytraps for rapid energy storage and release to capture prey (*18, 19*), and induces nanoscale cuticular wrinkles on petals to produce iridescence (*5, 6*). Buckling has also been proposed to drive formation and positioning of primordia on the shoot apical meristem (*20, 21*).

Buckling, however, is not always beneficial. Inflorescence weight and external forces such as wind or rain can compress and buckle the stems of cereal crops, thereby causing damage and reduced yield (*22*). Similarly, slender tree stems are also vulnerable to buckling (*23*). To counter this, plants need to allocate resources to increase stem thickness and stiffness (*24, 25*). While buckling generates diverse leaf shapes (*26, 27*), majority of the leaves suppress buckling, generating smooth leaf blades with a relatively flat structure that enhance light absorption and gas exchange for efficient photosynthesis (*28, 29*). Similarly, sepals, which are the outermost leaf-like floral organs, also exhibit smooth surfaces, which is important for all four sepals to properly enclose and protect the developing flower bud (*30*). However, what growth mechanisms and tissue properties underlie the development of smooth organ morphology remain relatively unknown.

Using *Arabidopsis thaliana* (henceforth Arabidopsis) sepals as the model organ, we identified and characterized a gain-of-function mutant of the transcription factor *ASYMMETRIC LEAVES 2* (*AS2*), *as2-7D*, which exhibits folds and outgrowths on the outer epidermis of the sepal. We used

78   *as2-7D* as a tool to determine what cell growth patterns and mechanical properties of the tissue
79   facilitate the formation of smooth sepals in the wild type.
80
81   **RESULTS**
82
83   **Ectopic expression of *ASYMMETRIC LEAVES 2* (*AS2*) disrupts sepal smoothness**
84
To determine how the sepals achieve smooth morphology, we performed a forward genetic
screen (*31*), and identified an Arabidopsis mutant that disrupts sepal smoothness. This mutant
exhibits an irregular morphology with folds and outgrowths (Fig. 1, A to F). The folds comprise
ridges and invaginations, whereas the outgrowths, which are pointed epidermal projections, are
typically located at the end of the folds (Fig. 1F and fig. S1, A and B). These folds and
outgrowths are visible only on the outer (abaxial) epidermis, while the inner (adaxial) epidermis
is smooth like the wild type (Fig. 1, E to H, fig. S1, A and B and movie S1). Positional cloning
identified a G-to-A point mutation 1,484 bps upstream of the translational start site of
ASYMMETRIC LEAVES 2 (*AS2*) (Fig. 1I). *AS2* encodes a LATERAL ORGAN BOUNDARIES
(LOB) domain transcription factor that plays a critical role in leaf development, particularly in
specifying the upper (adaxial) layer of the leaf (*32-34*). An identical mutation in the same locus
was identified previously (*35*). However, the fold and outgrowth phenotype in the sepals was not
reported (*35*), since the phenotype is very subtle in the Columbia (Col-0) background compared
to what we observe in the Landsberg *erecta* (L*er*-0) background (hereafter referred to as wild
type) (Fig. 1, A to D and fig. S1, C to F). Previously, Wu et. al. (*35*) demonstrated that this
mutation in the *AS2* promoter disrupts the binding site of the repressor KANADI, a transcription
factor involved in specifying cell fate on the lower (abaxial) side of the leaf (*36, 37*). They
further showed that this mutation leads to ectopic *AS2* expression on the abaxial side, while
normally in the wild type, the *AS2* expression domain is restricted to the adaxial side (Fig.1I). To
verify that this mutation also leads to ectopic *AS2* expression in sepals of the L*er*-0 background,
we drove GFP expression under the wild type *AS2* promoter (*pAS2^{WT}::GUS-GFP*) and mutant
as2-7D AS2 promoter (*pAS2^{as2-7D}::GUS-GFP*), in the wild type (L*er*-0) background. While the
AS2 promoter from wild type drives GFP specifically on the inner sepal epidermal layer (Fig. 1, J
and K), the mutant *AS2* promoter drives GFP on both the inner and outer epidermal layers (Fig.
1L). The mutant *AS2* promoter drove GFP expression evenly throughout the outer epidermis, and
no instance of patchy *AS2* expression patterns were observed on the outer epidermis. In addition,
AS2 is still expressed in its regular expression domain on the inner epidermis in *as2-7D* mutants
(Fig. 1L). This analysis confirms that the mutation causes a gain-of-function through ectopic *AS2*
expression on the outer epidermis, consistent with the dominant nature of this allele (see
Materials and methods). Therefore, following the convention of previous reports (*35, 38*), we
named the mutant allele *as2-7D*.
116

To test whether ectopic *AS2* expression on the outer epidermal layer is sufficient to generate the
folds and outgrowths, we drove *AS2* expression under epidermal specific *MERISTEM LAYER 1*
(*ML1*) promoter (*pML1::AS2*), which ectopically expresses *AS2* on both inner and outer
epidermis of the wild type. All the transgenic plants form lumps and outgrowths on the outer
epidermis of the sepals, similar to *as2-7D*, which suggests that ectopic outer epidermal
expression of *AS2* is sufficient to replicate the *as2-7D* phenotype (Fig. 1, M to P). Additionally,
we do observe folds and outgrowths form on the epidermis of the leaves and fruits (fig. S1, G to
N), consistent with previous reports (*34*), indicating a potential stronger effect of constitutive
epidermal *AS2* expression in the transgenics than in the *as2-7D* gain-of-function mutant. In
summary, the mutation in *as2-7D* induces ectopic *AS2* expression on the outer epidermal layer,
which is sufficient to disrupt smooth sepal morphology by generating folds and outgrowths on
the outer epidermal layer.
129
130   **The *as2-7D* outer epidermis appears to buckle first and then form outgrowths**
131
To understand how ectopic *AS2* expression on the outer epidermis disrupts smooth sepal
morphology, we live-imaged wild type and *as2-7D* flowers from early sepal developmental
stages (Fig. 2, A and B, fig. S2, A to D, and movie S2). Initially, the outer epidermis of both the
wild type and *as2-7D* sepals are smooth (Fig. 2, A and B; 0-24h). While the outer epidermis of
wild type sepals remains smooth throughout live imaging, *as2-7D* starts to develop a fold in the
middle of the sepal blade at 48 hours (Fig. 2, B and C). As the flower develops, folding
intensifies further, forming several inward invaginations and outward ridges throughout the outer
epidermis of *as2-7D* mutant (Fig. 2, B to D and movie S2). As the folds intensify over time, we
begin to see the development of phenotypically distinct pointed outgrowths. The outgrowths
become apparent only in the later stages as the folds intensify, which suggests that the formation
of folds precedes the emergence of the outgrowths (Fig. 2, B to D and fig. S2, C and D).
143
We considered two hypotheses for how the folds form: (1) due to localized growth at each ridge
or (2) due to mechanical buckling caused by a surface-wide growth disruption. To test the first
hypothesis, we compared the cell growth rates and division rates between wild type and *as2-7D*
mutant on the outer sepal epidermis every 48 hours over 96 hours (Fig. 2, E to H and fig. S2, E
to N). We do not see any localized fast growth or cell division in the region where the first fold is
formed in *as2-7D* at 48 hours (Fig. 2, C to F). In the initial stages (0-48 hrs), the wild type
exhibits the stereotypical basipetal growth pattern described earlier (*39*), exhibiting fast cell
growth at the tip, moderate growth in the center, and slow growth towards the base. In contrast,
the cells grow at relatively homogeneous rates throughout the outer epidermis of *as2-7D* (Fig. 2,
E and F; 0-48 hrs). Although the pattern of growth is different, we observe no difference in the
average cell growth rates between the wild type and *as2-7D* (fig. S2M). This pattern is also
reflected in cell division rates, where we observe a basipetal pattern in wild type and a more
uniform pattern throughout the sepal epidermis in *as2-7D* (Fig. 2, G and H; 0-48 hrs). Our results

suggest that folds form as a result of surface-wide disruption of cell growth patterns, and not
because of localized fast cell growth and division at the location of the ridge. In the later time
point (48-96 hrs) the growth is still relatively uniform along the places where the *as2-7D* folds
form. In contrast to folds, we do observe localized growth and division later at sites where
outgrowths form (Fig. 2F and fig. S2, G and H; 48-96 hrs). We therefore considered the
hypothesis that the folds on *as2-7D* outer epidermis may form through mechanical buckling
caused by surface-wide growth defects.
164
165   **Modeling predicts two criteria required for buckling**
166
Based on our observations of *as2-7D* development, we formulated a simple two-dimensional
(2D) analytical model to determine the basic conditions that could give rise to buckling (see
Materials and methods). Our model indicates that buckling can occur if the following two
conditions are simultaneously fulfilled. Firstly, if the average specified growth rate (the cell-
autonomous growth rate achieved if the layer were isolated) of the outer and inner epidermis is
higher than that of the mesophyll, then the epidermal layers will be under compression, which
favors buckling in the epidermal layers. Secondly, considering that the mesophyll is known to be
softer than the epidermis (*40*), a significantly lower stiffness of the outer epidermis compared to
the inner epidermis may cause the outer epidermis specifically to buckle, without causing any
buckling of the inner epidermis.
177
We further tested this with simulations on a simple geometry: a cross-sectional representation of
a growing elastic system, comprising of the outer epidermis, the mesophyll, the inner epidermis,
and the edges (Fig. 2, I to K, fig. S3 and S4). In these simulations, only the outer epidermis was
allowed to grow in width, which is equivalent to the outer epidermis having a higher specified
growth rate than the other two layers. Under these conditions, when the outer and inner
epidermis exhibit similar stiffness, the tissue curves while maintaining a smooth morphology for
both the outer and inner epidermis (Fig. 2I). However, when the outer epidermis is softer than
the inner epidermis, the outer epidermis buckles, while the inner epidermis shows no change in
morphology (Fig. 2J). In contrast, increasing the stiffness of the outer epidermis relative to the
inner epidermis restores the smooth morphology of the outer epidermis, while the inner
epidermal morphology remains unaffected (Fig. 2K). In summary, although our simple analytical
model and simulations do not account for all the complex dynamics of tissue growth, it guides
our investigations towards testing whether differences in growth and tissue stiffness properties
between the outer and inner epidermal layers contribute to buckling in *as2-7D*.
192
193   **Aligned of cell growth orientation between epidermal layers is essential for the formation**
194   **of smooth sepals**
195

To test the first of our model predictions, we live-imaged the wild type and *as2-7D* sepals
through the pre-buckling stages (Fig. 3, A and B), and extracted the inner and outer epidermal
layers of the same growing sepals by image processing (Fig. 3, C and D). We found that cells
throughout the wild type inner epidermis grow significantly faster than the outer epidermal cells
(Fig. 3E and fig. S5, A, B, and E). We also observe fast cell divisions on the inner epidermis
(Fig. 3G and fig. S5, F, G, and J). The fast growth of the inner epidermis was surprising, since
for the sepals to eventually grow flat, one would intuitively expect the cells on both epidermal
layers to grow at similar rates. However, during the initial developmental stages, the sepals are
highly curved along the longitudinal direction, due to the smaller surface area of the inner
compared to the outer epidermis. Hence, to transition from an initially curved shape to a
subsequently flat structure, the sepal requires faster growth in the inner epidermis compared to
the outer epidermis. (Fig. 3C and D, Fig. 4, and movie S2). Like the wild type, the inner
epidermal cells of *as2-7D* mutant also grow faster than outer epidermal cells; however, the inner
epidermal cells of *as2-7D* do not grow as fast as the wild type (Fig. 3, E and F and fig. S5, A to
E). In concert with the growth rate, the cell division rate is also lower on the inner epidermis of
as2-7D compared to the wild type (Fig. 3, G and H and fig. S5, F to J).
212
Since buckling is a surface-wide phenomenon in our model, we analyzed the differences in
whole inner versus outer epidermal growth rates. On an epidermis wide scale, we found that the
outer epidermis of the wild type and *as2-7D* grow at the same rate (Fig. 4, A, B and E). This
observation is consistent with the growth rate data of individual cell lineages, where the average
outer epidermal cell growth rates of the wild type and *as2-7D* show no significant difference
(Fig. 4, A, B and E and fig. S5E). However, the inner epidermis of *as2-7D* grows significantly
slower than the wild type, which also aligns with the growth rate data of individual cell lineages
on the inner epidermis (Fig. 4, C, D and E and fig. S5E). Our findings indicate that the rate at
which the inner epidermis of *as2-7D* grows is insufficient to keep up with the increasing outer
epidermal surface area (Fig. 4F).
223
We then asked whether outer epidermal buckling in *as2-7D* could potentially be attributed to
local hypergrowth of the mesophyll cells underneath the outer epidermis. To test this, we
processed the sepal images using Convoluted Neural Network (see Materials and methods) (*41*)
to improve the conspicuity of mesophyll cell membranes (fig. S6, A to H). We analyzed the
growth rates of the outermost layer of mesophyll cells which contains fewer airspaces in early
stages, thus allowing us to lineage track the cells over time. Our findings revealed that the spatial
growth patterns of the mesophyll cells of both wild type and *as2-7D* mimic their respective outer
epidermal cell growth patterns (fig. S6, I and J). Specifically, the wild type mesophyll cells
exhibit the same basipetal growth gradient as that of the wild type outer epidermal cells (Fig. 3E
and fig. S6I). The mesophyll cells of *as2-7D* exhibit homogeneous cell growth rates, similar to
the *as2-7D* outer epidermis (Fig. 3F and fig. S6J). Comparable to the outer epidermal cell growth
patterns, no significant difference was found in the average growth rate of the mesophyll cells

between the wild type and *as2-7D* (fig. S6M). Notably, there was no evidence of local
hypergrowth in the central region of the sepal blade, where the initial buckling occurs (fig. S6J).
To test our model predictions, we then compared the magnitude of growth of the outer and inner
epidermal cells relative to the mesophyll (fig. S6N). We found that, the average growth rates of
the inner and outer epidermal cells significantly exceed that of the mesophyll cells for both wild
type and *as2-7D* (fig. S6N), which aligns with our model predictions, suggesting that the
epidermal layers are under compression.
243
Since cell growth directions are also critical for determining organ shapes (*42-44*), we assessed
the principal direction of maximal cell growth (PDG$_{max}$) in the wild type and *as2-7D* mutant. The
wild type cells on both epidermal layers grow in a highly aligned manner along the sepal
longitudinal axis (Fig. 3I). In *as2-7D*, although inner epidermal cells grow parallel to the sepal
longitudinal axis, the outer epidermal cells exhibit highly misaligned growth, with the majority
of PDG$_{max}$ tensors oriented in the transverse direction (Fig. 3J). Similar to the outer epidermal
growth directions, the wild type mesophyll cells grow longitudinally, while the mesophyll cells
of *as2-7D* exhibit misaligned transverse growth (fig. S6, K and L). We quantified the growth
directions of the epidermal cells by calculating the deviation angle of PDG$_{max}$ tensors relative to
the sepal's longitudinal axis (Fig. 3K). The heatmaps reveal minimal deviation in cell growth
directions of the wild type epidermal layers and the inner epidermal layer of *as2-7D* (Fig. 3, L
and M and fig. S5, K to O). In contrast, the outer layer of *as2-7D* displayed a high degree of
deviation in growth directions (Fig. 3M and fig. S5, K to O). Since microtubules are known to
align perpendicular to the maximal cell growth direction (*45*), we then tested the microtubule
orientation pattern in comparable regions of the wild type and *as2-7D* outer epidermis (Fig. 3, N
and O). Microtubules show the typical transverse alignment pattern in wild type, which is
perpendicular to sepal longitudinal growth direction (Fig. 3N). Conversely, the microtubules are
largely misoriented in the *as2-7D,* and in certain instances, mirror the transverse cell growth
pattern (yellow arrowheads, Fig. 3O).
263
Taken together, in the *as2-7D* mutant, buckling occurs on the outer epidermis because it
overgrows in width compared to the inner epidermis. Our observations align with the simulation
prediction that transverse expansion of the outer epidermis, constrained along the edges by the
inner epidermis would favor buckling. Overall, our data indicates that anisotropic growth in an
aligned manner on both the epidermal layers is required for maintaining smooth sepal
morphology.
270
Since we hypothesized that the transversal growth orientation of the outer epidermal cells causes
the outer epidermis to buckle, we tested whether prompting the cells to grow along the
longitudinal direction can rescue the *as2-7D* phenotype. Overexpressing cyclin-dependent kinase
(CDK) inhibitor Kip-related protein 1 (KRP1) throughout the epidermis (*pML1::KRP1*) at
moderate levels trigger epidermal cells to skip division and undergo endoreduplication early

during development, resulting in increased giant cell formation (*46, 47*). Giant cells, exclusive to
the outer epidermis, exhibit the unique property of growing in a highly anisotropic manner (*46*)
along the longitudinal sepal axis (Fig. 5, A, C and E and G, fig. S7, A and C, and movie S3).
Therefore, we wondered whether increasing giant cell formation can restore directional growth
in *as2-7D* and consequently reduce buckling. Indeed, overexpressing *KRP1* in *as2-7D*
(*pML1::KRP1 as2-7D*) results in a notable smoothing of the outer sepal epidermis (Fig. 5, A to
H, fig. S7, A to D and movie S3). As expected, overexpressing KRP1 increases the number and
size of the giant cells, which grow in a highly aligned manner along the longitudinal sepal
direction (Fig. 5, E to P and movie S3). Analysis of the growth directions show that the PDG$_{max}$
tensors of *pML1::KRP1 as2-7D* outer epidermal cells are similar to those of *pML1::KRP1*,
wherein the transverse growth orientation exhibited by *as2-7D* is largely re-oriented
longitudinally (Fig. 5, M to P and fig. S7, E to L). In addition to reduced buckling, the
characteristic outgrowths of the *as2-7D* outer epidermis are also nearly absent in *pML1::KRP1*
as2-7D (Fig. 5, E to H), suggesting that buckling is required for the formation of the outgrowths.
While there is no obvious association between the location of giant cells and small cells and the
buckles in the *as2-7D* mutant, it is possible that the reduction of cell division in *pML1::KRP1*
as2-7D, in addition to the growth orientation contributes to the inhibition of buckling. Overall,
consistent with our hypothesis, our findings suggest that transverse growth of the outer epidermal
cells may be a key driver of buckling in *as2-7D*.
295

296   **Reduced stiffness of the outer epidermis compared to the inner epidermis is associated with**
297   **buckling of the outer epidermis in *as2-7D* mutants**
298

The second condition of our 2D analytical model states that buckling can be restricted to the
outer epidermis if it is softer than the inner epidermis. To test the relative stiffness of the outer
and the inner epidermal layers, we performed osmotic treatments (*48*) and compared the
shrinkage of the wild type and *as2-7D* epidermal layers. When exposed to a salt solution, cells
undergo plasmolysis and no longer exert turgor pressure on the cell wall, causing the cell wall to
contract (Fig. 5, Q and R). The extent of cell wall shrinkage gives a proxy measure for cell wall
stiffness, with low shrinkage indicating the wall is stiff (high Young's modulus) and high
shrinkage indicating the wall is soft (low young's modulus) (*48*). We exposed early sepals to salt
solution and measured the epidermal shrinkage (Fig. 5, S and T, fig. S7, M to P, and see
Materials and methods). We found that the wild type epidermal inner and outer layers exhibit
similar shrinkage, indicating comparable stiffness (Fig. 5, U and Y). In contrast, the outer
epidermis of *as2-7D* shrank more than the inner epidermis, indicating that *as2-7D* outer
epidermis is softer (Fig. 5, V and Y). This observation is also consistent with the second
condition of our analytical model, which proposes that reduced stiffness of the outer epidermis is
an essential factor for buckling (Fig. 2, I and J). We then wondered whether increasing giant cell
formation affects the stiffness of the outer epidermis. To test this, we performed osmotic
treatments on *ML1::KRP1* and *ML1::KRP1 as2-7D*. Indeed, we found increased stiffness of the

outer epidermis in both *ML1::KRP1* and *ML1::KRP1 as2-7D* (Fig. 5, W to Y), which aligns with
our simulation predictions that increasing outer epidermal stiffness may restore smooth
epidermal morphology (Fig. 2K).
319
Taken together, our results suggest that aligning growth along the longitudinal direction,
decreasing cell division, and increasing outer epidermal stiffness can rescue the *as2-7D*
phenotype. However, we cannot also decouple the possibility that accelerated differentiation
resulting from ectopic *KRP1* expression could also potentially contribute as an additional factor
to the rescue. Our findings collectively indicate that relative stiffness and growth orientations of
the epidermal layers play a critical role in shaping sepal morphology. Specifically, unidirectional
growth and comparable stiffness across both epidermal layers are essential criteria for smooth
sepal formation as observed in the wild type.
328
329 **Buckles in early *as2-7D* sepals are not caused by an apparent disruption of abaxial-adaxial**
330 **patterning**
331
It has long been established that the juxtaposition of inner (adaxial) and outer (abaxial) identity
leads to the growth of the leaf blade (*36, 49-53*). Ectopic boundary formation between these
identities can result in outgrowths on the leaf blade that disrupt the smooth leaf surface (*49-52*).
An ectopic boundary can emerge when the inner identity-promoting transcription factors are mis-
expressed on the outer side, creating a juxtaposition between inner and outer identity (*54*). Our
evidence strongly supports our hypothesis that the folds observed in the outer epidermis of *as2-*
7D in the initial stages are in fact buckles caused due to disrupted growth patterns and tissue
properties. However, since *AS2* is known to promote inner epidermal identity (*32-34*), we tested
the alternative hypothesis that the initial folds observed on the outer epidermis of *as2-7D* are
caused by ectopic juxtaposition of inner and outer identity. We therefore looked at the expression
pattern of *REVOLUTA* (*REV*) as an inner identity transcription factor (*55*), and *FILAMENTOUS*
FLOWER (*FIL*) as an outer identity transcription factor (*56*). We observe no difference in *REV*
and *FIL* expression patterns between wild type and *as2-7D*, and particularly, no ectopic
juxtaposition between inner and outer gene expression was formed in the *as2-7D* sepal surface
(Fig. 6, A to H). Since REV promotes adaxial identity in conjunction with AS2, we tested
whether REV is necessary or sufficient for the *as2-7D* phenotype. Ectopic *AS2* expression in the
rev-6 mutant (*pML1::AS2 rev-6*) leads to folds and outgrowth formation similar to *pML1::AS2* in
wild type (Fig. 1, M to P, fig. S1, G to N, Fig. 6, I to L and fig. S8, A to I), indicating REV is not
necessary for either the buckles or the outgrowths. Conversely, ectopic *REV* expression in wild
type sepals is insufficient to generate buckles or outgrowths (fig. S8, J and K). Further, ectopic
REV expression in *as2-7D* also does not enhance the phenotype (fig. S8, L and M).
353
Since abaxial identity promoting KANADI transcription factors directly repress *AS2*, and the
disruption of KANADI binding site leads to ectopic *AS2* expression in *as2-7D* (*35*), we assessed

the sepal phenotypes of *kanadi* loss of function mutants (*kan1-2 kan2-1*). Consistent with
previous reports (*33, 36, 37*), the *kan1-2 kan2-1* double mutants display severe floral
development and phyllotaxy defects, characterized by irregular initiation and arrangement of
sepals, petals and stamens, and conspicuously enlarged carpels (fig. S9, A and B, E to L). In
particular, the *kan1-2 kan2-1* sepals from early developmental stages are strongly radialized with
little sepal lamina, and they fail to enclose the floral bud (fig. S9, F to L). These phenotypes are
distinct from those observed in *as2-7D* mutants and *pML1::AS2* in wild type transgenics, which
show no such defects (Fig. 1 B, M to P and Fig. 2, A and B). While the mature *kan1-2 kan2-1*
sepals are curved downwards, indicative of strong adaxialization, we do not observe the strong
folds and associated outgrowths on the outer epidermis of *kan1-2 kan2-1* sepals unlike *as2-7D*
(Fig. 1 E and F, fig. S9, C and D). However, as reported previously (*32, 33, 36, 37, 57*), we do
observe the folds and outgrowths on the abaxial epidermis of the *kan1-2 kan2*-1 leaves, similar to
AS2 constitutive overexpression phenotypes in the leaves (fig. S9, M to O). The positions of
outgrowth formation typically overlap with the region where the folds form (fig. S9O), which
suggests a potential link between fold formation and ectopic lamina outgrowth.

Overall, our investigations indicate that the outer epidermal buckling exhibited by *as2-7D* sepals
is not caused by an obvious inner-outer polarity disruption which is consistent with our
hypothesis that buckling in *as2-7D* sepals is indeed attributable to tissue-wide growth defects.
However, our investigations do not rule out the possibility that other polarity-associated genes
are disrupted in *as2-7D*.

Buckling in *as2-7D* generates outgrowths via ectopic PIN convergence driving auxin
maxima formation

In addition to the buckles, we observe the emergence of pointed outgrowths at the ends of the
buckles in the later stages in *as2-7D* sepals (Fig. 1F). While constitutive *AS2* overexpression has
previously been reported to cause outgrowths (*34*), what local signal triggers the formation of
these outgrowths at specific sites remains elusive. In previously characterized outgrowths,
expression of auxin biosynthesis genes has been reported (*58, 59*), suggesting auxin plays a role.

The polar localization of PIN-FORMED 1 (PIN1) protein, an auxin efflux transporter, in the
plasma membrane on a particular side of the cell determines the directionality of auxin flow.
When PIN1 polarity from multiple neighboring cells converges to the same location, auxin
accumulates at that site, creating an auxin maximum (*60*). PIN1 was previously shown to be
expressed in the outgrowths formed on the abaxial epidermis of *kanadi* mutant leaves, indicating
that outgrowth formation involves PIN1 (*59, 61*). To test whether outgrowths formed on the
outer epidermis of *as2-7D* sepals also involve PIN1 convergence, we analyzed PIN1-GFP
localization patterns in the wild type and *as2-7D* mutant. In the inflorescence and early-stage
flowers, the PIN1-GFP accumulation pattern in *as2-7D* is similar to that of the wild type (Fig. 7,

A and B). However, in the later stages we observe PIN1-GFP convergence specifically in the
sites where the outgrowths are formed (Fig. 7, C and D, fig. S10, A and B). In contrast, no PIN
convergence sites are observed in the middle of the wild type sepals (Fig. 7, C and D, fig. S10, A
and B). We also observe the expression of auxin-responsive promoter DR5 specifically where
outgrowths are formed in *as2-7D* (fig. S10, C and D), which is consistent with the fact that *PIN1*
convergence is essential for creating local auxin peaks (*62*).
402
The polarity of PIN1 is mechano-sensitive in nature and responds to mechanical stress (*63-65*).
In the later stages when *as2-7D* mutants exhibit both buckles and outgrowths, we observe PIN1
convergence specifically in the outgrowths and not the buckles (Fig. 7, C and D, fig. S10, A and
B). Since buckling inherently alters mechanical stress distributions (*66, 67*), and buckling
precedes outgrowth formation, we hypothesized that buckling induced mechanical stress causes
PIN1 convergence subsequently leading to outgrowth formation in *as2-7D*. To test our
hypothesis, we live imaged growing *pPIN1::PIN1-GFP as2-7D* sepals from initial stages prior to
buckling, through the stages where the outer epidermis buckles and outgrowths are evident (Fig.
8). Initially, when the sepal outer epidermis uniformly curves outwards, no ectopic PIN1
convergence is observed on the sepal blade (Fig. 8, A to D). As the sepal grows, while buckle
formation occurs in the middle of the sepal blade, we do not observe PIN1 expression in the
region where the buckle forms (Fig. 8, E and F). Although outgrowth formation is not evident at
this stage, we do observe PIN1 convergence at the tip of the buckle, which marks the site of
outgrowth initiation (Fig. 8, E to H). In the following time point, an outgrowth is evident in the
region where PIN1 convergence site was formed in the preceding stage (Fig. 8, I to L). As the
outgrowth progresses, the site of PIN1 convergence shifts towards the growing tip of the
outgrowth, while PIN1 expression continues to be absent in the region where the sepal buckles
(Fig. 8, M to P). Collectively, our observations suggest that the tissue first buckles and then PIN1
convergence occurs at the end of a buckle, which precedes the formation of the outgrowths. Our
data is consistent with the hypothesis that buckling induced mechanical stress may cause PIN1 to
converge and initiate an outgrowth.
424
We next tested whether PIN-mediated polar auxin transport is required specifically for the
formation of outgrowths and not buckles. We treated growing *as2-7D* inflorescences with polar
auxin transport inhibitor N-1-naphthylphthalamic acid (NPA) and compared the phenotype to
mock-treated samples. While *as2-7D* live-imaged in the presence of NPA exhibited
characteristic buckles, the formation of outgrowths was largely attenuated (Fig. 7, E and F, fig.
S10, E to H, and movie S4). Buckling continues to occur since the characteristic growth patterns
and misdirected sideways PDG_{max} tensors leading to buckling remain unaltered with NPA
treatment (Fig. 7, G to J). However, the localized fast growth leading to outgrowth formation
was largely reduced in the NPA-treated samples (Fig. 7, K and L). Our results indicate that polar
auxin transport is critical for the formation of outgrowth characterized by localized fast growth.
Overall, our results are consistent with our hypothesis that mechanical buckling causes ectopic

PIN1 convergence, generating local auxin maxima formation and fast growth, leading to

outgrowth formation.

DISCUSSION

While buckling is critical for generating complex organ shapes required for specialized

functional roles (*1-6, 15, 16*), buckling is not always advantageous, and needs to be

circumvented to generate smooth organs like sepals (*68*). Using Arabidopsis sepals as the model,

our study suggests that unidirectional growth of both the outer and inner epidermis along the

sepal longitudinal axis, and similar tissue stiffness across the epidermal layers are key drivers of

smooth sepal formation. We show that ectopic *AS2* expression on the outer epidermis of *as2-7D*

creates an imbalance of growth and tissue stiffness across outer and inner epidermal layers,

resulting in outer epidermal buckling.

Overexpression of *AS2* leading to the formation of ectopic outgrowths has been documented by

previous studies (*32, 34, 69*). However, we identify buckling mediated fold formation as an

upstream morphological deformity that occurs prior to outgrowth initiation in the sepals. PIN1

convergence sites form specifically at the tip of buckles, marking the site of outgrowth initiation.

Since 1995 (*52*), the formation of ectopic outgrowths has primarily been attributed to

juxtaposition of abaxial-adaxial identities. It is possible that the outgrowths observed in the later

stages, classically characterized as ectopic laminas, exhibit juxtaposed abaxial-adaxial identities

as they continue to develop further. However, our study investigated the factors that lead to early

stage buckling in *as2-7D* sepals, and how buckling triggers initiation of outgrowth via ectopic

PIN1 convergence. Probably due to the growth dependent nature of the phenotype and its

occurrence at very early stages of development, the buckling phenomenon was previously

uncharacterized.

Although one would intuitively expect *AS2* ectopic expression to affect only the outer epidermis,

we find the growth rate and stiffness of the inner epidermis, where *AS2* is normally expressed is

affected as well in *as2-7D*. Changes in inner epidermal growth properties in *as2-7D* could be

attributed to the fact that the outer and inner epidermis are connected via the mesophyll and at

the sepal margins. Altering growth patterns in the outer epidermis creates mechanical constraints

on the inner epidermis, thereby affecting its growth indirectly. Simply put, the outer epidermis of

as2-7D buckles as the sideways growth of the outer epidermis is limited by the inner epidermis.

Conceptually, this resembles trying to fit a cloth inside a box. If the cloth fits perfectly along the

edges, it remains smooth; otherwise, the cloth must wrinkle (buckle) to fit inside the box. Our

results show how the growth of an epidermal cell layer (or speculatively an animal epithelium

(*70*)) relative to the tissue to which it is attached, drives organ shape.

While how molecular cues influence tissue mechanics has been relatively well studied, how
tissue mechanics inform molecular signals remain elusive. Our study shows that, not only does
AS2 influence tissue mechanics generating buckles, but also that buckles feed back into
molecular signals. We show that buckling influences the polarity of the PIN1-auxin efflux
transporter. Our findings support the previous observations that mechanical perturbations affect
PIN1 polarity (*63, 65, 71*). We further show that buckling is sufficient to converge PIN1 polarity
causing the formation of an auxin maxima to subsequently trigger epidermal outgrowth
initiation. There is a long-standing debate as to whether primordial initiation in the SAM is also
influenced by buckling, a theory originally proposed by Paul Green (*20, 21, 72*). Contradicting
evidence shows that PIN1 polarizes before primordia bumps are visible (*73*). Our collective
evidence suggests that buckling might be sufficient to converge PIN1 and generate outgrowth.
486

ACKNOWLEDGEMENTS

We thank Isabella Burda, Frances Clark, Shuyao Kong, Byron Rusnak, Maura Zimmermann,
Erich Schwarz, Phoebe Roeder, Scott D. Emr and Stéphane Verger for their valuable comments
and suggestions on the manuscript. We thank Olivier Hamant for the *pPDF1::mCitrine-MBD*
microtubule marker plasmid. We thank John Paul Alvarez and John Bowman for sharing *kan1-2*
kan2-1/+ seeds. We thank Soren Strauss for his guidance with image processing. We thank
Mingyuan Zhu for his input on osmotic treatment experiments. We thank Trinh Duy Chi for his
input on imaging adult sepals. Finally, we thank Arabidopsis Biological Resource Center for
providing seed stocks for this work.

FUNDING
Research reported in this publication was supported by the National Institute of General Medical
Sciences of the National Institutes of Health under Award Number R01GM134037 to AHKR.
French National Research Agency (ANR, grant ANR-21-CE30-0039-01 GrowFlat, to AB). 2023
Sam and Nancy Fleming Research Fellowship to ASY. National Natural Science Foundation of
China grant no. 32270867 and Hundred-Talent Program of Zhejiang University to LH.
Cornell China Center and Zhejiang University for a joint seed fund grant to AHKR and LH.
IMB was supported by NSF REU Award (1850796).

The authors acknowledge the use of facilities and instrumentation supported by NSF through the
Cornell University Materials Research Science and Engineering Center DMR-1719875. The
content is solely the responsibility of the authors and does not necessarily represent the official
views of the National Institutes of Health or other funders.
510

AUTHOR CONTRIBUTIONS

Conceptualization: AHKR, ASY, LH. Experimentation: ASY, LH, PMK, XH, MH. Imaging:
ASY, LH, MH, PMK. Image processing: ASY, LH, PMK, IMB, AMG, RSS. Data analysis:
ASY, LH, AK. Analytical modeling: AK, AB. Simulations: AK, MP, AB. Funding acquisition:
AHKR, AB, LH. Project administration: AHKR, AB. Supervision: AHKR, AB, LH. Writing –
original draft: ASY, LH, AHKR. Writing – review & editing: ASY, LH, PMK, AK, XH, IMB,
MH, AMG, RSS, AB and AHKR.

COMPETING INTERESTS
Authors declare that they have no competing interests.

DATA AND MATERIALS AVAILABILITY
The raw datasets used for making the graphs are provided in Supplementary Data S1. Primers
used for this study are available in Table S1 in Supplementary materials. The transgenic lines for
live imaging purposes generated in this study have been deposited with the Arabidopsis
Biological Resource Center (ABRC), associated with the following accession numbers: CS73559
(pLH13 L*er*-0), CS73560 (pLH13 *as2-7D*), CS73561 (pLH13 *pML1::KRP1* L*er*-0) and
CS73562 (pLH13 *pML1::KRP1 as2-7D*). The meshes and images are publicly available at osf.io
(https://osf.io/p5q39/). The R codes used for generating the plots are available at Github
(https://github.com/Avilash1993/R_codes_for_plots_as2-7D_sepal_buckling) under GNU
General Public License v3.0. Numerical model implementation and the codes for exploration of
the parameter space are available under LGPL license at
https://gitlab.inria.fr/mosaic/publications/growth_buckling.

Fig. 1. The ectopic expression of *ASYMMETRIC LEAVES 2* (*AS2*) on the outer epidermal layer disrupts sepal smoothness in *as2-7D* mutant.

(**A** and **B**) Whole inflorescences of (**A**) Wild type (WT; L*er*-0) and **B**) *as2-7D* mutant (in L*er*-0 background). Yellow arrowheads indicate outgrowth. Scalebar, 1 cm; representative images of *n* = 3 inflorescences for both genotypes.

(**C** and **D**) Mature flowers (stage 14 (*74*)) of (**C**) WT and (**D**) *as2-7D* mutant. WT sepals (green leaf-like organs) are marked with the letter "S" in white (C). Flowers are enclosed within four sepals. Note the outgrowths (yellow arrowheads) on the outer epidermis of *as2-7D* sepals (D). Scale bar, 1 cm; representative images of *n* = 3 flowers.

(**E** and **F**) Scanning electron microscopy (SEM) images of the outer epidermis of (**E**) WT and
(**F**) *as2-7D* mutant sepals. Note the folds (*****) and outgrowths (yellow arrowheads) on the *as2-7D*
outer epidermis. Scale bar, 100 μm; representative images of *n* = 5 for WT and *n* = 10 for *as2-*
7D.
(**G** and **H**) Confocal microscopy images of the inner epidermis of (**G**) WT and (**H**) *as2-7D*
mutant. The images are representative of T2s harboring pLH13 (*35S::mCitrine-RCI2A*), a yellow
fluorescent protein-based cell membrane marker (*75*). Scale bar, 100 μm; representative images
of *n* = 3 for both genotypes.
(**I**) Schematic diagram representing the genetic mutation in *as2-7D*. Gene body of *AS2* showing
the G-to-A point mutation (red arrow) at position 1484 bps upstream of ATG in *as2-7D* (Top
half). The *AS2* expression (red dots) domain in the WT and *as2-7D* sepals is shown (Bottom
half).
(**J**) Anatomical diagram representing (Top) an early stage 6 flower and (Bottom) an optical
longitudinal section through the flower. The letter "S" in black represents sepals. The outer and
inner epidermal layers of the outer sepal are marked.
(**K** and **L**) Longitudinal section of stage 6 flowers of WT. Cell walls are stained with propidium-
iodide (PI, greyscale). Images show the GFP expression pattern (green) driven by (**K**) WT *AS2*
promoter (*pAS2^{WT}*) and (**L**) *AS2* promoter from *as2-7D* (*pAS2^{as2-7D}*). Arrows indicate the outer
and inner epidermal layers of the outer sepal. Scale bar = 20 μm; *n* = 3.
(**M**) Whole inflorescence of a representative transgenic (T1) harboring *pML1::AS2* (*AS2*
expression driven by *ATML1* epidermal specific promoter) transgene in WT. Yellow arrows
mark the outgrowths on the sepal epidermis. Independent T1s assessed = 28, all of which show
similar levels of outgrowth formation. Representative image of *n* = 3 inflorescences.
(**N** to **P**) Confocal images of PI-stained (greyscale) inflorescences of (**N**) WT, (**O**) representative
transgenic (T1) harboring *pML1::AS2* transgene in WT, and (**P**) *as2-7D* mutant. WT outer
epidermis is smooth, whereas *pML1::AS2* exhibits outgrowths (yellow arrowheads) like *as2-7D*.
Scale bar, 50 μm; *n* = 3 inflorescences per genotype.
575

Fig. 2. The *as2-7D* sepal outer epidermal morphology transitions from smooth to buckles and subsequently outgrowths.

(**A** and **B**) Representative live-imaging series of (**A**) a WT flower and (**B**) an *as2-7D* mutant flower starting from developmental stage 4 over a course of 96 hours, imaged at 24-hour intervals ($n = 3$). Images capture morphological changes of the outer sepal. Both genotypes harbor the pLH13 (*35S::mCitrine-RCI2A*) cell membrane marker (greyscale). Trichomes were removed from the outer sepal epidermis by image processing to better visualization the sepal morphology. Scale bar, 50 μm.

(**C**) Tilted view of *as2-7D* mutant flower at the 48-hour time point. Note the appearance of the out-of-plane protrusion (ridge) in the middle of the sepal, circled in white. Scale bar, 50 μm.

(**D**) The *as2-7D* mutant flower at the 96-hour time point. Note the dramatic increase in the number and extent of ridges (red circles) and invaginations (white circles), characteristic of buckles, on the outer sepal epidermis. Yellow arrowheads indicate outgrowths. Scale bar, 50 μm.

(**E** and **F**) Cell area growth rate over a 48-hour period for the (**E**) WT and (**F**) *as2-7D* mutant sepals. Growth rate is calculated as the ratio of cell lineage area at the later time point divided by that of the earlier time point. The heatmap is projected onto the later time point. Red to blue gradient shows fast to slow growth. The growth rate over 0-48 hours (left) and 48-96 hours (right) is shown. Arrowheads point to the outgrowths. Scale bar, 50 μm.

(**G** and **H**) Cell division rate over a 48-hour period for the (**G**) WT and (**H**) *as2-7D*. Cell division rate is calculated based on the number of daughter cells per lineage, represented as a heatmap projected onto the later time point. Red to blue gradient shows more to less division. The cell lineages were tracked using MorphoGraphX software (*76*). Segmented cells are outlined in black. Representative heatmaps for one of the three replicates ($n = 3$) for WT and *as2-7D* are shown. Scale bar, 50 μm.

(**I** to **K**) Simulations based on the analytical model (see Materials and methods), providing visual illustrations of the conditions that may favor buckling of the outer epidermis. The simulations illustrate change in tissue morphology along the cross-section when the outer epidermis was allowed to grow in width over two time points. The elastic modulus (E) of the outer epidermis (E_o), inner epidermis (E_i), mesophyll (E_m) and edges (E_{end}) are shown. Initial growth tensor along the width for all the four components were set to 1, with the outer epidermal growth rate set to 0.1, while the growth rates of the other components was set to 0. (**I**) Simulation outcomes when the outer and inner epidermis have similar stiffness (E_o and $E_i = 10^2$), causing the tissue to curve. (**J**) When the stiffness of the outer epidermis ($E_o = 10^2$) is less than that of the inner epidermis ($E_o = 10^6$), the outer epidermis buckles, while the inner epidermis remains unaffected. (**K**) Simulation outcomes wherein the stiffness of the outer epidermis ($E_o = 10^6$) is more than that of the inner epidermis ($E_o = 10^2$), leading to a slight increase in tissue curvature over time.

Fig. 3. Unidirectional growth on both epidermal layers is essential for smooth sepal development.

(**A** and **B**) Representative confocal images of the (**A**) WT and (**B**) *as2-7D* mutant live-imaged with a high laser intensity (~16-20%) every 24 hours. Both the genotypes harbor pLH13 (*35S::mCitrine-RCI2A*) cell membrane marker (green). Scale bar, 20 μm. One of the three representative replicates (*n* = 3) is shown.

(**C** and **D**) Segmented cells in grey superimposed onto the plasma membrane signal (also in grey) projected on mesh surfaces generated using MorphoGraphX (*76*). Mesh surfaces for 0 and 48-hour time points of the outer (left) and inner (right) epidermis of the (**C**) WT and (**D**) *as2-7D* mutant are shown. Note the strikingly reduced number of cells on the inner versus outer surface at the 0-hour time point. Scale bar, 20 μm; *n* = 3.

(**E** and **F**) Cell growth rates over a 48-hour period for the (**E**) WT and (**F**) *as2-7D* mutants are shown. The heatmap has been projected onto the 48-hour time-point surface. In each panel, the growth rate of cells on the outer surface (left) and inner surface (right) is shown. Red to blue heatmap gradient shows fast to slow growth. Scale bar, 20 μm; *n* = 3.

(**G** and **H**) Cell division rates over a 48-hour period for the (**G**) WT and (**H**) *as2-7D* mutant on the outer surface (left) and inner surface (right) are shown. Segmented cells are outlined in black. Red to blue gradient shows more to less/no cell division respectively. Scale bar, 20 μm; *n* = 3.

(**I** and **J**) Principal directions of maximal growth (PDG$_{max}$, yellow) of the cells on the outer (left) and inner (right) epidermis of (**K**) WT and (**L**) *as2-7D*. Segmented cells in grey are superimposed on the plasma membrane signal (also in grey). Length of any given PDG$_{max}$ tensor is directly proportional to the cell growth rate in that direction. A blue arrow indicates the longitudinal sepal axis. Note the misoriented PDG$_{max}$ tensors on the *as2-7D* outer epidermis. Scale bar, 20 μm; *n* = 3.

(**K**) A schematic diagram depicting the calculation of the angle of deviation between PDG$_{max}$ tensor and the longitudinal axis to obtain a quantitative measure of the deviation.

(**L** and **M**) Heatmaps show the angle of deviation of the PDG$_{max}$ tensor from the sepal longitudinal axis for the outer (left) and inner (right) epidermal cells of the (**L**) WT and (**M**) *as2-7D* mutant. Scale for the angle of deviation ranges from 90º (highly deviated, red) to 0º (no deviation, blue). Scale bar, 20 μm; *n* = 3.

(**N** and **O**) Cortical microtubule orientation (greyscale) from comparable regions of stage 6 flower outer epidermis for the (**I**) WT and (**J**) *as2-7D* mutant is shown. These plants harbor *pPDF1::mCitrine-MBD*, a YFP based microtubule marker (*77*). Images show MorphoGraphX generated meshes from PI-stained images, onto which the microtubule signal has been projected. Segmented cells are outlined in cyan. White arrows point towards the longitudinal axis of the sepals. Microtubules in the WT are oriented perpendicular to the sepal longitudinal axis, whereas, in *as2-7D*, the microtubules are misoriented. Yellow arrowheads point towards the misoriented microtubules. Scale bar, 10 μm; *n* = 3.

Fig. 4. Slower growth on the inner epidermis leads to overgrowth of the outer epidermis in *as2-7D*.

(**A** to **D**) Heatmaps represent growth rates of (**A**) WT outer, (**B**) *as2-7D* outer, (**C**) WT inner, and (**D**) *as2-7D* inner epidermal surface at every 24-hour interval, over a 48-hour live imaging period. Growth rate is projected onto the later time point. Growth rate for the region demarcated in cyan at 0-hour time point was tracked over time. Heatmap: red, fast growth, and blue, slow growth. Representative heatmaps for *n* = 3 biological replicates are shown. Scalebar, 20 μm. (**E**) Graph shows epidermal surface growth rates every 24 hours for the WT (black) and *as2-7D* (red) is shown (*n* = 3). Lowercase letters on top of the data points represent significant differences based on one-way ANOVA with Tukey's HSD post-hoc test (*p* < 0.05, similar letters represent no significant difference).

670
671
Fig. 5. Aligning growth along the longitudinal direction and increasing outer epidermal
stiffness is sufficient to rescue buckling in *as2-7D*.

(**A** to **D**) Representative stage 14 flowers of (**A**) WT, (**B**) *as2-7D* mutant, (**C**) *AtML1::KRP1,* and
(**D**) *AtML1::KRP1 as2-7D*. The WT sepals are marked with the letter "S" in white (A). Yellow
arrowheads point to outgrowths in *as2-7D* (B). Note the smooth sepal of *AtML1::KRP1 as2-7D*
compared to *as2-7D*. Scale bar, 1 cm; *n* = 3 flowers for all genotypes.
(**E** to **H**) Confocal images of the outer epidermis of (**E**) WT, (**F**) *as2-7D* mutant, (**G**)
AtML1::KRP1, and (**H**) *AtML1::KRP1 as2-7D* stage 14 sepals. The sepals were stained with PI
(grey, cell walls). Note the increased number of highly elongated giant cells in *AtML1::KRP1*
and *AtML1::KRP1 as2-7D* sepal, and the smooth outer epidermis of *AtML1::KRP1 as2-7D* sepal
compared to the highly rugose morphology of *as2-7D*. Scale bar, 200 µm; *n* = 3.
(**I** to **L**) Magnified insets of the (**I**) WT, (**J**) *as2-7D* mutant, (**K**) *AtML1::KRP1,* and (**L**)
AtML1::KRP1 as2-7D sepal for the regions marked by white boxes in panels E-H respectively.
Asterisks denote a typical giant cell. Scale bar, 200 µm; *n* = 3.
(**M** to **P**) The principal direction of growth (PDG$_{max}$) for the outer epidermal cells of the (**M**)
WT, (**N**) *as2-7D,* (**O**) *AtML1::KRP1,* and (**P**) *AtML1::KRP1 as2-7D* sepals. Yellow lines mark
the PDG$_{max}$ tensors calculated over a 48-hour period, projected onto the later time point. For WT
and *as2-7D*, the flower images used are the same as in Figure 2. Segmented cells in grey are
superimposed onto the plasma membrane signal (also in grey). Scale bar, 20 µm; *n* = 3.
(**Q** and **R**) Schematic diagrams demonstrating the concept of cell wall stiffness assessment by
osmotic treatment assays (*48*). (**Q**) A typical plant epidermal cell before osmotic treatment. The
cell wall and cell membrane are shown in black and green respectively. (**R**) A cell that has
undergone plasmolysis in a hyperosmotic solution (0.35 M NaCl).
(**S** and **T**) Anatomical diagrams demonstrate using osmotic treatments to estimate epidermal
stiffness. Longitudinal sections of an early stage 6 flower (**S**) before and (**T**) after osmotic
treatment are shown. The outer (green) and inner (blue) sepal epidermis are marked. Arrowheads
(red) demonstrate aerial shrinkage.
(**U** to **X**) Heatmaps show the aerial shrinkage (in percentage) of a longitudinal sepal cross-section
for the inner and outer epidermis of (**U**) WT, (**V**) *as2-7D* mutant, (**W**) *AtML1::KRP1* and (**X**)
AtML1::KRP1 as2-7D, post osmotic treatment in 0.35M NaCl solution. Shrinkage was computed
as the decrease in area of a comparable cross-sectional region from an early stage 6 flower before
and after osmotic treatment. Heatmaps are projected on the post-treatment surface. More
shrinkage is represented in red and less in blue. Scale bar, 20 µm. Heatmaps for each genotype
are representative of 5 independent replicates (*n* = 5).
(**Y**) Graph showing the percentage aerial shrinkage post osmotic treatment. Data across 5
independent replicates each for the WT (black dots), *as2-7D* (red dots), *AtML1::KRP1* (grey
dots), and *AtML1::KRP1 as2-7D* (pink dots) for both the outer and inner epidermis are shown.
The letters above the plots represent significant differences based on one-way analysis of
variance (ANOVA) with Tukey's honest significant difference (HSD) post hoc correction ($p <$
0.05). Similar letters indicate no significant difference.
712

713
714
Fig. 6. Buckling in *as2-7D* is not caused due to disruption of outer (abaxial) – inner
(adaxial) polarity.
(**A** to **D**) REVOLUTA (REV) protein reporter (*proREV::REV-mGFP*) shows nuclear localization
of REV, restricted exclusively to the inner epidermis of WT and *as2-7D* sepals. (**A** and **B**)
Confocal images of inflorescence meristems show adaxial specific REV expression (yellow
arrowheads) in the young sepals of (**A**) WT and (**B**) *as2-7D* mutant. (**C** and **D**) Representative
middle stage (Stage 8-9) flowers of (**C**) WT and (**D**) *as2-7D* show no detectable REV expression
on the outer epidermis of the sepals in the folds or outgrowths. Images of representative T3s (*n* =
3) show mGFP in green and chlorophyll in greyscale. Scalebar, 50 µm.
(**E** to **H**) *FILAMENTOUS FLOWER* (*FIL*) promoter reporter (*proFIL::mCherry-RCI2A*)
expression pattern is restricted to the outer sepal epidermis and shows no difference between the

WT and *as2-7D* mutant. (**E** and **F**) Confocal images of inflorescence meristems show *FIL*

expression pattern in the early floral developmental stages of (**E**) WT and (**F**) *as2-7D* mutant.

The membrane localized mCherry is shown in green and chlorophyll in greyscale. (**G** and **H**)

Representative middle stage (Stage 7-8) flowers of (**G**) WT and (**H**) *as2-7D* mutant showing the

FIL promoter reporter signal (mCherry, green). Images are representative of three T3s ($n = 3$).

Scalebar, 50 µm.

(**I** to **L**) Whole inflorescence images of (**I**) WT, (**J**) *rev-6* mutant, (**K**) a representative transgenic

(T1) harboring *pML1::AS2* transgene in *rev-6* and (**L**) *as2-7D* mutant. Note the outgrowths

(yellow arrows) on the sepal outer epidermis of *pML1::AS2 rev-6* are similar to those of *as2-7D*

and *pML1::AS2* in WT (Fig. 1M). Multiple independent T1s for *pML1::AS2* in *rev-6* were

assayed ($n = 19$), all of which show similar outgrowth formation on the sepal outer epidermis in

the *rev-6* background.

pPIN1::PIN1-GFP

Inflorescence
A — WT
B — *as2-7D*

GFP
PI

Stage 8 Flower
C — WT

Stage 8 Flower
D — *as2-7D*

GFP
PI

E — *as2-7D* (Mock)
0 hr   24 hr   48 hr   72 hr   96 hr
p35S::mCitrine-RCI2A

F — *as2-7D* (NPA treatment)
0 hr   24 hr   48 hr   72 hr   96 hr
p35S::mCitrine-RCI2A

PDGs (0 - 48 hrs)
G — *as2-7D* (Mock)
H — *as2-7D* (NPA)

Cell growth Rate (0 - 48 hrs)
I — *as2-7D* (Mock)
J — *as2-7D* (NPA)

Cell growth Rate (48 - 96 hrs)
K — *as2-7D* (Mock)
L — *as2-7D* (NPA)

Growth rate heatmap scale
Fast
Slow

Fig. 7. Mechanical buckling drives ectopic PIN convergence creating auxin maxima, causing fast local growth and subsequent outgrowth formation.

(**A** to **D**) Confocal stack images showing the expression pattern of auxin efflux carrier protein PIN-FORMED 1 (PIN1) in the WT and *as2-7D*. (**A** and **B**) Expression pattern of PIN1 protein reporter (*pPIN1::PIN1-GFP* (*78*)) in the inflorescence meristem of (**A**) the WT, (**B**) *as2-7D* mutant. The *PIN1* expression pattern in *as2-7D* is identical to WT in the young floral development stages, wherein PIN1 polarity is directed towards the insipient primordia and growing sepal edges, as described previously (*78*). Scalebar, 50 µm. (**C** and **D**) Expression pattern of *PIN1* in a representative middle stage (stage 8) flower of the (**C**) WT and (**D**) *as2-7D* mutant. Note the expression of *PIN1* at the growing tip of the outgrowths (yellow arrows) in *as2-7D*, whereas no *PIN1* expression is detected in the blade region where the epidermis buckles. In panels A to D, the images are representative of three *pPIN1::PIN1-GFP* and *pPIN1::PIN1-GFP as2-7D* plants (*n* = 3). PI is shown in greyscale and GFP in green. Trichomes were removed from the outer sepal epidermis by image processing. Scale bar, 50 µm.

(**E** to **F**) Representative live-imaging series of (**E**) Mock treated *as2-7D* and (**F**) 10µM NPA treated *as2-7D* developing flowers over 96 hours, imaged at every 24-hour interval (representative of *n* = 3 flowers). Note the development of outgrowths from 72-hour time-point onwards (yellow arrowheads) in the mock-treated *as2-7D*, which are absent in NPA treated *as2-7D* flowers. Plants harbor pLH13 (*35S::mCitrine-RCI2A*) cell membrane marker (greyscale). Scale bar, 50 µm.

(**G** to **H**) The PDG_{max} tensors (yellow) of the cells on the outer epidermis of (**G**) mock-treated and (**H**) 10µM NPA treated *as2-7D* over the first 48-hour growth period for the representative samples shown in panels E to F. Characteristic misoriented PDG_{max} tensors (relative to the longitudinal sepal axis) on the outer epidermis of *as2-7D* remain unaffected in the NPA-treated samples, thus driving buckling in later stages. Scale bar, 50 µm.

(**I** to **L**) Outer epidermal cell growth rates over (**I** and **J**) 0–48-hour growth period and (**K** and **L**) 48–96-hour growth period of the mock and 10µM NPA treated *as2-7D* flowers respectively. Heatmaps are projected onto the later time point. Fast (red) to slow (blue) growth rates are shown. In panel K, note the fast cell growth driving outgrowth formation (yellow arrowheads) which is absent in NPA treated sample (panel L). Scale bar, 50 µm.

Fig. 8. PIN1-GFP converges at the end of a buckle where an outgrowth subsequently forms in *as2-7D* mutants.

Confocal live imaging series of a *pPIN1::PIN1-GFP as2-7D* flower, imaged every 12 hours starting from stage 3 (representative of *n = 3*). PIN1-GFP (green) and chlorophyll autofluorescence (gray) are shown. We analyzed the curvature (minimal curvature), in which

warm colors denote out-of-plane (bulging) and cool colors represent inward (saddling)
deformations. Scalebars, 20 μm.
(**A** to **D**) Confocal image of *pPIN1::PIN1-GFP as2-7D* stage 4 flower at 24-hour time point (**A**),
and the analysis of its minimal curvature (**B**) is shown. At this stage, the sepal exhibits uniform
outward curvature. The inset (**C**) and the cross-section through the inset (**D**) for the region
highlighted in panel A (yellow box) depicts the site where an outgrowth will form in the later
stages. Note the absence of *PIN1-GFP* expression at this site (white arrow).
(**E** to **H**) The *pPIN1::PIN1-GFP as2-7D* flower at the 36-hour time point as it transitions into
stages 6 (**E**), and its minimal curvature (**F**) is shown. At this stage, the initial buckle (marked by
a white circle) appears as an out-of-plane fold in the center of the sepal blade (deep red on the
curvature). No *PIN1* expression is observed at one end of the buckle. The inset (**G**) and the
cross-section through the inset (**H**) for the region highlighted in panel **E** (yellow box) is shown.
As the outer layer buckles, *PIN1* expression begins to converge in the highlighted region (white
arrow), marking the site of the later outgrowth.
(**I** to **L**) The *pPIN1::PIN1-GFP as2-7D* flower at the 48-hour time point (**I**), and its minimal
curvature (**J**) is shown. At this stage, the outgrowth begins to form (yellow arrow pointing
towards the red bulge) at the end of the buckle. The inset (**K**) and its corresponding cross-section
(**L**) for the region highlighted in panel E (yellow box) shows the *PIN1* expression associated
with the insipient outgrowth (white arrow).
(**M** to **P**) The *pPIN1::PIN1-GFP as2-7D* flower at the 84-hour time point as it transitions into
stages 9 (**M**), and its minimal curvature (**N**) is shown. The inset (**O**) and the cross-section
through the inset (**P**) for the region highlighted in panel **M** (yellow box) where the outgrowth has
already formed, is shown. By this stage, *PIN1* expression (white arrow) converges towards the
growing tip of the outgrowth.

MATERIALS AND METHODS

Plant Materials

Arabidopsis thaliana genotypes used in this study are in Landsberg *erecta* (L*er*) background, which is referred to as the wild type (WT), unless otherwise mentioned. The mutants *rev-6* (*55*) and *as2-5D* mutant (in Col-0 background) (*35*) have been previously described, and were obtained from Arabidopsis Biological Resource Center (ABRC). The *kan1-2 kan2-1* mutant previously described (*79*) was a kind gift from John Alvarez (Monash University). The plants are homozygous for *kan1-2* while *kan2-1* is segregating; the phenotypes were assayed for plants that are homozygous for both *kan1-2* and *kan2-1*. The *as2-7D* mutant in the L*er*-0 background was identified from ethyl methanesulfonate (EMS) based forward genetic screen. Briefly, M2 seeds (second generation) obtained from self-fertilized EMS mutagenized L*er* plants were purchased from Lehle Seeds (*80*). M2 plants were screened under a stereo microscope for sepal defects (*46, 81*). The *as2-7D* mutant was isolated based on the lumpy sepal phenotype. A mapping population was generated by crossing the *as2-7D* mutant with a Columbia (Col-0) accession plant. Standard map-based cloning techniques were used to clone the underlying mutation (*82*). The *as2-7D* mutation is a G to A change at base 1,484 upstream of the ATG translational start site of AT1G65620/*AS2*. To reduce background mutations, the *as2-7D* mutant was backcrossed twice to wild type L*er* plants twice before further analysis. In the backcrosses, we found that *as2-7D* mutant behaves as a dominant allele as the F1s exhibit the *as2-7D* phenotype, although it is not as strong. The *as2-7D* mutation can be PCR-genotyped by amplifying with primer pairs oLH108 and oLH324 at 55°C annealing temperature, followed by digesting the product with EcoRI to produce an 82-bp WT product or a 111-bp mutant product. Primers are listed in Supplementary Table S1.

Plant Growth

Plants were grown in Percival (Percival-Scientific) growth chambers programmed for long day conditions (16-hour light/8-hour dark cycles). Growth temperature (22ºC), relative humidity (65%) and fluorescence intensity (~100 µmol m^{-1}s^{-1}) were maintained.

For soil-based experiments, plants were grown on soil premixed with Vermaculite (LM111). For stratification, the trays were placed in a 4 ºC cold room for three days prior to moving them to the growth chambers. For BASTA (glufosinate ammonium) selection, the soil was autoclaved twice (Program 3- 15 min sterilization, 30 min dry, Primus ST-1 Autoclave (Model PSS5-B-MSSD)), and sterilized seeds were planted at a density of 50-100 seeds per pot. For live-imaging and adult-stage phenotyping experiments, one plant was grown per pot (6.5×8.5 inches) to facilitate healthy vegetative growth.

For plate-based experiments, seeds were washed twice with a sterilization solution containing
70% ethanol + 0.1% Triton X-100, replacing the solution after each wash. Two final rounds of
washes were provided with 100% ethanol. Seeds were spread on filter paper (Whatman 1,
diameter 90 mm) and allowed to dry inside a sterile safety cabinet (Class II, Type A2, Labgard).
Dried seeds were evenly distributed on freshly prepared MS plates (1/2 MS basal salt mixture,
0.05% MES w/v, 1% sucrose w/v, pH ~ 5.8 (adjusted using KOH) containing 0.8% phytoagar.
The plates were sealed with 3M micropore tape, wrapped in aluminum foil, and stratified for
three days at 4 ºC before being moved to the growth chamber.
864
865 **Phenotyping whole inflorescences, flowers, and plants**
866
867 Whole inflorescence and stage 14 flower images were taken either using Canon PowerShot A640
868 10MP Digital camera attached to Zeiss Stemi 2000-C stereo microscope or Accu-Scope Excelis
869 4K UHD camera attached to Zeiss Stemi 508 stereo microscope. For inflorescence images, and
870 Stage 14 flowers an optical zoom of 1.0 and 1.6 were used respectively. Whole plant images
871 were captured using iPhone XR with a 26 mm focal length and aperture of f/1.8.
872
873 **Schematic diagrams**
874
875 The schematic diagrams shown in Fig. 1I and Fig. 4 Q to T were hand-drawn using the
876 Notability application in Apple iPad (model A1701).
877
878 **Genetic crosses**
879
To generate *pATML1::KRP1 as2-7D*, the *pATML1::KRP1* in L*er* background (*46, 47*) were
crossed into *as2-7D* mutants. Segregation analysis and confocal microscopy was used to find
pATML1::KRP1 as2-7D plants. The *pPIN1::PIN1-GFP DR5::VENUS* (*78*) plants and
pPIN1::PIN1-GFP pin1-4 (*78*) plants were crossed into *as2-7D* mutants. The *pPIN1::PIN1-GFP*
pin1-4 plants (control) exhibit a mild phyllotaxy defect. Genetic segregation analysis and
confocal microscopy was used to find *DR5::VENUS as2-7D* and *pPIN1::PIN1-GFP as2-7D*
plants.
887
888 **Cloning overexpression constructs and reporters**
889 The pLH13 (*p35S::mCitrine-RCI2A,* hygromycin resistance) and pAR169 (*pATML1::RCI2A-*
890 *mCtrine*, basta resistance) plasma membrane markers have been described previously (*46, 83*).
891 The *pPDF1::mCitrine-MBD* microtubule marker (*77*) was a kind gift from Olivier Hamant (RDP
892 institute, ENS de Lyon, France).
893
For cloning *pML1::AS2*, the AT1G65620/*AS2* coding sequence (600 bps) was amplified from
Col-0 cDNA using primers oLH111 and oLH112 by Phusion PCR (NEB), and cloned into

pENTR™/D-TOPO™ (Invitrogen™). The *AS2* coding sequence was cloned into the *pATML1*
containing gateway vector pAR176 (Basta resistance (*46*)) by LR reaction.

For *pML1::REV*, the AT5G60690/*REV* full length coding sequence spanning 2,529 bp was
amplified from Col-0 inflorescence cDNA using primer pairs oAS_005 and oAS_006. The
resulting PCR product was subcloned into pCR™8TOPO™ vector (Invitrogen™) by TA cloning
(*84*). LR reaction was used to clone the *REV* coding sequence into pAR176.

To clone *pAS2 ^{WT}::GFP-GUS*, the 4.1 kb upstream promoter region and part of the first exon of
AS2 was PCR amplified from L*er*-0 genomic DNA using primer pairs oLH347 and oLH370, and
cloned into pENTR™/D-TOPO™ vector (Invitrogen™). The resultant vector was LR
recombined into the gateway vector pBGWFS7 (*85*). For cloning *pAS2^{as2-7D}::GFP-GUS,* the 4.1
kb upstream promoter region and part of the first exon of *AS2* was PCR amplified from *as2-7D*
genomic DNA, and the subsequent cloning steps were similar to cloning *pAS2::GFP-GUS.*

For *proREV::REV-GFP*, the 7.75 kb *proREV::REV* fragment (amplified from L*er*-0) and 1.08
kb mGFP-T35S fragment amplified from *pML1::H2B-mGFP* (pAR180 (*46*)) were assembled
into the PmeI site of pMOA33 (Kanamycin resistance).

For cloning *proFIL::mCherry-RCI2A*, a 3.92 kb *FIL* promoter fragment was amplified from L*er*-
0. This fragment was assembled with a 2.19 kb *omega-mCherry-RCI2A-T35S* fragment
amplified from *p35S::mCherry-RCI2A* (pMZ11 (*86*)) into the SacI and SpeI site of pMOA34
destination vector (hygromycin plant resistance, (*87*)). All the intermediate and final constructs
were verified for sequence and directionality by sequencing. Primers are listed in Supplementary
Table S1.

Plant transformation and selection

The binary vector constructs were transformed into *Agrobacterium tumefaciens* GV3101 strain
(Rifampicin and Gentamycin resistant) by triparental mating method (*88*). Briefly, two *E.coli*
strains, one containing the binary vector and the other containing the helper (conjugative)
plasmid, and *Agrobacterium* strain GV3101 were mated together. The GV3101 clones were
streaked onto Rifampicin + Gentamycin + binary vector specific antibiotic plates for selection,
and positive clones were isolated by colony PCR. The T-DNAs were transformed into plants by
Agrobacterium-mediated floral dipping. Transformants were selected either on ½ MS media
plates (for Hygromycin and Kanamycin selection) or on soil (for Basta selection). For selecting
pLH13 in WT and *as2-7D*, *proFIL::mCherry-RCI2A* in WT and *as2-7D* T1s, a final
concentration of 25 -30 µg/mL Hygromycin was used. For selecting *proREV::REV-mGFP* in
WT and *as2-7D* T1s, a final concentration of 50 µg/mL Kanamycin was used. Healthy, non-
chlorotic plants were transplanted to the soil post two weeks of growth. For selecting *pML1::AS2*

in WT and in *rev-6*, *pML1::REV* in WT and in *as2-7D*, *pAS2^{WT}::GFP-GUS* and *pAS2^{as2-}*
^{7D}::GFP-GUS in WT, *pPDF1::mCitrine-MBD* in WT and in *as2-7D* T1s, BASTA selection was
performed on soil grown plants. Seedlings aged 1 to 2 weeks were selected with 100 µg/mL
BASTA solution by spraying, and non-chlorotic plants were transplanted to fresh pots after a
week. At least twenty independent T1 plants were characterized for each construct. For
fluorescent reporter lines, T1 plant was chosen for each line based on fluorescence. Confocal
microscopy was used to assess the stability of fluorescence in every generation.
943
944 **Flower staging**
945
946 Floral staging was performed based on Smyth et al (*74*). Briefly, sepal primordia barely arise in
947 Stage 3, and overlie the floral meristem in Stage 4. Sepals enclose the floral bud completely by
948 late stage 6. Flowers beyond stage 6 were staged based on size and morphology (*30*).
949
950 **Sepal selected for microscopy**
951
For flower imaging across experiments (live imaging and single time point reporter imaging), we
focused primarily on the outer sepal of the flower, unless specified. Briefly, flowers are enclosed
within four sepals, wherein the foremost sepal, which faces away from the meristem is denoted
as the outer sepal, whereas the sepal that is diametrically opposite to the outer sepal and facing
towards the meristem is denoted as the inner sepal. The remaining two sepals positioned on
either side are the lateral sepals. For a detailed description, please refer to Roeder et. al (*30*).
958
959 **Scanning electron microscopy (SEM)**
960
961 Scanning electron microscopy was performed as described previously (*46*). Briefly, Stage 14
962 flowers were fixed in an FAA solution (50% ethanol, 5% acetic acid, and 3.7% formaldehyde)
963 for 4 hours, dehydrated using an ethanol series, and critical point dried. Sepals were dissected,
964 sputter-coated with platinum palladium and then imaged using a LEICA 440 scanning electron
965 microscope.
966
967 **Confocal Microscopy**
968
Live imaging
For live imaging sepal development, primary inflorescences from six- to eight-week-old plants
containing the *p35S::mCitrine-RCI2A* membrane marker (pLH13 in WT, *as2-7D*, *pML1::KRP1,*
and *pML1::KRP1as2-7D* T2s) were excised. Dissection was performed as described previously
(*86*), with modifications. Briefly, siliques were removed using surgical scissors (Artman
Instruments). Flowers older than stage 7 were dissected using Dumont tweezers (Electron
Microscopy Sciences, Style 5) under a stereo microscope (ZEISS Stemi 2000). Dissected

inflorescences were inserted upright into small petri-plates containing live imaging media (1/2
MS media supplemented with 1X Gamborg Vitamin mix and 1000-fold diluted plant
preservative mixture) with 1.2% w/v phytoagar. The plates were filled with autoclaved water and
allowed to stand for two-three hours for the dissected inflorescence to rehydrate. Residual buds
were dissected down to Stage 3-4 in water using a needle (BD Micro-Fine^{TM} IV Insulin
Syringes). Inflorescences were moved to fresh media plates and transferred to the growth
chamber for at least 24 hours for recovery from dissection, prior to imaging. Stage 4 flowers
were selected as the 0- hour time point across all live imaging series.
984
Post recovery, dissected tissues for live imaging were positioned at an angle to ensure the outer
sepal of the chosen stage 4 flower directly faces the objective of the confocal microscope. The
flowers were submerged in MillQ water containing 0.01% v/v SILWET L-77 surfactant and
imaged using Zeiss 710 upright confocal microscope using 20X water dipping objective (Plan-
Apochromat 20x/1.0 DIC D=0.17 M27 75mm). The following settings were used: mCitrine
(yellow fluorescent protein), excitation laser 514 nm (intensity- 2.4-6.0%), emission collection
range 519-596 nm and a z-step of 0.415-0.420 μm. For high laser intensity based live imaging of
pLH13 in WT and in *as2-7D* to visualize the adaxial sepal epidermis, imaging settings for
mCitrine used: excitation laser 514 nm (intensity- 16-18%), emission collection range 519-596
nm and a z-step of 0.244 μm. Post imaging, the plates were returned to the growth chamber and
the process was repeated for each time-point. Inflorescence tissues were transferred to fresh
media every 48 hours to prevent fungal contamination.
997
998 **NPA treatment assay**
999
Primary inflorescences of six- to eight-week-old pLH13 in *as2-7D* T2s were dissected down to
stage 3 and were divided into two groups of 5-6 samples. One group of samples were placed in ½
MS live imaging media plates containing 10 μM NPA. The other group, serving as the control,
was placed in media plates containing an equivalent amount of DMSO (20 μL DMSO per 100
ml media). The tissues were dissected 48 hours prior to imaging. Post dissection, the tissues were
treated every 12 hours with a 100 μL liquid solution, applied directly above the inflorescence
tissue by pipetting. For the 10 μM NPA treatment samples, the liquid solution comprised of 1 μL
of 1M NPA+19 μL DMSO+10 μL Silwet in 100 ml water, whereas the control (mock) samples
were treated with 20 μL DMSO+10 μL Silwet in 100 ml water. The aim of administering liquid
solution was to enhance the effectiveness of the treatment, which was performed every 12 hours
throughout the experimental duration. Stage 4 flowers, 48 hours post dissection were selected as
the 0- hour time point for imaging mock and 10 μM NPA treated samples. Live imaging was
performed as described above, with the samples being moved to fresh mock or 10 μM NPA
plates every 48 hours. Imaging settings used: mCitrine (yellow fluorescent protein), excitation
laser 514 nm (intensity -2.6-2.8%), emission collection range 519-596 nm, and a z-step of 0.415
μm.

1016
1017 **Propidium Iodide (PI) cell wall staining and imaging**
1018
A 5 mg/ml stock solution was prepared for PI staining, with 5 mgs of PI (Sigma) solubilized in 1
mL of MilliQ water. The stock solution was stored in a foil-covered 1.7 ml microcentrifuge tube
and refrigerated at 4ºC. For staining, stock solution was five-fold diluted with water containing
0.01% v/v Silwet surfactant (PI working solution). Whole inflorescences and flowers below
Stage 10 were stained for 6-8 minutes, stage 14 sepals were stained for 12 minutes, and leaves
were stained for 25 minutes. Care was taken to ensure that the tissue was fully submerged in the
staining solution, and no air bubbles surrounded the tissue. Following staining, the tissue was
washed in a fresh solution containing just 0.01% v/v Silwet, prior to imaging. Fresh PI working
solution was prepared before every experiment. For imaging PI-stained inflorescences of WT,
as2-7D and the transgenics (T1s) of *pML1::AS2* in WT, *pML1::REV* in WT, *pML1::REV* in *as2-*
7D and the inflorescences, flowers of *kan1-2 kan 2-1* mutants, the following settings were used:
excitation 514 nm (laser intensity -18-22%), emission 566-650 nm. For imaging sections of PI-
stained *kan1-2 kan 2-1* mutant leaves, samples were excited with 514 nm (laser intensity - 70%)
and emission was collected between 566-643 nm range.
1033
1034 **Imaging reporters**
1035
For single time point imaging of the reporters, inflorescences were dissected down to stage 3-4.
The recovery period was extended to 48 hours to allow ample time for stabilization of the
reporter signal. The inflorescences were placed upright in the live imaging media plates with the
meristem facing the objective. Prior to taking images of late-stage flowers, the inflorescences
were allowed to grow for a few days. For the reporters imaged together with PI, fresh
inflorescences were dissected for imaging late-stage flowers. Images were taken with Zeiss 710
upright confocal microscope using 20X water dipping objective.
1043
The following settings specific for each of the reporters were used. For *pAS2 ^{WT}::GFP-GUS* and
pAS2 $^{as2-7D}$::GFP-GUS images, PI - excitation laser 594 nm (intensity- 2%), emission range 599-
650 nm and GFP - excitation laser 488 nm (intensity-20%), emission range 493-550 nm, with z-
step of 0.240 μm. For *proREV::REV-GFP*, Chlorophyll and GFP were excited with 488 nm laser
(intensity- 40%), emission range was collected between 647-722 nm for Cholorophyll and 493-
550 nm for GFP. Z-step, 0.240 μm (inflorescences), 0.4 μm (flowers). For *proFIL::mCherry-*
RCI2A, Chlorophyll and mCherry were excited with 561 nm laser (intensity- 12-15%), emission
range was collected between 655-722 nm for Chlorophyll and 578-650 nm for mCherry. Z-step,
0.240 μm. For *pPIN1::PIN1-GFP* single time-point images, PI - excitation laser 594 nm
(intensity- 2%), emission collection range 599-650 nm and GFP - excitation laser 488 nm
(intensity- 25-30%), emission collection range 493-550 nm, with z-steps of 0.240-0.5 μm. For
pPIN1::PIN1-GFP as2-7D live-imaging, GFP – excitation laser 488 nm (intensity – 20-25%),

emission collection range 493-538 nm and Chlorophyll – excitation laser 488 nm (intensity – 20-25%), emission collection range 647-722 nm, with z-steps of 0.5 μm. For *pDR5::VENUS*, samples were excited with an argon laser (514 nm), and data were collected in the YFP (520–565 nm) channelFor *pPDF1::mCitrine-MBD,* PI and mCitrine were excited with 514 nm (intensity-30-32%), and emission range was collected between 603-643 nm for PI and 519-567 nm for mCitrine. Z-step, 0.2 μm.

Imaging adult sepals

For imaging fully mature sepals, all flowers were dissected off the inflorescence except for a single stage 14 flowers. All the internal floral organs from the stage 14 flower were carefully dissected off, leaving just the outer sepal (*30*) attached to the pedicel. The sepal was wetted in 0.01% v/v Silwet water and mounted between two coverslips filled with 0.01% v/v Silwet water. To prevent the sepal from drying out during imaging, the coverslips were sealed with vacuum grease. Sepals were imaged using 10X objective (C-Apochromat 10X/0.45 W M27). Three-horizontally tiled images were taken with an overlap of 5% and were auto-stitched with a threshold of 0.70 in the ZEISS ZEN software. For imaging fully mature sepals of pLH13 in WT and in *as2-7D*, mCitrine was excited with 514 nm laser (intensity- 20-22%) and emission was collected at 519-621 nm. For imaging PI stained sepals of WT, *as2-7D*, *pML1::KRP1* and *pML1::KRP1 as2-7D*, PI was excited at 514 nm (intensity- 27-30%) and emission was collected at 566-648 nm. For epidermal structure observation, mature flowers harboring pAR169 (*pATML1::RCI2A-mCtrine*) were transverse sectioned using a razor blade. Both mCitrine and chlorophyll were excited using 514 nm laser, and emission was collected at 520-565 nm for mCitrine and 651-700 nm for chlorophyll.

Osmotic treatments for measuring epidermal stiffness

Osmotic treatments were performed as described previously (*31, 48*), with modifications. In brief, primarily inflorescences from pLH13 in WT, pLH13 in *as2-7D*, pLH13 in *pML1::KRP1* and pLH13 in *pML1::KRP1 as2-7D* six to eight week old plants were dissected down to stage 3 to 4, and placed in ½ MS live imaging media plate. Dissected tissues were allowed to recover for 24 hours in the growth chamber. Early stage 6 buds were precisely chosen for osmotic treatment, since this is the stage right before *as2-7D* outer epidermis starts buckling.

Each bud (corresponding to one replicate) was subjected to osmotic treatment individually and was imaged pre- and post-treatment. Pre-treatment imaging involved staining the bud with a PI working solution for precisely 6 minutes. After washing off the PI stain, the tissue was transferred to a fresh media plate and was oriented to focus on the outer sepal. The tissue was submerged in 0.01% v/v Silwet water, and the pre-treatment image was captured. Imaging

1095 settings used for mCitrine – excitation 514 nm (intensity- 20-25%), emission 519-567 nm and PI
1096 - excitation 514 nm (intensity- 20-25%), emission 603-643 nm.
1097
For osmotic treatment, the bud was immersed in 0.35M NaCl solution containing 0.01% v/v
Silwet for exactly 20 minutes. Post treatment, the bud was re-stained in a PI working solution
prepared in 0.35M NaCl (1 part 5 mg/mL PI (solubilized in 0.35M NaCl) and 4 parts 0.35M
NaCl solution containing 0.01% v/v Silwet)) for 10 minutes, leading to a cumulative osmotic
treatment duration of 30 minutes. This timeframe was found to be optimum for inducing
plasmolysis in epidermal cells of early stage 6 flowers, wherein the plasma membrane gets
visibly detached from the cell wall at the corners. Re-staining the buds with PI is necessary since
NaCl solution erodes off the PI stain. The bud was finally re-positioned in a fresh media plate
and submerged in 0.35M NaCl solution. Before image acquisition, the tissue was allowed to
stabilize for a few minutes due to its increased fragility post osmotic treatment. The post-
treatment image acquisition setting was as follows: mCitrine – excitation 514 nm (intensity- 25-
35%), emission 519-567 nm and PI - excitation 514 nm (intensity- 25-35%), emission 603-643
nm. This procedure was iterated for all samples individually.
1111
1112 **Image processing**
1113
Converting raw images to tiffs, taking screenshots and videos
The raw images, generated either in lsm or czi formats from ZEN imaging software, were
converted to tiff files in ImageJ distribution Fiji. For multichannel images, images were split
based on channels, which were saved individually. The MorphoGraphX (MGX) (*76*) software
was used for image processing and taking the screenshots using the built-in screenshot option in
MGX. For taking screenshots of the reporter lines, the reporter image was loaded onto Stack 1,
and either chlorophyll or PI was loaded onto Stack 2. Colormaps for Stack 1 and Stack 2 were
set to green and gray respectively. Flower images and meristems were rotated in the desired
orientation. The voxels not associated with the flower or meristem of interest, were removed
using the Voxel Edit tool in MGX. In addition, trichomes from the outer sepal epidermis were
removed in a similar manner, for better visualization of the sepal surface. Following this,
screenshots were captured. From flower images, trichomes on the outer sepal were removed by
image processing for better visualization of the sepal epidermis. The brightness and contrast
were uniformly adjusted across samples for any given experiment. Screenshots of MGX
generated heatmaps were taken using the default brightness and contrast settings, without any
adjustments.
1130
Videos were captured as a series of two-dimensional tiffs files using the built-in "Record movie"
option in MGX. The 3D stack images were rotated in the MGX window for the desired duration
and time points. For capturing videos involving two genotypes/treatments, the images were
loaded onto Stack 1 and Stack 2 and rotated independently. The stack of tiffs generated was

loaded onto Fiji and saved as a .avi file without compression. The .avi files were assembled and
labeled in Adobe Premiere Pro 2023.
1137
1138   **Generating surfaces for analyzing outer epidermal growth**
1139
Making the surfaces and analysis of growth data was performed in MGX as previously described
in Zhu et. al. and Hong et. al. (*31, 89*), with modifications. The following process was used to
generate the surfaces for analyzing cell growth parameters on the outer epidermis of WT, *as2-*
7D, pML1::KRP1, pML1::KRP1 as2-7D, as2-7D mock and 10 µM NPA treatment samples.
Briefly, the tiffs were imported onto Stack 1 Main Store in the software. Trichomes on the outer
epidermis were removed using the voxel edit tool to get a better surface. Since the outer
epidermal morphology of *as2-7D* becomes increasingly irregular in later stages, Level Sets (*90*)
was used to extract the structure representing the general flower shape. For this, Level Sets was
initialized with an up/down threshold value of 3 to 4. The structure was allowed to evolve (Level
Set Evolve) with default parameters, except for "view steps", which was set to 1 for visualization
at more frequent intervals. For performing Edge Detect Angle, the mask was oriented such that
the outer sepal directly faces the viewer. The Edge Detect Angle process was then run with
default parameters to get a finer structure that closely resembles the image. The holes generated
in the structure (which corresponds to places in the image with low membrane marker signal)
were closed using Stack/Morphology/Closing - X/Y/Z radius of 3 to 8. A 2.5D surface fitted
with a triangular mesh was created from the structure using Marching Cube Surface with a cube
size of 3 to 5 µm, followed by smoothing once (Smooth Mesh, passes = 10) and subdividing
twice. The membrane signal was projected onto the 2.5D surface with a min. and max. distance
range of 1 to 8 µm from the image.
1159
1160   **Generating surfaces for analyzing outer vs inner epidermal growth in WT and *as2-7D***
1161
To obtain the inner epidermal surface, clipping plane 1 (View Tab/Clip 1), set to a width of 10 to
15 µm, was used to clip the flower image along the longitudinal direction. The voxel editing tool
was used to carefully remove the voxels below the inner epidermis of the sepal within the
clipping plane region. The clipping plane was then moved to the adjacent region and the same
process was repeated, while tilting the plane to ensure voxels from the inner epidermis itself are
not removed. Once the voxels below the inner epidermis were removed throughout, the edited
tiff was copied back to the main stack and saved. To generate the surface for the inner epidermis,
the Z axis of the stack was reversed (Stack/Canvas/Reverse Axes). The stack was blurred using
Gaussian blur (X/Y/Z sigma (µm) = 2 to 2.5), and structure was extracted using Edge Detect
with a threshold of 2000 to 3000. A triangular mesh surface was created using a cube size of 3
µm, followed by smoothing once (passes = 10) and subdividing twice. Plasma membrane signal
from the inner surface was projected onto the mesh surface with a min. and max. distance range
of 3 to 8 µm.

1175

1176 To obtain the outer epidermal surface, the processed tiff, prior to reversing the axes, was used.
1177 To process the stack, Gaussian blur (X/Y/Z sigma (µm) = 1 to 1.5) followed by Edge detect
1178 (threshold of 8000 to 10,000) were used sequentially. Mesh surface was created as described
1179 above, and the membrane signal was projected within a 2 to 4 µm distance range from the image.
1180

1181 **Generating surfaces for analyzing growth of outermost mesophyll layer in WT and *as2-7D***

1182

1183 To enhance the membrane marker signal for the mesophyll cells, we utilized a deep learning
1184 approach based on a Convolutional Neural Network (CNN). We used the inbuilt CNN Unet 3D
1185 prediction algorithm VijayanUnet.pt (*41*) in MorphoGraphX to precisely predict the cell outlines.
1186 The algorithm then uses these predictions for enhancing and de-noising the membrane marker
1187 signal and generates a new image based on the predictions.

1188

For both WT and *as2-7D* mutants harboring pLH13 membrane marker, the CNN algorithm was
run on the pre-processed confocal images of the outer sepal, where the parts of the flower not
associated with the outer sepal were already removed by voxel editing (see 'Generating surfaces
for analyzing outer vs inner epidermal growth in WT and *as2-7D*' section in Materials and
Methods). Using the pre-processed sepal images instead of whole flowers for running the CNN
algorithm largely reduced computational processing times. Briefly, for WT and *as2-7D* mutants,
CNN-generated images of the sepals for each time point were obtained by running
Stack/CNN/UNet3D Prediction/VijayanUnet.pt using default parameters. The CNN Unet3D
prediction process usually also generates voxels unrelated to the sepals, which were removed
using the 'Voxel edit' tool. These edited CNN-generated images were then used to run
Stack/Mesh Interaction/Annihilate function, to remove the voxels associated with the outer
epidermis and the inner layers and obtain the outermost mesophyll layer (layer closest to the
outer epidermis). For sepals at 0-hour and 24-hour time points, the outermost mesophyll layer
was extracted by setting the minimum distance values for Annihilate function to 5/6 and the
maximum distance values to 15. For the 48-hour time points, the minimum and maximum
distance value for the Annihilate function was set to 6 and 20 respectively. The resulting images
generated post running Annihilate function were used to generate the 2.5D surface for the
outermost mesophyll layer. Briefly, for each time-point and genotype, the annihilated image
stack was loaded onto Stack 1 and blurred using Gaussian Blur, with X/Y/Z sigma (µm) value of
3. The surface was extracted using Edge Detect threshold of 2000 to 3000. The mesh surface was
created using a cube size of 3 µm, followed by smoothing once (passes = 10) and subdividing
twice. The membrane signal was projected onto the mesh surface with a minimum and maximum
distance range of 7 to 8 µm.

1212

1213

1214 **Lineage tracking**

1215
For meshes across all time points, individual cells were manually seeded. Cells were segmented
using the watershed algorithm, and unlabeled vertices were fixed by running Cell Mesh/Fix
Corner Triangles. Cell lineages were parent tracked to the earlier time point to analyze cell
growth. Parent (lineage) tracking involves transferring the parent labels from cells in the earlier
time point onto the daughter cells at the later time point. For this, the meshes corresponding to
the earlier (Mesh 1) and later (Mesh 2) time points were opened side by side on the same MGX
window. A few daughter cells in Mesh 2 were manually parent labeled, and using these as
landmarks, the remaining daughter cells were auto parent labeled (by running
Deformation/Morphing/Set Correct Parents followed by Deformation/Mesh 2D/Auto Parent
Labelling 2D). Errors in segmentation or parent tracking were manually corrected. Parent
tracking was checked for accuracy using Cell Axis/PDG/Check Correspondence, which checks
whether the cell junctions are identical across time points. The final parent labels were saved as
.csv files. A custom R script (combinelineages.r) was run to automatically correlate the lineage
information through multiple time points, followed by manual correction.
1230
1231 **Analyzing cell growth parameters**
1232
Meshes corresponding to the earlier (Mesh 1) and later time points (Mesh 2) were loaded side by
side in the same MGX window, with parent labels loaded onto Mesh 2. Cell growth rate was
calculated based on the total area of the daughter cells in the second time point divided by the
area of the parent cell in the earlier time point. Growth rate heatmaps were generated using Heat
Map/Heat Map Classic/Heat Map with the Type set to Area. The cell division rate was calculated
based on the total number of daughters formed in the second time point for each cell in the
earlier time point. Cell division heatmaps were generated using Lineage Tracking/Heat Map
Proliferation. The values for growth rate and cell division rate were exported to .csv files.
1241
The Principal Direction of Growth (PDG) uses information of corresponding junctions between
the earlier and later time points, to compute the direction of maximum and minimum growth of
each individual cell in the earlier time point (*76*). In the figures, the PDGs represent the
maximum growth direction (PDG$_{max}$) tensor (yellow lines), the length of which is directly
proportional to the magnitude along the direction of maximum growth. The PDGs were
calculated for the cells in the earlier time point (using the information of parent labels from the
later time point), the PDG$_{max}$ tensors were projected onto the later time point for visualization.
For displaying PDG$_{max}$, the values for line width and line scale were set to 5 and 3 respectively.
1250
The Angle of Deviation of the PDG$_{max}$ tensor was calculated relative to the longitudinal axis of
the sepal. The sepal longitudinal axis was set using a Bezier line on the earlier time point, since
PDGs and deviation angles are calculated on the earlier time point. Briefly, a new Bezier grid
(Misc/Measure/New Bezier) was created, which was then collapsed into a Bezier line

(Misc/Bezier/Collapse Bezier Points). The supporting points on the Bezier line were placed on
the mesh in a manner such that the Bezier line is parallel to the sepal longitudinal direction, and
the line shape fits the curvature of the sepal. The custom axis for each cell was set using this
Bezier line (Cell Axis/Custom/Create Bezier Line Directions) followed by smoothing (Smooth
Custom Directions). The PDGs were loaded onto the mesh, and the angle of deviation of the
PDG_{max} from custom X direction (which represents the longitudinal axis) was calculated using
Cell Axis/Custom/Custom Direction Angle. The values of the angle of deviation heatmap were
exported as .csv files.
1263
1264   **Analyzing osmotic treatment induced epidermal shrinkage**
1265
Our osmotic treatment conditions were optimized such that the cell walls shrank, but the cell
membrane barely plasmolyzed. Post osmotic treatment, the membrane just visibly pulled away
from the cell wall corners, thereby rendering the wall and membrane area almost comparable.
Since the PI signal (corresponding to the cell wall) is very weak post-treatment, especially for the
inner layers, the PI and mCitrine (membrane marker) images were merged as a single image
MGX (Stack/Multistack/Merge Stacks).
1272
For analyzing epidermal stiffness, we assumed that the surface tension across genotypes is
comparable. The shrinkage in area of the epidermal layers along a comparable longitudinal
section was analyzed to infer tissue stiffness. For this, the pre- and post-treatment merged stacks
for any given sample were opened side by side in the same MGX window. Using clipping plane
1 (Clip 1), longitudinal cross-section corresponding to the same regions were extracted from the
flower images. The tiffs were processed using Levels Sets to extract the global shape. Mesh
surfaces were made with a cube size of 3 µm, which was smoothed once (10 passes) and
subdivided twice. Signal was projected with a 1.5 to 3 µm distance range from the image. From
the cross-section surface, the outer epidermis and inner epidermis were seeded and segmented as
single "cells". Care was taken to ensure that the epidermal region being seeded was comparable
between the pre- and post-treatment surfaces. For measuring shrinkage, parent labels from the
pre-treatment mesh were transferred onto the post-treatment mesh, following which the
shrinkage ratios were quantified (Mesh/Heatmap/Heatmap Classic). The areal shrinkage values
were stored as a .csv file, which was converted to percentages for generating heatmaps shown in
the figures.
1288
1289   **Computing minimum curvature**
1290
In MorphographX, minimal curvature analysis quantifies the least degree of curvature at any
given location on the surface. The output is displayed in the form of a heatmap where warm
colors represent regions of higher curvature or outward bulging, whereas cool colors represent
regions of lower curvature, such as inward deformations (saddles). For calculating the minimal

curvature for the sepals from the live imaging series of *pPIN1::PIN1-GFP as2-7D* flowers, we
used the images obtained from the chlorophyll autofluorescence channel to construct the mesh
surfaces. Briefly, the chlorophyll image corresponding to a given flower (for which the minimal
curvature is to be analyzed) was loaded onto Stack 1. The structure representing the general
shape of the flower was extracted using Level Sets (Stack/Lyon/Init Level Set), with an up and
down threshold values set to 4. The structure was allowed to evolve (Stack/Lyon/Level Set
Evolve) with default parameters. The triangular 2.5D mesh was created using
Mesh/Creation/Marching Cube Surface with 5 µm cube size, followed by one round of
smoothing (Mesh/Structure/Smooth Mesh, passes = 5) and subdividing
(Mesh/Structure/subdivide). Minimal curvature was then calculated using Mesh/Signal/Project
Mesh Curvature, selecting Type = Minimal and setting the neighborhood radius value to 50 µm.
For visualizing the resulting heatmap, the transfer function was set to "Jet".

1307
1308

1309 **Analytical model for a growing trilayer**

1310

System and assumptions. We consider a growing elastic system made of three layers,
corresponding to the inner epidermis (denoted i), mesophyll (m), and outer epidermis (o) of a
lateral organ. We consider that the three layers may grow at different (specified or target) rates
and have different elastic moduli. To simplify the formulae, we assume that the epidermal layers
have similar thicknesses. We also assume that the Poisson ratios of all layers are equal (value v).
Finally, we neglect the bending modulus of each layer considering their small thickness. Note,
these three assumptions do not affect the qualitative behavior of the system.

We assumed E_i, E_m, and E_o be the values of elastic modulus, h, h_m, and h the values of
thickness,and g_i, g_m, and g_o as the values of strain associated with target (specified) growth rates
for the inner epidermis, the mesophyll, and the outer epidermis, respectively.

1321

The differences in growth rates between the layers may lead to changes in system geometry. As
the three layers are bound to each other, they have approximately the same curvature c
(corresponding to the radius of curvature measured at the middle of the mesophyll), while the
strain in the inner and outer layers can be deduced from the strain in the middle layer ε using
curvatures and assuming that the reference state is flat. From kinematics of the system, the
elastic (residual) strains in the three layers are given by

1328
$$\varepsilon_i = \varepsilon - \frac{1}{2}(h_m + h)c - g_i$$

1329
$$\varepsilon_m = \varepsilon - g_m$$

1330
$$\varepsilon_o = \varepsilon + \frac{1}{2}(h_m + h)c - g_o$$

1331 The elastic energy of the system takes the form

$$1332 \quad E = \frac{1}{2(1-v^2)}[E_i h \varepsilon_i^2 + E_m h_m \varepsilon_m^2 + E_o h \varepsilon_o^2]$$

1333

1334   Minimizing the elastic energy with respect to strain and curvature yields their values at
1335   equilibrium. In particular, using the notation

$$1336 \quad A = \frac{1}{(4E_i E_o h + E_m(E_i + E_o)h_m)} > 0$$

1337   we obtain the equilibrium values for the strain

$$1338 \quad \varepsilon = A(2E_i E_o h(g_i + g_o) + E_m(E_i + E_o)h_m g_m)$$

1339   and the curvature

$$1340 \quad c = \frac{2A}{h+h_m}\left(2E_i E_o h(g_o - g_i) + E_m E_o h_m(g_o - g_m) - E_m E_i h_m(g_i - g_m)\right).$$

1341   This curvature is in general non-zero, the system curves with the abaxial side on the convex side
1342   as soon as the abaxial growth rate is sufficiently larger than the adaxial growth rate.

1343   To determine whether buckling of the abaxial epidermis may occur, we compute the elastic
1344   (residual) strain in this layer at equilibrium:

$$1345 \quad \varepsilon_o = AE_m E_i h_m(2g_m - g_i - g_o).$$

1346 ε_o can become compressive (<0) only if $2g_m - g_i - g_o < 0$. When it falls beyond a threshold
1347   (91), this leads to buckling of the abaxial side, with patterns as described (92), depending on the
1348   anisotropy of the residual strain.

1349   Interestingly, the elastic strain of the adaxial epidermis,

$$1350 \quad \varepsilon_i = AE_m E_o h_m(2g_m - g_i - g_o)$$

always has the same sign as that of the abaxial epidermis. This means that either the two
epidermal layers are in compression, or the two epidermal layers are in tension. A situation
where one epidermis would be in tension and the other would be in compression does not
correspond to mechanical equilibrium. Nevertheless, the amount of compression may differ
between the two epidermal layers: the ratio of the elastic deformations is related to the ratio of
Young's moduli

$$1357 \quad \frac{\varepsilon_o}{\varepsilon_i} = \frac{E_i}{E_o}.$$

1358   Accordingly, it is possible that only one of the two epidermal layers reaches the buckling
1359   threshold. For instance, a much smaller Young's modulus for the outer epidermis compared to
1360   the inner epidermis would lead to buckling of the outer epidermis exclusively.

1361

1362   Finally, we mention that the elastic strain in the mesophyll has an opposite sign

$$1363 \quad \varepsilon_m = -2AE_i E_o h(2g_m - g_i - g_o).$$

1364   This means that as soon as the epidermis is under compression, the mesophyll is under tension
1365   and vice versa.

1366

1367   **Numerical model for a growing layered sepal**

1368

We again considered a growing elastic system comprising of three layers (inner epidermis (i),
mesophyll (m), and outer epidermis (o)). To better represent the geometry of the transversal
section of a sepal, we also considered two lateral semi-circles (the edges) to connect the inner
and outer epidermis (Fig. 2, I to K). These semi-circles are denoted "end", while the region
enclosed by the epidermal layers and the edges denote the mesophyll of the trilayer. This
geometry is characterized by three parameters: the epidermal layer thickness h, the mesophyll
thickness h_m and the width of the trilayer part L.

We adopt the framework of morpho-elasticity (*93*), assuming that the residual stresses are solely
created by a local growth deformation tensor F_g, and the observed deformation gradient can be
decomposed multiplicatively on the elastic and the growth deformation tensors. In this
framework we consider a quadratic energy density in the Green-Lagrange strain tensor E_e,

1380
$$W = \tfrac{1}{2} E_e : H : E_e,$$

1381   where H, the elasticity tensor of rank four, represents the mechanical properties of the material.
1382   The proposed model assumes all regions of the geometry to be isotropic materials. Therefore, the
1383   elasticity tensor field H, is characterized by independent parameters: the Young's moduli of the
1384   four regions (E_i, E_m, E_o and E_{end}), and the common Poisson ratio v. An equation for the time
1385   evolution of the growth tensor F_g completes the model,

1386
$$\begin{aligned} \frac{\partial F_g}{\partial t} &= G F_g, & t &> 0 \\ F_g &= 1, & t &= 0 \end{aligned},$$

1387   with G the growth rate tensor. We consider here that the growth rate has both an imposed target
1388   (or specified) component G_T, as well as a component that stands for an influence of the elastic
1389   strain on growth,

1390
$$G = G_T + C E_e,$$

with C being the extensibility (assumed to be scalar). We assume that the target growth is
unidirectional along the trilayer, so that the target growth rates γ_i, γ_m, γ_o and γ_{end} for the inner
epidermis, the mesophyll, the outer epidermis, and the edge respectively are the only
independent parameters encoded in the target growth rate tensor G_T. As for the extensibility C, we
have four additional parameters C_i, C_m, C_o and C_{end} corresponding to the four regions.

In comparison with the analytical model presented above, all geometrical and mechanical
parameters are the same, while the growth parameters g_i, g_m, and g_o can be seen as the relative
growth cumulated until time t, which can be approximated as when $C = 0$.

The model of the growing layered sepal was solved using the Finite Element Method and
implemented with the FEniCS computing platform (*94*) with its Python interface, with a special
use of the BVPy Python library (*95*).

1402

We simulated the deformation of the defined geometry when the outer epidermis is growing (
$\gamma_o \neq 0$), while the other regions do not ($\gamma_i = \gamma_m = \gamma_{end} = 0$). This is one of the simplest situations,
when the mean growth rate of the two epidermal layers is higher than the growth rate of the
mesophyll, and according to the predictions of the analytical model the epidermal layers are
under compression. This first condition for buckling being satisfied, we studied whether buckling
indeed occurs depending on the moduli of the mesophyll and epidermal layers (fig. S3). To
validate the approach, we also compared the equilibrium curvature predicted by the analytical
model to the mean midline curvature of the deformed sepal in the numerical model (see fig. S4).

1411

For all our simulations we considered fixed values for the length $L = 10$, and thickness $2h + h_m =$
5 of the sepal, the target growth rate of the outer epidermis $\gamma_o = 0.1$, and the Poisson ratio $\nu =$
0.49. The value of the extensibility is chosen low for all regions but relatively higher for the
mesophyll region, $C_o = C_i = C_{end} = 0.01$ and $C_m = 0.1$. The Young's modulus of the mesophyll is
chosen as the lowest among all regions, $E_m = 1$, while the Young's modulus of the edge
epidermis is considered to be a high value, $E_{end} = 10^6$. As for the Young moduli of the outer and
inner epidermis, we performed an exploration of the parameter space defined by these two
extreme values, $(E_o, E_i) \in [1, 10^6] \times [1, 10^6]$. We present the results of this analysis for different
values of the epidermis thickness, $h \in \{0.1, 0.2, 0.3, 0.4\}$ (fig. S3 and S4).

1421

1422

1423 **Graph plotting**

1424

1425 Graphs were generated in R (version 4.2.2 (2022-10-31 ucrt)) using ggplot, raw data for which is
1426 provided in Data S1.

 SUPPLEMENTARY FIGURES AND LEGENDS

1428

Fig. S1. *AS2* overexpression causes characteristic ridges, invaginations, and outgrowths on
the outer epidermis, associated with Main Fig. 1.
(**A** and **B**) Representative optical cross-sections of Stage 11 flowers of (**A**) the WT and (**B**) *as2-*
7D mutant. Images are representative of T2s harboring pAR169 (*pATML1::RCI2A-mCitrine*), an
epidermal layer specific plasma membrane marker. Green and grey represent epidermal cells and
chlorophyll containing mesophyll cells respectively. WT sepals (A) have a smooth shape, while
the *as2-7D* sepal (B) outer epidermis exhibit outward protrusions (ridges), inward indents
(invaginations) and pointed epidermal extensions (outgrowths). Note that the outgrowths lack
mesophyll cells (grey) underneath whereas mesophylls occupy the space beneath ridges and
invaginations.
(**C** and **D**) Whole inflorescence images of (**C**) Col-0 (control) and (**D**) *as2-5D* (in Col-0, isolated
by Wu et. al (*35*)). Images are representative of *n* = 3 inflorescences. Scalebar, 1 cm.
(**E** and **F**) Stage 14 flowers of (**E**) Col-0 and (**F**) *as2-5D* mutant, representative of *n* = 3 flowers.
Yellow arrowhead points toward an outgrowth. Note the mild nature of outgrowth formation in
as2-5D mutant as compared to *as2-7D* (Fig. 1). Scalebar, 1 cm.
(**G** to **J**) Representative six-week-old adult plants of (**G**) WT, (**H**) *as2-7D* and (**I-J**) *pML1::AS2*
in WT T1s. The transgenics show strong (17/28 plants, panel I) to extreme (9/28 plants, panel J)
developmental defects, categorized qualitatively by the extent of leaf deformity. The transgenics
exhibit outgrowths on the lower (abaxial) surface of the leaves as well, in addition to severely
stunted growth. Scalebar, 1 cm.
(**K** to **N**) Fully formed siliques of (**K**) WT, (**L**) *as2-7D*, (**M-N**) *pML1::AS2* in WT T1s. Note the
outgrowths on the silique epidermis. In panels M and N, siliques are representative of strong and
extreme categories, described for panels I and J respectively. Three siliques on each panel
correspond to three independent plants. Total number of independent transgenics (T1s) assessed
n = 28.

1454
1455 **Fig. S2. The outer epidermis of *as2-7D* exhibits loss of stereotypical growth patterns,**
1456 **associated with Main Fig. 2.**

(**A** to **D**) Confocal stack images of (**A**) WT replicate 2, (**B**) WT replicate 3, (**C**) *as2-7D* replicate 2 and (**D**) *as2-7D* replicate 3 developing flowers, transformed with the pLH13 (*35S::mCitrine-RCI2A*) membrane marker. Images correspond to the 48-hour (left) and 96-hour (right) time points from WT and *as2-7D* live imaging experiment (*n* = 3). Yellow arrowheads show outgrowths. Trichomes were removed from the outer sepal epidermis by image processing. Scalebar, 50 μm.

(**E** to **H**) Cell area growth rates of (**E**) WT replicate 2, (**F**) WT replicate 3, (**G**) *as2-7D* replicate 2 and (**H**) *as2-7D* replicate 3. The heatmap is projected onto the later time point. Red to blue gradient shows fast to slow growth. Yellow arrowheads point towards the outgrowth which exhibit localized fast growth. Scalebar, 50 μm.

(**I** to **L**) The cell division rates of (**I**) WT replicate 2, (**J**) WT replicate 3, (**K**) *as2-7D* replicate 2 and (**L**) *as2-7D* replicate 3. Red to blue gradient shows more to less division. Cell division over 0-48 hours is shown on the left and over 48-96 hours on the right. Scalebar, 50 μm.

(**M** and **N**) Graphical representation of (**M**) cell growth rates and (**N**) cell division rates across all three independent biological replicates of the WT (black dots) and *as2-7D* mutant (red dots). Dot represents the growth rate (M) and division rate (N) of a given cell lineage. Dot plots are superimposed with box plots, where the central line indicates the median. Left and right panels display data for 0-48 and 48-96 hours, respectively. Note the reduced spatial variability in the growth rates of *as2-7D* (panel M). Lines above data points indicate statistical comparisons. The *p-values* for Bonferroni corrected pairwise t-test (two-tailed) are shown. *p-values*: ***<0.0001, **<0.001, *<0.05, ns = non-significant. Exact *p-values* for growth rates: WT vs *as2-7D* for 0-48 hours, 0.9884; WT vs *as2-7D* for 48-96 hours, 1.565172×10^{-93}. Exact *p-values* for cell division rates: WT vs *as2-7D* for 0-48 hours, 7.800489×10^{-05}; WT vs *as2-7D* for 48 - 96 hours, 9.47882×10^{-07}.

1481
Fig. S3. Numerical simulations for a sepal cross-section with the outer epidermis growing
in width, showing that buckling is determined by the relative Young's moduli of the outer
and inner epidermis, associated with Main Fig. 2.

Parameter space exploration for the numerical model of the growing sepal with three layers,
corresponding to the outer epidermis (blue), mesophyll (yellow) and the inner epidermis (orange)
as shown in Fig. 2, I to K. As a proxy for detecting whether outer epidermis buckles or not, we
monitor the number of curvature maxima on the surface of the outer epidermis. A value for
maxima higher than one indicates the occurrence of a buckling event. Block rows in the grid
correspond to different epidermal thickness values $h \in \{0.1, 0.2, 0.3, 0.4\}$, while block columns
correspond to different time points during growth. In each block the Young's moduli of the outer
and inner epidermis are variable, $(E_o, E_i) \in [1, 10^6] \times [1, 10^6]$. All other parameters are kept
constant: $L = 10$, $2h + h_m = 5$, $\gamma_o = 0.1$, $\gamma_i = \gamma_m = \gamma_{end} = 0$, $\nu = 0.49$, $C_o = C_i = C_{end} =$
0.01, $C_m = 0.1$, $E_m = 1$, $E_{end} = 10^6$. Holes in the grid are simulations for which the numerical
solver diverged for the given parameter set and time point.

1496

1497

Fig. S4. Cross validation of the analytical and numerical models for the growing layered
sepal, associated with Main Fig. 2.
(**A**) Equilibrium curvature value predicted by the analytical tri-layer model. (**B**) Mean midline
curvature of the deformed sepal in the numerical model. (**A** and **B**) Parameter values and the
organization of the grids are same as described in fig. S3. The curvature values in the numerical
model deviate from the analytical model prediction primarily when the inner epidermis is much
softer than the outer epidermis and the edges. We attribute this deviation to the difference in
geometry between the two models, namely the presence of a rigid edge in the numerical model,
that connects the inner and outer epidermal layers.

1507
Fig. S5. Slower growth on the inner epidermis together with misdirected outer epidermal
growth trigger buckling in *as2-7D*, associated with Main Fig. 3.
(**A** to **D**) Growth rate heatmaps for (**A**) WT replicate 2, (**B**) WT replicate 3, (**C**) *as2-7D* replicate
2 and (**D**) *as2-7D* replicate 3 developing sepals. Red to blue heatmap gradient shows fast to slow

growth. Outer (left) and inner (right) epidermal growth rates over 48 hours is projected as a
heatmap onto the later time point. Scalebar, 20 µm.
(**E**) Graph shows cell growth rates over a 48-hour period for the WT (black) and *as2-7D* (red)
mutant (*n* = 3 biological replicates). Each dot represents the growth rate for a given cell lineage.
Dot plots are overlayed with boxplots, where the central line represents the median. The *p-values*
for Bonferroni corrected pairwise t-test (two-tailed) are shown. *p-values*: ***<0.0001, **<0.001,
*<0.05, ns = non-significant. Exact *p*-values for epidermal cell growth rate: WT outer vs inner,
5.751359×10^{-35}; *as2-7D* outer vs inner, 3.039502×10^{-25}; WT vs *as2-7D* outer, 0.1011; WT vs
as2-7D inner, 8.496122×10^{-16}.
(**F** to **I**) Cell division heatmaps for (**F**) WT replicate 2, (**G**) WT replicate 3, (**H**) *as2-7D* replicate
2 and (**I**) *as2-7D* replicate 3. Red to blue color gradient shows more to less cell division. The
data is presented in the same format as described for panels A-D. Scalebar, 20 µm.
(**J**) Graph shows cell division rates for the WT and *as2-7D* mutant (*n* = 3 biological replicates)
Dots represent the number of cell divisions for each cell lineage. Data presentation and statistical
comparisons are same as described above for panel E. Exact *p*-values for epidermal cell division
rates: WT outer vs inner, 4.08502×10^{-28}; *as2-7D* outer vs inner, 0.004; WT vs *as2-7D* outer,
3.677026×10^{-23}; WT vs *as2-7D* inner, 1.865456×10^{-06}.
(**K** to **N**) Heatmap representing the angle of deviation of PDG_{max} tensors from the sepal
longitudinal axis for (**K**) WT replicate 2, (**L**) WT replicate 3, (**M**) *as2-7D* replicate 2 and (**N**)
as2-7D replicate 3. Scalebar represents a deviation of 90º in red and 0º (no deviation) in blue.
Scalebar, 20 µm.
(**O**) Graph showing the angle of deviation for the WT and *as2-7D* mutant (*n* = 3 biological
replicates). Each dot represents the angle of deviation for a given cell lineage. Data
representation and statistical comparisons are shown in the same format as described for panels E
and J. Exact *p-values* for epidermal angle of deviation: WT outer vs inner, 3.793354×10^{-05}; *as2-*
7D outer vs inner, 1.572509×10^{-20}; WT vs *as2-7D* outer, 8.052118×10^{-10}; WT vs *as2-7D*
inner, 0.349.
1539
1540

Fig. S6. Growth patterns of the mesophyll cells mimic that of the outer epidermis for both WT and *as2-7D*, associated with Main Fig. 3.

(**A** to **B**) Processed confocal images of (**A**) the WT and (**B**) *as2-7D* mutant stage 7 sepals, harboring the pLH13 membrane marker, at the 48-hour time point (representative of $n = 3$). Scalebars, 20 µm.

(**C** to **D**) View of the longitudinal cross section through the center of the stage 7 sepal blade for (**C**) the WT and (**D**) *as2-7D* mutant (representative of $n = 3$). Scalebars, 20 µm.

(**E** to **F**) Longitudinal sections of stage 7 sepal images of (**E**) the WT and (**F**) *as2-7D* mutant processed with CNN (Unet display/Vijayan.Unet algorithm, see Materials and Methods), to enhance the cell outlines (representative of $n = 3$). The gray areas mark the mesophyll layer closest to the outer epidermis (outer mesophyll), generated after running (Mesh Interaction/Annihilate) process in MorphographX to remove the outer epidermis. The annihilated image (gray) was overlayed on the CNN generated image (green) for comparison. The outer and inner epidermal layers are shown (yellow arrows). Scalebars, 20 µm.

(**G** to **H**) Plasma membrane signal from the outer mesophyll layer of (**G**) to WT and (**H**) the *as2-7D* mutant projected onto the mesh surface generated using MorphographX. Scalebars, 20 µm.

(**I** to **J**) Growth rates of the mesophyll cells over a 48-hour period for (**I**) the WT and (**J**) *as2-7D* mutant are shown (representative of $n = 3$). Red to blue heatmap gradient shows fast to slow growth. Scalebars, 20 µm.

(**K** to **L**) Principal directions of maximal growth (PDGmax, yellow) of the mesophyll cells of (**K**) WT and (**L**) *as2-7D* mutant are shown. Segmented cells in grey are superimposed on the plasma membrane signal (also in grey). Note that both the growth rates (panels I to J) and growth directions of the mesophyll cells are similar to those of the outer epidermis, as shown in Main Figure 3. Scalebars, 20 µm.

(**M**) Graph shows the cell growth rates of the mesophylls cells over a 48-hour period for the WT (black) and *as2-7D* (red) mutant ($n = 3$ biological replicates). Each dot represents the growth rate for a given cell lineage. The dot plots are overlayed with box plots, where the central line represents the median. The p-value from Tukey's test (two-tailed) is shown. *p-values*: ns = non-significant ($p > 0.05$).

(**N**) Graph shows the average cell growth rates of the outer epidermis (Outer), inner epidermis (inner), average of the outer and inner epidermis and the mesophyll cells of the WT (black) and *as2-7D* (red) mutant. Each dot represents the average growth rate across all cell lineages in each replicate ($n = 3$ biological replicates). The lines represent the mean growth rate across three biological replicates. The p-value from Tukey's test (two-tailed) is shown. *p-values*: *<0.05, ns = non-significant. Exact p-values for WT average (Outer & Inner) vs Mesophyll, 0.0036; WT average (Outer & Inner) vs Mesophyll, 0.0036.

Inner epidermis

WT   *as2-7D*   *AtML1::KRP1*   *AtML1::KRP1 as2-7D*

Floral remains
post dissection

WT Rep 2   *as2-7D* Rep 2   WT Rep 3   *as2-7D* Rep 3

AtML1::KRP1 Rep 2   *AtML1::KRP1 as2-7D* Rep 2   *AtML1::KRP1* Rep 3   *AtML1::KRP1 as2-7D* Rep 3

WT flower   Longitudinal cross-section meshes

Before treatment   After treatment   Before treatment   After treatment

1579

Fig. S7. Analysis of cell growth direction and stiffness due to *KRP1* overexpression in WT and *as2-7D*, associated with Main Fig. 5.

(**A** to **D**) Confocal images of the inner epidermis of (**A**) WT, (**B**) *as2-7D* mutant, (**C**) *AtML1::KRP1* and (**D**) *AtML1::KRP1 as2-7D* stage 14 sepals. The sepals were stained with PI. Note the smooth morphology of the inner epidermis across all genotypes. Scalebar, 200 μm, images representative of $n = 3$ adult sepals.

(**E** to **L**) Principal direction of growth (PDG$_{max}$) for the outer epidermal cells of (**E**) WT Replicate 2, (**F**) *as2-7D* mutant Replicate 2, (**G**) *AtML1::KRP1* Replicate 2, (**H**) *AtML1::KRP1 as2-7D* Replicate 2, (**I**) WT Replicate 3, (**J**) *as2-7D* mutant Replicate 3, (**K**) *AtML1::KRP1* Replicate 3 and (**L**) *AtML1::KRP1 as2-7D* Replicate 3 sepals. PDG$_{max}$ tensors (yellow) calculated over a 48-hour live imaging period starting from stage 4, projected on a later time point surface. Segmented cells in grey are superimposed onto the plasma membrane signal (also in grey). Scalebar, 20 μm, $n = 3$.

(**M** and **N**) Representative confocal images of WT ($n = 5$) early stage 6 flower (**M**) before and (**N**) after osmotic treatment in 0.35M NaCl. The PI (cell wall stain) and membrane marker (pLH13, *35S::mCitrine-RCI2A*) signals have been merged together and shown. The highlighted region of the outer sepal (white rectangle) indicates the comparable longitudinal section extracted for analyzing epidermal shrinkage.

(**O** and **P**) MorphoGraphX generated surface of the longitudinal section (**O**) before treatment and (**P**) after osmotic treatment, extracted from flower images shown in panels M and N respectively. Comparable regions of the outer and inner sepal epidermis were segmented (outlined in cyan) to calculate the percentage areal shrinkage.

1602

Fig. S8. Ectopic *AS2* expression mediated buckles and subsequent outgrowth formation

occur independent of *REVOLUTA*, associated with Main Fig. 6.

(**A**) Representation of the previously characterized meristem initiation defect phenotype of *rev-6*
mutant. Images show an excised lateral inflorescence meristem of the WT (left) and *rev-6* mutant
(right). Majority of the cauline leaves axils (white arrows) in the *rev-6* mutant are bare, as they
fail to produce axillary meristem, as described by Otsuga et. al. (*55*). Scalebar, 1cm.
(**B** to **D**) Fully developed siliques of (**B**) *rev-6* and (**C** and **D**) *pML1::AS2* in rev-6 T1s. No of
independent T1s assessed *n* = 19. The three siliques correspond to three independent plants. The
numbers represent the total number of plants showing strong (12/19 plants, panel C) to extreme
(7/19 plants, panel D) phenotypes, categorized based on the extent of leaf deformity in the adult
vegetative stage.
(**E** to **I**) Representative six-week-old adult plants of (**E**) WT, (**F**) *rev-6*, (**G**) *as2-7D* and (**H** and
I) *pML1::AS2* in rev-6 T1s. Total number of T1s assessed *n* = 19. The transgenics show strong
(12/19 plants, panel H) to extreme (7/19 plants, panel I) defects in leaf development,
characterized by outgrowth formation on the abaxial epidermis and stunted growth. Scalebar, 1
cm.
(**J** to **M**) Confocal images of PI-stained (greyscale) inflorescences of (**J**) WT, (**K**) representative
pML1::REV in WT T1, (**L**) *as2-7D* mutant, and (**M**) representative *pML1::REV* in *as2-7D* T1.
Note that the transgenics ectopically expressing *REV* looks like their respective backgrounds in
either the WT (panels J and K) or *as2-7D* (panels L and M). Yellow arrowheads point to the
outgrowths. For each *pML1::REV* transgenics in WT and in *as2-7D* backgrounds, number of
independent T1s assessed, *n* = 28. Scalebar, 50 μm.

1637
Fig. S9. KANADI loss-of-function mutant (*kan1-2 kan2-1*) exhibits floral development
defects which are distinct from *as2-7D*, associated with Main Fig. 6.
(**A**) Whole inflorescence images of *kan1-2 kan 2-1*. The image is representative of *n* = 3
inflorescences. Scalebar, 1 mm.
(**B**) Stage 14 flower of *kan1-2 kan 2-1*. The image is representative of *n* = 3 inflorescences.
Scalebar, 1 mm.

(**C**) Confocal image of PI-stained (greyscale) *kan1-2 kan 2-1* mature sepal showing the outer

(abaxial) epidermis. The image is representative of $n = 3$ sepals. Scalebar, 100 µm. Note the

twisted and curled morphology of the sepal.

(**D**) Inset of *kan1-2 kan 2-1* mature sepal outlined in panel (C). Note the folds on the abaxial

epidermis of *kan1-2 kan 2-1* sepal, which are less severe as compared to *as2-7D*. Scalebar, 50

µm.

(**E**) Confocal image of PI-stained (greyscale) *kan1-2 kan 2-1* inflorescence. The image is

representative of $n = 3$ inflorescences.

(**F** to **L**) Confocal images of PI-stained (greyscale) *kan1-2 kan 2-1* flowers of different

developmental stages (staged based on floral organ development in accordance with Smyth et al.,

1990 (*74*). The sepals are shown by white arrows. All the floral organs are morphologically

deformed. Note that the sepals fail to enclose the floral meristem from initial stage onwards and

exhibit a narrow blade development in the later stages. Parts of the inflorescence tissue not

associated with the flowers were removed by image processing. Scalebar, 20 µm.

(**M**) Representative five-week-old adult *kan1-2 kan 2-1* plant. The image is representative of $n =$

3 plants. Scalebar, 1 cm.

(**N**) Representative rosette leaf of *kan1-2 kan 2-1*. Yellow arrowheads show the outgrowths. The

image is representative of $n = 3$ leaves. Scalebar, 1 cm.

(**O**) Confocal image showing a region of the abaxial epidermis of kan*1-2 kan 2-1* leaf (PI-

stained, grayscale). Yellow arrows show the outgrowth, while the white arrows denote the folds

on the leaf. Note that the outgrowth forms in the region where the folds form. Scalebar, 200 µm.

1665

1667   **Fig. S10. Outgrowths on *as2-7D* outer epidermis are formed due to buckling mediated**
1668   **PIN1 convergence and subsequent auxin maxima formation, associated with Main Fig. 7.**
1669

(**A** and **B**) Confocal images show the expression pattern of PIN1 protein reporter (*pPIN1::PIN1-*
GFP) in **(A)** WT and **(B)** *as2-7D* inflorescence meristems. Note the expression of *PIN1* directed
towards the emerging outgrowths on the outer epidermis of *as2-7D* older flowers. Scalebar, 50
μm, $n = 3$.
(**C** and **D**) Expression pattern of auxin response reporter (*pDR5::VENUS*) in stage 11 flower of
as2-7D mutant. Shown is (**C**) VENUS signal by itself (green) and (**D**) VENUS signal merged
with PI (grey). Note the presence of DR5 signal specifically in the outgrowths (yellow
arrowheads). Scalebar, 100 μm, $n = 3$.
(**E** to **H**) Representative live-imaging series showing the second replicate of (**E**) Mock treated,
(**F**) 10μM NPA treated developing flowers, and third replicate of (**G**) Mock treated, (**H**) 10μM
NPA treated developing flowers of *as2-7D* mutant. Images taken at every 24-hour interval over
96-hour period. Note the development of outgrowths from 72-hour time-point onwards (yellow
arrowheads) in the mock treated *as2-7D* samples (panels E and G), which are absent in NPA
treated *as2-7D* flowers (panels F and H). The *as2-7D* flowers harbor pLH13 (*35S::mCitrine-*
RCI2A) cell membrane marker (greyscale). Trichomes were removed from the outer sepal
epidermis by image processing. Scalebar, 50 μm.
1686
1687
1688
1689

Supplementary Table S1: List of primers used in this study.

Supplementary Data S1: Raw data for making the plots

Supplementary Movie S1: Confocal images of the outer and inner epidermis of adult sepals of the WT (left) and *as2-7D* mutant (right). Genotypes harbor pLH13 (*p35S::mCitrine-RCI2A*) plasma membrane marker. Scalebar, 200 μm.

Supplementary Movie S2: Confocal images of developing WT (left) and *as2-7D* mutant (right) flowers imaged every 24 hours, over 96-hour time period. Genotypes harbor pLH13 (*p35S::mCitrine-RCI2A*) plasma membrane marker. Trichomes were removed from the outer sepal epidermis by image processing for better visualization of sepal morphology. Scalebar, 50 μm.

Supplementary Movie S3: Confocal images show the outer epidermis of adult sepals of WT (top left), *as2-7D* (top right), *pML1::KRP1* (bottom left) and *pML1::KRP1 as2-7D* (bottom right). Note the largely rescued sepal smoothness in *pML1::KRP1 as2-7D* compared to *as2-7D*, due to an increase in the longitudinally aligned giant cells. Sepals were stained with Propidium-iodide, which stains the cell walls. Scalebar, 200 μm.

Supplementary Movie S4: Confocal images of developing *as2-7D* mutant flowers grown in the presence of mock (left) or 10μM NPA (right). Images were captured every 24 hours over the 96 hour time period. The *as2-7D* mutant harbors pLH13 (*p35S::mCitrine-RCI2A*) plasma membrane marker. Trichomes were removed from the outer sepal epidermis by image processing. Note the reduction in outgrowth formation in 10μM NPA-treated samples despite buckling. Scalebar, 50 μm.

REFERENCES AND NOTES

1.   D. P. Richman, R. M. Stewart, J. W. Hutchinson, V. S. Caviness, Jr., Mechanical model of brain convolutional development. *Science* **189**, 18-21 (1975).
2.   E. Hannezo, J. Prost, J. F. Joanny, Instabilities of monolayered epithelia: shape and structure of villi and crypts. *Phys Rev Lett* **107**, 078104 (2011).
3.   T. Savin *et al.*, On the growth and form of the gut. *Nature* **476**, 57-62 (2011).
4.   A. E. Shyer *et al.*, Villification: how the gut gets its villi. *Science* **342**, 212-218 (2013).
5.   R. L. Antoniou Kourounioti *et al.*, Buckling as an origin of ordered cuticular patterns in flower petals. *Journal of The Royal Society Interface* **10**, 20120847 (2013).
6.   E. Moyroud *et al.*, Cuticle chemistry drives the development of diffraction gratings on the surface of Hibiscus trionum petals. *Current Biology* **32**, 5323-5334.e5326 (2022).

7.   C. H. Waddington, Preliminary Notes on the Development of the Wings in Normal and
Mutant Strains of Drosophila. *Proc Natl Acad Sci U S A* **25**, 299-307 (1939).

8.   J. A. Thomson, On Growth and Form. *Nature* **100**, 21-22 (1917).

9.   H. Tsukaya, Leaf shape: genetic controls and environmental factors. *Int J Dev Biol* **49**,
547-555 (2005).

10.   H. Tsukaya, Mechanism of leaf-shape determination. *Annu Rev Plant Biol* **57**, 477-496
(2006).

11.   A. Boudaoud, An introduction to the mechanics of morphogenesis for plant biologists.
Trends Plant Sci **15**, 353-360 (2010).

12.   B. Li, Y.-P. Cao, X.-Q. Feng, H. Gao, Mechanics of morphological instabilities and
surface wrinkling in soft materials: a review. *Soft Matter* **8**, 5728-5745 (2012).

13.   C. M. Nelson, On Buckling Morphogenesis. *J Biomech Eng* **138**, 021005 (2016).

14.   A. E. Shyer *et al.*, Emergent cellular self-organization and mechanosensation initiate
follicle pattern in the avian skin. *Science* **357**, 811-815 (2017).

15.   V. D. Varner, J. P. Gleghorn, E. Miller, D. C. Radisky, C. M. Nelson, Mechanically
patterning the embryonic airway epithelium. *Proceedings of the National Academy of*
Sciences **112**, 9230-9235 (2015).

16.   J. L. Silverberg *et al.*, 3D imaging and mechanical modeling of helical buckling in
Medicago truncatula plant roots. *Proc Natl Acad Sci U S A* **109**, 16794-16799 (2012).

17.   J. Yan, B. Wang, Y. Zhou, S. Hao, Resistance from agar medium impacts the helical
growth of Arabidopsis primary roots. *Journal of the Mechanical Behavior of Biomedical*
Materials **85**, 43-50 (2018).

18.   R. Sachse *et al.*, Snapping mechanics of the Venus flytrap (Dionaea muscipula). *Proc*
Natl Acad Sci U S A **117**, 16035-16042 (2020).

19.   Y. Forterre, J. M. Skotheim, J. Dumais, L. Mahadevan, How the Venus flytrap snaps.
Nature **433**, 421-425 (2005).

20.   P. B. Green, Pattern Formation in Shoots: A Likely Role for Minimal Energy
Configurations of the Tunica. *International Journal of Plant Sciences* **153**, S59-S75
(1992).

21.   J. Dumais, C. R. Steele, New Evidence for the Role of Mechanical Forces in the Shoot
Apical Meristem. *Journal of Plant Growth Regulation* **19**, 7-18 (2000).

22.   J. Żebrowski, Dynamic behaviour of inflorescence-bearing Triticale and Triticum stems.
Planta **207**, 410-417 (1999).

23.   M. Fournier, J. Dlouhá, G. Jaouen, T. Almeras, Integrative biomechanics for tree
ecology: beyond wood density and strength. *Journal of Experimental Botany* **64**, 4793-
4815 (2013).

24.   S. M. Chapotin, J. H. Razanameharizaka, N. M. Holbrook, A biomechanical perspective
on the role of large stem volume and high water content in baobab trees (Adansonia spp.;
Bombacaceae). *Am J Bot* **93**, 1251-1264 (2006).

25.   R. Méndez-Alonzo *et al.*, Root biomechanics in Rhizophora mangle: anatomy,
1770   morphology and ecology of mangrove's flying buttresses. *Annals of Botany* **115**, 833-840
1771   (2015).

1772   26.   C. Huang, Z. Wang, D. Quinn, S. Suresh, K. J. Hsia, Differential growth and shape
1773   formation in plant organs. *Proceedings of the National Academy of Sciences* **115**, 12359-
1774   12364 (2018).

1775   27.   J. Derr, R. Bastien, É. Couturier, S. Douady, Fluttering of growing leaves as a way to
1776   reach flatness: experimental evidence on <i>Persea americana</i>. *Journal of The Royal*
1777   *Society Interface* **15**, 20170595 (2018).

1778   28.   C. D. Whitewoods, Riddled with holes: Understanding air space formation in plant
1779   leaves. *PLoS Biol* **19**, e3001475 (2021).

1780   29.   J. R. Evans, Improving photosynthesis. *Plant Physiol* **162**, 1780-1793 (2013).

1781   30.   A. H. K. Roeder, Arabidopsis sepals: A model system for the emergent process of
1782   morphogenesis. *Quantitative Plant Biology* **2**, e14 (2021).

1783   31.   L. Hong *et al.*, Variable Cell Growth Yields Reproducible OrganDevelopment through
1784   Spatiotemporal Averaging. *Dev Cell* **38**, 15-32 (2016).

1785   32.   H. Iwakawa *et al.*, The ASYMMETRIC LEAVES2 gene of Arabidopsis thaliana,
1786   required for formation of a symmetric flat leaf lamina, encodes a member of a novel
1787   family of proteins characterized by cysteine repeats and a leucine zipper. *Plant Cell*
1788   *Physiol* **43**, 467-478 (2002).

1789   33.   J. L. Bowman, Y. Eshed, S. F. Baum, Establishment of polarity in angiosperm lateral
1790   organs. *Trends Genet* **18**, 134-141 (2002).

1791   34.   W. C. Lin, B. Shuai, P. S. Springer, The Arabidopsis LATERAL ORGAN
1792   BOUNDARIES-domain gene ASYMMETRIC LEAVES2 functions in the repression of
1793   KNOX gene expression and in adaxial-abaxial patterning. *Plant Cell* **15**, 2241-2252
1794   (2003).

1795   35.   G. Wu *et al.*, KANADI1 regulates adaxial-abaxial polarity in Arabidopsis by directly
1796   repressing the transcription of ASYMMETRIC LEAVES2. *Proc Natl Acad Sci U S A*
1797   **105**, 16392-16397 (2008).

1798   36.   Y. Eshed, S. F. Baum, J. V. Perea, J. L. Bowman, Establishment of polarity in lateral
1799   organs of plants. *Curr Biol* **11**, 1251-1260 (2001).

1800   37.   R. A. Kerstetter, K. Bollman, R. A. Taylor, K. Bomblies, R. S. Poethig, KANADI
1801   regulates organ polarity in Arabidopsis. *Nature* **411**, 706-709 (2001).

1802   38.   X. Lin *et al.*, LFR is functionally associated with AS2 to mediate leaf development in
1803   Arabidopsis. *Plant J*, (2018).

1804   39.   N. Hervieux *et al.*, A Mechanical Feedback Restricts Sepal Growth and Shape in
1805   Arabidopsis. *Curr Biol*, (2016).

1806   40.   Y. Onoda, F. Schieving, N. P. Anten, A novel method of measuring leaf epidermis and
1807   mesophyll stiffness shows the ubiquitous nature of the sandwich structure of leaf laminas
1808   in broad-leaved angiosperm species. *J Exp Bot* **66**, 2487-2499 (2015).

41. A. Vijayan *et al.*, A digital 3D reference atlas reveals cellular growth patterns shaping the
Arabidopsis ovule. *eLife* **10**, e63262 (2021).
42. C. Mollier *et al.*, Spatial consistency of cell growth direction during organ morphogenesis
requires CELLULOSE SYNTHASE INTERACTIVE1. *Cell Rep* **42**, 112689 (2023).
43. S. Harmansa, A. Erlich, C. Eloy, G. Zurlo, T. Lecuit, Growth anisotropy of the
extracellular matrix shapes a developing organ. *Nature Communications* **14**, 1220 (2023).
44. L. Hong *et al.*, Heterogeneity and Robustness in Plant Morphogenesis: From Cells to
Organs. *Annual Review of Plant Biology* **69**, 469-495 (2018).
45. A. W. Sylvester, M. H. Williams, P. B. Green, Orientation of cortical microtubules
correlates with cell shape and division direction. *Protoplasma* **153**, 91-103 (1989).
46. A. H. Roeder *et al.*, Variability in the control of cell division underlies sepal epidermal
patterning in Arabidopsis thaliana. *PLoS Biol* **8**, e1000367 (2010).
47. S. M. Bemis, K. U. Torii, Autonomy of cell proliferation and developmental programs
during Arabidopsis aboveground organ morphogenesis. *Dev Biol* **304**, 367-381 (2007).
48. A. Sapala, R. S. Smith, Osmotic Treatment for Quantifying Cell Wall Elasticity in the
Sepal of Arabidopsis thaliana. *Methods Mol Biol* **2094**, 101-112 (2020).
49. T. Yamaguchi, A. Nukazuka, H. Tsukaya, Leaf adaxial-abaxial polarity specification and
lamina outgrowth: evolution and development. *Plant Cell Physiol* **53**, 1180-1194 (2012).
50. K. Fukushima, M. Hasebe, Adaxial–abaxial polarity: The developmental basis of leaf
shape diversity. *genesis* **52**, 1-18 (2014).
51. A. Y. Husbands, D. H. Chitwood, Y. Plavskin, M. C. Timmermans, Signals and
prepatterns: new insights into organ polarity in plants. *Genes Dev* **23**, 1986-1997 (2009).
52. R. Waites, A. Hudson, phantastica: a gene required for dorsoventrality of leaves in
Antirrhinum majus. *Development* **121**, 2143-2154 (1995).
53. J. R. McConnell, M. K. Barton, Leaf polarity and meristem formation in Arabidopsis.
Development **125**, 2935-2942 (1998).
54. M. P. Caggiano *et al.*, Cell type boundaries organize plant development. *eLife* **6**, e27421
(2017).
55. D. Otsuga, B. DeGuzman, M. J. Prigge, G. N. Drews, S. E. Clark, REVOLUTA regulates
meristem initiation at lateral positions. *Plant J* **25**, 223-236 (2001).
56. K. Watanabe, K. Okada, Two discrete cis elements control the Abaxial side-specific
expression of the FILAMENTOUS FLOWER gene in Arabidopsis. *Plant Cell* **15**, 2592-
2602 (2003).
57. E. Semiarti *et al.*, The ASYMMETRIC LEAVES2 gene of Arabidopsis thaliana regulates
formation of a symmetric lamina, establishment of venation and repression of meristem-
related homeobox genes in leaves. *Development* **128**, 1771-1783 (2001).
58. W. Wang *et al.*, YUCCA genes are expressed in response to leaf adaxial-abaxial
juxtaposition and are required for leaf margin development. *Plant Physiol* **157**, 1805-
1819 (2011).

59. K. Abley, S. Sauret-Güeto, A. F. Marée, E. Coen, Formation of polarity convergences
1849   underlying shoot outgrowths. *Elife* **5**, (2016).

1850 60. L. Gälweiler *et al.*, Regulation of polar auxin transport by AtPIN1 in Arabidopsis
1851   vascular tissue. *Science* **282**, 2226-2230 (1998).

1852 61. A. Izhaki, J. L. Bowman, KANADI and Class III HD-Zip Gene Families Regulate
1853   Embryo Patterning and Modulate Auxin Flow during Embryogenesis in Arabidopsis. *The*
1854   *Plant Cell* **19**, 495-508 (2007).

1855 62. D. Reinhardt *et al.*, Regulation of phyllotaxis by polar auxin transport. *Nature* **426**, 255-
1856   260 (2003).

1857 63. M. G. Heisler *et al.*, Alignment between PIN1 polarity and microtubule orientation in the
1858   shoot apical meristem reveals a tight coupling between morphogenesis and auxin
1859   transport. *PLoS Biol* **8**, e1000516 (2010).

1860 64. O. Hamant, E. M. Meyerowitz, J. Traas, Is cell polarity under mechanical control in
1861   plants? *Plant Signal Behav* **6**, 137-139 (2011).

1862 65. N. Nakayama *et al.*, Mechanical Regulation of Auxin-Mediated Growth. *Current Biology*
1863   **22**, 1468-1476 (2012).

1864 66. A. M. R. Kabir *et al.*, Buckling of Microtubules on a 2D Elastic Medium. *Scientific*
1865   *Reports* **5**, 17222 (2015).

1866 67. A. Sanyal, H. C. Han, Artery buckling affects the mechanical stress in atherosclerotic
1867   plaques. *Biomed Eng Online* **14 Suppl 1**, S4 (2015).

1868 68. A. H. K. Roeder, Arabidopsis sepals: A model system for the emergent process of
1869   morphogenesis. *Quant Plant Biol* **2**, (2021).

1870 69. L. Xu *et al.*, Novel as1 and as2 defects in leaf adaxial-abaxial polarity reveal the
1871   requirement for ASYMMETRIC LEAVES1 and 2 and ERECTA functions in specifying
1872   leaf adaxial identity. *Development* **130**, 4097-4107 (2003).

1873 70. S. Friesen, I. K. Hariharan, Coordinated growth of linked epithelia is mediated by the
1874   Hippo pathway. *bioRxiv*, 2023.2002.2026.530099 (2023).

1875 71. S. A. Braybrook, A. Peaucelle, Mechano-chemical aspects of organ formation in
1876   Arabidopsis thaliana: the relationship between auxin and pectin. *PLoS One* **8**, e57813
1877   (2013).

1878 72. P. B. Green, Expression of pattern in plants: combining molecular and calculus-based
1879   biophysical paradigms. *Am J Bot* **86**, 1059-1076 (1999).

1880 73. C. Kuhlemeier, Phyllotaxis. *Trends Plant Sci* **12**, 143-150 (2007).

1881 74. D. R. Smyth, J. L. Bowman, E. M. Meyerowitz, Early flower development in
1882   Arabidopsis. *The Plant Cell* **2**, 755-767 (1990).

1883 75. D. O. Robinson *et al.*, Ploidy and Size at Multiple Scales in the Arabidopsis Sepal. *The*
1884   *Plant Cell* **30**, 2308-2329 (2018).

1885 76. S. Strauss *et al.*, Using positional information to provide context for biological image
1886   analysis with MorphoGraphX 2.0. *eLife* **11**, e72601 (2022).

77.   A. Armezzani *et al.*, Transcriptional induction of cell wall remodelling genes is coupled
to microtubule-driven growth isotropy at the shoot apex in Arabidopsis. *Development*
145, (2018).
78.   M. G. Heisler *et al.*, Patterns of Auxin Transport and Gene Expression during
Primordium Development Revealed by Live Imaging of the Arabidopsis Inflorescence
Meristem. *Current Biology* **15**, 1899-1911 (2005).
79.   Y. Eshed, S. F. Baum, J. V. Perea, J. L. Bowman, Establishment of polarity in lateral
organs of plants. *Current Biology* **11**, 1251-1260 (2001).
80.   L. Hong, J. Brown, N. A. Segerson, J. K. Rose, A. H. Roeder, CUTIN SYNTHASE 2
Maintains Progressively Developing Cuticular Ridges in Arabidopsis Sepals. *Mol Plant*
10, 560-574 (2017).
81.   A. H. Roeder, A. Cunha, C. K. Ohno, E. M. Meyerowitz, Cell cycle regulates cell type in
the Arabidopsis sepal. *Development* **139**, 4416-4427 (2012).
82.   W. Lukowitz, C. S. Gillmor, W.-R. d. Scheible, Positional Cloning in Arabidopsis. Why
It Feels Good to Have a Genome Initiative Working for You1. *Plant Physiology* **123**,
795-806 (2000).
83.   D. O. Robinson *et al.*, Ploidy and Size at Multiple Scales in the Arabidopsis Sepal. *Plant*
Cell **30**, 2308-2329 (2018).
84.   M. Y. Zhou, C. E. Gomez-Sanchez, Universal TA cloning. *Curr Issues Mol Biol* **2**, 1-7
(2000).
85.   M. Karimi, D. Inze, A. Depicker, GATEWAY vectors for Agrobacterium-mediated plant
transformation. *Trends Plant Sci* **7**, 193-195 (2002).
86.   M. Zhu *et al.*, Robust organ size requires robust timing of initiation orchestrated by
focused auxin and cytokinin signalling. *Nat Plants* **6**, 686-698 (2020).
87.   P. J. Barrell, A. J. Conner, Minimal T-DNA vectors suitable for agricultural deployment
of transgenic plants. *BioTechniques* **41**, 708-710 (2006).
88.   A. A. Wise, Z. Liu, A. N. Binns, Three methods for the introduction of foreign DNA into
Agrobacterium. *Methods Mol Biol* **343**, 43-53 (2006).
89.   M. Zhu, A. H. Roeder, Plants are better engineers: the complexity of plant organ
morphogenesis. *Curr Opin Genet Dev* **63**, 16-23 (2020).
90.   A. Kiss *et al.*, Segmentation of 3D images of plant tissues at multiple scales using the
level set method. *Plant Methods* **13**, 114 (2017).
91.   Z. Y. Huang, W. Hong, Z. Suo, Nonlinear analyses of wrinkles in a film bonded to a
compliant substrate. *Journal of the Mechanics and Physics of Solids* **53**, 2101-2118
(2005).
92.   B. Audoly, A. Boudaoud, Buckling of a stiff film bound to a compliant substrate—Part I::
Formulation, linear stability of cylindrical patterns, secondary bifurcations. *Journal of the*
Mechanics and Physics of Solids **56**, 2401-2421 (2008).
93.   A. Goriely, *Morpho-elasticity: The mechanics and mathematics of biological growth*.
(Springer New York, 2016).

94.   M. Alnæs *et al.*, The FEniCS project version 1.5. **3**, (2015).

95.   F. Gacon, C. Godin, O. Ali, BVPy: A FEniCS-based Python package to ease the

expression and study of boundary value problems in Biology. *Journal of Open Source*

Software **6**, 1-6 (2021).

1931