AmpliDiff: An Optimized Amplicon Sequencing Approach to Estimating Lineage Abundances in Viral Metagenomes

Jasper van Bemmelen[1]   Davida S. Smyth[2]
Jasmijn A. Baaijens[1,3]

[1] Intelligent Systems Department, Delft University of Technology, Netherlands;
[2] Texas A&M University San Antonio, USA;
[3] Department of Biomedical Informatics, Harvard Medical School, USA
Corresponding author: `j.a.baaijens@tudelft.nl`

Abstract

Metagenomic profiling algorithms commonly rely on genomic differences between lineages, strains or species to infer the relative abundances of sequences present in a sample. This observation plays an important role in the analysis of diverse microbial communities, where targeted sequencing of 16S and 18S rRNA, both well-known hypervariable genomic regions, have led to insights in microbial diversity and the discovery of novel organisms. However, the variable nature of discriminatory regions can also act as a double-edged sword, as the sought after variability can make it difficult to design primers for their amplification through PCR. Moreover, the most variable regions are not necessarily the most informative regions for the purpose of differentiation; one should focus on regions which maximize the number of lineages that can be distinguished. Here we present AmpliDiff, a computational tool that simultaneously finds such highly discriminatory genomic regions, as well as primers allowing for the amplification of these regions. We show that regions and primers found by AmpliDiff can be used to accurately estimate relative abundances of SARS-CoV-2 lineages, for example in wastewater sequencing data. We obtain mean absolute prediction errors that are comparable with using whole genome information to estimate relative abundances. Furthermore, our results show that AmpliDiff is robust against incomplete input data, and that primers designed by AmpliDiff continue to bind to genomes originating from months after the primers were selected. With AmpliDiff we provide an effective and efficient alternative to whole genome sequencing for estimating lineage abundances in viral metagenomes.

Background

Studying the composition of metagenomic samples through genome sequencing has led to insights into diverse microbial communities [46, 11, 42]. While whole (meta)genome sequencing is frequently used for this purpose, not all regions of the genome are equally informative. If the goal is to distinguish between different lineages, strains, variants or species, we should focus on those regions which allow us to discriminate between the genomes of interest. For example, many studies focus only on 16S or 18S regions for metagenomic profiling [39, 38]. Such a targeted sequencing approach has two main advantages: first, it is cost-effective as it requires fewer sequencing reagents per sample, and second, it reduces the computational burden in downstream processing.

Targeted sequencing, otherwise known as amplicon sequencing, is a method in which genetic material of interest in (meta)genomic samples is amplified, using *Polymerase Chain Reaction* (*PCR*) for example, and later sequenced. A key step in the design of amplicon sequencing experiments is finding suitable regions of interest along with primers in order to perform the experiments. Depending on the application, the objective can be to identify the presence of particular genetic markers [13], or to differentiate between different species that are present [52, 57] for example. Moreover, if one is interested in *Whole Genome Sequencing* (*WGS*) it is also possible to design primers such that the resulting PCR products span the entirety of the genome(s) of interest.

Amplicon sequencing is widely applied in the context of *Wastewater-Based Epidemiology* (*WBE*) [1, 24, 37, 51, 56]. In wastewater-based epidemiology, the objective is to study population-scale phenomena such as trends in infections based on substance or pathogen concentrations found in wastewater samples [10]. Since this method is non-invasive and captures information from the entire population within an area, it is less prone to biases that occur in clinically obtained samples [3]. However, wastewater samples can suffer from problems like low concentrations of genetic material, environmental RNA degradation, the presence of inhibitors, and high fragmentation rates [43]. For these reasons, amplification by PCR is a crucial step in wastewater analysis.

The importance of WBE itself has become apparent during the SARS-CoV-2 pandemic. A multitude of studies have confirmed that SARS-CoV-2 can be detected in wastewater samples [5] and that we can estimate COVID-19 case numbers from such samples [33, 40, 41, 58]. Moreover, we can estimate the relative abundances of different SARS-CoV-2 lineages (and therefore different variants) from wastewater samples, using amplicon-based sequencing [4, 15, 28, 30, 47, 48, 61], further emphasizing the applicability of amplicon-based sequencing.

The key step in performing amplicon-based sequencing is determining a set of primers such that the genetic material of interest can be amplified [14]. Many physicochemical constraints (e.g. melting temperature, inter-primer interaction, intra-primer interaction) have to be considered, for which several different primer designing tools (e.g. Primer3 [54], Primer-BLAST [60], PriMux [22] and openPrimeR [34]) have been developed that take all constraints into account. These tools take as input a set of reference genomes and a region of interest and attempt to find a set of primers such that the region of interest can be amplified in as many reference genomes as possible. In addition, tools like PrimalScheme [44] have the option to find a near-minimal size set of primers such that the resulting amplicons would cover the entire genome enabling amplicon-based WGS.

While these tools have been successfully applied in practice, they only find primers once a region of interest, possibly the whole genome, has been provided. This suffices when these regions are known a priori, for example in 16S sequencing [39] and 18S sequencing [18]. However, when this is not the case there is a need for tools that can find regions of interest, along with primers simultaneously. Applications can range from the detection of pathogens [13], or alternatively facilitating the estimation of relative abundances of species present in a sample by focusing on variable regions in the genomes [50]. In the latter case, the focus should be on finding primers such that the resulting amplicons are different for different species in the sample. These steps should be considered simultaneously, as potential amplicons are only useable when feasible primers flanking them can be found. To our knowledge, there is no tool that integrates finding amplicons and corresponding primers for optimal differentiation between different genomic classes.

Here we introduce a new methodology, called AmpliDiff, that identifies discriminatory regions in a set of genomes and simultaneously finds primers such that these regions can be amplified through PCR. By considering multiple input genomes, AmpliDiff can find primers in genomic regions that are conserved within the input genomes, such that the resulting PCR products are able to discriminate between different genomes (or classes thereof). We demonstrate AmpliDiff by computing amplicons and corresponding primers for SARS-CoV-2 genomes. The effectiveness of these amplicons is assessed by evaluating the abundance estimation accuracy for different SARS-CoV-2 lineages on simulated data, comparing estimates on only selected amplicons versus whole genome sequencing. Finally, by showing that these primers continue to bind to genomes discovered months after they were generated, we show that AmpliDiff provides an efficient and economic alternative to whole genome

amplification that can be applied in the context of metagenomic profiling.

Results

An overview of AmpliDiff

AmpliDiff finds discriminatory amplicons with corresponding primers in three steps: (i) extracting a set of candidate primers, (ii) finding candidate amplicons and determining their ability to discriminate, (iii) greedily selecting amplicons and validating whether it is possible to find a set of primers to amplify the amplicon. Figure 1 outlines the overall methodology of AmpliDiff which we will briefly describe here, with a more thorough explanation available in the Methods section.

The input to AmpliDiff consists of multiple aligned genomes, each with a corresponding class. AmpliDiff then first extracts every physicochemically feasible primer of a given length and stores them in a *primer database* (Figure 1(i)) to be used in component (iii). Next, candidate amplicons are found and their differentiability is determined (Figure 1(ii)). We define the *differentiability* of an amplicon as the number of pairs of genomes of different classes that it can differentiate. In its final component, AmpliDiff heuristically tries to find a minimal set of amplicons with corresponding primers to differentiate between all the pairs of genomes. As can be seen in Figure 1(iii), this is done by iteratively finding the most differentiable amplicon and checking if it is possible to find forward and reverse primers to amplify it in the input genomes. Defining the *amplifiability* of an amplicon as the relative number of genomes in which both a forward and reverse primer flanking the amplicon can be found, AmpliDiff only accepts amplicons that meet a user-provided minimal required amplifiability. If this amplifiability is met, AmpliDiff minimizes the number of primer pairs needed to amplify the amplicon and adds the amplicon along with a minimal set of primers to the solution. Afterward, the differentiability of the remaining candidate amplicons is updated to correct for the pairs of genomes that can already be differentiated, and the amplicons are sorted (highest to lowest) based on the newly calculated differentiabilities. When it is not possible to amplify the amplicon in enough of the input genomes, the amplicon is discarded and the next most differentiable amplicon is considered. This cycle is repeated until either: (i) every pair of genomes can be differentiated, (ii) a predefined number of amplicons has been found, or (iii) there are no more candidate amplicons. The final output of AmpliDiff consists of the found set of amplicons (coordinates based on MSA) along with the corresponding forward and reverse primers.

Figure 1: Outline of the workflow of AmpliDiff. **(0)** Pre-processing consists of building a reference set from a database of genomes (e.g. GISAID) and applying multiple sequence alignment to obtain multiple aligned genomes. **(i)** AmpliDiff builds a database of feasible primers that can be considered during the greedy amplicon selection (iii) component. **(ii)** Feasible amplicon candidates are extracted from the multiple aligned genomes, and for every amplicon, the pairs of genomes it can differentiate are stored. **(iii)** In its final component, AmpliDiff greedily finds the most discriminatory amplicon, retrieves flanking primers, and determines if a (sub)set of primers can be found to amplify the amplicon in the reference genomes. If this is possible, the set of primers is minimized and the amplicon and primers are added to the solution, after which the discriminatory power of amplicons is updated and the amplicons are sorted accordingly. If amplification is not possible, the amplicon is discarded and the next most discriminatory amplicon is considered. This process is repeated until (a) all pairs of genomes can be differentiated, (b) a set number of amplicons is found, or (c) no more feasible amplicons exist. Finally, AmpliDiff outputs the selected amplicons and corresponding primer sets.

AmpliDiff finds highly differentiable amplicons in SARS-CoV-2 genomes

Using AmpliDiff we have determined a set of 10 amplicons, of widths 200 and 400 (corresponding to Illumina iSeq and MiSeq sequencing respectively) by using a global reference set of 2749 SARS-CoV-2 genomes over all 1837 lineages (1-7 genomes per lineage) that existed at the time the reference set was constructed (18 August 2022). The reference set was constructed by following the pre-processing steps described in the VLQ pipeline

[4], using all of the available genomes on GISAID [31], and were multiple aligned with MAFFT [29]. Every genome was supplemented with a corresponding lineage assigned through the Pango lineage system [45]. These lineages consist of closely related SARS-CoV-2 genomes which are defined by key phylogenetic markers and shared mutations, and the objective is to find amplicons that allow us to differentiate between these lineages.

Figure 2(a) shows the relative differentiability of all amplicons of widths 200 and 400 that have at least 50 nucleotides before and after them. From the figure we first observe that, given a starting position in the genome, a 400-width amplicon is always at least as differentiable as a 200-width amplicon since the 400-width amplicon contains all of the information contained in a 200-width amplicon and we thus expect that 400 width amplicons are generally more differentiable than 200 width amplicons. It can also be seen that both the ORF1a and ORF1b regions are relatively conserved between the reference genomes and that regions in the 3'-end of the genome are generally the most differentiable with amplicons reaching up to 93.3% differentiability. It is likely that different amplicons can differentiate between the same pairs of genomes, and as such the figure does not tell us anything about which amplicons to combine in order to attain a shared differentiability of 100%. Moreover, it is also not guaranteed that every amplicon is practically feasible since we would need to be able to find primers flanking the amplicons in order to amplify these regions through PCR, which could be difficult for the most differentiable amplicons, as these are generally located in ill-conserved regions of the genome. In these cases, the width of the differentiable region can become a limiting factor since a width larger than the amplicon width might prevent us from finding feasible primers flanking the amplicon.

Running AmpliDiff with the restriction to finding 10 amplicons (widths 200 and 400) with a minimal required amplifiability of 95% yields the amplicons shown in Figure 2(b). For both amplicon widths, we see that the first amplicon is chosen in the gene that encodes the nucleocapsid protein, which provides a relative differentiability of 67.1% and 70.8% for amplicons of widths 200 and 400 respectively. Subsequent amplicons are somewhat spread over the genome, but in both cases, we observe that at least one amplicon in the spike gene was selected. In the context of wastewater sequencing, this region often suffers from low sequencing depths due to problems with amplification such as amplicon drop-out which, in practice, makes it a difficult target to sequence with consistent results [6]. One potential explanation for this can be seen in Figure 2(a) where we observe that the variation between different SARS-CoV-2 lineages in the spike gene is spread over a relatively wide region. This could indicate that it is difficult to find amplicons and corresponding primers that collectively cover the full gene, even though it appears to be highly differentiable.

An additional interesting observation is that we find overlapping amplicons for both widths. While this may seem counter-intuitive, a possible explanation for this is the 95% minimal amplifiability condition. This makes it possible that AmpliDiff will first find an amplicon and require it to be amplifiable in 95% of the input genomes. If, at a later stage, there is still some differentiability to be found in the remaining 5% of the genomes in this region, AmpliDiff can try to find an overlapping amplicon to cover this. Alternatively, the differentiability is contained in a region wider than the pre-defined amplicon width in which case at least 2 overlapping amplicons would be required to capture it. This latter seems to be the case, as even when requiring an amplifiability of 100% overlapping amplicons are found (Figure 6) in the Supplementary Material.

It can also be seen that for both the 200 and 400-amplicon width solutions, the marginal differentiability seems to drop exponentially with every additional amplicon. Conceptually this makes sense as we expect every subsequent amplicon to be less differentiable than the previous since this is how the greedy algorithm picks amplicons. As a consequence, we see that the first 5 amplicons already allow us to differentiate between 81.6% and 88.7% of the pairs of genomes of different lineage for widths 200 and 400 respectively. Any further amplicons contribute approximately 0.4-1% to the cumulative relative differentiability indicating that, depending on the required resolution, it can be sufficient to use up to 5 amplicons.

The effect of different minimal required amplifiabilities

The amplicons considered so far are based on a minimal required amplifiability of 95%. To get an idea of how the minimal amplifiability affects the amplicons selected, and in particular the number of primers required we also ran AmpliDiff with minimal required amplifiabilities of 90%, 92.5%, 97.5%, 99.9% and 100%. The resulting cumulative differentiabilities for every respective configuration can be found in Figures 1-6 in the Supplementary Material. For minimal amplifiabilities below 99.9% we see that roughly the same regions were selected, whereas for higher required amplifiabilities the strictness in requirements becomes a bottleneck, and we observe that different amplicons are chosen. Nevertheless, we see that the initial amplicon is always located in the nucleocapsid gene and that the first few amplicons contribute the most to the cumulative differentiability. An interesting observation that can be seen in both the 99.9% and 100% minimal amplifiability results, is that the first amplicon has a larger relative differentiability than the first amplicons for the lower amplifiabilities. Even though any solution allowed for these higher amplifiability requirements is also valid when the required amplifiability is lower, this is due to the fact that our primer minimization model includes a trade-off between the number of primer pairs selected and the additional differentiability gained by adding them due to the amplicon being amplifiable in more input genomes. As a consequence, the model can decide to not add primer pairs if they allow for only a marginal increase in differentiability, hence attaining a slightly lower differentiability even when the same amplicon is added.

The impact of requiring a higher minimal amplifiability on the number of primer pairs can be seen in Table 1 in the Supplementary Material where, for every configuration, we show the minimal number of forward and reverse primers required to amplify the amplicon found by AmpliDiff. As can be seen from the Table, results are mostly similar for amplifiabilities below 97.5% with generally only a single primer pair being needed for the amplification of an amplicon in the corresponding number of input genomes. For minimal required amplifiabilities of 97.5%, 99.9%, and 100%, however, we see that both the number of forward and reverse primers substantially increase compared to lower amplifiabilities, with the 100% minimal amplifiability amplicons generally requiring the largest number of primers.

(a)

(b)

Figure 2: Overview of the differentiability of potential amplicons, and the amplicons selected by AmpliDiff on the global SARS-CoV-2 dataset. (a) Relative differentiability of all amplicons of widths 200 (blue) and 400 (orange) that have at least 50 nucleotides before or after them. (b) Cumulative relative differentiability of the ten best amplicons (both 200 and 400 widths) found by AmpliDiff.

Benchmarking AmpliDiff amplicons versus WGS

We show the effectiveness of amplicons selected by AmpliDiff by comparing abundance estimation results for SARS-CoV-2 lineages based on AmpliDiff amplicons to abundance estimation results based on whole genomes. To this end, we have generated two datasets: one for Texas and one for the Netherlands. Both datasets consist of a *reference set* (not the one used for AmpliDiff) which contains the genomes that the abundance estimator uses as reference material, and a *simulation set* that contains genomes to simulate reads from. For both datasets, the reference genomes are drawn from all genomes available on GISAID of the corresponding location between May and October of 2022. The simulation sets consist of all high quality genomes from November 2022. A more detailed explanation of the setup for the simulation study can be found in the Methods section.

We perform abundance estimation using VLQ [4], a pipeline for estimating relative abundance of viral lineages from wastewater sequencing data. VLQ employs kallisto [7] to assign to every genome in a set of reference genomes a relative abundance based on the likelihood that reads originate from that genome. The relative abundance of a lineage is then obtained by aggregating relative abundances of all genomes that are part of the lineage.

In the whole genome-based simulations we assume that read coverage and read depth are both uniform over the simulation genomes. From a practical point of view, this is an ideal but highly unrealistic situation, particularly so if we consider the context of wastewater sequencing [4]. Hence the whole genome-based results should be considered as idealized results only serving as a benchmark for the amplicon-based abundance estimations.

For both regions, we ran 20 simulations for both amplicon widths using the top 1, 2, 5 and 10 amplicons respectively. The results are shown in Figure 3, from which it can be seen that the abundance estimations show very little variance, indicating that results are consistent in all configurations. In general, we see that the *Mean Absolute Prediction Errors* (MAPEs) range between 0.6% and 1.3% with the whole genome-based approach usually achieving the smallest MAPE. The exception to this latter observation is the Texas dataset using 5 or 10 amplicons with width 400, or 10 amplicons with width 200. In these cases, we see that the amplicon-based approach outperforms the whole genome-based approach by a small margin.

Interestingly, we see that both 200 and 400-width amplicons yield similar results in the Texas dataset, whereas the 400-width amplicons strongly outperform the 200-width amplicons in the Netherlands dataset. In both regions we also see that using two amplicons of width 200 yields slightly worse MAPEs than using a single amplicon of width 200, indicating that the second amplicon does not provide more information and rather interferes with Kallisto's ability to assign relative abundances. Besides this, we do see that adding amplicons generally reduces the MAPE albeit at slow rates.

In contrast to the Texas dataset, we see a substantial discrepancy between estimation results in the Netherlands dataset. This is particularly prominent for the amplicons of width 200 where, even with 10 amplicons, the whole genome-based approach strongly outperforms the amplicon-based approach which can potentially be explained by the limited discriminatory power of short amplicons.

Alternatively, this discrepancy can also be due to the fact that nearly all of the genomes used in the simulation study are (sub)lineages or recombinant lineages of the B.1.1.529 lineage (Omicron). Omicron has, since its first documentation in November 2021, rapidly become the dominant variant nearly everywhere [36], which explains its prominence in these experiments. Omicron sublineages can be particularly hard to distinguish in metagenomic samples due to the large number of closely related sublineages. Moreover, as most sublineages of B.1.1.529 in the simulation datasets did not even exist (at least in the PANGO nomenclature) when the global reference set was determined, the amplicons used here aren't necessarily designed to differentiate between them.

To see whether this is actually the case, we show in Figures 4(a)-(d) the prediction errors made by all approaches for all lineages, but with the non-recombinant direct sublineages of B.1.1.529 considered as a single lineage (i.e. any lineage starting with B.1.1.529.1 is assigned to B.1.1.529.1). In the figures, it can be seen that at this resolution, most errors are made due to wrongly estimating the abundances of B.1.1.529 sublineages.

In addition, it can also be seen that the amplicon-based approach using 10 amplicons performs on par with the whole genome-based approach suggesting that, if the resolution is to mainly discriminate between different direct sublineages of B.1.1.529 and other lineages, the 10 amplicons found by AmpliDiff are able to do so. This observation is further supported by the results in Table 1 which shows the MAPE at this resolution for all approaches in both the Texas and Netherlands datasets. In the table, it can be seen that at this resolution, the amplicon-based approach using 10 amplicons actually outperforms the whole genome-based approach in all cases which provides further evidence that the amplicon-based results are affected by the Omicron sublineages. This shows that amplicons selected by AmpliDiff can discriminate reasonably well between sublineages of B.1.1.529, and outperforms the idealized whole genome-based approach on the resolution of discriminating between direct descendants of B.1.1.529

	200 width					400 width				
Amplicons	1	2	5	10	WGS	1	2	5	10	WGS
MAPE (TX)	3.619 (0.001)	1.391 (0.001)	2.729 (0.004)	0.638 (0.006)	3.674 (0.020)	5.049 (0.001)	3.842 (0.017)	1.960 (0.021)	0.936 (0.008)	3.718 (0.022)
MAPE (NL)	2.468 (0.005)	5.757 (0.002)	2.277 (0.008)	0.760 (0.011)	0.845 (0.028)	1.060 (0.053)	1.476 (0.036)	1.515 (0.161)	0.734 (0.012)	0.967 (0.039)

Table 1: Mean absolute prediction errors for both the amplicon-based approach (1, 2, 5, 10 amplicons) and the whole genome-based approach using fragment lengths of 200 and 400 over 20 simulations in both datasets when lineages are cut-off and aggregated at most 3 sublineage levels. In parentheses the standard deviations are given.

Figure 3: Mean absolute prediction errors for abundance estimations in both datasets for both amplicon widths using varying numbers of amplicons. **(a)** Texas dataset using amplicons of width 200. **(b)** Texas dataset using amplicons of width 400. **(c)** Netherlands dataset using amplicons of width 200. **(d)** Netherlands dataset using amplicons of width 400.

Figure 4: Prediction errors for both the amplicon-based approach (1, 2, 5, 10 amplicons) and the whole genome-based approach using sequencing fragments of lengths 200 and 400 in both datasets. The x-axes show the lineages (cut-off at most 3 sublineage levels) and the y-axis is the prediction error made by every approach.

AmpliDiff is robust against missing data

A problem with reference-based approaches in metagenomics is the selection of reference genomes included in an analysis. Due to the exponential growth of available genomes in genome databases [8], it might be necessary to concentrate only on a subset of genomes that are important for the problem that is being studied. On the other hand, there are also situations, such as the COVID-19 pandemic, where available information is limited and it might be impossible to find a reference set that is actually representative of the circulating population. In both scenarios, it is important to select an appropriate set of references that captures the diversity in the population, while not being excessively large as this can lead to inflated computational requirements. As a result, when designing reference sets it is possible that informative reference genomes are left out, either due to limitations in computational resources or because of scarcity in the available information. Here we consider what happens when AmpliDiff is presented with an incomplete reference set and the effect it has on the amplicons selected by AmpliDiff. For this, we take 15 subsamples of sizes 1500 (54.6%), 2000 (72.8%), and 2500 (90.9%) respectively from the reference set used here (2749 reference genomes) and run AmpliDiff with the same settings as before.

The results are shown in Figures 5(a) and (b) where we can see how often a nucleotide position is covered by an amplicon in a subsample, relative to the total number of subsamples, for amplicons of width 200 and 400 respectively with a minimal amplifiability requirement of 95%. As a general observation we see that, regardless of the number of excluded reference sequences and amplicon length, the majority of the results agree on which areas of the genome to cover. Moreover, the regions that are covered in at least 80% of these experiments also

appear to overlap with the initial amplicons selected based on the full reference set. The variation in covered nucleotide positions that can be observed, however, stems from amplicons selected in the later stages of the greedy cycle. Such amplicons generally contribute only marginally to the total differentiability of the selected amplicons, and thus when reference genomes are left out it is to be expected that these amplicons will be more spread out over the genome as it is likely that there are multiple candidate amplicons that share a similar differentiability. Nevertheless, these results indicate that AmpliDiff is robust against missing reference genomes, even when nearly half of the reference genomes are excluded.

Figure 5: Assessment of the robustness of amplicons selected by AmpliDiff through subsampling subsets of sizes 1500, 2000, 2500 of the reference genomes 15 times for both amplicon widths. **(a)** shows for every position (in the aligned genomes) how often it was contained in an amplicon of width 200 in each of the subsampling experiments, whereas **(b)** shows the same results for amplicons of width 400.

Primers and amplicons found by AmpliDiff remain viable over time

A crucial aspect of amplicon design for the sake of discriminating between genomes, is that amplicons remain amplifiable on genomes that are not present in the dataset used to determine the amplicons. This aspect is conceptually analogous to the problem of overfitting in the context of machine learning where a model is trained on a training set but validated on a validation or test set of data that is unobserved during the training stages. The ability to perform well on a test set indicates that the model is not overfitted for the training data, but is able to generalize to data yet unobserved. In our context, this translates to finding out whether amplicons found by AmpliDiff are still amplifiable on genomes that were not used in the reference set fed to AmpliDiff. As amplicons are only amplifiable in a genome if both a forward and reverse primer flanking the amplicon will bind to the genome this implies we can check how well the amplicons determined by AmpliDiff work on "new" genomes, by checking for every amplicon whether both a forward and reverse primer of the corresponding primer set bind. To validate how well the amplicons found so far amplify in new genomes, we have downloaded high-quality genomes from GISAID from September 2022 to May 2023 and checked the amplifiability of every amplicon in these genomes.

The amplifiability per month of every amplicon based on the 95% minimal amplifiablity requirement is shown in Figures 6 **(a)** and **(b)** for amplicon widths 200 and 400 respectively. From Figure 6**(a)** it can be observed that every 200-width amplicon retains an amplifiability of at least 95% in every month, with the exception of the final amplicon (17,042-17,242). This amplicon, located in ORF1b, appears to become significantly less amplifiable over time reaching an amplifiability of approximately 70% in April 2023. Nevertheless, the amplifiability seems to increase again in May 2023 and as this is the final amplicon selected by AmpliDiff it is expected to not have a significant effect on the ability to perform abundance estimation using the selected amplicons.

For amplicons of width 400 (Figure 6**(b)**) we see more variation in amplifiability, particularly in amplicons 6, 7, and 9. The pattern in amplifiability is also of interest, with amplicons 6 and 9 seeming to become less

amplifiable over time, whereas amplicon 7 seems to regain some of its amplifiability in May 2023, similar to amplicon 10 in the 200-width amplicons. As this again seems to affect less differentiable amplicons, we also expect this to not strongly affect the effectiveness of abundance estimation using these amplicons.

Amplifiability results for amplicons based on different minimal amplifiability requirements are shown in Figures 7-12 in the Supplementary Material. In these figures, we see similar trends in amplifiability which is due to the fact that the same amplicons are chosen for different minimal required amplifiabilities. The exception to this are amplicons selected when the amplifiability is required to be 99.9% or 100%. In these cases, we observe that a single amplicon tends to be less amplifiable in both amplicon widths. Together, these observations suggest that the amplicons and primers found by AmpliDiff remain usable over time, with only a few relatively weakly differentiable amplicons experiencing some dropout.

Figure 6: Fraction of genomes per month in which amplicons bind. **(a)** Amplifiability of 200-width amplicons generated with the 95% amplifiability requirement in the genomes from September 2022 to May 2023. **(b)** Amplifiability of 400 width amplicons generated with the 95% amplifiability requirement in the genomes from September 2022 to May 2023.

Computational requirements

All the results in this paper were obtained by running AmpliDiff on a High-Performance Computing (HPC) cluster, using 200G RAM and 12 CPUs. Table 2 shows the average running time (over 15 runs) of AmpliDiff on 1500, 2000, and 2500 genomes for a minimal required amplifiability of 95%. In the table, we can see that the runtime of every step tends to increase with the number of included reference sequences. This is a natural consequence of the fact that each of the steps goes over all of the reference genomes.

If we assume that determining the feasibility of a primer of length k takes $O(f(k))$ time, the primer database

construction needs to parse every k-mer (k being equal to the primer length) in each of the reference genomes resulting in a theoretical runtime of $O(|\mathcal{G}| \cdot |g^*| \cdot f(k))$, where $|\mathcal{G}|$ is the number of included reference genomes and $|g^*|$ is the length of the largest genome. This is, however, a highly inflated runtime since the primer database keeps track of every k-mer that occurs by storing them in a dictionary. As a consequence, every subsequent time a k-mer occurs, it does not have to be checked for all its properties, thus reducing the expected runtime in practice.

Determining the feasible amplicons is done by considering every window of the amplicon size (either 200 or 400 here), and calculating the number of misalignment characters. This can be achieved in $O(|\overline{g}|)$ time for every multiple aligned genome $\overline{g} \in \mathcal{G}$ as every character has to be considered exactly once. Calculating the cumulative number of misalignment characters can also be done in $O(|\overline{g}|)$ per genome which results in a theoretical runtime of $O(|\mathcal{G}| \cdot |\overline{g}|)$ for determining the feasibility of amplicons. In a similar fashion, calculating the amplicon differentiability is done by moving a window with a size equal to the amplicon length over all pairs of genomes of different class and calculating the number of positions where a pair disagree, which yields a runtime of $O(|\mathcal{G}|^2 \cdot |\overline{g}|)$. This term would seem to dominate the runtime of constructing the primer database, but we see from Table 2 that this is not the case in practice. While this may seem counter-intuitive, it is a consequence of the factor $f(k)$ in the runtime of the primer database construction which accounts for checking all of the primer attributes, such as melting temperature, GC-content, and self-dimerization risk among others.

Finally, we should comment on the runtimes for the greedy algorithm. In contrast to the greedy algorithm for the set cover problem, it is difficult to provide a theoretical runtime for the greedy amplicon picking algorithm. This is due to two factors. First, we do not know which amplicons will have appropriate flanking primers a priori, which would require one to make assumptions about the likelihood of this being the case in order to come up with theoretical runtimes. The second-factor ties into this, as currently, AmpliDiff determines whether an amplicon can be amplified in enough reference genomes by solving a mixed integer linear program (MILP). While we do not have results on the hardness of the corresponding MILP, its relation to the set cover problem and maximum coverage problem does suggest that it is likely to be NP-hard. Consequently, the runtime of deciding whether an amplicon can be added to the solution, and minimizing the number of primer pairs used for this relies on solving the MILP through a generic solver (Gurobi [17]), which we can assume to have an exponential runtime in the number of variables and/or constraints in the model. This explains why, in Table 2, we see that the greedy algorithm strongly dominates the overall runtime, with runtimes seemingly increasing exponentially with the number of included genomes.

| | Amplicon width 200 | | | | Amplicon width 400 | | | |
Task	1500	2000	2500	2749*	1500	2000	2500	2749*
Constructing primer database	390s (119s)	682s (292s)	695s (179s)	673s	446s (131s)	548s (118s)	695s (178s)	818s
Determining feasible amplicons	100s (22s)	123s (23s)	164s (42s)	144s	96s (22s)	123s (29s)	163s (35s)	142s
Calculating amplicon differentiability	151s (36s)	274s (30s)	440s (102s)	526s	194s (33s)	351s (50s)	560s (127s)	602s
Greedy algorithm	10,825s (2,878s)	16,089s (3,062s)	23,239s (4,926s)	23,651s	8,941s (2,376s)	13,358s (3,085s)	22,121s (4,134s)	27,014s

Table 2: Average running time of AmpliDiff on an HPC cluster using 200G RAM and 12 CPU cores for different numbers of included genomes. Standard deviations are shown in parentheses, with the exception of the full reference set running times for which we performed only a single run.

Discussion

Here, we introduce AmpliDiff, a computational tool designed to find discriminatory genomic regions along with primers with the objective of discriminating between different lineages (or classes) of genomes. Without requiring prior knowledge of the input genomes beyond their respective lineages, AmpliDiff avoids relying on pre-defined regions of interest such as the 16S rRNA gene in bacteria and archaea [59], or the internal transcribed

spacer (ITS) region in fungi [49], and instead can find combinations of genomic regions that collectively allow for the differentiation between genomes. By focusing only on discriminatory genomic regions, AmpliDiff can provide an effective and economic alternative to amplification-based whole genome sequencing in the context of metagenomic profiling, which can be susceptible to problems such as amplification bias [2] and often requires a substantial number of primer pairs. Through a simulation study, we show that the amplicons found by AmpliDiff can be used to obtain accurate and consistent abundance estimations for SARS-CoV-2 genomes. Moreover, by comparing these abundance estimations with abundance estimations based on a whole genome sequencing approach (which can require up to hundreds of primer pairs), we find that the amplicon-based results can generally achieve similar accuracy, despite only using up to 10 amplicons.

Although AmpliDiff is based on a reference database, we see that amplicons selected by AmpliDiff remain highly discriminatory even when the reference set used to determine the amplicons is not up to date. In particular, we observe that lineages get redefined over time, leading to misleading input for AmpliDiff: the amplicon selection approach may fail to distinguish between sequences from different lineages as these were initially assigned to the same lineage. Hence, results could improve even further if the lineage assignment at the time of amplicon selection is stable.

A shortcoming of our benchmarking study is that we compare abundance estimations based on AmpliDiff amplicons, which are processed to take the possibility of primers not binding (and hence amplicons not being amplified) into account, with abundance estimations based on reads that cover the full genomes with uniform sequencing depth. Achieving uniform coverage through amplicon-based sequencing can be difficult to realize [2], which means that the WGS results are highly unrealistic to obtain in practice. This could be addressed by considering a similar setup where primer binding is taken into account during read simulation. The ARTIC primers have been widely used in practice and are regularly updated to account for new mutations occurring in the SARS-CoV-2 genome [35]. However, in our experiments we check for exact primer binding, which the ARTIC primers are not designed for and under our conditions the ARTIC v4.1 amplicons would not amplify. If we were to incorporate less stringent primer binding conditions, we would expect to see two effects: 1) AmpliDiff amplicons may amplify better hence results will likely improve, and 2) the WGS-based results would likely suffer from issues such as amplicon dropout.

It is also of particular importance to experimentally show that primers found through primer design tools, AmpliDiff included, work in practice. This could be validated through in vitro experiments using synthethic populations and spiked-in samples of known variant composition. Such an approach would also lend itself to evaluate abundance estimation accuracy using AmpliDiff amplicons in practice. It would be of interest to then compare this against abundance estimations based on ARTIC primers and to verify the effectiveness of primers and amplicons found by AmpliDiff in the context of SARS-CoV-2.

In addition, it would also be of interest to apply AmpliDiff to sequence databases for other species, such as Influenza, SARS, measles, polio or HIV. A prerequish2ite for any such analysis is a database with sufficiently many high-quality genomes. By using AmpliDiff for different species, we will gain additional insights into AmpliDiff's effectiveness and limitations, allowing for further improvements to our methodology.

There are several aspects in AmpliDiff that can be improved. For example, the input genomes are pre-processed using MSA; since this is an NP-hard problem, we use a heuristic which can lead to sub-optimal alignments. Moreover, MSA is not well suited for genomes with extensive recombination or repetitiveness. Alternatively, pangenome representations (e.g. variation graphs [16] or colored de Bruijn graphs [23]) could serve as a data structure for AmpliDiff to work with. Another area for improvement is the fixed-length primer selection. In the current implementation, allowing variable-length primers would lead to a combinatorial explosion of the number of possible primer pairs. Finally, our primer feasibility algorithm relies on solving a MILP, which is related to the set cover problem and is therefore expected to be NP-hard. To address this problem efficiently, dedicated heuristics are needed.

Conclusions

With AmpliDiff we introduce a computational tool to find genomic regions along with primers that can be used in the context of metagenomic profiling. Through a simulation study on SARS-CoV-2, we have shown that using amplicons found by AmpliDiff gives accurate and consistent abundance estimates of SARS-CoV-2 lineages, competitive with results based on WGS. Thus, AmpliDiff provides an effective and economic alternative to whole genome-based metagenomic profiling.

Methods

AmpliDiff methodology

The input to AmpliDiff consists of a reference set of multiple aligned genomes $\mathcal{G}$, an amplicon width L_A, a primer width k, a primer search window length L_0, a maximum misalignment character threshold $L_\emptyset$, a minimal differentiability threshold D and a minimal amplifiability threshold γ. In addition, AmpliDiff also accepts a list of classes for every genome in $\mathcal{G}$, a trade-off parameter β that is used to model the trade-off between adding primer pairs and the additional sequences in which an amplicon can be amplified and finally a set of physicochemical properties that the primers have to satisfy. As shown in Figure 1, AmpliDiff consists of three steps: (i) a pre-processing step that builds a database of feasible primers for the reference genomes, (ii) an amplicon pre-processing step that extracts feasible amplicons and calculates their differentiability, and (iii) a greedy amplicon selection procedure in which iteratively finds the most differentiable amplicon and adds it to the solution if primers can be found to amplify the amplicon. Here we give a description of these steps.

Step (i): constructing the feasible primer database

The objective of AmpliDiff is to find amplicons along with primers such that they can be amplified with PCR. For this, it is necessary to consider the potential primer candidates (i.e. k-mers) that can be used from the genomes in $\mathcal{G}$. By parsing the un-aligned reference genomes, which can be obtained by removing misalignment characters, AmpliDiff extracts all k-mers, which are filtered according to the criteria listed in the Supplementary Material. These criteria include GC-content ranges, melting temperature ranges, risk of hairpin formation and risk of self-dimerization, which are properties that are commonly considered in state-of-the-art primer design tools such as Primer3 [54], OpenPrimeR [34] and PriMux [22] as they are likely to affect the effectiveness of primers [14] and can be estimated from the nucleotide sequences of the primers. Potential primers that do not meet these criteria, as well as primers that occur at multiple locations in a single genome, are deemed infeasible in order to prevent generating unwanted byproducts during PCR.

After extracting all of the primer candidates, AmpliDiff filters the infeasible primers from the primer database resulting in a set of feasible primers $\mathcal{P}$ that can be partitioned into forward primers $\mathcal{P}_F$ and reverse primers $\mathcal{P}_R$. An important property of AmpliDiff is that it allows reference genomes to contain degenerate nucleotides represented by their corresponding IUPAC code [25]. When a genome contains a degenerate k-mer, AmpliDiff will try to disambiguate it by considering every possible representation of the k-mer. This approach differs from conventional approaches where degenerate nucleotides are either replaced by pseudo-random nucleotides or kept degenerate which potentially results in degenerate primers. If a k-mer occurring in a genome is too degenerate (i.e. has too many representative non-degenerate k-mers), this step is omitted, and the k-mer is skipped. The default threshold is 4^5 which corresponds to at most 1024 different representations of a single k-mer, although this threshold can be changed by the user. Doing so always results in a set of non-degenerate primers which, through post-processing, could be turned into degenerate primers if required.

Step (ii): amplicon pre-processing

By requiring that the input genomes $\mathcal{G}$ are multiple aligned, it is guaranteed that similar and dissimilar regions in the genomes are aligned. Another consequence of multiple alignments is that all of the genomes in $\mathcal{G}$ have the same length, which we will denote L_G. In what follows we will also adhere to the following notation. Given a genome $g \in \mathcal{G}$, we define $g[i]$ as the nucleotide at position i in genome g (starting with index 0). Similarly, for natural numbers $i < j \leq L_G$, we let $g[i:j] = g[i, i+1, \ldots, j-1]$ denote the substring of g starting at position i and ending before position j. Using this notation and the fact that all genomes in $\mathcal{G}$ have the same length, we can define an amplicon A as an interval $[i : i + L_A]$, where L_A is the length of the amplicon. To simplify notation we will write $g[A] := g[i : i + L_A]$ to refer to the substring defined by the interval $A = [i : i + L_A]$, in genome g.

In order for multiple aligned genomes to have the same length, they are padded by gap characters ('-') indicating that a genome contains an indel at a given position, relative to the genomes it is multiple aligned against. The problem with this, however, is that multiple sequence alignment is an NP-hard problem under general conditions [26]. As a result, most multiple sequence alignment programs (e.g. MAFFT [29] or ClustalW [53]) heuristically find good solutions which are not guaranteed to be optimal. These methods generally provide sub-optimal alignments in practice, which can have additional gaps that could potentially lead to finding discriminatory regions based on falsely identified regions with indels. With AmpliDiff we combat this problem in two ways.

First, we include a threshold, $L_\emptyset$ (default value equal to 10% of the amplicon length), for the maximum number of gap characters that may occur in an amplicon in all of the reference genomes. More concretely, we define $\emptyset(g[i:j]) := |\{lin[i, i+1, \ldots, j-2, j-1] : g[l] = \text{'-'}\}|$ as the number of gap characters in genome g, on the interval $[i : j]$. Then, by requiring that $\emptyset(g[A]) \leq L_\emptyset \ \forall g \in \mathcal{G}$ for all considered amplicons A we enforce the constraint that any input genome can only contain up to $L_\emptyset$ gap characters. By selecting an appropriate value for $L_\emptyset$ (default is 10% of the amplicon length), we remove amplicons that are likely affected by an inflated number of gap characters due to the sub-optimal multiple sequence alignment. Additionally, this also enforces that the final PCR fragments will have approximately the same length.

In addition to classifying amplicons as infeasible due to large gaps, we apply another filter on amplicons in order to make sure that space directly flanking amplicons is available to find primers in. This is achieved by requiring that the starting position of amplicons is at least $i_{\text{start}} := \min\{i \in \{0, \ldots, L_G\} : \emptyset^c(g[0:i]) \geq L_s \ \forall g \in \mathcal{G}\}$, and the ending position is smaller than $i_{\text{end}} := \max\{i \in \{0, \ldots, L_G - 1\} : \emptyset^c(g[i : L_G]) \geq L_s \ \forall g \in \mathcal{G}\}$, as then every genome in $\mathcal{G}$ has at least L_G positions to find primers flanking an amplicon. Together with the previous condition, this yields a set of feasible amplicon candidates $\mathcal{A} := \{[i : i + L_A] : i \geq i_{\text{start}} \ \wedge \ i < i_{\text{end}} \ \wedge \emptyset(g[A]) \leq L_\emptyset \ \forall g \in \mathcal{G}\}$.

The second way in which we alleviate the effect of sub-optimal alignments, is by incorporating a minimal differentiability threshold D (default value is 0) that controls when AmpliDiff considers a pair of genomes to be differentiated by an amplicon. Given an amplicon $A = [i : i + L_A]$ and two genomes $g, g' \in \mathcal{G}$ such that $g \not\sim_C g'$ (here $\sim$ is used as an equivalence relation between classes of genomes in $\mathcal{G}$), AmpliDiff considers the genomes to be differentiated by A if and only if $|\{j \in \{i, \ldots, i + L_A - 1\} : g[j] \neq g'[j]\}| > D$. In otherwords, an amplicon can only differentiate between a pair of genomes if they disagree at at least $D + 1$ nucleotide positions, thereby preventing AmpliDiff to differentiate genomes based on unnecessarily introduced gaps due to sub-optimal alignment. Importantly, as input genomes can contain degenerate characters, AmpliDiff only considers two nucleotides $g[i]$ and $g'[i]$ to disagree if there is no overlap in the respective representations of $g[i]$ and $g'[i]$. This serves as a countermeasure against incorrectly calling relative Single Nucleotide Polymorphisms (SNPs) due to uncertainty about the nucleotides occurring in a genome.

Once the set of feasible amplicons, $\mathcal{A}$, has been determined, AmpliDiff continues by calculating the discriminatory power of amplicon candidates. For this, we define $\delta(A; D) := \{(g, g') \subseteq \mathcal{G} : |\{j \in \{i, \ldots, i + L_A - 1\} :$

$g[j] \neq g'[j]\}| > D\}$ as the set of genomes that can be differentiated by amplicon A under minimal differentiability threshold D, such that the *differentiability* of A is defined as $\Delta(A; D) := |\delta(A; D)|$. In the final step of component (ii), AmpliDiff calculates $\delta(A; D)$ and $\Delta(A; D)$ for every amplicon $A \in \mathcal{A}$, which will serve as the starting point for the greedy amplicon selection component.

Step (iii): greedy amplicon selection

In the final component, AmpliDiff finds a near minimal set of amplicons with corresponding primers, to differentiate between as many pairs of genomes with different classes. The methodology employed by AmpliDiff is based on casting the amplicon finding problem as the *Set Cover Problem* (SCP) when looking to differentiate between all pairs of genomes of different classes, or the *Maximum Coverage Problem* (MCP) when looking to find a fixed number of amplicons that differentiate between as many pairs of class disjoint genomes as possible. These coverage problems are NP-hard combinatorial optimization problems [27, 32] that have been well-studied in the field of Operations Research. Instances of these coverage problems are generally given by a set of elements $\mathcal{U}$, often called the universe, a collection of subsets of the universe: $\mathcal{C} \subseteq 2^{\mathcal{U}}$ such that $\cup_{S \in \mathcal{C}} S = \mathcal{U}$ and a positive rational cost function $c : \mathcal{C} \to \mathbb{Q}_{++}$ that assigns a cost to every set in $\mathcal{C}$. In the SCP the objective is to find a minimal cost sub-collection $C^* \subseteq \mathcal{C}$ such that $\cup_{S \in \mathcal{C}^*} = \mathcal{U}$ whereas in the MCP the objective is to find a sub-collection of at most a given size covering as many elements of $\mathcal{U}$ as possible.

To see how these coverage problems related to the problem at hand, we can define our universe $\mathcal{U}$ as the set of all pairs of genomes of different classes: $\mathcal{U} := \{\{g, g'\} \subseteq \mathcal{G} : g \not\sim_C g'\}$. Then, defining an amplicon as the set of genomes it can differentiate and assigning every amplicon a cost of 1, we directly obtain an instance of the SCP, or an instance of the MPC if a maximum number of allowed amplicons is also given. As a consequence, we can use algorithms that have specifically been developed for these types of problems [9, 12, 19, 20, 62], to solve our amplicon finding problem. However, in our case, we also need to consider whether an amplicon can be amplified in enough of the genomes. Specifically, we require that an amplicon $A \in \mathcal{A}$ has both a corresponding forward and reverse primer in at least $\gamma\%$ of the input genomes. The problem with this, however, is that we do not know a priori which amplicons are able to satisfy this condition, which means that this has to be determined for every individual amplicon. As this is computationally infeasible to determine for all candidate amplicons beforehand due to having to check primer interactions as well, we have to resort to a more ad hoc approach where, whenever an amplicon is considered, we determine its amplifiability to see if it exceeds the minimal amplifiability condition. For this reason, we have chosen to base AmpliDiff on the greedy algorithm for the SCP and MCP [12] which iteratively adds the set with the largest number of uncovered elements to the solution. This algorithm has been shown to be an approximation algorithm with approximation factor $(\sum_{i=1}^{|\mathcal{U}|} \frac{1}{i}) \leq (1 + ln(|\mathcal{U}|))$, which means that if we were to determine which amplicons satisfy the minimal amplifiability constraint beforehand, we would obtain an approximately optimal solution for our problem as well.

In the case of AmpliDiff, the greedy amplicon algorithm works as shown in Figure 1**(iii)**. First, the candidate amplicons are sorted based on their differentiability (highest to lowest), and the most differentiable amplicon is picked, with ties broken arbitrarily. Using the primer database built-in component (i), AmpliDiff finds feasible primers flanking this amplicon in all of the input genomes by solving a *Mixed Integer Linear Program* (MILP) that determines if it is possible to find primers that guarantee that the amplicon can be amplified in at least $\gamma\%$ of the input genomes (solving is done through Gurobi [17]). The MILP checks whether it is possible to find enough forward and reverse primers that have approximately the same melting temperature (maximum difference of 5° Celsius) and do not pose a significant risk of having primer-primer interactions. Full details can be found in the Supplementary Material.

When an amplicon does not meet the minimal amplifiability requirement, it is discarded and the next most

differentiable amplicon is considered. Alternatively, when an amplicon meets the requirement, the same MILP is solved again, but now with the objective of finding a minimal set of primer pairs that enable the amplification of this amplicon in at least $\gamma\%$ of the genomes. If γ is smaller than 100%, it is allowed for an amplicon to not be amplifiable (i.e. have binding forward and reverse primers) in all of the reference genomes. In the general primer design problem, this is not much of a problem, but in AmpliDiff the objective is to differentiate between pairs of class disjoint genomes. As a consequence, we can no longer use generic primer design tools, as these consider every reference genome as equivalent. For example, suppose there are three genomes g_1, g_2, g_3 such that $g_1 \sim_C g_2 \nsim_C g_3$, and it is only possible to amplify an amplicon A with $\delta(A; D) = \{(g_1, g_3), (g_2, g_3)\}$ in at most two of these genomes. Generic primer design tools would in this case not care about whether the amplicon could be amplified in g_1 and g_2, g_1 and g_3 or g_2 and g_3. However, in our case, it would be preferred to amplify the amplicon in g_3 and either of the other genomes, as this makes it more likely that this amplicon can provide evidence that the class of genome g_3 is present in a sample.

To deal with this problem we have added an auxiliary step in the greedy amplicon selection where we minimize the number of primer pairs required to amplify the candidate amplicon, which we know is guaranteed to meet the amplifiability requirement, while simultaneously maximizing the number of pairs of genomes, differentiated by the amplicon, in which the amplicon can be amplified (see the Primer feasibility models section in the Supplementary Material). The trade-off between these conflicting objectives is modeled through the trade-off parameter β (default value 0.05) which allows adding additional primer pairs only if they increase the number of class disjoint genome pairs in which the amplicon can be amplified by at least $100 \cdot \beta\%$ of the total differentiability of the amplicon, as long as the amplifiability requirement is met. When the minimal required amplifiability is set to 100%, this trade-off is omitted since the amplicon is already required to be amplifiable in every input genome, and the number of required primer pairs is directly minimized instead.

After solving the primer minimization problem, the amplicon, along with the computed minimal set of primer pairs is added to the solution. Then $\delta(A; D)$ is updated for the remaining amplicons in order to account for the pairs that can be differentiated by the added amplicon. Any amplicons with zero amplifiability are removed and amplicons are sorted again based on their differentiability. The full amplicon selection loop continues until either all pairs of genomes with different classes can be differentiated, a given maximum number of amplicons have been found, or no more feasible amplicons exist, after which the found amplicons and their corresponding primer pairs are returned.

Assessment on SARS-CoV-2

Amplicon generation

We have generated sets of 10 amplicons of different widths and with different minimal amplifiability requirements by using a global reference set of SARS-CoV-2 genomes. This reference set was constructed from all available genomes on the GISAID database [31] up until 18 August 2022, by following the reference set construction procedure detailed in [4] which resulted in a set of 2,749 reference genomes distributed over 1,837 Pango lineages with 1-7 genomes per lineage. These genomes were multiple aligned using MAFFT [29] to obtain the input genomes for AmpliDiff. Then, by running AmpliDiff with $L_A \in \{200, 400\}$, $k = 25$, $L_0 = 50$, $L_\emptyset = 0.1 \cdot L_A$, $D = 1$, $\gamma \in \{90\%, 92.5\%, 95\%, 97.5\%, 99.9\%, 100\%\}$, $\beta \in \{0.05, 0.10\}$ and a maximum primer degeneracy of 1024 we obtained multiple sets of 10 amplicons of widths 200 and 400, requiring a minimal amplifiability of 90%, 92.5%, 95%, 97.5%, 99.9%, and 100% respectively for a total of 12 amplicon sets (β did not affect results) which can be found in Figures 1-6 in the Supplementary Material.

Simulation study

To evaluate how well amplicons and primers selected by AmpliDiff work in the context of abundance estimation, we have created two benchmarking datasets: one with genomes from the Netherlands (constructed on 13 April 2023) and another with genomes from Texas (constructed on 6 March 2023), on which we estimate relative abundances of different lineages using the found amplicons as well as using whole genomes by following the VLQ pipeline in [4]. Both datasets consist of a *reference dataset* with genomes spanning the months of May until October 2022 that is used by Kallisto as references for the abundance estimations, and a *simulation dataset* with genomes from November 2022 which is used to simulate reads from and estimate abundances for (20 simulations per configuration for a total of $(12 + 2) \times 20 = 280$ simulations). The reference genomes were filtered by following the same pre-processing steps as before, with the adjustment that they were not checked for ambiguous nucleotides, and the simulation genomes were not filtered. For the whole genome-based abundance estimations, the reference genomes are directly passed to Kallisto. For the amplicon-based abundance estimations, we further process these reference genomes by extracting amplicons that are considered amplifiable in the respective genomes (see next subsection) and concatenating them with sequences of 200 A nucleotides in between amplicons in order to avoid falsely mapping reads. This way, only the genomic regions corresponding to amplicons that we expect to bind are considered during the abundance estimations, and, as opposed to using the full reference genomes, these amplicon-based representations of the genomes are used as references by Kallisto.

Read simulations were done with ART [21] in default mode for the whole genome-based reads, and in amplicon mode for the amplicon-based reads. Similar to processing the amplicon-based reference set, we have only simulated reads from amplicons that were amplifiable according to the corresponding primers, whereas reads for the whole genome were simulated uniformly over the simulation genomes. For the fragments of length 200 (corresponding to 200 width amplicons), we simulated 2×150bp Illumina HiSeqX TrueSeq sequencing reads with an average fragment length of 200 and a standard deviation of 10 for the non-amplicon-based reads. Similarly, for fragments of length 400 (corresponding to 400 width amplicons), we simulated 2×250bp Illumina MiSeq v3 reads with an average fragment length of 400 and a standard deviation of 10 for the non-amplicon-based reads. Since it is to be expected that reads according to the amplicons have much higher sequencing depth, we set the read depth to 1000 for the amplicon-based read simulations, and to 100 for the whole genome-based simulations. The lineage abundances in the simulated samples correspond to the relative abundances of lineages in the simulation datasets, where the (estimated) relative abundance for a lineage is calculated by aggregating over all sequences assigned to the corresponding lineage and filtering out lineages with an estimated abundance below 0.1%.

The performance of abundance estimations is determined by the *Mean Absolute Prediction Error* which for abundances $\{\phi_l\}_{l \in \mathcal{L}}$ and estimated abundances $\{\hat{\phi}_l\}_{l \in \mathcal{L}}$ is defined as $\text{MAPE} := \frac{1}{|L|} \sum_{l \in \mathcal{L}} |\phi_l - \hat{\phi}_l|$ with $\mathcal{L}$ equal to the set of all lineages occurring in the reference set or the simulation set.

Primer binding and amplicon extraction

In our experiments, we assume that primers only bind to genomes if they match a genome exactly, as this is how primers are generated during component (i) of AmpliDiff. Since it is possible that primers bind to genomes in the presence of nucleotide mismatches in practice, this likely provides a conservative estimate of how amplifiable the amplicons found by AmpliDiff work in practice. However, when looking for nucleotide matches, we do allow up to a predefined number of degenerate nucleotides (10 by default) in the primer binding site which are considered to bind if their respective representations match the corresponding nucleotides in the primers. Although this approach may allow primers to bind to locations in the genome due to the presence of degenerate nucleotides, this should not pose much of a problem if the genomes considered are of high quality.

Moreover, this approach also avoids discarding potential primer binding sites as a consequence of the presence of degenerate nucleotides in the input genomes.

Availablility of data and materials

The code for AmpliDiff, and the simulated reads (.fastq files) used for benchmarking are publicly available (MIT license) at https://github.com/JaspervB-tud/AmpliDiff [55]

Acknowledgements

We gratefully acknowledge all data contributors, i.e., the Authors and their Originating laboratories responsible for obtaining the specimens, and their Submitting laboratories for generating the genetic sequence and metadata and sharing via the GISAID Initiative, on which this research is based.

References

[1] Warish Ahmed, Christopher Staley, Jatinder Sidhu, Michael Sadowsky, and Simon Toze. Amplicon-based profiling of bacteria in raw and secondary treated wastewater from treatment plants across Australia. *Applied microbiology and biotechnology*, 101:1253–1266, 2017.

[2] Daniel Aird, Michael G Ross, Wei-Sheng Chen, Maxwell Danielsson, Timothy Fennell, Carsten Russ, David B Jaffe, Chad Nusbaum, and Andreas Gnirke. Analyzing and minimizing PCR amplification bias in Illumina sequencing libraries. *Genome biology*, 12(2):1–14, 2011.

[3] Najla F Alsiri, Meshal A Alhadhoud, and Shea Palmer. The impact of the COVID-19 on research. *Journal of clinical epidemiology*, 129:124–125, 2021.

[4] Jasmijn A Baaijens, Alessandro Zulli, Isabel M Ott, Ioanna Nika, Mart J van der Lugt, Mary E Petrone, Tara Alpert, Joseph R Fauver, Chaney C Kalinich, Chantal BF Vogels, et al. Lineage abundance estimation for SARS-CoV-2 in wastewater using transcriptome quantification techniques. *Genome biology*, 23(1):1–20, 2022.

[5] Aaron Bivins, Justin Greaves, Robert Fischer, Kwe Claude Yinda, Warish Ahmed, Masaaki Kitajima, Vincent J Munster, and Kyle Bibby. Persistence of SARS-CoV-2 in water and wastewater. *Environmental Science & Technology Letters*, 7(12):937–942, 2020.

[6] Loïc Borcard, Sonja Gempeler, Miguel A Terrazos Miani, Christian Baumann, Carole Grädel, Ronald Dijkman, Franziska Suter-Riniker, Stephen L Leib, Pascal Bittel, Stefan Neuenschwander, et al. Investigating the extent of primer dropout in SARS-CoV-2 genome sequences during the early circulation of delta variants. *Frontiers in Virology*, 2:840952, 2022.

[7] Nicolas L Bray, Harold Pimentel, Páll Melsted, and Lior Pachter. Near-optimal probabilistic rna-seq quantification. *Nature biotechnology*, 34(5):525–527, 2016.

[8] Florian P Breitwieser, Jennifer Lu, and Steven L Salzberg. A review of methods and databases for metagenomic classification and assembly. *Briefings in bioinformatics*, 20(4):1125–1136, 2019.

[9] Alberto Caprara, Paolo Toth, and Matteo Fischetti. Algorithms for the set covering problem. *Annals of Operations Research*, 98(1-4):353–371, 2000.

[10] Phil M Choi, Ben J Tscharke, Erica Donner, Jake W O'Brien, Sharon C Grant, Sarit L Kaserzon, Rachel Mackie, Elissa O'Malley, Nicholas D Crosbie, Kevin V Thomas, et al. Wastewater-based epidemiology biomarkers: past, present and future. *TrAC Trends in Analytical Chemistry*, 105:453–469, 2018.

[11] Yongliang Chu, Silong Sun, Yufen Huang, Qiang Gao, Xuefeng Xie, Peng Wang, Junxia Li, Lifeng Liang, Xiaohong He, Yiqi Jiang, et al. Metagenomic analysis revealed the potential role of gut microbiome in gout. *npj Biofilms and Microbiomes*, 7(1):66, 2021.

[12] Vasek Chvatal. A greedy heuristic for the set-covering problem. *Mathematics of operations research*, 4(3):233–235, 1979.

[13] Victor M Corman, Olfert Landt, Marco Kaiser, Richard Molenkamp, Adam Meijer, Daniel KW Chu, Tobias Bleicker, Sebastian Brünink, Julia Schneider, Marie Luisa Schmidt, et al. Detection of 2019 novel coronavirus (2019-ncov) by real-time RT-PCR. *Eurosurveillance*, 25(3):2000045, 2020.

[14] CW Dieffenbach, TM Lowe, GS Dveksler, et al. General concepts for PCR primer design. *PCR methods appl*, 3(3):S30–S37, 1993.

[15] Askar Gafurov, Andrej Baláž, Fabian Amman, Kristína Boršová, Viktória Čabanová, Boris Klempa, Andreas Bergthaler, Tomáš Vinař, and Broňa Brejová. VirPool: model-based estimation of SARS-CoV-2 variant proportions in wastewater samples. *BMC bioinformatics*, 23(1):1–18, 2022.

[16] Erik Garrison, Jouni Sirén, Adam M Novak, Glenn Hickey, Jordan M Eizenga, Eric T Dawson, William Jones, Shilpa Garg, Charles Markello, Michael F Lin, et al. Variation graph toolkit improves read mapping by representing genetic variation in the reference. *Nature biotechnology*, 36(9):875–879, 2018.

[17] Gurobi Optimization, LLC. Gurobi Optimizer Reference Manual, 2023.

[18] Kenan Hadziavdic, Katrine Lekang, Anders Lanzen, Inge Jonassen, Eric M Thompson, and Christofer Troedsson. Characterization of the 18S rRNA gene for designing universal eukaryote specific primers. *PloS one*, 9(2):e87624, 2014.

[19] Refael Hassin and Asaf Levin. A better-than-greedy approximation algorithm for the minimum set cover problem. *SIAM Journal on Computing*, 35(1):189–200, 2005.

[20] Dorit S Hochbaum. Approximation algorithms for the set covering and vertex cover problems. *SIAM Journal on computing*, 11(3):555–556, 1982.

[21] Weichun Huang, Leping Li, Jason R Myers, and Gabor T Marth. ART: a next-generation sequencing read simulator. *Bioinformatics*, 28(4):593–594, 2012.

[22] David A Hysom, Pejman Naraghi-Arani, Maher Elsheikh, A Celena Carrillo, Peter L Williams, and Shea N Gardner. Skip the alignment: degenerate, multiplex primer and probe design using K-mer matching instead of alignments. *PloS one*, 7(4):e34560, 2012.

[23] Zamin Iqbal, Mario Caccamo, Isaac Turner, Paul Flicek, and Gil McVean. De novo assembly and genotyping of variants using colored de Bruijn graphs. *Nature genetics*, 44(2):226–232, 2012.

[24] Ray Izquierdo-Lara, Goffe Elsinga, Leo Heijnen, Bas B Oude Munnink, Claudia ME Schapendonk, David Nieuwenhuijse, Matthijs Kon, Lu Lu, Frank M Aarestrup, Samantha Lycett, et al. Monitoring SARS-CoV-2 circulation and diversity through community wastewater sequencing, the Netherlands and Belgium. *Emerging infectious diseases*, 27(5):1405, 2021.

[25] Andrew D Johnson. An extended IUPAC nomenclature code for polymorphic nucleic acids. *Bioinformatics*, 26(10):1386–1389, 2010.

[26] Winfried Just. Computational complexity of multiple sequence alignment with SP-score. *Journal of computational biology*, 8(6):615–623, 2001.

[27] Richard M Karp. Reducibility among combinatorial problems. In *Complexity of computer computations*, pages 85–103. Springer, 1972.

[28] Smruthi Karthikeyan, Joshua I Levy, Peter De Hoff, Greg Humphrey, Amanda Birmingham, Kristen Jepsen, Sawyer Farmer, Helena M Tubb, Tommy Valles, Caitlin E Tribelhorn, et al. Wastewater sequencing reveals early cryptic SARS-CoV-2 variant transmission. *Nature*, 609(7925):101–108, 2022.

[29] Kazutaka Katoh, Kazuharu Misawa, Kei-ichi Kuma, and Takashi Miyata. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic acids research*, 30(14):3059–3066, 2002.

[30] Tunc Kayikcioglu, Jasmine Amirzadegan, Hugh Rand, Bereket Tesfaldet, Ruth E Timme, and James B Pettengill. Performance of methods for SARS-CoV-2 variant detection and abundance estimation within mixed population samples. *PeerJ*, 11:e14596, 2023.

[31] Shruti Khare, Céline Gurry, Lucas Freitas, Mark B Schultz, Gunter Bach, Amadou Diallo, Nancy Akite, Joses Ho, Raphael TC Lee, Winston Yeo, et al. GISAID's role in pandemic response. *China CDC Weekly*, 3(49):1049, 2021.

[32] Samir Khuller, Anna Moss, and Joseph Seffi Naor. The budgeted maximum coverage problem. *Information processing letters*, 70(1):39–45, 1999.

[33] Amy E Kirby, Maroya Spalding Walters, Wiley C Jennings, Rebecca Fugitt, Nathan LaCross, Mia Mattioli, Zachary A Marsh, Virginia A Roberts, Jeffrey W Mercante, Jonathan Yoder, et al. Using wastewater surveillance data to support the COVID-19 response—United States, 2020–2021. *Morbidity and Mortality Weekly Report*, 70(36):1242, 2021.

[34] Christoph Kreer, Matthias Döring, Nathalie Lehnen, Meryem S Ercanoglu, Lutz Gieselmann, Domnica Luca, Kanika Jain, Philipp Schommers, Nico Pfeifer, and Florian Klein. openPrimeR for multiplex amplification of highly diverse templates. *Journal of Immunological Methods*, 480:112752, 2020.

[35] Arnold W Lambisia, Khadija S Mohammed, Timothy O Makori, Leonard Ndwiga, Maureen W Mburu, John M Morobe, Edidah O Moraa, Jennifer Musyoki, Nickson Murunga, Jane N Mwangi, et al. Optimization of the SARS-CoV-2 ARTIC network V4 primers and whole genome sequencing protocol. *Frontiers in Medicine*, 9:836728, 2022.

[36] Trang Thi Bich Le, Tamilarasy Vasanthakumaran, Hau Nguyen Thi Hien, I-Chun Hung, Mai Ngoc Luu, Zeeshan Ali Khan, Nguyen Thanh An, Van Phu Tran, Wei Jun Lee, Jeza Muhamad Abdul Aziz, et al. SARS-CoV-2 Omicron and its current known unknowns: A narrative review. *Reviews in Medical Virology*, 33(1):e2398, 2023.

[37] Esther G Lou, Nicolae Sapoval, Camille McCall, Lauren Bauhs, Russell Carlson-Stadler, Prashant Kalvapalle, Yanlai Lai, Kyle Palmer, Ryker Penn, Whitney Rich, et al. Direct comparison of RT-ddPCR and targeted amplicon sequencing for SARS-CoV-2 mutation monitoring in wastewater. *Science of The Total Environment*, 833:155059, 2022.

[38] Priyaranjan Mishra, Nilam J Tulsani, Subhash J Jakhesara, Nishant A Dafale, Niteen V Patil, Hemant J Purohit, Prakash G Koringa, and Chaitanya G Joshi. Exploring the eukaryotic diversity in rumen of indian camel (camelus dromedarius) using 18S rRNA amplicon sequencing. *Archives of Microbiology*, 202:1861–1872, 2020.

[39] Laura Moreno-Mesonero, María Antonia Ferrús, and Yolanda Moreno. Determination of the bacterial microbiome of free-living amoebae isolated from wastewater by 16S rRNA amplicon-based sequencing. *Environmental Research*, 190:109987, 2020.

[40] Artem Nemudryi, Anna Nemudraia, Tanner Wiegand, Kevin Surya, Murat Buyukyoruk, Calvin Cicha, Karl K Vanderwood, Royce Wilkinson, and Blake Wiedenheft. Temporal detection and phylogenetic assessment of SARS-CoV-2 in municipal wastewater. *Cell Reports Medicine*, 1(6):100098, 2020.

[41] Jordan Peccia, Alessandro Zulli, Doug E Brackney, Nathan D Grubaugh, Edward H Kaplan, Arnau Casanovas-Massana, Albert I Ko, Amyn A Malik, Dennis Wang, Mike Wang, et al. Measurement of SARS-CoV-2 RNA in wastewater tracks community infection dynamics. *Nature biotechnology*, 38(10):1164–1167, 2020.

[42] Danielle Peterson, Kevin S Bonham, Sophie Rowland, Cassandra W Pattanayak, Resonance Consortium, and Vanja Klepac-Ceraj. Comparative analysis of 16s rRNA gene and metagenome sequencing in pediatric gut microbiomes. *Frontiers in microbiology*, 12:670336, 2021.

[43] David Polo, Marcos Quintela-Baluja, Alexander Corbishley, Davey L Jones, Andrew C Singer, David W Graham, and Jesús L Romalde. Making waves: Wastewater-based epidemiology for COVID-19–approaches and challenges for surveillance and prediction. *Water research*, 186:116404, 2020.

[44] Joshua Quick, Nathan D Grubaugh, Steven T Pullan, Ingra M Claro, Andrew D Smith, Karthik Gangavarapu, Glenn Oliveira, Refugio Robles-Sikisaka, Thomas F Rogers, Nathan A Beutler, et al. Multiplex PCR method for MinION and Illumina sequencing of Zika and other virus genomes directly from clinical samples. *Nature protocols*, 12(6):1261–1276, 2017.

[45] Andrew Rambaut, Edward C Holmes, Áine O'Toole, Verity Hill, John T McCrone, Christopher Ruis, Louis du Plessis, and Oliver G Pybus. A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic epidemiology. *Nature microbiology*, 5(11):1403–1407, 2020.

[46] Philipp Rausch, Malte Rühlemann, Britt M Hermes, Shauni Doms, Tal Dagan, Katja Dierking, Hanna Domin, Sebastian Fraune, Jakob Von Frieling, Ute Hentschel, et al. Comparative analysis of amplicon and metagenomic sequencing methods reveals key features in the evolution of animal metaorganisms. *Microbiome*, 7(1):1–19, 2019.

[47] Géraldine Rios, Caroline Lacoux, Vianney Leclercq, Anna Diamant, Kévin Lebrigand, Adèle Lazuka, Emmanuel Soyeux, Sébastien Lacroix, Julien Fassy, Aurélie Couesnon, et al. Monitoring SARS-CoV-2 variants alterations in Nice neighborhoods by wastewater nanopore sequencing. *The Lancet Regional Health–Europe*, 10, 2021.

[48] Nicolae Sapoval, Yunxi Liu, Esther G Lou, Loren Hopkins, Katherine B Ensor, Rebecca Schneider, Lauren B Stadler, and Todd J Treangen. Enabling accurate and early detection of recently emerged SARS-CoV-2 variants of concern in wastewater. *Nature Communications*, 14(1):2834, 2023.

[49] Conrad L Schoch, Keith A Seifert, Sabine Huhndorf, Vincent Robert, John L Spouge, C André Levesque, Wen Chen, Fungal Barcoding Consortium, Fungal Barcoding Consortium Author List, Elena Bolchacova, et al. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for fungi. *Proceedings of the national academy of Sciences*, 109(16):6241–6246, 2012.

[50] Davida S Smyth, Monica Trujillo, Devon A Gregory, Kristen Cheung, Anna Gao, Maddie Graham, Yue Guan, Caitlyn Guldenpfennig, Irene Hoxie, Sherin Kannoly, et al. Tracking cryptic SARS-CoV-2 lineages detected in NYC wastewater. *Nature communications*, 13(1):1–9, 2022.

[51] Christen Rune Stensvold, Marianne Lebbad, Anette Hansen, Jessica Beser, Salem Belkessa, Lee O'Brien Andersen, and C Graham Clark. Differentiation of Blastocystis and parasitic archamoebids encountered in untreated wastewater samples by amplicon-based next-generation sequencing. *Parasite epidemiology and control*, 9:e00131, 2020.

[52] D Lee Taylor, William A Walters, Niall J Lennon, James Bochicchio, Andrew Krohn, J Gregory Caporaso, and Taina Pennanen. Accurate estimation of fungal diversity and abundance through improved lineage-specific primers optimized for Illumina amplicon sequencing. *Applied and environmental microbiology*, 82(24):7217–7226, 2016.

[53] Julie D Thompson, Toby J Gibson, and Des G Higgins. Multiple sequence alignment using ClustalW and ClustalX. *Current protocols in bioinformatics*, (1):2–3, 2003.

[54] Andreas Untergasser, Ioana Cutcutache, Triinu Koressaar, Jian Ye, Brant C Faircloth, Maido Remm, and Steven G Rozen. Primer3—new capabilities and interfaces. *Nucleic acids research*, 40(15):e115–e115, 2012.

[55] Jasper van Bemmelen. Amplidiff. https://github.com/JaspervB-tud/AmpliDiff, 2022.

[56] Holger Volkmann, Thomas Schwartz, Petra Bischoff, Silke Kirchen, and Ursula Obst. Detection of clinically relevant antibiotic-resistance genes in municipal wastewater using real-time PCR (TaqMan). *Journal of microbiological methods*, 56(2):277–286, 2004.

[57] Yong Wang and Pei-Yuan Qian. Conservative fragments in bacterial 16S rRNA genes and primer design for 16S ribosomal DNA amplicons in metagenomic studies. *PloS one*, 4(10):e7401, 2009.

[58] Amy Xiao, Fuqing Wu, Mary Bushman, Jianbo Zhang, Maxim Imakaev, Peter R Chai, Claire Duvallet, Noriko Endo, Timothy B Erickson, Federica Armas, et al. Metrics to relate COVID-19 wastewater data to clinical testing dynamics. *Water research*, 212:118070, 2022.

[59] Pablo Yarza, Pelin Yilmaz, Elmar Pruesse, Frank Oliver Glöckner, Wolfgang Ludwig, Karl-Heinz Schleifer, William B Whitman, Jean Euzéby, Rudolf Amann, and Ramon Rosselló-Móra. Uniting the classification of cultured and uncultured bacteria and archaea using 16S rRNA gene sequences. *Nature Reviews Microbiology*, 12(9):635–645, 2014.

[60] Jian Ye, George Coulouris, Irena Zaretskaya, Ioana Cutcutache, Steve Rozen, and Thomas L Madden. Primer-BLAST: a tool to design target-specific primers for polymerase chain reaction. *BMC bioinformatics*, 13(1):1–11, 2012.

[61] Alexander T Yu, Bridgette Hughes, Marlene K Wolfe, Tomas Leon, Dorothea Duong, Angela Rabe, Lauren C Kennedy, Sindhu Ravuri, Bradley J White, Krista R Wigginton, et al. Estimating relative abundance of 2 SARS-CoV-2 variants through wastewater surveillance at 2 large metropolitan sites, United States. *Emerging Infectious Diseases*, 28(5):940–947, 2022.

[62] Xin-Yuan Zhang, Jun Zhang, Yue-Jiao Gong, Zhi-Hui Zhan, Wei-Neng Chen, and Yun Li. Kuhn–Munkres parallel genetic algorithm for the set cover problem and its application to large-scale wireless sensor networks. *IEEE Transactions on Evolutionary Computation*, 20(5):695–710, 2015.