Spatial patterns of within-thalamus anatomical connectivity vary across the cortical hierarchy in humans and macaques

Amber M. Howell[a,b,c,1], Shaun Warrington[f], Clara Fonteneau[a,b], Youngsun Cho[a,b], Stamatios N. Sotiropoulos[f,g,h], John D. Murray[a,b,c,e,2], and Alan Anticevic[a,b,c,d,2]

[a]Department of Psychiatry, Yale University School of Medicine, New Haven, CT 06511, USA
[b]Division of Neurocognition, Neurocomputation, & Neurogenetics (N3), Yale University School of Medicine, New Haven, Connecticut, 06511, USA
[c]Interdepartmental Neuroscience Program, Yale University, New Haven, Connecticut, 06511, USA
[d]Department of Psychology, Yale University, New Haven, Connecticut, 06511, USA
[e]Physics, Yale University, New Haven, Connecticut, 06511, USA
[f]Sir Peter Mansfield Imaging Centre, School of Medicine, University of Nottingham, Nottingham, UK
[g]Wellcome Centre for Integrative Neuroimaging, University of Oxford, Oxford, UK
[h]National Institute for Health Research (NIHR) Nottingham Biomedical Research Centre, Queens Medical Centre, Nottingham, UK

The thalamus is composed of functionally and structurally distinct nuclei. Previous studies have indicated that certain cortical areas may project across multiple thalamic nuclei, potentially allowing them to modulate distributed information flow. However, there is a lack of quantitative investigations into anatomical connectivity patterns within the thalamus. Consequently, it remains unknown if cortical areas exhibit differences in the spread of their thalamic connectivity patterns. To address this knowledge gap, we used diffusion magnetic resonance imaging (dMRI) to compute brain-wide probabilistic tractography using data from 828 healthy adults collected by the Human Connectome Project. Additionally, we examined post-mortem data from six macaque monkeys to assess cross-species generalizability. To measure the spatial spread of anatomical connectivity patterns within the thalamus, we developed an innovative framework that quantifies the spatial properties of each cortical area's within-thalamus connectivity patterns. We then leveraged resting-state, myelin, and human neural gene expression data to test if the spread of within-thalamus connectivity patterns varied along the cortical hierarchy. These results revealed two broad cortico-thalamic tractography motifs: 1) a sensorimotor cortical motif characterized by focal thalamic connections targeting posterolateral thalamus, which potentially supports fast, feed-forward information flow; and 2) an associative cortical motif characterized by diffuse thalamic connections targeting anteromedial thalamus, which potentially supports slower, feed-back information flow. These results were consistent among human subjects and were also observed in macaques, indicating generalizability. In summary, these findings demonstrate that sensorimotor and association cortical areas exhibit distinct spatial connectivity patterns within the thalamus, which may support functionally-distinct cortico-thalamic information transmission.

thalamus | spatial topography | thalamo-cortical | diffusion MRI | anatomical connectivity | tractography

Correspondence: *amber.howell@yale.edu*

Introduction

Mapping the anatomical connections of the brain is a fundamental goal in neuroscience, as these pathways tether different brain areas together and impose constraints on their functional interactions. The thalamus is a central subcortical brain structure that is extensively connected to the entire cortex through long-range white matter fiber bundles (1). These bundles form parallel cortico-thalamic circuits, which enable the thalamus to relay, sustain, and coordinate information across the entire cortex (2–4). Notably, the thalamus lacks reciprocal excitatory connections (5). Thus, sensorimotor and associative neural computations involving the thalamus heavily rely on its long-range inputs from and outputs to the cortex (6). Mapping these long-range connections can provide critical insight into the role of the thalamus in shaping cortical information flow and the neural basis of higher-order cognitive functions, which are critically reliant on the long-range interactions between the thalamus and cortex in vertebrates (2, 7–12).

Research on the anatomy of cortico-thalamic connectivity date back to the early 19th century, yet we still lack a comprehensive understanding of the organizing principles of these connections, particularly in humans (see 13 and 14 for review). The traditional view of the thalamus has largely been informed by its histologically-defined nuclear structure (6). This view has also been supported by evidence showing that certain cortical areas project to individual thalamic nuclei. However, some cortical areas exhibit widespread connections across the thalamus, projecting to multiple thalamic nuclei (15–21). One proposed role for such cross-nuclei targeting is to allow for spatial overlap of connections from different cortical areas, thereby facilitating information integration within the thalamus (22–24). However, knowledge of how within-thalamus connectivity patterns vary across cortical areas, especially in humans, remains incomplete.

Primate studies investigating cortico-thalamic circuitry have primarily relied on anatomical tracer data in monkeys (e.g., 18–20, 25–27). However, such invasive studies cannot be replicated in humans. Fortunately, advancements in magnetic resonance imaging (MRI) have enabled the examination of white matter pathways *in vivo* using diffusion MRI (dMRI), which measures the diffusion properties of water molecules in brain tissue (28–30). These properties are then

used by tractography algorithms to reconstruct white matter pathways, known as streamlines (31–34).

State-of-the-art tractography techniques can now map streamlines at high spatial resolutions to reveal connectivity patterns within the thalamus (15, 16, 35–37). Tractography studies have unveiled a diverse array of cortico-thalamic circuits. These circuits are characterized by distinct origins, targets, strengths, and microstructural profiles, which are optimized for specialized roles in information transmission (15, 37–40). Emerging evidence suggests that certain cortical areas may exhibit more widespread connectivity patterns within the thalamus (37), which may grant them privileged access for integrating or modulating whole-brain functional interactions (41). However, it is uncertain whether the spread of within-thalamus connectivity patterns systematically varies across cortical areas in humans and what implications such variation may have for information processing within cortico-thalamic systems.

Furthermore, there are few studies that have directly compared cortico-thalamic anatomical circuitry between humans and non-human primates using tractography methods (e.g., 40, 42). Such studies form a bridge to the existing macaque tract tracing literature. They also provide validation for human dMRI findings, as macaque dMRI can be collected at much higher resolutions, without confounds such as motion artifacts (43).

The aim of this study was to investigate the spatial patterns of anatomical connectivity within the thalamus in both humans and non-human primates. To this end, we leveraged probabilistic tractography derived from 3T diffusion data for 828 healthy human adults from the Human Connectome Project (HCP) and 7T diffusion data from six postmortem macaque monkeys. We then developed an innovative approach to quantify the spatial properties of anatomical connectivity patterns. We next examined intercortical differences in the spread of within-thalamus connectivity patterns. These findings provide convergent evidence using resting-state, myelin, and human neural gene expression data to demonstrate distinct patterns of thalamic connectivity between sensory and associative cortical areas in humans. Additionally, we show that these differences are generalized in macaques. Overall, our findings highlight that sensorimotor and association cortical areas exhibit distinct anatomical connectivity patterns within the thalamus, which may underlie functionally distinct cortico-thalamic computations.

Results

Cortical areas differ in the spread of their thalamic connectivity patterns. We first examined whether there are systematic differences in the spatial spread of anatomical connectivity patterns within the thalamus across cortical areas. To achieve this, we developed a thresholding framework to investigate the spatial properties of brain connectivity patterns (**Fig. 1A**). Briefly, the framework involved iteratively excluding a percentage of thalamic voxels based on their streamline count, which represents the likelihood that a voxel is connected to the given cortical area. For each corti-

cal parcel, we then calculated the average pairwise Euclidean distance (ED) between these 'surviving' thalamic voxels. To obtain a single measure for each cortical area, each cortical area's ED values across all thresholds were used as input into principal components analysis (PCA), yielding a loading value for the first principal component (ED_{pc1} loadings; see SI Appendix for additional methodological details). ED_{pc1} loadings serve as an index of the spatial spread of thalamic connectivity patterns for each cortical area.

We applied the thresholding framework to dMRI-derived probabilistic tractography data from healthy adults from the Human Connectome Project (n=828; see SI Appendix **Fig. S1** for preprocessing steps). Here, we parcellated dense (grey-ordinate to grey-ordinate) streamline count connectivity data along one dimension to generate a cortico-thalamic 'connectome'. This connectome represents the anatomical connectivity between 360 cortical parcels and all thalamic voxels. Only ipsilateral thalamic voxels were considered, unless otherwise specified in supplementary analyses. Each cortical parcel exhibited a distinct thalamic connectivity pattern, as exemplified by motor area 1 (M1; top magenta panel) and dorsolateral prefrontal cortex (DLPFC; bottom cyan panel) (**Fig. 1B**). Thalamic voxels displayed varying probabilities of connectivity to each cortical area, where warmer colors indicated a higher likelihood of connectivity relative to cooler colors.

For each cortical parcel, we iteratively calculated the average pairwise ED between between thalamic voxels, after progressively excluding a percentage (0 to 99%) of thalamic voxels with the lowest streamline counts (**Fig. S2** shows the surviving thalamic voxels for a subset of thresholds). This process generated a matrix of ED values for each cortical parcel and threshold combination (360x100) (**Fig. 1C**).

We then conducted PCA using the ED matrix as input. The first principal component accounted for almost all of the variation of ED across cortical parcels (92%) (**Fig. 1D**). Cortical parcels with lower ED_{pc1} loadings included bilateral anterior cingulate areas (such as areas 25 and 24), bilateral precuneus, and right temporal cortex. In contrast, cortical parcels with higher ED_{pc1} loadings included bilateral visual areas, right somatosensory cortex, and left entorhinal cortex (refer to **Fig. S3** for visualizations of their respective thalamic connectivity patterns).

We then compared ED_{pc1} loadings with alternative measures of connectivity spread. Based on the qualitative observation that some cortical parcels had ED values that remained higher across conservative thresholds, the standard deviation of ED across thresholds was calculated for each cortical parcel, denoted as ED_σ (**Fig. 1E**). Remarkably, the ED_{pc1} and ED_σ measures exhibited an almost perfect correlation ($r_s = 0.99$; $p_{sa} < 0.001$), with statistical significance determined using spatial-auto-correlation (SA) preserving surrogate maps (see SI Appendix for further details) (44). The high correlation between ED_{pc1} and ED_σ was consistently observed across subjects (mean = 0.99, max = 0.99, min = 0.94) (**Fig. 1F**).

Moreover, we explored other alternative measures to cap-

A Thresholded Metric Reduction Workflow (TMRW)

DWI-derived anatomical connectivity → Surviving thalamic voxels for each cortical parcel → Parcel by threshold ED matrix → ED_{pc1} map

Rowwise threshold · Euclidean distance (ED) calculation · Perform PCA

Cortical parcel | Thalamic voxel
Streamline count (SC)
0 25 50 75 99 % voxels excluded
0 — 99 ED
Loading

B Exemplar anatomical connectivity patterns
M1
DLPFC
Thalamus
SC 0 — 5e-07

C 100 ED values for each cortical parcel
ED 8 5 2
— R. M1
— R. DLPFC
0 — 99
Threshold (% voxels excluded)

Perform PCA
Summary operation
Calculate ED σ for each parcel

D ED_{pc1}: 92% variance
Diffuse --> Focal
-0.28 | 1.29
Loading

E ED_σ
0.12 | 1.34
Standard deviation (σ)

$r_s = 0.99$

F
r_s 1.00 0.95
• subject
Correlation between ED_{pc1} and ED_σ

G Correlation between ED_{pc1} and ED
r_s 1 0 -1
0 — 99
Threshold (% voxels excluded)

Fig. 1. Workflow for quantifying the spread of each cortical area's thalamic anatomical connectivity pattern using Euclidean distance (ED). **(A)** Schematic overview of the thresholding and ED calculation framework applied to group-level and individual-level human probabilistic tractography data (n = 828). ED was used to measure the spread of each cortical parcel's anatomical connectivity pattern within ipsilateral thalamus (see **Fig. S8** for bilateral calculation). **(B)** Thalamic connectivity patterns for right motor area 1 (M1) and dorsolateral prefrontal cortex (DLPFC). M1 and DLPFC illustrate focal and diffuse thalamic connectivity patterns, respectively. **(C)** ED matrix depicting pairwise distances calculated between surviving thalamic voxels for 100 thresholds. Cortical parcels with more diffuse thalamic connectivity patterns exhibited higher ED values across thresholds (e.g., DLPFC). **(D)** Cortical ED_{pc1} loading map obtained through principal component analysis (PCA) of the ED matrix. The first PC accounted for 92% of the variance. **(E)** Cortical ED_σ map representing the standard deviation of ED across 100 thresholds for each cortical parcel. ED_σ values were highly correlated with ED_{pc1} loadings (r_s = 0.99; p_{sa} < 0.001), with p-values estimated using spatial-autocorrelation (sa) preserving surrogate brain maps (44). r_s : Spearman rho. **(F)** Correlations between ED_{pc1} and ED_σ were highly consistent across subjects. **(G)** ED_{pc1} loadings and ED values negatively correlated at more conservative thresholds, indicating that higher ED_{pc1} loadings correspond to more focal thalamic connectivity patterns.

ture thalamic connectivity spread (see **Fig. S4**). However, based on the superior agreement with ED calculated at more conservative thresholds, we selected ED_{pc1} loadings for further analysis. These alternative measures largely replicated the main findings of this study (**Fig. S4**).

We next examined whether the ED_{pc1} measure can distinguish between focal and diffuse thalamic connectivity patterns. For this purpose, we compared ED_{pc1} loadings with the ED values calculated at individual thresholds (**Fig. 1G**). We observed a strong negative correlation between ED_{pc1} and ED at more conservative thresholds (e.g., 78-99%; **Fig. S5**). Additionally, we investigated the relationship between ED_{pc1} loadings and the mean and standard deviation of ED across different threshold ranges. Similar to the correlation observed with individual thresholds, these measures also displayed a high correlation with ED_{pc1} loadings at more conservative thresholds (**Fig. S6**). In essence, cortical parcels exhibiting focal thalamic connectivity patterns were characterized by higher ED_{pc1} loadings compared to those with diffuse thalamic connectivity patterns.

We proceeded to examine the properties of bilateral (**Fig. S8**) and contralateral (**Fig. S9**) thalamic connectivity patterns. Bilateral ED_{pc1} loadings exhibited a strong correlation with ipsilateral loadings (r_s = 0.91). Conversely, contralateral loadings demonstrated a weaker, yet still significant, correlation with ipsilateral loadings (r_s = 0.57). Furthermore, bilateral ED_{pc1} loadings successfully differentiated cortical parcels with unilateral and bilateral thalamic connectivity patterns (**Fig. S8**), indicating that the ED_{pc1} measure can distinguish between focal and diffuse thalamic connectivity patterns within and across hemispheres.

Finally, we conducted extensive control analyses to en-

sure the robustness of these findings. Firstly, we found that ED_{pc1} loadings were not associated with intersubject variation in motion (**Fig. S14**) or volumes of gray or white matter in the cortex, subcortex, or thalamus (**Fig. S15**). Additionally, intercortical variation in ED_{pc1} did not correspond with average streamline count (Mean SC) or streamline length (**Fig. S10**), anatomical overlap with thalamic fractional anisotropy and mean diffusivity values (**Fig. S11**), or cortical geometry, distortion, or bias (**Fig. S12**). ED_{pc1} loadings also remained largely unchanged after accounting for volume differences between the left and right thalamus (**Fig. S13**) and intercortical variation of cortical curvature (**Fig. S16**).

Lastly, we also calculated isotropy, or evenness of spread, to capture thalamic connectivity pattern shape (**Fig. S7**), which also did not correspond highly with ED_{pc1} loadings.

Sensory cortical parcels have more focal thalamic connectivity patterns relative to association cortical parcels. Animal studies have provided evidence that cortico-thalamic anatomical connectivity is hierarchically-organized, with a progression from sensory to association cortical areas (46–49). Building upon this, we hypothesized that the spread of within-thalamus connectivity patterns would also vary along the cortical hierarchy in humans. To test this hypothesis, we compared ED_{pc1} loadings between sensory and association cortical parcels using both network and gradient approaches (**Fig. 2**). Overall, these results show that sensory cortical parcels have more focal thalamic connectivity patterns relative to association cortical parcels.

First, we assigned each cortical parcel to specific networks based on the work of Ji et al. (45), which categorized

Fig. 2. Differences in thalamic connectivity patterns between sensory and association cortical parcels. **(A)** Resting-state functional connectivity networks identified by Ji et al. (45). The 360 cortical parcels were assigned to twelve functional networks, including eight association and four sensory networks. **(B)** ED_{pc1} loadings for each network, representing the average ED_{pc1} loading within the network for each subject. Barplots show the mean and standard error. **(C)** Sensory networks exhibited significantly higher ED_{pc1} loadings compared to association networks (two-sided Wilcoxon signed-rank test; *** p < 0.001). **(D)** Cortical myelin map calculated by averaging T1w/T2w values across subjects. **(E)** Cortical principal functional gradient loading ($RSFC_{pc1}$) map derived from PCA on cortico-cortical resting-state functional connectivity data. Sensory cortical parcels displayed higher T1w/T2w values and $RSFC_{pc1}$ loadings compared to association cortical parcels. **(F,G)** Correlations between ED_{pc1} loadings and T1w/T2w values, as well as $RSFC_{pc1}$ loadings, across the cortex. **(H,I)** On average, a moderate relationship was observed between ED_{pc1} loadings and T1w/T2w values, as well as $RSFC_{pc1}$ loadings, across subjects.

the 360 cortical parcels into twelve functional resting-state networks, encompassing sensorimotor (referred to as 'sensory') and higher-order associative (referred to as 'association') networks (**Fig. 2A**). ED_{pc1} loadings exhibited variations within and across these networks (**Fig. 2B**). Notably, sensory cortical networks demonstrated higher ED_{pc1} loadings (median=0.93) compared to association cortical parcels (median=0.68) (Wilcoxon signed-rank test: W=0, p=3.76e-137) (**Fig. 2C**).

Recently, novel methods have emerged to characterize brain organization along gradients, which exhibit smooth spatial transitions in brain features (50). Many of these

gradients demonstrate hierarchical variations from sensory to association cortical areas, including the T1w/T2w ratio, a proxy measure of cortical myelin (51), and the principal resting-state functional gradient ($RSFC_{pc1}$), which is derived from cortico-cortical resting-state blood-oxygen-level-dependent (BOLD) functional connectivity (52). T1w/T2w values and $RSFC_{pc1}$ loadings are higher in sensory unimodal cortical parcels compared to associative transmodal cortical parcels (**Fig. 2D,E**). Additionally, they significantly correlate with one another ($r_s = 0.51$; **Fig. S17**A).

ED_{pc1} loadings demonstrated a strong positive correlation with both T1w/T2w values ($r_s = 0.48$, $p_{sa} = 0.026$) (**Fig. 2F**) and $RSFC_{pc1}$ loadings ($r_s = 0.59$, $p_{sa} = 0.001$) across cortex (**Fig. 2G**). On average, the correlations between each subject's ED_{pc1} loadings and T1w/T2w values (mean: 0.35, median: 0.36, SEM = 0.003, STD = 0.09) and $RSFC_{pc1}$ loading (mean: 0.45, median: 0.45, SEM = 0.003, STD = 0.09) were moderate (**Fig. 2H,(I)**. However, there were instances where a weak relationship was observed between these cortical maps. This discrepancy may be attributed to the presence of biologically implausible connections originating from certain cortical parcels (**Fig. S18**), which was associated with a weaker correlation between ED_{pc1} loadings and T1w/T2w values, as well as $RSFC_{pc1}$ loadings, across subjects (**Fig. S19**).

The finding that ED_{pc1} loadings significantly correlated with T1w/T2w values, as well as $RSFC_{pc1}$ loadings, was replicated using bilateral thalamic connectivity patterns (**Fig. S8**), but not contralateral patterns (**Fig. S9**). This finding was replicated using an alternate tractography seeding strategy (**Fig. S20**), dense connectivity data (**Fig. S21**), and structurally and functionally defined thalamic masks (**Fig. S22**). Lastly, at more conservative thresholds, ED calculated at individual thresholds (**Fig. S5**) and the mean and standard deviation of ED calculated for a range of thresholds (**Fig. S6**) demonstrated variations between sensory and association cortical parcels.

To determine specificity, we tested the correspondence between ED_{pc1} loadings and the anteroposterior cortical gradient or the secondary principal functional gradient, which reflects functional specialization along a sensory-association-motor cortical axis (52). Neither cortical gradient significantly correlated with ED_{pc1} loadings (**Fig. S23**).

Cortical parcels with focal thalamic connectivity patterns preferentially couple to sensorimotor thalamus. Thalamic nuclei are known to play a role in various sensorimotor and associative cognitive processes (9, 54). We hypothesized that cortical parcels with diffuse thalamic connectivity patterns would anatomically couple with associative thalamic nuclei, while those with focal connectivity patterns would anatomically couple with sensorimotor thalamic nuclei. To test this hypothesis, we compared ED_{pc1} loadings to quantitative indices that capture anatomical coupling across the thalamic subregions. These findings demonstrate that cortical parcels with more focal thalamic connectivity patterns primarily couple with thalamic subregions associated with sensorimotor processes, predominantly located in the lateral

Fig. 3. Mapping the spread of cortico-thalamic connectivity patterns onto anatomical coupling with the anteromedial-posterolateral thalamic gradient. **(A)** Axial view displaying 28 histologically-defined thalamic nuclei from the Morel thalamic atlas (53). **(B)** An anatomical connectivity matrix between each cortical parcel and each thalamic nucleus. The y-axis is sorted such that cortical parcels with more focal thalamic connectivity patterns (higher ED_{pc1} loadings) are positioned near the top. **(C-D)** The correlation between ED_{pc1} loadings and Mean SC values was calculated for each thalamic nucleus in each subject. Higher values indicate thalamic nuclei that preferentially couple with cortical parcels exhibiting focal thalamic connectivity patterns. **(C)** Cortical parcels with focal thalamic connectivity patterns preferentially coupled with posterior nuclei, followed by lateral, anterior, and medial nuclei (Friedman test; *** p = 0.001 for Nemenyi post-hoc tests). Each dot represents the averaged r_s value of a subject across thalamic nuclei. **(D)** Cortical parcels with focal thalamic connectivity patterns preferentially coupled with first-order thalamic nuclei (two-sided Wilcoxon signed-rank test). **(E)** The overlap between each cortical parcel's anatomical connectivity along the anteromedial-posterolateral thalamic spatial gradient ($PL\text{-}AM_t$) was calculated for each cortical parcel. Across cortex, higher values denote cortical parcels that preferentially couple with anteromedial thalamus ($PL\text{-}AM_c$). **(F)** Cortical parcels with higher ED_{pc1} loadings preferentially coupled with the posterolateral thalamus, resulting in lower $PL\text{-}AM_c$ rho values. **(G)** On average, a moderate relationship was observed between ED_{pc1} loadings and $PL\text{-}AM_c$ values across subjects.

and posterior thalamus, referred to as 'first-order' nuclei. Interestingly, cortical parcels with diffuse thalamic connectivity patterns only preferentially coupled with a few nuclei, such as the mediodorsal nucleus (**Fig. S24**). In contrast, 'higher-order' medial and anterior nuclei exhibited more balanced coupling to cortical parcels with both focal and diffuse thalamic connectivity patterns.

First, we investigated the relationship between ED_{pc1} loadings and anatomical coupling to discrete thalamic nuclei. For this analysis, we used the Morel histological thalamic atlas to segment the thalamus into 28 thalamic nuclei

(**Fig. 3A**). Previous studies have proposed different classifications of these nuclei based on their spatial proximity (e.g., anterior, medial, posterior, and lateral groups), primary input sources (e.g., higher-order vs first-order; (55)), or molecular architecture (e.g., primary, secondary, and tertiary; (56)). (**Fig. S25**). See **Table 1** for all nuclei labels and subgroup assignments. We tested if anatomical overlap across thalamic nuclei, or their associated classes, corresponded with ED_{pc1} loadings across the cortex.

First, we constructed an anatomical connectivity matrix to assess the strength of connectivity between each thalamic nucleus and every cortical parcel (**Fig. 3B**). Each element in the matrix represents the mean streamline count (Mean SC) between a cortical parcel and a thalamic nucleus, normalized within each nucleus. The cortical parcels with the highest group-averaged ED_{pc1} loadings are positioned at the top.

Anterior and medial thalamic nuclei are commonly associated with higher-order cognitive functions, while lateral and posterior nuclei are primarily associated with sensorimotor cognition (8, 41, 57, 58). Qualitatively, we observed that medial and anterior nuclei coupled to cortical parcels exhibiting both focal and diffuse thalamic connectivity patterns. Posterior nuclei preferentially coupled to those with focal thalamic connectivity patterns. Lateral nuclei displayed a split pattern, with ventroposterior thalamic nuclei preferentially coupling to cortical parcels with focal patterns, while ventrolateral, ventroanterior, and ventromedial nuclei preferred those with diffuse patterns.

To quantify these relationships, we correlated ED_{pc1} loadings with Mean SC values across cortex separately for each thalamic nucleus. We then compared these values between the four classes of thalamic nuclei. A Friedman test revealed a significant difference in the correlations between ED_{pc1} loadings and Mean SC across the posterior (median: 0.27), lateral (median: 0.15), anterior (median: 0.1), and medial (median: 0.01) thalamic nuclei (χ^2 (3) = 1165, p < .001) (**Fig. 3C**). Post-hoc Nemenyi tests indicated significant differences between all group comparisons (*** p = 0.001). We also examined the correlation between Mean SC and ED_{pc1} for each of the 28 thalamic nuclei, which showed modest correlations that did not survive correction for multiple comparisons (**Fig. S24**).

Previous studies have classified thalamic nuclei based on their driving inputs. First-order nuclei, which receive primary sensory inputs, transmit sensory information to the cortex. On the other hand, higher-order thalamic nuclei receive sensory inputs and inputs from layer 5 of cortex, facilitating trans-thalamic information processing (55, 59). We hypothesized that cortical parcels with focal thalamic connectivity patterns would preferentially couple with first-order thalamic nuclei, while those with diffuse thalamic connectivity patterns would preferentially couple with higher-order thalamic nuclei. We observed a stronger correlation between ED_{pc1} loadings and Mean SC for first-order (median: 0.20) compared to higher-order (median: 0.11) thalamic nuclei (Wilcoxon signed-rank test; W: 2067, p = 6.47e-134). These findings indicate that cortical parcels with diffuse thalamic

connectivity patterns preferentially couple with both first-order and higher-order thalamic nuclei, while those with focal thalamic connectivity patterns preferentially couple with first-order nuclei. We also examined the correspondence between ED_{pc1} loadings and Mean SC within thalamic classes defined based on gene expression data (56), which largely replicated these results **(Fig. S25)**. It is worth noting that other classifications of thalamo-cortical nuclei exist (e.g., 60), which we did not consider in this study.

Previous research has demonstrated that cortico-thalamic connectivity may continuously vary along the anteroposterior and mediolateral thalamic axes axes (16, 21, 61). Based on these data, we hypothesized that the thalamic connectivity spread would also continuously vary along these axes. To investigate this hypothesis, we examined cortico-thalamic anatomical connectivity along Cartesian spatial gradients within the thalamus.

First, we calculated the position of each thalamic voxel along the anteromedial-posterolateral spatial gradient. Next, we computed the Spearman correlation between a thalamic voxel's position along the anteromedial-posterolateral gradient ($PL\text{-}AM_t$) and its streamline count for each cortical parcel, representing its coupling strength ($PL\text{-}AM_c$) (schematized in **Fig. S26**) **(Fig. 3E)**. We then correlated $PL\text{-}AM_c$ values with ED_{pc1} loadings **(Fig. 3F)**. These findings revealed that cortical parcels with focal thalamic connectivity patterns exhibited a significant preference for coupling with the posterolateral thalamus ($PL\text{-}AM_c$; r_s = -0.57, p_{sa} = 0.004) compared to cortical parcels with diffuse thalamic connectivity patterns, which was largely consistent across subjects (mean: -0.53, median: -0.54, SEM = 0.004, STD = 0.12) **(Fig. S28I)**.

Additionally, we conducted specificity analyses using the dorsoventral gradient and combinations of the anteroposterior, mediolateral, and dorsoventral gradients **(Fig. S28)**. Across subjects, the average correlation between ED_{pc1} loadings and overlap across the anteromedial-posterolateral thalamic spatial gradient was the highest compared to other thalamic spatial gradients **(Fig. 3G)**.

Thalamic connectivity spread is associated with distinct cortical information processing roles. Given that the the spread of thalamic connectivity patterns varied along the cortical hierarchy, we hypothesized that thalamic connectivity spread would correspond to different information processing roles in the cortex. These findings support this hypothesis, revealing that cortical parcels with focal thalamic connectivity patterns are associated with faster, relay-like feed-forward information flow, whereas parcels with diffuse thalamic connectivity patterns are associated with slower, modulatory feed-back information flow.

We compared the spread of thalamic connectivity patterns in each cortical parcel to their anatomically coupling across two thalamic subpopulations: 'core' and 'matrix'. Thalamic neurons of the 'core' type project focally to middle cortical layers, while 'matrix' type neurons project diffusely to superficial cortical layers (5, 64). The 'core' fibers are associated with relay-type, feed-forward information flow, suited for

Fig. 4. Relationship between thalamic connectivity spread and overlap with thalamic subpopulations and intrinsic timescale across cortex. **(A)** Thalamic gradient reflecting the relative mRNA expression of Parvalbumin (PVALB) and Calbindin (CALB1)-expressing thalamic subpopulations (CP_t) (62). Warmer colors indicate higher CALB1 expression relative to PVALB. **(B)** CP_t and $PL\text{-}AM_t$ values strongly correlated with one another. **(C)** Cortical map showing the overlap between anatomical connectivity and CP_t values (CP_c). Warmer colors denote preferential coupling with CALB1-expressing thalamic populations. **(D)** ED_{pc1} loadings and CP_c values significantly correlated with one another. **(E)** Cortical map of z-transformed intrinsic timescale (τ), calculated from group-averaged resting-state data (63). **(F)** ED_{pc1} loadings and z-transformed τ values exhibited a trending correspondence when accounting for spatial-autocorrelation.

sensory processing, while the 'matrix' fibers support slower, modulatory information flow, ideal for associative processing (5, 64). Recent BOLD-derived functional connectivity work in humans demonstrated that sensory cortical areas functionally couple to 'core' thalamus while associative cortical areas functionally couple to 'matrix' thalamus (62). Building upon these findings, we hypothesized that cortical parcels with more focal thalamic connectivity patterns would anatomically couple with 'core' thalamus, while parcels with more diffuse thalamic connectivity patterns would anatomically couple with 'matrix' thalamus.

To examine this, we used the relative mRNA expression levels of calcium-binding proteins Parvalbumin (PVALB; 'core' thalamus) and Calbindin (CALB1; 'matrix' thalamus) to index each thalamic voxel's position along the core-matrix thalamic gradient (CP_t) (62) **(Fig. 4A)**. Remarkably, the CP_t gradient demonstrated a strong correlation with the anteromedial-posterolateral thalamic gradient (r_s = 0.84) **(Fig. 4B)**. While other calcium-binding proteins, like Calretinin, are expressed by thalamic neurons as well (65), we did not consider them in this study.

We then examined the correlation between thalamic streamline counts for each cortical parcel and their corresponding CP_t values, resulting in a cortical map of CP_c values **(Fig. 4C)**. We found that cortical parcels with higher ED_{pc1} loadings also displayed higher CP_c values **(Fig. 4D)**. In essence, cortical parcels with focal thalamic connectivity patterns exhibited preferential coupling with 'core' thalamus, while parcels with diffuse thalamic patterns exhibited

Fig. 5. Generalized pattern of cortical variation in thalamic connectivity spread observed in macaques. **(A)** Parcellation of macaque cortex based on the Markov Atlas (67). Exemplar thalamic connectivity patterns for M1 (area F1) and DLPFC (area 9/46d) are displayed in the magenta and cyan panels, respectively. **(B)** ED matrix. **(C)** Cortical map showing ED_{pc1} loadings. **(D)** Cortical myelin map, calculated by averaging T1w/T2w values across 30 macaque monkeys (68). **(E)** Across cortex, ED_{pc1} loadings positively correlated with T1w/T2w values at the group level and **(F)** at the subject level. **(G)** Cortical map depicting anatomical coupling with the anteromedial-posterolateral thalamic spatial gradient ($PL\text{-}AM_t$). Higher values indicate cortical parcels that preferentially couple with anteromedial thalamus ($PL\text{-}AM_c$). **(H)** Cortical parcels with higher ED_{pc1} loadings preferentially coupled with posterolateral thalamus. The red dashed line represents parcels with balanced coupling along the $PL\text{-}AM_t$ gradient. **(I)** On average, ED_{pc1} loadings and $PL\text{-}AM_c$ value negatively corresponded.

preferential coupling with 'matrix' thalamus ($r_s = 0.57$, $p_{sa} = 0.009$). Moreover, consistent with previous research on cortico-thalamic functional connectivity, association cortical parcels exhibited higher anatomical CP_c values compared to sensory cortical parcels (62) (**Fig. S29**).

Cortical parcels are known to exhibit differences in the timescales of their intrinsic BOLD fluctuations, with feed-forward processing parcels operating at relatively faster timescales compared to those involved in feedback processing (63, 66). In line with this, we hypothesized that cortical parcels with more diffuse thalamic connectivity patterns would also exhibit longer intrinsic timescales. To investigate this, we compared the spread of thalamic connectivity patterns in each cortical parcel to their intrinsic timescale (τ) derived from resting-state functional connectivity (**Fig. 4E**) (63). This analysis revealed a modest correlation between ED_{pc1} loadings and τ values, indicating that cortical parcels with focal thalamic connectivity patterns operated at faster timescales during rest. However, this relationship only showed a trending significance when accounting for spatial auto-correlation ($r_s = -0.38$, $p_{sa} = 0.09$) (**Fig. 4F**).

Generalized hierarchical variation of thalamic connectivity spread in macaques. Following previous work comparing cortico-thalamic connectivity patterns between humans and macaques (40), we investigated within-thalamus connectivity patterns in macaque monkeys. We hypothesized that macaque thalamic connectivity spread would show similarities to the human data, albeit to a lesser extent. To test this, we analyzed tractography data from six post-mortem macaque brains, obtained from 7T diffusion MRI scans. Our results revealed a similar organization of thalamo-cortical connectivity in macaques compared to humans.

We parcellated macaque tractography data to derive connectivity between 128 cortical parcels from the Markov atlas across all ipsilateral thalamic voxels (67). Macaque M1 (area F1; magenta panel) projected to the lateral portion of the thalamus, while DLPFC (area 9/46d; cyan panel) projected to medial and anterior thalamic regions (**Fig. 5A**). Comparing ED values across thresholds, we observed greater similarity between M1 and DLPFC in macaque compared to humans. We then followed the framework developed in humans to calculate macaque ED_{pc1} loadings, which accounted for 82% of the variance (**Fig. 5B-C**).

To examine the variation of ED_{pc1} loadings between sensory and association cortical parcels in macaques, we correlated macaque ED_{pc1} loadings and group-averaged T1w/T2w values obtained from (68) (**Fig. 5D**). We observed a strong positive correlation between ED_{pc1} loadings and T1w/T2w values at the group level ($r_s = 0.52$, $p_{sa} = 0.024$) (**Fig. 5E**), which was consistent on average across subjects (mean: 0.36, median: 0.34, SEM = 0.06, STD = 0.15). Surprisingly, the median value was not significantly different from the human data (**Fig. 5F**; **Fig. S30D**).

We also conducted further analysis to investigate whether cortical parcels with focal thalamic connectivity patterns in macaques preferentially coupled with the posterolateral thalamus, similar to the findings in humans. Consistent with the human data, macaque ED_{pc1} loadings showed a strong negative correlation with $PL\text{-}AM_c$ values at both the group level ($r_s = -0.57$; $p_{sa} = 0.003$) and the subject-level (mean: -0.27, median: -0.33, SEM = 0.086, STD = 0.21) **Fig. 5G-I**. This association was more pronounced compared to other thalamic spatial gradients (**Fig. S31**). Lastly, the correlation between ED_{pc1} loadings and $PL\text{-}AM_c$ values showed a trending difference between humans and macaques, while no significant differences were observed for any other thalamic gradients (**Fig. S30E**).

The relationship between macaque ED_{pc1} and ED_σ was slightly weaker and more nonlinear compared to the human data, likely due to cortical parcels with exceptionally high ED values, resulting in an underestimation of thalamic connectivity spread by the ED_σ measure. Conversely, the ED_{pc1} measure accurately captured the spread of thalamic connectivity patterns for these cortical parcels (**Fig. S30A-C**).

Discussion

This study contributes to the rich body of literature investigating the organization of cortico-thalamic systems in hu-

mans and non-human primates. By employing dMRI-derived tractography across species, we tested if different cortical areas show distinct spatial patterns of connectivity within the primate thalamus. This is critical to establish the anatomical architecture of how information flows within distinct cortico-thalamic systems. Here, we show that thalamic connectivity patterns systematically varied across cortical areas in humans and macaques. Our results implicate distinct tractography motifs corresponding to sensorimotor and association cortico-thalamic circuits. These motifs were consistent across people and generalized in macaques. Collectively, this study offers convergent evidence that different cortical areas exhibit systematic hierarchical variation in how they spatially project onto the thalamus, which may support distinct computations across cortico-thalamic loops.

Spatial properties of thalamic connectivity patterns vary along the cortical hierarchy. Here we replicate findings from prior tracer and tractography studies, providing confirmation that each cortical area exhibits a distinct pattern of anatomical connectivity within the thalamus (15, 16, 18–20, 25–27, 37, 40). Moreover, this study's findings are consistent with previous animal studies that demonstrate a correspondence between thalamic organization and the sensory-association cortical hierarchy (5, 46–49, 55). Here we build on this body of work to systematically characterize the variation of within-thalamus connectivity spread across the entire cortex. The core innovation here is that we demonstrate that sensory cortical areas exhibit more focal thalamic connectivity patterns relative to association cortical areas. Our findings suggest that the spatial distribution of within-thalamus connectivity patterns is a key distinguishing feature of corticothalamic circuits, potentially associated with their roles in sensorimotor and higher-order associative information processing. In line with this notion, other quantitative measures derived from dMRI, including connectivity strength and microstructure, do not exhibit significant variation along the cortical hierarchy.

We offer convergent evidence for this phenomenon by demonstrating that cortical areas that exhibit focal projections target posterolateral thalamus, whereas cortical areas that exhibit diffuse projections generally target anteromedial thalamus (55). This finding is consistent with previous studies showing the organization of thalamic functional connectivity, microstructure, and gene expression along the antero-posterior and mediolateral thalamic axis (16, 21, 56, 69–71). Prior work has shown that axon guidance cues along the anteromedial-posterolateral thalamic gradient shape the topography of thalamo-cortical connections (72). Powell et al. provide insight into the molecular mechanisms that may shape cortex-to-thalamus connectivity patterns. Here, it will be vital to characterize the mechanisms that give rise to hierarchical differences in the spatial spread of within-thalamus connectivity patterns, which may be critical for distinct information flow across cortico-thalamic systems and compromised in neurodevelopmental disorders (73–75).

Building on human quantitative dMRI, we show convergent effects in macaques, which mitigate limitations re-

lated to lower resolution, shorter collection time, and motion bias in humans (43). Using macaque dMRI data, previous studies have identified both similarities and differences in cortico-thalamic anatomical connectivity patterns relative to humans (76). Despite notable inter-species differences in many cortical areas, here we present data that support a generalized cortex-wide hierarchical organization of within-thalamus connectivity spread across both humans and macaques. The integration of macaque tractography and tracer data in cortico-thalamic systems remains limited (77, 78), but future studies incorporating tracer data could provide more specific insights into cortico-thalamic system organization by considering directionality. For example, it has been previously suggested that feed-forward non-reciprocal cortex-to-thalamus signals may allow information from segregated cortical pathways to converge within the thalamus (23, 24). Consequently, cortical areas with more spatial spread in their thalamic connectivity may receive a greater number of thalamic feedback projections, perhaps enabling such areas to perform broader computational modulation across distributed circuits. These hypotheses will need to be tested in future animal studies.

In addition to area-level projection patterns onto the thalamus discussed above, dMRI-derived tractography data can address gaps in the non-human primate tracer literature. For instance, it can reveal patterns regarding contralateral cortico-thalamic connections. These connections have received less attention in macaque tracer studies (19, 79, 80), yet hold potential for investigating the influence of cortico-thalamo-cortical circuits on inter-hemispheric communication.

Implications of large-scale cortico-thalamic projection patterns for understanding levels of anatomical architecture. Cortex-to-thalamus and thalamus-to-cortex neuronal fibers exhibit distinct patterns in their axonal projections (5, 19, 64). We hypothesized these distinct patterns of neuronal connectivity may also be reflected in the patterns of large-scale white matter tract terminations within the thalamus. Specifically, 'core' neurons display focal projections to middle cortical layers, while 'matrix' neurons exhibit diffuse projections to superficial cortical layers that often span architectonic boundaries (5). Moreover, prefrontal neurons have been observed to exhibit either dense, focal projections to the ipsilateral thalamus or sparse, diffuse projections to bilateral thalamus (19, 81). Therefore, the spread of axonal projections distinguishes both cortical and thalamic fibers within cortico-thalamic systems.

In this study, we show that the spread of cortical terminations within the thalamus distinguishes cortico-thalamic white matter tracts. Furthermore, we linked human neural gene expression and tractography to show that cortical areas with focal and diffuse thalamic connectivity patterns preferentially couple with PVALB-expressing 'core' and CALB1-expressing 'matrix' thalamus, respectively. This finding highlights a mirrored anatomical principle of variation between individual neuronal fibers and large-scale tracts within cortico-thalamic systems. This finding suggests that cortico-

thalamic white matter tracts may be composed of individual neuronal fibers with similar termination patterns.

This finding aligns with theoretical proposals that hypothesize similar principles governing connectivity at multiple levels of analysis (82, 83). Moreover, it implies that core principles of variation in thalamic fibers provide insights into the properties of large-scale white matter tracts. For instance, thalamic neurons exhibit variations in the spread of their axonal projections across cortical layers (5, 64). Future studies employing dMRI tractography can investigate if this effect extends to the spread of streamline terminations from thalamic nuclei to individual cortical areas.

Additionally, previous studies have shown that cortical projections onto the contralateral thalamus primarily involve non-reciprocal connections (9, 19). We found that cortical areas with stronger contralateral thalamic connectivity display more focal thalamic connectivity patterns across the contralateral thalamus. This suggests that cortical areas may relay information to the opposite hemisphere via feed-forward connections onto the contralateral thalamus.

Interrogating cortico-thalamic anatomical connectivity to better understand the role of the thalamus in shaping brain-wide information flow. While early studies suggested that the thalamus primarily relays information through parallel and segregated circuits from the sensory periphery, accumulating empirical and computational evidence supports the notion that the thalamus is a crucial integration hub capable of coordinating and sustaining signals across the cortex (11, 12, 49, 84–86). The structural properties of cortico-thalamic circuits invariably constrain the types of computations these circuits can support (6), and prior work has established a relationship between the spread of axonal terminations in cortico-thalamic systems and both feed-forward and feed-back information flow (5, 19). While the present findings do not directly support directional interpretations, our data raise the possibility that diffuse thalamic connectivity patterns may be associated with slower, feed-back brain dynamics that may support integrative information transmission, whereas focal thalamic connectivity patterns may be associated with faster, feed-forward brain dynamics that may support the relay of sensory information.

Complementary functional neuroimaging studies have revealed areas of integration and segregation within the human thalamus (58, 87, 88). The anatomical basis for thalamic integration is not fully understood, but it has been proposed that overlapping terminations may contribute to this process (22, 23, 37). It is hypothesized that overlapping thalamic connections emerging from segregated cortical circuits may enable the integration of information across different cortical areas (41). In this study, we demonstrate that certain cortical areas, such as the anterior cingulate and precuneus, exhibit diffuse thalamic connectivity, allowing them to impact thalamo-cortical information in a distributed manner. Specifically, the anterior cingulate demonstrates widespread connections across the thalamus, aligning with observations by Phillips et al. that area 24 exhibits diffuse anatomical terminations across the mediodorsal nucleus of the thalamus (37).

This property has been hypothesized to support the integration of signals within the prefrontal cortex (41).

Circuits connecting the cortex, thalamus, and the basal ganglia have also been implicated in integrative functions (23, 89). The patterns of anatomical connections between the basal ganglia and thalamus may also exhibit variation in their spatial properties. Future studies can better elucidate the variation in the spatial properties of anatomical connections between the thalamus and basal ganglia and how such variation may correspond to integrated vs. segregated cortico-basal ganglia-thalamo-cortical functional interactions

Our findings indicate that cortical areas with diffuse thalamic connectivity patterns couple to both anteromedial and posterolateral thalamic subregions. Surprisingly, these results did not support the hypothesis that 'higher-order' thalamic nuclei preferentially couple with cortical areas displaying diffuse thalamic connectivity. Instead, these nuclei display more balanced coupling to cortical areas with both focal and diffuse thalamic connectivity patterns. Previous studies have shown that 'higher-order' thalamic nuclei receive input from cortical and subcortical regions (90, 91). This could be one mechanism for how the integration of higher-order and first-order signals in the thalamus may support complex cognitive functions (92).

Our findings indicate that cortical areas with diffuse thalamic connectivity patterns are coupled to both anteromedial and posterolateral thalamic subregions. Surprisingly, these results did not support the hypothesis that 'higher-order' thalamic nuclei preferentially couple with cortical areas displaying diffuse thalamic connectivity. Instead, these nuclei showed more balanced coupling to cortical areas with both focal and diffuse thalamic connectivity patterns. Previous studies have demonstrated that 'higher-order' thalamic nuclei receive input from cortical and subcortical regions (90, 91). This mechanism may explain how the integration of higher-order and first-order signals in the thalamus supports complex cognitive functions (92).

Although we did not directly compare cortico-thalamic structure and function in this study, we offer an anatomical framework to complement the findings of a previous investigation on cortico-thalamic functional coupling (62). Muller et al. (62) examined cortico-thalamic BOLD-derived functional connectivity and demonstrated that sensory cortical areas exhibited stronger functional coupling with 'core' thalamus, whereas association cortical areas exhibited stronger functional coupling with 'matrix' thalamus, and this dichotomy was found to align with patterns of whole-brain dynamics. However, the specific ways in which thalamo-cortical anatomical connectivity constraints may shape functional connectivity remain unknown. A possible future direction would be to investigate how thalamo-cortical anatomical connections contribute to both intra- and inter-hemispheric functional interactions.

Lastly, these data imply that cortical areas with diffuse thalamic connectivity patterns may selectively overlap with thalamic subregions implicated in functional integration and multimodal cognitive processes (58, 88). This observation

warrants future investigations in combination with functional modalities.

Study limitations. While powerful, probabilistic tractography has notable limitations, such as the possibility of producing false positives and lacking directionality information ([93], [94]). In this study, we did not find any specific confounding factors systematically influencing the patterns of thalamic connectivity derived from tractography. However, we did observe that cortical curvature and sulcal depth were positively correlated with an increased mean streamline count, potentially reflecting gyral bias.

Furthermore, we observed significant individual variation in thalamic connectivity patterns. Notably, we identified biologically unlikely tractography patterns in some subjects. The nature of these differences, whether they represent true individual variation or false positive connections, remains unclear, but it is likely a combination of both. While these variations did not appear to produce a spurious difference in thalamic connectivity spread between sensory and association cortical areas, these data are consistent with previous reports suggesting that interpretations of tractography data have inherent limitations (e.g., ([93], [94])).

Importantly, tractography-based anatomical connectivity has shown strong agreement with invasive tract tracing studies conducted in monkeys, providing validation for its use ([43], [95]–[101]). However, it is worth noting that the majority of these investigations have primarily focused on cortico-cortical connections and connectivity at the areal level. In comparison, studies directly assessing the correspondence of tractography-derived connectivity with tracer-derived connectivity in thalamo-cortical systems are limited ([94], [102]). Consequently, future research should prioritize examining the correspondence between tracer-based and tractography-derived anatomical terminations within subcortical gray matter structures while also investigating factors that may contribute to the presence of biologically implausible connections across different individuals. This is vital for a comprehensive understanding of inter-individual variation in anatomical connectivity and its implications for cognition and behavior in both health and disease.

Conclusions

The thalamus plays a key role in various neural functions involving multiple sensory modalities ([8], [41], [103]). Dysfunction of the thalamus has been linked to severe neuropsychiatric disorders such as psychosis ([104]–[109]), and the symptoms of these disorders have been associated with abnormal anatomical cortico-thalamic connectivity ([73]–[75]). However, our understanding of the contribution of the thalamus to healthy whole-brain information transmission and its dysfunction in diseases has been hindered by limited knowledge of the underlying circuitry, especially in humans ([6], [110]).

Since the first *in vivo* examinations of cortico-thalamic anatomy in humans ([35], [36]), neuroimaging studies have made significant progress in mapping thalamo-cortical circuitry and investigating its role in shaping whole-brain func-

tional interactions ([54], [111]). This study provides quantitative evidence that the spatial properties of within-thalamus anatomical connectivity patterns exhibit variation across cortical areas, following established hierarchical principles of cortical organization, which may reflect variations associated with different types of information processing. Here we demonstrate that an in-depth investigation of cortico-thalamic anatomical circuitry can offer insights into how the thalamus supports segregated and integrated information flow throughout the brain, which may underlie the computations enabling higher-order cognition in humans.

ACKNOWLEDGEMENTS
We thank Jie Lisa Ji, Grega Repovš, Jure Demsar, and Takuya Ito for assistance with data processing and analysis and Amy F.T. Arnsten, Jane R. Taylor, John H. Krystal, Maxwell Shinn, Rachel Cooper, Jacob A. Miller, and Warren W. Pettine for helpful discussions. This work was supported by the Guber Foundation (AMH) and the National Institute of Mental Health (Grant Nos. U01 MH121766-03, R01 MH112189) (AA). Data collection and sharing for this project was provided by the Human Connectome Project (HCP) and the PRIMatE Data Exchange (PRIME-DE).

Bibliography

1. S Murray Sherman and R W Guillery. Distinct functions for direct and transthalamic corticocortical connections. *J Neurophysiol*, 106(3):1068–1077, Sep 2011. ISSN 1522-1598 (Electronic); 0022-3077 (Linking). doi: 10.1152/jn.00429.2011.

2. Scott S Bolkan, Joseph M Stujenske, Sebastien Parnaudeau, Timothy J Spellman, Caroline Rauffenbart, Atheir I Abbas, Alexander Z Harris, Joshua A Gordon, and Christoph Kellendonk. Thalamic projections sustain prefrontal activity during working memory maintenance. *Nat Neurosci*, 20(7):987–996, Jul 2017. ISSN 1546-1726 (Electronic); 1097-6256 (Print); 1097-6256 (Linking). doi: 10.1038/nn.4568.

3. Rajeev V Rikhye, Ralf D Wimmer, and Michael M Halassa. Toward an integrative theory of thalamic function. *Annu Rev Neurosci*, 41:163–183, Jul 2018. ISSN 1545-4126 (Electronic); 0147-006X (Linking). doi: 10.1146/annurev-neuro-080317-062144.

4. Miho Nakajima and Michael M Halassa. Thalamic control of functional cortical connectivity. *Curr Opin Neurobiol*, 44:127–131, Jun 2017. ISSN 1873-6882 (Electronic); 0959-4388 (Print); 0959-4388 (Linking). doi: 10.1016/j.conb.2017.04.001.

5. E G Jones. The thalamic matrix and thalamocortical synchrony. *Trends Neurosci*, 24 (10):595–601, Oct 2001. ISSN 0166-2236 (Print); 0166-2236 (Linking). doi: 10.1016/s0166-2236(00)01922-6.

6. Michael M Halassa and S Murray Sherman. Thalamocortical circuit motifs: A general framework. *Neuron*, 103(5):762–770, Sep 2019. ISSN 1097-4199 (Electronic); 0896-6273 (Print); 0896-6273 (Linking). doi: 10.1016/j.neuron.2019.06.005.

7. Dheeraj S Roy, Ying Zhang, Michael M Halassa, and Guoping Feng. Thalamic subnetworks as units of function. *Nat Neurosci*, 25(2):140–153, Feb 2022. ISSN 1546-1726 (Electronic); 1097-6256 (Print); 1097-6256 (Linking). doi: 10.1038/s41593-021-00996-1.

8. Yuri B Saalmann and Sabine Kastner. The cognitive thalamus. *Front Syst Neurosci*, 9: 39, 2015. ISSN 1662-5137 (Print); 1662-5137 (Electronic); 1662-5137 (Linking). doi: 10.3389/fnsys.2015.00039.

9. Mathieu Wolff and Seralynne D. Vann. The cognitive thalamus as a gateway to mental representations. *Journal of Neuroscience*, 39(1):3–14, 2019. ISSN 0270-6474. doi: 10.1523/JNEUROSCI.0479-18.2018.

10. Weiguo Yang, Sri Laasya Tipparaju, Guang Chen, and Nuo Li. Thalamus-driven functional populations in frontal cortex support decision-making. *Nat Neurosci*, 25(10):1339–1352, Oct 2022. ISSN 1546-1726 (Electronic); 1097-6256 (Print); 1097-6256 (Linking). doi: 10.1038/s41593-022-01171-w.

11. L. Ian Schmitt, Ralf D. Wimmer, Miho Nakajima, Michael Happ, Sima Mofakham, and Michael M. Halassa. Thalamic amplification of cortical connectivity sustains attentional control. *Nature*, 545(7653):219–223, 2017. doi: 10.1038/nature22073.

12. Zengcai V Guo, Hidehiko K Inagaki, Kayvon Daie, Shaul Druckmann, Charles R Gerfen, and Karel Svoboda. Maintenance of persistent activity in a frontal thalamocortical loop. *Nature*, 545(7653):181–186, May 2017. ISSN 1476-4687 (Electronic); 0028-0836 (Print); 0028-0836 (Linking). doi: 10.1038/nature22324.

13. Miguel Ángel García-Cabezas, Isabel Pérez-Santos, and Carmen Cavada. Mapping the primate thalamus: historical perspective and modern approaches for defining nuclei. *Brain Struct Funct*, 228(5):1125–1151, Jun 2023. ISSN 1863-2661 (Electronic); 1863-2653 (Print); 1863-2653 (Linking). doi: 10.1007/s00429-022-02594-4.

14. Jean-Christophe Cassel and Anne Pereira de Vasconcelos. Routes of the thalamus through the history of neuroanatomy. *Neuroscience Biobehavioral Reviews*, 125:442–465, 2021. ISSN 0149-7634. doi: https://doi.org/10.1016/j.neubiorev.2021.03.001.

15. Dongyang Zhang, Abraham Z Snyder, Joshua S Shimony, Michael D Fox, and Marcus E Raichle. Noninvasive functional and structural connectivity mapping of the human thalamocortical system. *Cereb Cortex*, 20(5):1187–1194, May 2010. ISSN 1460-2199 (Electronic); 1047-3211 (Print); 1047-3211 (Linking). doi: 10.1093/cercor/bhp182.

16. Christian Lambert, Henry Simon, Jordan Colman, and Thomas R. Barrick. Defining thalamic nuclei and topographic connectivity gradients in vivo. *NeuroImage*, 158:466–479, 2017. ISSN 1053-8119. doi: https://doi.org/10.1016/j.neuroimage.2016.08.028.

17. D. Xiao, B. Zikopoulos, and H. Barbas. Laminar and modular organization of prefrontal

projections to multiple thalamic nuclei. *Neuroscience*, 161(4):1067–1081, 2009. ISSN 0306-4522. doi: https://doi.org/10.1016/j.neuroscience.2009.04.034.

18. H Barbas, T H Henion, and C R Dermon. Diverse thalamic projections to the prefrontal cortex in the rhesus monkey. *J Comp Neurol*, 313(1):65–94, Nov 1991. ISSN 0021-9967 (Print); 0021-9967 (Linking). doi: 10.1002/cne.903130106.

19. T M Preuss and P S Goldman-Rakic. Crossed corticothalamic and thalamocortical connections of macaque prefrontal cortex. *J Comp Neurol*, 257(2):269–281, Mar 1987. ISSN 0021-9967 (Print); 0021-9967 (Linking). doi: 10.1002/cne.902570211.

20. L M Romanski, M Giguere, J F Bates, and P S Goldman-Rakic. Topographic organization of medial pulvinar connections with the prefrontal cortex in the rhesus monkey. *J Comp Neurol*, 379(3):313–332, Mar 1997. ISSN 0021-9967 (Print); 0021-9967 (Linking).

21. Stuart Oldham and Gareth Ball. A phylogenetically-conserved axis of thalamocortical connectivity in the human brain. *bioRxiv*, page 2022.11.15.516574, 01 2022. doi: 10.1101/2022.11.15.516574.

22. Bogdan Draganski, Ferath Kherif, Stefan Klöppel, Philip A Cook, Daniel C Alexander, Geoff J M Parker, Ralf Deichmann, John Ashburner, and Richard S J Frackowiak. Evidence for segregated and integrative connectivity patterns in the human basal ganglia. *J Neurosci*, 28(28):7143–7152, Jul 2008. ISSN 1529-2401 (Electronic); 0270-6474; 0270-6474 (Linking). doi: 10.1523/JNEUROSCI.1486-08.2008.

23. Suzanne N Haber and Roberta Calzavara. The cortico-basal ganglia integrative network: the role of the thalamus. *Brain Res Bull*, 78(2-3):69–74, Feb 2009. ISSN 1873-2747 (Electronic); 0361-9230 (Print); 0361-9230 (Linking). doi: 10.1016/j.brainresbull.2008.09.013.

24. Nikolaus R McFarland and Suzanne N Haber. Thalamic relay nuclei of the basal ganglia form both reciprocal and nonreciprocal cortical connections, linking multiple frontal cortical areas. *J Neurosci*, 22(18):8117–8132, Sep 2002. ISSN 1529-2401 (Electronic); 0270-6474 (Print); 0270-6474 (Linking). doi: 10.1523/JNEUROSCI.22-18-08117.2002.

25. P S Goldman-Rakic and L J Porrino. The primate mediodorsal (md) nucleus and its projection to the frontal lobe. *J Comp Neurol*, 242(4):535–560, Dec 1985. ISSN 0021-9967 (Print); 0021-9967 (Linking). doi: 10.1002/cne.902420406.

26. D F Siwek and D N Pandya. Prefrontal projections to the mediodorsal nucleus of the thalamus in the rhesus monkey. *J Comp Neurol*, 312(4):509–524, Oct 1991. ISSN 0021-9967 (Print); 0021-9967 (Linking). doi: 10.1002/cne.903120403.

27. L D Selemon and P S Goldman-Rakic. Common cortical and subcortical targets of the dorsolateral prefrontal and posterior parietal cortices in the rhesus monkey: evidence for a distributed neural network subserving spatially guided behavior. *J Neurosci*, 8(11):4049–4068, Nov 1988. ISSN 0270-6474 (Print); 1529-2401 (Electronic); 0270-6474 (Linking). doi: 10.1523/JNEUROSCI.08-11-04049.1988.

28. Saad Jbabdi, Stamatios N Sotiropoulos, Suzanne N Haber, David C Van Essen, and Timothy E Behrens. Measuring macroscopic brain connections in vivo. *Nat Neurosci*, 18(11):1546–1555, Nov 2015. ISSN 1546-1726 (Electronic); 1097-6256 (Linking). doi: 10.1038/nn.4134.

29. P J Basser, J Mattiello, and D LeBihan. Mr diffusion tensor spectroscopy and imaging. *Biophys J*, 66(1):259–267, Jan 1994. ISSN 0006-3495 (Print); 1542-0086 (Electronic); 0006-3495 (Linking). doi: 10.1016/S0006-3495(94)80775-1.

30. Denis Le Bihan. Diffusion mri: what water tells us about the brain. *EMBO Mol Med*, 6(5):569–573, May 2014. ISSN 1757-4684 (Electronic); 1757-4676 (Print); 1757-4676 (Linking). doi: 10.1002/emmm.201404055.

31. T.E.J. Behrens, H. Johansen Berg, S. Jbabdi, M.F.S. Rushworth, and M.W. Woolrich. Probabilistic diffusion tractography with multiple fibre orientations: What can we gain? *NeuroImage*, 34(1):144 – 155, 2007. ISSN 1053-8119. doi: https://doi.org/10.1016/j.neuroimage.2006.09.018.

32. Stamatios N Sotiropoulos and Andrew Zalesky. Building connectomes using diffusion mri: why, how and but. *NMR Biomed*, 32(4):e3752, Apr 2019. ISSN 1099-1492 (Electronic); 0952-3480 (Print); 0952-3480 (Linking). doi: 10.1002/nbm.3752.

33. Saad Jbabdi and Heidi Johansen-Berg. Tractography: where do we go from here? *Brain Connect*, 1(3):169–183, 2011. ISSN 2158-0022 (Electronic); 2158-0014 (Print); 2158-0014 (Linking). doi: 10.1089/brain.2011.0033.

34. Fan Zhang, Alessandro Daducci, Yong He, Simona Schiavi, Caio Seguin, Robert E Smith, Chun-Hung Yeh, Tengda Zhao, and Lauren J O'Donnell. Quantitative mapping of the brain's structural connectivity using diffusion mri tractography: A review. *Neuroimage*, 249: 118870, Apr 2022. ISSN 1095-9572 (Electronic); 1053-8119 (Print); 1053-8119 (Linking). doi: 10.1016/j.neuroimage.2021.118870.

35. Heidi Johansen-Berg, Timothy E J Behrens, Emma Sillery, Olga Ciccarelli, Alan J Thompson, Stephen M Smith, and Paul M Matthews. Functional-anatomical validation and individual variation of diffusion tractography-based segmentation of the human thalamus. *Cereb Cortex*, 15(1):31–39, Jan 2005. ISSN 1047-3211 (Print); 1047-3211 (Linking). doi: 10.1093/cercor/bhh105.

36. T E J Behrens, H Johansen-Berg, M W Woolrich, S M Smith, C A M Wheeler-Kingshott, P A Boulby, G J Barker, E L Sillery, K Sheehan, O Ciccarelli, A J Thompson, J M Brady, and P M Matthews. Non-invasive mapping of connections between human thalamus and cortex using diffusion imaging. *Nat Neurosci*, 6(7):750–757, Jul 2003. ISSN 1097-6256 (Print); 1097-6256 (Linking). doi: 10.1038/nn1075.

37. Jessica M Phillips, Lesenia R Fish, Niranjan A Kambi, Michelle J Redinbaugh, Sounak Mohanta, Steven R Kecskemeti, and Yuri B Saalmann. Topographic organization of connections between prefrontal cortex and mediodorsal thalamus: Evidence for a general principle of indirect thalamic pathways between directly connected cortical areas. *Neuroimage*, 189:832–846, Apr 2019. ISSN 1095-9572 (Electronic); 1053-8119 (Print); 1053-8119 (Linking). doi: 10.1016/j.neuroimage.2019.01.078.

38. András Jakab, Rémi Blanc, and Ervin L Berényi. Mapping changes of in vivo connectivity patterns in the human mediodorsal thalamus: correlations with higher cognitive and executive functions. *Brain Imaging Behav*, 6(3):472–483, Sep 2012. ISSN 1931-7565 (Electronic); 1931-7557 (Linking). doi: 10.1007/s11682-012-9172-5.

39. Monica Giraldo-Chica, Baxter P Rogers, Stephen M Damon, Bennett A Landman, and Neil D Woodward. Prefrontal-thalamic anatomical connectivity and executive cognitive function in schizophrenia. *Biol Psychiatry*, 83(6):509–517, Mar 2018. ISSN 1873-2402

(Electronic); 0006-3223 (Print); 0006-3223 (Linking). doi: 10.1016/j.biopsych.2017.09.022.

40. Johannes C Klein, Matthew F S Rushworth, Timothy E J Behrens, Clare E Mackay, Alex J de Crespigny, Helen D'Arceuil, and Heidi Johansen-Berg. Topography of connections between human prefrontal cortex and mediodorsal thalamus studied with diffusion tractography. *Neuroimage*, 51(2):555–564, Jun 2010. ISSN 1095-9572 (Electronic); 1053-8119 (Print); 1053-8119 (Linking). doi: 10.1016/j.neuroimage.2010.02.062.

41. Jessica M Phillips, Niranjan A Kambi, Michelle J Redinbaugh, Sounak Mohanta, and Yuri B Saalmann. Disentangling the influences of multiple thalamic nuclei on prefrontal cortex and cognitive control. *Neurosci Biobehav Rev*, 128:487–510, Sep 2021. ISSN 1873-7528 (Electronic); 0149-7634 (Print); 0149-7634 (Linking). doi: 10.1016/j.neubiorev.2021.06.042.

42. Paula L. Croxson, Heidi Johansen-Berg, Timothy E. J. Behrens, Matthew D. Robson, Mark A. Pinsk, Charles G. Gross, Wolfgang Richter, Marlene C. Richter, Sabine Kastner, and Matthew F. S. Rushworth. Quantitative investigation of connections of the prefrontal cortex in the human and macaque using probabilistic diffusion tractography. *Journal of Neuroscience*, 25(39):8854–8866, 2005. ISSN 0270-6474. doi: 10.1523/JNEUROSCI.1311-05.2005.

43. Anastasia Yendiki, Manisha Aggarwal, Markus Axer, Amy F D Howard, Anne-Marie van Cappellen van Walsum, and Suzanne N Haber. Post mortem mapping of connectional anatomy for the validation of diffusion mri. *Neuroimage*, 256:119146, Aug 2022. ISSN 1095-9572 (Electronic); 1053-8119 (Print); 1053-8119 (Linking). doi: 10.1016/j.neuroimage.2022.119146.

44. Joshua B. Burt, Markus Helmer, Maxwell Shinn, Alan Anticevic, and John D. Murray. Generative modeling of brain maps with spatial autocorrelation. *NeuroImage*, 220:117038, 2020. ISSN 1053-8119. doi: https://doi.org/10.1016/j.neuroimage.2020.117038.

45. Jie Lisa Ji, Marjolein Spronk, Kaustubh Kulkarni, Grega Repovš, Alan Anticevic, and Michael W Cole. Mapping the human brain's cortical-subcortical functional network organization. *NeuroImage*, 185:35–57, 2019.

46. Julie A Harris, Stefan Mihalas, Karla E Hirokawa, Jennifer D Whitesell, Hannah Choi, Amy Bernard, Phillip Bohn, Shiella Caldejon, Linzy Casal, Andrew Cho, Aaron Feiner, David Feng, Nathalie Gaudreault, Charles R Gerfen, Nile Graddis, Peter A Groblewski, Alex M Henry, Anh Ho, Robert Howard, Joseph E Knox, Leonard Kuan, Xiuli Kuang, Jerome Lecoq, Phil Lesnar, Yaoyao Li, Jennifer Luviano, Stephen McConoughey, Marty T Mortrud, Maitham Naeemi, Lydia Ng, Seung Wook Oh, Benjamin Ouellette, Elise Shen, Staci A Sorensen, Wayne Wakeman, Quanxin Wang, Yun Wang, Ali Williford, John W Phillips, Allan R Jones, Christof Koch, and Hongkui Zeng. Hierarchical organization of cortical and thalamic connectivity. *Nature*, 575(7781):195–202, Nov 2019. ISSN 1476-4687 (Electronic); 0028-0836 (Print); 0028-0836 (Linking). doi: 10.1038/s41586-019-1716-z.

47. Guofen Ma, Yanmei Liu, Lizhao Wang, Zhongyi Xiao, Kun Song, Yanjie Wang, Wanling Peng, Xiaotong Liu, Ziyue Wang, Sen Jin, Zi Tao, Chengyu T Li, Tianle Xu, Fuqiang Xu, Min Xu, and Siyu Zhang. Hierarchy in sensory processing reflected by innervation balance on cortical interneurons. *Sci Adv*, 7(20), May 2021. ISSN 2375-2548 (Electronic); 2375-2548 (Linking). doi: 10.1126/sciadv.abf5676.

48. Reza Abbas Farishta, Denis Boire, and Christian Casanova. Hierarchical Organization of Corticothalamic Projections to the Pulvinar. *Cerebral Cortex Communications*, 1(1), 07 2020. ISSN 2632-7376. doi: 10.1093/texcom/tgaa030. tgaa030.

49. Arghya Mukherjee, Navdeep Bajwa, Norman H Lam, César Porrero, Francisco Clasca, and Michael M Halassa. Variation of connectivity across exemplar sensory and associative thalamocortical loops in the mouse. *eLife*, 9:e62554, oct 2020. ISSN 2050-084X. doi: 10.7554/eLife.62554.

50. Boris C Bernhardt, Jonathan Smallwood, Shella Keilholz, and Daniel S Margulies. Gradients in brain organization. *Neuroimage*, 251:118987, Feb 2022. ISSN 1095-9572 (Electronic); 1053-8119 (Linking). doi: 10.1016/j.neuroimage.2022.118987.

51. Joshua B Burt, Murat Demirtaş, William J Eckner, Natasha M Navejar, Jie Lisa Ji, William J Martin, Alberto Bernacchia, Alan Anticevic, and John D Murray. Hierarchy of transcriptomic specialization across human cortex captured by structural neuroimaging topography. *Nat Neurosci*, 21(9):1251–1259, Sep 2018. ISSN 1546-1726 (Electronic); 1097-6256 (Print); 1097-6256 (Linking). doi: 10.1038/s41593-018-0195-0.

52. Daniel S. Margulies, Satrajit S. Ghosh, Alexandros Goulas, Marcel Falkiewicz, Julia M. Huntenburg, Georg Langs, Gleb Bezgin, Simon B. Eickhoff, F. Xavier Castellanos, Michael Petrides, Elizabeth Jefferies, and Jonathan Smallwood. Situating the default-mode network along a principal gradient of macroscale cortical organization. *Proceedings of the National Academy of Sciences*, 113(44):12574–12579, 2016. ISSN 0027-8424. doi: 10.1073/pnas.1608282113.

53. Axel Krauth, Remi Blanc, Alejandra Poveda, Daniel Jeanmonod, Anne Morel, and Gábor Székely. A mean three-dimensional atlas of the human thalamus: generation from multiple histological data. *Neuroimage*, 49(3):2053–2062, Feb 2010. ISSN 1095-9572 (Electronic); 1053-8119 (Print); 1053-8119 (Linking). doi: 10.1016/j.neuroimage.2009.10.042.

54. S Murray Sherman. Functioning of circuits connecting thalamus and cortex. *Compr Physiol*, 7(2):713–739, Mar 2017. ISSN 2040-4603 (Electronic); 2040-4603 (Linking). doi: 10.1002/cphy.c160032.

55. S Murray Sherman and R W Guillery. The role of the thalamus in the flow of information to the cortex. *Philos Trans R Soc Lond B Biol Sci*, 357(1428):1695–1708, Dec 2002. ISSN 0962-8436 (Print); 1471-2970 (Electronic); 0962-8436 (Linking). doi: 10.1098/rstb.2002.1161.

56. James W Phillips, Anton Schulmann, Erina Hara, Johan Winnubst, Chenghao Liu, Vera Valakh, Lihua Wang, Brenda C Shields, Wyatt Korff, Jayaram Chandrashekar, Andrew L Lemire, Brett Mensh, Joshua T Dudman, Sacha B Nelson, and Adam W Hantman. A repeated molecular architecture across thalamic pathways. *Nat Neurosci*, 22(11):1925–1935, Nov 2019. ISSN 1546-1726 (Electronic); 1097-6256 (Print); 1097-6256 (Linking). doi: 10.1038/s41593-019-0483-3.

57. Yuri B Saalmann. Intralaminar and medial thalamic influence on cortical synchrony, information transmission and cognition. *Front Syst Neurosci*, 8:83, 2014. ISSN 1662-5137 (Print); 1662-5137 (Electronic); 1662-5137 (Linking). doi: 10.3389/fnsys.2014.00083.

58. Kazuya Kawabata, Epifanio Bagarinao, Hirohisa Watanabe, Satoshi Maesawa, Daisuke Mori, Kazuhiro Hara, Reiko Ohdake, Michihito Masuda, Aya Ogura, Toshiyasu Kato, Shuji Koyama, Masahisa Katsuno, Toshihiko Wakabayashi, Masafumi Kuzuya, Minoru

Hoshiyama, Haruo Isoda, Shinji Naganawa, Norio Ozaki, and Gen Sobue. Bridging large-scale cortical networks: Integrative and function-specific hubs in the thalamus. *iScience*, 24(10):103106, Oct 2021. ISSN 2589-0042 (Electronic); 2589-0042 (Linking). doi: 10.1016/j.isci.2021.103106.

59. S Murray Sherman. The thalamus is more than just a relay. *Curr Opin Neurobiol*, 17(4): 417–422, Aug 2007. ISSN 0959-4388 (Print); 1873-6882 (Electronic); 0959-4388 (Linking). doi: 10.1016/j.conb.2007.07.003.

60. Francisco Clascá, Pablo Rubio-Garrido, and Denis Jabaudon. Unveiling the diversity of thalamocortical neuron subtypes. *Eur J Neurosci*, 35(10):1524–1532, May 2012. ISSN 1460-9568 (Electronic); 0953-816X (Linking). doi: 10.1111/j.1460-9568.2012.08033.x.

61. Siqi Yang, Yao Meng, Jiao Li, Bing Li, Yun-Shuang Fan, Huafu Chen, and Wei Liao. The thalamic functional gradient and its relationship to structural basis and cognitive relevance. *NeuroImage*, 218:116960, 2020. ISSN 1053-8119. doi: https://doi.org/10.1016/j.neuroimage.2020.116960.

62. Eli J. Müller, Brandon Munn, Luke J. Hearne, Jared B. Smith, Ben Fulcher, Aurina Arnatkevičiūtė, Daniel J. Lurie, Luca Cocchi, and James M. Shine. Core and matrix thalamic subpopulations relate to spatio-temporal cortical connectivity gradients. *NeuroImage*, 222: 117224, 2020. ISSN 1053-8119. doi: https://doi.org/10.1016/j.neuroimage.2020.117224.

63. Takuya Ito, Luke J Hearne, and Michael W Cole. A cortical hierarchy of localized and distributed processes revealed via dissociation of task activations, connectivity changes, and intrinsic timescales. *Neuroimage*, 221:117141, Nov 2020. ISSN 1095-9572 (Electronic); 1053-8119 (Print); 1053-8119 (Linking). doi: 10.1016/j.neuroimage.2020.117141.

64. Edward G Jones. Synchrony in the interconnected circuitry of the thalamus and cerebral cortex. *Ann N Y Acad Sci*, 1157:10–23, Mar 2009. ISSN 1749-6632 (Electronic); 0077-8923 (Linking). doi: 10.1111/j.1749-6632.2009.04534.x.

65. M.C. Münkle, H.J. Waldvogel, and R.L.M. Faull. The distribution of calbindin, calretinin and parvalbumin immunoreactivity in the human thalamus. *Journal of Chemical Neuroanatomy*, 19(3):155–173, 2000. ISSN 0891-0618. doi: https://doi.org/10.1016/S0891-0618(00)00060-0.

66. John D Murray, Alan Anticevic, Mark Gancsos, Megan Ichinose, Philip R Corlett, John H Krystal, and Xiao-Jing Wang. Linking microcircuit dysfunction to cognitive impairment: effects of disinhibition associated with schizophrenia in a cortical working memory model. *Cereb Cortex*, 24(4):859–872, Apr 2014. ISSN 1460-2199 (Electronic); 1047-3211 (Print); 1047-3211 (Linking). doi: 10.1093/cercor/bhs370.

67. N T Markov, M M Ercsey-Ravasz, A R Ribeiro Gomes, C Lamy, L Magrou, J Vezoli, P Misery, A Falchier, R Quilodran, M A Gariel, J Sallet, R Gamanut, C Huissoud, S Clavagnier, P Giroud, D Sappey-Marinier, P Barone, C Dehay, Z Toroczkai, K Knoblauch, D C Van Essen, and H Kennedy. A weighted and directed interareal connectivity matrix for macaque cerebral cortex. *Cereb Cortex*, 24(1):17–36, Jan 2014. ISSN 1460-2199 (Electronic); 1047-3211 (Print); 1047-3211 (Linking). doi: 10.1093/cercor/bhs270.

68. Takuya Hayashi, Yujie Hou, Matthew F Glasser, Joonas A Autio, Kenneth Knoblauch, Miho Inoue-Murayama, Tim Coalson, Essa Yacoub, Stephen Smith, Henry Kennedy, and David C Van Essen. The nonhuman primate neuroimaging and neuroanatomy project. *NeuroImage*, 229:117726, 2021. ISSN 1053-8119. doi: https://doi.org/10.1016/j.neuroimage.2021.117726.

69. Siqi Yang, Yao Meng, Jiao Li, Bing Li, Yun-Shuang Fan, Huafu Chen, and Wei Liao. The thalamic functional gradient and its relationship to structural basis and cognitive relevance. *Neuroimage*, 218:116960, Sep 2020. ISSN 1095-9572 (Electronic); 1053-8119 (Linking). doi: 10.1016/j.neuroimage.2020.116960.

70. W Brysch, I Brysch, O D Creutzfeldt, R Schlingensiepen, and K H Schlingensiepen. The topology of the thalamo-cortical projections in the marmoset monkey (callithrix jacchus). *Exp Brain Res*, 81(1):1–17, 1990. ISSN 0014-4819 (Print); 0014-4819 (Linking). doi: 10.1007/BF00230095.

71. J C Höhl-Abrahão and O D Creutzfeldt. Topographical mapping of the thalamocortical projections in rodents and comparison with that in primates. *Exp Brain Res*, 87(2):283–294, 1991. ISSN 0014-4819 (Print); 0014-4819 (Linking). doi: 10.1007/BF00231845.

72. Ashton W Powell, Takayuki Sassa, Yongqin Wu, Marc Tessier-Lavigne, and Franck Polleux. Topography of thalamic projections requires attractive and repulsive functions of netrin-1 in the ventral telencephalon. *PLoS Biol*, 6(5):e116, May 2008. ISSN 1545-7885 (Electronic); 1544-9173 (Print); 1544-9173 (Linking). doi: 10.1371/journal.pbio.0060116.

73. Monica Giraldo-Chica and Neil D Woodward. Review of thalamocortical resting-state fmri studies in schizophrenia. *Schizophr Res*, 180:58–63, Feb 2017. ISSN 1573-2509 (Electronic); 0920-9964 (Print); 0920-9964 (Linking). doi: 10.1016/j.schres.2016.08.005.

74. Julia M Sheffield, Anna S Huang, Baxter P Rogers, Monica Giraldo-Chica, Bennett A Landman, Jennifer Urbano Blackford, Stephan Heckers, and Neil D Woodward. Thalamocortical anatomical connectivity in schizophrenia and psychotic bipolar disorder. *Schizophr Bull*, 46(5):1062–1071, Mar 2020. ISSN 1745-1701 (Electronic); 0586-7614 (Print); 0586-7614 (Linking). doi: 10.1093/schbul/sbaa022.

75. Suzanne N Avery, Anna S Huang, Julia M Sheffield, Baxter P Rogers, Simon Vandekar, Alan Anticevic, and Neil D Woodward. Development of thalamocortical structural connectivity in typically developing and psychosis spectrum youths. *Biol Psychiatry Cogn Neurosci Neuroimaging*, 7(8):782–792, Aug 2022. ISSN 2451-9030 (Electronic); 2451-9022 (Print); 2451-9022 (Linking). doi: 10.1016/j.bpsc.2021.09.009.

76. Nadia Blostein, Gabriel A. Devenyi, Sejal Patel, Raihaan Patel, Stephanie Tullo, Eric Plitman, Manuela Costantino, Ross Markello, Olivier Parent, Saashi A. Bedford, Chet C. Sherwood, William D Hopkins, Jakob Seidlitz, Armin Raznahan, and M. Mallar Chakravarty. Variation in subcortical anatomy: relating interspecies differences, heritability, and brain-behavior relationships. *bioRxiv*, page 2022.04.11.487874, 01 2022. doi: 10.1101/2022.04.11.487874.

77. Mingchao Yan, Wenwen Yu, Qian Lv, Qiming Lv, Tingting Bo, Xiaoyu Chen, Yilin Liu, Yafeng Zhan, Shengyao Yan, Xiangyu Shen, Baofeng Yang, Qiming Hu, Jiangli Yu, Zilong Qiu, Yuanjing Feng, Xiao-Yong Zhang, He Wang, Fuqiang Xu, and Zheng Wang. Mapping brain-wide excitatory projectome of primate prefrontal cortex at submicron resolution and comparison with diffusion tractography. *eLife*, 11:e72534, may 2022. ISSN 2050-084X. doi: 10.7554/eLife.72534.

78. Gleb Bezgin, Ana Solodkin, Rembrandt Bakker, Petra Ritter, and Anthony R. McIntosh. Mapping complementary features of cross-species structural connectivity to construct realistic "virtual brains". *Human Brain Mapping*, 38(4):2080–2093, 2017. doi: https://doi.org/10.1002/hbm.23506.

79. C R Dermon and H Barbas. Contralateral thalamic projections predominantly reach transitional cortices in the rhesus monkey. *J Comp Neurol*, 344(4):508–531, Jun 1994. ISSN 0021-9967 (Print); 0021-9967 (Linking). doi: 10.1002/cne.903440403.

80. Tolulope Adeyelu, Tanya Gandhi, and Charles C Lee. Crossed connections from insular cortex to the contralateral thalamus. *Front Neural Circuits*, 15:710925, 2021. ISSN 1662-5110 (Electronic); 1662-5110 (Linking). doi: 10.3389/fncir.2021.710925.

81. Madge E. Scheibel and Arnold B. Scheibel. Structural organization of nonspecific thalamic nuclei and their projection toward cortex. *Brain Research*, 6(1):60–94, 1967. ISSN 0006-8993. doi: https://doi.org/10.1016/0006-8993(67)90183-7. Forebrain Inhibitory Mechanisms.

82. Lianne H Scholtens, Rory Pijnenburg, Siemon C de Lange, Inge Huitinga, and Martijn P van den Heuvel. Common microscale and macroscale principles of connectivity in the human brain. *J Neurosci*, 42(20):4147–4163, May 2022. ISSN 1529-2401 (Electronic); 0270-6474 (Print). doi: 10.1523/JNEUROSCI.1572-21.2022.

83. Martijn P van den Heuvel and Olaf Sporns. Rich-club organization of the human connectome. *J Neurosci*, 31(44):15775–15786, Nov 2011. ISSN 1529-2401 (Electronic); 0270-6474 (Print); 0270-6474 (Linking). doi: 10.1523/JNEUROSCI.3539-11.2011.

84. Alexander Groh, Hajnalka Bokor, Rebecca A Mease, Viktor M Plattner, Balázs Hangya, Albrecht Stroh, Martin Deschenes, and László Acsády. Convergence of cortical and sensory driver inputs on single thalamocortical cells. *Cereb Cortex*, 24(12):3167–3179, Dec 2014. ISSN 1460-2199 (Electronic); 1047-3211 (Print); 1047-3211 (Linking). doi: 10.1093/cercor/bht173.

85. Qinglong L. Gu, Norman H. Lam, Michael M. Halassa, and John D. Murray. Circuit mechanisms of top-down attentional control in a thalamic reticular model. *bioRxiv*, page 2020.09.16.300749, 01 2020. doi: 10.1101/2020.09.16.300749.

86. Jorge Jaramillo, Jorge F. Mejias, and Xiao-Jing Wang. Engagement of pulvino-cortical feedforward and feedback pathways in cognitive computations. *Neuron*, 101(2):321 – 336.e9, 2019. ISSN 0896-6273. doi: https://doi.org/10.1016/j.neuron.2018.11.023.

87. Deanna J Greene, Scott Marek, Evan M Gordon, Joshua S Siegel, Caterina Gratton, Timothy O Laumann, Adrian W Gilmore, Jeffrey J Berg, Annie L Nguyen, Donna Dierker, Andrew N Van, Mario Ortega, Dillan J Newbold, Jacqueline M Hampton, Ashley N Nielsen, Kathleen B McDermott, Jarod L Roland, Scott A Norris, Steven M Nelson, Abraham Z Snyder, Bradley L Schlaggar, Steven E Petersen, and Nico U F Dosenbach. Integrative and network-specific connectivity of the basal ganglia and thalamus defined in individuals. *Neuron*, 105(4):742–758, Feb 2020. ISSN 1097-4199 (Electronic); 0896-6273 (Print); 0896-6273 (Linking). doi: 10.1016/j.neuron.2019.11.012.

88. Kai Hwang, Maxwell A Bertolero, William B Liu, and Mark D'Esposito. The human thalamus is an integrative hub for functional brain networks. *J Neurosci*, 37(23):5594–5607, Jun 2017. ISSN 1529-2401 (Electronic); 0270-6474 (Print); 0270-6474 (Linking). doi: 10.1523/JNEUROSCI.0067-17.2017.

89. Suzanne Haber and Nikolaus R. Mcfarland. The place of the thalamus in frontal cortical-basal ganglia circuits. *The Neuroscientist*, 7(4):315–324, 2001. doi: 10.1177/107385840100700408. PMID: 11488397.

90. Riccardo Beltramo and Massimo Scanziani. A collicular visual cortex: Neocortical space for an ancient midbrain visual structure. *Science*, 363(6422):64–69, Jan 2019. ISSN 1095-9203 (Electronic); 0036-8075 (Linking). doi: 10.1126/science.aau7052.

91. Robert H. Wurtz, Marc A. Sommer, and James Cavanaugh. Drivers from the deep: the contribution of collicular input to thalamocortical processing. In *Cortical Function: a View from the Thalamus*, volume 149 of *Progress in Brain Research*, pages 207–225. Elsevier, 2005. doi: https://doi.org/10.1016/S0079-6123(05)49015-9.

92. M. Wolff, S. Morceau, R. Folkard, J. Martin-Cortecero, and A. Groh. A thalamic bridge from sensory perception to cognition. *Neuroscience Biobehavioral Reviews*, 120:222–235, 2021. ISSN 0149-7634. doi: https://doi.org/10.1016/j.neubiorev.2020.11.013.

93. Klaus H. Maier-Hein, Peter F. Neher, Jean-Christophe Houde, Marc-Alexandre Côté, Eleftherios Garyfallidis, Jidan Zhong, Maxime Chamberland, Fang-Cheng Yeh, Ying-Chia Lin, Qing Ji, Wilburn E. Reddick, John O. Glass, David Qixiang Chen, Yuanjing Feng, Chengfeng Gao, Ye Wu, Jieyan Ma, Renjie He, Qiang Li, Carl-Fredrik Westin, Samuel Deslauriers-Gauthier, J. Omar Ocegueda González, Michael Paquette, Samuel St-Jean, Gabriel Girard, François Rheault, Jasmeen Sidhu, Chantal M. W. Tax, Fenghua Guo, Hamed Y. Mesri, Szabolcs Dávid, Martijn Froeling, Anneriet M. Heemskerk, Alexander Leemans, Arnaud Boré, Basile Pinsard, Christophe Bedetti, Matthieu Desrosiers, Simona Brambati, Julien Doyon, Alessia Sarica, Roberta Vasta, Antonio Cerasa, Aldo Quattrone, Jason Yeatman, Ali R. Khan, Wes Hodges, Simon Alexander, David Romascano, Muhamed Barakovic, Anna Auría, Oscar Esteban, Alia Lemkaddem, Jean-Philippe Thiran, H. Ertan Cetingul, Benjamin L. Odry, Boris Mailhe, Mariappan S. Nadar, Fabrizio Pizzagalli, Gautam Prasad, Julio E. Villalon-Reina, Justin Galvis, Paul M. Thompson, Francisco De Santiago Requejo, Pedro Luque Laguna, Luis Miguel Lacerda, Rachel Barrett, Flavio Dell'Acqua, Marco Catani, Laurent Petit, Emmanuel Caruyer, Alessandro Daducci, Tim B. Dyrby, Tim Holland-Letz, Claus C. Hilgetag, Bram Stieltjes, and Maxime Descoteaux. The challenge of mapping the human connectome based on diffusion tractography. *Nature Communications*, 8(1):1349, 2017. doi: 10.1038/s41467-017-01285-x.

94. Kurt G Schilling, Laurent Petit, Francois Rheault, Samuel Remedios, Carlo Pierpaoli, Adam W Anderson, Bennett A Landman, and Maxime Descoteaux. Brain connections derived from diffusion mri tractography can be highly anatomically accurate-if we know where white matter pathways start, where they end, and where they do not go. *Brain Struct Funct*, 225(8):2387–2402, Nov 2020. ISSN 1863-2661 (Electronic); 1863-2653 (Print); 1863-2653 (Linking). doi: 10.1007/s00429-020-02129-z.

95. Céline Delettre, Arnaud Messé, Leigh-Anne Dell, Ophélie Foubet, Katja Heuer, Benoit Larrat, Sebastien Meriaux, Jean-Francois Mangin, Isabel Reillo, Camino de Juan Romero, Victor Borrell, Roberto Toro, and Claus C Hilgetag. Comparison between diffusion mri tractography and histological tract-tracing of cortico-cortical structural connectivity in the ferret brain. *Netw Neurosci*, 3(4):1038–1050, 2019. ISSN 2472-1751 (Electronic); 2472-1751 (Linking). doi: 10.1162/netn{_}a{_}00098.

96. Chad J Donahue, Stamatios N Sotiropoulos, Saad Jbabdi, Moises Hernandez-Fernandez, Timothy E Behrens, Tim B Dyrby, Timothy Coalson, Henry Kennedy, Kenneth Knoblauch, David C Van Essen, and Matthew F Glasser. Using diffusion tractography to predict cortical connection strength and distance: A quantitative comparison with tracers in the monkey. *J Neurosci*, 36(25):6758–6770, Jun 2016. ISSN 1529-2401 (Electronic); 0270-6474 (Print); 0270-6474 (Linking). doi: 10.1523/JNEUROSCI.0493-16.2016.

97. Martijn P van den Heuvel, Marcel A de Reus, Lisa Feldman Barrett, Lianne H Scholtens, Fraukje M T Coopmans, Ruben Schmidt, Todd M Preuss, James K Rilling, and Longchuan Li. Comparison of diffusion tractography and tract-tracing measures of connectivity strength in rhesus macaque connectome. *Hum Brain Mapp*, 36(8):3064–3075, Aug 2015. ISSN 1097-0193 (Electronic); 1065-9471 (Print); 1065-9471 (Linking). doi: 10.1002/hbm.22828.

98. Giorgia Grisot, Suzanne N. Haber, and Anastasia Yendiki. Diffusion mri and anatomic tracing in the same brain reveal common failure modes of tractography. *NeuroImage*, 239: 118300, 2021. ISSN 1053-8119. doi: https://doi.org/10.1016/j.neuroimage.2021.118300.

99. Kurt G. Schilling, Yurui Gao, Iwona Stepniewska, Vaibhav Janve, Bennett A. Landman, and Adam W. Anderson. Anatomical accuracy of standard-practice tractography algorithms in the motor system - a histological validation in the squirrel monkey brain. *Magnetic Resonance Imaging*, 55:7–25, 2019. ISSN 0730-725X. doi: https://doi.org/10.1016/j.mri.2018.09.004.

100. Julien Dauguet, Sharon Peled, Vladimir Berezovskii, Thierry Delzescaux, Simon K. Warfield, Richard Born, and Carl-Fredrik Westin. Comparison of fiber tracts derived from in-vivo dti tractography with 3d histological neural tract tracer reconstruction on a macaque brain. *NeuroImage*, 37(2):530–538, 2007. ISSN 1053-8119. doi: https://doi.org/10.1016/j.neuroimage.2007.04.067.

101. Yurui Gao, Ann S Choe, Iwona Stepniewska, Xia Li, Malcolm J Avison, and Adam W Anderson. Validation of dti tractography-based measures of primary motor area connectivity in the squirrel monkey brain. *PLoS One*, 8(10):e75065, 2013. ISSN 1932-6203 (Electronic); 1932-6203 (Linking). doi: 10.1371/journal.pone.0075065.

102. Cibu Thomas, Frank Q Ye, M Okan Irfanoglu, Pooja Modi, Kadharbatcha S Saleem, David A Leopold, and Carlo Pierpaoli. Anatomical accuracy of brain connections derived from diffusion mri tractography is inherently limited. *Proc Natl Acad Sci U S A*, 111(46): 16574–16579, Nov 2014. ISSN 1091-6490 (Electronic); 0027-8424 (Print); 0027-8424 (Linking). doi: 10.1073/pnas.1405672111.

103. Kai Hwang, James M Shine, Joel Bruss, Daniel Tranel, and Aaron Boes. Neuropsychological evidence of multi-domain network hubs in the human thalamus. *eLife*, 10:e69480, oct 2021. ISSN 2050-084X. doi: 10.7554/eLife.69480.

104. Alan Anticevic, Genevieve Yang, Aleksandar Savic, John D Murray, Michael W Cole, Grega Repovs, Godfrey D Pearlson, and David C Glahn. Mediodorsal and visual thalamic connectivity differ in schizophrenia and bipolar disorder with and without psychosis history. *Schizophr Bull*, 40(6):1227–1243, Nov 2014. ISSN 1745-1701 (Electronic); 0586-7614 (Print); 0586-7614 (Linking). doi: 10.1093/schbul/sbu100.

105. Alan Anticevic, Kristen Haut, John D Murray, Grega Repovs, Genevieve J Yang, Caroline Diehl, Sarah C McEwen, Carrie E Bearden, Jean Addington, Bradley Goodyear, Kristin S Cadenhead, Heline Mirzakhanian, Barbara A Cornblatt, Doreen Olvet, Daniel H Mathalon, Thomas H McGlashan, Diana O Perkins, Aysenil Belger, Larry J Seidman, Ming T Tsuang, Theo G M van Erp, Elaine F Walker, Stephan Hamann, Scott W Woods, Maolin Qiu, and Tyrone D Cannon. Association of thalamic dysconnectivity and conversion to psychosis in youth and young adults at elevated clinical risk. *JAMA Psychiatry*, 72(9):882–891, Sep 2015. ISSN 2168-6238 (Electronic); 2168-622X (Print); 2168-622X (Linking). doi: 10.1001/jamapsychiatry.2015.0566.

106. John D Murray and Alan Anticevic. Toward understanding thalamocortical dysfunction in schizophrenia through computational models of neural circuit dynamics. *Schizophr Res*, 180:70–77, Feb 2017. ISSN 1573-2509 (Electronic); 0920-9964 (Print); 0920-9964 (Linking). doi: 10.1016/j.schres.2016.10.021.

107. Neil D Woodward, Haleh Karbasforoushan, and Stephan Heckers. Thalamocortical dysconnectivity in schizophrenia. *Am J Psychiatry*, 169(10):1092–1099, Oct 2012. ISSN 1535-7228 (Electronic); 0002-953X (Print); 0002-953X (Linking). doi: 10.1176/appi.ajp.2012.12010056.

108. Neil D Woodward and Stephan Heckers. Mapping thalamocortical functional connectivity in chronic and early stages of psychotic disorders. *Biol Psychiatry*, 79(12):1016–1025, Jun 2016. ISSN 1873-2402 (Electronic); 0006-3223 (Print); 0006-3223 (Linking). doi: 10.1016/j.biopsych.2015.06.026.

109. Wu Jeong Hwang, Yoo Bin Kwak, Kang Ik K Cho, Tae Young Lee, Harin Oh, Minji Ha, Minah Kim, and Jun Soo Kwon. Thalamic connectivity system across psychiatric disorders: Current status and clinical implications. *Biol Psychiatry Glob Open Sci*, 2(4):332–340, Oct 2022. ISSN 2667-1743 (Electronic); 2667-1743 (Linking). doi: 10.1016/j.bpsgos.2021.09.008.

110. Peter T. Bell and James M. Shine. Subcortical contributions to large-scale network communication. *Neuroscience Biobehavioral Reviews*, 71:313–322, 2016. ISSN 0149-7634. doi: https://doi.org/10.1016/j.neubiorev.2016.08.036.

111. James M. Shine. The thalamus integrates the macrosystems of the brain to facilitate complex, adaptive brain network dynamics. *Progress in Neurobiology*, 199:101951, 2021. ISSN 0301-0082. doi: https://doi.org/10.1016/j.pneurobio.2020.101951.

112. J Bourassa, D Pinault, and M Deschênes. Corticothalamic projections from the cortical barrel field to the somatosensory thalamus in rats: a single-fibre study using biocytin as an anterograde tracer. *Eur J Neurosci*, 7(1):19–30, Jan 1995. ISSN 0953-816X (Print); 0953-816X (Linking). doi: 10.1111/j.1460-9568.1995.tb01016.x.

113. Matthew F. Glasser, Timothy S. Coalson, Emma C. Robinson, Carl D. Hacker, John Harwell, Essa Yacoub, Kamil Ugurbil, Jesper Andersson, Christian F. Beckmann, Mark Jenkinson, Stephen M. Smith, and David C. Van Essen. A multi-modal parcellation of human cerebral cortex. *Nature*, 536(7615):171–178, 2016. doi: 10.1038/nature18933.

114. B T Thomas Yeo, Fenna M Krienen, Jorge Sepulcre, Mert R Sabuncu, Danial Lashkari, Marisa Hollinshead, Joshua L Roffman, Jordan W Smoller, Lilla Zöllei, Jonathan R Polimeni, Bruce Fischl, Hesheng Liu, and Randy L Buckner. The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J Neurophysiol*, 106(3):1125–1165, Sep 2011. ISSN 1522-1598 (Electronic); 0022-3077 (Print); 0022-3077 (Linking). doi: 10.1152/jn.00338.2011.

115. Ye Tian, Daniel S Margulies, Michael Breakspear, and Andrew Zalesky. Topographic organization of the human subcortex unveiled with functional connectivity gradients. *Nat Neurosci*, 23(11):1421–1432, Nov 2020. ISSN 1546-1726 (Electronic); 1097-6256 (Linking). doi: 10.1038/s41593-020-00711-6.