Improving PD-1 blockade plus chemotherapy for complete remission of lung cancer

by nanoPDLIM2

Fan Sun[1], Pengrong Yan[1], Yadong Xiao[1, 2, 3, 4], Hongqiao Zhang[3],

Steven D. Shapiro[2, 4], Gutian Xiao[1, 3, *], and Zhaoxia Qu[1, 3, *]

[1]UPMC Hillman Cancer Center, Department of Microbiology and Molecular Genetics, University of Pittsburgh School of Medicine, Pittsburgh, PA 15213, USA. [2]Division of Pulmonary, Allergy, and Critical Care Medicine, Department of Medicine, University of Pittsburgh School of Medicine, Pittsburgh, PA 15213, USA. [3]Norris Comprehensive Cancer Center, Hastings Center for Pulmonary Research, Department of Molecular Microbiology and Immunology, University of Southern California Keck School of Medicine, Los Angeles, CA 90033, USA. [4]Department of Medicine, University of Southern California Keck School of Medicine, Los Angeles, CA 90033, USA.

[*]Correspondence should be addressed to Zhaoxia Qu (Email: Zhaoxia.Qu@med.usc.edu; Phone: 323-442-6932) or Gutian Xiao (Email: Gutian.Xiao@med.usc.edu; Phone: 323-442-7985)

Running Title: NanoPDLIM2-chemo-immuno-therapy in lung cancer

Keywords: PD-1; PDLIM2; immune checkpoint; ICIs; nanotherapy; combination therapy; lung cancer; preclinical studies; mouse models

Disclosure of Potential Conflicts of Interest: The authors declare no potential conflicts of interest.

Abstract

Background: Immune checkpoint inhibitors (ICIs) and their combination with other therapies such as chemotherapy, fail in most cancer patients. We previously identified the PDZ-LIM domain-containing protein 2 (PDLIM2) as a *bona fide* tumor suppressor that is repressed in lung cancer to drive cancer and its chemo and immunotherapy resistance, suggesting a new target for lung cancer therapy improvement.

Methods: Human clinical samples and data were used to investigate *PDLIM2* genetic and epigenetic changes in lung cancer. Using an endogenous mouse lung cancer model faithfully recapitulating refractory human lung cancer and a clinically feasible nano-delivery system, we investigated the therapeutic efficacy, action mechanism, and safety of systemically administrated PDLIM2 expression plasmids encapsulated in nanoparticles (nanoPDLIM2) and its combination with PD-1 antibody and chemotherapeutic drugs.

Results: PDLIM2 repression in human lung cancer involves both genetic deletion and epigenetic alteration. NanoPDLIM2 showed low toxicity, high tumor specificity, antitumor activity, and greatly improved the efficacy of anti-PD-1 and chemotherapeutic drugs, with complete tumor remission in most mice and substantial tumor reduction in the remaining mice by their triple combination. Mechanistically, nanoPDLIM2 increased major histocompatibility complex class I (MHC-I) expression, suppressed multi-drug resistance 1 (MDR1) induction and survival genes and other tumor-related genes expression in tumor cells, and enhanced lymphocyte tumor infiltration, turning the cold tumors hot and sensitive to ICIs and rendering them vulnerable to chemotherapeutic drugs and activated tumor-infiltrating lymphocytes (TILs) including those unleashed by ICIs.

Conclusions: These studies established a clinically applicable PDLIM2-based combination therapy with great efficacy for lung cancer and possibly other cold cancers.

Introduction

Lung cancer is the leading cause of cancer-related deaths in both men and women with a 5-year survival rate of only 22% [1]. Although these outcomes may be improved by immune checkpoint blockade therapy involving the disruption of the binding of programmed cell death 1 (PD-1, also known as CD279) on tumor-infiltrating lymphocytes (TILs) to programmed death receptor ligand 1 (PD-L1, also known as B7-H1 or CD274) on tumor and tumor-associated cells, most lung cancer patients still fail the therapy, with a response rate of only about 20% [2]. In general, this revolutionary immunotherapy works better against 'hot' tumors, which have abundant TILs, strong immunogenicity and sufficient PD-L1 expression. Unfortunately, most tumors are 'cold', with low T-cell infiltration, weak immunogenicity and minimal PD-L1 expression, and show weak response to immune checkpoint inhibitors (ICIs) [3, 4].

Cold lung tumors without targetable oncogenic drivers are treated with chemotherapy as the standard approach [5]. However, the response rate to this conventional cancer therapy is also low and resistance often occurs after an initial response, with an overall survival (OS) of about 12-17 months. Given its roles in inducing TILs and immunogenicity, in particular PD-L1 expression, to turn cold tumors hot, chemotherapy can be an ideal candidate for combination with PD-1/PD-L1 blockade to improve therapeutic efficacy [6]. Indeed, combination treatment with ICIs and chemotherapeutic drugs shows synergy and better efficacy in both preclinical animal models and clinical trials of lung cancer and other cancers [6-8]. However, even with the combination of ICIs and chemotherapy, tumors in animals do not remit completely and the objective response rate (ORR) of lung cancer patients only reaches 33%-49.7%, with a median progression-free survival (PFS) of just 5.1-9 months and a median OS of 13-22 months [7, 8]. Thus, further improvement over the chemo-immunotherapy is direly needed.

Our recent human and mouse studies have shown that most lung tumors not only have low TILs and

decreased PD-L1, but also down-regulate major histocompatibility complex class I (MHC-I), evading recognition and attack by CD8[+] T cells, including those unleashed by ICIs and/or recruited by chemotherapy [6, 9, 10]. Following our previous cell line studies [11-17], we have established the PDZ-LIM domain-containing protein PDLIM2, also known as SLIM or mystique [18-20], as a *bona fide* tumor suppressor and its repression as a causative driver of lung cancer and resistance to ICIs and chemotherapeutic agents [6]. While PDLIM2 is epigenetically repressed in human lung cancer, associating with therapeutic resistance and poor prognosis, its global or lung epithelial-specific deletion in mice leads to lung cancer development, chemoresistance, and complete resistance to anti-PD-1 and epigenetic drugs. One most important function of PDLIM2 is to promote the ubiquitination and proteasomal degradation of nuclear signal transducer and activator of transcription 3 (STAT3) and nuclear factor-κB (NF-κB) RelA (also known as p65), two master transcription factors that function as proto-oncogenes in lung and many other cancers [6, 10-13, 21-27]. PDLIM2 repression in tumor cells thus results in the persistent activation of STAT3 and RelA, leading to MHC-I downregulation and high expression of tumor growth-related genes [6]. It also leads to strong induction of multi-drug resistance 1 (MDR1) for acquired chemo-resistance, as chemotherapy further enhances RelA activation for MDR1 transcription in PDLIM2 deficient tumor cells [6].

In this study, we examined whether and how PDLIM2 can be targeted to treat lung cancer in a faithful mouse model of human lung cancer. In particular, we tested whether systemic administration of nanoparticle-encapsulated PDLIM2-expression plasmids (nanoPDLIM2) could enhance the efficacy of anti-PD-1 and/or chemotherapeutic drugs. We also examined whether PDLIM2 repression in human lung cancer involves genetic deletion and its relationship with epigenetic silencing. These studies indicate that besides epigenetic repression, loss of heterozygosity (LOH) contributes to *PDLIM2* downregulation in about 58% of human lung tumors, and that *Pdlim2* heterozygous deletion (*Pdlim2*[+/-]) mice develop spontaneous tumors in lung and other organs. Notably, systemic administration of

100   nanoPDLIM2 reverses the phenotypes caused by PDLIM2 repression and induces complete remission of

101   all lung tumors in most mice without further increasing toxicity when combined with anti-PD-1 and

102   chemotherapeutic drugs. These findings provide a firm basis to combine ICIs and chemotherapeutic

103   drugs with PDLIM2-targeted therapy for the treatment of lung and other cancers.

Material and Methods

Animals and lung carcinogenesis

We have complied with all relevant ethical regulations for animal testing and research. The animal

experiments were performed in accordance with the US National Institutes of Health (NIH) Guidelines

on the Use of Laboratory Animals. All animals were maintained under pathogen-free conditions and

used according to protocols approved by the Institutional Animal Care and Use Committee (IACAUC)

of the University of Pittsburgh. *Pdlim2*$^{flx/flx}$/*SP-C*-rtTA$^{tg/-}$/(tetO)7CMV-Cre$^{tg/tg}$ (ΔSPC) mice under a

pure FVB/N background and *Pdlim2*$^{-/-}$ mice under a pure BALB/c background have been described

before [6, 19, 21, 22, 28]. For lung carcinogenesis, six- to eight-week-old ΔSPC and wild type FVB/N

mice were intraperitoneally (i.p.) injected with urethane (1 mg/g body weight, Sigma-Aldrich, St. Louis,

MO, USA) once a week for six consecutive weeks [6], followed by different treatments as shown in the

figures. Mice were euthanized at six weeks post urethane treatment for examination of lung tumors,

inflammation, and treatment-induced toxicity. Surface tumors in mouse lungs were counted blinded

under a dissecting microscope, and tumor diameters were measured by microcalipers.

Preparation of PDLIM2-expression plasmid or empty vector plasmid nanoparticles

The polyethylenimine (PEI)-based nanoparticles (*in vivo*-jetPEI) (Polyplus Transfection, New York,

NY, USA) and plasmid DNA complexes at a nitrogen-to-phosphate ratio of 8 (N/P=8) were prepared

according to the manufacturer's instructions. Briefly, 25□µg of pCMV-myc-*Pdlim2* or empty vector

plasmids in 100 μl of a 5% glucose solution were mixed with the *in vivo*-jetPEI reagent (4 μl) diluted

into 100 μl of a 5% glucose solution. After 15 minutes of incubation at room temperature, the mixed

solution (200 μl/mouse) was injected intravenously (i.v.) via the tail vein. PDLIM2-expression plasmid

or empty vector plasmid DNA in tissues were measured by PCR assays targeting Amp-R in the genome

of these plasmids.

Histology and immunohistochemistry (IHC) analysis

Lung, liver, kidney, and spleen tissues were excised, fixed in formalin, embedded in paraffin, and cut

into 4-μm-thick sections. Sections were stained with H&E or subjected to sequential incubations with

the indicated primary antibodies, biotinylated secondary antibodies and streptavidin-HRP [6].

Antibodies were listed in Supplementary Table S1.

***In vivo* BrdU labeling**

Mice were i.p. injected with 50 mg/kg BrdU (Sigma-Aldrich, St. Louis, MO, USA) 24 h prior to

euthanasia. Mouse lung tissue sections were stained with anti-BrdU (Sigma-Aldrich, St. Louis, MO,

USA). BrdU labeling index was calculated as the percentage of labeled cells per total cells counted (>

500 cells in each counted tumor-containing area).

Flow cytometry (FACS) analysis

The cells were isolated from mouse lungs using collagenase/dispase digestion followed by filtration with

70 μm Nylon cell strainer and red blood cell lysis; then the cells were incubated with the antibodies

against cell surface antigens after blocked with αCD16/CD32. The cells were then fixed with

paraformaldehyde (2%), permeabilized and incubated with antibodies against intracellular antigens if

needed. For interferon-γ (IFNγ) staining, cells were treated with phorbol 12-myristate 13-acetate (PMA,

50 ng/ml), ionomycin (1 μM), brefeldin A (BFA, 3 μg/ml) and monensin (2 μM) for 4 h before

they were stained for FACS analysis. Data were acquired and analyzed by Accuri C6 or LSRFortessa I

(BD Biosciences) and FlowJo software [29].

Quantitative polymerase chain reaction (qPCR) analysis

The indicated tissues or cells were subjected to DNA or RNA extraction, RNA reverse transcription and

real-time PCR using trizol, reverse transcriptase, and Power SYBR Green PCR Master Mix (Thermo

Fisher Scientific, Waltham, MA USA) according to manufacturer's protocol. Primer pairs used for

qPCR were listed in Supplementary Table S2.

Microsatellite and gene-specific PCR-based LOH analysis of *PDLIM2*

Genomic DNAs were isolated from human lung tumors and their matched normal tissues using the

PureLink Genomic DNA Purification Kit (Invitrogen, Carlsbad, CA, USA), and subjected to semi-

quantitative PCR using the primers specific for the microsatellite markers D8S1786 and D8S1752 that

straddle the *PDLIM2* genetic locus or the *PDLIM2* genetic locus itself. Primer pairs used for the assays

were listed in Supplementary Table S2.

Statistical analysis

One-way ANOVA power analysis was used to determine the minimum sample size. Animals were

randomly assigned to different treatment groups. Measurements were taken from distinct samples.

Student's t test (two tailed) and one-way ANOVA/Tukey's or two-way ANOVA/Sidak's test were used

to assess significance of differences between two groups and multiple comparisons, respectively.

Pearson's correlation test was used to assess association between *PDLIM2* expression with its promoter

methylation or genetic deletion and the overlap between *PDLIM2* promoter methylation and genetic

deletion. All bars in figures represent means $\pm$ standard error of the mean (SEM). The *p* values are

indicated as $*p < 0.05$, $**p < 0.01$, ns, not statistically significant, except for those shown in figures.

The *p* values □<□0.05 and 0.01 are considered statistically significant and highly statistically significant,

respectively. Grubbs' and ROUT outlier tests were established a priori. There were no exclusions in the

analyses.

Results

Both epigenetic repression and genetic deletion of *PDLIM2* in lung cancer

Using 40% of the expression level in matched normal lung tissues as the cut-off in the analysis of the

Cancer Genome Atlas (TCGA) data, we found that *PDLIM2* was repressed in over 75% of human lung

tumors in a prior study [6]. If using 50% as the cut-off, *PDLIM2* was repressed in about 93% of human

lung tumors (Figure 1*A*). Using 125% of the methylation level of the *PDLIM2* promoter in normal lung

tissues as the cut-off to analyze TCGA database, over 70% of human lung tumors were found to have

hypermethylation of the *PDLIM2* promoter (Figure 1*A*, 1*B*), further validating our previous finding of

epigenetic silencing as the main mechanism underlying PDLIM2 repression [6].

In line with *PDLIM2*'s location on chromosome 8p21.3, a frequent LOH region in lung and other tumors

[30-34], analysis of TCGA database revealed that *PDLIM2* expression was positively associated with its

gene copy numbers, and that over 58% of human lung tumors had genetic deletion of the *PDLIM2* gene

if copy number variation (CNV) of -0.1 was used as the cut-off for the gene deletion (Figure 1*C*).

Further analysis indicated that about 44% of human lung tumors simultaneously harbored the promoter

hypermethylation and LOH of the *PDLIM2* gene, and approximately 26% and 14% of them only having

the promoter hypermethylation or LOH of the *PDLIM2* gene, respectively (Figure 1*D*). Around 16% of

lung tumors possessed no such epigenetic or genetic alterations in *PDLIM2*. These findings were

confirmed by microsatellite and gene-specific PCR-based LOH analysis of human primary lung tumor

tissues and cell lines (Figure 1*E*, 1*F*; Supplementary Table S3). These data suggested that PDLIM2

repression in lung cancer involves both epigenetic silencing and genetic deletion.

To determine the significance of *PDLIM2* LOH in lung tumors, we examined whether *Pdlim2*
heterozygous deletion leads to decreased PDLIM2 expression and spontaneous tumors in mice. Unlike
its absolute absence in *Pdlim2* homozygous deletion (*Pdlim2*$^{-/-}$) mice, PDLIM2 was detected in the lung
of *Pdlim2*$^{+/-}$ mice, but at a much lower level compared to wild type (WT) mice (Supplementary Figure
S1*A*). Importantly, *Pdlim2*$^{+/-}$ mice, like *Pdlim2*$^{-/-}$ mice, also developed spontaneous tumors (Figure 1*G*).
Of note, over 50% of tumors developed in *Pdlim2*$^{-/-}$ and *Pdlim2*$^{+/-}$ mice were lung tumors (Figure 1*H*;
Supplementary Figure S1*B*). These data are also highly consistent with the fact that PDLIM2 is
ubiquitously expressed under physiological conditions, with the highest level in the lung and lung
epithelial cells in particular [6, 18-20]. Thus, PDLIM2 is a haploinsufficient tumor suppressor that is
particularly important for lung tumor suppression.

Efficacy of systemic administration of nanoPDLIM2 in refractory lung cancer

Although reversal of PDLIM2 epigenetic repression by epigenetic drugs to restore PDLIM2 expression
in cancer cells may be used to treat lung cancer [6], it is logical that this approach cannot be applied to
lung tumors involving *PDLIM2* LOH, which accounts for about 58% of all lung cancer cases. To
overcome this limitation and expand PDLIM2-targeted therapy to all lung tumors with PDLIM2
repression regardless of the involved mechanisms, we tested the therapeutic efficacy of systemic
administration of PDLIM2-expression plasmids encapsulated by the clinically feasible *in vivo*-jetPEI
[35-39]. To this end, we employed mouse lung tumors induced by urethane, a faithful model of human
lung cancer and adenocarcinoma (AC) in particular [6, 23, 24, 40-43]. Urethane is a chemical
carcinogen present in fermented food, alcoholic beverage and also cigarette smoke, the predominant risk
factor accounting for about 90% of human lung cancer cases [44]. Like its human counterpart, the
murine lung cancer induced by urethane also shares PDLIM2 repression, in addition to their similarities
in histology, genetics, molecular biology and immunology [6, 9, 23, 24, 40-43]. WT mice with lung

tumors induced by urethane were i.v. injected with nanoparticle-encapsulated *Pdlim2* plasmids or empty

vector plasmids. Six weeks post the initial treatment of nanotherapy, mice were euthanized and lung

tissues were collected (Figure 2*A*). Compared to the vector plasmid mock-treated group (Vec),

nanoPDLIM2-treated mice had a significantly reduced tumor burden in their lungs, although the

decrease in tumor numbers was not statistically significant (Figure 2*B*).

In line with the tumor reduction, nanoPDLIM2 administration decreased nuclear RelA and STAT3, a

hallmark of NF-κB and STAT3 activation, accordingly reduced the expression of their downstream cell

survival gene Bcl-xL and cell proliferation gene Cyclin D1, increased apoptosis and decreased

proliferation of lung cancer cells (Figure 2*C*-2*E*). These data indicated that intravenous administration

of nanoPDLIM2 shows efficacy in suppressing oncogenic RelA and STAT3 activation and in treating

lung cancer.

High tumor specificity and low toxicity of systemic administration of nanoPDLIM2

To characterize the potential new lung cancer therapy, we examined the expression levels of PDLIM2 in

lung tumors and several organs, including the liver, kidney, and spleen. Consistent with the treatment

efficacy, a high level of PDLIM2 was detected in the lung tumors from mice treated with the PDLIM2

plasmid nanoparticles one week post nanoPDLIM2 treatment, whereas no obvious PDLIM2 was found

in the lung tumors from mice treated with the control plasmid nanoparticles (Figure 3*A*, 3*B*). It should

be pointed out that the tumor delivery efficiency of PDLIM2-expression or control plasmid

nanoparticles was similarly high (Figure 3*B*). However, either PDLIM2-expression or control plasmids

were hardly detected in other organs/tissues of the same mice, including liver, kidney and spleen. Of

note, in the first two days after injection, both empty vector and PDLIM2-expression plasmids were also

detected at high levels in those tissues but quickly cleared afterward (data not shown), consistent with

the well-documented tumor-specific enrichment of nanoparticles [37, 45-47]. Consistently, ectopic

Pdlim2 was not detected, and the levels of PDLIM2 proteins were comparable in these tissues (Figure

3*B*, 3*C*).

In further support of the lung tumor-specific delivery, the i.v. injection of either PDLIM2-expression or

control plasmid nanoparticles showed no obvious toxicity to animals, as evidenced by no significant

changes in the animal body weight and histology of all organs/tissues we examined, including spleen,

kidney and liver (Figure 3*D*, *E*). Mouse appearance and behaviors, such as eating, drinking, defecating,

urinating, sniffing, grooming, and digging, were not different between PDLIM2-expression and control

plasmid nanoparticle groups (data not shown). Taken together, these data suggested the therapeutic

efficacy and low toxicity of PDLIM2-based nanotherapy in the mouse model of refractory lung cancer.

Synergy of nanoPDLIM2 with chemotherapy in lung cancer treatment

Given the role of PDLIM2 in inhibiting the expression of cell survival and proliferation genes in tumor

cells, which contribute to chemoresistance, we tested whether nanoPDLIM2 increases the efficacy of

chemotherapy in the mouse model of lung cancer. Treatment with carboplatin and paclitaxel, two

chemotherapeutic drugs that are often used together as the first-line treatment for lung and many other

cancers, led to significant decrease in tumor number and tumor burden (Figure 4*A*). Importantly,

combination with nanoPDLIM2 further significantly decreased both tumor number and tumor burden,

suggesting a promising synergy between PDLIM2 nanotherapy and chemotherapy in lung cancer

treatment. Consistently, significantly higher tumor cell apoptosis was detected in mice treated with the

combination therapy, in comparison to those with nanoPDLIM2 or chemotherapy alone (Figure 4*B*).

Another important mechanism contributing to the synergy between these two therapies involves

nanoPDLIM2 blockade of the acquired chemoresistance of lung cancer cells. Chemotherapy induced

strong RelA activation/nuclear expression and MDR1 expression (Figure 4*C*, 4*D*). NanoPDLIM2 not

only repressed the constitutive activation of RelA in cancer cells but also prevented the strong induction

of RelA activation and MDR1 expression by the chemotherapy. Thus, PDLIM2 nanotherapy improves

the therapeutic efficacy of chemotherapy through blocking both intrinsic and acquired chemoresistance

of lung cancer cells.

279 NanoPDLIM2 enhancement of ICI's efficacy in lung cancer treatment

High expression of cell survival genes also renders tumor cells resistant to the tumoricidal activity of

cytotoxic T lymphocytes (CTLs), including those unleashed by ICIs. PDLIM2 nanotherapy should also

enhance the efficacy of immunotherapy, given its ability in suppressing the expression of cell survival

genes. Furthermore, NanoPDLIM2 increased the number of TILs and the expression of MHC-I, the

most important and core components of immunotherapies including PD-1 immune checkpoint blockade

therapy (Figure 5*A*, 5*B*). In line with our previous studies [6], PD-1 blocking antibody showed some

efficacies in the mouse model of lung cancer, as evidenced by the significant decrease in tumor burden

and tumor number (Figure 5*C*). Combining with PDLIM2 nanotherapy further significantly decreased

tumor burden, although the decrease in tumor number failed to reach statistical significance.

Consistently, significant increases in both CD4$^+$ and CD8$^+$ TILs, CD8$^+$ CTL activation and lung tumor

cell death were detected (Figure 5*D*-5*F*).

292 Complete remission of all lung tumors in most mice by the combination treatment of

293 nanoPDLIM2 and anti-PD-1 and chemotherapeutic drugs

Like most human lung tumors, lung tumors in our animal model exhibit decreased expression of PD-L1

on cell surface, in additional to the low expression of MHC-I and low number of TILs [6, 9]. In line

with our previous finding that PD-L1 expression is largely independent of PDLIM2 [6], nanoPDLIM2

failed to induce PD-L1 expression in lung tumors, which was in contrast to chemotherapy (Figure 6*A*).

This may explain why the enhancing effect of nanoPDLIM2 on PD-1 blockade therapy is only

moderate.

Although it dramatically increases TILs as well [6], chemotherapy also induced the expression of PD-L1

on cancer cells (Figure 6A), which presumably protects cancer cells from immune attack and thereby

restricts further efficacy improvement of its combination with PDLIM2 nanotherapy.   But it may

increase the sensitivity of cancer cells to PD-1 blockade therapy, particularly in combination with

nanoPDLIM2.   Indeed, chemotherapy and PD-1 blockade therapy showed a promising synergy in

reducing both tumor number and tumor burden in comparison to the individual treatment of

chemotherapy or PD-1 blockade therapy [6].   The synergy was largely blocked when *Pdlim2* was

genetically deleted from lung cancer cells (ΔSPC) (Figure 6B, 6C), suggesting an important role of

PDLIM2 in the combination therapy of chemotherapeutic drugs and anti-PD-1.   Whereas PDLIM2

nanotherapy increased MHC-I expression on tumor cells (Figure 5B), which is critical for better

recognition and killing of tumor cells by CD8$^+$ CTLs, chemotherapy alone or its combination with anti-

PD-1 failed to do so (Figure 6D).   Accordingly, combination of anti-PD-1 and chemotherapeutic drugs,

like the combination therapies of nanoPDLIM2 and anti-PD-1 or chemotherapeutic drugs, also failed to

induce a complete remission of lung tumors in any mice.

Given their overlapping roles in increasing TILs and in particular their complement roles in inducing

MHC-I and PD-L1 expression on tumor cells, which turn tumors hot and more sensitive to PD-1

blockade, combination of nanoPDLIM2 and chemotherapeutic drugs with anti-PD-1 is expected to show

better efficacy compared to combining only two of them.   Indeed, combination of all three showed the

strongest effect on reducing tumor burden and numbers, and caused complete remission of all lung

tumors in 60% of mice (Figure 6E) and much more shrinkage of lung tumors in remaining mice in

comparison to those treated with two combined therapies.   In line with the high therapeutic efficacy, the

triple combination therapy significantly increased the numbers and/or activation of CD4$^+$ and CD8$^+$ T

cells in the lung, compared to the combination of chemotherapeutic drugs and anti-PD-1 (Figure *6F*,

6G).

Of note, consistent with the undetectable toxicity of PDLIM2 nanotherapy, its co-treatment did not

further increase the toxicity of anti-PD-1 and chemotherapeutic drugs.   There was no obvious

histological difference of major organs, including the liver, lung, kidney, and spleen (Supplementary

Figure S2). Moreover, no significant differences in animal body weights were observed by additional

nanoPDLIM2, in comparison to the mice received the combinational treatment of anti-PD-1 and

chemotherapeutic drugs in the presence or absence of empty vector plasmid nanoparticles (Figure *6H*).

These data suggested a novel combination therapy with very high therapeutic efficacy and no increased

toxicity for lung cancer, particularly refractory lung cancer.

Discussion

PD-1/PD-L1 blockade immunotherapy has recently joined chemotherapy as a standard treatment for

lung and several other cancers [2-4]. While some patients have shown dramatic responses, most patients

do not benefit from this novel treatment. Currently, various combinations of ICIs with other therapies,

in particular its combination with chemotherapeutic drugs are being extensively tested in both

preclinical and clinical trial studies to expand the benefit of this innovative therapy [6-8]. Although a

promising synergy and better efficacy has been observed in both preclinical animal models and human

clinical trial studies [6-8], significant further improvement is direly needed. Using an authentic mouse

model of lung cancer, we show, for the first time, that PDLIM2 nanotherapy shows efficacy and high

safety, and more importantly, induces complete remission of all lung tumors in most animals when it is

combined with anti-PD-1 and chemotherapeutic drugs.

Most human lung tumors as well as lung tumors in our animal model have low numbers of TILs and

decreased expression of PD-L1 and MHC-I on the cell surface [6, 9], all of which are important

mechanisms contributing to the resistance to PD-1 blockade therapy. Through inducing immunogenic

cell death (ICD) of cancer cells [8], chemotherapy can increase TILs and PD-L1 expression on tumor

cells, and thereby synergize with anti-PD-1. However, chemotherapy cannot induce MHC-I expression,

which limits further improvement of its synergy with immune therapy for complete cancer remission.

Also, tumor cells usually express high levels of survival genes against the tumoricidal effects of

chemotherapeutic drugs and of CTLs, including those activated by chemotherapy and unleashed by ICIs.

On the other hand, PDLIM2 nanotherapy induces MHC-I expression and lymphocyte tumor infiltration

but does not up-regulate PD-L1. Moreover, PDLIM2 nanotherapy prevents the induction of MDR1 and

the expression of tumor-related genes and in particular cell survival genes, further sensitizing tumor

cells to the cytotoxicity of chemotherapeutic drugs and immune cells including those recruited by

chemotherapy and unleashed by PD-1 blockade. Because of these important functions of PDLIM2

nanotherapy, PDLIM2 nanotherapy improved the efficacy of chemotherapy and PD-1 blockade therapy,

and in combination with chemotherapy and PD-1 blockade therapy, resulted in complete cancer

remission in most of the animals and dramatic tumor reduction in the remaining mice.

Another important clinical characteristic of PDLIM2 nanotherapy is its tumor-specificity and high safety

profile. It delivered PDLIM2-expression plasmids to lung tumor tissues and showed undetectable

toxicity in the animal model. Its combination does not further increase the toxicity of anti-PD-1 and

chemotherapeutic drugs. This is in sharp contrast to the Food and Drug Administration (FDA)-approved

epigenetic drugs, which can restore PDLIM2 expression in cancer cells with PDLIM2 epigenetic

repression [6, 10-15]. Epigenetic drug treatment leads to body weight loss of animals (Supplementary

Figure S3*A*).   The toxicity is much worse when epigenetic drugs are combined with chemotherapy

(Supplementary Figure S3).

Besides its side effects, epigenetic therapy should have much more limited application, even though it

shows better efficacy as a monotherapy or combined with chemotherapeutic drugs or anti-PD-1 in

comparison to PDLIM2 nanotherapy (Supplementary Figure S4).   While nanoPDLIM2-based

combination therapies could be applicable to all lung tumors with PDLIM2 repression regardless of the

mechanisms involved, epigenetic therapy may be used to treat about 26% of lung tumors with PDLIM2

epigenetic repression only.   About 58% of lung tumors harboring PDLIM2 LOH are not suitable to

epigenetic therapies, although most of them are also with epigenetic alterations of the *pdlim2* gene.   Of

note, the therapeutic efficacy of epigenetic therapy depends on PDLIM2 expression [6], and PDLIM2

heterozygous loss causes spontaneous lung and other cancers.   Further, the efficacy and toxicity of the

combination of epigenetic agents with both anti-PD-1 and chemotherapeutic drugs has not been

examined yet.

Conclusions

In summary, the presented data identify genetic deletions as a major mechanism other than epigenetic

alterations for PDLIM2 repression in human lung cancer, and PDLIM2 as a haploinsufficient tumor

suppressor particularly important for suppressing lung cancer and therapy resistance.   More importantly,

these preclinical data establish a novel combination treatment of nanoPDLIM2, anti-PD-1 and

chemotherapeutic drugs that induces complete remission of all lung tumors in most animals and is also

with high safety profile.   We believe that these knowledges are applicable to other cancers, because

PDLIM2 repression has also been linked to numerous human cancers other than lung cancer.

Abbreviations

AC: adenocarcinoma; BFA: brefeldin A; CNV: copy number variation; CTL: cytotoxic T lymphocyte; FACS: fluorescent activated cell sorting; FDA: Food and Drug Administration; IFNγ☐interferon-γ; ICD: immunogenic cell death; ICI: immune checkpoint inhibitor; IHC: immunohistochemistry; i.p.: intraperitoneal; i.v.: intravenous; LOH: loss of heterozygosity; MDR1: multi-drug resistance 1; MHC-I: major histocompatibility complex class I; nanoPDLIM2: nanoparticle-encapsulated PDLIM2-expression plasmids; NF-κB: nuclear factor-κB; NIH: National Institutes of Health; ORR: objective response rate; OS: overall survival; PD-1: programmed cell death 1; PD-L1: programmed death ligand 1; PD-L2: programmed death ligand 2; PDLIM2: PDZ-LIM domain-containing protein 2; PEI: polyethylenimine; PFS: progression-free survival; PMA: phorbol 12-myristate 13-acetate; qPCR: quantitative polymerase chain reaction; SEM: standard error of the mean; STAT3: signal transducer and activator of transcription 3; TCGA: The Cancer Genome Atlas; TIL: tumor-infiltrating lymphocyte; WT: wild type.

Acknowledgments

The authors thank Dr. M. J. Grusby (Harvard School of Public Health) and Dr. J. A. Whitsett (University of Cincinnati College of Medicine) for providing PDLIM2$^{-/-}$ and SP-C-rtTA$^{tg/-}$/(tetO)7CMV-Cre$^{tg/tg}$ mice, respectively. The authors also thank Dr. W. Ma and Dr. S. Li (University of Pittsburgh) for their suggestions on the nanoparticle preparation, and Dr. L. H. Rigatti (University of Pittsburgh) for her histological diagnosis of mouse tissues.

Funding

This study was financially supported in part by the NIH National Institute of General Medical Sciences (NIGMS) grant R01 GM144890, National Cancer Institute (NCI) grants R01 CA172090 and R01 CA258614, R21 CA259706, American Cancer Society (ACS) Research Scholar grant RSG-19-166-01-TBG, and American Lung Association (ALA) Lung Cancer Discovery Award 821321.

Availability of data and materials

TCGA lung adenocarcinoma, lung squamous cell carcinoma, and lung cancer data we analyzed were obtained from https://portal.gdc.cancer.gov/projects. All data generated or analyzed during this study are included in this article and its additional files. Extra data that support the findings of this study are available from the corresponding authors upon reasonable request.

Author contributions

Z.Q. and G.X. conceived and designed the study, led and contributed to all aspects of the analysis, and wrote the manuscript. S.D.S. contributed to the overall experimental design and data analysis and interpretation, provided advice and constructive feedback, and edited the manuscript. F.S. performed and analyzed all the experimental assays. H.Z. contributed to data analysis and edited the manuscript. P.Y. originally identified PDLIM2 LOH in human lung tumors by gene-specific PCR-based LOH analysis. Y.X. contributed to mouse clone maintenance and IHC staining.

Ethics approval and consent to participate

All animal studies were approved by the Institutional Animal Care and Use Committee of the University of Pittsburgh and carried out in accordance with NIH guidelines on animal care.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

[1]   Siegel RL, Miller KD, Fuchs HE, Jemal A (2022) Cancer statistics, 2022. *CA Cancer J Clin* **72**,
7-33.
[2]   Doroshow DB, Sanmamed MF, Hastings K, Politi K, Rimm DL, Chen L, Melero I, Schalper
KA, Herbst RS (2019) Immunotherapy in Non-Small Cell Lung Cancer: Facts and Hopes. *Clin*
Cancer Res **25**, 4592-4602.
[3]   Zou W, Wolchok JD, Chen L (2016) PD-L1 (B7-H1) and PD-1 pathway blockade for cancer
therapy: Mechanisms, response biomarkers, and combinations. *Sci Transl Med* **8**, 328rv324.
[4]   Zappasodi R, Merghoub T, Wolchok JD (2018) Emerging Concepts for Immune Checkpoint
Blockade-Based Combination Therapies. *Cancer Cell* **34**, 690.
[5]   Baxevanos P, Mountzios G (2018) Novel chemotherapy regimens for advanced lung cancer:
have we reached a plateau? *Ann Transl Med* **6**, 139.
[6]   Sun F, Li L, Yan P, Zhou J, Shapiro SD, Xiao G, Qu Z (2019) Causative role of PDLIM2
epigenetic repression in lung cancer and therapeutic resistance. *Nat Commun* **10**, 5324.
[7]   Garassino MC, Gadgeel S, Esteban E, Felip E, Speranza G, Domine M, Hochmair MJ, Powell S,
Cheng SY, Bischoff HG, Peled N, Reck M, Hui R, Garon EB, Boyer M, Wei Z, Burke T,
Pietanza MC, Rodriguez-Abreu D (2020) Patient-reported outcomes following pembrolizumab
or placebo plus pemetrexed and platinum in patients with previously untreated, metastatic, non-
squamous non-small-cell lung cancer (KEYNOTE-189): a multicentre, double-blind,
randomised, placebo-controlled, phase 3 trial. *Lancet Oncol* **21**, 387-397.
[8]   Leonetti A, Wever B, Mazzaschi G, Assaraf YG, Rolfo C, Quaini F, Tiseo M, Giovannetti E
(2019) Molecular basis and rationale for combining immune checkpoint inhibitors with
chemotherapy in non-small cell lung cancer. *Drug Resist Updat* **46**, 100644.
[9]   Sun F, Guo ZS, Gregory AD, Shapiro SD, Xiao G, Qu Z (2020) Dual but not single PD-1 or
TIM-3 blockade enhances oncolytic virotherapy in refractory lung cancer. *J Immunother Cancer*
8.
[10] Guo ZS, Qu Z (2021) PDLIM2: Signaling pathways and functions in cancer suppression and
host immunity. *Biochim Biophys Acta Rev Cancer* **1876**, 188630.
[11] Qu Z, Fu J, Yan P, Hu J, Cheng SY, Xiao G (2010) Epigenetic repression of PDZ-LIM domain-
containing protein 2: implications for the biology and treatment of breast cancer. *J Biol Chem*
285, 11786-11792.
[12] Qu Z, Yan P, Fu J, Jiang J, Grusby MJ, Smithgall TE, Xiao G (2010) DNA methylation-
dependent repression of PDZ-LIM domain-containing protein 2 in colon cancer and its role as a
potential therapeutic target. *Cancer Res* **70**, 1766-1772.
[13] Sun F, Xiao Y, Qu Z (2015) Oncovirus Kaposi sarcoma herpesvirus (KSHV) represses tumor
suppressor PDLIM2 to persistently activate nuclear factor kappaB (NF-kappaB) and STAT3
transcription factors for tumorigenesis and tumor maintenance. *J Biol Chem* **290**, 7362-7368.
[14] Yan P, Qu Z, Ishikawa C, Mori N, Xiao G (2009) Human T-cell leukemia virus type I-mediated
repression of PDZ-LIM domain-containing protein 2 involves DNA methylation but independent
of the viral oncoprotein tax. *Neoplasia* **11**, 1036-1041.
[15] Vanoirbeek E, Eelen G, Verlinden L, Carmeliet G, Mathieu C, Bouillon R, O'Connor R, Xiao G,
Verstuyf A (2014) PDLIM2 expression is driven by vitamin D and is involved in the pro-
adhesion, and anti-migration and -invasion activity of vitamin D. *Oncogene* **33**, 1904-1911.
[16] Yan P, Fu J, Qu Z, Li S, Tanaka T, Grusby MJ, Xiao G (2009) PDLIM2 suppresses human T-
cell leukemia virus type I Tax-mediated tumorigenesis by targeting Tax into the nuclear matrix
for proteasomal degradation. *Blood* **113**, 4370-4380.
[17] Fu J, Yan P, Li S, Qu Z, Xiao G (2010) Molecular determinants of PDLIM2 in suppressing
HTLV-I Tax-mediated tumorigenesis. *Oncogene* **29**, 6499-6507.

[18] Torrado M, Senatorov VV, Trivedi R, Fariss RN, Tomarev SI (2004) Pdlim2, a novel PDZ-LIM
domain protein, interacts with alpha-actinins and filamin A. *Invest Ophthalmol Vis Sci* **45**, 3955-
3963.
[19] Tanaka T, Soriano MA, Grusby MJ (2005) SLIM is a nuclear ubiquitin E3 ligase that negatively
regulates STAT signaling. *Immunity* **22**, 729-736.
[20] Loughran G, Healy NC, Kiely PA, Huigsloot M, Kedersha NL, O'Connor R (2005) Mystique is a
new insulin-like growth factor-I-regulated PDZ-LIM domain protein that promotes cell
attachment and migration and suppresses Anchorage-independent growth. *Mol Biol Cell* **16**,
1811-1822.
[21] Tanaka T, Grusby MJ, Kaisho T (2007) PDLIM2-mediated termination of transcription factor
NF-kappaB activation by intranuclear sequestration and degradation of the p65 subunit. *Nat*
Immunol **8**, 584-591.
[22] Qu Z, Fu J, Ma H, Zhou J, Jin M, Mapara MY, Grusby MJ, Xiao G (2012) PDLIM2 restricts Th1
and Th17 differentiation and prevents autoimmune disease. *Cell Biosci* **2**, 23.
[23] Zhou J, Qu Z, Yan S, Sun F, Whitsett JA, Shapiro SD, Xiao G (2015) Differential roles of
STAT3 in the initiation and growth of lung cancer. *Oncogene* **34**, 3804-3814.
[24] Qu Z, Sun F, Zhou J, Li L, Shapiro SD, Xiao G (2015) Interleukin-6 Prevents the Initiation but
Enhances the Progression of Lung Cancer. *Cancer Res* **75**, 3209-3215.
[25] Steinbrecher KA, Harmel-Laws E, Sitcheran R, Baldwin AS (2008) Loss of epithelial RelA
results in deregulated intestinal proliferative/apoptotic homeostasis and susceptibility to
inflammation. *J Immunol* **180**, 2588-2599.
[26] Yu H, Pardoll D, Jove R (2009) STATs in cancer inflammation and immunity: a leading role for
STAT3. *Nat Rev Cancer* **9**, 798-809.
[27] Xiao G, Fu J (2011) NF-kappaB and cancer: a paradigm of Yin-Yang. *Am J Cancer Res* **1**, 192-
221.
[28] Li L, Sun F, Han L, Liu X, Xiao Y, Gregory AD, Shapiro SD, Xiao G, Qu Z (2021) PDLIM2
repression by ROS in alveolar macrophages promotes lung tumorigenesis. *JCI Insight* **6**.
[29] Sun F, Li L, Xiao Y, Gregory AD, Shapiro SD, Xiao G, Qu Z (2021) Alveolar Macrophages
Inherently Express Programmed Death-1 Ligand 1 for Optimal Protective Immunity and
Tolerance. *J Immunol* **207**, 110-114.
[30] Wistuba, II, Behrens C, Virmani AK, Milchgrub S, Syed S, Lam S, Mackay B, Minna JD,
Gazdar AF (1999) Allelic losses at chromosome 8p21-23 are early and frequent events in the
pathogenesis of lung cancer. *Cancer Res* **59**, 1973-1979.
[31] Kang J (2015) Genomic alterations on 8p21-p23 are the most frequent genetic events in stage I
squamous cell carcinoma of the lung. *Exp Ther Med* **9**, 345-350.
[32] Macartney-Coxson DP, Hood KA, Shi HJ, Ward T, Wiles A, O'Connor R, Hall DA, Lea RA,
Royds JA, Stubbs RS, Rooker S (2008) Metastatic susceptibility locus, an 8p hot-spot for tumour
progression disrupted in colorectal liver metastases: 13 candidate genes examined at the DNA,
mRNA and protein level. *BMC Cancer* **8**, 187.
[33] Swalwell JI, Vocke CD, Yang Y, Walker JR, Grouse L, Myers SH, Gillespie JW, Bostwick DG,
Duray PH, Linehan WM, Emmert-Buck MR (2002) Determination of a minimal deletion interval
on chromosome band 8p21 in sporadic prostate cancer. *Genes Chromosomes Cancer* **33**, 201-
205.
[34] Wurster KD, Hummel F, Richter J, Giefing M, Hartmann S, Hansmann ML, Kreher S, Kochert
K, Krappmann D, Klapper W, Hummel M, Wenzel SS, Lenz G, Janz M, Dorken B, Siebert R,
Mathas S (2017) Inactivation of the putative ubiquitin-E3 ligase PDLIM2 in classical Hodgkin
and anaplastic large cell lymphoma. *Leukemia* **31**, 602-613.

[35] Matouk I, Raveh E, Ohana P, Lail RA, Gershtain E, Gilon M, De Groot N, Czerniak A, Hochberg A (2013) The increasing complexity of the oncofetal h19 gene locus: functional dissection and therapeutic intervention. *Int J Mol Sci* **14**, 4298-4316.

[36] Buscail L, Bournet B, Vernejoul F, Cambois G, Lulka H, Hanoun N, Dufresne M, Meulle A, Vignolle-Vidoni A, Ligat L, Saint-Laurent N, Pont F, Dejean S, Gayral M, Martins F, Torrisani J, Barbey O, Gross F, Guimbaud R, Otal P, Lopez F, Tiraby G, Cordelier P (2015) First-in-man phase 1 clinical trial of gene therapy for advanced pancreatic cancer: safety, biodistribution, and preliminary clinical findings. *Mol Ther* **23**, 779-789.

[37] Nyamay'Antu A, Dumont M, Kedinger V, Erbacher p (2019) Non-Viral Vector Mediated Gene Delivery: the Outsider to Watch Out For in Gene Therapy. *Cell & Gene Therapy Insights* **5**, 7.

[38] Bonnet ME, Erbacher P, Bolcato-Bellemin AL (2008) Systemic delivery of DNA or siRNA mediated by linear polyethylenimine (L-PEI) does not induce an inflammatory response. *Pharm Res* **25**, 2972-2982.

[39] Yang J, Hendricks W, Liu G, McCaffery JM, Kinzler KW, Huso DL, Vogelstein B, Zhou S (2013) A nanoparticle formulation that selectively transfects metastatic tumors in mice. *Proc Natl Acad Sci U S A* **110**, 14717-14722.

[40] Kellar A, Egan C, Morris D (2015) Preclinical Murine Models for Lung Cancer: Clinical Trial Applications. *Biomed Res Int* **2015**, 621324.

[41] Zhou J, Qu Z, Sun F, Han L, Li L, Yan S, Stabile LP, Chen LF, Siegfried JM, Xiao G (2017) Myeloid STAT3 Promotes Lung Tumorigenesis by Transforming Tumor Immunosurveillance into Tumor-Promoting Inflammation. *Cancer Immunol Res* **5**, 257-268.

[42] Li L, Han L, Sun F, Zhou J, Ohaegbulam KC, Tang X, Zang X, Steinbrecher KA, Qu Z, Xiao G (2018) NF-kappaB RelA renders tumor-associated macrophages resistant to and capable of directly suppressing CD8(+) T cells for tumor promotion. *Oncoimmunology* **7**, e1435250.

[43] Sun F, Qu Z, Xiao Y, Zhou J, Burns TF, Stabile LP, Siegfried JM, Xiao G (2016) NF-kappaB1 p105 suppresses lung tumorigenesis through the Tpl2 kinase but independently of its NF-kappaB function. *Oncogene* **35**, 2299-2310.

[44] Hecht SS (2002) Cigarette smoking and lung cancer: chemical mechanisms and approaches to prevention. *Lancet Oncol* **3**, 461-469.

[45] Commisso C, Davidson SM, Soydaner-Azeloglu RG, Parker SJ, Kamphorst JJ, Hackett S, Grabocka E, Nofal M, Drebin JA, Thompson CB, Rabinowitz JD, Metallo CM, Vander Heiden MG, Bar-Sagi D (2013) Macropinocytosis of protein is an amino acid supply route in Ras-transformed cells. *Nature* **497**, 633-637.

[46] Huang JL, Jiang G, Song QX, Gu X, Hu M, Wang XL, Song HH, Chen LP, Lin YY, Jiang D, Chen J, Feng JF, Qiu YM, Jiang JY, Jiang XG, Chen HZ, Gao XL (2017) Lipoprotein-biomimetic nanostructure enables efficient targeting delivery of siRNA to Ras-activated glioblastoma cells via macropinocytosis. *Nat Commun* **8**, 15144.

[47] Shi J, Kantoff PW, Wooster R, Farokhzad OC (2017) Cancer nanomedicine: progress, challenges and opportunities. *Nat Rev Cancer* **17**, 20-37.

Figure legends

Figure 1. PDLIM2 repression in human lung cancer involves both epigenetic alteration and genetic deletion, and *Pdlim2* genetic deletion in mice leads to development of spontaneous tumors, majorly lung tumors. (A) TCGA data showing *PDLIM2* repression in over 90% of lung tumors if using 50% of the expression level in normal lung tissues as the cut-off (NL, n = 110; Tumor, n = 1019). (B) TCGA data showing *PDLIM2* promoter hypermethylation and expression repression (dashed box) in over 70% of lung tumors when using 125% of the methylation level in normal lung tissues as the cut-off (n = 827). (C) TCGA data showing positive associations between *PDLIM2* expression and its gene copy numbers as well as *PDLIM2* genetic deletion and expression repression (dashed box) in about 58% of lung tumors using the copy number variation of -0.1 as the cut-off (n = 1010). (D) TCGA data showing simultaneous promoter hypermethylation and genomic deletion of *PDLIM2* (dashed box) in about 44% of lung tumors (n = 816). (E) Microsatellite-PCR showing *PDLIM2* loss in human lung tumors (n = 21). (F) qPCR showing *PDLIM2* loss in human lung cancer cell lines with known copy number of the *PDLIM2* gene (n = 25). (G) Kaplan-Meier tumor-free survival curve showing increased spontaneous tumors in *Pdlim2*$^{-/-}$ and *Pdlim2*$^{+/-}$ mice compared to WT mice. Gehan-Breslow-Wilcoxon test was performed. *p < 0.05. (H) Percentage of tumor types spontaneously developed in *Pdlim2*$^{-/-}$ and *Pdlim2*$^{+/-}$ mice showing a majority of lung tumors.

Figure 2. Systemic administration of PDLIM2 plasmid nanoparticles shows efficacy in mouse model of refractory lung cancer. (A) Schedule of lung cancer induction and treatment. (B) Urethane model showing efficacy of intravenous administration of PDLIM2-expression plasmid nanoparticles for refractory lung cancer (n ≥ 6). Nanoparticles with an empty vector plasmid (Vec) that was employed to express PDLIM2 were used as a control. (C) IHC staining showing decreased nuclear expression of STAT3 and RelA in lung tumors by PDLIM2 nanotherapy (n = 6). (D) IHC staining showing decreased Bcl-xL and increased apoptosis marker cleaved caspase -3 in lung tumors by PDLIM2 nanotherapy (n =

6). (E) IHC staining showing decreased Cyclin D1 and proliferation (BrdU incorporation) in lung

tumors by PDLIM2 nanotherapy (n = 6). Scale bar in (C-E), 20 μm. Student's *t* test was performed

(two tailed, unpaired) and data represent means ±□SEM in (B-E). **$p < 0.01$; ns, not statistically

significant.

Figure 3. PDLIM2 nanotherapy shows high tumor specificity and low toxicity. (A) IHC staining

showing high PDLIM2 re-expression in lung tumors after PDLIM2 nanotherapy. (B) PCR and IB

assays showing lung tumor-specific plasmids delivery and PDLIM2 expression by PDLIM2

nanotherapy (n = 3). (C) IHC staining showing comparable expression of PDLIM2 in the indicated

tissues of mice treated with PDLIM2 expression plasmid or empty vector plasmid nanoparticles. (D) No

significant changes in animal body weight by nanoPDLIM2 (n = 5). (E) H&E staining showing no

noticeable changes in major organs by nanoPDLIM2. Scale bar: (A and C) 20 μm; (E) 50 μm.

Figure 4. PDLIM2 nanotherapy renders lung cancers more vulnerable to chemotherapy. (A) Urethane

model showing synergy of PDLIM2 nanotherapy and chemotherapy in lung cancer treatment (n ≥ 5). (B)

IHC staining showing increased lung tumor cell apoptosis by PDLIM2 nanotherapy, chemotherapy, and

further increase by their combination (n = 6). (C) IHC staining showing RelA activation by

chemotherapy and blockage of chemo activation of RelA by PDLIM2 nanotherapy (n = 6). (D) IHC

staining showing strong MDR1 induction by chemotherapy and blockage of MDR1 induction by

PDLIM2 nanotherapy (n = 6). Scale bar in (B-D), 20 μm. Student's *t* test was performed (two tailed,

unpaired) and data represent means ± SEM. *$p < 0.05$; **$p < 0.01$; ns, not statistically significant.

Figure 5. PDLIM2 nanotherapy increases the efficacy of PD-1 blockade immunotherapy for refractory

lung cancer. (A) IHC staining showing increased TILs in lung tumors by PDLIM2 nanotherapy (n = 6).

(B) FACS showing increased MHC-I expression in lung tumor cells by PDLIM2 nanotherapy (n = 4).

(C) Urethane model showing PDLIM2 nanotherapy enhancing PD-1 immunotherapy efficacy in lung cancer treatment (n ≥ 5). (D) IHC staining showing increased TILs by PDLIM2 nanotherapy in the context of immunotherapy (n = 6). (E) FACS showing increased activation of CD8$^+$ T cells by PDLIM2 nanotherapy in the context of immunotherapy (n = 3). (F) IHC staining showing increased lung tumor cell apoptosis by PDLIM2 nanotherapy, immunotherapy, and further increase by their combination (n = 6). Scale bar in (A, D, F), 20 μm. Student's *t* test was performed (two tailed, unpaired) and data represent means ±□SEM. **$p < 0.01$; ns, not statistically significant.

Figure 6. Combination of PDLIM2 nanotherapy, chemotherapy and immunotherapy shows great efficacy in lung cancer treatment. (A) IHC staining showing PD-L1 induction by chemotherapy but not PDLIM2 nanotherapy. (B) Schedule of lung cancer induction and treatment. (C) Urethane model showing high resistance of lung tumors to the chemo and αPD-1 combination therapy in lung epithelial specific PDLIM2 deletion mice (ΔSPC) (n ≥ 4). (D) IHC staining showing no MHC-I induction by chemotherapy, PD-1 immunotherapy or their combination (n = 6). (E) Tumor examination showing complete remission of all lung tumors in 60% of mice by combination of the three therapies (n ≥ 5). (F) IHC staining showing increased TILs by PDLIM2 nanotherapy in mice treated with anti-PD-1 and chemotherapeutic drugs (n = 6). (G) FACS analysis showing increased lung CD8$^+$ T-cell activation by PDLIM2 nanotherapy in mice treated with anti-PD-1 and chemotherapeutic drugs (n = 5). (H) No significant effect of PDLIM2 nanotherapy on the body weight of mice treated with anti-PD-1 and chemotherapeutic drugs (n = 5). Scale bar in (A and F), 20 μm. Student's *t* test was performed (two tailed, unpaired) and data represent means ±□SEM in (**c-g**). *$p < 0.05$; **$p < 0.01$; ns, not statistically significant.

661   Figure 2
662

666   Figure 3

671   Figure 4

675   Figure 5

Supplementary Table S1.   Antibodies Used

Antibody	Company	Cat. No.	Dose	Usage
Anti-PDLIM2	Everest Biotech, Ramona, CA, USA	EB11878	1:400	IHC
Anti-Cleaved Caspase 3	Cell Signaling Technology, Danvers, MA, USA	9661	1:200	IHC
Anti-Bcl-xL	Cell Signaling Technology, Danvers, MA, USA	2764	1:2400	IHC
Anti-Cyclin D1	Santa Cruz Biotechnology, Dallas, TX, USA	sc-450	1:200	IHC
Anti-BrdU	Sigma-Aldrich, St. Louis, MO, USA	B2531	1:500	IHC
Anti-RelA	Cell Signaling Technology, Danvers, MA, USA	8242	1:500	IHC
Anti-STAT3	Cell Signaling Technology, Danvers, MA, USA	4904	1:1000	IHC
Anti-MDR1	Cell Signaling Technology, Danvers, MA, USA	13978	1:800	IHC
Anti-CD4	Abcam, Cambridge, MA, USA	ab183685	1:1000	IHC
Anti-CD8	Abcam, Cambridge, MA, USA	ab209775	1:1000	IHC
Anti-PD-L1	Cell Signaling Technology, Danvers, MA, USA	64988	1:200	IHC
Anti-MHC-I	Abcam, Cambridge, MA, USA	ab15681	1:200	IHC
Anti-mouse IgG Biotinylated	Vector Laboratories, Burlingame, CA, USA	BMK-2202	1:200	IHC
Goat anti-rat IgG Biotinylated	Vector Laboratories, Burlingame, CA, USA	BA9401	1:100	IHC
Rabbit anti-goat IgG Biotinylated	Santa Cruz Biotechnology, Dallas, TX, USA	sc-2774	1:200	IHC
Goat anti-rabbit IgG Biotinylated	Dako, Carpinteria, CA, USA	E0432	1:200	IHC
Anti-Hsp90	Santa Cruz Biotechnology, Dallas, TX, USA	sc-13119	1:1000	WB
Anti-c-Myc	Santa Cruz Biotechnology, Dallas, TX, USA	sc-40	1:1000	WB
Anti-CD16/CD32	eBioscience, San Diego, CA, USA	14-0161-85	1.0 µl per sample	FACS
Anti-CD45 FITC	Biolegend, San Diego, CA, USA	103107	0.5 µl per sample	FACS
Anti-EpCAM PE	eBioscience, San Diego, CA, USA	12-5791-82	0.625 µl per sample	FACS
Anti-MHC-I APC	eBioscience, San Diego, CA, USA	17-5998-82	1.25 µl per sample	FACS
Anti-CD3 PE	eBioscience, San Diego, CA, USA	12-0031-83	2.5 µl per sample	FACS
Anti-CD4 PE-Cy7	eBioscience, San Diego, CA, USA	25-0042-82	1.25 µl per sample	FACS
Anti-CD8a APC	eBioscience, San Diego, CA, USA	17-0081-83	0.625 µl per sample	FACS
Anti-IFNγ FITC	eBioscience, San Diego, CA, USA	11-7311-82	1 µl per sample	FACS
Anti-PD-1	BioXcell, West Lebanon, NH, USA	BE0273	200 µg/mouse/time	*in vivo* blockade

Supplementary Table S2. Primers Used

684
685

Gene	Species	Accession number	Forward (5' to 3')	Reverse (5' to 3')	Usage
D8S1786	human		CGAAAGATTGAGACCCCAT	GTTTCCACACCGAAGCC	Microsatellite PCR
D8S1752	human		TCCTGGATCAGGCAGAAA	TCAGAGTTGGGTGAGCGA	Microsatellite PCR
LINE-1	human	M80343.1	AAAGCCGCTCAACTACATGG	TGCTTTGAATGCGTCCCAGAG	RT-PCR
PDLIM2	human	NG_030435.1	CAAAGCCAAGGACGCTGACC	GCTCTGGCGGATCTTGCTCT	RT-PCR
pdlim2	mouse	Mutant primer	GCTACCCGTGATATTGCTGAAGAG		Genotyping PCR
		WT primer	GTCTTCTCTTCCTCAGGCATGGCG		
		Common primer	GCCAACATTGCCCCAGAAGACTCC		
Amp-R	Plasmid		AACTTTATCCGCCTCCAT	GAAGCCATACCAAACGAC	Plasmid detection

Supplementary Table S3. PDLIM2 RNA level and copy number in human lung cancer cell lines

	PDLIM2 gene	
	Relative RNA level	Copy number
NL-20	1.000	2.000
CALU-6	0.010	1.021
H727	0.020	1.590
H460	0.001	1.341
H2122	0.042	2.038
H23	0.128	0.376
H358	7.260	1.244
SKLU-1	0.390	1.685
SW1573	0.010	0.842
CALU-1	1.950	1.293
128-88T	0.049	1.767
H1299	0.003	0.243
273T	0.815	2.172
H827	1.017	1.237
H1650	2.168	0.679
H3255	5.118	0.587
343T	1.365	0.919
CALU-3	0.377	1.290
H1435	1.198	0.915
H1793	6.833	1.637
H596	0.195	0.782
H838	0.080	1.740
H1838	2.360	1.603
A549	0.010	5.265
H1975	1.086	2.507

691 Supplement Figure S1

692
693
694 Figure S1. Mice with *Pdlim2* deletion develop spontaneous tumors, majorly lung tumors. A, IHC
695 assays showing decreased and complete loss of PDLIM2 protein expression in the lungs of *Pdlim2*$^{+/-}$
696 (HT) and *Pdlim2*$^{-/-}$ (KO) mice, respectively. Scale bar, 20 µm. B, H&E staining of the lung tissues
697 showing spontaneous lung tumors in *Pdlim2*$^{+/-}$ and *Pdlim2*$^{-/-}$ mice. Scale bar, 20 µm.
698

699   Supplement Figure S2

700
701
702   Figure S2. PDLIM2 nanotherapy causes no obvious pathological changes in major organs. H&E staining
703   showing comparable patterns in lung, liver, kidney and spleen between Vec and PDLIM2 group in the
704   context of combinational chemotherapy and PD-1 blockade immunotherapy (n = 5). Scale bar: 50 µm.
705

706   Supplement Figure S3

707
708
Figure S3. Epigenetic drugs cause body weight loss in mice with lung cancer. Mice were i.p. injected
with urethane (1 g/kg) for 6 weeks to induce lung tumors, and then treated for 6 weeks with 5-aza-dC
and MS-275 (Epi, i.p., 1 mg/kg each, twice per week), carboplatin and paclitaxel (Chemo, i.p., 30 mg/kg
and 15 mg/kg, respectively, once per week), or their combinations. A, Body weight change between
prior to the first treatment and the euthanasia endpoint. B, Percentage of mice with peritoneal effusion
(n ≥ 4).
715

716   Supplement Figure S4

717
Figure S4. Epigenetic drugs show better efficacy than PDLIM2 nanotherapy in lung cancer treatment.
Mice were i.p. injected with urethane (1 g/kg) for 6 weeks to induce lung tumors, and then treated with
5-aza-dC and MS-275 (Epi, i.p., 1 mg/kg each, twice per week), carboplatin and paclitaxel (Chemo, i.p.,
30 mg/kg and 15 mg/kg, respectively, once per week), anti-PD-1 antibody (aPD-1, i.p., 200 µg/mouse,
three times per week), PDLIM2-expression plasmids containing nanoparticles (i.v., 25 µg
plasmid/mouse, once per week), or their combinations for 6 weeks before they were sacrificed for tumor
examination (n ≥ 4).

725
726