Gut microbiota and faecal cytokine profiles of rural Cambodian infants and

links to early life nutrition and diarrhoeal episodes.

Matthew J Dalby[1,2], Raymond Kiu[1,2], Iliana R Serghiou[1], Asuka Miyazaki[4], Holly Acford-

Palmer[1], Rathavy Tung[6], Shabhonam Caim[1], Sarah Phillips[1], Magdalena Kujawska[1, 2,3],

Mitsuaki Matsui[4], Azusa Iwamoto[5], Bunsreng Taking[7], Sharon E Cox[4,8,9], Lindsay J Hall[1,2,3,10].

[1]Food, Microbiome & Health, Quadram Institute Bioscience, Norwich Research Park, Norwich, NR4

7UQ, UK

[2]Institute of Microbiology and Infection, College of Medical and Dental Sciences, University of

Birmingham, Birmingham, B15 2TT, UK

[3]Intestinal Microbiome, School of Life Sciences, ZIEL – Institute for Food & Health, Technical

University of Munich, Freising, 80333, Germany

[4]School of Tropical Medicine & Global Health, Nagasaki University, Nagasaki, Japan.

[5]Bureau of International Health Cooperation, National Centre for Global Health and Medicine, Tokyo,

Japan

[6]National Maternal and Child Health Centre, Ministry of Health, Phnom Penh, Cambodia

[7]Kampong Cham Provincial Health Department, Ministry of Health, Kampong Cham, Cambodia

[8]London School of Hygiene and Tropical Medicine, Keppel Street, London, WC1E 7HT, UK

[9]Institute of Tropical Medicine, Nagasaki University, Nagasaki, Japan

[10]Norwich Medical School, University of East Anglia, Norwich Research Park, Norwich NR4 7TJ, UK

Abstract

The gut microbiota of infants in low-middle income countries like Cambodia remain underrepresented in microbiome research. This study aimed to explore the faecal gut microbiota composition and faecal cytokine profiles in a cohort of infants living in a rural province of Cambodia and explore the impact of sample storage conditions and infant environment on microbiota composition. Faecal samples collected at three time points from 32 infants (96 samples in total) after 7 months of age were analysed using 16S rRNA amplicon sequencing to determine the composition of the microbiota. Bacterial strains were isolated and subjected to whole genome sequencing and genomic analysis and concentrations of faecal cytokines were also measured. Initially, we compared the effects of two sample collection methods due to the challenges of faecal sample collection and storage in a rural location. Storage of faecal samples in a DNA preservation solution retained a greater abundance of *Bacteroides*. Analyses of microbiota composition of samples stored in DNA preservation solution indicated that *Bifidobacterium* was the most abundant genus with *Bifidobacterium longum* the most abundant species, particularly in breastfed infants. Most infants had detectable pathogenic taxa indicating frequent pathogen exposure, with *Shigella* and *Klebsiella* more abundant in infants with recent diarrheal illness. We did not detect antibiotic-associated perturbations in the gut microbiota, and no associations were found between the gut microbiota and infant growth. Genomic analysis of isolated strains indicated the presence of gene clusters encoding the ability to digest human milk oligosaccharides in *B. longum* and *Bifidobacterium breve* isolates. The presence of antibiotic-resistant genes was also identified in potentially pathogenic species, as well as in beneficial genera including *Bifidobacterium*. Faecal cytokine analysis showed higher concentrations of Interlukin-1alpha and vascular endothelial growth factor in breastfed infants, which may influence the infant gut mucosal immune system. This study provides insights into an underrepresented population of rural Cambodian infants, emphasising the impact of pathogen exposure and breastfeeding on gut microbiota composition and faecal immune profiles.

Introduction

Research on the gut microbiota has primarily focussed on populations in high income countries (e.g. Europe and North America) (Abdill et al., 2022), and the problems and limitations this bias is introducing to microbiome research is gaining increasing attention (Browne et al., 2024). Although more recent studies have increased representation of populations in low- and low-middle income countries (LMICs), infant studies lag behind those working on adult cohorts (Porras and Brito 2019). This is important for defining what is a 'healthy' human gut microbiota (Lackey et al., 2019) across individuals in LMICs (Porras and Brito, 2019), and because optimal infant development is particularly important for infants from LMICs due to increased risk of (mal)nutrition and infection. While a small proportion of microbiota samples studied globally so far have come from South Asia, including India and Bangladesh (Abdill et al., 2022), research into the gut microbiota of people living outside these countries remains very limited. Moreover, due to the logistics of sample collection and storage in rural areas, research in LMICs has often focused on individuals in urban areas, neglecting key infant populations. Cambodia in particular has a scarcity of microbiota studies and given that this country has such a high prevalence of infants classified as stunted/wasted, and has amongst the highest infant mortality rates in Southeast Asia, with infections accounting for many of these deaths (Cambodia Demographic and Health Survey, 2022), new studies targeting Cambodian infant populations are needed. Previous research focusing on the gut-associated microbiota of people in Cambodia has been limited with studies focusing on particular bacterial pathogens, such as antimicrobial resistance in *Enterobacterales* in faecal samples from babies and children (Auguet et al., 2021), the epidemiology of antibiotic resistance in *Escherichia coli* and *Klebsiella pneumoniae* in children (van Aartsen et al., 2019), and effects of iron supplementation in Cambodian women (Finlayson-Trick et al., 2023). However, analysis of the composition of the gut microbiota as a whole in infants from Cambodia has not previously been published.

While the nutritional status of infants is improving in Cambodia, children under the age of 5 in rural areas still face challenges, with 22% suffering linear growth faltering or 'stunting' (Cambodia Demographic and Health Survey, 2022). This is also true for infants (<1 year) in rural Cambodia who also suffer from similar levels of growth stunting at 19.2% (Miyazaki et al., 2021), with a high proportion also lacking access to appropriate hygiene facilities. This is coupled with a high proportion reportedly experiencing symptoms of diarrhoea, fever or cough and frequent use of unregulated antibiotic treatment to treat these childhood illnesses (Miyazaki et al., 2020). Malnutrition has previously been linked to differences in the gut microbiota of children from several different low income countries (Robertson, 2020; Surono et al., 2021), although research into these associations is lacking for children from Cambodia.

The establishment of the gut microbiota during infancy is important aspect of healthy infant

development with roles in immune, metabolic, endocrine, and other host developmental pathways

(Robertson et al., 2019). Vaginally-delivered, breast fed infants are characterised by their very high

abundance of the bacterial genus *Bifidobacterium*. This keystone microbiota member orchestrates

wider microbiome structuring and modulates host physiology through the breakdown of complex

dietary components such as human milk oligosaccharides (HMOs) in breast milk and plant-based

carbohydrates, leading to production of key metabolites that facilitate immune system development

and maturation and cognitive responses (Henrick et al., 2021). Previous research indicates that LMIC

infant microbiomes are characterised by *Bifidobacterium* species and strains, although these are

often different genetic variants when compared to those infants from high income country settings

(Lackey et al., 2019; Taft et al., 2022).

Infants born into resource-poor environments can have difficulties in establishing a healthy gut

microbial ecosystem (Lang & MAL-ED, 2018) due to multiple factors such as birth and breast feeding

practices, antibiotic exposure, maternal microbiota, diet, socio-economic status, host genetics,

agricultural dependence, urbanisation, access to clean water and hygiene practices (Porras and

Brito, 2019; Robertson et al., 2019). Infant microbiomes of those in LMICs are reportedly higher in

Segatella (previously *Prevotella*) species, associated with both positive and negative health

outcomes, and may correlate with higher plant based, fibre rich diets (Porras and Brito 2019). The

phylum Pseudomonadota (previously Proteobacteria) that includes a variety of pathogenic and

opportunistic pathogens, is highly prevalent in microbiomes of those in LMICs (Porras and Brito

2019). These factors may negatively affect the composition of the early infant gut microbiota

(Robertson et al., 2019), and may drive these observed microbiome differences across populations

(Porras and Brito 2019).

In this study we carried out an exploratory analysis of the gut microbiota in faecal samples collected

from 32 infants that are part of the Nutrition for Health of Aka-chan (baby in Japanese) and Mamas'

cohort (NHAM) based in Cambodia, which included background data on infant diet, health, and living

conditions. This provides an opportunity to explore the composition of the gut microbiota in rural

Cambodian infants that has not previously been investigated. The samples from this cohort are

unusual due to the rural location in which these infants live, with poor access to hygiene facilities,

high prevalence of pathogen exposure, and frequent antibiotic usage. We analysed the effects of

faecal sample storage on the composition of the microbiota of samples collected under suboptimal

conditions. Then described the infant faecal microbiota composition and explored its associations

with environmental factors such as infant diet and growth. Individual faecal bacteria cultured and

isolated and their genomes sequenced to identify the presence of antibiotic resistance and HMO

genes. Finally, the concentrations of faecal cytokines were measured to identify any associations with the gut microbiota. We aimed to explore the composition of gut microbiota of these infants and its associations with their environment.

Results

The longitudinal infant faecal samples analysed in this study were sourced from 32 infants representing a subset of infants that are part of the NHAM birth cohort based in the rural Kampong Cham province in southeast Cambodia, and approximately 60 kilometres from the provincial capital Kampong Cham (Miyazaki et al., 2020, 2021) (**Table 1**). Samples were collected from infants living in a riverside community stretching along approximately 15.5 km of riverbank of the Mekong River from Khbop Ta Nguon to Peam Koh Sna (Miyazaki et al., 2021). The infants were living in a rural area, with most families being small-scale farmers.

Variable	Infants
n	32
Age (days) (mean (SD))	214.9 (37.3)
sex = Female (%)	15 (46.9)
Breast feeding = No (%)	9 (28.1)
Complementary feeding = No (%)	17 (53.1)
Feeding bottle use = Yes (%)	21 (65.6)
Low birth weight (<2.5kg) (%)	
Normal	29 (90.6)
Unknown	2 (6.2)
Low	1 (3.1)
Growth = Stunted (%)	3 (9.4)
Length-for-age z-score (mean (SD))	-0.9 (0.8)
Underweight z-score (mean (SD))	-0.6 (0.8)
Weight-for-length (mean (SD))	0.0 (1.0)
Hand washing score (mean (SD))	5.2 (1.6)
Improved toilet facility = Yes (%)	22 (68.8)
Household toilet facility = Septic tank (%)	22 (68.8)
Number in household (mean (SD))	5.8 (2.2)
Antibiotic use (previous 7 days) = No (%)	28 (87.5)
Any recent illness (previous 7 days) = No (%)	11 (34.4)
Diarrheal illness (previous 7 days) = No (%)	26 (81.2)
Fever (previous 7 days) = No (%)	17 (53.1)
Cough (previous 7 days) = No (%)	20 (62.5)

Table 1: Infant baseline characteristics.

Sample storage conditions influence gut microbiota profiles

Sample collection and storage is challenging in areas without the facilities to immediately freeze
faecal samples after collection. To compare the preservation of DNA in samples under different
conditions, half of each faecal sample was immediately placed into a DNA preservation solution
(DNAShield), with the remaining half remaining untreated. The faecal samples were collected in a
rural location with approximately 2 days between the sample being produced and final laboratory
storage. The untreated half of each sample was transported in iceboxes and then initially frozen at -
20°C and then later stored at -80°C. The half of each sample stored in DNAShield were transported
at ambient temperature and then stored refrigerated at 4°C. Three, biweekly samples from 32
infants were collected (total of 96 samples), with half split between each storage condition. Both
frozen and DNAShield paired infant gut microbiota samples underwent 16S rRNA amplicon
sequencing to analyse their bacterial composition.

Out of the 96 paired samples, a total of 94 were successfully sequenced when stored in DNAShield,
while only 70 were successfully sequenced from Frozen samples (**Supplementary File 1**). There were
69 pairs of samples with successful 16S rRNA amplicon sequencing from both the Frozen and
DNAShield portion of the same faecal sample. The resulting bacterial genus composition was
compared between these 69 pairs of samples (**Figure 1**). While the total number of bacterial genera
detected was similar between frozen and DNAShield samples (**Figure 1A**), microbiota diversity
calculated using the Shannon index and Inverse Simpson index was higher in DNAShield samples
than paired frozen samples (**Figure 1B-C**). *Bacteroides*, *Escherichia*, *Parabacteroides, and*
Lachnoclostridium were the four bacterial genera with the largest median difference in relative
abundance between storage conditions that were higher in DNAShield samples compared to frozen
samples (**Figure 1D-G; Supplementary File 1**). *Bacteroides* showed the largest difference in favour of
DNAShield storage with a median relative abundance of 0.4% in frozen samples and 8.6% in
DNAShield samples (**Figure 1D; Supplementary File 1**). The relative abundance of *Bifidobacterium*,
Streptococcus, *Enterococcus,* and *Actinomyces* were the genera with the largest difference and
significantly higher abundance in favour of frozen samples (**Figure 1H-K; Supplementary File 1**).
Bifidobacterium comprised a median relative abundance of 63% in frozen samples and 46% in
DNAShield samples (**Figure 1H; Supplementary File 1**). Further analysis of bacterial microbiota
composition in this study was performed using samples stored in DNAShield due to the larger
number of samples with successful sequencing results and the higher abundance of individual
bacterial genera surviving in DNAShield stored samples.

These results indicate that sample storage conditions selectively can alter the abundance of different
genera in analysis of bacterial composition with significantly higher abundance of *Bacteroides* and
Parabacteroides remaining when samples were stored in DNA preservation solution. This shows that
DNA preservation solution can preserve genus abundance in final composition analysis even when
samples are initially transported at ambient temperatures and then stored refrigerated and not
frozen.

Figure 1. Differences in genus abundance between paired frozen samples and DNAShield stored samples. (A) The number of genera detected in each sample; (B) The Shannon diversity index of each sample; (C) The Inverse Simpson diversity index of each sample; Relative abundance between paired sampled after different storage conditions of (D) *Bacteroides*; (E) *Escherichia*; (F) *Parabacteroides* ; (G) *Lachnoclostridium;* (H) *Bifidobacterium*; (I) *Streptococcus*; (J) *Enterococcus*; (K) *Actinomyces*. Box plots show median and interquartile range with circles showing individual samples. Dotted lines link together paired Frozen and DNAShield samples from the same infant sample. Statistical significance between groups was tested using a Wilcoxon signed rank test with correction for multiple testing amongst bacteria genera using the false discovery rate method.

Bacterial genus composition of the infant gut microbiota is influenced by breastfeeding and

diarrheal illness

The gut microbiota of infants was dominated by *Bifidobacterium,* which comprised a median relative

abundance of 48.7% (**Figure 2A**). In addition, other genera including *Bacteroides, Blautia,*

Erysipelatoclostridium, Escherichia, Lactobacillus, Megamonas, Parabacteroides, and Streptococcus,

comprised the top ten most abundant genera, and formed the majority of the gut microbiota in

these infants (**Figure 2A**).

An exploratory analysis was carried out using metadata available for the infants from which samples

were collected. This included cross-sectional data collected at the start of the study before collection

of the faecal sample at time point 1 and additional information collected at each sample collection

visit (**Supplementary File 2**). The exploratory analyses used non-metric multidimensional scaling

(NMDS) and sample beta-diversity to cluster the samples by the similarity of their overall microbiota

composition at time point 1. Associations between the infant metadata with the beta-diversity of

the gut microbiota was analysed using Envfit and significant associations illustrated on the NMDS

plot (**Figure 2B**). Only breast-feeding status was significantly associated with microbiota beta-

diversity with clear separation in clusters between infants breast fed and those infants not breast

fed (**Figure 2B**).

To identify associations between the relative abundance of individual genera and infant metadata an

analysis using the MaAsLin2 package was carried out that included the samples from all three time

points, with the time point variable included as a random effect (**Figure 2C; Supplementary File 3**).

When background metadata variables were analysed individually as a fixed effect, both

breastfeeding status and recent diarrhoeal illness within the previous seven days showed significant

associations with genus abundance, which was maintained when both variables were included

together as a multivariable analysis (**Figure 2C**). Genera with differential abundance that was

significantly associated with breast-feeding and/or diarrheal illness are shown as a heat map with

positive and negative symbols indicating the direction of association (**Figure 2C**). The genera

positively associated with breast-feeding were *Bifidobacterium* and *Streptococcus* (**Figure 2C; Figure**

2D), whilst those with the strongest negative associations were *Klebsiella* and *Clostridium* (**Figure 2C**

; Figure 2E). Breast-feeding status remained unchanged for individual infants across the three time

points. At time point 1 the mean age of breast fed infants was 207 days and non-breast fed infants

214 days. Including age as a variable in the analysis did not alter the associations with breastfeeding.

Other measures of infant diet, the use of a feeding bottle or complementary feeding, were not

associated with genus abundance. MaAsLin2 analysis also showed that that recent diarrheal illness

within the previous seven days was associated with a lower relative abundance of *Bacteroides*

(**Figure 2F**) and higher relative abundance of *Clostridium, Klebsiella* and *Shigella* (**Figure 2G-H**).

Shigella was detected in 78% of samples, *Klebsiella* in 82% and *Clostridium* in 65% indicating that

these infants have a high prevalence of these potentially pathogenic genera in their gut microbiota

(**Supplementary File 4**). Breast feeding was the only variable associated with differences in microbial

diversity with the total number of bacterial genus (**Figure 3I**) and the Shannon diversity lower in

breast fed infants than non-breast fed infants (**Figure 3J**).

Only three infants included in the analysis were classified as having stunted growth with a height-

for-age z-score of less than 2 and this was not associated with microbiota composition. Tested as

continuous measures, length-for-age z-score , underweight z-score, and weight-for-length z-score

were also not associated with microbiota composition. However, this number of infants was too

small for robust testing preventing any strong conclusions being drawn from this result. Measures of

household hygiene facilities and total number of people in the household were not associated with

microbiota composition. Of particular note was the lack of an association between genus

composition and recent antibiotic use within the previous 7 days (**Supplementary File 3**).

These results indicate that breast feeding was the most influential factor shaping the gut microbiota

composition in favour of increased *Bifidobacterium* abundance. These results also show that recent

diarrheal illness was associated with higher relative abundance of the potentially pathogenic genera

Clostridium, Klebsiella and *Shigella*.

Figure 2. Genus composition and associations of genus composition with breast feeding status and

diarrheal illness. (A) The ten most abundant bacterial genera across all samples ordered on the x-

axis by *Bifidobacterium* abundance in each infant; (B) Non-metric multidimensional scaling (NMDS)

plot representing beta-diversity of infant genus microbiota at time point 1 clustered by similarity and

coloured by breastfeeding status with arrows showing top ten bacterial genus diving separation of

samples; (C) Heat map showing significant associations determined by MaAsLin2 analysis between

genus and breast feeding, diarrheal illness and time point; (D) The relative abundance of the genus *Bifidobacterium* by breastfeeding status; (E) The relative abundance of the genus *Klebsiella* by breastfeeding status; (F) The relative abundance of the genus *Bacteroides* by breastfeeding status; (G) The relative abundance of the genus *Shigella* by diarrheal illness within the previous 7 days; (H) The relative abundance of the genus *Klebsiella* by diarrheal illness within the previous 7 days; (I) The number of genera by breast feeding status; (J) The Shannon diversity index by breast feeding status; Points on plots show individual infant data and dotted lines on spaghetti plots link individual infants across time points with the bold line showing the group median.

Bacterial species composition of the infant gut microbiota is influenced by breastfeeding status

The same analysis carried out on genus level composition was carried out on the data at the species level with the limitation that species classifications using 16S rRNA amplicon data are less robust than genus (**Supplementary File 5**). The gut microbiota at the species level was dominated by a single *Bifidobacterium* species classified as *Bifidobacterium longum* (**Figure 3A**). Other abundant species detected were members of those genera identified in (**Figure 2A**) with two other species of *Bifidobacterium* identified as *B. breve* and *B. bifidum*, while *Escherichia coli*, *Klebsiella pneumoniae*, *Bacteroides fragilis* and an unclassified *Streptococcus* comprised the other abundant species present.

Repeating previous exploratory MaAsLin2 analysis of the species level 16S rRNA amplicon sequencing data identified the same association of breast-feeding status on microbiota composition with a stronger effect with species level data (**Figure 3C; Supplementary File 6**). Species with relative abundance that was statistically significantly associated with breast feeding and recent diarrheal illness are shown as a heat map with positive and negative symbols indicating the direction of association (**Figure 2C**). *Bifidobacterium longum* showed the strongest positive association with breast-feeding, with consistently higher median relative abundance in breastfed infants compared to those not breastfed (**Figure 3D**) while *Klebsiella pneumoniae* showed the strongest negative association with breast feeding (**Figure 3E**). The species *Shigella flexneri* (**Figure 3F**) and *Clostridium perfringens* (**Figure 3G**) were higher in samples taken after recent diarrheal illness. Breast feeding was the only variable associated with differences in microbial diversity at the species level of classification. The total number of bacterial species (**Figure 3H**) and the Shannon diversity was lower in breast fed infants than non-breast fed infants (**Figure 3I**). *Shigella flexneri* was detected in 49% of samples and *Clostridium perfringens* detected in 24% of samples indicating that these infants have a high prevalence of these potentially pathogenic genera in their gut microbiota (**Supplementary File 7**).

284 As described at the genus level, classification as stunted with a height-for-age z-score of less that 2

285 was not associated with gut microbiota species composition. Similarly, length-for-age z-score ,

286 underweight z-score, and weight-for-length z-score as continuous measures were also not

287 associated with species composition. Hygiene measures of household toilet facilities and total

288 number of people in the household were again not associated with species composition. Also noted

289 was the lack of an association between the gut microbiota composition and recent antibiotic use

290 (**Supplementary File 6**).

291 These species level results support the finding that breast feeding status was an major influence on

292 the gut microbiota composition and associated with the abundance of *Bifidobacterium longum*.

293 Recent diarrheal illness was associated with higher abundance bacterial species including *Shigella*

294 *flexneri* and *Clostridium perfringens* that are known to be potential causes of diarrheal illness.

Figure 3. Species composition and association of species composition with breast feeding and diarrheal illness. (A) The top ten most abundant bacterial species across all samples at time point 1 ordered by total *Bifidobacterium* species abundance; (B) Non-metric multidimensional scaling (NMDS) plot representing beta-diversity of infant species microbiota at time point 1 clustered by similarity and coloured by breastfeeding status with arrows showing top five bacterial genus diving

separation of samples; (C) Heat map showing significant associations determined by MaAsLin2

analysis between species and breast feeding, diarrheal illness and time point; (D) The relative

abundance of the *Bifidobacterium longum* by breastfeeding status; (E) The relative abundance of the

Klebsiella pneumoniae by breastfeeding status; (F) The relative abundance of the *Shigella flexneri* by

diarrheal illness within the previous 7 days; (G) The relative abundance of *Clostridium perfringens* by

diarrheal illness within the previous 7 days; (H) The number of species by breast feeding status; (I)

The Shannon diversity index by breast feeding status, Points on plots show individual infant data and

dotted lines on spaghetti plots link individual infants across time points with the bold line showing

the group median.

Infants harbour bacterial strains that encode diverse human milk oligosaccharide and antimicrobial resistance genes

The frozen faecal samples that were not stored in DNA preservation solution were used to culture

and isolate individual colonies for whole genome sequencing to gain further insight into the species

of bacteria in the gut microbiota of these infants. Genomic analysis was carried out on the

sequenced isolates to identify the species (and strains), and for the presence of virulence traits,

antimicrobial resistance genes and HMO gene clusters. From the bacterial isolates genome

sequenced 2 were identified as *Bacteroides*, 6 as *Bifidobacterium*, 1 as *Ruminococcus*, 3 as

Enterococcus, 21 as *Clostridium*, 3 as *Clostridioides*, and 2 as *Paraclostridium* (**Figure 4A**).

Bifidobacterium isolates comprised three species including *B. pseudocatenulatum*, *B. breve*, and *B.*

longum. Both *B. breve* and *B. longum* isolates contained complete HMO gene clusters, with none in

B. pseudocatenulatum (**Figure 4B**). This indicates the ability of the *B. breve* and *B. longum* present to

digest the oligosaccharides present in human breast milk, with only *B. longum* showing higher

relative abundance in breastfed infants compared to non-breast fed (**Figure 4C**). Comparing the four

genome-sequenced bacterial isolates identified as *Bifidobacterium longum* using Average Nucleotide

Identity with a 98% cut-off, three were identified as *B. longum* subsp. *longum* and one as *B. longum*

subsp. *suis* (**Supplementary File 8; Supplementary Figure 1**).

Genomic analysis showed the presence of antimicrobial resistance genes across all strains isolated

(**Figure 4D**). Of particular note was the presence of the antimicrobial resistance genes *erm*(X), *tet*(O),

tet(O/23/O), *tet*(Q), and *tet*(W) in the genomes of isolated *Bifidobacterium* species (**Figure 4B**),

which provide resistance to erythromycin (*erm*) and tetracycline (*tet*) respectively. The identification

of antimicrobial resistance genes in the *Bifidobacterium* strains present may link with the lack of any

association between gut microbiota composition and those infants receiving antibiotics, as shown

here with NMDS plots showing all samples combined (**Figure 4E**). The relative abundance of

Bifidobacterium was unchanged in samples from infants receiving antibiotics (**Figure 4F**). Amongst

this group of infants the most commonly received antibiotic was amoxycillin, with individual

instances of cefixime, and cefadroxile also received (**Supplementary File 2**).

Two *Clostridium perfringens* strains were isolated and genome-sequenced. Both contained five toxin

producing genes and six colonisation factor genes that relate to the ability to cause infection and

disease in the gut (**Figure 4G**). This confirms the presence of *C. perfringens* that was identified using

16S rRNA amplicon sequencing and the presence of toxin and colonisation factor genes supports a

potential role in its association with cases of infant diarrhoea.

Figure 4. Genomic analysis of isolates for species determination, human milk oligosaccharide (HMO) gene clusters, antimicrobial resistance genes, and virulence genes. (A) Tree showing species and relatedness of whole genome-sequenced isolates; (B) Presence of HMO gene clusters in *Bifidobacterium* isolates; (C) Antimicrobial resistance genes and virulence associated genes identified

in the isolate genomes; (D) NMDS plot showing all clustered faecal samples highlighted with recent

antibiotic use; (E) *Bifidobacterium* genus abundance by antibiotic use; (F) Presence of toxin genes

and colonisation factors in *Clostridium perfringens* isolates.

Faecal cytokines profiles are associated with breastfeeding status

The early life gut microbiota is known to play a key role in shaping gut and mucosal immune

responses. A multi-plex assay including a panel of thirty cytokines was used to determine

concentrations in the infant faecal samples (**Supplementary File 9**).

A linear mixed model incorporating infant metadata variables as a fixed effect and the infant as a

random effect was used as an exploratory analysis of associations between each cytokine and

metadata variables in all sample across the three time points (**Supplementary File 10**). The most

robust and consistent association was identified between two cytokines and breast feeding (**Figure**

5A-C). The cytokines Interleukin-1-alpha (IL-1α) (**Figure 5A**) and Vascular endothelial growth factor

(VEGF) (**Figure 5B**) were at significantly higher concentration in breast-fed infants. Both cytokines

showed substantial variation in concentration across time points within samples from the same

infant. IL-1α (**Figure 5A**) and VEGF (**Figure 5B**) were also the two cytokines with the highest

concentrations in the faecal samples across all infants.

To identify whether differences in major faecal cytokines by breastfeeding status were linked to

differences seen in the microbiota composition a correlation analysis was carried out between faecal

cytokine concentrations and the relative abundance of the 5 most abundant genus (**Figure 5C**) and

species (**Supplementary Figure 2**) in the microbiota. Weakly consistent positive associations were

seen between VEGF and *Lactobacillus* (**Figure 5D**) and *Streptococcus* (**Figure 5E**) across the three

time points, significant at two time points after correction for multiple testing. There were no

consistent correlations between the concentration of other cytokines and relative abundance of

individual bacterial genus or species, including *Bifidobacterium*.

Figure 5. Infant faecal cytokines concentration and associations with breast feeding status. (A) Individual infant IL-1α concentration by breastfeeding status across time points; (B) Individual infant VEGF concentration by breastfeeding status across time points; (C) Pearson correlations between faecal cytokine concentrations and five highest abundance genus with variables in both datasets individually normalised using the bestNormalize package; (D) Scatter plot of VEGF and *Lactobacillus* correlations labelled for each time point; (E) Scatter plot of VEGF vs *Streptococcus* correlations labelled for each time point.

Discussion

This study represents the first time that the infant gut microbiota composition of a cohort of rural Cambodian infants has been analysed in detail. We show that shortage of samples under rural conditions can radically reduce the abundance of certain genus within the samples. Infants had a gut microbiota dominated by the genus *Bifidobacterium,* shaped to a large degree by breastfeeding, which also impacts faecal cytokine profiles. Previous gut microbiota research in Cambodia has included investigation into antimicrobial resistance genes in Enterobacterales (Auguet et al., 2021) the epidemiology of antibiotic resistance in *Escherichia coli* and *Klebsiella pneumoniae* (van Aartsen et al., 2019), and effects of iron supplementation on individual genus abundance in women (Finlayson-Trick et al., 2023). This present study goes further and examines in-depth genus and selected species and strain composition in Cambodian infants for the first time.

Our study results represent an unusual example of different sample storage conditions as these samples were stored without immediate freezing. Overall the effect of storage conditions on the composition of the microbiota was small but a higher relative abundance of *Bacteroides* in particular was seen in samples stored in DNA preservation solution and a lower abundance of common species including *Bifidobacterium* due to the proportional nature of the microbiota data. *Bacteroides* represents an important genus in the infant gut microbiota, transferred from the mother during delivery and has been found to be higher in vaginally delivered infants (Matharu et al., 2022) with some *Bacteroides* species having the ability to consume components of breast milk (Kijner et al., 2022). The relatively high proportion of *Bacteroides* in these infants and their almost disappearance from samples without DNA preservation solution shows the importance of a storage method that preserves this significant component of the infant microbiota.

Previous research has investigated the effects of storage conditions on microbiota analysis and reported alterations in *Bacteroides* abundance, although this has focused on storage conditions while frozen or number of freeze thaw cycles (Pérez-Burillo et al., 2021; Poulsen et al., 2021). Room temperature storage of faecal samples for up to 72 hours has been reported to reduce genus including *Bacteroides* in adult faecal samples over time (Roesch et al., 2009). While defrosting frozen samples for three hours before analysis reduced *Bacteroides* abundance by half (Cardona et al., 2012). The duration of cool but unfrozen temperatures that samples in the present study were exposed to appear to have resulted in the particular low of *Bacteroides*. However, the unpreserved samples were also important for culturing and isolating live bacteria for genomic sequencing, highlighting the importance of having samples stored under both conditions.

Bifidobacterium was present at lower abundance in non-breast fed infants and not breast feeding
was associated with higher abundance of potentially pathogenic bacterial genus like *Klebsiella* and
Clostridium. Although the numbers of infants not breast fed was too low to draw strong conclusions,
previous research has indicated that increased *Bifidobacterium* in the preterm infant gut can help
reduce the proportions of potential pathogens (Alcon-Giner et al., 2020). This is linked to the ability
of *Bifidobacterium* to metabolise HMOs in breast milk and reduce gut pH to make it a more
inhospitable environment for overgrowth of opportunistic pathogens. *Bifidobacterium longum* was
the dominant species in this cohort of infants and *B. longum* was the only *Bifidobacterium* species
with higher abundance in breast fed infants. From the four *B. longum* isolates genome-sequenced,
all contained one or two HMO gene clusters indicating that they were potentially capable of
metabolising the sugars present in breast milk. Interestingly, three of the *B. longum* isolates were
confirmed to be *B. longum* subsp. *longum*, which do not typically dominate the gut microbiota of
infants in developed western countries nor other LMICs with high rates of breast feeding. It was
recently reported that *B. longum* subsp. *infantis* was the dominant *Bifidobacterium* in 1 to 2 month
old infants from the Gambia and Bangladesh with *B. longum* subsp. *longum* only forming a minor
proportion of the microbiota (Taft et al., 2022). The isolation of *B. longum* subsp. *longum* from
Cambodian samples may be due to sampling at an older age of around 7 months, with recent
research from Bangladesh indicating that a transition occurs from *B. longum* subsp. *infantis* to *B.*
longum subsp. *longum* with increasing infant age (Vatanen et al., 2022). One *B. longum* isolate was
identified as most closely matching *B. longum* subsp. *suis*, a subspecies originally isolated from pigs
(Matteuzzi et al., 1971) and found to be a dominant member of the piglet gut microbiota (Zani et al.,
1974). *Bifidobacterium* isolates corresponding to *B. longum* subsp. *suis* were recently identified in
Bangladeshi infants (Vatanen et al., 2022), which suggests that this may be an under-recognised
member of the infant gut microbiota. From only a single isolate we cannot determine the proportion
of *B. longum* subsp. *suis* within Cambodian samples, and this could represent the transient
colonisation of this bacterium from close proximity to pigs or that this is a resident of the gut
microbiota in this region. The presence of an HMO gene cluster capable of metabolising breast milk
sugars indicates that this *B. longum* subsp. *suis* isolate may be capable of metabolising breast milk
HMOs. The dominant abundance of the species *B. longum* in breastfed infants underline the central
effect of diet on the gut microbiota composition of these infants, overwhelming any other
environmental factor in this small cohort.

Previous research has identified associations between restricted infant growth and stunting due to
malnutrition and differences in the infant gut microbiota (Monira et al., 2011; Surono et al., 2021).
We did not find any associations between the gut microbiota and infants growth. This was

potentially due to the small cohort of infants involved with only three out of the thirty infants
included showing stunted growth and the relatively young age of the infants starting at an average
of 7 months of age. Previous research has looked at older children (Monira et al., 2011) or older
infants with a higher proportion of the infants suffering from stunting (Surono et al., 2021). This
indicates that in this small cohort of infants it is the presence or absence of breast milk having the
largest effect, potentially overwhelming the ability to detect other environmental influences.

In Cambodia, and many other LMICs, infants can be exposed to pathogens through provision of
contaminated drinking water in the first months of life (Poirot et al., 2018). Whilst we show that
there are high proportions of beneficial bacteria in most samples, pathogenic taxa including *Shigella*,
Escherichia, and *Clostridium* genus were also detected in most infants. Although *Shigella flexneri* and
Clostridium perfringens abundance in the gut was associated with recent diarrhoea these bacterial
species were also detectable in a large minority of infant samples indicating environmental exposure
to these organisms and potential asymptomatic carriage.

Antibiotics can significantly alter the profile of the early gut colonization (Alcon-Giner, et al., 2019)
resulting in reduced diversity (Gallego et al. 2019) and susceptibility to other infections (Porras and
Brita 2019). Notably, in Cambodia there is widespread, unregulated and inappropriate antibiotic use,
and these selection pressures may further facilitate the dissemination of AMR genes (van Aartsen et
al., 2019). An estimate of one-third of the Cambodian population is < 18 years old, and this group
may be a significant reservoir for the spread of antimicrobial resistant organisms (van Aartsen et al.,
2019). Indeed, we observed multiple antibiotic resistance genes in potentially pathogenic species,
including those associated with diarrhoeal episodes. We also observed presence of antimicrobial
resistance genes in *Bifidobacterium*. Although this may seem unexpected, as *Bifidobacterium* is very
rarely reported to be antibiotic resistant, a recent report from the neighbouring country of Vietnam
indicates that *Bifidobacterium* isolated from healthy infants harbour antibiotic resistance genes
(Chung The et al., 2021). We also found that Cambodian *Bifidobacterium* isolates contained
resistance genes to the antibiotics erythromycin and tetracycline. This may link to the fact that this
cohort of infants has previously been reported to be exposed to frequent and unregulated use of
antibiotics (Miyazaki et al., 2020), thus potentially driving AMR dynamics in beneficial members of
the infant gut microbiota. However, although amoxicillin was the most commonly used antibiotic in
this cohort, we did not observe any beta-lactam resistance genes in the isolates. Previous studies
have indicated that while it appears there is high and unregulated use of antibiotics within the
Cambodian population – they may be ineffective due to inappropriate storage thus reducing their
effectiveness. However, the widespread presence of resistance genes in other isolates may point
towards active selection through exposure to antibiotics. The apparent maintenance of

Bifidobacterium abundance despite antibiotic treatment may point to the presence of other novel
resistance genes in these *Bifidobacterium*, which if identified may have future potential for
combining effective probiotic therapy with antibiotic treatment.

One genus notable by its absence from this cohort of Cambodian infants is the genus *Segatella*.
Previously found to be a dominant member of the gut microbiota in rural West Africa while rare in
Italian children (De Filippo et al., 2010). More recently this high prevalence of *Segatella* in children in
low income countries has been confirmed in The Gambia, Kenya, Mali, and Bangladesh (Pop et al.,
2014). Whilst *Segatella* abundance was lower in children from Bangladesh it was still detectable in
children of a similar age to this cohort of Cambodian infants (Pop et al., 2014). In contrast
Cambodian infants contained *Bacteroides*, a genus previously associated with the gut microbiota of
European and US inhabitants (De Filippo et al., 2010). What dietary differences may account for this
difference cannot be determined in this study, however, it does show that further investigation is
needed into underrepresented populations. Whether the presence of *Bacteroides* and absence of
Segatella persists into adulthood remains unknown as composition of the adult gut microbiota in
this region remains still to be investigated.

Consistent differences in the faecal concentrations of the cytokines IL-1α and VEGF point to
potential effects of breastfeeding on the infant gut mucosal immune system. VEGF has been
reported to be secreted by the mammary glands, and breast milk contains high concentrations of
free VEGF that can bind to receptors on the surface of cells in the infant intestine (Kobata et al.,
2008; Siafakas et al., 1999; Vuorela et al., 2000). Reduced VEGF in preterm infants is suspected to be
involved in necrotising enterocolitis which is associated with overgrowth of opportunistic pathogens
including *C. perfringens* (Kiu et al., 2023; Sabnis et al., 2015). While the role of VEGF originating from
breast milk in term infants remains unexplored this growth factors could exert growth-promoting
and protective effects on the infant gut (Kobata et al., 2008). While the role of IL-1α in the infant gut
has not been explored, it has also been reported to be present in breast milk (Zanardo et al., 2005)
however, whether this has any effects the infant gut is currently unknown. Although we are unable
to demonstrate a breast milk origin from these results, the presence of higher concentrations of
these three faecal cytokines in breast fed infants does suggest that they may be derived from breast
milk, with a potential to influence infant development that warrants further investigation.

While there was little consistent association across time points between the gut microbiota at the
genus level and faecal cytokines, a weak positive correlation was noted between *Lactobacillus* and
Streptococcus and the faecal cytokine VEGF. As *Streptococcus* abundance was higher in breastfed
infants this correlation is potentially linked to breastfeeding. As both microbiota composition and

faecal cytokines is variable with time it may not be possible to identify links between factors in a
limited sample size of infants. Interestingly, despite being the genus most differentiated by breast
feeding status there no consistent correlations between *Bifidobacterium* IL-1α or VEGF. Indeed,
many species and strains from this genus are known to positively influence the gut and systemic
immune system – including *B. longum,* which is proposed to be in part mediated by metabolites
produced as a result of HMO degradation e.g. acetate (Henrick et al., 2021; Laursen et al., 2021).
However, further studies including larger number of infants would need to be performed to explore
microbial-immune interactions in more detail. In addition, the use of the frozen samples for faecal
cytokine analysis may be limited as it is unknown how such storage conditions may impact the
degradation of the cytokines tested.

We show that this cohort of infants from rural Cambodia have a *Bifidobacterium* dominated gut
microbiota (including atypical species) that is primarily shaped by breastfeeding. This highlights the
influence of breast feeding in supporting a beneficial infant microbiota in rural conditions. As
Bacteroides represents a common and important genus in the human gut the effects of sample
storage are an important consideration when immediate freezing of samples is not possible. The lack
of impact of antibiotic use may indicates either widespread antibiotic resistance or the use of
ineffective antibiotics. The finding that faecal cytokines are shaped by breast feeding and could
potentially originate from breast milk is a finding with unknown but potentially interesting
implications. This study adds significantly to the available data on the composition of infant gut
microbiota communities in a previously neglected and difficult to sample geographical area of the
world. Moreover, given Cambodian profiles are somewhat different from other infant cohorts,
including LMICs, studies such as these shows the importance of looking at infants from
underrepresented areas for defining what can constitute a normal infant microbiota.

Materials & Methods

Study design & ethics

Samples were collected from a sub-group of infants enrolled in NHAM cohort based in villages along the Mekong river in rural Kampong Cham province in Cambodia. This registered all infants born between April 2016 and March 2019 with the objective of investigating causes of stunting. Full ethical approvals including National Ethical Committee for Health Research in the Ministry of Health, Cambodia (016) and the Ethical Committee of the National Centre for Global Health and Medicine in Japan (NCGM-G-001870-01), and methods for the collection of detailed meta-data for infants included in this sub-study are as previously described (Miyazaki et al., 2021, 2020). Weight and length were measured to the nearest 0.1kg and 1 mm by trained field workers at each home visit using a weighing scale (Seca 877, Seca, Hamburg, Germany) and a length board (Seca 417, Seca).

Sample collection

Infant faecal samples were analysed from a subset of 32 infants out of 47 enrolled in a detailed 3-month longitudinal follow-up study (Miyazaki et al., 2021, 2020) collected between March and July in 2017. Mothers/primary care-givers and their infants were visited by study field workers every 14 days. Three samples were analysed from each infant for a total of 96 individual samples. The trained field worker gave instructions on how to collect and equipped the caregivers with the material for the child stool collection prior to routine home visits. Once stool sample collection by mothers or caregivers was completed, the sample was kept in relatively cooler place in the house or in cooler bags with ice until the field worker visited to collect them. The majority of the samples were collected by the field worker within 10 hours of the time of child defecation reported by mothers (maximum: 47 hours and minimum: 15 min among all sample). Each sample collected was divided in two and half stored in DNAShield and stored unaltered at ambient temperatures until delivered to laboratory. The other half of the sample was stored in iceboxes with refrigerant and transported to a -20° freezer on the same day and once delivered was stored at -80°. The first sample was collected after an initial visit where cross-sectional metadata was collected. Subsequent samples were collected at approximately two week intervals along with further individual visit metadata. Underweight z-scores (weight-for-age), weight-for-length z-scores, length-for-age z-score were calculated previously (Miyazaki et al., 2021). A hand washing score was generated as a composite score combining the frequency and consistency of handwashing with soap. Improved toilet facility was defined as having access to a toilet flushing to a septic tank or pit latrine compared to unimproved having only access to a pit latrine or no toilet facility. Household toilet facility was defined as having a toilet within the home flushing to a septic tank.

DNA extraction of preterm stool samples

FastDNA Spin Kit for Soil (MP) was used to extract DNA from preterm faeces following manufacturer

instructions, with extended 3 min bead-beating. DNA concentration and quality were quantified

using a Qubit® 2.0 fluorometer (Invitrogen).

16S rRNA amplicon sequencing of faecal samples

16S rRNA region (V1-V2) primers were used for library construction as described previously (Alcon-

Giner et al., 2020). This set of primers allowed the amplification of one 16S rRNA gene sequencing

library containing 96 different samples. PCR conditions used were: cycle of 94°C 3 min and 25 cycles

of 94°C for 45 s, 55°C for 15 s and 72°C for 30 s. Sequencing of the 16S rRNA gene libraries was

performed using Illumina MiSeq platform with 300 bp paired end reads.

16S rRNA amplicon data analysis

The raw sequence reads obtained were subjected to quality filtering using trim_galore v0.4.3. The

resulting filtered sequence reads were then mapped against the 16S rRNA SILVA database

(SILVA_132_SSURef_tax_silva) using BLASTn v2.2.25+ with a maximum e-value threshold of 10e-3.

This mapping process was performed separately for both pairs of sequences (Quast et al., 2013).

After completing the BLASTn alignments, the output files were annotated using the paired-end

protocol of MEGAN6, employing the default Lowest Common Ancestor (LCA) parameters (Huson et

al., 2016; Huson and Mitra, 2012).

Statistical analysis was conducted using R version 4.2.2 (2022-10-31 ucrt) and RStudio

(2023.06.0+421). To determine the number of reads required to adequately represent the

microbiota in each sample, rarefaction curves were generated using the vegan package in R. For the

analysis at the genus level, the sample with the smallest number of reads contained 37,306 classified

reads, which was deemed an acceptable minimum. For the species level analysis, the sample with

the second lowest number of reads contained 9,317 reads, which was considered an acceptable

minimum. The 16S rRNA gene sequence data were then normalized by subsampling using the rarefy

function in the phyloseq package, resulting in an even depth of 37,306 reads for the genus data and

9,317 reads for the species data. One sample was excluded from the species data analysis as it only

contained 3,965 reads.

To examine differences in the relative abundance of genera among the 69 infant fecal samples,

which had 16S rRNA sequence data obtained from both samples stored in DNAShield solution and

Frozen, a Wilcoxon rank sum test was performed in R. This test was chosen due to the non-

parametric nature of the majority of the genus abundance data. To account for multiple testing, a

Benjamini & Hochberg false discovery rate (FDR) method was applied, and a p-value of less than 0.05
was considered statistically significant for adjusted p-values.

The overall composition of the 16S rRNA gene sequence data was analyzed using NMDS (Non-metric
multidimensional scaling) plots. These plots were generated in R Studio using the vegan package
version 2.6-4, employing a Bray-Curtis dissimilarity calculation. To determine the statistical
significance of differences in microbial community structure based on infant metadata variables,
permutational MANOVA was conducted using the Adonis function from the vegan R package version
2.6-4.

Microbial diversity at both the genus and species taxa levels was calculated using the vegan package
version 2.6-4. The number of genera or species was determined for each sample, and the Shannon
diversity index was calculated accordingly. Linear mixed-effect models, utilizing the "lmer" function
in R, were employed to test the associations between the relative abundance of genera or species
and metadata variables. The number of genera or species, as well as the Shannon diversity index,
were assessed against each individual metadata variable, with Time Point included as a fixed effect
and Patient ID included as a random effect in the model.

To test for differential abundance of genera and species in relation to available background
metadata variables for the infant samples, the Maaslin2 package (Microbiome Multivariable
Associations with Linear Models) (Mallick et al., 2021), was utilized. The Maaslin2 analysis was
performed with default parameters, including Benjamini-Hochberg FDR multiple testing correction
and a significance threshold of 0.25. No minimum prevalence or abundance cut-offs were applied,
and the data input was previously normalized. Each metadata variable was individually tested,
incorporating Time Point and Patient ID as random effect grouping variables in a mixed-effect
model. Positive associations were identified between Breast Feeding and Diarrheal Illness. These
two variables were subsequently tested together as fixed effects, with Time Point and Patient ID
included as random effects in the final analysis.

Anaerobic bacterial culturing and isolation from infant faecal samples

Anaerobic bacterial culturing and isolation was performed on seventeen Cambodian infant faecal
samples (C151, C152, C153, C154, C172, C176, C185, C186, C203, C224, C230, C232, C243, C257,
C267, C281, and C287) following (Browne, 2018; Browne et al., 2016; Forster et al., 2019). Sample
preparation was performed under anaerobic conditions. YCFA media and PBS was pre-reduced in an
anaerobic cabinet at 37 °C for a minimum of 12hs before culturing. Two samples, weighing 100mg
each, were prepared from each faecal sample. The first sample was homogenised in reduced PBS,

serially diluted, and plated directly onto the YCFA agar. The second sample was ethanol treated

using 70% ethanol under ambient aerobic conditions for 1h to kill vegetative cells; the sample was

then washed three times with reduced PBS to remove the ethanol, homogenised in the reduced PBS,

serially diluted and plated onto YCFA + Sodium taurocholate agar to stimulate spore germination. All

agar plates were placed in the anaerobic cabinet at 37°C for 48hs. Single colonies were picked and

streaked onto pre-reduced YCFA media to isolate and purify the colonies; YCFA media was pre-

reduced in the anaerobic cabinet at 37 °C for a minimum of 12hs before isolating colonies. Colonies

from the non-ethanol treated samples were streaked onto the reduced YCFA agar plates, and

colonies from the ethanol treated samples were streaked onto the reduced YCFA + Sodium

taurocholate agar plates. Plates were then left to incubate in the anaerobic cabinet at 37°C for 48-

72hs. This was carried out a second time to make sure isolates were pure. Falcon tubes containing

reduced YCFA broth was inoculated with a single colony from the YCFA and YCFA + Sodium

taurocholate agar plates, and left to incubate in the anaerobic cabinet at 37°C for 48hs, with

bacterial stocks prepared in 1.5ml cryovials with 30% glycerol YCFA broth, and stored at -80°C.

Additionally, two infant faecal samples C203 and C232 were processed for targeted *Bifidobacterium*

isolation. Briefly, serially diluted aliquots of faecal homogenates ($10^{-1} - 10^{-6}$ in PBS) were plated onto

MRS agar supplemented with 50mg/l mupirocin and 0.5g/l L-cysteine HCL and incubated in an

anaerobic cabinet at 37 °C for 48h. Colonies with desired morphology were randomly selected and

re-streaked on MRS agar to purity. Pure cultures were prepared in 1.5 ml cryovials with 30% glycerol

Reinforced Clostridial Medium (RCM) and stored at −80 °C.

Whole Genome Sequencing (WGS)

DNA extractions were performed using the MPbio Fast DNA Spin Kit for soil and bead beating (but

with an extended 3 min bead beating time). DNA quantification was carried out using the Qubit 2.0

fluorometer. DNA from pure bacterial cultures was sequenced at Wellcome Trust Sanger Institute

using 96-plex Illumina HiSeq 2500 platform to generate 125 bp paired end reads as described

previously (Harris et al., 2010).

Genomic analysis

Paired-end raw sequence reads (fastq) were quality-filtered (-q 20) via fastp v0.20.0 (Chen et al.,

2018) prior to de novo genome assembly using SPAdes v3.14.1 (Bankevich et al., 2012) at default

parameters. Next, contigs smaller than 500bp were discarded in each genome assembly. All genome

assemblies were subjected to contamination check via CheckM v1.1.3 (Parks et al., 2015),

contaminated (contamination >5%) and/or incomplete (completeness <90%) genome assemblies

were excluded from further analyses (n=5). All genomes were identified taxonomically using gtdb-tk

v1.5.1 (Chaumeil et al., 2020), using ANI cut-off 95% against type strains of each species.

These high-quality genome assemblies (n=71) were then dereplicated at ANI 99.9% (cut-off for

strain level) using dRep v3.2.2 (Olm et al., 2017) prior to generation of a distance tree using

Mashtree v1.2.0 (Katz et al., 2019) on default parameters. Tree was annotated in iTOL v6.5.8

(Letunic and Bork, 2019).

These genomes were also subjected to sequence search analysis via ABRicate v1.0.1 (Seemann,

2023) based on ResFinder v4.0 database (Bortolaia et al., 2020) (--minid=95 and --mincov=90) to

determine antimicrobial resistance (AMR) profiles, and VFDB database (Liu et al., 2019) (--minid=90

and –mincov=90) to determine virulence profiles. Furthermore, HMO clusters were detected using

an in-house HMO sequence database via ARBricate at –minid=70 (Lawson et al., 2020). In addition,

C. perfringens virulence profiles were predicted via database TOXIper v1.1 using ABRicate at –

minid=90 (Kiu, 2023).

For *Bifidobacterium longum* subspecies analysis, *B. longum* strains were compared with four

subspecies type strains namely subspecies *infantis*, subspecies *longum*, subspecies *suillum* and

subspecies *suis*, to generate a core gene alignment via Panaroo v1.2.8 (Tonkin-Hill et al., 2020),

resulting in 1,212 core genes, which core-gene alignment constructed by MAFFT (Katoh and

Standley, 2013) were SNP-extracted (via SNP-sites v2.3.3)(Page et al., 2016) and subsequently used

to generate a phylogenetic tree via IQ-TREE v2.0.5 (Minh et al., 2020, p. 2) with ultra-fast bootstrap

replicates -B 1000 and automatic evolutionary model selection option -m TEST (best fit model was

determined to be GTR+F+ASC+G4). Subspecies identification was performed using fastANI v1.3 (Jain

et al., 2018) at cut-off 98%.

Faecal cytokine measurement

Faecal samples were homogenized with PBS using a FastPrep® Bead Beater (4.0m/s, 3min),

centrifuged (14,000rpm, 15min) and 25µl of supernatant was used for the assay. Samples were

analysed using MULTI-SPOT™ plates, MESO Quickplex SQ120 and discovery workbench software

according to the manufacturer's protocol. Pre-coated immunoassays V-PLEX Proinflammatory Panel

1 (human) and V-PLEX Cytokine Panel 1 (human) were used to detect a set of 20 different cytokines:

IFNγ, IL-1β, IL-2, IL-4, IL-6, IL-8, IL-10, IL-12p70, IL-13, TNFα, GMCSF, IL-1α, IL-5, 273 IL-7, IL-12p40, IL-

15, IL-16, IL-17A, TNFβ, and VEGF-A.

Data and Code Availability

Bacterial pure genome assemblies are now deposited at NCBI GenBank under project accession
PRJNA958097 while 16S rRNA amplicon sequencing raw reads are deposited in NCBI SRA under
project PRJNA958098.

Statistical analysis

Associations between faecal cytokine concentrations and metadata variables were tested using the
MaAsLin2 package (Mallick et al., 2021), a differences in concentration against background metadata
variables available for the infant samples. For faecal cytokine analysis MaAsLin2 was run with default
parameters including multiple testing correction using Benjamini-Hochberg FDR with a significance
threshold of 0.25, except: no minimum prevalence or abundance cut-offs were applied, normalised
methods used wans "CLR", and the transformation "LOG". Each metadata variable was tested
individually with Time Point and with Patient ID included as a Random Effect grouping variable to
run a mixed effect model. Positive associations were identified with Breast Feeding and Diarrheal
Illness and these two variables where tested together as Fixed Effects with Time Point and Patient ID
included as a Random Effect in the final analysis.

Correlations between faecal cytokine concentration and the relative abundance of genus and species
was tested using a Pearson correlation with correction for multiple testing using Benjamini &
Hochberg FDR correcting between both cytokines and time point. Both cytokine concentration and
genus/species abundance were first normalised using the bestNormalize package in R.

Author contributions

Conceptualisation, L.J.H and S.E.C.; Methodology, M.J.D, R.K., I.R.S, A.M., HA-P., R.T., S.C., S.P., M.K.,
M.M., A.I., and B.T; Software, M.J.D., R.K., S.C., and M.K.; Validation, M.J.D, K.S., S.E.C., and L.J.H.;
Formal analysis, M.J.D, R.K., A.M, S.E.C., and L.J.H.; Investigation, M.J.D, R.K., I.R.S, A.M., HA-P., R.T.,
S.C., S.P., M.K., M.M., A.I., B.T., S.E.C., and L.J.H.; Resources, M.J.D., A.M., R.T., M.M., A.I., B.T., S.E.C.,
and L.J.H; Data Curation, M.J.D., R.K., A.M., S.E.C.; Writing - Original Draft, M.J.D. and L.J.H.; Writing -
Review and Editing, M.J.D, R.K., I.R.S, A.M., HA-P., R.T., S.C., S.P., M.K., M.M., A.I., B.T., S.E.C, and
L.J.H; Visualisation, M.J.D and R.K.; Supervision, S.E.C,. and L.J.H.; Project Administration, M.J.D.,
S.E.C, and L.J.H.; Funding Acquisition, S.E.C, and L.J.H.

Competing interest statement

735 The authors declare that they have no competing interests.

Acknowledgments

738   We thank the families for participating in the study.

Funding information

741   L.J.H. is supported by Wellcome Trust Investigator Awards 100974/C/13/Z and 220876/Z/20/Z; the
742   Biotechnology and Biological Sciences Research Council (BBSRC), Institute Strategic Programme Gut
743   Microbes and Health BB/R012490/1, and its constituent projects BBS/E/F/000PR10353 and
744   BBS/E/F/000PR10356.

References

Abdill, R.J., Adamowicz, E.M., Blekhman, R., 2022. Public human microbiome data are dominated by
highly developed countries. PLoS Biol 20, e3001536.
https://doi.org/10.1371/journal.pbio.3001536

Alcon-Giner, C., Dalby, M.J., Caim, S., Ketskemety, J., Shaw, A., Sim, K., Lawson, M.A.E., Kiu, R.,
Leclaire, C., Chalklen, L., Kujawska, M., Mitra, S., Fardus-Reid, F., Belteki, G., McColl, K.,
Swann, J.R., Kroll, J.S., Clarke, P., Hall, L.J., 2020. Microbiota Supplementation with
Bifidobacterium and Lactobacillus Modifies the Preterm Infant Gut Microbiota and
Metabolome: An Observational Study. Cell Rep Med 1, 100077.
https://doi.org/10.1016/j.xcrm.2020.100077

Auguet, O.T., Niehus, R., Gweon, H.S., Berkley, J.A., Waichungo, J., Njim, T., Edgeworth, J.D., Batra,
R., Chau, K., Swann, J., Walker, S.A., Peto, T.E.A., Crook, D.W., Lamble, S., Turner, P., Cooper,
B.S., Stoesser, N., 2021. Population-level faecal metagenomic profiling as a tool to predict
antimicrobial resistance in Enterobacterales isolates causing invasive infections: An
exploratory study across Cambodia, Kenya, and the UK. EClinicalMedicine 36, 100910.
https://doi.org/10.1016/j.eclinm.2021.100910

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M., Nikolenko,
S.I., Pham, S., Prjibelski, A.D., Pyshkin, A.V., Sirotkin, A.V., Vyahhi, N., Tesler, G., Alekseyev,
M.A., Pevzner, P.A., 2012. SPAdes: a new genome assembly algorithm and its applications to
single-cell sequencing. J Comput Biol 19, 455–477. https://doi.org/10.1089/cmb.2012.0021

Bortolaia, V., Kaas, R.S., Ruppe, E., Roberts, M.C., Schwarz, S., Cattoir, V., Philippon, A., Allesoe, R.L.,
Rebelo, A.R., Florensa, A.F., Fagelhauer, L., Chakraborty, T., Neumann, B., Werner, G.,
Bender, J.K., Stingl, K., Nguyen, M., Coppens, J., Xavier, B.B., Malhotra-Kumar, S., Westh, H.,
Pinholt, M., Anjum, M.F., Duggett, N.A., Kempf, I., Nykäsenoja, S., Olkkola, S., Wieczorek, K.,
Amaro, A., Clemente, L., Mossong, J., Losch, S., Ragimbeau, C., Lund, O., Aarestrup, F.M.,
2020. ResFinder 4.0 for predictions of phenotypes from genotypes. Journal of Antimicrobial
Chemotherapy 75, 3491–3500. https://doi.org/10.1093/jac/dkaa345

Browne, H.P., 2018. Development of anaerobic culturing methods to study the evolution and
transmission of the human intestinal microbiota (doctoral). London School of Hygiene &
Tropical Medicine. https://doi.org/10.17037/PUBS.04647551

Browne, H.P., Forster, S.C., Anonye, B.O., Kumar, N., Neville, B.A., Stares, M.D., Goulding, D., Lawley,
T.D., 2016. Culturing of "unculturable" human microbiota reveals novel taxa and extensive
sporulation. Nature 533, 543–546. https://doi.org/10.1038/nature17645

Browne, H.P., Iqbal, N.T., Osman, M., Tigoi, C., Lawley, T.D., Gordon, J.I., Ahmed, T., Kariuki, S., 2024.
Boosting microbiome science worldwide could save millions of children's lives. Nature 625,
237–240. https://doi.org/10.1038/d41586-024-00017-8

Cambodia Demographic and Health Survey, 2022. Cambodia Demographic and Health Survey [WWW
Document]. URL https://dhsprogram.com/publications/publication-PR136-Preliminary-
Reports-Key-Indicators-Reports.cfm (accessed 1.19.23).

Cardona, S., Eck, A., Cassellas, M., Gallart, M., Alastrue, C., Dore, J., Azpiroz, F., Roca, J., Guarner, F.,
Manichanh, C., 2012. Storage conditions of intestinal microbiota matter in metagenomic
analysis. BMC Microbiology 12, 158. https://doi.org/10.1186/1471-2180-12-158

Chaumeil, P.-A., Mussig, A.J., Hugenholtz, P., Parks, D.H., 2020. GTDB-Tk: a toolkit to classify
genomes with the Genome Taxonomy Database. Bioinformatics 36, 1925–1927.
https://doi.org/10.1093/bioinformatics/btz848

Chen, S., Zhou, Y., Chen, Y., Gu, J., 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics 34, i884–i890. https://doi.org/10.1093/bioinformatics/bty560

Chung The, H., Nguyen Ngoc Minh, C., Tran Thi Hong, C., Nguyen Thi Nguyen, T., Pike, L.J., Zellmer,
C., Pham Duc, T., Tran, T.-A., Ha Thanh, T., Van, M.P., Thwaites, G.E., Rabaa, M.A., Hall, L.J.,
Baker, S., 2021. Exploring the Genomic Diversity and Antimicrobial Susceptibility of
Bifidobacterium pseudocatenulatum in a Vietnamese Population. Microbiology Spectrum 9,
e00526-21. https://doi.org/10.1128/Spectrum.00526-21

De Filippo, C., Cavalieri, D., Di Paola, M., Ramazzotti, M., Poullet, J.B., Massart, S., Collini, S.,
Pieraccini, G., Lionetti, P., 2010. Impact of diet in shaping gut microbiota revealed by a

comparative study in children from Europe and rural Africa. Proc. Natl. Acad. Sci. U.S.A. 107,

14691–14696. https://doi.org/10.1073/pnas.1005963107

Finlayson-Trick, E., Nearing, J., Fischer, J.A., Ma, Y., Wang, S., Krouen, H., Goldfarb, D.M., Karakochuk,

C.D., 2023. The Effect of Oral Iron Supplementation on Gut Microbial Composition: a

Secondary Analysis of a Double-Blind, Randomized Controlled Trial among Cambodian

Women of Reproductive Age. Microbiol Spectr 11, e0527322.

https://doi.org/10.1128/spectrum.05273-22

Forster, S.C., Kumar, N., Anonye, B.O., Almeida, A., Viciani, E., Stares, M.D., Dunn, M., Mkandawire,

T.T., Zhu, A., Shao, Y., Pike, L.J., Louie, T., Browne, H.P., Mitchell, A.L., Neville, B.A., Finn, R.D.,

Lawley, T.D., 2019. A human gut bacterial genome and culture collection for improved

metagenomic analyses. Nat Biotechnol 37, 186–192. https://doi.org/10.1038/s41587-018-

0009-7

Harris, S.R., Feil, E.J., Holden, M.T.G., Quail, M.A., Nickerson, E.K., Chantratita, N., Gardete, S.,

Tavares, A., Day, N., Lindsay, J.A., Edgeworth, J.D., de Lencastre, H., Parkhill, J., Peacock, S.J.,

Bentley, S.D., 2010. Evolution of MRSA during hospital transmission and intercontinental

spread. Science 327, 469–474. https://doi.org/10.1126/science.1182395

Henrick, B.M., Rodriguez, L., Lakshmikanth, T., Pou, C., Henckel, E., Arzoomand, A., Olin, A., Wang, J.,

Mikes, J., Tan, Z., Chen, Y., Ehrlich, A.M., Bernhardsson, A.K., Mugabo, C.H., Ambrosiani, Y.,

Gustafsson, A., Chew, S., Brown, H.K., Prambs, J., Bohlin, K., Mitchell, R.D., Underwood,

M.A., Smilowitz, J.T., German, J.B., Frese, S.A., Brodin, P., 2021. Bifidobacteria-mediated

immune system imprinting early in life. Cell 184, 3884-3898.e11.

https://doi.org/10.1016/j.cell.2021.05.030

Huson, D.H., Beier, S., Flade, I., Górska, A., El-Hadidi, M., Mitra, S., Ruscheweyh, H.-J., Tappu, R.,

2016. MEGAN Community Edition - Interactive Exploration and Analysis of Large-Scale

Microbiome Sequencing Data. PLOS Computational Biology 12, e1004957.

https://doi.org/10.1371/journal.pcbi.1004957

Huson, D.H., Mitra, S., 2012. Introduction to the Analysis of Environmental Sequences:

Metagenomics with MEGAN, in: Anisimova, M. (Ed.), Evolutionary Genomics: Statistical and

Computational Methods, Volume 2, Methods in Molecular Biology. Humana Press, Totowa,

NJ, pp. 415–429. https://doi.org/10.1007/978-1-61779-585-5_17

Jain, C., Rodriguez-R, L.M., Phillippy, A.M., Konstantinidis, K.T., Aluru, S., 2018. High throughput ANI

analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat Commun 9, 5114.

https://doi.org/10.1038/s41467-018-07641-9

Katoh, K., Standley, D.M., 2013. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. Molecular Biology and Evolution 30, 772–780.

https://doi.org/10.1093/molbev/mst010

Katz, L.S., Griswold, T., Morrison, S.S., Caravas, J.A., Zhang, S., den Bakker, H.C., Deng, X., Carleton,

H.A., 2019. Mashtree: a rapid comparison of whole genome sequence files. J Open Source

Softw 4. https://doi.org/10.21105/joss.01762

Kijner, S., Cher, A., Yassour, M., 2022. The Infant Gut Commensal Bacteroides dorei Presents a

Generalized Transcriptional Response to Various Human Milk Oligosaccharides. Frontiers in

Cellular and Infection Microbiology 12.

Kiu, R., 2023. TOXIper: Toxinotype assignment of Clostridium perfringens via ABRicate.

Kiu, R., Shaw, A.G., Sim, K., Acuna-Gonzalez, A., Price, C.A., Bedwell, H., Dreger, S.A., Fowler, W.J.,

Cornwell, E., Pickard, D., Belteki, G., Malsom, J., Phillips, S., Young, G.R., Schofield, Z., Alcon-

Giner, C., Berrington, J.E., Stewart, C.J., Dougan, G., Clarke, P., Douce, G., Robinson, S.D.,

Kroll, J.S., Hall, L.J., 2023. Particular genomic and virulence traits associated with preterm

infant-derived toxigenic Clostridium perfringens strains. Nat Microbiol 8, 1160–1175.

https://doi.org/10.1038/s41564-023-01385-z

Kobata, R., Tsukahara, H., Ohshima, Y., Ohta, N., Tokuriki, S., Tamura, S., Mayumi, M., 2008. High

levels of growth factors in human breast milk. Early Human Development 84, 67–69.

https://doi.org/10.1016/j.earlhumdev.2007.07.005

Lackey, K.A., Williams, J.E., Meehan, C.L., Zachek, J.A., Benda, E.D., Price, W.J., Foster, J.A., Sellen,

D.W., Kamau-Mbuthia, E.W., Kamundia, E.W., Mbugua, S., Moore, S.E., Prentice, A.M., K.,

D.G., Kvist, L.J., Otoo, G.E., García-Carral, C., Jiménez, E., Ruiz, L., Rodríguez, J.M., Pareja,

R.G., Bode, L., McGuire, M.A., McGuire, M.K., 2019. What's Normal? Microbiomes in Human

Milk and Infant Feces Are Related to Each Other but Vary Geographically: The INSPIRE Study.

Front. Nutr. 6, 45. https://doi.org/10.3389/fnut.2019.00045

Laursen, M.F., Sakanaka, M., von Burg, N., Mörbe, U., Andersen, D., Moll, J.M., Pekmez, C.T.,

Rivollier, A., Michaelsen, K.F., Mølgaard, C., Lind, M.V., Dragsted, L.O., Katayama, T.,

Frandsen, H.L., Vinggaard, A.M., Bahl, M.I., Brix, S., Agace, W., Licht, T.R., Roager, H.M.,

2021. Bifidobacterium species associated with breastfeeding produce aromatic lactic acids in

the infant gut. Nat Microbiol 6, 1367–1382. https://doi.org/10.1038/s41564-021-00970-4

Lawson, M.A.E., O'Neill, I.J., Kujawska, M., Gowrinadh Javvadi, S., Wijeyesekera, A., Flegg, Z.,

Chalklen, L., Hall, L.J., 2020. Breast milk-derived human milk oligosaccharides promote

Bifidobacterium interactions within a single ecosystem. ISME J 14, 635–648.

https://doi.org/10.1038/s41396-019-0553-2

Letunic, I., Bork, P., 2019. Interactive Tree Of Life (iTOL) v4: recent updates and new developments.
Nucleic Acids Research 47, W256–W259. https://doi.org/10.1093/nar/gkz239

Liu, B., Zheng, D., Jin, Q., Chen, L., Yang, J., 2019. VFDB 2019: a comparative pathogenomic platform
with an interactive web interface. Nucleic Acids Research 47, D687–D692.
https://doi.org/10.1093/nar/gky1080

Mallick, H., Rahnavard, A., McIver, L.J., Ma, S., Zhang, Y., Nguyen, L.H., Tickle, T.L., Weingart, G., Ren,
B., Schwager, E.H., Chatterjee, S., Thompson, K.N., Wilkinson, J.E., Subramanian, A., Lu, Y.,
Waldron, L., Paulson, J.N., Franzosa, E.A., Bravo, H.C., Huttenhower, C., 2021. Multivariable
association discovery in population-scale meta-omics studies. PLOS Computational Biology
17, e1009442. https://doi.org/10.1371/journal.pcbi.1009442

Matharu, D., Ponsero, A.J., Dikareva, E., Korpela, K., Kolho, K.-L., de Vos, W.M., Salonen, A., 2022.
Bacteroides abundance drives birth mode dependent infant gut microbiota developmental
trajectories. Frontiers in Microbiology 13.

Matteuzzi, D., Crociani, F., Zani, O., Trovatelli, L.D., 1971. Bifidobacterium suis n. sp.: A new species
of the genus Bifidobacterium isolated from pig faces. Zeitschrift für allgemeine Mikrobiologie
11, 387–395. https://doi.org/10.1002/jobm.19710110504

Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A., Lanfear,
R., 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the
Genomic Era. Molecular Biology and Evolution 37, 1530–1534.
https://doi.org/10.1093/molbev/msaa015

Miyazaki, A., Matsui, M., Tung, R., Taing, B., White, L.V., Iwamoto, A., Cox, S.E., 2021. Determinants
of growth measurements in rural Cambodian infants: a cross-sectional study. Int Health 13,
49–56. https://doi.org/10.1093/inthealth/ihaa018

Miyazaki, A., Tung, R., Taing, B., Matsui, M., Iwamoto, A., Cox, S.E., 2020. Frequent unregulated use
of antibiotics in rural Cambodian infants. Transactions of The Royal Society of Tropical
Medicine and Hygiene 114, 401–407. https://doi.org/10.1093/trstmh/traa020

Monira, S., Nakamura, S., Gotoh, K., Izutsu, K., Watanabe, H., Alam, N., Endtz, H., Cravioto, A., Ali,
Sk., Nakaya, T., Horii, T., Iida, T., Alam, M., 2011. Gut Microbiota of Healthy and
Malnourished Children in Bangladesh. Frontiers in Microbiology 2.

Olm, M.R., Brown, C.T., Brooks, B., Banfield, J.F., 2017. dRep: a tool for fast and accurate genomic
comparisons that enables improved genome recovery from metagenomes through de-
replication. ISME J 11, 2864–2868. https://doi.org/10.1038/ismej.2017.126

Page, A.J., Taylor, B., Delaney, A.J., Soares, J., Seemann, T., Keane, J.A., Harris, S.R., 2016. SNP-sites:
rapid efficient extraction of SNPs from multi-FASTA alignments. Microb Genom 2, e000056.
https://doi.org/10.1099/mgen.0.000056

Parks, D.H., Imelfort, M., Skennerton, C.T., Hugenholtz, P., Tyson, G.W., 2015. CheckM: assessing the
quality of microbial genomes recovered from isolates, single cells, and metagenomes.
Genome Res. 25, 1043–1055. https://doi.org/10.1101/gr.186072.114

Pérez-Burillo, S., Hinojosa-Nogueira, D., Navajas-Porras, B., Blasco, T., Balzerani, F., Lerma-Aguilera,
A., León, D., Pastoriza, S., Apaolaza, I., Planes, F.J., Francino, M.P., Rufián-Henares, J.Á., 2021.
Effect of Freezing on Gut Microbiota Composition and Functionality for In Vitro
Fermentation Experiments. Nutrients 13, 2207. https://doi.org/10.3390/nu13072207

Pop, M., Walker, A.W., Paulson, J., Lindsay, B., Antonio, M., Hossain, M.A., Oundo, J., Tamboura, B.,
Mai, V., Astrovskaya, I., Bravo, H.C., Rance, R., Stares, M., Levine, M.M., Panchalingam, S.,
Kotloff, K., Ikumapayi, U.N., Ebruke, C., Adeyemi, M., Ahmed, D., Ahmed, F., Alam, M.T.,
Amin, R., Siddiqui, S., Ochieng, J.B., Ouma, E., Juma, J., Mailu, E., Omore, R., Morris, J.G.,
Breiman, R.F., Saha, D., Parkhill, J., Nataro, J.P., Stine, O.C., 2014. Diarrhea in young children
from low-income countries leads to large-scale alterations in intestinal microbiota
composition. Genome Biology 15, R76. https://doi.org/10.1186/gb-2014-15-6-r76

Porras, A.M., Brito, I.L., 2019. The internationalization of human microbiome research. Current
Opinion in Microbiology 50, 50–55. https://doi.org/10.1016/j.mib.2019.09.012

Poulsen, C.S., Kaas, R.S., Aarestrup, F.M., Pamp, S.J., 2021. Standard Sample Storage Conditions Have
an Impact on Inferred Microbiome Composition and Antimicrobial Resistance Patterns.
Microbiology Spectrum 9, e01387-21. https://doi.org/10.1128/Spectrum.01387-21

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner, F.O., 2013.
The SILVA ribosomal RNA gene database project: improved data processing and web-based
tools. Nucleic Acids Research 41, D590–D596. https://doi.org/10.1093/nar/gks1219

Robertson, R.C., 2020. The Gut Microbiome in Child Malnutrition, in: Michaelsen, K.F., Neufeld, L.M.,
Prentice, A.M. (Eds.), Nestlé Nutrition Institute Workshop Series. S. Karger AG, pp. 133–144.
https://doi.org/10.1159/000503352

Robertson, R.C., Manges, A.R., Finlay, B.B., Prendergast, A.J., 2019. The Human Microbiome and
Child Growth – First 1000 Days and Beyond. Trends in Microbiology 27, 131–147.
https://doi.org/10.1016/j.tim.2018.09.008

Roesch, L.F.W., Casella, G., Simell, O., Krischer, J., Wasserfall, C.H., Schatz, D., Atkinson, M.A., Neu, J.,
Triplett, E.W., 2009. Influence of Fecal Sample Storage on Bacterial Community Diversity.
Open Microbiol J 3, 40–46. https://doi.org/10.2174/1874285800903010040

Sabnis, A., Carrasco, R., Liu, S.X.L., Yan, X., Managlia, E., Chou, P.M., Tan, X.-D., De Plaen, I.G., 2015.
Intestinal Vascular Endothelial Growth Factor Is Decreased in Necrotizing Enterocolitis.
Neonatology 107, 191–198. https://doi.org/10.1159/000368879
Seemann, T., 2023. ABRicate: Mass screening of contigs for antimicrobial and virulence genes.
Siafakas, C.G., Anatolitou, F., Fusunyan, R.D., Walker, W.A., Sanderson, I.R., 1999. Vascular
endothelial growth factor (VEGF) is present in human breast milk and its receptor is present
on intestinal epithelial cells. Pediatr Res 45, 652–657. https://doi.org/10.1203/00006450-
199905010-00007
Surono, I.S., Widiyanti, D., Kusumo, P.D., Venema, K., 2021. Gut microbiota profile of Indonesian
stunted children and children with normal nutritional status. PLOS ONE 16, e0245399.
https://doi.org/10.1371/journal.pone.0245399
Taft, D.H., Lewis, Z.T., Nguyen, N., Ho, S., Masarweh, C., Dunne-Castagna, V., Tancredi, D.J., Huda,
M.N., Stephensen, C.B., Hinde, K., von Mutius, E., Kirjavainen, P.V., Dalphin, J.-C., Lauener,
R., Riedler, J., Smilowitz, J.T., German, J.B., Morrow, A.L., Mills, D.A., 2022. Bifidobacterium
Species Colonization in Infancy: A Global Cross-Sectional Comparison by Population History
of Breastfeeding. Nutrients 14, 1423. https://doi.org/10.3390/nu14071423
Tonkin-Hill, G., MacAlasdair, N., Ruis, C., Weimann, A., Horesh, G., Lees, J.A., Gladstone, R.A., Lo, S.,
Beaudoin, C., Floto, R.A., Frost, S.D.W., Corander, J., Bentley, S.D., Parkhill, J., 2020.
Producing polished prokaryotic pangenomes with the Panaroo pipeline. Genome Biology 21,
180. https://doi.org/10.1186/s13059-020-02090-4
van Aartsen, J.J., Moore, C.E., Parry, C.M., Turner, P., Phot, N., Mao, S., Suy, K., Davies, T., Giess, A.,
Sheppard, A.E., Peto, T.E.A., Day, N.P.J., Crook, D.W., Walker, A.S., Stoesser, N., 2019.
Epidemiology of paediatric gastrointestinal colonisation by extended spectrum
cephalosporin-resistant Escherichia coli and Klebsiella pneumoniae isolates in north-west
Cambodia. BMC Microbiol 19, 59. https://doi.org/10.1186/s12866-019-1431-9
Vatanen, T., Ang, Q.Y., Siegwald, L., Sarker, S.A., Le Roy, C.I., Duboux, S., Delannoy-Bruno, O., Ngom-
Bru, C., Boulangé, C.L., Stražar, M., Avila-Pacheco, J., Deik, A., Pierce, K., Bullock, K., Dennis,
C., Sultana, S., Sayed, S., Rahman, M., Ahmed, T., Modesto, M., Mattarelli, P., Clish, C.B.,
Vlamakis, H., Plichta, D.R., Sakwinska, O., Xavier, R.J., 2022. A distinct clade of
Bifidobacterium longum in the gut of Bangladeshi children thrives during weaning. Cell 185,
4280-4297.e12. https://doi.org/10.1016/j.cell.2022.10.011
Vuorela, P., Andersson, S., Carpén, O., Ylikorkala, O., Halmesmäki, E., 2000. Unbound vascular
endothelial growth factor and its receptors in breast, human milk, and newborn intestine.

The American Journal of Clinical Nutrition 72, 1196–1201.

https://doi.org/10.1093/ajcn/72.5.1196

Zanardo, V., Nicolussi, S., Cavallin, S., Trevisanuto, D., Barbato, A., Faggian, D., Favaro, F., Plebani, M.,

2005. Effect of Maternal Smoking on Breast Milk Interleukin-1α, β-Endorphin, and Leptin

Concentrations. Environ Health Perspect 113, 1410–1413. https://doi.org/10.1289/ehp.7702

Zani, G., Biavati, B., Crociani, F., Matteuzzi, D., 1974. Bifidobacteria from the Faeces of Piglets.

Journal of Applied Bacteriology 37, 537–547. https://doi.org/10.1111/j.1365-

2672.1974.tb00479.x

Supplementary Files

Supplementary file 1. Genus differences between DNAShield and Frozen storage conditions.

Supplementary File 2. Metadata and Genus Sequence Data

Supplementary file 3. Genus data Maaslin2 analysis results.

Supplementary file 4. Genus prevalence in samples.

Supplementary file 5. Combined sample metadata and species sequencing data for all samples.

Supplementary file 6. Species data Maaslin2 analysis results.

Supplementary file 7. Species prevalence in samples.

Supplementary File 8. Pairwise Average Nucleotide Identity (ANI)

Supplementary file 9. Metadata and Faecal Cytokine Concentrations.

Supplementary file 10. MaAsLin2 Cytokine Results

Supplementary Figures:

Supplementary Figure 1. The subspecies of *Bifidobacterium longum* isolates determined using

average nucleotide identity with a 98% cut-off.

Supplementary Figure 2. Correlations between cytokines and Genus or Species relative abundance.

1000

1001