Coding dynamics of the striatal networks during learning

Maxime Villet[1†]**, Patricia Reynaud-Bouret**[2†]**, Julien Poitreau**[3]**, Jacopo Baldi**[1,2]**, Sophie Jaffard**[2]**, Ashwin James**[4]**, Alexandre Muzy**[4]**, Francesca Sargolini**[3‡]**, Ingrid Bethus**[1‡]

[1]Université Côte d'Azur, CNRS, IPMC and NeuroMod; [2]Université Côte d'Azur, CNRS, LJAD and NeuroMod; [3]Laboratoire de Neurosciences Cognitives, UMR 7291, CNRS, Aix-Marseille Univ.; [4]Université Côte d'Azur, CNRS, I3S and NeuroMod

*For correspondence:
bethus@ipmc.cnrs.fr (IB); Patricia.
Reynaud-Bouret@univ-cotedazur.fr
(PRB)

[†]These authors contributed
equally to this work.
[‡]These authors also contributed
equally to this work.

Present address: [1]Université Côte
d'Azur, CNRS, IPMC, France;
[2]Université Côte d'Azur, CNRS,
LJAD, France; [3]Université Côte
d'Azur, CNRS, I3S, France;
[4]Laboratoire de Neurosciences
Cognitives, UMR 7291, CNRS,
Aix-Marseille Univ.

Abstract   The rat dorsomedial (DMS) and dorsolateral striatum (DMS), equivalent to caudate nucleus and putamen in primates, are generally required for goal-directed and habit behaviour, respectively. This functional dichotomy is well established in instrumental conditioning, but it is less clear whether it also exists in non-instrumental learning. In this study we investigated this issue by recording DMS and DLS single neuron activity in rats performing a continuous alternation task in which no cue instruction is provided. We first applied a classical analytical approach to identify task-related activity based on the modifications of single neuron firing rate in relation to specific task events or maze trajectories. We then used an innovative approach based on Hawkes process to reconstruct a directed connectivity graph of neurons simultaneously recorded in each learning session, that was used to decode animal behavior. This approach enabled us to better unravel the role of DMS and DLS neural networks across learning stages, from the acquisition to the optimization of the behavioral strategy. We showed that DMS and DLS display different task-related activity throughout learning stages, and the proportion of coding neurons over time decreases in the DMS and increases in the DLS. Despite theses major differences, the decoding power of both networks increases during learning. These results suggest that both DMS and DLS are engaged during all learning stages, but the two neural networks gradually reorganize in different ways, in contrast with the common assumption of a gradual shift from DMS to DLS activity across learning stages.

Introduction

During learning process, it is commonly believed that the goal directed behaviour, controlled by flexible action-outcome contingencies, is sustained by the activity of the dorsomedial portion of the striatum - DMS, whereas the activation of the dorsolateral striatum - DLS - sustains inflexible motor routines, stimulus-response learning and more generally habit behaviour (*Balleine and O'Doherty, 2010*; *Balleine and Dickinson, 1998*; *Dickinson and Balleine, 1994*; *Yin et al., 2005*; *Dickinson et al., 1995*; *Graybiel, 1998*; *Packard and Knowlton, 2002*; *Yin et al., 2004*; *Hilario et al., 2012*). Such functional dichotomy is also supported by the anatomical organization of the striatal cortical projections. Cortical inputs from the sensorimotor and premotor cortices project somatotopically to the DLS, whereas associative and prefrontal cortices project to the DMS (*Hunnicutt et al., 2016*; *Hintiryan et al., 2016*; *Hooks et al., 2018*) with fewer inputs from somatosensory cortical areas (*Reig and Silberberg, 2014*).

However DMS-DLS functional dissociation is mainly observed in instrumental conditioning (*Yin et al., 2004*, *2005*; *Balleine et al., 2009*), and it is less straightforward in non-instrumental learning.

For example, *Featherstone and McDonald* (*2004*) reported that both DLS and DMS lesions spare performance in a conditional-place preference task, whereas *Corbit and Janak* (*2010*) showed that both DLS and DMS lesions impaired the formation of Pavlovian associations. Moreover, some authors suggested that the DMS might play a role in reversal learning and cognitive flexibility, rather than mediating a specific learning strategy (*Ragozzino et al., 2002*; *Okada et al., 2014*).

Many studies have attempted to resolve the DMS-DLS controversy using the dual-solution plus maze task, in which both inflexible (i.e. response-guided) and flexible (i.e. spatial) learning strategies can be implemented. In this paradigm, rats typically prefer a spatial strategy in the early stages of training and gradually shift toward a response strategy following additional training. In accordance with results in instrumental conditioning, spatial strategy was shown to rely on DMS activity, while response strategy is mediated by the DLS (*De Leonibus et al., 2011*; *Yin et al., 2006*). So far, it is still unclear whether this gradual change in learning strategy is supported by a cooperation or a competition between the two striatal regions (*Bicanski and Burgess, 2020*). Some studies have indeed demonstrated that the DMS and the DLS may work in competition (*Malvaez et al., 2018*), since DLS lesion facilitates the learning of DMS-dependent-goal-directed spatial strategies (*Moussa et al., 2011*), and DMS lesion accelerates the acquisition of inflexible motor sequences (*Turner et al., 2022*).

Electrophysiological approaches (i.e. neural recordings in freely behaving animals) allow to go further in characterizing the interplay between the two subregions during learning processes, but the results are not fully consistent. Both DMS and DLS are active simultaneously during instrumental learning, but in some cases task-related activity is highly similar (*Stalnaker et al., 2010*; *Kimchi and Laubach, 2009*) whereas in other cases it strongly diverges (*Gremel and Costa, 2013*; *Cunningham et al., 2021*; *Thorn et al., 2010*; *Her et al., 2016*). Graybiel and colleagues were the first to report that during learning of a cue-guided T-maze task, the DLS develops preferential firing at the start and end points of the maze ('task bracketing') (*Jog et al., 1999*; *Smith and Graybiel, 2013*; *Thorn et al., 2010*; *Barnes et al., 2005*) , and this result was confirmed in other behavioral paradigms (*Sales-Carbonell et al., 2018*; *Yin et al., 2009*) . Task bracketing is thought to underlie bracketing of a sequence of action chains on a task, possibly by representing recently experienced reward locations to guide procedural actions (*Cunningham et al., 2021*). In contrast to DLS, learning-related activity in DMS is more variable and strongly depends on the behavioural task. In a conditional T-maze task DMS displays maximal activity at cue onset and movement onset, in contrast to the strong DLS ensemble activity at action boundaries (*Thorn et al., 2010*). In a Hebb-Williams task, DMS activity (and particularly posterior DMS) was shown to correlate with changes in reward prediction (*Stalnaker et al., 2012*) or changes in the reward delivery contingency (*Regier et al., 2015*), but neither task-bracketing activity nor choice-related activity was shown (*Cunningham et al., 2021*). Finally, *Peters et al.* (*2021*) reported that in a visually-guided wheel-turning task, DMS activity mainly correlates with the visual stimulus while the DLS displays motor responses.

Altogether, these results, although highly diverse, suggest that both DMS and DLS may support learning processes through complementary task-related activity. If this is the case, the same results should be generalized to different learning paradigms. In this study we aimed at investigating this issue using a continuous spatial alternation task with no cue instruction, thus limiting experimentally-defined discrete actions that possibly influence 'task bracketing' activity. DMS and DLS single neuron activity was recorded from the first day rats were exposed to the task until they reached and maintained stable performance. To decode animal behaviour from neural activity we used classical analyses based on either firing rate or neural synchronization, but we also combined the two measures using Hawkes process to reconstruct directed connectivity graphs of neurons. This approach enabled us to better unravel the coding capacity of both regions in the course of learning. We showed that DMS and DLS display different task-related activity throughout learning stages. The proportion of DMS coding neurons decreases whereas the number of DLS coding neurons increases over time, but this changes do not influence the capacity to decode animal behaviour from DMS and DLS neural networks. These results suggest that both DMS and DLS neural

94 networks are engaged during all learning stages but they gradually reorganize in different ways, in
95 contrast with the common assumption of a gradual shift from DMS to DLS across learning stages.

Results

Behavioral performance

Animal performance during each session was expressed as the number of paths traveled per
minute. Paths consisted of 2 correct and 10 incorrect trajectories (see Methods Section and Figure
1C-D) and were defined based on a previous study (*James et al., 2023*). Performance evolution
across training sessions was similar in all rats, with a gradual increase of the number of correct
trajectories and a decrease of the number of incorrect trajectories (Figure 1B). Based on the rate
of correct and incorrect paths (i.e. number of paths traveled per minute), we identified four succes-
sive stages of learning: stage 1, with a similar rate of correct and incorrect paths; stage 2, showing
a progressive increase of the rate of correct paths and a progressive decrease of the rate of incor-
rect paths; stage 3, in which the rate of correct and incorrect paths stabilizes; and finally stage 4,
showing almost exclusively correct paths.

Task-event coding neurons

We recorded 293 neurons in 4 DMS-implanted animals and 357 neurons in 3 DLS-implanted an-
imals over a total of 206 training sessions (Supplementary Figure 6). On the basis of the firing
rate and waveform properties (i.e., peak-valley distance and peak width at mid-height of the peak),
cluster analysis revealed the presence of two clear clusters (*Berke et al., 2004*) (see Methods sec-
tion and Supplementary Figure 8). Neurons with low firing rates and large waveforms were classi-
fied as putative medium spiny neurons (MSNs, GABA-ergic projection neurons), whereas the other
group was identified as putative fast-spiking interneurons (FSIs, GABA-ergic interneurons). A total
of 451 MSNs and 199 FSIs were found (Supplementary Figure 8). A chi-square test of independence
showed no significant difference in the proportion of FSI/MSNs between DMS and DLS (p-val. =
0.97).

To analyze possible differences in firing activity in DMS and DLS neurons across learning stages,
we first used a classical analytical approach and categorized neurons as coding task-events based
on their firing activity in specific time windows. We identified six 500ms events: the two onsets
O1 and O2 corresponding to the beginning of each path, the two intersections A and B of the
maze and the two rewards R1 and R2, corresponding to the end of the correct paths (just before
receiving the reward) (see Supplementary Figure 7 for a precise definition). Each of the six 500ms
events was divided into five 100ms intervals. We classified as task-event coding each neuron whose
firing activity in one 100ms interval varied significantly from all other intervals using a chi-squared
detection method (see Methods Section for a description of the statistical identification procedure).

We found 65,5% of task-event coding neurons in DMS (16% non coding neurons) and 38% in
DLS (24% non coding). 18.5% of DMS and 38% of DLS could not be classified because of too low
firing rate. The Pearson's Chi-squared test showed that the proportion of coding neurons is signifi-
cantly higher in the DMS compared to the DLS (p-val.=7.57e-06). The results from the general linear
model (with references: DLS and MSN) confirmed this effect (see the first column of Table 1). The
proportion of task-event coding neurons in the DMS is greater than in the DLS (adj. p-val.=0.06).
Interestingly, this proportion is stable across learning stages except for the stage 4 during which
the proportion of DLS task-event coding neurons increases (adj. p-val.=0.07 with positive coeffi-
cient (coeff. = 0.68) at learning stage 4), while the proportion of DMS task-event coding neurons
decreases (cross-effect adj. p-val. = 0.03, with negative coefficient (coeff = 0.57-1.86=-1.29)). Fi-
nally, in the DLS, MSN tend to code for task-events more than FSI (adj. p-val=0.03 with a negative
coefficient (coeff. = -0.52)), whereas in the DMS the difference between FSI and MSN is marginal
(cross-effect adj. p-val = 0.08 with coeff.= -0.52+0.57 = 0.05).
Figure 2.A shows the normalized average firing activity and the corresponding confidence in-

Figure 1. Behavioral data. **A**: Sketch of the maze with the two intersections A and B and the two feeders locations 1 and 2. **B**: Number of paths per minutes across learning stages. Each color corresponds to a path drawn with the same color in **C**. **C**: All the 12 considered paths with the 2 correct paths in the first column and the 10 incorrect paths in the second. Each path begins with an Onset O (1 or 2 depending on the feeder location). Only the correct paths end with a Reward R (1 or 2 depending on the feeder location). Under each path, the corresponding average duration in seconds for all rats and all sessions is shown.

Coding type		Task-event	Left-Right Turns	Full Paths
$\frac{\#\text{Coding}}{\#\text{Non Coding}}$ DMS|DLS		$\frac{192 \mid 136}{47 \mid 87}$	$\frac{102 \mid 42}{144 \mid 172}$	$\frac{226 \mid 182}{61 \mid 161}$
DMS Coeff. (adj. pval)		1.27 (0.06)	1.12 (0.05)	0.86 (0.03)
FSI Coeff. (adj. pval)		-0.52 (0.03)	-	-
Stage 2 Coeff. (adj. pval)		-	-	-
Stage 3 Coeff. (adj. pval)		-	0.97 (0.06)	-
Stage 4 Coeff. (adj. pval)		0.68 (0.07)	0.98 (0.06)	-
Cross-Effect DMS:FSI Coeff. (adj. pval)		0.57 (0.08)	-	-
Cross-Effect DMS:Stage 2 Coeff. (adj. pval)		-	-	-
Cross-Effect DMS:Stage 3 Coeff. (adj. pval)		-	-1.06 (0.08)	-
Cross-Effect DMS:Stage 4 Coeff. (adj. pval)		-1.86 (0.03)	-	-0.82 (0.08)

Table 1. Task-event coding neurons, neurons coding for left and right turns and neurons coding for full paths. First Line: On the top, number of coding neurons and below, number of non coding neurons, in the DMS (left) and DLS (right). The remaining lines report the results of a generalized linear model (see method section). The reference corresponds to DLS, MSN and learning stage 1. All p-values have been adjusted for multiplicity by Benjamini-Hochberg method. Only adjusted p-values < 0.1 corresponding to a False Discovery Rate less than 10% have been reported with their associated coefficient.

Figure 2. Z-score and Entropy. **A**: Average z-score of task-event coding neurons in DMS (red) and DLS (blue) in the four learning stages. Confidence bands at level 0.05 corrected for multiplicity over the 8 curves and the 30 time intervals (Bonferroni) are represented in dashed line. **B**: Difference in Z-score entropy between learning stages 2-to-4 and stage 1 (reference) for DMS and DLS. Confidence intervals at level 0.05 corrected for multiplicity over the 4 stages and 2 regions are pictured by a vertical line.

tervals at the different task-events, for neurons in the DMS (red) and DLS (blue) during the four learning stages. At the beginning of learning, during stage 1, the activity of DMS and DLS neurons decreases similarly during the R1 and R2 events. In stage 2, as behavioral performance improves, DMS and DLS neural activity starts to diverge. DMS neurons increase their firing rate particularly at the intersection B of the maze, before the rat turns into the left or right arm, whereas no significant change was observed at the intersection A, yet implying similar turning behavior. This result possibly indicates an involvement of the DMS in the implementation of the learning rule. As for the DLS, neurons change firing rate specifically at the reward areas during both R and O events. These modifications are similar to the DLS 'task bracketing' activity reported in previous studies (*Thorn et al., 2010*; *Cunningham et al., 2021*). Finally, no specific task-related activity is observed in the DLS during the later stages of learning, whereas the DMS tends to show some activity at action boundaries similar to DLS, during stage 4, as also shown by *Vandaele et al.* (*2019*).

To test whether these changes in neural activity during task-events may reflect changes in coding capacity across learning stages, we estimated the z-score entropy from the firing rates of the task-event coding neurons in the DMS and DLS. A decrease in entropy reflects a decrease in the randomness of the (renormalized) firing rate in the neural population, which is classically interpreted as an increase in the coding ability of the neuronal population (*Thorn et al., 2010*). As shown in Figure 2.B, the entropy of both DMS and DLS firing activity increases between stages 1 and 3, which may suggest a decrease in coding capacity during task acquisition for both areas. However, in contrast to the DMS, the entropy of the DLS tends to decrease in stage 4 to a level similar to that observed in stage 1. This may indicate that the DLS, but not the DMS, is involved in late learning stages following overtraining, as previously suggested (*Tang et al., 2009*). However, given the overlapping confidence intervals, it is difficult to draw a clear conclusion. It should also be noted that in this task animals are free to perform any trajectory and no specific instructions are given at any time. Thus, an event-based analysis may not be the most appropriate to reveal learning-related changes in neural activities. We therefore analyzed DMS and DLS capacity to code for specific paths (rather than events) across learning stages, as presented in the next paragraph.

Neurons coding for left and right turns

In classical event-based analysis, a change in firing rate is considered as a code for a specific event in the task (i.e., movement onset, reward, choice), but not for the task itself (here for example dissociating left and right turns). For this reason, we compared the firing activity of each neuron in the central stem (from A to B intersections) between left-turn and right-turn trials, including correct and incorrect paths. If the firing activity is significantly different between left-turn and right-turn trials, the neuron is classified as a left/right turns coding neuron (see the Methods section for a full description of the statistical method). Table 1 (second column) summarizes the results of the statistical analysis. As for the task-event coding neurons, we detected significantly more left/right turns coding neurons in the DMS than in the DLS (adj. p-val= 0.05), but no difference between FSI and MSN. Moreover the proportion of coding neurons tends to increase during learning stages 3 and 4 in the DLS (coeff. 0.97 or 0.98 and adj. p-val=0.06), whereas it is less clear for the DMS (stage 3: cross effect adj. p-val. = 0.08, coeff =0.97-1.06=-0.09).

We then focus on neural synchronization as a marker of task encoding (*Grammont and Riehle, 2003*; *Avital et al., 2013*) and we calculated for each pair of DMS or DLS neurons recorded simultaneously in each session, the probability of being synchronized at intersections A and B. We analyzed a total of 1821 pairs of neurons. The cumulative distribution functions (c.d.f.) of all p-values (one per neuron pair) are plotted in Figure 3 for DMS (red) and DLS (blue) neurons, at each learning stage. If the neuron pairs are independent, the c.d.f. should be below the diagonal; any positive deviation from the diagonal, especially for small p-values, reflects a lack of independence (i.e., the neurons are synchronized). The Kolmogorov-Smirnov test revealed a significant deviation from the diagonal for DMS neurons at stage 2, particularly at intersection B (p.adj = 8.6 e-03), whereas no significant effect was observed for DLS neurons at all stages. We then compared the c.d.f. of each

Figure 3. Synchronization analysis at intersections A and B. Synchronization between pairs of neurons was detected in 500ms time windows centered on the two maze intersections (A and B). Coincidences count was statistically compared to the empirical distribution of 50000 trial permutations. Cumulative distribution functions (c.d.f.) of the obtained p-values have been plotted as a function of the brain region (red for DMS and blue for DLS), intersection A or B and learning stages (color darkens as learning stage increases). A one sided Kolmogorov-Smirnov test of uniformity have been performed to compare all c.d.f. curves (16 in total). In addition, for each learning stage and maze intersection a two-sample one sided Kolmogorov-Smirnov test has been performed. All p-values were adjusted for multiplicity by Benjamini-Hochberg method. Note that c.d.f. of DMS p-values in stage 2 at intersection B is significantly above the diagonal (p.adj.= $8.6 \cdot 10^{-3}$). Moreover, in stage 2 c.d.f. of DMS p-values at A and B are significantly above c.d.f. of DLS p-values (intersection A: adj. p-val. = 0.027; intersection B: adj. p-val. = 1.7 e-04)

¹⁹² learning stage between DMS and DLS to assess whether DMS neurons were more synchronized
¹⁹³ than DLS neurons. We found that in stage 2, the c.d.f. for DMS neurons was significantly higher
¹⁹⁴ than the c.d.f. for DLS neurons at both intersections A (adj. p-val. = 0.027) and B (adj. p-val.=1.71e-
¹⁹⁵ 04). These results indicate that, overall, DMS neurons tend to fire simultaneously more than DLS
¹⁹⁶ neurons during initial learning. In addition, DMS neurons are significantly synchronized at inter-
¹⁹⁷ section B of the maze, before the rat turns into the left or right goal arm, especially during initial
¹⁹⁸ learning.

¹⁹⁹ Overall, these results demonstrate that DMS neurons code for left and right turns , particularly
²⁰⁰ during initial learning, an activity that may support the acquisition of the task rule. This hypothesis
²⁰¹ is reinforced by the observation that DMS neurons are synchronized during initial learning, par-
²⁰² ticularly before the rat chooses the left or right goal arm. In contrast, no specific synchronization
²⁰³ pattern was observed in the DLS at any learning phase.

²⁰⁴

Decoding animal behavior using Hawkes model

²⁰⁵

²⁰⁶ We then asked whether the previously identified coding properties were decisive for decoding an-
²⁰⁷ imal behavior. In other words, we asked whether the information provided by neural activity was
²⁰⁸ sufficient to guess the path taken by the rat (among the 12 identified). Neural activity is tradition-
²⁰⁹ ally modeled by a Poisson process based on the firing rate. However, we have shown that the
²¹⁰ synchronization of neurons, in addition to the firing rate, also reflects their coding capacity: we
²¹¹ therefore need a decoding model that takes into account both single neuron firing activity and
²¹² neural interactions. We chose to use the multivariate Hawkes model, which includes neural firing
²¹³ rate, as well as the strength and directionality of the interactions between neurons(*Lambert et al.,*
²¹⁴ *2018*; *Reynaud-Bouret et al., 2021*). Since our ultimate goal was to compare path decoding proper-
²¹⁵ ties between DMS and DLS, we limited the analysis to the neurons showing significantly different
²¹⁶ firing activity between any two paths (181 DLS neurons and 226 DMS neurons) (see third column
²¹⁷ of Table 1). The proportion of path coding neurons in the DMS is again greater than in the DLS
²¹⁸ (adj.p-val. = 0.03), and tends to decrease over learning stages (cross-effect adj. p-val.=0.08 with
²¹⁹ coeff.=-0.82) similarly to task-event and left/right turns coding neurons (see Table 1). Since no
²²⁰ significant difference was observed between MSN and FSI, both were included in the analysis.

²²¹ Figure 4 shows the interaction graphs reconstructed from the Hawkes model versus the Poisson
²²² model (vertical line) during a representative session for a DMS- (red) and a DLS-implanted rat (blue).
²²³ The reconstructed graphs for two different paths from the same session are shown on the same
²²⁴ line. The color code represents either the firing rate for the Poisson model or the spontaneous part
²²⁵ for the Hawkes model (i.e., baseline activity with all interaction functions canceled). For the Hawkes
²²⁶ model, the directionality and strength of the interactions between two neurons are represented
²²⁷ by the direction and color of the arrows, respectively. In this example, we can clearly notice that,
²²⁸ for the same set of neurons, the graphs for the two paths are different while their firing activity is
²²⁹ roughly the same.

²³⁰ To evaluate the performance of the two models in decoding animal behavior, we first calculated
²³¹ the difference between the decoding power of the Poisson model, the Hawkes model, and a ran-
²³² dom guess. Decoding power is defined as the percentage of correct identification of the traveled
²³³ paths, calculated by estimating the models on 2/3 of the paths in each training session and testing
²³⁴ the model predictions on the remaining 1/3. As shown in Figure 5.A, the Hawkes model has bet-
²³⁵ ter decoding power than the Poisson model, which predicts rat performance as well as a random
²³⁶ guess. This may be due to the short path lengths (see Figure 1.B) combined with a fairly low firing
²³⁷ rate of the neurons, which results in the prediction being based on 10-20 spikes per neuron in a
²³⁸ given session. By taking into account other parameters of neuronal activity (i.e., the direction and
²³⁹ strength of neuronal interactions), the Hawkes model is able to decode more efficiently the paths
²⁴⁰ taken by the rats.

²⁴¹ We then compared the decoding power computed by the Hawkes model in DMS and DLS across

Figure 4. Examples of estimated network models for DMS (in red) and DLS (blue) neurons recorded from one DMS-implanted rat and one DLS-implanted rat during two training sessions (session 11 for DMS and session 6 for DLS) and two different paths (a correct path on the left and a straight path on the right). Each node represents one neuron. The color of the node represents either the average firing rate for the Poisson model (first vertical line on the left) and the spontaneous part for Hawkes model (with all interactions canceled). The color and directionality of the lines represent the interaction function. Right panel: color code for the interaction function, with the number of spikes modified by this interaction.

A

B

Figure 5. Decoding power. **A**: Boxplot showing the difference in decoding power (see Methods section for the precise definition) between the random guess (1/Nb of Paths), the Poisson model and the Hawkes model, for all rats and training sessions. **B**: Boxplot showing the Hawkes decoding power for DMS and DLS neurons across learning stages (red for DMS and blue for DLS). A beta regression with probit link is performed to compare DMS and DLS across learning stages. All p-values have been adjusted for multiplicity with Benjamini-Hochberg method. The colored dots represent the value predicted by the regression model for each boxplot. Significant adjusted p-values are indicated in the figure.

learning stages (Figure 5.B). To do so, we performed a beta regression with probit link (and explana-
tory variables: 'number of paths', 'number of neurons', 'average firing rate', 'DMS-DLS', 'learning
stage' and 'DMS-DLS'/'learning stage' cross-effects) to disentangle the effects of interest ('DMS-DLS'
and 'learning stage') from other auxiliary effects we need to consider. The results show that, in gen-
eral, the decoding power in the later learning stages is greater compared to stage 1 (stage 3: adj.
p-val.=2.11e-02 ; stage 4: adj. p-val.=1.29e-02). The number of paths (adj. p-val.=4.7e-12) and neu-
rons (adj. p-val.=2.3e-07), as well as their average firing rate (p.adj= 1.3e-02) also have a significant
effect, but no cross effect was observed. As for the comparison between DMS and DLS, the results
show that the decoding power of DMS is globally greater than that of DLS (p.adj=2.8e-03), and par-
ticularly during learning stage 2 (adj. p-val.=3.92e-06). Note that the network size effect (number
of neurons) has been taken into account in the model and cannot explain such difference. These
results show that both the DMS and DLS neural networks undergo a gradual reorganization during
learning, since the decoding power of the network increases as the rats performance improves in
both DMS and DLS. However, it should be noted that DMS shows greater decoding power relative
to the DLS during initial learning, possibly indicating that the acquisition of a spatially guided task
rule without any temporal or cue instruction requires DMS neural activity.

Discussion
In this study we aimed at characterizing the functional difference between DMS and DLS in a spon-
taneous and continuous spatial alternation task, in which rats learn to execute spatial trajectories
in the absence of any specific instruction. Our results show:
a) more coding neurons (for events, paths and left/right turns) and stronger decoding power in
the DMS compared to the DLS during all learning stages, suggesting that the DMS is more involved
than the DLS in learning and performing a spatial alternation strategy;
b) at the level of single neurons, DMS and DLS coding properties strongly differ at the beginning
of learning, with the DMS mainly activated (and synchronized) at the maze intersection before the

rat turns into one of the goal arms, and the DLS activating at action boundaries; with extensive
training both DMS and DLS tend to activate at action boundaries ;
c) at the level of neural population, DMS and DLS network properties as well as their modification
over learning highly diverge since the proportion of coding neurons across learning increases in
the DLS and decreases in the DMS, and the evolution of the entropy of the population firing rate
follows opposite trends in the two areas;
d) despite these differences, the decoding power of the network increases across learning stages
in DMS and DLS, suggesting an involvement of both striatal structures in supporting behavioral
performance from acquisition to extensive training.
Altogether these results demonstrate that at the level of DMS and DLS neural networks, the task-
related neural coding and the ability to decode animal behavior are decorrelated, since DMS de-
creases and DLS increases its coding capacity across learning, but their ability to decode animal
behavior increases in both structures. This suggests that the two striatal areas provide complemen-
tary information necessary to support both the acquisition and the implementation/optimization
of a behavioral strategy.
282
DMS vs DLS single neuron coding properties across learning stages
The DMS is generally involved in explicit action-outcome associations whereas the DLS has a
major role in skill learning and habit formation (*Balleine and O'Doherty, 2010*). Accordingly, the DLS
was shown to activate at action boundaries or start/stop sequences of a task (*Jog et al., 1999*; *Yin
et al., 2009*), whereas DMS activity is more variable but tends to focus at cue (instruction) presenta-
tion as a prediction for future actions or rewards (*Thorn et al., 2010*; *Peters et al., 2021*). Here we
demonstrate that in a continuous and non cued T maze alternation task, DMS shows the greatest
proportion of coding neurons across all learning stages. This is observed using a classical analytical
approach based on single-unit firing rate, and was confirmed for all types of task-related coding (i.e.
task event coding, path coding or left/right turn coding). Interestingly, during initial learning DMS
neurons tend to activate and synchronize at the intersection of the maze just before the rat choose
to turn into the left or right arm. Neural synchronization has been related to stimulus encoding
in different cortical areas (*Grammont and Riehle, 2003*; *Avital et al., 2013*) and can be possibly
generalized to the encoding of abstract information such as the learning rule. Therefore, the DMS
activity pattern we observe at maze intersection possibly reflects an implication of the DMS in the
acquisition of the spatial rule or could even sustains deliberative processes (i.e. decision making).
It should be noted that previous studies have ascribed a similar role to the hippocampus and the
prefrontal cortex as well as their interaction (*Hassabis et al., 2007*; *Benchenane et al., 2010*; *Spiers
and Gilbert, 2015*). Our result supports the hypothesis that DMS could be part of a network medi-
ating decision making in spatially guided behaviors (*Redish, 2016*).
In contrast to the DMS, DLS neurons during early learning show mainly activation at action bound-
aries, as previously described in a cued (i.e. non spatial) T maze (*Jog et al., 1999*; *Thorn et al., 2010*)
and in go-nogo tasks (*Jin and Costa, 2010*). This 'bracketing activity' is supposed to improve the ac-
quisition of behavioral routines and promote habit learning. Here we observe a similar 'bracketing'
activity in the DLS but in the absence of any specific instruction, thus suggesting that a decompo-
sition of the global task in multiple sub-tasks is at play even in the absence of any clear start-stop
signals. This hypothesis is sustained by a recent study in which we showed that in the continuous
T-maze task rats tend to use a behavioral strategy that consists in chunking sub-actions to perform
complete paths (*James et al., 2023*). Altogether these results point to a role of the DLS in identifying
sub-actions or sub-tasks not exclusively in the context of habit learning.
Thus, at the beginning of learning when rats are acquiring the learning rule, DMS and DLS provide
different information that possibly allows to choose which behavioral strategy will be performed.
Accordingly, we have previously shown that both DMS and DLS are involved in the acquisition of a
spatial alternation task and compete for the control of the behavioral response at the beginning of
learning, since DMS lesions impaired and DLS lesions accelerated the acquisition of the behavioral

strategy (*Moussa et al., 2011*). However, as learning progresses, DMS and DLS single neurons show
similar coding properties since both tend to activate at action boundaries in learning stage 4. Using
a lever-press task designed to promote skill learning, *Vandaele et al.* (*2019*) also reported that both
DMS and DLS activate at the start/end of action sequences following over training. However, since
start and end were also marked by overt cues (lever insertion and retraction), the activity in the
two areas could still diverge after over training with DMS reflecting cue coding and DLS reflecting
action boundaries (*Thorn et al., 2010*). This hypothesis is nonetheless confuted by our result show-
ing similar DMS and DLS activity at action boundaries in a task with no explicit cue instruction, thus
suggesting that single neurons in the two areas code similar information in later training stages.
Despite the progressive similarity in single neuron coding properties, the activity of the neural pop-
ulation in the DMS and DLS are highly divergent throughout learning stages. Indeed, we observe
that the proportion of coding neurons decreases in the DMS and increases in the DLS across learn-
ing sessions. Similarly, the entropy of the neural ensemble firing rate increases in the DMS and
decreases in DLS. Taken together these results indicate that DMS and DLS neural networks diverge
across learning, with the DLS progressively increasing its coding capacity and the DMS following an
opposite trend. It is generally believed that the optimization of a learning strategy relies on a grad-
ual shift of activation from the DMS to the DLS neural networks. Such strategy optimization does
not imply necessarily the development of habitual actions, since greater DLS activation compared
to the DMS is also observed following over-training of non-habitual behavioral responses (*Peters*
et al., 2021). Our results show that the two networks evolve in a way that could be in accordance
with this hypothesis. However, the proportion of coding neurons is still greater in DMS after over
training compared to the DLS, and this result may indicate a more complex interaction between
DMS and DLS across learning.
341
Decoding rat behavior from DMS/DLS network activities using Hawkes processes
Another approach to infer neural network function in learning processes is to decode animal
behavior from neural activity. Therefore, instead of looking for the conditions for which neural
activity changes, we analyzed how well we can reconstruct animal performance based on neural
activity. To do so, we used a classical Poisson model based exclusively on single neuron firing rate,
and the Hawkes model that takes into account both single neuron firing rate and the interactions
between neurons as well as the directionality of this interaction. This model allows to reconstruct
a graph of functional connectivity between simultaneously recorded neurons, by analyzing how
much of the activity of a neuron is explained by the activity of the surrounding neurons. There-
fore, both the strength of the connections between neurons and the direction of the interaction
are represented in the graph. We observe that compared to the Poisson model the Hawkes model
better decodes animal performance, indicating that neural interaction is a strong component of
the striatal network function.
Using this analytical approach we show that despite the progressive decrease in the proportion of
coding neurons, the DMS network has higher decoding power than the DLS throughout learning
stages. This result supports the hypothesis of a more selective implication of the DMS compared
to the DLS in a continuous spatial alternation (*Moussa et al., 2011*). However, it should be noted
that the decoding power of both DMS and DLS increases over training, suggesting that as learn-
ing progresses neural activity in both areas conveys information necessary to support behavioral
performance. It is interesting to note that the evolution of task-responsive neurons and decod-
ing power across learning stages can be decorrelated. Specifically, the proportion of task-related
coding neurons decreases in the DMS but the network itself exhibits enhanced decoding capacity.
Conversely, the number of coding neurons and the decoding power increases over time in the DLS.
Thus, an increasing proportion of coding neurons does not necessarily imply an increased ability
to decode animal behavior from the same neural activities.
In this context, it is necessary to define which behavior is decoded. Here we consider the decoded
performance in terms of elementary paths, as described in Figure 1. In a recent study we have

shown that rats learn a continuous alternation strategy by subdividing each of the identified paths into sub-paths or sub-actions (*James et al., 2023*). Thus, the learning strategy consisted in chunking the different sub-actions, similarly to what was previously suggested for the acquisition of complex tasks (*Botvinick et al., 2009*). We have tried to use the Hawkes model on the different sub-actions. However, given their short duration, the number of spikes per trial was not sufficient to perform the analysis. In the future it would be interesting to design behavioral experiments where the sub-actions are long enough to allow to use the Hawkes decoding approach. Alternatively, it would be useful to increase the number of recorded neurons. However, it should be noted that we obtained very good decoding results despite a relative low number of neurons recorded per session, which indicates that the striatal network is highly redundant (*Reynaud-Bouret et al., 2021*).

Conclusions

In summary, we show that inferring neural network functions based exclusively on task-related firing activity is insufficient. By incorporating the Hawkes model to decode behavior from neural network activities, we were able to achieve a more comprehensive description of DMS and DLS functions across learning. Altogether our results show that the coding pattern evolves differently in the two striatal areas over training, but both supports learning from early to late stages. This was shown in the context of spatial tasks. It will be interesting to use the same analytical approach in instrumental tasks to test whether these striatal network properties could be generalized to all types of learning processes.

It remains to be determined which is the exact nature of DMS and DLS interactions during learning. One possibility is that DMS and DLS activity supports distinct learning strategies and control action selection through a competitive mechanism. The results we obtained at early stages of learning supports this hypothesis, but the increased engagement of the two areas across learning stages is not coherent. Alternatively, we may speculate that either DMS- and DLS- mediated action selection process occurs exclusively at the beginning of the learning, or that action selection is implemented by the downstream basal ganglia circuits based on the complementary information provided by the DMS and the DLS. Once the behavioral strategy has been implemented, both striatal areas may be implicated in maintain the performance, that may explain why decoding power in both areas increases over training.

Methods and Materials

Animals and surgery

Seven male Long-Evans rats (Janvier, Le Genest-St-Isles, France) weighing 300-350 g were housed in individual cages (40 cm long x 26 cm wide x 16 cm high) with food and water ad libitum and maintained in a temperature-controlled room (20°C +/- 2). One week after their arrival, animals were handled daily by the experimenter for 7 days. 4 animals were then implanted with tetrodes aimed at either the left (3 rats) or the right (1 rat) dorsomedial striatum (DMS) at the following coordinates: AP: ±1mm, ML: ±2.2 mm from the midline, DV: -3mm below the dura. Two animals were implanted with tetrodes aimed at the left dorsolateral striatum (DLS) and one rat was implanted bilaterally at the following coordinates: AP: ±1mm, ML: ±3.7 mm from the midline, DV: -3mm below the dura 6). The surgery was performed under sterile conditions and under general anaesthesia (Ketamine 75 mg/kg (Imalgene 1000, Merial, France)/Medetomidine 0.25 mg/kg (Domitor, Janssen, France)). As postoperative treatment, the rats were injected with antibiotic (Clamoxyl, 150 mg/kg) and analgesic (Tolfedine, 4 mg/kg). After surgery, the rats were given 5-7 days of recovery. They were then subjected to a food deprivation program that kept them at 90 % of their body weight during behavioral testing. All experiments were performed in accordance with the National Institute of Health's Guide for Care and Use of Laboratory Animals (NIH Publication no. 80-23) revised in 1996 for the UK Animals (Scientific Procedures) Act of 1986 and associated guidelines or the Policy on Ethics approved by the Society for Neuroscience in November 1989 and amended in Novem-

418 ber 1993 and under veterinary and National Ethical Committee supervision (French Agriculture
419 Ministry Authorization)

Microdrives and recording setup

Four tetrodes formed a bundle threaded through a piece of stainless-steel tubing. Each tetrode
consisted of four twisted 25µm nichrome wires. The connector, tubing and wires could be moved
downwards by turning the drive screw assemblies cemented to the skull. Cable was connected to
the rat's headstage, which contained a field effect transistor amplifier for each wire. The signals
from each tetrode wire were amplified 10 000 times, bandpass filtered between 0.3 and 6 kHz
with Neuralynx amplifiers (Neuralynx, Bozeman, MT, USA), digitised (32 kHz) and stored by the
DataWave Sciworks acquisition system (DataWave Technologies, Longmont, CO, USA). A red light-
emitting diode (LED) attached to the head assembly was used to determine the position of the rats.
The LED was filmed by a CCD camera mounted on the ceiling above the maze, and its position was
tracked at 25 Hz by a digital point tracker.

T maze apparatus

The T-maze consisted of four 10 cm wide, grey-painted wooden tracks (with walls of 2 cm height
on each side), a 100 cm long central stem, a 100 cm long crossbar forming the two goal arms,
and two additional tracks each connecting the distal end of one goal arm to the base of the cen-
tral stem. The reward wells were located at the distal end of each choice arm. Food rewards (45
mg sugar pellets) were released from two feeders (MedAssociates) mounted above the wells and
activated by remote manual switches. The maze was elevated 40 cm off the ground on a metal
frame. The apparatus was illuminated by four symmetrical spotlights (40 W) mounted on the ceil-
ing. A centered radio above the maze was used to mask uncontrolled disturbing sounds and the
experimenter was located in an adjacent room. Behavioral training was performed as follows. Af-
ter one week of recovery period from surgery, the rats were familiarized with the maze in daily
20-minutes sessions for two days, during which they were allowed to freely explore the apparatus
and collect randomly scattered sugar pellets. Training began on the third day with either one or
two 20-minutes sessions per day. Animals were gently placed at the intersection point labelled 'A'
in Figure 1.A of the 'continuous T-maze' and let free to explore each arm as they wished. In order
to obtain a 15 mg sugar pellet, they had to run along the central stem and alternately enter the
left or right arm of choice. Rats were submitted to one training sessions per day. Tetrodes were
lowered by either 25 μm or 50 μm after each training session or every two sessions, for a maximal
length of 2mm (between 1 and 2mm, Supplementary Figure 6).

Data Analysis – behavior

Animal position (x/y coordinates) over time was sampled at 25 Hz. A preprocessing of the data,
consisting mainly in clearing rapid head movements and reflection artifacts of the camera, was
first performed. We identified 12 possible elementary paths (i.e. 2 correct paths and 10 incorrect
paths) that rats could travel along the maze (Figure 1.C). Note that some rare portions of the trajec-
tories were not classified as elementary paths and were removed from the analysis. These mainly
include animals jumping out of the maze or jumping in a non contiguous spot in the maze. Ani-
mal performance was expressed as the number of paths (either correct or incorrect) per minute
travelled by each rat during the training sessions. Based on rat performance, we identified four
successive learning stages : Stage 1, with a similar number of correct and incorrect paths; Stage 2,
showing a progressive increase of the number of correct paths and a progressive decrease of the
incorrect ones; Stage 3, in which the number of correct and incorrect paths stabilize; and finally
Stage 4, showing almost exclusively correct paths (Figure 1.C). The duration of each path type was
also scored.

Data analysis – neuronal activity

Neural data from the four DMS implanted rats were also used in another study investigating rat learning strategies using credit assignment models (*James et al., 2023*). All the results reported in the present study are original.

Spike Sorting

Spike sorting was performed manually using the graphical cluster-cutting software Offline Sorter (Plexon). Units selected for analysis had to be well discriminated clusters with spiking activity clearly dissociated from background noise. Units that were lost or whose waveform changed too much before the session was completed, were excluded. Units having interspike intervals <2 ms were removed due to poor isolation, as were cells with a peak firing rate <1 Hz. A total of 650 cell clusters was accepted (293 DMS and 357 DLS units). Since the tetrodes were lowered at the end of the recording session, neural activity in each session should be collected from different neurons. However, if two neurons recorded in two successive sessions showed similar characteristics in their firing patterns, waveforms and clusters, they were included only once.

MSN and FSI classification

For each neuron we computed the log of the global firing rate, the peak-valley distance (PV) (i.e. the distance between the peak and next minimal value) and the peak width at half-height (W) from the digitalized waveforms. These three variables were next centered and renormalized. Then we used the hierarchical clustering method with 'Ward 2' distance to build the dendrogram, showing clearly the existence of 2 clusters (Supplementary Figure 8). The k-means cluster algorithm with $k = 2$ was used to identify the clusters. The cluster with small firing rate and large PV and W corresponded to putative medium-spiny neurons (MSN), whereas points with large firing rate and small PV and W were allocated to the putative fast-spiking interneurons (FSI) cluster.

Identification of coding neurons

To identify coding neurons we performed three different analysis that were based on either temporal or spatial firing activity. In the first analysis, a task-event coding neuron was defined as a neuron showing significantly different firing activity during any of the six task-related temporal events (described below). For the second analysis (left/right coding neurons), we focused on the neural firing within the central stem of the maze with respect to whether the rat will subsequently chose the left or right goal arm. In this case, a coding neuron should activate differentially for left and right turns. The third analysis allowed us to categorize path coding neurons as neurons showing significantly different firing activity between any two of the 12 paths described in the behavioral section. Path coding neurons were then used in the decoding analysis. The results from these analysis were used to compare the proportion of coding neurons across learning stages, brain areas (DMS - DLS) and types of neurons (FSI - MSN). Whatever the type of coding, the detection of coding neuron is the same and is described hereafter.

Chi-squared method for detection of coding neurons. The purpose of this statistical analysis is to test whether the firing activity is constant over all possible conditions (null hypothesis) or if it is significantly different in at least one given condition (i.e. a 100ms bin or a particular path) with respect to all other possible conditions. These possible conditions will include the 30 temporal bins defining the six task events in the first analysis (i.e. five 100ms-bins per event), or the two left and right paths for the second analysis, or the paths taken in a session for the third analysis. Given N_k the number of spikes produced in a given condition $k = k, ..., K$, we compute the p-value of the

chi-square test that decides if $(N_1, ..., N_K)$ is a multinomial distribution of parameters

$$p = \left(\frac{d_1}{d_1 + ... + d_K}, ..., \frac{d_K}{d_1 + ... + d_K} \right)$$

and

$$n = N_1 + ... + N_K,$$

with d_k, the duration of the condition k in the session. The test gives a valid p-value if $np_k \geq 5$ for
all coordinates k. Therefore if it is not the case, we can either drop a condition or group together
conditions. If despite that, there is still not enough spikes, the corresponding neuron was removed
from the analysis.
All neurons with a Benjamini-Hochberg (BH) adjusted p-value lower than 0.05 were declared as
coding for a particular set of conditions. To assess the difference in the proportion of coding neu-
rons (see Table 1) we then used a generalized linear model (Bernoulli with probit link) to explain the
coding character as a function of brain region, neuron type and learning stages (with cross effects
Brain region-Learning stage and Brain region-neuron type). Each time, DLS, MSN and Stage 1 were
used as references. In a preliminary study, firing rates and all other possible cross-effects were
added to the model, but since no significant results even without correction were found, they have
been discarded. The fact that the firing rate does not impact the coding character is in adequation
with the chi-square detection method: as soon as there are enough spikes in total in the session
to use the method, there is no bias *per se* toward largest firing rates, once the FSI character has
been taken into account.

• *Task-event coding neurons.* To identify event-based coding properties, we analyzed neural fir-
ing during specific time windows in four areas of the maze : the two intersections (depicted
as A and B in Figure 1.A) and the two reward areas (depicted as R1 and R2 in Figure 1.A).
We defined six 500ms time events based on the passages through the different areas (see
Supplementary Figure 7). For the two intersections A and B, we extracted the central 500 ms
time window of the trajectory within the two areas. For the reward areas we defined two
different events: the reward obtention (R1 and R2 for the left and right goal arm) including
the first 500ms after the rat entered the reward areas, and the movement onset (O1 and O2
for the left and right goal arm) including the last 500ms before the rat left the reward areas
(Supplementary Figure 7). For each neuron we calculated the average firing rate during the
500ms events by refining the time window in 5 bins of 100ms. A task-event coding neuron
is a neuron that has been detected by the chi-square method to show a firing rate activity in
one 100ms bin that is different over all 30 time bins (i. e. 5 bins for each of the 6 events). If
the condition $np \geq 5$ is not fulfilled, we have grouped the 100ms bins into 6 conditions, corre-
sponding to 6 time windows of 500 ms around each event (intersections, rewards, onsets). If
the condition is still unfulfilled the neuron is discarded.
         The z-score of an event coding neuron n in a 100ms bin of time t is given by

$$z_{n,t} = \frac{f_{n,t} - \bar{f}_{n,.}}{\sigma(f_{n,.})},$$

539   where $f_{n,t}$ is the firing rate of neuron n in bin t, $\bar{f}_{n,.}$ is the average over the bins and $\sigma(f_{n,.})$ the
540   corresponding standard deviation over the bins.
541   Then, we averaged the normalized (z-score) activity of DMS and DLS event coding neurons
542   over each neuronal population - brain region*stage - (see Figure 2.A).
         We also estimated the entropy of the spike probability distribution in each population from
         the ensemble z-scores across learning stages by counting the number $\mathbf{Nz}_I$ of $z_{n,t}$ in a given
         population that appears in an interval I of z-score given by segmenting $[-3.5, 5.5]$ in bins of

0.5 length, then the entropy of the distribution of z-score was estimated by

$$\sum_{I\,s.t.\mathbf{Nz}_I \neq 0} \frac{\mathbf{Nz}_I}{\mathbf{Nz}_{tot}} \log\left(\frac{\mathbf{Nz}_I}{0.5\mathbf{Nz}_{tot}}\right),$$

where $\mathbf{Nz}_{tot}$ is the total number of z-score in the population. Note that $\frac{\mathbf{Nz}_I}{0.5\mathbf{Nz}_{tot}}$ estimates the density of the z-score distribution inside interval I.

- *Neurons coding for left and right turns.* Here we focused on DMS and DLS neural firing activity in the central stem of the maze (between A and B intersections, Figure 1.A) with respect to whether the rat will subsequently choose the left or right goal arm. Here the chi-square method is performed only with two conditions and there is no grouping or dropping of conditions if $np \geq 5$ is not fulfilled. In this case, the neuron is just discarded.
- *Neurons coding for paths.* Firing activity in DMS and DLS during any of the twelve possible paths was calculated. We use again the chi-square method to identify significant spike activity in one of the twelve paths over all paths. If any of the path k does not satisfy $np_k \geq 5$, this condition is simply left out. This is in particular the case if $d_k = 0$, meaning that this path is not taken by the rat during the session. If even with only two paths left, the condition $np \geq 5$ is not satisfied, the neuron is discarded.

Neuronal synchronization at maze intersections

For each learning session, we performed a trial shuffling / permutation analysis to detect synchronization of spiking activity between pairs of simultaneously recorded neurons at the intersection areas of the maze A and B (see *Albert et al.* (*2015*, 2016)). More precisely, for any pair of neurons simultaneously recorded, we isolated the number of times the rat passed through A or B and for each passage we extracted the central 500 ms window, defined as a trial. The number of coincident spiking activity observed within 20ms in each trial was compared to the empirical distribution obtained by computing 50 000 permutations of the trials, to obtain a p-value. We then calculated the cumulative distribution functions (c.d.f.) of all p-values from pairs of neurons in the DMS and the DLS across learning stages. We quantified the deviation of the c.d.f. from the diagonal by a Kolmogorov-Smirnov test of uniformity and the comparison between DMS and DLS c.d.f.s by a two-sample Kolmogorov-Smirnov test.

Decoding behavior using Poisson and Hawkes models

Decoding of rat behavior (i.e. choosing a particular path among the 12 identified, Figure 1.C) was implemented using a Poisson and Hawkes model. We chose Poisson model since neural firing is generally modeled in that way. In addition, we used the less conventional multivariate Hawkes model because it includes not only firing rate but also the strength and the directionality of the interaction between pairs of neurons (*Lambert et al., 2018*; *Reynaud-Bouret et al., 2021*). Since we focused on path decoding, we included in the analysis exclusively DMS and DLS path coding neurons. In particular, this prevents the method to be diluted by a different proportion of non coding neurons in each area.

The Poisson model is defined by an intensity (instantaneous firing rate) of neuron n in condition (path) k given by

$$\lambda_{n,k}(t) = \nu_{n,k}.$$

The parameter $\nu_{n,k}$ represents the firing rate of neuron n in condition k. Hence if N is the total number of path coding neurons in the session, the Poisson model has N parameters per condition.

The Hawkes model is defined by an intensity of neuron n in condition (path) k given by

$$\lambda_{n,k}(t) = \mu_{n,k} + \sum_{m=1}^{N} \sum_{T\,\text{spikes of neuron }m, T<t} h_{m\to n}(t - T).$$

582 The parameter $\mu_{n,k}$ represents the spontaneous firing rate of neuron n in condition k. In addition,
583 the interaction $h_{m \to n}$ quantifies the impact of one spike of neuron m at time T on the apparition
584 of a spike on neuron n. The functions $h_{m \to n}$ are decomposed into 6 plateaux of length 10 ms each
585 associated with a different value $a_{m,n,1}, ..., a_{m,n,6}$, so that the value of $h_{m \to n}$ between 10 and 20 ms is
586 for instance $a_{m,n,2}$. The strength of an interaction is then measured by $|h_{m \to n}| = \sum_{i=1}^{6} |a_{m,n,i}|$. In total,
587 $N + 6N^2$ parameters per condition are necessary for a Hawkes process. In particular, it includes
588 self-interactions.

In each model, the parameters are estimated by minimizing the least-square contrast that is

$$C(\lambda_{n,k}) = -2 \sum_{T \text{ spikes of neuron } n} \lambda_{n,k}(T) + \int \lambda_{n,k}(t)^2 dt.$$

589 More precisely, for Figure 4, the sum and integral above are considered over all time T or t such
590 that condition k holds. The estimated parameters $\nu_{n,k}$ for the Poisson and $\mu_{n,k}$ for the Hawkes are
591 given by the color of the nodes. The strength of the interaction $h_{m \to n}$ is given by the color of the
592 edge $m \to n$.

The computation of the decoding power is more intricate. In each session, we consider only the paths that are taken at least 3 times. We defined as a trial each single path (and the corresponding neural activity) taken by the rat. So for condition (path) we have at least 3 trials. For each condition k, we computed estimators of the two models by using only 2/3 of the trials, leading to estimators of the intensity $\hat{\lambda}_{n,k}$. To do so, it is sufficient to restrict the integral and summation in the least-square contrast $C(\lambda_{n,k}) = C_{2/3}(\lambda_{n,k})$ to time inside 2/3 of the trials. For each condition k, it remains 1/3 of the trials. For each remaining trial, we compute the least-square contrast $C(\hat{\lambda}_{n,k'}) = C_{trial}(\hat{\lambda}_{n,k'})$, for all condition k', where now the time in the sum and integrals in C is restricted to this precise trial. Therefore $C_{trial}(\hat{\lambda}_{n,k'})$ measures the similarity between the model estimated in the condition k' and the spike train of the trial under inspection, which is in condition k. The estimated condition of a given trial is therefore given by

$$\hat{k} = \arg\min_{k'} C_{trial}(\hat{\lambda}_{n,k'}).$$

593 The decoding power of a model is the proportion of good guess, that is the proportion of trials for
594 which $\hat{k} = k$.

595 The decoding power strongly depends on the number of paths (Nb Paths) within a session.
596 Indeed, if there is not enough information in one path to decode rat behavior, the decoding power
597 should be equivalent to 1/ Nb Paths. Therefore, we used this value as reference (i.e. random guess),
598 that was used to assess the differences in decoding power between Poisson and Hawkes model in
599 Figure 5.

600 Then , we calculated the decoding power for each striatal region across the four learning stages.
601 As previously noted, the decoding power depends on the number of paths (Nb Paths) but also on
602 the number of (path coding) neurons (Nb Neurons) and their average firing rates (Firing). Moreover,
603 the encoding power is a quantity in $[0, 1]$ that is best modeled by a beta variable. Therefore we
604 performed a beta regression with probit link to test the effect of Nb Paths, Nb Neurons, Firing,
605 Striatal Region (i.e. DMS/DLS), Learning Stage and cross effects Region/Stage, to disentangle the
606 effects of interest (Region, Stage) and the other auxiliary effects that we need to take into account.

Acknowledgments

607

608 This work was supported by the French government, through the UCA^{Jedi} and 3IA Côte d'Azur In-
609 vestissements d'Avenir managed by the National Research Agency (ANR-15- IDEX-01 and ANR-19-
610 P3IA-0002), by the interdisciplinary Institute for Modeling in Neuroscience and Cognition (Neuro-
611 Mod) of the Université Côte d'Azur and directly by the National Research Agency ANR-08-JCJC-0125-
612 01 and ANR-19-CE40-0024 with the ChaMaNe project. It is part of the Computabrain project.

References

Albert M, Bouret Y, Fromont M, Reynaud-Bouret P. Bootstrap and permutation tests of independence for point processes. . 2015; .

Albert M, Bouret Y, Fromont M, Reynaud-Bouret P. Surrogate data methods based on a shuffling of the trials for synchrony detection: the centering issue. Neural Computation. 2016; 28(11):2352–2392.

Avital A, Finkes I, Katabi S, Zhu Y, Teixeiria K, Prut Y, Bergman H. Encoding by Synchronization in the Primate Striatum. Jounal of Neuroscience. 2013; 33(11):4854–66.

Balleine BW, Dickinson A. Goal-directed instrumental action: contingency and incentive learning and their cortical substrates. Neuropharmacology. 1998; 37(4-5):407–419.

Balleine BW, Liljeholm M, Ostlund SB. The integrative function of the basal ganglia in instrumental conditioning. Behavioural Brain Research. 2009; 199(1):43–52.

Balleine BW, O'Doherty JP. Human and rodent homologies in action control: corticostriatal determinants of goal-directed and habitual action. Neuropsychopharmacology. 2010; 35(1):48–69.

Barnes TD, Kubota Y, Hu D, Jin DZ, Graybiel AM. Activity of striatal neurons reflects dynamic encoding and recoding of procedural memories. Nature. 2005; 437(7062):1158–1161.

Benchenane K, Peyrache A, Khamassi M, Thierney PL, Gioanni Y, Battaglia FP, Wiener SI. Coherent Theta Oscillations and Reorganization of Spike Timing in the Hippocampal- Prefrontal Network upon Learning. Neuron. 2010; 66(6):921–36.

Berke JD, Okatam N, Skurski J, Eichenbaum HB. Oscillatory entrainment of striatal neurons in freely moving rats. Neuron. 2004; 43(6):883–96.

Bicanski A, Burgess N. Neuronal vector coding in spatial cognition. Nature Reviews Neuroscience. 2020; 21(9):453–470.

Botvinick MM, Niv Y, Barto AG. Hierarchically organized behavior and its neural foundations: A reinforcement learning perspective. Cognition. 2009; 113:262–280.

Corbit LH, Janak PH. Posterior dorsomedial striatum is critical for both selective instrumental and Pavlovian reward learning. Eur J Neurosci. 2010; 31:1312–1321.

Cunningham PJ, Regier PS, Redish AD. Dorsolateral striatal task-initiation bursts represent past experiences more than future action plans. Journal of Neuroscience. 2021; 41(38):8051–8064.

De Leonibus E, Costantini VJA, Massaro A, Mandolesi G, Vanni V, Luvisetto S, Pavone F, Oliverio A, Mele A. Cognitive and neural determinants of response strategy in the dual-solution plus-maze task. Learning & Memory. 2011; 18(4):241–244.

Dickinson A, Balleine B. Motivational control of goal-directed action. Animal learning and behavior. 1994; 22(1):1–18.

Dickinson A, Balleine B, Watt A, Gonzalez F, Boakes RA. Motivational control after extended instrumental training. Animal Learning & Behavior. 1995; 23:197–206.

Featherstone RE, McDonald RJ. Dorsal striatum and stimulus-response learning: lesions of the dorsolateral, but not dorsomedial, striatum impair acquisition of a stimulus-response-based instrumental discrimination task, while sparing conditioned place preference learning. Neuroscience. 2004; 124(1):23–31.

Grammont F, Riehle A. Spike synchronisation and firing rate in a popula- tion of motor cortical neurons in relation to movement direction and reaction time. Biological Cybernetics. 2003; 88:360–373.

Graybiel AM. The basal ganglia and chunking of action repertoires. Neurobiology of learning and memory. 1998; 70(1-2):119–136.

Gremel CM, Costa RM. Orbitofrontal and striatal circuits dynamically encode the shift between goal-directed and habitual actions. Nature communications. 2013; 4(1):2264.

Hassabis D, Kumaran D, Vann SD, Maguire EA. atients with hippocampal amnesia cannot imagine experiences. Proc Natl Acad Sci USA. 2007; 104(5):1726–31.

Her ES, Huh N, Kim J, Jung MW. Neuronal activity in dorsomedial and dorsolateral striatum under the requirement for temporal credit assignment. Scientific Reports. 2016; 6(1):1–11.

Hilario M, Holloway T, Jin X, Costa RM. Different dorsal striatum circuits mediate action discrimination and action generalization. European Journal of Neuroscience. 2012; 35(7):1105–1114.

Hintiryan H, Foster NN, Bowman I, Bay M, Song MY, Gou L, Yamashita S, Bienkowski MS, Zingg B, Zhu M, et al. The mouse cortico-striatal projectome. Nature Neuroscience. 2016; 19(8):1100–1114.

Hooks BM, Papale AE, Paletzki RF, Feroze MW, Eastwood BS, Couey JJ, Winnubst J, Chandrashekar J, Gerfen CR. Topographic precision in sensory and motor corticostriatal projections varies across cell type and cortical area. Nature Communications. 2018; 9(1):3549.

Hunnicutt BJ, Jongbloets BC, Birdsong WT, Gertz KJ, Zhong H, Mao T. A comprehensive excitatory input map of the striatum reveals novel functional organization. eLife. 2016; 5:e19103.

James A, Reynaud-Bouret P, Mezzadri G, Sargolini F, Bethus I, Muzy A. Strategy inference during learning via cognitive activity-based credit assignment models. Scientific reports. 2023; .

Jin X, Costa RM. Start/stop signals emerge in nigrostriatal circuits during sequence learning. Nature. 2010; 466(7305):457–462.

Jog MS, Kubota Y, Connolly CI, Hillegaart V, Graybiel AM. Building neural representations of habits. Science. 1999; 286(5445):1745–1749.

Kimchi EY, Laubach M. The dorsomedial striatum reflects response bias during learning. Jounal of Neuroscience. 2009; 29(47):14891–902.

Lambert RC, Tuleau-Malot C, Bessaih T, Rivoirard V, Bouret Y, Leresche N, Reynaud-Bouret P. Reconstructing the functional connectivity of multiple spike trains using Hawkes models. Journal of neuroscience methods. 2018; 297:9–21.

Malvaez M, Greenfield VY, Matheos DP, Angelillis NA, Murphy MD, Kennedy PJ, Wood MA, Wassum KM. Habits are negatively regulated by histone deacetylase 3 in the dorsal striatum. Biological Psychiatry. 2018; 84(5):383–392.

Moussa R, Poucet B, Amalric M, Sargolini F. Contributions of dorsal striatal subregions to spatial alternation behavior. Learning & Memory. 2011; 18(7):444–451.

Okada K, Nishizawa K, Fukabori R, Kai N, Shiota A, Ueda M, Tsutsui Y, Sakata S, Matsushita N, Kobayashi K. Enhanced flexibility of place discrimination learning by targeting striatal cholinergic interneurons. Nature Communications. 2014; 5(1):3778.

Packard MG, Knowlton BJ. Learning and memory functions of the basal ganglia. Annual review of neuroscience. 2002; 25(1):563–593.

Peters AJ, Fabre JMJ, Steinmetz NA, Harris KD, Carandini M. Striatal activity topographically reflects cortical activity. Nature. 2021; 591(7850):420–425.

Ragozzino ME, Ragozzino KE, Mizumori SJY, Kesner RP. Role of the dorsomedial striatum in behavioral flexibility for response and visual cue discrimination learning. Behavioral Neuroscience. 2002; 116(1):105.

Redish AD. Vicarious trial and error. Nature Neuroscience. 2016; 17(3):147–59.

Regier PS, Amemiya S, Redish AD. Hippocampus and subregions of the dorsal striatum respond differently to a behavioral strategy change on a spatial navigation task. Journal of Neurophysiology. 2015; .

Reig R, Silberberg G. Multisensory integration in the mouse striatum. Neuron. 2014; 83(5):1200–1212.

Reynaud-Bouret P, Muzy A, Bethus I. Strategy inference during learning via cognitive activity-based credit assignment models. Comptes rendus de l'Académie des Sciences -Series I- Mathematics. 2021; 359(4):481–492.

Sales-Carbonell C, Taouali W, Khalki L, Pasquet MO, Petit LF, Moreau T, Rueda-Orozco PE, Robbe D. No discrete start/stop signals in the dorsal striatum of mice performing a learned action. Current Biology. 2018; 28(19):3044–3055.

705 **Smith KS**, Graybiel AM. A dual operator view of habitual behavior reflecting cortical and striatal dynamics.
706 Neuron. 2013; 79(2):361–374.

707 **Spiers HJ**, Gilbert SJ. Solving the detour problem in navigation: a model of prefrontal and hippocampal inter-
708 actions. Frontiers in human neuroscience. 2015; p. 125.

709 **Stalnaker TA**, Calhoon GC, Ogawa M, Roesch M, Schoenbaum G. Reward prediction error signaling in posterior
710 dorsomedial striatum is action specific. Jounal of Neuroscience. 2012; 32(30):10296–305.

711 **Stalnaker TA**, Calhoon GG, Ogawa M, Roesch MR, Schoenbaum G. Neural correlates of stimulus-response
712 and response-outcome associations in dorsolateral versus dorsomedial striatum. Frontiers in integrative
713 neuroscience. 2010; p. 12.

714 **Tang CC**, Root DH, Duke DC, Zhu Y, Teixeiria K, Sisi M, Barker DJ, West MO. Decreased Firing of Striatal Neu-
715 rons Related to Licking during Acquisition and Overtraining of a Licking Task. Jounal of Neuroscience. 2009;
716 29(44):13952–961.

717 **Thorn CA**, Atallah H, Howe M, Graybiel AM. Differential dynamics of activity changes in dorsolateral and dor-
718 somedial striatal loops during learning. Neuron. 2010; 66(5):781–795.

719 **Turner KM**, Svegborn A, Langguth M, McKenzie C, Robbins TW. Opposing roles of the dorsolateral and dor-
720 somedial striatum in the acquisition of skilled action sequencing in rats. Journal of Neuroscience. 2022;
721 42(10):2039–2051.

722 **Vandaele Y**, Mahajan NR, Ottenheimer DJ, Richard JM, Mysore SP, Janak PH. Distinct recruitment of dorsomedial
723 and dorsolateral striatum erodes with extended training. eLife. 2019; 8:e49536.

724 **Yin HH**, Knowlton BJ, Balleine BW. Lesions of dorsolateral striatum preserve outcome expectancy but disrupt
725 habit formation in instrumental learning. European journal of neuroscience. 2004; 19(1):181–189.

726 **Yin HH**, Ostlund SB, Knowlton BJ, Balleine BW. The role of the dorsomedial striatum in instrumental condition-
727 ing. European Journal of Neuroscience. 2005; 22(2):513–523.

728 **Yin HH**, Knowlton BJ, Balleine BW. Inactivation of dorsolateral striatum enhances sensitivity to changes in the
729 action–outcome contingency in instrumental conditioning. Behavioural brain research. 2006; 166(2):189–
730 196.

731 **Yin HH**, Mulcare SP, Hilário MRF, Clouse E, Holloway T, Davis MI, Hansson AC, Lovinger DM, Costa RM. Dynamic
732 reorganization of striatal circuits during the acquisition and consolidation of a skill. Nature Neuroscience.
733 2009; 12(3):333–341.

A

Figure 6. Reproduction of tetrode positions and recording sites in the DMS and DLS. **A**: Photos of Nisl-stained coronal sections from two representative rats implanted in the DLS (left panel) and DMS (right panel). The tetrode tracks are visible in both sections. The vertical black bars represent the lengths of striatal tissu from which neural recordings were performed. **B**: Representation of the recording sites from all rats (DLS in blue and DMS in red)

A   **B**

Figure 7. Timeline of a typical training session. **A**: The maze has been divided in boxes. The black line represents the trajectory of the animal during one example session. The feeders are located in the boxes 1 and 2. Maze intersections are located in boxes A and B. **B**: Each 500ms task event (centered on the onsets, intersections and reward locations) was divided in five 100ms bins. This leads to a total of 6 * 5 = 30 time bins.

A

B

Figure 8. FSI-MSN classification. **A**: Dendogram of the hierarchical clustering (method "Ward D2") of neurons based on waveform (wf) properties (Peak-Valley distance (PV), Width at mid-height (W)) and firing rate. Two clusters are identified using K-means method (with $K = 2$). **B**: 3D representation of wf properties and firing rate of the two clusters. Fast Spiking Interneurons (FSI) and Medium Spiny Neurons (MSN) are represented in orange and green, respectively. A total of 451 MSN and 199 FSI have been identified. No significant difference in the proportion of FSI/MSN was observed between DMS and DLS (p-value = 0.97).