A cline within an ecotype of the yellow monkeyflower, *Mimulus guttatus*

Thomas Zambiasi[1,2,3*], David B. Lowry[1,4,5]

[1]Department of Plant Biology, Michigan State University, East Lansing, MI, USA

[2]Department of Biology, Indiana University, Bloomington, IN, USA

[3]Evolution, Ecology, and Behavior Program, Indiana University, Bloomington, IN, USA

[4]Evolution, Ecology, and Behavior Program, Michigan State University, East Lansing, MI, USA

[5]Plant Resilience Institute, Michigan State University, East Lansing, MI, USA

*For correspondence:

Thomas Zambiasi tzambias@iu.edu

Word count: 4963

Tables: 1

Figures: 1

Color figures: 2

Supplementary information: 3 Tables

Manuscript received _______; revision accepted _______.

RUNNING HEAD: A cline within an ecotype

ABSTRACT

Premise: A key goal of evolutionary biologists is to understand how and why adaptive genetic

variation is partitioned within species. In the yellow monkeyflower, *Mimulus guttatus* (syn.

Erythranthe guttata), coastal perennial populations collectively constitute a single genetically

and morphologically differentiated ecotype. While the distinctiveness of the coastal ecotype has

now been well documented, there is also variation in environmental factors across the range of

the coastal ecotype that could drive differentiation among its component populations in a more

continuous way.

Methods: Based on previous observations of a potential cline within this ecotype, we quantified

plant height across coastal perennial accessions from 69 total populations in two greenhouse

common garden experiments. To evaluate possible environmental factors driving the relationship

between plant height and latitude, we regressed height against multiple climatic factors,

including temperature, precipitation, and coastal wind speeds.

Results: In both experiments, plant height was negatively correlated with latitude. *Mimulus*

height correlated positively with annual precipitation and with mean wind speed, to a lesser

degree, and negatively with annual mean temperature. We hypothesize that one or more of these

factors drove clinal variation within the coastal ecotype.

Conclusion: Overall, our study illustrates the complexity of how the distribution of

environmental variation can both simultaneously drive the evolution of distinct ecotypes as well

as continuous clines within those ecotypes. These results are discussed in the context of the

classic criticisms of ecotypes being intermediates in the process of speciation.

KEY WORDS: Local adaptation, salt spray, *Erythranthe guttata*, wind, speciation

INTRODUCTION

A century ago, Turesson (1922a,b) recognized that adaptive phenotypic variation was often

highly divergent in correlated sets of traits when comparing groups of populations derived from

different habitats. Turesson referred to these groups of phenotypically similar, locally-adapted

populations as ecotypes (Turesson 1922a,b; Gregor et al. 1936). Several researchers following

Turesson found the concept of ecotypes useful for describing the distribution of adaptive genetic

variation within species (Gregor et al. 1936; Clausen et al. 1948; Clausen 1951; Böcher 1967;

Heywood 2011). Further, these researchers viewed ecotypes as an intermediate stage in the

process of speciation and a way to directly link natural selection to the formation of new species

(reviewed in Lowry 2012). However, other researchers doubted the validity of Turesson's

ecotype concept and argued that clines (Huxley 1938) best represented how adaptive genetic

variation is partitioned across the geographic distribution of a species' range (Langlet 1963;

1971; Stebbins 1980).

While some recent authors have concluded that the cline versus ecotype controversy has been

resolved because adaptive genetic variation can be partitioned different ways depending on the

distribution of selective factors across the range of a species (Linhart and Grant 1996; Briggs and

Walters 2016), multiple questions still remain. Foremost among these questions is whether

ecotypes do indeed represent intermediate stages in the formation of new species. Both Quinn

(1978) and Levin (1993) argued that there was no way that ecotypes could be intermediate stages

in the process of speciation because, in their view, formation of geographically widespread

ecotypes would require uniform habitats to also be widespread across large geographic regions.

This led Levin to conclude that most speciation in plants occurs in localized, isolated populations

(Levin 1993). However, if it can be shown that ecotypes can evolve despite a lack of uniformity

of habitat conditions, it would support the hypothesis that widespread ecotypes are indeed a stage

in the process of the formation of new species.

In this study, we examined whether there is a clinal distribution of phenotypic variation within a

widespread ecotype of the yellow monkeyflower, *Mimulus guttatus* (syn. *Erythrante guttata*;

Barker et al. 2012). *M. guttatus* exhibits high levels of genetic and phenotypic variation across its

extensive range across western North America, which makes it a particularly useful system for

studying local adaptation (Vickery 1978; Lowry et al. 2008). Coastal perennial populations of *M.*

guttatus, occurring from southern California to southern Washington State, collectively comprise

a distinct ecotype based on population genetic data (Lowry et al. 2008; Twyford and Friedman

2015). This ecotype has been designated as a variety (*M. guttatus var. grandis* Greene) or species

(*M. grandis* syn. *E. grandis*; Barker et al. 2012, Nesom 2012, 2014) by taxonomists. Despite

these taxonomic designations, due to its morphological distinctiveness, the coastal perennial

ecotype is highly interfertile with other populations of *M. guttatus* (Lowry and Willis 2010).

Further, there were only four fixed genetic differences between the coastal perennial and inland

annual populations out of 29,693,578 SNPs surveyed in a population genetic study of this system

(Gould et al. 2017). Based on these results, we believe that it is best to think of the coastal

perennial populations as intraspecific genetic variation within the *Mimulus guttatus* species

complex (Lowry and Willis 2010; Twyford and Friedman 2015; Gould et al. 2017).

Several previous studies of local adaptation in *M. guttatus* have found striking genetically-based

phenotypic differences between the coastal perennial ecotype and nearby inland populations in

the western United States (Vickery 1952; Hall et al. 2006, 2010; Lowry et al. 2008; Oneal et al.

2014). Many of the inland populations of *M. guttatus* have an annual life-history, especially

those that occur in inland habitats that completely dry out during summer months (Vickery 1952;

Clausen and Heisey 1958; Wu et al. 2008; Lowry et al. 2008). In contrast, the coastal perennial

ecotype occurs in coastal habitats that are regularly damp and foggy during summer months

(Vickery 1952; Clausen and Heisey 1958; Hitchcock and Cronquist 1973; Hall and Willis 2006;

Lowry et al. 2008). In greenhouse common garden experiments, accessions derived from coastal

habitats had thicker stems, many more prostrate stoloniferous branches, later flowering times,

and produced far more herbivore defensive compounds than nearby inland annual populations

(Hall et al. 2006, 2010; Lowry et al. 2008; Lowry and Willis 2010; Oneal et al. 2014; Lowry et

al. 2019).

While there is both phenotypic and genetic divergence between ecotypes of *M. guttatus*, it is

unclear to what extent genetic variation is present within individual ecotypes – either due to

adaptation to widespread environmental gradients or idiosyncratic with regard to geographic

space. However, a previous pilot study suggested the possibility that there may be a cline in plant

height within the coastal ecotype (Lowry 2010). Since that initial study, dozens of additional

coastal *M. guttatus* populations have been collected, which provides an opportunity for a more

comprehensive survey of variation over the entire geographic range of the coastal perennial

ecotype.

In this study, we tested the hypothesis that clinal phenotypic variation can occur within ecotypes

of a species. Specifically, we investigated several populations of coastal perennial *M. guttatus*

from across the ecotype's range to determine whether there was indeed a latitudinal cline in plant

height. To establish potential sources of selection, we examined whether there is a relationship

between the phenotypes of the accessions and the climates from where they were collected in

nature. Through these efforts, we attempted to understand whether clinal trait variation occurred

within a distinct ecotype, which would support the hypothesis that the evolution of

geographically widespread ecotypes does not require a widespread distribution of uniform

habitats, as argued by Quinn (1978) and Levin (1993).

METHODS

Germplasm for experiments –

The seeds used in these experiments originated from natural population collections made

between 2004 and 2016. Seeds were either collected directly from plants in the field or collected

from plants that had been brought from the field back to the greenhouses at Duke University

(Durham, North Carolina, USA) or Michigan State University (East Lansing, Michigan, USA).

For each population, we collected location description information, habitat description, and GPS

coordinates. Photos of the plants in the field for each population have been posted on iNaturalist

and vouchers for a subset of the populations were deposited at the California Academy of

Sciences. Permits were obtained for collections on public lands, where permission was required.

All maternal families used in this study are currently stored at Michigan State University in

airtight containers with silica gel as a desiccant.

Greenhouse Experiments –

To determine whether there was clinal variation in *M. guttatus* height, we conducted two

independent experiments in two different controlled greenhouse environments.

For the first experiment, seeds collected from coastal perennial *M. guttatus* were planted

according to maternal family ($N = 135$ seed families, 56 populations) in 3.5 sq. in. pots with

SUREMIX (Michigan Grower Products, Inc., Galesburg, Michigan, USA). We randomized these

pots into flats (17 pots per flat) and stratified the seeds in a dark cold room (set to 4º C) for a

week to encourage more uniform germination before being transferred to the greenhouses at

Michigan State University. All germinating seedlings were removed from each pot, except for

the centermost plant. Plants were bottom-watered approximately every other day throughout the

remainder of the experiment. Plants were grown in the greenhouses with 16 h of supplemental

lighting and room temperature set to 22º C. We measured plant height in centimeters for each

plant at the time of first flower. Approximate germination dates and flowering dates were also

recorded.

Based on promising results from the first experiment, we conducted a second greenhouse

experiment to determine if the initial results would be replicable with a larger sample size (more

populations and individuals) and in a different greenhouse. Seeds from coastal perennial *M.*

guttatus ($N = 402$ seed families, 66 populations) were planted (SUREMIX, 3.5 sq. in. pots)

without prior cold stratification, directly in Plant Ecology Field Lab greenhouse at Kellogg

Biological Station (KBS, Hickory Corners, MI, USA). Each pot contained seeds from a single

maternal family and were randomized into flats in the same way as the first experiment. Again,

pots were thinned to one plant and remaining individuals were watered every other day

throughout the experiment. Germination and flowering dates were recorded, and plant height was

measured (in cm) at the time of first flowering.

Climate Data –

To investigate potential environmental drivers of adaptive phenotypic variation, we extracted

climatic data for each population's collection location of origin from the WorldClim Bioclimatic

variables database (Fick and Hijmans 2017). This data was imported and compiled using R

(v.3.6.0, R Core Team 2019).

In addition to the Bioclim data, we extracted oceanic wind speed data from along the US Pacific

Coast using the National Renewable Energy Laboratory's Wind Prospector website, which

featured interactive online maps of wind speeds surveyed by NREL (King et al. 2014; Draxl et

al. 2015a,b). We used Pacific Monthly Offshore Wind Speed map layers to collect wind speeds

for each month from the site nearest to shore (8 km offshore at the closest; Draxl et al. 2015b) at

the latitudes of each of the coastal perennial populations. The wind speeds found through the

NREL Wind Prospector were the averages for each 1.2 km X 1.2 km offshore site and were

modeled at 100 m (Draxl et al. 2017). Due to unavailability of data outside of the contiguous US,

populations found in British Columbia and Alaska (denoted by population codes ALA and LBG)

were not included in climate data analyses. Monthly wind speed averages accessible via NREL

were based on observations and statistical analyses over the course of seven years (King et al.

2014; Draxl et al. 2015a,b), which we used to calculate quarterly and annual wind speeds for

each population collection location. We then matched these climate variables with their

corresponding coordinates and population codes for further analysis.

Statistical Analysis –

To test whether there was a relationship between plant height and the latitude of origin (location

from which each population accession was collected), we conducted linear regression modeling

using the R software package (v.3.6.0, R Core Team 2019). We conducted independent analyses

for each of the two greenhouse experiments.

Although we predicted that latitude would correlate with plant height across space, we wanted to

evaluate which climatic variables correlated with latitude might ultimately be driving observed

clinal patterns. Climate variables are often highly correlated with each other, so we conducted a

principal components analysis (PCA) to reduce the dimensionality of the data prior to further

analyses. Climate data, described above, was extracted for each population and we performed the

PCA in R. For both sets of greenhouse data, we utilized all principal components (PCs) with an

eigenvalue greater than 1, following the Kaiser criterion (Kaiser 1960). We then conducted

regression analyses of plant height against each of these PCs, again using linear models in R. All

data organization, analysis, and visualization were conducted in R Studio.

RESULTS

Plant height was negatively correlated with latitude –

Overall, we found a significant relationship between population latitude and plant height in both

Greenhouse Experiment 1 (Multiple $R^2 = 0.24$, $F_{1,54} = 16.71$, $P < 0.001$, $N = 56$ populations; 135

individuals) and Greenhouse Experiment 2 ($R^2 = 0.37$, $F_{1,64} = 36.83$, $P < 0.001$, $N = 66$

populations; 402 individuals). In both experiments, coastal perennial *M. guttatus* height was

negatively correlated with latitude (Figures 1, 2). Based on our linear modeling, we found that

every degree latitude further north translated to roughly 1 cm decrease in plant height under

greenhouse conditions (Experiment 1 slope = -0.95; Experiment 2 slope = -1.12).

202   ***The relationship of climatic factors and plant height for Experiment 1 –***

To gain further insight into the possible causes of the negative relationship between latitude and

plant height, we conducted principal component analyses of climate factors. For Greenhouse

Experiment 1, we focused our analyses on climate PC1 through PC4, as all of them had

eigenvalues greater than 1. These 4 PCs accounted for 94.55% of cumulative variance in the

climate factors; PC1 explained 59.03% of the variation, followed by PC2 at 19.57%, PC3 at

11.13%, and PC4 at 4.82%. We found that PC1 had a significant negative relationship with plant

height (*Multiple R²* = 0.22, $F_{1,54} = 14.97$, $P < 0.001$, slope = -0.72). The other three PCs were not

significantly correlated with plant height ($P > 0.05$).

Given that PC1 had a significant negative correlation with height, we investigated which climate

variables were most highly loaded on this PC axis. The most heavily loaded variables on PC1

were Annual Precipitation, Average Annual Temperature, Precipitation of the Wettest Quarter,

and Precipitation of the Coldest Quarter. Annual Temperature was negatively correlated with

PC1, while all others were positively correlated. While not as heavily loaded, Average Wind

Speeds in Quarter 1 and Quarter 4 were both positively correlated with PC1. Additional variables

that were significantly loaded on PC1 included Precipitation of the Wettest Month, Precipitation

Seasonality, Precipitation of the Warmest Quarter, and Average Temperature of the Coldest

Quarter. More specific information regarding loadings and coordinates of individual climate

variables on PC1 can be found in Table 1.

222   ***The relationship of climatic factors and plant height for Experiment 2 –***

For Greenhouse Experiment 2, we found that PC1 through PC6 all had eigenvalues greater than

1. In total, those 6 PCs accounted for 98.33% of cumulative variance in climate factors. PC1

explained the most variation at 55.93%, followed by PC2 with 22.94%, PC3 with 10.88%, PC4

with 3.87%, PC5 with 2.65%, and PC6 with 2.07%. PC1 was significantly negatively correlated

with plant height (*Multiple R^2* = 0.39, $F_{1,62}$ = 39.53, $P < 0.001$, slope = -1.25). As in Experiment

1, none of the other 5 PCs were significantly correlated with plant height ($P > 0.05$).

As we did for Experiment 1, we evaluated which climatic factors loaded onto PC1. Heavily

loaded climate variables included Annual Precipitation and Average Annual Temperature, in

addition to Precipitation Seasonality, Precipitation of the Warmest Quarter, and Precipitation of

the Wettest Quarter. Of these variables, Average Annual Temperature and Precipitation

Seasonality were negatively correlated with PC1, while the others were positively correlated

(Table 2). The Average Wind Speed in Quarters 1 and 4 were also loaded on PC1, but not as

strongly as the other factors.

237   ***Latitude strongly predicts climate for coastal populations –***

While latitude can be a weak predictor of climate, especially for continental populations, we

suspected that it might be highly correlated with climate in our study, as all populations were

distributed latitudinally along a coastline. To assess how well latitude predicts climate for these

populations, we regressed the extracted principal components from both greenhouse experiments

against latitude. These analyses confirmed that many of the climate variables involved were

highly correlated with latitude. Using the principal components from Experiment 1, we found

that PC1 had a very strong positive relationship with latitude (*Multiple R²* = 0.93, $F_{1,54}$ = 689, *P*

< 0.001, slope = 0.76). The other PCs were not significantly correlated with latitude (*P* > 0.05).

Just as for Experiment 1, PC1 of Experiment 2 was the only PC significantly associated with

latitude (*Multiple R²* = 0.91, $F_{1,62}$ = 652.1, *P* < 0.001, slope = 0.85). To determine how well

variation in individual climatic factors was predicted by latitude, we regressed all the climate

factors that were highly loaded on PC1 individually against latitude. Those results are reported in

Table 1.

252   *Patterns of wind speed along the west coast of North America –*

Plants are well known to respond to wind through changes in growth form (Grace 1977;

Gardiner et al. 2016; Telewski 2021). This, combined with the observation of wind speed

loading onto the first PC of climate, led us to evaluate the patterns of wind speed variation more

carefully across the populations that we studied. We found that, in general, annual wind speeds

increase with latitude (*Multiple R²* = 0.06, $F_{1,76}$ = 4.465, *P* = 0.038, slope = 0.07). Average wind

speeds in Quarter 1 (*Multiple R²* = 0.46, $F_{1,76}$ = 65.9, *P* < 0.001, slope = 0.19) and Quarter 4

(*Multiple R²* = 0.50, $F_{1,76}$ = 75.32, *P* < 0.001, slope = 0.27) of the year follow this pattern closely,

although Quarter 2 wind speeds (*Multiple R²* = 0.20, $F_{1,76}$ = 19.1, *P* < 0.001, slope = -0.20)

showed a negative relationship with latitude. Quarter 3 wind speeds were not correlated with

latitude (*Multiple R²* = 0.01, $F_{1,76}$ = 0.57, *P* > 0.05, slope = 0.05). A key realization from this

analysis is that patterns of wind speed variation shift dramatically across seasons and have a

striking non-linear relationship with latitude. This nonlinear relationship can be observed in

visualizations of annual and quarterly average wind speeds: after initially increasing with

latitude, average wind speeds generally decrease with latitude for populations north of southern

Oregon (42º N) (Figure 3). Across seasons, wind speeds slow down during the summer much

more in the southern end of the range of the coastal populations, while remaining high in

northern California and southern Oregon throughout the summer months.

DISCUSSION

In this study, we found a significant negative relationship between latitude and genetically-based

differences in plant height within the coastal ecotype of *M. guttatus*. This clinal pattern was

likely driven by adaptation to climate variables that are correlated with latitude along the coast of

the Pacific Ocean. Populations of *M. guttatus* collected from the southern end of the range were

repeatedly found to be taller in common garden experiments than more northern populations.

These southern populations generally experience higher temperatures, less precipitation, and

lower velocity winds than more northern populations, especially during summer months. The

results of this study further reinforce the idea that adaptive genetic variation can be

simultaneously partitioned in disjunct (among ecotypes) and continuous (clinal) ways. Overall,

our results have implications for the evolution of adaptive genetic variation in this system, as

well as general implications for the longstanding criticism of the hypothesis that ecotypes are

intermediates in the evolution of plant species.

The finding of clines within ecotypes helps shape our understanding of the evolution of

new plant species –

Understanding the patterns by which genetic variation is partitioned among populations within a

species across a geographic landscape has been a matter of considerable debate, especially in the

classic plant evolutionary biological literature (reviewed in Lowry 2012). It is reasonable to

hypothesize that the partitioning of that genetic variation is a function of how environmental

variation is partitioned across the same landscape. However, several previous researchers have

argued against the existence of ecotypes and that only clinal variation exists within species

(Langlet 1971; Quinn 1978; Stebbins 1980; Levin 1993).

A series of prior studies has clearly established that the coastal populations of *Mimulus guttatus*

from central California to southern Washington state collectively comprise a genetically distinct

geographically widespread ecotype (Lowry et al. 2008; Lowry and Willis 2010; Oneal et al.

2014; Twyford and Friedman 2015). This coastal perennial ecotype clusters as a distinct group

from the rest of the *Mimulus guttatus* species complex based on both a set of morphological traits

and allele frequencies of loci across the genome (Lowry 2012; Oneal et al. 2014; Twyford

andFriedman 2015) but has not accumulated appreciable divergence through fixed genetic

differences (Gould et al. 2017).

While genomic data has allowed us to address the classic criticism about the existence of

geographically widespread ecotypes made by Langlet (1971) and others, it is the discovery of

clines within ecotypes that allows us to address doubts about how those ecotypes could come

into existence in the first place. Both Quinn (1978) and Levin (1993) expressed doubt about the

evolution of geographically widespread ecotypes evolving within species because they

hypothesized that uniform habitats would also necessarily be widespread to facilitate the

concerted evolution of new widespread ecotypes. Levin (1993) took this argument a step further

and argued that most speciation in plants must be local and thus, follow a model of peripatric

speciation. Levin's argument was built off those of Raven (1980), who concluded that there was

no logical way that geographically widespread ecotypes or subspecies could eventually evolve

into distinct species.

The level of environmental change across the range of the coastal ecotype of *M. guttatus*

demonstrates that habitats with uniform environments are not necessary for the evolution of

geographically widespread ecotypes. Further, the finding of a cline within the coastal ecotype

suggests that that nonuniformity of climate across the range of an ecotype can drive adaptive

phenotypic variation within an ecotype, while other shared climate factors simultaneously

maintain the divergence of that ecotype from the rest of a species complex. Indeed, this should

not be a surprising conclusion, as clines within coastal ecotypes/species have been observed in

the classic literature for plant systems such as *Plantago maritima* (Gregor 1930, 1938) and

Achillea borealis (Clausen et al. 1948).

Overall, the conclusion that widespread ecotypes exist as intermediates in the evolution of new

species fits the emerging modern paradigm of speciation based on recent results from genomic

studies. In those studies, there is increasing evidence that introgression (i.e., gene flow) occurs at

various stages during the process of species splitting into distinct phylogenetic lineages

(Reviewed in Mallet et al. 2015; Taylor and Larson 2019; Edelman and Mallet 2021). This

model of speciation with gene flow is not the same as classic models of sympatric speciation,

which require consistent gene flow throughout the entire process of speciation (Coyne andOrr

2004). Instead, recent genomic studies suggest that it takes considerable time, with successive

bouts of sympatry and allopatry for distinct, fully reproductively isolated phylogenetic lineages

to become established (Mallet et al. 2015; Taylor and Lason 2019; Edelman and Malet 2021).

Further, a recent review by Levin (2012) supports the long period over which the speciation

process is completed in plants. During the long period over which new lineages evolve,

ecotypes/subspecies will evolve clines and locally adapted populations within their boundaries.

Many of these ecotypes/subspecies will go extinct or merge fully back into broader species

complexes. Others will eventually become fully reproductively isolated lineages, but only after

multiple bouts of gene flow with other members of the species complex. In the case of the

coastal perennial ecotype of *M. guttatus*, the population genetic evidence suggests that it was

derived at a single point in time from the rest of the species complex and then spread along the

Pacific coast, adapting to local environments as it spread (Lowry et al. 2008; Twyford &

Frianddman 2015). During and following the spread of the coastal ecotype, there has likely been

appreciable gene flow with other parts of the species complex, which is further supported by

there only being four fixed nucleotide differences between these groups out of millions of SNPs

(Gould et al. 2017).

The environmental factors responsible for the cline in plant height *–*

While we have demonstrated that a cline in plant height does occur within the coastal ecotype of

Mimulus guttatus, the question remains as to what environmental factors have driven this cline.

Though our analysis identified multiple factors correlated with both latitude and plant height,

these factors must be carefully considered, as they are autocorrelated with each other and not all

of them may have impact on the selective advantage of plant height.

Other researchers studying the evolution of ecotypes in the same region have found similar

patterns in plant height variation along the Pacific coast of North America. These studies were

far less comprehensive, but are useful, as they suggest that the patterns observed within the

coastal ecotype of *M. guttatus* are not unique. For example, Clausen et al. (1948) found similar

genetic variation with the coastal ecotype of *Achillea borealis*, when comparing populations in

Northern California, where populations north of the San Francisco Bay were shorter in stature

than those from farther south in California in common garden experiments.

While temperature and precipitation were most highly correlated with latitude and thus, plant

height in our study, it is difficult to conceptualize a mechanistic hypothesis of how those factors

would drive the evolution of the cline in plant height. In contrast, the spatial-temporal

distribution of winds speeds along the Pacific coast, while less closely correlated with latitude,

due to its non-linear distribution, is more likely to have driven the cline in plant height. Onshore

wind from the ocean can be intense along the Pacific Coast, depending on the location and time

of year (Grace 1977; Gilliam 2002). The intensity of wind speed increases with the distance

above the ground (Grace 1977; Jaffe and Forbes 1993). While wind can be damaging by itself, it

is particularly damaging to plants along ocean coastlines because it contains aerosolized oceanic

salt spray (Boyce 1954; Barbour 1978; Cheplick and Demetri 1999; Griffiths et al. 2006; Kouali

et al. 2017; Du and Hesp 2020). Unlike stress caused by soil salinity, salt spray primarily

damages plants by landing directly on aboveground tissues and entering through stomata and

small cracks in the cuticle (Grieve andPitman 1978; Du and Hesp 2020). Boyce (1954)

recognized that this compact growth form is advantageous for coastal plants because it helps

them to avoid the impacts of salt spray by being more sheltered from the wind. Prior research has

demonstrated that the coastal perennial ecotype of *M. guttatus* is consistently more tolerant to

salt spray in laboratory assays than the inland annual ecotype (Lowry et al. 2009). However,

these assays could not assess how the combination of wind speeds and salt spray could drive the

evolution of plant architecture.

We hypothesize that the evolution of the cline in height of the coastal perennial ecotype is a

function of the interaction of time and space of wind speeds along the Pacific coast. Across the

range of the ecotype, wind speeds and wave energy that drive salt spray are greater in the spring

(April, May) than the summer (July, August) (Gilliam 2002; Garcia-Reyes and Largier 2012).

However, those winds speeds dissipate most during the summer in the region south of the San

Francisco, especially in more protected areas, like the Monterey Bay (Figure 2b), where plants in

the field reach up to two meters in height (Lowry *personal observation*). In Northern California

and Southern Oregon, high coastal windspeeds remain elevated throughout the summer months,

which coincides with when this species flowers. Since reproductive structures are often the most

sensitive to salt spray (Griffiths et al. 2006), maintaining a compact growth form when flowering

under higher windspeed/salt spray conditions is likely to be advantageous. While this hypothesis

works well to explain the results for populations from central California to southern Oregon, it does

not hold up for central and Northern Oregon, where windspeeds are generally reduced during

summer months. This suggests that other environmental factors are responsible for maintaining a

compact stature in this region or that these populations are shorter in stature due to evolutionary

history. Evolutionary history could be responsible for the short stature of the northernmost

populations, if they are descended from populations from windier regions to the south. Alternatively,

the northern region of Oregon could have had higher wind speeds in the recent past, which drove

their present-day short stature. Future population genomic studies could establish whether the

observed geographic patterns could be a result of evolutionary history. Further, tests to establish

whether reduced height is an adaptation to wind and salt spray could be conducted by growing

populations of various heights in a common garden experiment at a windy coastal location.

ACKNOWLEDGEMENTS

This work is dedicated to the memory of Don Levin, whose papers and conversations challenged

us to think hard about the nature of ecotypes and how we should expect variation to be

partitioned among groups during plant speciation. We would like to thank Mauricio

Swartzendruber, Marty Schmidt, Simran Patel, Thomas Young, and Will Burke (Michigan State

University) for assisting with plant care and data collection. Nate Emery, Anne-Sophie Bohrer,

Linnea Fraser, and Milagros del Pilar Jimenez-Hernandez provided advice with analysis using R.

Caroline Draxl of NREL also shared her insight regarding the collection of offshore wind data.

This work was funded by National Science Foundation Division of Integrative Organismal

Systems Grants to DBL (IOS-1855927 and IOS-2153100).

AUTHOR CONTRIBUTIONS

DBL conceived of and oversaw the experiments and analyses. TZ conducted the experiments and analyses. Both authors wrote the manuscript.

DATA AVAILABILITY STATEMENT

All data will be made available at publication, either as supplementary information or on Dryad.

LITERATURE CITED

Barbour, M. G. 1978. Salt spray as a microenvironmental factor in the distribution of beach plants at Point Reyes, California. *Oecologia* 32: 213-224.

Barker, W. R., G. L. Nesom, P. M. Beardsle, and N. S. Fraga. 2012 A taxonomic conspectus of Phrymaceae: a narrowed circumscription for Mimulus, new and resurrected genera, and new names and combinations. *Phytoneuron* 39: 1-60.

Böcher, T. W. 1967. Continuous variation and taxonomy. *Taxon* 16: 255-258.

Boyce, S. G. 1954. The salt spray community. *Ecological Monographs* 24: 29– 67.

Briggs, D., and S. M. Walters. 2016. Plant variation and evolution. Cambridge University Press, Cambridge, UK.

Cheplick, G. P., and H. Demetri. 1999. Impact of saltwater spray and sand deposition on the coastal annual *Triplasis purpurea* (Poaceae). *American Journal of Botany* 86: 703–710.

Clausen, J. 1951. Stages in the evolution of plant species. Cornell University Press, Ithaca, NY,

USA.

Clausen, J., D. D. Keck, and W. M. Hiesey. 1948. Experimental studies on the nature of species.

III. Environmental responses of climatic races of *Achillea*. Carnegie Institution of

Washington, Washington, DC, USA.

Clausen, J., and W. M. Heisey. 1958. Experimental studies on the nature of species. IV. Genetic

structure of ecological races. Carnegie Institution of Washington, Washington, DC, USA.

Coyne, J. A., and H. A. Orr. 2004. Speciation. Sinauer Associates, Inc, Sunderland, MA, USA.

Draxl, C., B. M. Hodge, A. Clifton, and J. McCaa. 2015a. Overview and Meteorological

Validation of the Wind Integration National Dataset Toolkit (Technical Report,

NREL/TP-5000-61740). National Renewable Energy Laboratory, Golden, CO, USA.

Draxl, C., B. M. Hodge, A. Clifton, and J. McCaa. 2015b. The Wind Integration National

Dataset (WIND) Toolkit. *Applied Energy* 151: 355-366.

Draxl, C., C. Philips, G. Scott, and W. Musial. 2017. "Characterization of Offshore Wind

Resources in the Contiguous United States." Presented at the 2017 NAWEA Conference,

September 29, Ames, Iowa, USA. Retrieved July 18, 2022.

Du, J., and P. A. Hesp. 2020. Salt spray distribution and its impact on vegetation zonation on

coastal dunes: a review. *Estuaries and Coasts* 43: 1885-1907.

Edelman, N. B., and J. Mallet. 2021. Prevalence and adaptive impact of introgression. *Annual*

Review of Genetics 55: 265-283.

Fick, S. E., and R. J. Hijmans. 2017. WorldClim 2: new 1km spatial resolution climate surfaces

for global land areas. *International Journal of Climatology* 37: 4302-4315.

García-Reyes, M., and J. L. Largier. 2012. Seasonality of coastal upwelling off central and

northern California: New insights, including temporal and spatial variability. *Journal of*

Geophysical Research 117: C03028.

Gardiner, B., P. Berry, and B. Moulia. 2016. Review: Wind impacts on plant growth, mechanics

and damage. *Plant Science* 245: 94-118.

Gilliam, H. 2002. Weather of the San Francisco Bay Region. University of California Press,

Berkeley, CA, USA.

Gould, B. A., Y. Chen, and D. B. Lowry. 2017. Pooled ecotype sequencing reveals candidate

genetic mechanisms for adaptive differentiation and reproductive isolation. *Molecular*

Ecology 26: 163-177.

Grace, J. 1977. Plant Response to Wind. Academic Press, New York, NY, USA.

Gregor, J. W. 1930. Experiments on the genetics of wild populations. I. *Plantago maritima*.

Journal of Genetics 22: 15-25.

Gregor, J. W. 1938. Experimental taxonomy II: Initial population differentiation in *Plantago*

maritima L. of Britain. *New Phytologist* 37: 15-49.

Gregor, J. W., V. McM. Davey, and J. M. S. Lang. 1936. Experimental taxonomy I.

Experimental garden technique in relation to the recognition of the small taxonomic

units. *New Phytologist* 35: 323–350.

Grieve, A. M., and M. G. Pitman. 1978. Salinity damage to Norfolk Island pines caused by

surfactants. III. Evidence for stomatal penetration as the pathway of salt entry to leaves.

Functional Plant Biology 5: 397-413.

Griffiths, M. E., R. P. Keith, and C. M. Orians. 2006. Direct and indirect effects of salt spray and

fire on coastal heathland plant physiology and community composition. *Rhodora* 108:

32–42.

Hall, M. C., C. J. Basten, and J. H. Willis. 2006. Pleiotropic quantitative trait loci contribute to

population divergence in traits associated with life-history variation in *Mimulus guttatus*.

Genetics 172: 1829-1844.

Hall, M. C., D. B. Lowry, and J. H. Willis. 2010. Is local adaptation in *Mimulus guttatus* caused

by trade-offs at individual loci? *Molecular Ecology* 19: 2739–2753.

Heywood, V. H. 2011. The genesis of IOPB: a personal memoir. *Taxon* 60: 320-323.

Hitchcock, C. L., and A. Cronquist. 1973. Flora of the Pacific Northwest. University of

Washington Press, Seattle, WA, USA.

Huxley, J. 1938. Clines: an auxiliary taxonomic principle. *Nature* 142: 219-220.

Jaffe, M. J., and S. Forbes. 1993 Thigmomorphogenesis: the effect of mechanical perturbation on

plants. *Plant Growth Regulation* 12: 313-324.

Kaiser, H. F. 1960. The application of electronic computers to factor analysis. *Educational and*

Psychological Measurement 20:141-151.

King, J., A. Clifton, and B. M. Hodge. 2014. Validation of Power Output for the WIND Toolkit
(Technical Report, NREL/TP-5D00-61714). National Renewable Energy Laboratory,
Golden, CO, USA.

Kouali, A., M. Mouradi, L. Latrach, A. Khadraji, O. Elasri, Z. Baicha, A. Berrichi, N. Kouddane,
and A. Boukroute. 2017. Study of effects of seawater salt spray on growth, chlorophyll
fluorescence and chlorophyll content in three coastal species of Morocco. *Journal of*
Materials and Environmental Sciences 8: 4385–4390.

Langlet, O. 1963. Patterns and terms of intraspecific ecological variability. *Nature* 200: 347–348.

Langlet, O. 1971. Two hundred years of genecology. *Taxon* 20: 653–722.

Levin, D. A. 1993. Local speciation in plants: The rule not the exception. *Systematic Botany* 18:
197-208.

Levin, D. A. 2012. The long wait for hybrid sterility in plants. *New Phytologist* 196: 666-670.

Linhart, Y. B., and M. C. Grant. 1996. Evolutionary significance of local genetic differentiation
in plants. *Annual Review of Ecology and Systematics* 27: 237–277.

Lowry, D. B. 2010. Integrating genetics, geography, and local adaptation to understand ecotype
formation in the yellow monkeyflower, *Mimulus guttatus.* PhD dissertation, Duke
University, Durham, NC, USA.

Lowry, D. B. 2012. Ecotypes and the controversy over stages in the formation of new species.
Biological Journal of the Linnean Society 106: 241–257.

Lowry, D. B., R. C. Rockwood, and J. H. Willis. 2008. Ecological reproductive isolation of coast
and inland races of *Mimulus guttatus*. *Evolution* 62: 2196–2214.

Lowry, D. B., M. C. Hall, D. E. Salt, and J. H. Willis. 2009. Genetic and physiological basis of

adaptive salt tolerance divergence between coastal and inland *Mimulus guttatus*. *New*

Phytologist 183: 776–788.

Lowry, D. B., and J. H. Willis. 2010. A widespread chromosomal inversion polymorphism

contributes to a major life-history transition, local adaptation, and reproductive

isolation. *PLoS Biology* 8: e1000500.

Lowry, D. B., D. Popovic, D. J. Brennan, and L. M. Holeski. 2019. Mechanisms of a locally

adaptive shift in allocation among growth, reproduction, and herbivore resistance in

Mimulus guttatus. *Evolution* 73: 1168-1181.

Mallet, J., N. Besansky, and M. W. Hahn. 2015. How reticulated are species? *Bioessays* 38: 140-

149.

Nesom, G. L. 2012. Taxonomy of *Erythranthe* sect. Simiola (Phrymaceae) in the USA and

Mexico. *Phytoneuron* 2012-40: 1–123.

Nesom, G. L. 2014. Further observations on relationships in the *Erythranthe guttata* group

(Phrymaceae). *Phytoneuron* 2014-93: 1–8

Oneal, E., D. B. Lowry, K. M. Wright, Z. Zhu, and J. H. Willis. 2014. Divergent population

structure and climate associations of a chromosomal inversion polymorphism across the

Mimulus guttatus species complex. *Molecular Ecology* 23: 2844-2860.

Quinn, J. A. 1978 Plant ecotypes: ecological or evolutionary units? *Bulletin of the Torrey*

Botanical Club 1: 58-64.

R Core Team. 2019. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria.

Raven, P. H. 1980. Hybridization and the nature of species in higher plants. *Canadian Botanical*

Association Bulletin 13: 3-10.

Stebbins, G. L. 1980. Botany and the synthetic theory of evolution. p139-152. In: Mayr E

Provine WB, eds. *Evolutionary synthesis: perspectives on the unification of biology.*

Harvard University Press, Cambridge, MA, USA.

Taylor, S. A., and E. L. Larson. 2019. Insights from genomes into the evolutionary importance

and prevalence of hybridization in nature. *Nature Ecology and Evolution* 3: 170-177.

Telewski, F. W. 2021. Mechanosensing and plant growth regulators elicited during the

thigmomorphogenetic response. *Frontiers in Forests and Global Change* 3: 574096.

Turesson, G. 1922a. The species and the variety as ecological units. *Hereditas* 3: 100-113.

Turesson, G. 1922b. The genotypic response of the plant species to habitat. *Hereditas* 3: 211-

350.

Twyford, A. D., and J. Friedman. 2015. Adaptive divergence in the monkey flower *Mimulus*

guttatus is maintained by a chromosomal inversion. *Evolution* 69: 1476-1486.

Vickery, R. K. 1952. A study of the genetic relationships in a sample of the *Mimulus guttatus*

complex. PhD Dissertation, Stanford University, Stanford, CA, USA.

Vickery, R. K. 1978. Case studies in the evolution of species complexes in *Mimulus*.

Evolutionary Biology 11: 405–507.

Wu, C. A., D. B. Lowry, A. M. Cooley, K. M. Wright, Y. W. Lee, and J. H. Willis. 2008.

Mimulus is an emerging model system for the integration of ecological and genomic

studies. *Heredity* 100: 220-230.

578 **TABLES**

Table 1. Regressions of climate factors that were highly-loaded on PC1 for both experiments against latitude

Greenhouse Experiment 1		Estimate	F	Multiple R²	P
	Annual Precipitation	0.045939	401.9	0.8816	< 2.2e-16
	Mean Annual Temperature	-2.3367	237.2	0.8146	< 2.2e-16
	Wettest Quarter Precipitation	0.09784	239.6	0.8161	< 2.2e-16
	Coldest Quarter Precipitation	0.10375	200.9	0.7882	< 2.2e-16
	Mean Wind Speed, 1st Quarter	2.1737	50.06	0.4811	3.11E-09
	Mean Wind Speed, 4th Quarter	1.6486	74.6	0.5801	9.37E-12
	Wettest Month Precipitation	0.2877	191.4	0.7799	< 2.2e-16
	Precipitation Seasonality	-0.27944	607.2	0.9183	< 2.2e-16
	Warmest Quarter Precipitation	0.58611	470.4	0.897	< 2.2e-16
	Mean Temperature, Coldest Quarter	-1.8891	163.2	0.7514	< 2.2e-16
Greenhouse Experiment 2					
	Annual Precipitation	0.048022	242.2	0.7962	< 2.2e-16
	Mean Annual Temperature	-2.538	315.9	0.8359	< 2.2e-16
	Precipitation Seasonality	0.29744	553.1	0.8992	< 2.2e-16
	Warmest Quarter Precipitation	0.60288	430.6	0.8741	< 2.2e-16
	Wettest Quarter Precipitation	0.10225	169.2	0.7318	< 2.2e-16
	Mean Wind Speed, 1st Quarter	2.3456	47.42	0.4334	3.38E-09
	Mean Wind Speed, 4th Quarter	1.8108	67.49	0.5212	1.68E-11
	Coldest Quarter Precipitation	0.108175	143.3	0.698	< 2.2e-16
	Driest Quarter Precipitation	0.5965	372.9	0.8574	< 2.2e-16
	Driest Month Precipitation	2.5381	223.7	0.783	< 2.2e-16
	Wettest Month Precipitation	0.30039	143.6	0.6984	< 2.2e-16
	Mean Temperature, Coldest Quarter	-1.8411	232.1	0.7892	< 2.2e-16

579 Note: F-statistic degrees of freedom for Greenhouse Experiment 1 were 1,54 and were 1,62 for Greenhouse Experiment 2

 FIGURES

590 **Figure 1.** Linear regression of the relationship between latitude of origin for *M. guttatus*

591 populations and the average heights of individuals of these populations grown in the

592 greenhouses. Indigo circle points represent populations raised in Greenhouse 1, and purple

593 triangle points represent populations raised in Greenhouse 2. Both regressions are statistically

594 significant (p < 0.01).

Figure 2. Variation in plant height in *Mimulus guttatus.* A) Divergence in height between plants

from the Central California coast (left) and Oregon (right) when grown in a greenhouse common

garden experiment. B) Tall plants growing along the Central California coast reach up to 2

meters in height. C) Plants growing along the coast in Oregon are more compact in growth form.

Figure 3. Mean wind speeds (m/s) along the United States Pacific Coast reported annually (A),

during January (B), April (C), and August (D). The dashed line indicates 42° N, the latitude of

the Oregon coast at which wind speeds start declining going further north.

SUPPLEMENTARY MATERIAL

Table S1. Principal Component Loadings and Coordinates for Greenhouse Experiment 1

Climate Variables	PC1 loading	PC2 loading	PC3 loading	PC4 loading	PC1 coordinate	PC2 coordinate	PC3 coordinate	PC4 coordinate
Q1Wind	4.7449265	3.8933108	1.8929443	0.0074429	0.8198760	-0.4276365	0.2248873	0.0092769
Q2Wind	0.0124223	2.3409782	29.7008855	0.0484955	-0.0419502	-0.3315995	0.8908012	0.0236802
Q3Wind	1.7739690	3.3327906	17.7481881	0.0103884	0.5013101	-0.3956574	0.6886096	-0.0109600
Q4Wind	5.2802762	3.1766903	1.0233375	0.0205459	0.8648917	-0.3862804	0.1653505	-0.0154133
WindAnnual	2.5906409	4.4450227	15.2435472	0.0000002	0.6058108	-0.4569327	0.6381740	-0.0000490
AnnTemp	6.3433504	0.2197840	0.0108660	4.1278805	-0.9479660	-0.1016046	-0.0170385	-0.2184728
DiurnalRange	3.4968500	4.8877912	5.3021721	9.5279702	-0.7038367	0.4791501	0.3763769	0.3319204
Isothermality	3.8984923	2.2127317	1.8480043	18.1269951	-0.7431590	-0.3223886	0.2222018	0.4578221
TempSeasonality	0.4632877	17.6388953	1.7888015	3.7936280	0.2561880	0.9102296	0.2186136	-0.2094407
MaxTemp_WarmMonth	3.9263041	7.0925750	3.3380579	0.3554186	-0.7458051	0.5771879	0.2986366	-0.0641068
MinTemp_ColdMonth	1.1783690	11.6069940	2.2681721	17.6298756	-0.4085771	-0.7383718	-0.2461694	-0.4515007
TempAnnRange	1.4526542	13.6196131	4.7440718	1.6514689	-0.4536433	0.7998299	0.3560178	0.1381876
TempWetQuarter	4.7505953	5.7312285	0.0955002	1.3391414	-0.8203656	-0.5188468	-0.0505124	-0.1244363
TempDryQuarter	3.9910093	4.8960362	1.1994531	11.2226697	-0.7519254	0.4795540	0.1790143	-0.3602317
TempWarmQuarter	4.4405749	3.9872096	0.6432115	10.3684736	-0.7931458	0.4327627	0.1310910	-0.3462511
TempColdQuarter	5.0049115	4.8110737	0.2660906	0.6330349	-0.8420378	-0.4753749	-0.0843162	-0.0855554
AnnPrecip	6.5362397	0.4393174	0.3745736	2.2974578	0.9622710	0.1436496	0.1000379	-0.1629887
PrecipWetMonth	5.9663180	0.4246475	1.0996666	5.9058571	0.9193621	0.1412308	0.1714062	-0.2613216
PrecipDryMonth	4.4475359	2.0116984	4.3691156	1.9567445	0.7937672	0.3073949	-0.3416590	0.1504184
PrecipSeasonality	5.9642127	0.3193915	1.0051339	1.2306553	-0.9191999	-0.1224833	0.1638732	-0.1192894
PrecipWetQuarter	6.2284665	0.4230799	0.9455099	4.0132959	0.9393425	0.1409699	0.1589385	-0.2154192
PrecipDryQuarter	5.5609895	1.1543451	2.3090199	0.4665059	0.8875840	0.2328537	-0.2483762	0.0734450
PrecipWarmQuarter	5.8628034	0.9038088	1.5353738	0.4709767	0.9113518	0.2060409	-0.2025364	0.0737961
PrecipColdQuarter	6.0848009	0.4309863	1.2483029	4.7950773	0.9284459	0.1422810	0.1826233	-0.2354678

Table S2. Principal Component Loadings and Coordinates for Greenhouse Experiment 2

Climate Variables	PC1 loading	PC2 loading	PC3 loading	PC4 loading	PC5 loading	PC6 loading	PC1 coordinate	PC2 coordinate	PC3 coordinate	PC4 coordinate	PC5 coordinate	PC6 coordinate
Q1Wind	4.3843848	3.2314270	2.4502873	0.7983780	1.8765040	22.3732323	0.7671537	0.4217511	-0.2529117	-0.0861679	-0.1091620	0.3333754
Q2Wind	0.3768277	3.6407226	24.8838252	1.0983227	1.9832131	2.7193330	-0.2249052	0.4476647	-0.8059702	-0.1010662	-0.1122229	-0.1162252
Q3Wind	0.8635182	5.4417680	18.1054543	0.0104486	12.0414096	0.4329545	0.3404582	0.5473046	-0.6874880	0.0098576	0.2765257	0.0463757
Q4Wind	5.1450571	3.1904171	1.4492831	0.0179842	0.0047425	12.2987882	0.8310424	0.4190664	-0.1945078	-0.0129326	0.0054878	0.2471726
WindAnnual	1.5963802	6.0948328	16.4460761	0.2694635	0.5216136	2.9070556	0.4629097	0.5792152	-0.6552266	-0.0500600	0.0575534	0.1201699
AnnTemp	6.7184529	0.1776023	0.0038761	5.3058906	4.8884824	0.5609606	-0.9496481	0.0988743	-0.0100591	0.2221366	0.1761911	0.0527880
DiurnalRange	3.7500835	3.9658549	4.8549101	9.5173513	5.1346629	3.5233552	-0.7094937	-0.4672265	-0.3560008	-0.2975083	0.1805730	-0.1322962
Isothermality	4.4205587	3.2689307	0.4681782	9.9016747	8.4104450	8.6835226	-0.7703120	0.4241915	-0.1105518	-0.3034557	0.2311034	-0.2076907
TempSeasonality	0.9537103	14.0636287	2.4165093	0.8244544	1.0626286	3.1155190	0.3577967	-0.8798482	-0.2511624	0.0875637	-0.0821462	0.1244040
MaxTemp_WarmMonth	3.0383943	8.3455266	4.4200214	0.2976827	0.5112156	0.1636737	-0.6386312	-0.6777756	-0.3396820	0.0526160	0.0569769	0.0285140
MinTemp_ColdMonth	1.6368959	9.7506077	1.5771838	20.1062109	0.3569091	0.3101932	-0.4687471	0.7326135	0.2029091	0.4324203	0.0476075	0.0392541
TempAnnRange	0.7308812	13.5203534	5.0073569	2.2282360	0.1019958	0.0074547	-0.3132214	-0.8626867	-0.3615469	-0.1439533	0.0254500	0.0060853
TempWetQuarter	4.8660637	5.2473663	0.1227338	2.2123170	3.8769347	0.1526979	-0.8081965	0.5374397	0.0566034	0.1434381	0.1569066	-0.0275414
TempDryQuarter	3.1071548	6.8505531	2.8384888	9.7881553	1.7452617	4.8230355	-0.6458171	-0.6140757	-0.2722099	0.3017112	0.1052755	0.1547852
TempWarmQuarter	3.6982025	5.7268566	1.9385112	10.3630140	3.2066364	3.6991858	-0.7045688	-0.5614579	-0.2249544	0.3104445	0.1426993	0.1355571
TempColdQuarter	5.1383409	4.4257274	0.2993206	1.7181994	5.6880054	0.0814673	-0.8304998	0.4935730	0.0883952	0.1264089	0.1900539	-0.0201169
AnnPrecip	6.7487340	0.1245649	0.7205310	3.5171208	0.3414570	6.0570145	0.9517858	-0.0828051	-0.1371471	0.1808567	0.0465656	-0.1734567
PrecipWetMonth	6.1215200	0.1188437	1.7608788	7.8431993	0.0643518	8.7286477	0.9064789	-0.0808811	-0.2144002	0.2700770	-0.0202152	-0.2082297
PrecipDryMonth	5.1432074	1.3473748	3.1582917	1.2706166	15.4562933	1.6896000	0.8308930	-0.2723349	0.2871353	-0.1087047	0.3132922	0.0916138
PrecipSeasonality	6.4834833	0.1558699	0.8209399	0.8501078	5.4040727	0.3903922	-0.9328939	0.0926276	-0.1463916	0.0889156	-0.1852497	-0.0440372
PrecipWetQuarter	6.3767695	0.1285835	1.5239047	5.5203626	0.0170095	8.1894952	0.9251847	-0.0841301	-0.1994524	0.2265817	-0.0103930	-0.2016962
PrecipDryQuarter	6.1080873	0.5747741	1.6439590	0.0523528	15.6983277	0.0274041	0.9054838	-0.1778720	0.2071600	-0.0220654	0.3157356	0.0116675
PrecipWarmQuarter	6.4077079	0.4915632	1.1108009	0.0590273	11.4423299	0.0083199	0.9274263	-0.1644935	0.1702858	-0.0234297	0.2695591	-0.0064288
PrecipColdQuarter	6.1855840	0.1162509	1.9786777	6.4294294	0.1654978	9.0566975	0.9112099	-0.0799940	-0.2272730	0.2445272	-0.0324185	-0.2121066

Table S3. *M. guttatus* population codes and coordinates

PopCode	Latitude	Longitude
ALA	58.00433	-135.743
BCB	36.06285	-121.592
BHE	38.30703	-123.058
BLN	42.83772	-124.56
BND	43.07971	-124.43
BON	37.00016	-122.182
BRK	42.06429	-124.301
BYR	43.34902	-124.346
CAB	39.3592	-123.817
CKI	45.24213	-123.969
CMD	40.40923	-124.392
CML	41.04636	-124.123
COV	40.02157	-124.069
CPT	36.9746	-121.945
CRD	36.30686	-121.85
CRU	35.84372	-121.404
CVG	38.3723	-123.055
DAV	37.02498	-122.218
DIL	38.25167	-122.968
DIS	46.3053	-124.072
FPT	37.80918	-122.475
FRA	39.0083	-123.694
GBM	41.37863	-124.07
GRE	37.07795	-122.268
GRZ	40.56885	-124.212
GWC	37.56394	-122.514
HEC	44.13536	-124.124
HOC	47.3854	-123.147
HUM	39.92726	-123.761
LAG	41.23157	-124.106
LBG	49.07185	-125.768
LJA	39.86053	-123.902
LOK	45.33825	-123.977
MTO	40.28343	-124.36
MTR	37.5517	-122.514
MXX	40.31068	-124.267
NAV	39.1869	-123.757
OPB	42.46402	-124.423
ORO	35.27335	-120.889
OSW	45.76108	-123.967
OTT	44.75224	-124.064
PCR	41.39856	-124.069
PGR	36.62799	-121.921
PPT	41.13709	-124.152
PTL	42.28225	-124.405
PTR	38.04861	-122.869
PTU	44.27916	-124.112
PUN	44.74484	-124.06
SCC	48.41262	-124.012
SDB	35.92127	-121.467
SIM	44.87659	-124.038
SRN	38.72593	-123.481
STB	37.89132	-122.635
SUR	36.33194	-121.889
SWB	39.03598	-123.691
TSU	41.75983	-124.222
USB	39.83218	-123.849
VIC	36.04834	-121.589
WCK	42.00256	-124.213
WCR	35.89838	-121.461
WEV	35.42313	-120.789
WOD	39.31239	-123.803
WPT	42.21213	-124.377
WPV	39.59383	-123.785
WSZ	36.96233	-122.076
WTB	38.40525	-123.096
CVR	38.3684	-123.061
MUR	37.87685	-122.558
OZF	38.95578	-123.66

617