Multiple clines within an ecotype of the yellow monkeyflower, *Mimulus guttatus*

Thomas Zambiasi[1,2,3*], David B. Lowry[1,4,5]

[1]Department of Plant Biology, Michigan State University, East Lansing, MI, USA

[2]Department of Biology, Indiana University, Bloomington, IN, USA

[3]Evolution, Ecology, and Behavior Program, Indiana University, Bloomington, IN, USA

[4]Evolution, Ecology, and Behavior Program, Michigan State University, East Lansing, MI, USA

[5]Plant Resilience Institute, Michigan State University, East Lansing, MI, USA

*For correspondence:

Thomas Zambiasi tzambias@iu.edu

Word count: 5133

Tables: 3

Figures: 1

Color figures: 4

Supplementary information: 6 Tables, 3 Color Figures

RUNNING HEAD: A cline within an ecotype

ABSTRACT

Premise: A key goal of evolutionary biologists is to understand how and why genetic variation is

partitioned within species. In the yellow monkeyflower, *Mimulus guttatus* (syn. *Erythranthe*

guttata), coastal perennial populations collectively constitute a single genetically and

morphologically differentiated ecotype compared to inland populations of *M. guttatus*. While the

distinctiveness of the coastal ecotype has now been well documented, there is also variation in

environmental factors across the range of the coastal ecotype that could drive differentiation

among its component populations in a more continuous way.

Methods: Based on previous observations of a potential cline within this ecotype, we quantified

plant height across coastal perennial accessions from 74 total populations in a greenhouse

common garden experiment. To evaluate possible environmental factors driving the relationship

between trait variation and latitude, we regressed height against multiple climatic factors,

including temperature, precipitation, and coastal wind speeds.

Results: Multiple traits were correlated with latitude of origin, but none more than plant height.

Plant height was negatively correlated with latitude and plants directly exposed to the open ocean

were shorter than those that were more protected from onshore coastal winds. Further analyses

revealed that height was correlated with climatic factors (precipitation, temperature, and

windspeeds) that were autocorrelated with latitude. We hypothesize that one or more of these

climatic factors drove the evolution of latitudinal clinal variation within the coastal ecotype.

Conclusion: Overall, our study illustrates the complexity of how the distribution of

environmental variation can simultaneously drive the evolution of distinct ecotypes as well as

continuous clines within those ecotypes.

KEY WORDS: Local adaptation, salt spray, *Erythranthe guttata*, wind

INTRODUCTION

Over a century ago, Turesson (1922a,b) recognized that adaptive phenotypic variation in plants

was often highly divergent in correlated sets of traits when comparing groups of populations

derived from different habitats. Turesson referred to these groups of phenotypically similar,

locally-adapted populations as ecotypes (Turesson 1922a,b; Gregor et al. 1936). Several

researchers following Turesson found the concept of ecotypes useful for describing the

distribution of adaptive genetic variation within species (Gregor et al. 1936; Clausen et al. 1948;

Clausen 1951; Böcher 1967; Heywood 2011). Further, these researchers viewed partially

reproductively isolated ecotypes as an intermediate stage in the process of speciation and a way

to directly link natural selection to the formation of new species (reviewed in Lowry 2012).

However, other researchers doubted the validity of Turesson's ecotype concept and argued that

clines (Huxley 1938) best represented how adaptive genetic variation is partitioned across the

geographic distribution of a species' range (Langlet 1963; 1971; Stebbins 1980).

While many contemporary biologists recognize the utility of concepts like ecotypes and/or clines

to describe the distribution of adaptive phenotypic variation within species, there is still a lack of

clarity of the full extent to which phenotypic variation is distributed across the complexity of the

natural landscape (Lowry et al. 2014; Briggs and Walters 2016). In particular, questions remain

about the extent to which geographically widespread ecotypes themselves may harbor clines in

phenotypic variation. This question was of major importance to the biosystematists like Clausen,

Keck, Hiesey, (Clausen et al. 1948) and their contemporaries (Gregor 1930, 1938), but has since

received limited attention.

In this study, we examined the distribution of phenotypic variation within a widespread ecotype

of the yellow monkeyflower, *Mimulus guttatus* (syn. *Erythrante guttata*; Barker et al. 2012)*. M.*

guttatus exhibits high levels of genetic and phenotypic variation across its extensive range

spanning western North America, which makes it a particularly useful system for studying how

local adaptation shapes the distribution of adaptive genetic variation across space (Vickery 1978;

Lowry et al. 2008). Coastal perennial populations of *M. guttatus*, occurring from southern

California to southern Washington State, collectively comprise a distinct ecotype based on

population genetic data (Lowry et al. 2008; Twyford and Friedman 2015). This ecotype has been

designated as a variety (*M. guttatus var. grandis* Greene) or species (*M. grandis* syn. *E. grandis*;

Barker et al. 2012, Nesom 2012, 2014) by taxonomists. Despite these taxonomic designations

due to their morphological distinctiveness, the coastal perennial ecotype is highly interfertile

with the inland annual ecotype of *M. guttatus* (Lowry and Willis 2010). Further, there were only

four fixed genetic differences between the coastal perennial and inland annual populations out of

29,693,578 SNPs surveyed in a population genetic study of this system (Gould et al. 2017).

Based on these results, we believe that it is best to think of the coastal perennial populations as

an intraspecific ecotype within the *Mimulus guttatus* species complex, which is the result of non-

fixed differences in allele frequencies at many loci across the genome (Lowry and Willis 2010;

Twyford and Friedman 2015; Gould et al. 2017).

Several previous studies of local adaptation in *M. guttatus* have found striking genetically-based

phenotypic differences between the coastal perennial ecotype and nearby inland populations in

the western United States (Vickery 1952; Hall et al. 2006, 2010; Lowry et al. 2008; Oneal et al.

2014). Inland *M. guttatus* populations of the coastal mountain ranges of western North America

primarily have an annual life-history, especially those that occur in inland habitats that

completely dry out during summer months (Vickery 1952; Clausen and Heisey 1958; Wu et al.

2008; Lowry et al. 2008). While inland populations in the coast ranges are annuals, perennial

populations do occur with some frequency in areas that maintain soil water through the summer

months (van Kleunen 2007; Oneal et al. 2014; Twyford & Friedman 2015). The occurrence of

inland perennial populations increases further inland, especially at higher elevations in the Sierra

Nevada and Cascade Mountain ranges (Coughlin et al. 2021). The level of morphological

differentiation between coastal perennial and inland annual ecotypes is immense (Lowry &

Willis 2008). Further, coastal perennial populations are separated genetically from inland

populations, both annual and perennial, in previously published principal component plots (Fig.

3 of Lowry 2012 and Fig. S2 of Twyford & Friedman 2015). In contrast to the inland annuals,

the coastal perennial ecotype occurs in coastal habitats that are regularly damp and foggy during

summer months (Vickery 1952; Clausen and Heisey 1958; Hitchcock and Cronquist 1973; Hall

and Willis 2006; Lowry et al. 2008). In greenhouse common garden experiments, accessions

derived from coastal habitats had thicker stems, many more prostrate stoloniferous branches,

later flowering times, and produced far more herbivore defensive compounds than nearby inland

annual populations (Hall et al. 2006, 2010; Lowry et al. 2008; Lowry and Willis 2010; Oneal et

al. 2014; Lowry et al. 2019).

Much of the trait differences between the inland annual and coastal perennial ecotypes contribute

to local adaptation (Lowry et al. 2008, 2019; Lowry & Willis 2010; Popovic & Lowry 2019) and

are caused by a few large effect loci, including a large chromosomal inversion polymorphism on

chromosome 8 (Lowry and Willis 2010; Oneal et al. 2014; Twyford & Friedman 2015; Lowry et

al. 2019). Inland perennial populations of *M. guttatus* consistently have the same orientation of

the chromosomal inversion as coastal perennial populations (Lowry & Willis 2010). Indeed, one

of the few regions of the genome that is highly differentiated between inland annual and inland

perennial populations is the chromosome 8 inversion (Oneal et al. 2014; Twyford & Friedman

2015). In this study, we focus on phenotypic variation only among coastal perennial populations,

which we assume all share the same orientation chromosome 8 orientation based on previous

studies (Lowry and Willis 2008; Oneal et al. 2014; Twyford & Friedman 2015; Gould et al.

2017; Kollar et al. 2023).

While there is both phenotypic and genetic divergence between ecotypes of *M. guttatus*, it is

unclear to what extent genetic variation is present within individual ecotypes – either due to

adaptation to widespread environmental gradients or idiosyncratic with regard to geographic

space. However, a previous pilot study suggested the possibility that there may be a latitudinal

cline in plant height within the coastal ecotype (Lowry 2010). Since that initial study, dozens of

additional coastal *M. guttatus* populations have been collected (Fig. 1), which provides an

opportunity for a more comprehensive survey of variation over the entire geographic range of the

coastal perennial ecotype.

In this study, we tested the hypothesis that there is a latitudinal cline in phenotypic variation

within the coastal ecotype. We simultaneously evaluated the role of exposure of the coastal

perennial populations to the open ocean on trait variation. To establish potential sources of

selection, we examined whether there is a relationship between the phenotypes of the accessions

and the climates from where they were collected in nature. Through these efforts, we attempted

to understand whether clinal trait variation occurred within the distinctive coastal perennial

ecotype.

METHODS

Germplasm for experiments –

The seeds used in these experiments originated from natural population collections made

between 2004 and 2016. Seeds were either collected directly from plants in the field or collected

from plants that had been brought from the field back to the greenhouses at Duke University

(Durham, North Carolina, USA) or Michigan State University (East Lansing, Michigan, USA).

All seeds used for the experiment were grown for at least one generation in the greenhouse prior

to the study. For each population, we collected location description information, habitat

description, and GPS coordinates. Photos of the plants in the field for each population have been

posted on iNaturalist and vouchers for a subset of the populations were deposited at the

California Academy of Sciences. Permits were obtained for collections on public lands, where

permission was required. All maternal families used in this study are currently stored at

Michigan State University in airtight containers with silica gel as a desiccant.

Greenhouse Experiment –

To determine whether there was clinal variation in *M. guttatus* traits, we conducted an

experiment in a controlled greenhouse environment. We conducted the greenhouse experiment in

the Plant Ecology Field Lab greenhouse at Kellogg Biological Station (KBS, Hickory Corners,

MI, USA). Seeds from coastal perennial *M. guttatus* ($N = 423$ individuals, 418 seed families, 74

populations) were planted with SUREMIX (Michigan Grower Products, Inc., Galesburg,

Michigan, USA) in 3.5 sq. in. pots without prior cold stratification. We included a variable

number of seed families per population (Range = 1-15, Mean = 5.72 families) and all families

but five were represented by a single individual (BON-2, BON-4, CPT-1, CPT-5, and CRU-1

each had two individuals measured). More information on the populations used in this

experiment, plus the coordinates and maps of the original population locations can be found in

Figure 1 and Table S1 in the supplementary material (Appendix S1; see Supplemental Data with

this article). All germinating seedlings except the centermost plant were removed from the pot

and remaining individuals were watered every other day throughout the experiment. Germination

and flowering dates were recorded to determine flowering time; height of plant at first flower

(cm), width of the second set of true leaves (cm), corolla length (mm), and corolla width (mm)

were measured at the time of first flowering.

Climate Data –

To investigate potential environmental drivers of adaptive phenotypic variation, we extracted

climatic data for each population's collection location of origin from the WorldClim Bioclimatic

variables database (Fick and Hijmans 2017). This data was imported and compiled using R

(v4.2.1, R Core Team 2022) using the *stats* (v4.2.1), *nlme* (v3.1-157)*, raster* (v3.6-26), and *sp*

(v2.1-3) packages.

In addition to the Bioclim data, we extracted oceanic wind speed data from along the US Pacific

Coast using the National Renewable Energy Laboratory's Wind Prospector website, which

featured interactive online maps of wind speeds surveyed by NREL (King et al. 2014; Draxl et

al. 2015a,b). We used Pacific Monthly Offshore Wind Speed map layers to collect wind speeds

for each month from the site nearest to shore (8 km offshore at the closest; Draxl et al. 2015b) at

the latitudes of each of the coastal perennial populations. The wind speeds found through the

NREL Wind Prospector were the averages for each 1.2 km X 1.2 km offshore site and were

modeled at 100 m (Draxl et al. 2017). Due to unavailability of wind data outside of the

contiguous US, populations found in British Columbia and Alaska (denoted by population codes

ALA and LBG) were not included in climate data analyses. Monthly wind speed averages

accessible via NREL were based on observations and statistical analyses over the course of seven

years (King et al. 2014; Draxl et al. 2015a,b), which we used to calculate quarterly and annual

wind speeds for each population collection location. We then matched these climate variables

with their corresponding coordinates and population codes for further analysis.

Statistical Analysis –

Prior to data analyses, we calculated mean trait values for each population to avoid

pseudoreplication in our analyses. Our analyses included a binary factor to identify whether each

population was exposed to (1) or protected (0) from the open ocean. Exposed populations were

located on the cliffs, headlands, and seeps just adjacent to the open Pacific Ocean. In contrast,

protected populations were not directly exposed to the open ocean by being located in inland

roadside ditches and lagoons or on the coastlines of Washington, British Columbia, and Alaska

in locations where they were protected from the open ocean by other land masses. To test

whether there was a relationship between plant traits and the latitude of origin (location from

which each population accession was collected), we modeled trait values against latitude with the

binary predictor for ocean exposure via linear regression in R (v4.2.1, R Core Team 2022) using

the *stats* (v4.2.1) package, evaluated predictor significance via ANOVA (*car* package v3.1-2),

and visualized our results in *ggplot2* (v3.5.0). We modeled both additive and interactive

relationships between the predictors for each trait, but in each case found that the additive model

was a better descriptor of the data per AIC and ANOVA model comparison (both in the *stats*

package v4.2.1).

Although we predicted that latitude would correlate with plant height across space, we wanted to

evaluate which climatic variables correlated with latitude might ultimately be driving observed

clinal patterns. Climate variables are often highly correlated with each other, so we conducted a

principal components analysis (PCA) to reduce the dimensionality of the data prior to further

analyses. We extracted climate data, described above, for each population and performed the

PCA in R (v4.2.1) using the *factoextra* (v1.0.7) and *stats* (v4.2.1) packages. We evaluated all

principal components (PCs) with an eigenvalue greater than 1, following the Kaiser criterion

(Kaiser 1960). We then conducted regression analyses of population mean plant height against

each of these PCs individually and with all as factors in the same model, again including the

exposure factor and using linear models in the *stats* (v4.2.1) package. We also evaluated models

with interactions and polynomial fits and selected the best descriptors for the data via AIC and

ANOVA model comparison. For all regression analyses, we evaluated residuals with the

DHARMa (v.0.4.6) package and confirmed via visual inspection that residuals were normally

distributed for all models. All data organization, analysis, and visualization was conducted in R

Studio (2023.12.0+369 "Ocean Storm" Release, Posit Team 2023).

RESULTS

Trait correlations with latitude –

We found a significant negative relationship between latitude of population origin and plant height within the coastal perennial ecotype in our greenhouse experiment ($R^2 = 0.43$, $F_{2,71} = 26.83$, $P < 0.001$; Figs. 2, 3). Based on our linear modeling, we found that every degree latitude further north translated to a roughly 1 cm decrease in plant height under greenhouse conditions. We also found a significant difference in the height-latitude relationship depending on whether populations were highly exposed to the wind and salt spray from the open Pacific Ocean. In general, protected populations were taller than those exposed to the open ocean (Table 1, Fig. 2). In addition to the strong relationship with height, we found significant, but much weaker relationships between population latitude and width of the second set of true leaves ($R^2 = 0.12$, $F_{2,71} = 4.83$, $P = 0.011$), corolla width ($R^2 = 0.12$, $F_{2,63} = 4.38$, $P = 0.017$), and corolla length ($R^2 = 0.17$, $F_{2,63} = 6.51$, $P = 0.003$). Leaf width generally decreased with latitude, as did corolla width and length; none of these models included a significant term for ocean exposure (Table 1). The relationship of latitude to flowering time ($R^2 = 0.03$, $F_{2,71} = 0.97$, $P = 0.383$) was not significant. Because the correlation between plant height and latitude was so much stronger than for the other traits, we focused the rest of our analyses on the possible causes of this correlation.

The relationship of climatic factors and plant height –

In our greenhouse experiment, we found that climate PC1 through PC3 all had eigenvalues greater than 1. In total, those three PCs accounted for 89.99% of cumulative variance in climate factors. PC1 explained the most variation at 56.39%, followed by PC2 with 22.81%, and PC3 with 10.79%. PC1 was significantly negatively correlated with plant height in a linear regression

model (*PC1 estimate* = -1.09, *exposure estimate* = -5.69, *Multiple R²* = 0.52, $F_{2,62}$ = 33.05, *P* <

0.001). Due to the nonlinear relationship between wind speeds and latitude, we also assessed

nonlinear regression models and found that the relationship between PC1 and plant height was

best explained by a quadratic regression model as determined via AIC (*Multiple R²* = 0.57, $F_{3,61}$

= 27.23, *P* < 0.001; *AIC linear* = 402.52, *AIC quadratic* = 396.46). The quadratic regression

yielded the same qualitative relationships between mean population height and the climate

factors as the linear model; height decreased with increasing values of PC1 (Fig. 4). We also

evaluated regression models of the remaining PCs against plant height. While each showed a

significant relationship with height, PCs 2 and 3 only included significant terms for ocean

exposure (Table 2). However, our model with the PC1 square term and ocean exposure as

predictors was a better descriptor of the data based on AIC than models with PC2 and PC3

(Table 2). We also constructed a linear model with PCs 1-3 and exposure as predictors to

compare its fit with our quadratic PC1 model, but yet again found PC1 to best describe the data

via AIC (*AIC PC1* = 396.46, *AIC All PCs* = 396.82). Following our tests for model fit, we

focused on PC1's loadings for the remainder of our interpretations.

We evaluated which climatic factors loaded onto PC1. Heavily loaded climate variables included

Annual Precipitation, Average Annual Temperature, Precipitation Seasonality, Precipitation of

the Warmest Quarter, Precipitation of the Wettest Quarter, Precipitation of the Coldest Quarter,

Precipitation of the Wettest Month, and Precipitation of the Driest Quarter; Average Wind

Speeds in Quarter 1 (January, February, March) and Quarter 4 (October, November, December)

were not as heavily loaded, but were still represented on PC1 (Table S2). Higher values of PC1

indicated lower Average Annual Temperature and Precipitation Seasonality, but greater Annual

Precipitation, Precipitation of the Warmest Quarter, Precipitation of the Wettest Quarter,

Precipitation of the Coldest Quarter, Precipitation of the Wettest Month, Precipitation of the

Driest Quarter, and Average Wind Speeds in Quarter 1 and Quarter 4 (Table S2). Therefore, any

population at a location corresponding to a high value of PC1 experienced lower temperatures,

higher amounts and greater consistency of precipitation, and faster wind speeds than populations

at sites with a low PC1 value.

273   *Latitude strongly predicts climate for coastal populations –*

To assess how well latitude predicts climate at the sites of our study populations, we regressed

the extracted principal components from our greenhouse experiment against latitude. Our

analysis confirmed that many of the climate variables involved were highly correlated with

latitude. PC1 was the only principal component significantly associated with latitude and the two

variables had a positive relationship (*slope* = 0.89, *Multiple R²* = 0.92, $F_{1,63}$ = 704.9, P < 0.001).

To determine how well variation in individual climatic factors was predicted by latitude, we

regressed all the climate factors that were highly loaded on PC1 individually against latitude

(Table 3).

283   *Patterns of wind speed along the west coast of North America –*

Plants are well known to respond to wind through changes in growth form (Grace 1977;

Gardiner et al. 2016; Telewski 2021). This, combined with the observation of wind speed

loading onto the first PC of climate, led us to evaluate the patterns of wind speed variation more

carefully across the populations that we studied (Fig. 5). We found that, in general, annual wind

speeds increase with latitude (*slope* = 0.07, *Multiple R²* = 0.06, $F_{1,76}$ = 4.465, P = 0.038).

However, the relationship between wind speeds and latitude varied dramatically based on time of

year. Average wind speeds in Quarter 1 (*slope* = 0.19, *Multiple R²* = 0.46, $F_{1,76}$ = 65.9, *P* <

0.001) and Quarter 4 (*slope* = 0.27, *Multiple R²* = 0.50, $F_{1,76}$ = 75.32, *P* < 0.001) of the year were

strongly positively correlated with latitude. In contrast, Quarter 2 wind speeds (*slope* = -0.20,

Multiple R² = 0.20, $F_{1,76}$ = 19.1, *P* < 0.001) showed a negative relationship with latitude and

Quarter 3 wind speeds were not correlated with latitude (*slope* = 0.05, *Multiple R²* = 0.01, $F_{1,76}$ =

0.57, *P* > 0.05).

Finally, we constructed a linear model to evaluate how mean population height varied with

annual mean wind speeds for each population from which accessions were collected. We also

included the ocean exposure term in this model. While annual mean wind speed only had a

marginally significant effect on plant height (*estimate* = -1.29, *P* = 0.073), open ocean exposure

made a clear difference (*estimate* = -6.98, *P* < 0.001) and the model was a significant fit for the

data (*Multiple R²* = 0.26, $F_{2,62}$ = 10.95, *P* < 0.001). In general, *M. guttatus* populations exposed

to the open ocean were shorter than their protected counterparts (Fig. S1).

DISCUSSION

In this study, we found a significant role of latitude and exposure to the open ocean in predicting

genetically based trait variation among populations of the coastal perennial ecotype of *M.*

guttatus. The trait most affected by these two factors was plant height, which was smaller for

populations collected from locations exposed to the open ocean and was negatively correlated

with latitude. Multiple environmental factors may have contributed to the evolution of these

patterns of trait variation. These southern populations generally experience higher temperatures,

less precipitation, and lower velocity winds than more northern populations, but this can vary

based on the time of year. Further, populations exposed to the open ocean often experience

greater wind speeds and more toxic salt spray, which is a byproduct of the wave energy of the

ocean (Boyce 1954; Du & Hesp 2020). The results of this study further reinforce the idea that

genetically-based phenotypic variation can be simultaneously partitioned in disjunct (among

coastal perennial and inland annual ecotypes) and continuous, clinal ways (within ecotypes). We

found similar overall results in a pilot experiment (See Appendix S1), indicating that our

findings are replicable.

Phenotypic variation is partitioned at multiple levels within species

Understanding the patterns by which genetic variation is partitioned among populations within a

species across a geographic landscape has been a matter of considerable debate, especially in the

classic plant evolutionary biological literature (reviewed in Lowry 2012). It is reasonable to

hypothesize that the partitioning of that genetic variation is a function of how environmental

variation is partitioned across the same landscape. However, several previous researchers have

argued against the existence of ecotypes and that only clinal variation exists within species

(Langlet 1971; Quinn 1978; Stebbins 1980; Levin 1993).

A series of prior studies has clearly established that the coastal populations of *Mimulus guttatus*

from central California to southern Washington state collectively comprise a genetically distinct,

geographically widespread ecotype (Lowry et al. 2008; Lowry and Willis 2010; Oneal et al.

2014; Twyford and Friedman 2015). This coastal perennial ecotype clusters as a distinct group

from the rest of the *Mimulus guttatus* species complex based on both a set of morphological traits

and allele frequencies of loci across the genome (Lowry 2012; Oneal et al. 2014; Twyford and

Friedman 2015) but has not accumulated appreciable divergence through fixed genetic

differences (Gould et al. 2017). This study demonstrates that while the coastal perennial ecotype

is strikingly divergent from the inland annual ecotype, it also harbors genetically-based

phenotypic variation that is partitioned along two major environmental gradients. Latitudinal

clines within ecotypes/species have been observed in the classic literature for plant systems such

as *Plantago maritima* (Gregor 1930, 1938) and *Achillea borealis* (Clausen et al. 1948).

Similarly, differences in traits including plant height have been found recently for coastal plants

that occur on dunes versus headlands in *Senecio lautus* (Roda et al. 2013: James et al. 2021).

However, in contrast to *M. guttatus*, the evolution of short headland versus tall dune ecotypes in

S. latutus has occurred independently through repeated parallel evolution (Roda et al. 2013;

James et al. 2021) instead of occurring all within a single coastal ecotype. In the case of the

coastal perennial ecotype of *M. guttatus*, the population genetic evidence suggests that it was

derived at a single point in time from the rest of the species complex and then spread along the

Pacific coast, adapting to local environments as it spread (Lowry et al. 2008; Twyford &

Friedman 2015). During and following the spread of the coastal ecotype, there has likely been

appreciable gene flow with other parts of the species complex, which is further supported by

there only being four fixed nucleotide differences between these groups out of millions of SNPs

(Gould et al. 2017).

The environmental factors responsible for the geographic distribution of trait variation

While we have demonstrated that clines in plant height do occur within the coastal ecotype of

Mimulus guttatus, the question remains as to what environmental factors have driven these

clines. Though our analysis identified multiple factors correlated with both latitude and plant

height, these factors must be carefully considered, as they are autocorrelated with each other and

not all of them may have an impact on the selective advantage of plant height. Regardless of

which environmental variables dictate the shape of the height cline, we have shown that

exposure to the open ocean is predictive of shorter plant height in *M. guttatus* populations (Figs.

2, S1). These results are consistent with classic observations by Boyce (1954) and recent studies

in *Setaria viridis* (Itoh 2021; Itoh et al. 2024).

Other researchers studying the evolution of ecotypes in the same region have found similar

patterns in plant height variation along the Pacific coast of North America. These studies were

far less comprehensive, but are useful, as they suggest that the patterns observed within the

coastal ecotype of *M. guttatus* are not unique. For example, Clausen et al. (1948) found similar

genetic variation with the coastal ecotype of *Achillea borealis* when comparing populations in

Northern California, where populations north of the San Francisco Bay were shorter in stature

than those from farther south in California in common garden experiments.

The causes of the latitudinal cline of plant height cannot be determined by this study alone and

likely are complex. A global study of plant heights collected from an array of ecosystems found

that there is a strong negative relationship between latitude and plant height (Moles et al. 2009).

However, none of the environmental variables evaluated in that study were very strongly

correlated with plant height. For our intraspecific study, temperature and precipitation were the

most highly correlated with latitude. Both of these variables could directly or indirectly be

driving the evolution of plant height in this system. For example, the longer growing season in

the southern end of the range of the coastal perennial ecotype could lead to greater competition,

which would favor greater height. A positive relationship between temperature and plant height

has been observed within species for altitudinal gradients (Totland & Birks 1996; Macek & Leps

2008; Pellissier et al. 2010), but other factors could also drive the overall negative association of

elevation and height.

Perhaps the most compelling factor driving patterns of plant height is the combination of wind

and salt spray. While wind is less closely correlated with latitude than other environmental

factors, whether a population is exposed to the open ocean or is protected from salt spray by

being located slightly inland can have a large effect on the impacts of wind and salt spray on

plants (Barbour 1978; Du and Hesp 2020). Wind and salt spray have long been thought to drive

the evolution of the stature of coastal plant species. For example, Boyce (1954) recognized that

this compact growth form is advantageous for coastal plants because it helps them to avoid the

impacts of salt spray by being more sheltered from the wind. Onshore wind from the ocean can

be intense along the Pacific Coast, depending on the location and time of year (Grace 1977;

Gilliam 2002). The intensity of wind speed increases with the distance above the ground (Grace

1977; Jaffe and Forbes 1993). While wind can be damaging by itself, it is particularly damaging

to plants along ocean coastlines because it contains aerosolized oceanic salt spray (Boyce 1954;

Barbour 1978; Cheplick and Demetri 1999; Griffiths et al. 2006; Kouali et al. 2017; Du and Hesp

2020). In winter months (Quarters 1 and 4, e.g. Fig 5b), there is a strong correlation between

latitude and wind speeds, which could play a role in the evolution of shorter plant heights in

more northern populations of this perennial plant ecotype. However, wind speed patterns are

complex and their relationship with latitude shifts greatly over the year. Across the range of the

coastal perennial ecotype, wind speeds and wave energy that drive salt spray are greater in the

spring (April, May) than the summer (July, August) (Gilliam 2002; García-Reyes and Largier

2012). However, those winds speeds dissipate most during the summer in the region south of the

San Francisco, especially in more protected areas, like the southern Monterey Bay (Fig. 3b),

where plants in the field reach up to two meters in height (Lowry *personal observation*). In

Northern California and Southern Oregon, high coastal windspeeds remain elevated throughout

the summer months, which coincides with when this species flowers. Since reproductive

structures are often the most sensitive to salt spray (Griffiths et al. 2006), maintaining a compact

growth form when flowering under higher windspeed/salt spray conditions is likely to be

advantageous. Field experimentation will be necessary to deconstruct how the time of year

interacts with potential selection imposed by wind speeds and salt spray.

CONCLUSIONS

Overall, our study demonstrates that there is dramatic and likely important phenotypic variation

distributed within distinct ecotypes contained within species. Therefore, as in other species, like

switchgrass (*Panicum virgatum*; Lowry et al. 2014), it is critical to account for both ecotypic and

clinal variation when conceptualizing how functional genetic variation is distributed within a

species. The exact causes of the clines within the coastal perennial ecotype of *M. guttatus* are

still unclear, especially for trait correlations with latitude. However, the fact that exposure to the

open ocean is predictive of plant height, suggests that wind and salt spray may play a crucial role

in driving genetically-based variation in plant height in this system.

ACKNOWLEDGEMENTS

This work is dedicated to the memory of Don Levin, whose papers and conversations challenged

us to think hard about the nature of ecotypes. We would like to thank Mauricio Swartzendruber,

Marty Schmidt, Simran Patel, Thomas Young, and Will Burke (Michigan State University) for

assisting with plant care and data collection. We also want to thank all of the public agencies,

private landowners, and First Nations Peoples who provided us with permission over the years to

make collections for our research. Maddie James and two anonymous reviewers provided

excellent comments on multiple versions of this manuscript, which greatly helped to improve it.

Nate Emery, Anne-Sophie Bohrer, Linnea Fraser, Milagros del Pilar Jimenez-Hernandez, Jen

Lau, Mackenzie Caple, and Emma Boehm provided advice with analysis using R. Caroline Draxl

of NREL also shared her insight regarding the collection of offshore wind data. This work was

funded by National Science Foundation Division of Integrative Organismal Systems Grants to

DBL (IOS-1855927 and IOS-2153100).

AUTHOR CONTRIBUTIONS

DBL conceived of and oversaw the experiments and analyses. TZ conducted the experiments and

analyses. Both authors wrote the manuscript.

DATA AVAILABILITY STATEMENT

All data will be made available at publication, either as supplementary information or on Dryad.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at

the end of the article.

Appendix S1: Pilot experiment and supplementary tables/figures

LITERATURE CITED

Barbour, M. G. 1978. Salt spray as a microenvironmental factor in the distribution of beach

455   plants at Point Reyes, California. *Oecologia* 32: 213-224.

456   Barker, W. R., G. L. Nesom, P. M. Beardsle, and N. S. Fraga. 2012 A taxonomic conspectus of

457   Phrymaceae: a narrowed circumscription for Mimulus, new and resurrected genera, and

458   new names and combinations. *Phytoneuron* 39: 1-60.

459   Böcher, T. W. 1967. Continuous variation and taxonomy. *Taxon* 16: 255-258.

460   Boyce, S. G. 1954. The salt spray community. *Ecological Monographs* 24: 29– 67.

461   Briggs, D., and S. M. Walters. 2016. Plant variation and evolution. Cambridge University Press,

462   Cambridge, UK.

463   Cheplick, G. P., and H. Demetri. 1999. Impact of saltwater spray and sand deposition on the

464   coastal annual *Triplasis purpurea* (Poaceae). *American Journal of Botany* 86: 703–710.

Clausen, J. 1951. Stages in the evolution of plant species. Cornell University Press, Ithaca, NY,

USA.

Clausen, J., D. D. Keck, and W. M. Hiesey. 1948. Experimental studies on the nature of species.

III. Environmental responses of climatic races of *Achillea*. Carnegie Institution of

Washington, Washington, DC, USA.

Clausen, J., and W. M. Heisey. 1958. Experimental studies on the nature of species. IV. Genetic

structure of ecological races. Carnegie Institution of Washington, Washington, DC, USA.

Draxl, C., B. M. Hodge, A. Clifton, and J. McCaa. 2015a. Overview and Meteorological

Validation of the Wind Integration National Dataset Toolkit (Technical Report,

NREL/TP-5000-61740). National Renewable Energy Laboratory, Golden, CO, USA.

Draxl, C., B. M. Hodge, A. Clifton, and J. McCaa. 2015b. The Wind Integration National

Dataset (WIND) Toolkit. *Applied Energy* 151: 355-366.

Draxl, C., C. Philips, G. Scott, and W. Musial. 2017. "Characterization of Offshore Wind

Resources in the Contiguous United States." Presented at the 2017 NAWEA Conference,

September 29, Ames, Iowa, USA. Retrieved July 18, 2022.

Du, J., and P. A. Hesp. 2020. Salt spray distribution and its impact on vegetation zonation on

coastal dunes: a review. *Estuaries and Coasts* 43: 1885-1907.

Fick, S. E., and R. J. Hijmans. 2017. WorldClim 2: new 1km spatial resolution climate surfaces

for global land areas. *International Journal of Climatology* 37: 4302-4315.

García-Reyes, M., and J. L. Largier. 2012. Seasonality of coastal upwelling off central and

northern California: New insights, including temporal and spatial variability. *Journal of*

Geophysical Research 117: C03028.

Gardiner, B., P. Berry, and B. Moulia. 2016. Review: Wind impacts on plant growth, mechanics

and damage. *Plant Science* 245: 94-118.

Gilliam, H. 2002. Weather of the San Francisco Bay Region. University of California Press,

Berkeley, CA, USA.

Gould, B. A., Y. Chen, and D. B. Lowry. 2017. Pooled ecotype sequencing reveals candidate

genetic mechanisms for adaptive differentiation and reproductive isolation. *Molecular*

Ecology 26: 163-177.

Grace, J. 1977. Plant Response to Wind. Academic Press, New York, NY, USA.

Gregor, J. W. 1930. Experiments on the genetics of wild populations. I. *Plantago maritima*.

Journal of Genetics 22: 15-25.

Gregor, J. W. 1938. Experimental taxonomy II: Initial population differentiation in *Plantago*

maritima L. of Britain. *New Phytologist* 37: 15-49.

Gregor, J. W., V. McM. Davey, and J. M. S. Lang. 1936. Experimental taxonomy I.

Experimental garden technique in relation to the recognition of the small taxonomic

units. *New Phytologist* 35: 323–350.

Griffiths, M. E., R. P. Keith, and C. M. Orians. 2006. Direct and indirect effects of salt spray and

fire on coastal heathland plant physiology and community composition. *Rhodora* 108:

32–42.

Hall, M. C., C. J. Basten, and J. H. Willis. 2006. Pleiotropic quantitative trait loci contribute to

population divergence in traits associated with life-history variation in *Mimulus guttatus*.

Genetics 172: 1829-1844.

Hall, M. C., D. B. Lowry, and J. H. Willis. 2010. Is local adaptation in *Mimulus guttatus* caused

by trade-offs at individual loci? *Molecular Ecology* 19: 2739–2753.

Heywood, V. H. 2011. The genesis of IOPB: a personal memoir. *Taxon* 60: 320-323.

Hitchcock, C. L., and A. Cronquist. 1973. Flora of the Pacific Northwest. University of

Washington Press, Seattle, WA, USA.

Huxley, J. 1938. Clines: an auxiliary taxonomic principle. *Nature* 142: 219-220.

Itoh, M. 2021 Phenotypic variation and adaptation in morphology and salt spray tolerance in

coastal and inland populations of *Setaria viridis* in central Japan. *Weed Research* 61:199-

209.

Itoh M, Fukunaga K, Osako T. 2024. Local adaptation in parapatric and sympatric mosaic

coastal habitats through trait divergence of *Setaria viridis*. *Journal of Ecology* 112: 784-

799.

Jaffe, M. J., and S. Forbes. 1993 Thigmomorphogenesis: the effect of mechanical perturbation on

plants. *Plant Growth Regulation* 12: 313-324.

James, M. E., M. J. Wilkinson, D. M. Bernal, H. Liu, H. L. North, J. Engelstädter, and D. Ortiz-

Barrientos. 2021. Phenotypic and genotypic parallel evolution in parapatric ecotypes of

Senecio. *Evolution* 75: 3115-3131.

Kaiser, H. F. 1960. The application of electronic computers to factor analysis. *Educational and*

Psychological Measurement 20:141-151.

King, J., A. Clifton, and B. M. Hodge. 2014. Validation of Power Output for the WIND Toolkit

(Technical Report, NREL/TP-5D00-61714). National Renewable Energy Laboratory,

Golden, CO, USA.

Kouali, A., M. Mouradi, L. Latrach, A. Khadraji, O. Elasri, Z. Baicha, A. Berrichi, N. Kouddane,

and A. Boukroute. 2017. Study of effects of seawater salt spray on growth, chlorophyll

fluorescence and chlorophyll content in three coastal species of Morocco. *Journal of*

Materials and Environmental Sciences 8: 4385–4390.

Langlet, O. 1963. Patterns and terms of intraspecific ecological variability. *Nature* 200: 347–348.

Langlet, O. 1971. Two hundred years of genecology. *Taxon* 20: 653–722.

Levin, D. A. 1993. Local speciation in plants: The rule not the exception. *Systematic Botany* 18:

197-208.

Lowry, D. B. 2010. Integrating genetics, geography, and local adaptation to understand ecotype

formation in the yellow monkeyflower, *Mimulus guttatus.* PhD dissertation, Duke

University, Durham, NC, USA.

Lowry, D. B. 2012. Ecotypes and the controversy over stages in the formation of new species.

Biological Journal of the Linnean Society 106: 241–257.

Lowry, D. B., R. C. Rockwood, and J. H. Willis. 2008. Ecological reproductive isolation of coast

and inland races of *Mimulus guttatus*. *Evolution* 62: 2196–2214.

Lowry, D. B., M. C. Hall, D. E. Salt, and J. H. Willis. 2009. Genetic and physiological basis of

adaptive salt tolerance divergence between coastal and inland *Mimulus guttatus*. *New*

Phytologist 183: 776–788.

Lowry, D. B., and J. H. Willis. 2010. A widespread chromosomal inversion polymorphism

contributes to a major life-history transition, local adaptation, and reproductive

isolation. *PLoS Biology* 8: e1000500.

Lowry, D. B, K. D. Behrman, P. Grabowski, G. P. Morris, J. R. Kiniry, T. E. Juenger. 2014.

Adaptation between ecotypes and along environmental gradients in *Panicum virgatum*.

The American Naturalist 183: 682-692.

Lowry, D. B., D. Popovic, D. J. Brennan, and L. M. Holeski. 2019. Mechanisms of a locally

adaptive shift in allocation among growth, reproduction, and herbivore resistance in

Mimulus guttatus. *Evolution* 73: 1168-1181.

Macek, P., and Leps, J. 2008. Environmental correlates of growth traits of the stoloniferous plant

Potentilla palustris. *Evolutionary Ecology* 22: 419–435.

Nesom, G. L. 2012. Taxonomy of *Erythranthe* sect. Simiola (Phrymaceae) in the USA and

Mexico. *Phytoneuron* 2012-40: 1–123.

Nesom, G. L. 2014. Further observations on relationships in the *Erythranthe guttata* group

(Phrymaceae). *Phytoneuron* 2014-93: 1–8

Oneal, E., D. B. Lowry, K. M. Wright, Z. Zhu, and J. H. Willis. 2014. Divergent population

structure and climate associations of a chromosomal inversion polymorphism across the

Mimulus guttatus species complex. *Molecular Ecology* 23: 2844-2860.

Pellissier, L., B. Fournier, A. Guisan, and P. Vittoz P. 2010. Plant traits co-vary with altitude in

grasslands and forests in the European Alps. *Plant Ecology* 211: 351-365.

Posit Team. 2023. RStudio: Integrated Development Environment for R. Posit Software, PBC,

Boston, MA. URL http://www.posit.co/.

Quinn, J. A. 1978 Plant ecotypes: ecological or evolutionary units? *Bulletin of the Torrey*

Botanical Club 1: 58-64.

R Core Team. 2022. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Roda, F., L. Ambrose, G. M. Walter, H. L. Liu, A. Schaul, A. Lowe, P. B. Pelser, P. Prentis, L.

H. Rieseberg, and D. Ortiz-Barrientos. 2013. Genomic evidence for the parallel evolution

of coastal forms in the *Senecio lautus* complex. *Molecular Ecology* 22: 2941-2952.

Stebbins, G. L. 1980. Botany and the synthetic theory of evolution. p139-152. In: Mayr E

Provine WB, eds. *Evolutionary synthesis: perspectives on the unification of biology.*

Harvard University Press, Cambridge, MA, USA.

Telewski, F. W. 2021. Mechanosensing and plant growth regulators elicited during the

thigmomorphogenetic response. *Frontiers in Forests and Global Change* 3: 574096.

Totland, O., and Birks, H.J.B. 1996. Factors influencing inter-population variation in *Ranunculus*

acris seed production in an alpine area of southwestern Norway. *Ecography* 19: 269–278.

Turesson, G. 1922a. The species and the variety as ecological units. *Hereditas* 3: 100-113.

Turesson, G. 1922b. The genotypic response of the plant species to habitat. *Hereditas* 3: 211-

350.

Twyford, A. D., and J. Friedman. 2015. Adaptive divergence in the monkey flower *Mimulus*

guttatus is maintained by a chromosomal inversion. *Evolution* 69: 1476-1486.

Vickery, R. K. 1952. A study of the genetic relationships in a sample of the *Mimulus guttatus*

complex. PhD Dissertation, Stanford University, Stanford, CA, USA.

Vickery, R. K. 1978. Case studies in the evolution of species complexes in *Mimulus*.

Evolutionary Biology 11: 405–507.

Wu, C. A., D. B. Lowry, A. M. Cooley, K. M. Wright, Y. W. Lee, and J. H. Willis. 2008.

Mimulus is an emerging model system for the integration of ecological and genomic

studies. *Heredity* 100: 220-230.

TABLES

Table 1. ANOVA and linear regression summaries for *M. guttatus* traits in the main greenhouse. Asterisks denote significance at $p<0.05$ (*), $p<0.01$ (**), and $p<0.001$ (***).

Trait	Parameter Summaries						Full Model Summaries			
	Predictor	Estimate	Sum Sq.	df	F	P	Multiple R^2	df	F	P
Flowering Time (days)	Latitude	-0.0633	7.6	1,71	0.120	0.7301	0.0267	2,71	0.974	0.3825
	Exposure	2.3836	98.4	1,71	1.551	0.2171				
2nd Leaf Width (cm)	Latitude	-0.1633	50.56	1,71	9.661	0.0027**	0.1198	2,71	4.83	0.0108*
	Exposure	-0.3736	2.42	1,71	0.462	0.4990				
Corolla Width (mm)	Latitude	-0.2036	71.82	1,63	5.636	0.0207*	0.1221	2,63	4.383	0.0165*
	Exposure	1.0034	15.13	1,63	1.187	0.2800				
Corolla Length (mm)	Latitude	-0.2106	76.84	1,63	7.008	0.0102*	0.1713	2,63	6.511	0.0027**
	Exposure	1.4334	30.87	1,63	2.816	0.0983				
Height (cm)	Latitude	-0.8297	1305.94	1,71	39.017	2.738e-08***	0.4304	2,71	26.83	2.099e-09***
	Exposure	-7.0716	865.79	1,71	25.867	2.863e-06***				

Table 2. ANOVA, linear regression, and AIC summaries for models evaluating the relationship of selected principal components to *M. guttatus* height in the main greenhouse. Asterisks denote significance at p<0.5 (*), p<0.01 (**), and p<0.001 (***). Lowest AIC is bolded.

Principal Component	Parameter Summaries						Full Model Summaries				AIC
	Predictor	Estimate	Sum Sq.	df	F	P	Multiple R^2	df	F	P	
PC1	$(Dim.\ 1)^2$	13.923	1193.90	2,61	25.044	1.147e-08***	0.5725	3,61	27.23	2.679e-11***	**396.462**
	Exposure	-6.048	533.47	1,61	22.381	1.368e-05***					
PC2	Dim. 2	0.0520	0.94	1,62	0.022	0.883	0.2217	2,62	8.831	4.219e-04***	433.403
	Exposure	-7.0733	752.57	1,62	17.628	8.736e-05***					
PC3	Dim. 3	0.0663	0.71	1,62	0.017	0.898	0.2217	2,62	8.828	4.23e-4***	433.408
	Exposure	-7.0856	743.63	1,62	17.417	9.524e-05***					

Table 3. Regressions of climate factors that were highly-loaded on PC1 against latitude

Climate Factors	Estimate	F	Multiple R²	P
Annual Precipitation	0.0495	231	0.7857	< 2.2e-16
Coldest Quarter Precipitation	0.1116	136.4	0.684	< 2.2e-16
Driest Quarter Precipitation	0.6088	405.3	0.8655	< 2.2e-16
Mean Annual Temperature	-2.5849	350.2	0.8475	< 2.2e-16
Mean Wind Speed, 1st Quarter	2.4760	53.32	0.4584	5.94E-10
Mean Wind Speed, 4th Quarter	1.8632	64	0.5039	3.57E-11
Precipitation Seasonality	-0.3049	549.7	0.8972	< 2.2e-16
Warmest Quarter Precipitation	0.6092	483.8	0.8848	< 2.2e-16
Wettest Month Precipitation	0.3099	137.1	0.6852	< 2.2e-16
Wettest Quarter Precipitation	0.1054	159.6	0.7169	< 2.2e-16

Note: F-statistic degrees of freedom were all 1,63

FIGURES

610 **Figure 1.** Population locations for coastal perennial *M. guttatus* in a) Washington, British

611 Columbia, and Alaska; b) Oregon; c) Northern California; and d) Southern California. These

612 maps include populations grown in both the main and pilot studies.

Figure 2. Mean *M. guttatus* population height decreases significantly with latitude (*P* < 0.001),

and exposed populations have shorter average heights than protected populations (*P* < 0.001);

both population groups express the same relationship between height and latitude of origin.

Relationship of latitude of origin (in degrees) for *M. guttatus* populations and mean individual

heights (cm) of these populations was determined by linear regression.

Figure 3. Variation in plant height within the coastal perennial ecotype in *Mimulus guttatus.* A)

Divergence in height between plants from the Central California coast (left) and Oregon (right)

when grown in a greenhouse common garden experiment. B) Tall plants growing in a protected

area of the Monterey Bay along the Central California coast reach up to 2 meters in height. C)

Plants exposed to the open ocean in Oregon are much more compact in growth form.

Figure 4. Mean *M. guttatus* population height decreased significantly and non-linearly with

increasing values of climate PC1 in the main experiment ($P < 0.001$); exposed populations were

also significantly shorter than protected populations but maintained the same relationship to PC1

($P < 0.001$). The figure displays the quadratic relationship between climate PC1 and plant height,

which was the best fitting model as determined by AIC.

Figure 5. Mean wind speeds (m/s) along the United States Pacific Coast reported annually (A),

during January (B), April (C), and August (D). The dashed line indicates 42° N, the latitude of

the Oregon coast at which wind speeds start declining going further north.