Downregulation of a cell polarity protein potentiates Chikungunya Virus infection in host cells

Pushkar Tatiya[1#,] Ramesh Kumar[1$#,] Debajit Dey[1^], Yashika Ratra[2], Syed Yusuf Mian[2&,] Subhomoi Borkotoky[1], Bhawna[2] , Sanchi Arora[2], Kirti Suhag[1], Soumen Basak[2], Manidipa Banerjee[1*]

[1] Kusuma School of Biological Sciences, Indian Institute of Technology Delhi, Hauz Khas, New Delhi
[2] Signaling Systems Laboratory, National Institute of Immunology, Aruna Asaf Ali Marg, New Delhi
[$] Department of Chemistry and Biochemistry, University of Maryland, College Park
[^] Current address: Department of Biology, University of York
[&] Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of Public Health

# Equal contribution

*Correspondence: mbanerjee@bioschool.iitd.ac.in

Abstract

Viral manipulation of host cell morphology and behaviour represents an adaptive strategy that increases pathogen survival and replication efficiency. We establish a critical role for the downregulation of human Scribble, part of a cell polarity complex controlling morphology, apoptosis, and migration, for the potentiation of Chikungunya Virus (CHIKV) infection. A structural protein component (TF) of CHIKV targets Scribble, resulting in its punctation, ubiquitination, and subsequent degradation, probably through the proteasome pathway. The interaction between TF and Scribble is mediated *in vivo* by the PDZ domains of Scribble and a PDZ-domain binding motif (PBM) of TF, as shown by a combination of co-localization, co-immunoprecipitation, and proteomics analysis. Disruption of the Scribble polarity protein by shRNA mediated knockdown of Scribble potentiates CHIKV propagation. A mutated version of CHIKV generated through reverse genetics, which produces all structural proteins except TF, cannot affect Scribble to the same extent; and consequently displays reduced infectivity. Our work thus establishes a novel function for the TF component of alphaviruses in the modulation of host response potentiating virus transmission.

Keywords

6K;Chikungunya virus;PDZ domain;Scribble;TF

Introduction

Chikungunya Virus (CHIKV), a member of the alphavirus genus in the *Togaviridae* family, is a mosquito-borne human pathogen that affects almost 100 countries and causes ~3,000,000 infections per year globally (Wahid *et al.*, 2017). There are no targeted therapeutics or licensed preventive strategies for CHIKV and the treatment is primarily symptomatic (Bartholomeeusen *et al.*, 2023; Millsapps *et al.*, 2022). While the symptoms caused by this virus are self-limiting, there can be severe complications such as chronic, arthritis-like symptoms; and infections in the very young and elderly populations can be fatal. A clearer understanding of the functionalities of CHIKV encoded proteins, and identification of their host interaction partners, is crucial for the development of targeted therapeutics or attenuated vaccine candidates (Wong & Chu, 2018).

The CHIKV capsid, like that of all alphaviruses, is a 60-70 nm particle with T=4 icosahedral geometry (Sun *et al*, 2013; Kielian *et al.,*, 2010; Jose *et al.,*, 2009). The outer layer is composed of 240 heterodimers of envelope glycoproteins E1 and E2, embedded in a host-derived lipid envelope. While E2 is responsible for association with the host cellular receptor(s), E1 orchestrates the fusion of the viral and host lipid membranes (Wong & Chu, 2018). The inner layer of the particle consists of 240 copies of a multifunctional capsid protein (C) which encapsulates the genome (Chmielewski *et al.,*, 2022). Another envelope protein, E3, promotes the association between E1 and E2 during virus assembly (Yap *et al*, 2017).

The functionalities of the major structural proteins of CHIKV – E1, E2, E3, and C - are well investigated and established. However, a fifth protein – the small structural peptide "6K", remains relatively less explored (Wong & Chu, 2018). While 6K is categorized as a structural protein, its presence in the CHIKV capsid is yet unconfirmed and its role in the virus life cycle remains mysterious. It has been shown that recombinantly generated alphavirus 6K is an ion channel-forming protein (Dey *et al.*, 2019; Melton *et al.*, 2002). The 6K component from Ross River Virus (RRV) and Barmah Forest Virus (BFV) forms cation-specific channels in bilayer membranes in electrophysiology experiments (Melton *et al*, 2002). The expression of 6K in bacterial cells is detrimental to the growth of the cells, which is consistent with 6K possessing membrane-disrupting activity (Sanz *et al*, 1994; Melton *et al*, 2002). Bacterially produced CHIKV 6K, in conjunction with the expression partner Glutathione-S-transferase (GST), forms heterogeneous pores in the membrane *in vitro* (Dey *et al.*, 2019). *In vitro* liposome disruption experiments as well as cellular overexpression models have shown that CHIKV 6K tends to have

a strong affinity towards Endoplasmic Reticulum (ER) membranes and congregates in the ER.

CHIKV 6K, as well as Sindbis Virus (SV) 6K, co-localize with the glycoprotein E2. SV 6K co-

localizes with E2 in cytopathic vesicles on the way to the plasma membrane, which is the budding

site for alphaviruses (Dey *et al*, 2019; Brown *et al*, 2018; Elmasri *et al*, 2022). It has been

conjectured that 6K is required to facilitate concurrent transport of viral protein and genomic

complexes and promote association of glycoproteins during assembly, although there is no clear

connection between this functionality and the ion-channel activity of 6K at present.

The role of 6K in the alphavirus life cycle has been further complicated by the discovery of a

transframe variant "TF", which is generated due to ribosomal slippage within the 6K gene. TF is

produced at a 10-18 % frequency during viral protein translation (Firth *et al.*, 2008; Ramsey *et*

al., 2019) and shares the majority of the primary amino acid sequence with 6K but varies in the

C-terminal region. The production of TF results in premature termination of structural

polyprotein translation; thus, E1 is not generated along with TF (Figure 1A). Post-translational

modification, specifically palmitoylation, of the Sindbis Virus (SV) TF, is necessary for its

association with the plasma membrane (Ramsey *et al.*, 2017). Alphavirus capsids may contain

TF as a structural component, albeit in substoichiometric quantities (Snyder *et al.*, 2013; Firth *et*

al., 2008). The exact functionality of TF during alphavirus infections is not established; and it

remains a mysterious entity like 6K. The functionalities attributed to 6K in many previous works

may have been, to some limited extent, influenced by the coexistence of hybrid peptides

containing partial sections of TF, due to the non-removal of the ribosomal slippage site within

the 6K genomic sequence (Dey *et al.*, 2019;Ramsey & Mukhopadhyay, 2017).

Upon a close examination of the CHIKV TF amino acid sequence, we noted that the unique

region corresponding to the C-terminal region of TF contains a classical PDZ-domain binding

motif (PBM) (Figure 1B). The acronym PDZ originates from the first three proteins discovered

to contain this domain – the postsynaptic density protein 95 (PSD95), Drosophila disc large

tumour suppressor (Dlg1), and zonula occludens-1 protein (ZO-1) (Ye & Zhang, 2013; Barreda

et al., 2018). PDZ domains are protein-protein interaction domains found in practically all life

forms. *In silico* docking and simulation studies with the PBM at the C-terminus of TF suggested

that a potential binding partner for this motif is Scribble, part of a conserved set of proteins that

controls apico-basal as well as planar cell polarities and is also involved in the modulation of

tissue morphogenesis and apoptosis (Zhan *et al.*, 2008) (Thomas & Banks, 2018) (Figure 1B).

Previous studies have shown that Scribble is targeted by non-structural proteins from Influenza

Virus, Human T lymphotropic Virus-1 (HTLV-1), and Hepatitis C Virus (Javorsky *et al.*, 2023);

however, the involvement of Scribble in cellular infections by CHIKV or any other alphavirus is

so far unknown. In this study, we report that CHIKV infection results in more than 90%

abrogation of Scribble protein. Scribble is ubiquitinated and marked for degradation within 24

hours of CHIKV infection in HEK293T cells *in vitro*. We present evidence for the involvement

of TF in Scribble degradation, as transfection of TF or its production during CHIKV infection

results in the sequestration of Scribble into discrete cytosolic puncta. TF also colocalizes in these

puncta and can be immunoprecipitated with Scribble during CHIKV infection. The co-

localization as well as pulldown of TF with Scribble can be abrogated by mutations within the

PBM of TF, showing that the interaction between TF and Scribble is mediated by a PDZ-PBM

association. The knockdown of Scribble results in a significant enhancement in CHIKV

infectivity in cells. TF-mediated sequestration of Scribble accelerates virus propagation,

probably through negative regulation of cellular polarity and pro-apoptotic pathways controlled

by the Scribble complex. A version of CHIKV generated through reverse genetics, that does not

produce TF, has reduced ability to engage with Scribble and is inhibited in its propagation. Taken

together, our data attributes important functions in the contexts of host response modulation and

virus propagation to the enigmatic TF component of alphaviruses.

Results

The unique C-terminus of CHIKV TF contains a PDZ domain binding motif with potential for association with Scribble

The first 49 amino acids in the protein sequence of CHIKV 6K and TF are identical. This region is highly hydrophobic and capable of lipid membrane interaction as shown before (Dey *et al.*, 2019; Firth *et al.*, 2008). A ribosomal 212-1 frameshift at a slippery sequence corresponding to residues 48-49 within the 6K gene results in the production of TF with a unique C-terminal region of 27 amino acid residues which is primarily hydrophilic (Figure 1A). A motif search of this region indicated the presence of a putative PBM at the extreme C-terminus (Figure 1B). The Eukaryotic Linear Motif (ELM) resource (Kumar *et al.*, 2022) predicted that TF residues 71-76, with the sequence HGGSTV, constitute a class I PBM. This was confirmed by PDZPepInt, a PDZ domain identification software (Kundu *et al.*, 2014; Kundu & Backofen, 2014), which further predicted that residues 71-75 (HGGST) can constitute a potential binding motif for the host protein Scribble. Mammalian Scribble, a cell polarity component with four PDZ domains, is targeted by multiple viral proteins through PDZ-PBM interactions (Javorsky *et al.*, 2023). Particularly, domains 1 and 4 of Scribble have previously been reported to be high-affinity binders (Janezic *et al.*, 2019). We attempted to check the association of CHIKV TF with the PDZ binding domains of Scribble using *in silico* methods. A predicted structure of full-length TF (from the I-TASSER web server) was docked against PDZ domains 1, 2, 3, and 4 of Scribble (designated ScPDZ1, ScPDZ2, ScPDZ3, and ScPDZ4) using the ClusPro web server. The best-scoring models were selected for further analysis, and the binding interface was analyzed using Ligplot (Figure 1C, Supplementary Figure S1). MD simulations of TF in conjunction with ScPDZ1, ScPDZ2, ScPDZ3, and ScPDZ4 were subsequently carried out. In order to determine whether the interaction with the PBM of CHIKV TF is specific to the PDZ domains of Scribble, a PDZ domain from the mammalian High-temperature requirement A1 enzyme, a serine protease (HtrA1) (Wang & Nie, 2021) was also included in the study as a control. Several residues at the C-terminus of TF were found to be involved in mediating hydrogen bonds and hydrophobic interactions with residues of the PDZ domains 1 and 4 of Scribble (Table 1). An RMSF plot (Figure 1D) and RMSD (Supplementary Figure S2) analysis of the simulation trajectory showed that TF was stabilized by the Scribble PDZ domains. In particular, local fluctuations in the C-terminal tail of TF were significantly reduced upon interaction with ScPDZ1 (Figure 1D). These

fluctuations were most pronounced in simulations of TF alone. However, HtrA1 PDZ interaction had minimal effect on the TF C-tail fluctuations (Figure 1D). Higher binding energies were observed for the interaction of TF with the PDZ domains of Scribble in contrast to that of HtrA1 (ScPDZ1>ScPDZ4>>HtPDZ) (Supplementary Table S1). Specifically, TF-HtPDZ interaction was observed to be ~3.8 times weaker than that with ScPDZ4 and ~5.8 times weaker as compared to TF-ScPDZ1. The binding energies in all cases were primarily mediated by electrostatic and van der Waals interactions (Figure 1C, Supplementary Table S1). Thus, *in silico* analysis indicated that CHIKV TF preferentially binds to the PDZ domains 1 and 4 of Scribble, as compared to PDZ domains from other mammalian proteins. The association of TF with ScPDZ2 and ScPDZ3 was also analyzed (Supplementary Figure S1), and the binding energies were found to be at least 2-fold lower than that of TF-ScPDZ1 and 1.4 times lower than that of TF-ScPDZ4 (Supplementary Table S1). Further analysis of the TF interface residues with ScPDZ1 (Figure 1E) revealed that several C-terminal residues of TF, which are part of the putative PDZ-domain binding motif, contributed favourably towards binding with ScPDZ1.

Overexpressed CHIKV TF co-localizes with Scribble

In order to investigate interactions between TF and Scribble biochemically, a plasmid containing the cDNA of CHIKV TF, with an N-terminal Enhanced Green Fluorescent Protein (EGFP) tag (EGFP-TF), was transfected in HEK293T cells. To check for the specificity of the putative interaction, the cDNA of CHIKV 6K, which shares the N-terminal forty-nine residues with TF, was similarly fused with EGFP to form EGFP-6K and transfected. The ribosomal slippage site was mutated in the 6K construct to prevent the initiation of TF expression from this plasmid (Supplementary Table S2). Twenty-four hours post-transfection, cells were fixed and immuno-stained with a Scribble-specific antibody to establish the intracellular localization of Scribble in the presence of CHIKV TF or 6K (Figure 2A). In mock-transfected cells, Scribble was found to localize primarily to the cellular periphery near the plasma membrane. Similarly, Scribble localized to the plasma membrane in cells transfected with an empty vector (pEGFPC1), and no co-localization with EGFP was noted (Figure 2A). However, in cells transfected with EGFP-TF, the localization of Scribble was drastically altered. In these cells, EGFP-TF and Scribble were found to co-localize in the form of distinct puncta in the cytoplasm (Figure 2A). No such co-localization was noted upon transfection of cells with EGFP-6K. In order to establish co-

localization in a semi-quantitative fashion, we utilized intensity line profile analysis. The profile

graph displays the fluorescence signal intensity along the y-axis and the position or distance along

the x-axis for two fluorochromes. The overlapping intensity peaks clearly suggested that while

there is limited co-localization between 6K and Scribble, the co-localization between TF and

Scribble is substantial (Figure 2A and Supplementary Figure S3). We hypothesized that the

prominent punctation and mis-localization of Scribble is probably mediated by the unique C-

terminal region of TF, since the N-terminal regions of 6K and TF are identical (Figure 1A).

Given the conclusions of the *in-silico* studies, we further hypothesized that the association

between Scribble and TF is likely mediated by the PDZ domain(s) and PDZ-binding motifs of

the components.

The interaction between TF and Scribble is mediated by a PDZ-PBM interaction

In order to investigate the association between TF and Scribble, the putative PBM in CHIKV TF

(71-HGGSTV-76) was mutated to contain either two or four alanine substitutions

(HGAATV/AAAATV). The constructs were designated TF Mut1 and TF Mut 2, respectively.

The corresponding plasmids with the mutated EGFP-TF sequences were individually transfected

in HEK293T cells, and the effect of these mutants on Scribble localization was analyzed using

confocal microscopy.

We did not observe puncta formation by Scribble in cells transfected with the mutated EGFP-TF

constructs (Figure 2B). The fluorescence associated with TF Mut1 and TF Mut2 had a strikingly

different distribution compared to wildtype TF. No co-localization was detected between

Scribble and PBM mutated TF constructs visually and upon semi-quantification via line profile

analysis (Figure 2B and Supplementary Figure S3). To confirm that the TF PBM mutants were

being expressed in cells, the expression level of wildtype and mutated EGFP-TF constructs were

investigated via Western blots (Figure 2C). A slight reduction in the protein expression level of

EGFP-TF with the PBM mutated to HGAATV (TF Mut1) compared to wildtype was observed,

while the EGFP-TF construct with a quadruple alanine mutation (TF Mut2) displayed further

reduced expression. However, the molecular weight difference between EGFP only, EGFP-6K

and the EGFP-TF constructs clearly indicated the expression of the correct proteins in transfected

cells (Figure 2C). Further, the alteration in the cellular expression pattern of the mutated

constructs and the lack of co-localization with Scribble strongly suggested that the putative PDZ

binding motif HGGSTV is critical for the association between TF and Scribble (Figure 2A-C).

To probe this interaction further, we immunoprecipitated Scribble with a rabbit polyclonal

antibody from HEK293T cells transfected with EGFP only, EGFP-TF, EGFP-TFMut1 and

EGFP-TFMut2. The immunoprecipitated fractions were probed with an anti-Scribble antibody

and an anti-GFP antibody to identify TF and TF mutants (Figure 2D). We found that while EGFP-

TF could be immunoprecipitated with an anti-Scribble antibody; EGFP-TFMut1 or EGFP-

TFMut2 could not be pulled down along with Scribble. This strongly indicates that the

association between Scribble and TF is mediated via the PDZ domains of the former and the

PBM of the latter, since mutating the PBM abrogated the association.

To establish the differences in Scribble binding by wildtype vs PBM mutated versions of TF,

LC-MS/MS analysis of trypsin-digested fragments from the immunoprecipitated fractions was

carried out. While wildtype EGFP-TF had a substantial coverage in the fractions

immunoprecipitated by an anti-Scribble antibody, EGFP-TF Mut1 and EGFP-TF Mut2 had

significantly reduced coverage and fewer peptides, less than half of that observed for the wildtype

EGFP-TF (Table 2), suggesting considerably lower association of the mutated versions of TF

with Scribble. A list of all the proteins immunoprecipitated is shown in the supplementary file

(MS ANALYSIS). Taken together, co-localization studies, immunoprecipitation, and mass

spectrometry clearly establish that the *in vivo* association between Scribble and CHIKV TF is

driven by the interaction between the PDZ domain(s) of the former and the PBM at the unique

C-terminus of TF.

TF associates with Scribble during CHIKV infection: In order to determine if the association

of TF with Scribble has any functional relevance, the interaction was investigated during CHIKV

infection of HEK293T cells. An antibody was raised against the unique region of CHIKV TF,

by utilizing the C-terminal 27 amino acids coupled with Keyhole Limpet Hemocyanin (KLH) as

an antigen. The antibody, generated in rabbit, was tested for its ability to recognize TF in western

blots as well as in immunofluorescence assays (Supplementary Figure S4). The association of

the antibody with TF was found to be specific as it was able to recognize bacterially produced

and purified TF (conjugated with the Maltose Binding Protein, MBP), as well as EGFP-TF from

a HEK293T cell lysate, but was unable to bind to bacterially produced MBP-6K or overexpressed

EGFP-6K (Supplementary Figure S4).

The antibody against TF was utilized to visualize the localization and association of TF during

CHIKV infection. Confocal microscopy studies showed substantial production of TF during

CHIKV infection of HEK293T cells and significant TF co-localization with Scribble in punctate

forms (Figure 3A and Supplementary Figure S5). The co-localization was also validated by

intensity line profile analysis (Figure 3A). For comparison, CHIKV capsid protein and

glycoprotein E2 were also immunostained using specific antibodies (Figure 3B); however, there

was no co-localization of Scribble with either capsid protein C or E2. To understand the dynamics

of viral protein production and co-localization, CHIKV-infected cells were probed

simultaneously with E2 and Scribble antibodies or E2 and TF antibodies at 24 hours post-

infection. Interestingly, while E2 did not co-localize with Scribble even after 24 hours of

infection, substantial co-localization was noted between E2 and TF (Figure 3C).

The association between Scribble and TF during CHIKV infections was further established by

immunoprecipitation of Scribble using a polyclonal anti-Scribble antibody from CHIKV-infected

HEK293T cell lysates. Western blotting of the co-IP fractions with the TF unique-region specific

antibody clearly indicated the presence of TF (Figure 3D), thus establishing the association of

TF with Scribble during CHIKV infection.

CHIKV infection, or TF expression, results in degradation of Scribble

Multiple viruses have been shown to target Scribble for degradation in order to manipulate host

immune responses, cell junction integrity, and apoptosis pathways (Javorsky *et al.*, 2023). In

order to analyze the effect of CHIKV infection on the level of Scribble, the levels of Scribble

protein were detected by western blotting in HEK293T cells 24 and 48 h post infection (Figure

4A). A significant decrease in Scribble protein level was noted upon CHIKV infection in a time-

dependent manner. At 24 hours post-infection, Scribble levels were reduced to nearly 50%

compared to uninfected cells. The protein levels reduced to greater than 95% at 48 hpi (Figure

4A), while those in uninfected cells remained at the initial level. Thus, CHIKV infection appeared

to cause a drastic reduction in the levels of human Scribble protein in cell culture.

To confirm that Scribble degradation during CHIKV infection is triggered by the TF component,

the effect of overexpressed TF on Scribble protein expression level was also evaluated (Figure

4B). Since CHIKV-induced reduction in Scribble protein levels was the highest 48 h post

infection, the effect of TF, upon transfection, on Scribble expression was also studied at 48 h.

Transfection of HEK293T cells with the EGFP-TF plasmid, containing the wildtype sequence of TF, was found to reduce the level of Scribble protein by ~20%. The higher levels of Scribble degradation in CHIKV-infected versus TF-transfected cells is probably due to the different levels of TF expression in the infection versus the transfection model. The reduction in Scribble protein level was not elicited by transfection of EGFP-6K, or EGFP-TF Mut1 and EGFP-TF Mut2, (Figure 4B), indicating that this effect is triggered by TF, and is specific to the PBM at the C-terminus of TF.

The mode of degradation of Scribble was explored by co-staining CHIKV-infected cells with anti-Scribble and anti-Ubiquitin primary antibodies and appropriate secondary antibodies (Figure 4C). We found strong co-localization of the Scribble puncta with ubiquitin-specific antibodies in CHIKV-infected HEK293T cells, which suggests that CHIKV-mediated degradation of Scribble could be mediated by the Ubiquitin-Proteasome System (UPS). Similarly, in transfected cells overexpressing TF, co-localization of Scribble, Ubiquitin, and TF was detected in cytoplasmic puncta (Figure 5A). While some Scribble punctation, albeit at a lower level, was noted during overexpression of the double alanine PBM mutant (EGFP-TF-HGAATV), there was no co-localization of ubiquitin to these puncta (Figure 5A).

To further investigate the involvement of the UPS in Scribble degradation, ubiquitinated proteins in CHIKV infected or uninfected HEK293T cells were analyzed in the presence of the proteasome inhibitor MG-132 (Figure 5B). In uninfected cells, the addition of MG-132 had no effect on Scribble quantities. However, the addition of MG-132 restored the reduction in Scribble triggered by CHIKV infection.

Thus, taken together, our results suggest that during CHIKV infection, the TF component associates with Scribble through its C-terminal PBM (HGGSTV), resulting in the punctation and degradation of Scribble. The alterations in Scribble quantities upon the addition of MG132 suggest that the degradation of Scribble probably occurs through the UPS pathway.

Scribble knockdown enhances CHIKV infection

In order to investigate the significance of Scribble downregulation during CHIKV infection, the level of CHIKV propagation in Scribble knockdown cells was compared to that in untreated cells. Two short hairpin RNAs (shRNAs) were used for the knockdown of Scribble protein expression in HEK293T cells according to standard protocols. The effect of the shRNAs was analyzed 36

hours post-transfection using western blot analysis. Densitometric analysis, to quantitate the relative percentage of gene knockdown for each sample, indicated a relatively efficient Scribble protein knockdown of approximately 50% with both shRNA conditions (Figure 6A) . The degree of knockdown at the RNA level was also confirmed using quantitative real-time polymerase chain reaction (qRT-PCR) (Figure 6B). To ensure that any alteration observed in CHIKV infectivity upon Scribble knockdown was indeed because of a reduction in the quantity of Scribble, and not due to a difference in the number of viable cells as an effect of knockdown, cell viability was assessed using an MTT assay (Figure 6C). No significant change in cell viability was observed with Scribble knockdown.

The HEK293T cells with reduced Scribble expression were then infected with CHIKV (strain 119067) at an MOI of 0.1. The culture supernatants containing the new virus were harvested 24 h post infection and plaque assay was performed using standard methodologies in Vero cells. CHIKV infectivity was calculated by counting the number of plaques formed 48 h post-virus addition. We found that the virus titre, represented as the plaque forming unit per ml (pfu/ml), was significantly increased upon Scribble knockdown (Figure 6D). We also tested the levels of CHIKV RNA in infected HEK293T cells with Scribble knockdown by RT-PCR and found that the levels of CHIKV RNA were also markedly upregulated in these cells, compared to regular infected cells (Figure 6E). The increase in infectious CHIKV virions and genomic RNA was approximately 2-3 times higher upon Scribble knockdown. Taken together, our data strongly suggest that punctation and degradation of Scribble is a viral countermeasure to facilitate CHIKV propagation and is mediated by TF.

Infection with ΔTF CHIKV does not trigger Scribble punctation

To further investigate the role of TF during CHIKV infection, a mutated version of CHIKV was generated to produce all structural proteins including 6K, but not TF. An infectious clone of the CHIKV OPY-1 strain with an mCherry cassette for detection (mCherry-LR2006-OPY1) (Hallengärd et al, 2014) was utilized as a template to generate a silent mutation in the ribosomal slippage site, to abrogate the production of TF. Both infectious clones - WT CHIKV and ΔTF CHIKV - were transfected in HEK293T cells using the standard protocol for the formation of virus particles. The transfection supernatants were subjected to plaque assays and the viral plaques in each case were isolated and subjected to serial infections to increase the titre. We

confirmed, using specific antibodies, that while infection with WT CHIKV results in the production of TF as well as other structural proteins; infection with ΔTF CHIKV produces 6K, but not TF (Supplementary Figure. S9).

ΔTF CHIKV at 1 MOI was utilized to infect HEK293T cells, and the WT virus at the same MOI was utilized in control infections. The plaque assay consistently suggested lower cellular infectivity for ΔTF CHIKV compared to WT virus. qRT-PCR based quantification of viral RNA for the WT and ΔTF CHIKV virus showed higher viral RNA load in cells infected with ΔTF CHIKV 12 hpi(Figure 7A). This indicates production of significantly higher numbers of progeny virions that are release/assembly defective when TF is absent. The observation re-emphasizes the critical nature of cell polarity/adhesion disruption mediated by ubiquitination of scribble for CHIKV transmission in vitro. Cells infected with WT or ΔTF CHIKV were co-stained with anti-Scribble and anti-Ubiquitin antibodies to detect the extent of Scribble punctation and ubiquitin co-localization with Scribble (Figure 7B). Scribble appeared prominently in punctate form 24 hours post-infection in WT CHIKV-infected cells. Significant co-localization of Scribble and ubiquitin-specific antibodies was observed in WT CHIKV-infected HEK293T cells 12 hours post-infection, intensifying at 24 hours (Figure 7B). In contrast, while ΔTF CHIKV-infected cells did have limited amount of Scribble puncta, these did not co-localize with Ubiquitin (Figure 7B). Since Scribble and Ubiquitin co-localization was observed in WT mCherry-LR2006-OPY1 CHIKV 24 hours post-infection, cellular lysates from WT and ΔTF CHIKV OPY-1 infected cells were subjected to western blotting with anti-Scribble antibody at 24 hours p.i and 12 hours post this time point (Figure 7C). A significant, time-dependent, decrease in Scribble protein levels was observed in cells infected with WT CHIKV OPY-1 strain. In contrast, cells infected with ΔTF CHIKV OPY-1 showed Scribble levels slightly above normal (Figure 7C). Together, these results demonstrate that the TF component of CHIKV is crucial for effective ubiquitination and downregulation of Scribble, a process that cannot be executed by ΔTF CHIKV. Thus, the TF component of CHIKV is crucial for downregulating Scribble in infected cells, which enhances viral infectivity and propagation.

Discussion

Host-virus interactions play a crucial role in defining the trajectory of pathogenesis (Krishnakumar & Kannan, 2020). Key host molecules and pathways are consistently hijacked by viruses to ensure the propagation of infection. Some widely targeted host cellular co-factors include nuclear pore/envelope complexes, COPI vesicles, ribosomes and the translational pathway and signalling modulators (Ma-Lauer *et al.*, 2012). Increasing evidence suggests that the Scribble cell polarity complex is one of the central host factors targeted by multiple virus families (Javorsky *et al.*, 2023). Human Scribble is part of a large module of adapters, including several Dlg and Lgl proteins, which has a central role in regulating cell migration, adhesion, polarity, signalling, apoptosis, morphogenesis, and development (Bonello & Peifer, 2019; Santoni *et al.*, 2020; Stephens *et al.*, 2018). The members of this complex associate with a wide range of cellular factors to modulate cellular activities. Scribble is a member of the LAP (leucine-rich repeats and PDZ domain) family and contains numerous protein-protein interaction domains, including 16 Leucine-Rich Repeat domains (LRRs), 2 LAP-specific domains, and 4 PDZ domains (PDZ1-4). Loss of Scribble causes disruption of cellular adhesion and cell polarity, resulting in hyperproliferation (Bonello & Peifer, 2019). As a determinant of cellular growth, apoptosis and proliferation, Scribble has been a target of tumorigenic viruses such as Human Papillomavirus (HPV) and Human T-lymphotropic virus 1 (HTLV-1) (Arpin-André & Mesnard, 2007; Nakagawa & Huibregtse, 2000). Recently, effectors from several other viruses, such as the NS1 protein from Influenza A Virus (IAV), NS5 from Tick-borne encephalitis virus (TBEV), and the core protein from Hepatitis C Virus (HCV), have been shown to associate with the PDZ domains of Scribble through PBMs, resulting in mistargeting of Scribble and an increase in viral proliferation (Javorsky *et al.*, 2023). However, there have been no reports of Scribble modulation by alphaviruses so far. In this manuscript, we establish Scribble as a host target of CHIKV and identify the viral factor that induces the mislocalization and degradation of Scribble to enhance viral proliferation.

We found that CHIKV infection downregulates Scribble expression by as much as 95% in cultured cells in a time-dependent manner. Immunofluorescence analysis of CHIKV-infected cells showed clustering of Scribble into cytoplasmic puncta, and ubiquitin-mediated degradation. Degradation of Scribble via the Ubiquitin-Proteasome pathway was suggested by experimental

studies using a proteasome inhibitor, MG132. The cellular localization of Scribble is important for its functioning (Javorsky *et al.*, 2023; Zhan *et al.*, 2008; Stephens *et al.*, 2018), and mislocalization of Scribble and its sequestration in discrete puncta likely affect its regulatory role in cellular proliferation (Arpin-André & Mesnard, 2007; Liu *et al.*, 2010). To confirm the role of Scribble in CHIKV infection, shRNA-based downregulation of Scribble expression was carried out. A ~50% downregulation in Scribble levels resulted in 2-3 times increase in the virus titer and genomic RNA levels. This effect may be due to the elimination of the pro-apoptotic activity of Scribble, resulting in prolonged cell survival and consequent enhancement in viral replication (Liu *et al.*, 2010). Scribble is a cell polarity protein that is important for maintaining the integrity of cellular junctions, and disruption of cell junctions has been reported to promote virus egress and dissemination (Javorsky *et al.*, 2023; Bonello & Peifer, 2019; Torres-Flores & Arias, 2015; Thomas & Banks, 2018). While HEK293T cells do not contain conventional tight junctions, they do produce a number of cell adhesion molecules such as E-cadherin, occluding and ZO-1, in addition to Scribble (Inada *et al.,* 2016) Thus, the beneficial effect of Scribble knockdown on CHIKV replication might be due to alteration in cellular contacts, allowing enhancement of viral release. A more detailed analysis of the timeline of CHIKV-mediated Scribble downregulation and consequent spatio-temporal effects on cellular morphology, adhesion and signalling may be essential for understanding the functional basis for viral countermeasures to host response.

The small structural protein 6K, and its transframe variant TF, appear to have significant importance in the regulation of CHIKV morphology and virus-host interactions (Ramsey & Mukhopadhyay, 2017). It has been shown previously that CHIKV 6K, like its homologs from other alphaviruses, is an ion-channel forming protein (Dey *et al.*, 2019). It has been shown that inhibition of 6K activity may affect CHIKV assembly, resulting in the formation of morphologically unusual particles with reduced infectivity. Recent work has shown that 6K co-localizes with alphavirus glycoproteins in cytopathic vesicles (CPV-II), which may support its role in promoting viral assembly (Elmasri *et al.*, 2022). TF, in contrast, localizes at the plasma membrane and is packaged in virions (Firth *et al.*, 2008; Ramsey *et al.*, 2017). Our work attributes an essential role to TF in the modulation of host response and in the enhancement of viral proliferation. Several lines of evidence show that the TF component of CHIKV is primarily responsible for the downregulation of Scribble. The expression of TF in HEK293T cells by

transfection leads to the sequestration of Scribble into distinct puncta, ubiquitination, and
degradation, similar to that noted in CHIKV infected cells. It is possible that low tolerance for
TF in infected or transfected cells results in its sequestration in puncta and ubiquitination; while
Scribble is co-localized into the puncta due to its affinity for TF and is degraded simultaneously.
Sequestration and ubiquitination of Scribble are not evident in cells transfected with CHIKV 6K,
suggesting that TF, and not 6K, is the regulator of the human Scribble protein. TF can be
immunoprecipitated with an anti-Scribble antibody from cells infected with WT CHIKV,
indicating an association between the two components. Further, a ΔTF version of CHIKV OPY-
1 strain generated through reverse genetics, which produces 6K but not TF, cannot trigger
Scribble punctation or ubiquitination at the same levels during cellular infection. Infection of
cells with the ΔTF variant resulted in substantially enhanced viral RNA load, coupled with
reduced infectious particles in the supernatant, indicating that the lack of TF indeed causes
disruption in the release and propagation of CHIKV. The punctation and degradation of Scribble
by TF is likely essential for modulation of cellular polarity resulting in propagation of the virus.
Since TF and 6K share approximately two-thirds of their amino acid sequence, this functional
difference is likely to originate from the unique C-terminal region of TF, which contains the PDZ
domain binding motif (PBM) as a prominent feature.

Other viral proteins such as IAV NS1, TBEV NS5, HTLV tax, HCV NS4B and HPV-16 E6 are
known to associate with the PDZ domains of Scribble through their PBMs (Javorsky *et al.*, 2023).
In silico analysis of the unique C-terminus of TF indicated the presence of a Class I PBM which
could interact with the PDZ domains of Scribble. Mutational analysis of the predicted PBM
confirmed its involvement in regulating Scribble localization. The TF constructs with the PBM
mutated were mis-localized, as they were distributed throughout the cytoplasm, and were unable
to co-localize with or trigger the sequestration and ubiquitination of Scribble. Mutations in TF
PBM also resulted in somewhat decreased protein levels, particularly in the quadruple alanine
mutant, which could be due to the drastic nature of the mutation. However, the expressed protein
clearly showed an altered localization pattern distinct from wildtype TF. Further, while GFP-TF
with an intact PBM could be immunoprecipitated from transfected cells, TF constructs with
mutated PBMs could not be pulled down with an anti-Scribble antibody. MS analysis confirmed

the weakening of Scribble-TF association upon mutating the TF PBM, which suggests a central

role for the PDZ-PBM interaction in this association.

While transfection of WT TF in HEK293T cells caused mislocalization and ubiquitination of

Scribble; the reduction in Scribble protein quantity was marginal compared to the drastic

decrease observed during CHIKV infection. This could be because of temporal alterations in TF

expression in the transfection versus the infection model, or due to the differences in the quantity

of TF production in these cases. The onset of protein detection is usually delayed in transfected

compared to infected cells, and it is possible that in HEK293T cells expressing TF,

mislocalization and punctation of Scribble precede protein degradation, which occurs at later

time points. Also, we found that despite the high transfection efficiency of HEK293T cells, as

seen with empty vector producing EGFP, the expression efficiency of EGFP-TF or EGFP-6K by

transfection is relatively low, which could be due to the membrane permeabilization capability

of these components. It has been shown earlier that expression of 6K in bacterial cells impedes

the growth curve significantly, as the membrane activity is likely toxic to the cells (Melton *et al.*,

2002; Sanz *et al.*, 1994). Thus, reduced levels of TF expression in transfected HEK293T cells

may be a possible reason for decreased levels of Scribble degradation, compared to those

observed in CHIKV-infected cells. However, the decline in Scribble protein levels triggered by

transfection of EGFP-TF was absent in the case of PBM-mutated TF constructs, which suggests

the crucial role of TF PBM in PDZ-domain dependent association with Scribble. TF may have

other roles in other aspects of the alphavirus life cycle, such as assembly and/or budding.

However, detailed structural and functional analysis of ΔTF CHIKV is needed to understand the

contributions of TF to the assembly and budding processes of alphaviruses at the molecular level.

The PDZ domain-PBM interaction is one of the most abundant modes of protein-protein

interaction. The homologues of CHIKV TF from several other alphaviruses also have PBMs in

their unique C-terminal tails that can potentially associate with host partners (Supplementary

Figure S7). For example, the predicted TF sequences from O'nyong nyong Virus and Igbo-Ora

Virus have Class I PBMs at the C-termini that are almost identical to the PBM in CHIKV TF;

while TF sequences from Ross River Virus, Mayaro Virus and Semliki Forest Virus are predicted

to contain Class II PBMs (Table 3). Two prominent PBMs, in addition to the one at the C-

terminus of TF, are predicted in the CHIKV proteome (Table 4). Residues 418-423 in CHIKV

glycoprotein E2 are expected to constitute a Class I PBM, while the C-terminus of 6K is predicted

to have a Class II PBM. However, we did not note any co-localization of transiently expressed

6K with Scribble, therefore it is possible that this PBM may mediate interactions with other host

or viral proteins.

Human Scribble has four PDZ domains, and their association with several viral proteins has been

monitored using biophysical methods such as Isothermal Titration Calorimetry (ITC) and X-ray

Crystallography (Javorsky *et al.*, 2022a, 2022b). *In vitro*, IAV NS1 binds to all four PDZ domains

of Scribble, with PDZ4 being the tightest binder and PDZ1 being the weakest (Javorsky *et al.*,

2022b). TBEV NS5 shows strong binding to both PDZ4 and PDZ3 of Scribble (Werme *et al.*,

2008; Javorsky *et al.*, 2022a). Interestingly, the NS4B protein of HCV requires interaction with

multiple PDZ domains of Scribble for strong binding. Our study establishes that the association

between TF and Scribble is driven by PDZ-PBM interaction. Our *in silico* studies suggest that

PDZ1 of Scribble has the highest affinity for the PBM in TF, followed by PDZ4. Therefore,

either PDZ1 or both PDZ1 and PDZ4 of Scribble may be involved in the association of Scribble

with TF. However, existing studies clearly indicate that the structural determinants of PDZ-PBM

binding that mediate virus-host interactions are complex and vary significantly even among

similar binding pairs. Based on existing studies, it is possible that multiple PDZ motifs from

Scribble, or perhaps all four motifs, are required for optimal TF association. (Hu *et al.*, 2016). A

detailed *in vitro* biophysical and structural analysis of the CHIKV TF protein with full-length

Scribble, as well as the isolated PDZ domains, is essential in order to understand the binding

modalities, including minimal requirements for functional association and cooperativity between

different PDZ domains. It is to be noted that other host proteins that are part of the Scribble

complex, such as Dlg-1, also contain PDZ domains (Barreda *et al.*, 2020). The potential

association of alphavirus TF with other components in the Scribble cell polarity module also

requires further study. This will lead to a clearer understanding of host responses and viral

countermeasures during infection.

Materials and methods

Docking and Molecular Dynamics (MD) simulation

Since there is no experimental structure available for CHIKV TF, a structure predicted using the I-TASSER web server (Yang & Zhang, 2015; Zhang, 2008) was utilized for docking. The model with the highest c-score was selected and further refined using the GalaxyRefine web server (Seok *et al.*, 2021; Shin *et al.*, 2014; Ko *et al.*, 2012). The X-ray crystallographic structures of Scribble PDZ domains (1-4, referred to as ScPDZ1, ScPDZ2, ScPDZ3, and ScPDZ4) and mammalian HTRA1 PDZ (HtPDZ) domain were downloaded from the Protein Data Bank (PDB IDs: 5VWC, 1UJU, 7JO7, 4WYT and 2JOA respectively). All structures were energy minimized prior to docking and simulation studies.

Protein-protein docking between TF and the PDZ domains of Scribble and HTRA1 was carried out using the ClusPro web server (Desta *et al.*, 2020; Vajda *et al.*, 2017; Kozakov *et al.*, 2017, 2013). The binding interface and affinity of each docked complex were analyzed using LigPlot (Wallace *et al.*, 1995). All-atom MD simulation was carried out for each complex in triplicates as described. Briefly, GROMACS (Van Der Spoel *et al.*, 2005) was used to set up and execute the simulations. Each complex was solvated separately using the SPC water model (Mark & Nilsson, 2001) and neutralized using counter-ions. Periodic boundary condition was applied to each system. Energy minimization was executed using the steepest descent algorithm (Gautam, 2021). Position-restrained NVT-NPT ensemble equilibrations were carried out for 500 ps and 1 ns respectively. Finally, the systems were subjected to unrestrained dynamics for 100 ns. Nose-Hoover (Martyna *et al.*, 1992) and Parrinello-Rehmann (Parrinello & Rahman, 1981) were used for maintaining the temperature (300K) and pressure (1 bar) during simulations. Electrostatics was maintained using Reaction-Field (Tironi *et al.*, 1995) and LINCS (Hess *et al.*, 1997) was used as the constraints algorithm. A time-step of 2 femtoseconds and the Verlet cutoff scheme was applied. All simulations were carried out using the Gromos 54a7 force field. Trajectory analysis was carried out using the built-in GROMACS tools. VMD (Humphrey *et al.*, 1996) and UCSF Chimera (Pettersen *et al.*, 2004) were used as the visualization tools.

Molecular Mechanics Poisson-Boltzmann Surface Area (MM-PBSA) analysis

MM-PBSA was utilized to determine the Gibb's free energy (GFE) of protein-protein interactions. The script g_mmpbsa (Kumari *et al.*, 2014) was used for evaluating the GFE of TF

and PDZ domains from simulation trajectories. The following equations were used for the

analysis

$\Delta G\ bind = \Delta G\ complex - (\Delta G\ protein1 + \Delta G\ protein2)$

$\Delta GX = \Delta EMM + \Delta G\ solv - T\Delta SMM$

$\Delta EMM = \Delta E\ bonded + \Delta E\ nonbonded = \Delta E\ bonded + (\Delta E\ elec + \Delta E\ vdw)$

$\Delta G\ solv = \Delta G\ polar + \Delta G\ nonpolar$

where $\Delta G\ complex$ = total free energy of protein-protein complex

$\Delta G\ protein1$, $\Delta G\ protein2$ = free energies of proteins in solvent

ΔGX = free energy for each entity

$\Delta E\ MM$ = vacuum potential energy (combined bonded and nonbonded interactions)

$\Delta G\ solv$ = solvation free energy [polar ($\Delta G\ polar$), nonpolar ($\Delta G\ nonpolar$)]

$T\Delta S$ = entropic contribution of free energy in vacuum. T is temperature and ΔS is entropy

respectively.

$T\Delta S$ MM was not included in this study as g_mmpbsa does not calculate entropy (S). Hence, ΔG

bind was calculated as follows,

$\Delta G\ bind = (\Delta E\ elec + \Delta E\ vdw) + (\Delta G\ polar + \Delta G\ SASA)$

where, $\Delta G\ SASA = \Delta G\ nonpolar$ since nonpolar solvation energy was calculated based on the

SASA model in this study.

Cells and plasmids

Human embryonic kidney (HEK) 293T and Vero cells were grown in Dulbecco's modified Eagle

medium (DMEM) (Gibco) supplemented with 10 % fetal bovine serum (FBS) (Gibco) and 1 %

antibiotic antimycotic solution (Himedia). The cells were maintained at 37 °C with 5 % CO_2. The

cDNA sequences of CHIKV 6K and TF (Supplementary Figure S8) were cloned in the pEGFPC1

vector (Clontech) to generate fusion proteins tagged with the Enhanced Green Fluorescent

Protein (EGFP) at the N-terminus. The ribosomal slippery site UUUUUUA in the 6K cDNA

sequence was silently mutated to UUUCCTG to disrupt the consensus ribosomal frameshifting

sequence and produce only 6K without affecting its primary structure (Firth *et al.*, 2008). Alanine

substitution mutations were carried out to disrupt the PDZ domain binding motif within amino

acids 71-74 of TF. Two mutants – a) G73A S74A and b) H71A G72A G73A S74A, were

generated using sequential site-directed mutagenesis (Life Technologies) and were confirmed by

DNA sequencing (Supplementary Table S2). The ribosomal slippery was silently mutated to UUUCCTG in a similar way using site-directed mutagenesis and HPLC purified primers (Supplementary Table S2) to produce only 6K CHIKV - ΔTF CHIKV construct.

CHIKV infection

For all CHIKV infections, HEK293T cells were seeded 24 h prior to infection by CHIKV (strain 119067) (Singh *et al.*, 2018) at a MOI of 0.1. CHIKV was added to the cells post phosphate-buffered saline (PBS) washing for adsorption. After 90 min, cells were again washed with PBS, and complete media was added. Post-infection, cells were incubated for 24 - 48 h in normal cell growth conditions.

Mutated CHIKV generation: Vero E6 cells were seeded in a 6-well plate to reach 60% confluency. Cells were then transfected with 5µg of CHIKV plasmid (WT CHIKV), and the mutant generated (ΔTF CHIKV) with polyethyleneimine(PEI) in a ratio of 1:5. The cell supernatant was collected 48 hours post-transfection, centrifuged at 2000 rpm for 10 minutes at 4°C. The supernatant was carefully removed, aliquoted, and stored at -80°C. For virus propagation, Vero E6 cells were cultured up to 80% confluency in DMEM supplemented with 10% FBS. Before infection, cells were incubated in incomplete media for 2-3 hours. Thereafter, cells were infected with 0.01 MOI of CHIKV and observed for maximum cytopathic effect (40-48 hours post-infection). The supernatant was collected and centrifuged at 2,000 for 10 minutes at 4°C to remove cell debris. The virus supernatant was carefully separated without disturbing the cell pellet and stored at -80°C.

Western blotting

HEK293T cells infected with CHIKV were washed with PBS and lysed with RIPA lysis buffer supplemented with a protease inhibitor cocktail (ThermoFisher Scientific) as per the manufacturer's protocol. Total protein was quantified using the bicinchoninic acid (BCA) assay, followed by SDS-PAGE and western blotting with a primary antibody against Scribble (1:500, Santa Cruz sc55543), beta-actin (1:1000, Cell signaling Technology 4967), beta-tubulin (1:200, Santa Cruz sc55529), CHIKV E2 (1:3000, The Native antigen company MAB12130-200), and EGFP (1:3000, Bio Bharati BB-AB0065). Densitometric analysis was performed to quantify the

640   protein levels in western blots using the Fiji software as described elsewhere (Schindelin *et al.*,
641   2012; Davarinejad).
642

TF peptide production and generation of specific antibodies

A 15-amino acid peptide corresponding to the C-terminal region of TF was chemically
synthesized and coupled to a carrier protein, Keyhole Limpet Hemocyanin (KLH), via the N-
terminus (ThermoFisher Scientific). The KLH-TF peptide was dissolved in DMSO and stored at
–20ºC until use.

Polyclonal sera against TF were raised by immunizing two New Zealand white rabbits, aged 3-4
months, with the KLH-TF peptide emulsified in complete/incomplete Freund's adjuvant
(CFA/IFA) as per the following protocol. On Day 0 (priming), 1 mg of KLH-TF peptide was
emulsified with CFA and injected subcutaneously into the rabbits. Subsequently, four boosts (on
Days 21, 42, 63, 84) were given with 300 µg of the peptide emulsified in IFA intramuscularly.
Sera were collected two weeks post-immunization from boost 2 onwards and checked for anti-
TF antibodies using immunoblotting.
655

Immunoprecipitation

HEK293T cells were infected with CHIKV at an MOI of 0.1. Cell extracts were prepared 24
hours post-infection. A polyclonal antibody against Scribble (Invitrogen PA5-54821) was added
overnight to precleared cell extracts at 4 °C on a rocker, followed by incubation with protein A/G
agarose beads (Santa Cruz sc-2003) for 1 hour at 4 °C with gentle mixing. The beads were washed
thrice with lysis buffer (Tris HCl – 20 mM, KCl – 120 mM, EDTA – 0.2 mM, DTT – 1 mM, and
protease inhibitor cocktail). Post-washing, the beads were resuspended in SDS-PAGE buffer, and
the samples were analyzed in immunoblots with anti-Scribble and anti-TF antibodies.
664

MBP-tagged protein production in bacteria

The cDNA corresponding to the TF and 6K regions of CHIKV were cloned into the pMALc6T
vector (New England Biolabs), in conjunction with N-terminal Histidine and Maltose binding
protein (MBP) tags, and a TEV protease cleavage site. Recombinant expression of MBP-tagged
TF and 6K proteins was induced by the addition of Isopropyl ß-D-1-thiogalactopyranoside in *E.*
coli Rosetta pLysS cells at 16 °C at an OD 600 of 0.6. The expressed proteins were 2-step purified

by Ni-NTA affinity chromatography followed by size exclusion chromatography (SEC) in a

buffer containing 25 mM HEPES, 100 mM NaCl, and the detergent n-Dodecyl β-D maltoside

(DDM).

Confocal microscopy

HEK293T cells were cultured on glass coverslips and infected with CHIKV at an MOI of 0.1.

Twenty-four hours post-infection, cells were washed with chilled PBS and fixed with 4%

paraformaldehyde (PFA) for 10 min, followed by permeabilization with 0.1 % Triton X-100 in

PBS for 5 min and blocked with 5% bovine serum albumin for 60 min. Immunostaining was

performed at 4 °C for Scribble, Ubiquitin, and CHIKV proteins using anti-Scribble (1:500,

Invitrogen PA5-54821), anti-Ubiquitin (1:300, Santa Cruz sc-8017), anti-E2 (1:500, The Native

Antigen Company MAB12130-200), and anti-capsid (1:200, The Native Antigen Company

MAB12131-200) antibodies. Post-primary antibody incubation, Alexa fluor-labelled appropriate

secondary antibodies were added for 60 min. All washes were carried out using chilled PBS.

Nuclei were labelled with DAPI (Invitrogen D1306) for 10 min, and cells were mounted using

Prolong Glass Antifade Mountant (Invitrogen P36982). The slides were dried for a few minutes

and sealed with nail paint. Imaging was performed on a confocal microscope (Leica TCS SP8)

at a magnification of 63X using the oil immersion objective.

For transfection of pEGFPC1 constructs, HEK293T cells were seeded and grown at 37°C for 24

h, followed by the addition of plasmid DNA in conjunction with lipofectamine2000 (Invitrogen)

according to the manufacturer's protocol. Briefly, DNA and lipofectamine (1:3 w/v) were diluted

in Opti-MEM (Gibco) and incubated at room temperature (RT) for 20 min. The DNA-

lipofectamine complexes were used to replace the cellular growth medium for 5 h. Twenty-four

hours post-transfection, cells were washed thrice with PBS, fixed with 4% PFA, and processed

as above. Co-localization was estimated as the percentage of qualified double-positive cells

relative to the total number of infected or transfected cells as visualized in four distinct fields.

Mass spectroscopy

Sample Preparation: 25μg protein sample was first reduced with 5 mM TCEP and subsequently

alkylated with 50 mM iodoacetamide. The protein was then digested with Trypsin at a 1:50

Trypsin-to-lysate ratio for 16 hours at 37°C. After digestion, the mixture was purified using a

C18 silica cartridge and then concentrated by drying in a speed vac. The resulting dried pellet

was resuspended in buffer A, which consists of 2% acetonitrile and 0.1% formic acid.

Mass Spectrometric Analysis of Peptide Mixtures: For mass spectrometric analysis, all the

experiments were performed on an Easy-nLC-1000 system (Thermo Fisher Scientific) coupled

with an Orbitrap Exploris 240 mass spectrometer (Thermo Fisher Scientific) and equipped with

a nano electrospray ion source. 1ug of phosphopeptides sample was dissolved in buffer A

containing 2% acetonitrile/0.1% formic acid and resolved using Picofrit column (1.8-micron

resin, 15cm length). Gradient elution was performed with a 0–38% gradient of buffer B (80%

acetonitrile, 0.1% formic acid) at a flow rate of 500nl/min for 96mins, followed by 90% of buffer

B for 11 min and finally column equilibration for 3 minutes. Orbitrap Exploris 240 was used to

acquire MS spectra under the following conditions: Max IT = 60ms, AGC target = 300%; RF

Lens = 70%; R = 60K, mass range = 375−1500. MS2 data was collected using the following

conditions: Max IT= 60ms, R= 15K, AGC target 100%. MS/MS data was acquired using a data-

dependent top20 method dynamically choosing the most abundant precursor ions from the survey

scan, wherein dynamic exclusion was employed for 30s.

Data Processing: Sample was processed and RAW files generated was analyzed with Proteome

Discoverer (v2.5) against UniProt reference database. For dual Sequest and Amanda search, the

precursor and fragment mass tolerances were set at 10 ppm and 0.02 Da, respectively. The

protease used to generate peptides, i.e. enzyme specificity was set for trypsin/P (cleavage at the

C terminus of "K/R: unless followed by "P"). Carbamidomethyl on cysteine as fixed modification

and oxidation of methionine and N-terminal acetylation were considered as variable

modifications for database search. Both peptide spectrum match and protein false discovery rate

were set to 0.01 FDR.

shRNA mediated knockdown of Scribble

HEK293T cells were seeded and grown at 37 °C for 24 h, and then the Scribble short hairpin

RNA (shRNA) containing plasmid DNAs were transfected with lipofectamine 2000 as described

earlier. Four different shRNAs were used in different conditions, either alone or in combination,

to optimize Scribble knockdown. The effect of the knockdown was analyzed 36 hours post-

transfection using western blot analysis and the results were quantified via densitometry. Two

conditions, shRNA.C1: AAACCACAAAATAGAGTCT, and shRNA.C2: a combination of

AAACCACAAAATAGAGTCT and TGTACAAACATCACTAGTT, produced maximum and

consistent knockdown of ~50 %, and were utilized for further studies.

MTT assay

Assays with the tetrazolium dye 3-[4, 5-dimethylthiazol-2-yl]-2, 5-diphenyl tetrazolium bromide

(MTT) were performed as described in the product's manual (Himedia) with slight modifications.

Briefly, 10 µl MTT (5 mg/ml, Himedia) was added to cells, and the formazan crystals were

dissolved in dimethyl sulfoxide after 30 min. The absorbance measurements were recorded at

570 and 690 nm.

Quantitative real-time polymerase chain reaction (qRT-PCR):

CHIKV-infected HEK293T cells were washed with PBS, harvested in Trizol (Takara), and stored

at -80 °C. RNA was isolated from the samples as per the manufacturer's protocol, quantitated

using nanodrop (Thermo Scientific), and 5 µg RNA was reverse transcribed into cDNA using a

cDNA synthesis kit (Takara Bio) as per the manufacturer's instructions. The cDNA was used to

analyze human Scribble and CHIKV E1 expression levels by qRT-PCR using gene-specific

primers (Supplementary Table S2). PCR reactions were conducted on an ABI 7500 FAST

mastercycler, and the relative quantification of gene expression was carried out using the $\Delta\Delta Ct$

method with GAPDH as the housekeeping gene.

Plaque assay

Vero cells were seeded in a six-well plate in DMEM with 10% FBS (Gibco) and antibiotics

(Corning) and incubated for 24 h at 37°C, 5% CO_2. The cell growth media was then replaced

with serum-free media, and cells were incubated for 4-6 h. Subsequently, 10-fold dilutions (10^{-5}

to 10^{-7}) of the virus-containing culture supernatants of CHIKV-infected HEK293T cells, in

serum-free media, were used to replace the cellular growth media. Virus particles in the diluted

culture supernatant were allowed to infect Vero cells for 1 h with shaking every 15 min. Post 1

h, the cells were washed with PBS, and 3 ml of an overlay mix of 2% low melting agarose in 2X

DMEM was added to each well. The agarose was allowed to solidify at 4°C for 15 min, and cells

were then incubated at 37°C for 48 h. For plaque visualization, cells were fixed with 10 %

formaldehyde for 4-5 h at room temperature and stained with crystal violet for 5 min, followed

by washing with tap water. The plate was allowed to dry for a few minutes, and the plaques were

counted. The plaque-forming units (PFU) were calculated using the following formula: PFU/ml

= (number of plaques x dilution factor)/(inoculum volume). (Modify the plaque assay details to

include Sanchi's new data)

Acknowledgements: The work was funded by the Science and Engineering Research Board,

Government of India (grant number SPF/2021/000115) and Indian Council of Medical Research

(IIRPIG-2024-01-00529). The authors thank the Central Research Facility (CRF) and Sophisticated

Analytical & Technical Help Institutes (SATHI) foundation for the confocal microscopy facility. The

authors also extend their thanks to Dr. Anjali Kapoor, CRF, IIT Delhi, for her assistance during

confocal microscopy imaging. The authors also thank Prof. Manoj Menon, KSBS, IIT Delhi for

providing antibodies. RK and PT would also like to acknowledge IIT Delhi for providing fellowship

support.

Author contributions: Conceptualization: RK, DD, PT, MB; Methodology: PT, RK, DD, MB;

Investigation: RK, PT, DD, YR, SBK, B, SA, SYM, KS; Resources: SB, MB; Writing-original draft:

RK, PT, DD, MB; Writing- review & editing: RK, PT, MB; Visualization: RK, PT, MB; Supervision:

SB, MB. Project administration: MB; Funding acquisition: MB.

Conflict of interest:

The authors declare no conflict of interest.

References

References

Arpin-André C & Mesnard JM (2007) The PDZ domain-binding motif of the human T cell
leukemia virus type 1 tax protein induces mislocalization of the tumor suppressor hScrib in
T cells. *J Biol Chem* 282: 33132–33141

Barreda D, Gutiérrez-González LH, Martínez-Cordero E, Cabello-Gutiérrez C, Chacón-Salinas
R & Santos-Mendoza T (2020) The Scribble Complex PDZ Proteins in Immune Cell
Polarities. *J Immunol Res* 2020: 5649790

Barreda D, Sánchez-Galindo M, López-Flores J, Nava-Castro KE, Bobadilla K, Salgado-Aguayo
A & Santos-Mendoza T (2018) PDZ proteins are expressed and regulated in antigen-
presenting cells and are targets of influenza A virus. *J Leukoc Biol* 103: 731–738
doi:10.1002/JLB.4AB0517-184R

Bartholomeeusen K, Daniel M, LaBeaud DA, Gasque P, Peeling RW, Stephenson KE, Ng LFP
& Ariën KK (2023) Chikungunya fever. *Nat Rev Dis Prim* 9: 17

Bonello TT & Peifer M (2019) Scribble: A master scaffold in polarity, adhesion, synaptogenesis,
and proliferation. *J Cell Biol* 218: 742–756

Brown R, Wan J & Kielian M (2018) The Alphavirus Exit Pathway: What We Know and What
We Wish We Knew. *Viruses* 10: 89

Chmielewski D, Schmid MF, Simmons G, Jin J & Chiu W (2022) Chikungunya virus assembly
and budding visualized in situ using cryogenic electron tomography. *Nat Microbiol* 7: 1270–
1279

Davarinejad H Quantifications of Western Blots with ImageJ.

Desta IT, Porter KA, Xia B, Kozakov D & Vajda S (2020) Performance and Its Limits in Rigid
Body Protein-Protein Docking. *Structure* 28: 1071-1081.e3

Dey D, Siddiqui SI, Mamidi P, Ghosh S, Kumar CS, Chattopadhyay S, Ghosh S & Banerjee M
(2019) The effect of amantadine on an ion channel protein from Chikungunya virus. *PLoS
Negl Trop Dis* 13: e0007548

Elmasri Z, Negi V, Kuhn RJ & Jose J (2022) Requirement of a functional ion channel for Sindbis
virus glycoprotein transport, CPV-II formation, and efficient virus budding. *PLoS Pathog*
18: e1010892

Firth AE, Chung BYW, Fleeton MN & Atkins JF (2008) Discovery of frameshifting in
Alphavirus 6K resolves a 20-year enigma. *Virol J* 5: 108

Gautam B (2021) Energy Minimization. In *Homology Molecular Modeling - Perspectives and*

Applications, Trindade Maia R Maciel de Moraes Filho R & Campos M (eds) IntechOpen

Hallengärd D, Kakoulidou M, Lulla A, Kümmerer BM, Johansson DX, Mutso M, Lulla V,

Fazakerley JK, Roques P, Le Grand R, et al (2014) Novel Attenuated Chikungunya Vaccine

Candidates Elicit Protective Immunity in C57BL/6 mice. *J Virol* 88: 2858–2866

Hess B, Bekker H, Berendsen HJC & Fraaije JGEM (1997) LINCS: A linear constraint solver

for molecular simulations. *J Comput Chem* 18: 1463–1472

Hu B, Li S, Zhang Z, Xie S, Hu Y, Huang X & Zheng Y (2016) HCV NS4B targets Scribble for

proteasome-mediated degradation to facilitate cell transformation. *Tumor Biol* 37: 12387–

12396

Humphrey W, Dalke A & Schulten K (1996) VMD: Visual molecular dynamics. *J Mol Graph*

14: 33–38

Inada M, Izawa G, Kobayashi W & Ozawa M (2016) 293 cells express both epithelial as well as

mesenchymal cell adhesion molecules. Int J Mol Med 37: 1521–1527.

Javorsky A, Humbert PO & Kvansakul M (2022a) Molecular basis of Tick Born encephalitis

virus NS5 mediated subversion of apico-basal cell polarity signalling. *Biochem J* 479: 1303–

1315

Javorsky A, Humbert PO & Kvansakul M (2022b) Structural Basis of the Avian Influenza NS1

Protein Interactions with the Cell Polarity Regulator Scribble. *Viruses* 14: 583

Javorsky A, Humbert PO & Kvansakul M (2023) Viral subversion of the cell polarity regulator

Scribble. *Biochem Soc Trans* 51: 415–426

Jose J, Snyder JE & Kuhn RJ (2009) A structural and functional perspective of alphavirus

replication and assembly. *Future Microbiol* 4: 837–856

Kielian M, Chanel-Vos C & Liao M (2010) Alphavirus entry and membrane fusion. *Viruses* 2:

796–825

Ko J, Park H, Heo L & Seok C (2012) GalaxyWEB server for protein structure prediction and

refinement. *Nucleic Acids Res* 40: W294–W297

Kozakov D, Beglov D, Bohnuud T, Mottarella SE, Xia B, Hall DR & Vajda S (2013) How good

is automated protein docking? *Proteins Struct Funct Bioinforma* 81: 2159–2166

Kozakov D, Hall DR, Xia B, Porter KA, Padhorny D, Yueh C, Beglov D & Vajda S (2017) The

ClusPro web server for protein–protein docking. *Nat Protoc* 12: 255–278

Krishnakumar V & Kannan M (2020) Microbial Pathogenesis: Virus Pathogen–Host
Interactions. In *Model Organisms for Microbial Pathogenesis, Biofilm Formation and*
Antimicrobial Drug Discovery, Siddhardha B Dyavaiah M & Syed A (eds) pp 19–35.
Singapore: Springer, Singapore

Kumar M, Michael S, Alvarado-Valverde J, McRossed D Sign©száros B, Sámano-Sánchez H,
Zeke A, Dobson L, Lazar T, Örd M, Nagpal A, *et al* (2022) The Eukaryotic Linear Motif
resource: 2022 release. *Nucleic Acids Res* 50: D497–D508

Kumari R, Kumar R & Lynn A (2014) g_mmpbsa —A GROMACS Tool for High-Throughput
MM-PBSA Calculations. *J Chem Inf Model* 54: 1951–1962

Kundu K & Backofen R (2014) Cluster based prediction of PDZ-peptide interactions. *BMC*
Genomics 15: S5

Kundu K, Mann M, Costa F & Backofen R (2014) MoDPepInt: An interactive web server for
prediction of modular domain-peptide interactions. *Bioinformatics* 30: 2668–2669

Liu H, Golebiewski L, Dow EC, Krug RM, Javier RT & Rice AP (2010) The ESEV PDZ-Binding
Motif of the Avian Influenza A Virus NS1 Protein Protects Infected Cells from Apoptosis
by Directly Targeting Scribble. *J Virol* 84: 11164–11174

Ma-Lauer Y, Lei J, Hilgenfeld R & von Brunn A (2012) Virus–host interactomes — antiviral
drug discovery. *Curr Opin Virol* 2: 614–621

Mark P & Nilsson L (2001) Structure and Dynamics of the TIP3P, SPC, and SPC/E Water
Models at 298 K. *J Phys Chem A* 105: 9954–9960

Martyna GJ, Klein ML & Tuckerman M (1992) Nosé–Hoover chains: The canonical ensemble
via continuous dynamics. *J Chem Phys* 97: 2635–2643

Melton J V., Ewart GD, Weir RC, Board PG, Lee E & Gage PW (2002) Alphavirus 6K proteins
form ion channels. *J Biol Chem* 277: 46923–46931

Millsapps EM, Underwood EC & Barr KL (2022) Development and Application of Treatment
for Chikungunya Fever. *Res Rep Trop Med* 13: 55–66

Nakagawa S & Huibregtse JM (2000) Human Scribble (Vartul) Is Targeted for Ubiquitin-
Mediated Degradation by the High-Risk Papillomavirus E6 Proteins and the E6AP
Ubiquitin-Protein Ligase. *Mol Cell Biol* 20: 8244–8253

Parrinello M & Rahman A (1981) Polymorphic transitions in single crystals: A new molecular
dynamics method. *J Appl Phys* 52: 7182–7190

Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC & Ferrin TE (2004)
UCSF Chimera?A visualization system for exploratory research and analysis. *J Comput*
Chem 25: 1605–1612

Ramsey J, Chavez M & Mukhopadhyay S (2019) Domains of the TF protein important in
regulating its own palmitoylation. *Virology* 531: 31–39

Ramsey J & Mukhopadhyay S (2017) Disentangling the frames, the state of research on the
alphavirus 6K and TF proteins. *Viruses* 9: 228

Ramsey J, Renzi EC, Arnold RJ, Trinidad JC & Mukhopadhyay S (2017) Palmitoylation of
Sindbis Virus TF Protein Regulates Its Plasma Membrane Localization and Subsequent
Incorporation into Virions. *J Virol* 91: e02000-16

Santoni MJ, Kashyap R, Camoin L & Borg JP (2020) The Scribble family in cancer: twentieth
anniversary. *Oncogene* 39: 7019–7033

Sanz MA, Pérez L & Carrasco L (1994) Semliki forest virus 6K protein modifies membrane
permeability after inducible expression in Escherichia coli cells. *J Biol Chem* 269: 12106–
12110

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden
C, Saalfeld S, Schmid B, *et al* (2012) Fiji: An open-source platform for biological-image
analysis. *Nat Methods* 9: 676–682

Seok C, Baek M, Steinegger M, Park H, Lee GR & Won J (2021) Accurate protein structure
prediction: what comes next? *BIODESIGN* 9: 47–50

Shin W-H, Lee GR, Heo L, Lee H & Seok C (2014) Prediction of Protein Structure and
Interaction by GALAXY protein modeling programs. *Bio Des* 2: 1–11

Singh H, Mudgal R, Narwal M, Kaur R, Singh VA, Malik A, Chaudhary M & Tomar S (2018)
Chikungunya virus inhibition by peptidomimetic inhibitors targeting virus-specific cysteine
protease. *Biochimie* 149: 51–61

Snyder JE, Kulcsar KA, Schultz KLW, Riley CP, Neary JT, Marr S, Jose J, Griffin DE & Kuhn
RJ (2013) Functional Characterization of the Alphavirus TF Protein. *J Virol* 87: 8511–8523

Van Der Spoel D, Lindahl E, Hess B, Groenhof G, Mark AE & Berendsen HJC (2005)
GROMACS: Fast, flexible, and free. *J Comput Chem* 26: 1701–1718

Stephens R, Lim K, Portela M, Kvansakul M, Humbert PO & Richardson HE (2018) The
Scribble Cell Polarity Module in the Regulation of Cell Signaling in Tissue Development

and Tumorigenesis. *J Mol Biol* 430: 3585–3612

Sun S, Xiang Y, Akahata W, Holdaway H, Pal P, Zhang X, Diamond MS, Nabel GJ & Rossmann

MG (2013) Structural analyses at pseudo atomic resolution of Chikungunya virus and

antibodies show mechanisms of neutralization. *Elife* 2: e00435

Thomas M & Banks L (2018) PDZ domain proteins and virus infection: gateways to the cell. *J*

Mol Biol 430**,** 2896–2911.

Tironi IG, Sperb R, Smith PE & van Gunsteren WF (1995) A generalized reaction field method

for molecular dynamics simulations. *J Chem Phys* 102: 5451–5459

Torres-Flores J & Arias C (2015) Tight Junctions Go Viral! *Viruses* 7: 5145–5154

Vajda S, Yueh C, Beglov D, Bohnuud T, Mottarella SE, Xia B, Hall DR & Kozakov D (2017)

New additions to the ClusPro server motivated by CAPRI. *Proteins Struct Funct Bioinforma*

85: 435–444

Wahid B, Ali A, Rafique S & Idrees M (2017) Global expansion of chikungunya virus: mapping

the 64-year history. *Int J Infect Dis* 58: 69–76

Wallace AC, Laskowski RA & Thornton JM (1995) LIGPLOT: a program to generate schematic

diagrams of protein-ligand interactions. *"Protein Eng Des Sel* 8: 127–134

Wang Y & Nie G (2021) Overview of human HtrA family proteases and their distinctive

physiological roles and unique involvement in diseases, especially cancer and pregnancy

complications. *Int J Mol Sci* 22

Werme K, Wigerius M & Johansson M (2008) Tick-borne encephalitis virus NS5 associates with

membrane protein scribble and impairs interferon-stimulated JAK-STAT signalling. *Cell*

Microbiol 10: 696–712

Wong KZ & Chu JJH (2018) The interplay of viral and host factors in chikungunya virus

infection: Targets for antiviral strategies

Yang J & Zhang Y (2015) Protein Structure and Function Prediction Using I-TASSER. *Curr*

Protoc Bioinforma 52: 5.8.1–5.8.15

Yap ML, Klose T, Urakami A, Hasan SS, Akahata W & Rossmann MG (2017) Structural studies

of Chikungunya virus maturation. *Proc Natl Acad Sci* 114: 13703–13707

Ye F & Zhang M (2013) Structures and target recognition modes of PDZ domains: Recurring

themes and emerging pictures. *Biochem J* 455: 1–14

Zhan L, Rosenberg A, Bergami KC, Yu M, Xuan Z, Jaffe AB, Allred C & Muthuswamy SK

(2008) Deregulation of Scribble Promotes Mammary Tumorigenesis and Reveals a Role for

Cell Polarity in Carcinoma. *Cell* 135: 865–878

Zhang Y (2008) I-TASSER server for protein 3D structure prediction. *BMC Bioinformatics* 9:

40

FIGURE LEGENDS

Figure 1. CHIKV TF as a binding partner for human Scribble. The CHIKV RNA genome

has two open reading frames (ORFs) separated by a junctional region (J). The 5' ORF (ORF 1)

codes for four non-structural proteins (nsP1-4). The 3' ORF (ORF 2) codes for 5 structural

proteins, namely capsid (C), E3, E2, 6K, and E1. The 6K gene has a ribosomal slippery site which

leads to the production of a transframe (TF) protein (Firth *et al.,* 2008) (upper panel, A).

Schematic of 6K and TF sequences and the ribosome slip site (F at 48th and L at 49th position)

(lower panel, A). Potential PDZ binding motifs (PBMs) present in TF as predicted by different

softwares (B). Predicted interactions between TF and Scribble PDZ1, Scribble PDZ4, and

HTRA1 (C). RMSF plot for TF interactions with Scribble and HTRA1 (D). Binding of TF C-

terminal residues with ScPDZ1(E).

Figure 2. Indirect assessment of TF and Scribble association in transfected cells through

confocal microscopy . HEK293T cells were transfected with pEGFPC1 vector or pEGFPC1

plasmids carrying TF or 6K. Twenty-four hours post-transfection, cells were fixed, and

immunostaining was performed for Scribble (red). Inset shows zoomed images of an EGFP-TF

expressing cell. Arrowheads indicate Scribble puncta and co-localization with TF (A).

Mutational analysis of TF and its effect on Scribble localization (B). Expression levels of TF and

TF mutants in HEK293T cells. Twenty-four hours post-transfection, cells were lysed, and 6K

and TF expression were analyzed using western blot with anti-GFP antibody (C). L: protein

ladder, C1: pEGFPC1 empty vector, TF Mut1: EGFP-TF-HGAATV, TF Mut2: EGFP-TF-

AAAATV. Immunoprecipitation (IP) of Scribble using Scribble specific antibody (i) and co-IP

of TF-Scribble in transfected cells (C1, TF-WT, TF-Mut1 and TF-Mut2) (D). Fluorescent

intensity profiles of the lined areas are displayed alongside the merged image in all cases.

Figure 3. Assessment of TF-Scribble association in infected cells. HEK293T cells were

infected with CHIKV at 0.1 MOI. Co-localization of TF-Scribble during infection was analyzed

using TF and Scribble specific antibodies at the indicated time points (A), Co-localization;

Scribble, E2 and Capsid specific antibodies at the indicated time points (B), Co-localization;

Scribble, E2 and TF specific antibodies at the indicated time points (C), Immunoprecipitation

(IP) of Scribble using Scribble specific antibody and co-IP of TF-Scribble in infected cells (D).

Fluorescent intensity profiles of the lined areas are displayed alongside the merged image in all
cases.

Figure 4. Effect of CHIKV infection on Scribble localization and expression. HEK293T cells
were infected with CHIKV and Scribble levels were analyzed at 24 and 48 hpi using western blot
and quantitated using densitometry (A). Effect of TF on Scribble expression. HEK293T cells
were transfected with pEGFPC1-TF and its mutants. Only vector and pEGFPC1-6K were also
transfected, and Scribble levels were analyzed 48 h post-transfection using western blot and
quantitated using densitometry (B) Immunostaining for Ubiquitin (Ubiq) and Scribble in CHIKV
infected HEK293T cells at the indicated time points post-infection(C). Data represent mean ±
SEM of two biological replicates. Fluorescent intensity profiles of the lined areas are displayed
alongside the merged image in all cases.

Figure 5. Effect of TF on Scribble ubiquitination. HEK293T cells were transfected with
pEGFPC1 vector or pEGFPC1 plasmids carrying TF or 6K. Twenty-four hours post-transfection,
cells were fixed, and immunostaining was performed for Scribble (red) and Ubiquitin (green).
Inset shows a zoomed image of an EGFP-TF expressing cell. Arrowhead(white) indicates EGFP-
TF, scribble, and ubiq puncta and colocalization in the merged image (D). HEK293T cells were
infected with CHIKV, treated with MG132, and Scribble levels were analyzed at 24 hpi using
western blot (E). Data represent mean ± SEM of two biological replicates. Fluorescent intensity
profiles of the lined areas are displayed alongside the merged image in all cases.

1000

Figure 6. Effect of Scribble knockdown on CHIKV infection. Scrib shRNAs were transfected
into HEK293T cells, and knockdown was confirmed 36 hours post-transfection using western
blot and quantitated using densitometry (A). Scribble knockdown was also confirmed at RNA
level using real-time PCR (B). The effect of Scribble knockdown on cell viability was assessed
using MTT assay (C). Thirty-six hours post-transfection, cells were infected with CHIKV at MOI
of 0.1 for 24 hours. The effect of Scribble knockdown on CHIKV progeny virus was determined
using plaque assay (D) and CHIKV RNA levels using real-time PCR (E). Data represents mean
± SEM. **, $p \leq 0.01$; ***, $p \leq 0.001$, ns: non-significant (Student's t-test).

1009

Figure 7. ΔTF CHIKV is unable to cause Scribble punctation. A) The infectivity of WT and ΔTF CHIKV compared using plaque assay and by measuring respective RNA levels using real-time PCR B)Immunostaining for ubiquitin and Scribble in CHIKV-infected HEK293T cells 12 hours post-infection and 24 hours post-infection. Fluorescent intensity profiles of the lined areas are displayed alongside the merged image in all cases. C) Western blot depicting scribble expression in HEK 293T cells infected with 0.1 MOI WT mcherry CHIKV and ΔTF mcherry CHIKV at 24 hours post-infection and 36 hours post-infection.

 TABLES

 Table 1. Binding interface of docked complexes

System	CHIKV-TF (Hydrogen bonds)	CHIKV-TF (Hydrophobic interactions)
TF-ScPDZ1	Asp68, Glu70, Thr75	Trp11, Trp19, Leu23, Ala27, His71, Gly72, Gly73, Ser74
TF-ScPDZ4	Glu70, His71, Gly72, Thr75	Tyr3, Leu10, Trp11, Ala27, Ile30, Val31, Asn34, Leu38
TF-HtPDZ	Gln15, Asn66, Glu70, His71, Gly73	Leu10, Trp11, Leu20, Ser67, Asp68, Pro69, Gly72

1021

1022 **Table 2. Mass spectrometric analysis of immunoprecipitate using anti-Scribble Ab**

Sample	Accession	Description	Coverage [%]	# Peptides	# PSMs
C1	Q14160	Protein scribble homolog OS=Homo sapiens	58	88	771
	GFP	GFP	-	-	-
TF-WT	Q14160	Protein scribble homolog OS=Homo sapiens	58	87	683
	TF-GFP	TF-GFP WT	54	14	83
TF-MUT1	Q14160	Protein scribble homolog OS=Homo sapiens	*58*	*94*	*721*
	TF-GFP MUT1	TF-GFP MUT1	*28*	*7*	*38*
TF-MUT2	Q14160	Protein scribble homolog OS=Homo sapiens	*57*	*83*	*538*
	TF-GFP MUT2	TF-GFP MUT2	*21*	*6*	*41*

1023
1024

1025　**Table 3. Prediction of PBM in alphavirus TF using the ELM software** (Kumar *et al*, 2022)

Alphavirus	Region	PBM	Class
SINV	65-70	CATDTV	Class_3
ONNV	71-76	HGGSTV	Class_1
UNAV	69-74	HGGISV	Class_2
RRV	70-75	CGGVPV	Class_2
IGBO-ORBA	71-76	HGGSTV	Class_1
MAYV	70-75	PSGIPV	Class_2
SFV	70-75	RGGVPV	Class_2
EEEV	NA		
VEEV	NA		
MIDV	NA		

1026

1027　**Table 4. Prediction of PBM in CHIKV proteins using the ELM software** (Kumar *et al*,
1028　2022)

Protein	Region	PBM	Class
nsP1	NA		
nsP2	NA		
nsP3	NA		
nsP4	NA		
Capsid	NA		
E3	NA		
E2	418-423	VRTTKA	Class_1
6K	56-61	AHTVSA	Class_2
TF	71-76	HGGSTV	Class_1
E1	NA		

1029

A)

ORF 1 (Non-structural proteins) | ORF 2 (Structural proteins)

Cap | nsP1 | nsP2 | nsP3 | nsP4 | J | C | E3 | E2 | 6K | E1 | Poly (A)

C | E3 | E2 | 6K | E1 → Major product

C | E3 | E2 | TF → Minor product

Slippery sequence
UUUUUUA | 6K

...........LCNCLKLLPCCKTLA**FL**AVMSIGAHTV | E1 | HVTVIPNTVGV
.....
48 49
-1 ribosome frameshift
SRNEHRCPHCERVRTRNSDPE**HGGSTV** | TF

B)

SOFTWARE	REGION	PBM
PDZ PepInt	71-75	HGGST
ELM	71-76	HGGSTV

D) RMSF (nm) vs Residue plot; curves for CHIKV-TF, HtPDZ, ScPDZ1, ScPDZ4.

E) Binding Energy (kJ/mol) vs CHIKV-TF Residue (60-76) bar chart.

A)

B)

C)

D)

E)