Title: Temporal sensitivity for achromatic and chromatic flicker across the visual cortex

Running head: Visual temporal sensitivity

Author(s):
Carlyn Patterson Gentile[1,2], Manuel Spitschan[3,4], Huseyin O. Taskin[1], Andrew S. Bock[1], Geoffrey K. Aguirre[1]

Affiliation Information for All Authors:
[1]University of Pennsylvania
[2]Children's Hospital of Philadelphia
[3]Translational Sensory & Circadian Neuroscience, Max Planck Institute for Biological Cybernetics, Tübingen, Germany
[4]Chronobiology & Health, TUM Department of Sport and Health Sciences (TUM SG), Technical University of Munich, Munich, Germany

[*]**Corresponding Author:**
Geoffrey K Aguirre
3400 Spruce Street, Philadelphia, PA 19104
aguirreg@upenn.edu

Pages:
Figures:
Abstract (250 words max): 249
Introduction (650 words max):
Discussion (1,500 words max):

Conflicts of Interest: The authors declare that they have no relevant financial interests that relate to the research described in this paper.

Acknowledgements: none.

Abbreviations:

LGN	Lateral geniculate nucleus
TSF	Temporal Sensitivity Function
RGC	Retinal ganglion cell
V1	Primary visual cortex

Abstract

The classes of retinal ganglion cells (RGCs) receive different combinations of L, M, and S cone

inputs and give rise to one achromatic and two chromatic post-receptoral channels. Beyond the

retina, RGC outputs are subject to filtering and normalization along the geniculo-striate pathway,

ultimately producing the properties of human vision. The goal of the current study was to

determine temporal sensitivity across the three post-receptoral channels in subcortical and

cortical regions involved in vision, to better characterize post-retinal temporal processing. We

measured functional magnetic resonance imaging (MRI) responses at 7 Tesla from participants

viewing a high-contrast, flickering, spatially-uniform wide (~140°) field. Stimulus flicker

frequency varied logarithmically between 2 and 64 Hz and targeted the L+M+S, L–M, and S–

[L+M] cone combinations. These measurements were used to create temporal sensitivity

functions (TSFs) of primary visual cortex (V1) across eccentricity, and spatially averaged

responses from lateral geniculate nucleus (LGN), V2/V3, hV4, and MT. Functional MRI

responses reflected known properties of the visual system, including higher peak temporal

sensitivity to achromatic vs. chromatic stimuli, and low-pass filtering between the LGN and V1.

V1 had the slowest peak temporal sensitivity across cortical regions, which increased at higher

levels of the visual cortical hierarchy. Unexpectedly, peak temporal sensitivity decreased at

greater eccentricities in area V1, especially for achromatic stimuli. Comparison of measured

cortical responses to a model of integrated retinal output to our stimuli demonstrates that

extensive filtering and amplification is applied to post-retinal signals.

Significance Statement (120 words max)

We report the temporal sensitivity of human visual cortex across the three canonical post-

receptoral channels from central vision to the far periphery. Functional MRI measurements of

responses from the LGN, V1, and higher visual cortical areas demonstrate modification of

temporal sensitivity across the visual hierarchy. This includes amplification of chromatic signals

between the LGN and V1, and a steady increase in peak temporal sensitivity across visual

areas beyond area V1. Within V1, we unexpectedly find that peak temporal sensitivity slows in

the periphery, especially for achromatic stimuli. Comparison of our results to a model of retinal

output demonstrates the substantial post-retinal filtering at work, yielding greater uniformity of

responses across area V1 than would be predicted from unmodified retinal signals.

Introduction

Cone and retinal ganglion cell (RGC) density declines exponentially from the fovea to

the periphery. There is a corresponding and well-studied change in the spatial frequency

sensitivity (acuity) of human perception that reflects this change with eccentricity (Thibos et al.,

1996). Although less numerous, cell populations in the peripheral retina are found to have faster

kinetics (Seiple and Holopigian, 1996; Solomon, 2005), raising the possibility that visual

perception has a corresponding change in temporal frequency sensitivity with eccentricity.

Signals from the cones are combined into three post-receptoral channels (Figure 1a):

one achromatic or luminance (L+M) channel, and two chromatic channels, red-green (L-M), and

blue-yellow (S–[L+M]). The eccentricity dependence of temporal sensitivity has been most

studied for achromatic stimuli, and generally shows an improvement in threshold contrast

detection for higher frequency flicker in peripheral as compared to central vision (Snowden and

Hess, 1992). In foveal vision, chromatic channels favor slower frequencies as compared to the

achromatic channel (Swanson et al., 1987). Relatively little is known regarding how eccentricity

variation influences temporal sensitivity across the chromatic post-receptoral channels, although

one set of measurements made in the LM plane out to 14° found little variation in temporal

response (Gelfand and Horwitz, 2018).

The properties of threshold detection as measured with psychophysics may differ in

meaningful ways from the perceptual experience of supra-threshold stimuli (i.e., natural vision).

Neural responses to supra-threshold stimuli might be expected to reflect the properties of visual

perception. Electrophysiological studies in non-human primates demonstrate filtering of

temporal information at successive stages of the visual system; the largest decline in temporal

sensitivity occurs between the cones and RGCs, with additional filtering of high-temporal

frequency information between the lateral geniculate nucleus (LGN) and cortex (Hawken et al.,

1996; Horwitz, 2020, 2021). Consequently, we may expect that the temporal sensitivity of

human visual cortex—and presumably human supra-threshold perception—reflects a

substantial modification of the initial retinal signal.

Here we report measurements of the regional and eccentricity dependence of the

temporal sensitivity of human visual cortex. We studied two participants using a 7 Tesla scanner

while they viewed high-contrast, wide-field stimuli that targeted each of the three post-receptoral

channels, modulating across a range of flicker frequencies. Our results provide temporal

sensitivity functions across multiple brain regions representing stages of the visual pathway

from lateral geniculate nucleus (LGN) to extra-striate cortex. More detailed measurements

across eccentricity were also obtained for the primary visual cortex (area V1). Our findings

confirm several temporal filtering properties of the visual system that have been previously

reported, as well as normalization of signals in area V1 across post-receptoral channel and

eccentricity relative to the retina and LGN. Unexpectedly, we find a slowing of temporal

sensitivity to achromatic flicker at far eccentricities.

116
117 **Materials and Methods**

118
119 *Participants*

120   The two participants are authors of this report (P1, age 46; P2 age 32, both male). Both

121   participants had normal or corrected to normal acuity and normal color vision. This study was

122   approved by the University of Pennsylvania Institutional Review Board in accordance with the

123   Declaration of Helsinki, and all participants provided written consent.

Custom contact lens

The participants underwent clinical refraction and measurement of keratometry. These

measures were then used to design custom contact lenses for each participant that provided

either +36 or +32 diopters of magnification, with the latter selected to complement the –4 diopter

myopia for that participant. The contact lenses were Alden Optical, HP54 spherical soft lenses,

constructed from 54% Hioxifilcon D (sourced from Benz Research and Development). The lens

was 0.59 mm at its thickest point. During scanning, the participant wore a contact lens in the

right eye. Prior to each scanning session, the participant underwent pharmacologic dilation of

their right eye; the left eye was patched.

Spectral modulations

Stimuli were generated using a digital light synthesis engine (OneLight Spectra). This

device produces desired spectra through the use of a digital micro-mirror device chip, and was

configured to mix 56 independent primaries with a FWHM of ~16 nm at a refresh rate of 256 Hz.

Spectral modulations were designed to target specific photoreceptor classes alone or in

combination, using the silent substitution approach (Estévez and Spekreijse, 1982). The design

of these stimuli has been described in detail previously (Spitschan et al., 2015; Spitschan and

Woelders, 2018; Barnett et al., 2021). Briefly, a set of three modulations were created, which

targeted L–M, S, and LMS cone combinations; the last of these was a "light flux" modulation, in

which the entire stimulus spectrum was scaled. All three stimulus types were modulations

around the half-on background of the device.

The chromatic L–M and S modulations were designed to minimize differences in

contrast between the fovea and the periphery. This was achieved by treating the cone

fundamentals at 2 and 10 degrees (CIE, 2006) as different classes of photoreceptors, and

tailoring modulations to produce equivalent contrast on these two classes (Barnett et al., 2021).

The cone fundamentals were adjusted for age-related differences in lens density (CIE, 2006),

and the transmittance spectrum of the contact lens material was measured and incorporated

into the model of pre-receptoral filtering.

Spectroradiometric measurements of the stimuli were made before and after each

experimental session (PhotoResearch PR-670). The nominal maximum contrast of the

modulation was approximately 90% for LMS, 50% for S, and 8% for L–M. Supplementary Figure

1 and Supplementary Table 1 provide the predicted contrast of the stimuli as a function of

eccentricity and in the presence of imprecision in device control. The chromaticity of the

background was x = 0.4025, y = 0.4486, luminance Y = ~400 cd/m^2; chromaticity and luminance

computed with respect to the XYZ 10° physiologically relevant color matching functions

(https://cvrl.org). Assuming a 7mm diameter of the pharmacologically dilated eye, the stimulus

background was ~4200 Trolands.

The stimuli were conveyed via a fiber-optic cable to a custom-made, MR-compatible eye

piece that was positioned over the right eye of the participant. After passing through ultra-violet

and heat-absorbing glass, light was back-projected on a 35 mm diameter, opal diffusing glass.

The stimulus surface was positioned ~10 mm from the eye. This geometry provided a uniform

stimulus field of ~60° radial eccentricity. The +36 diopter contact lens produced approximately

x1.18 spectacle magnification. The resulting stimulus field was therefore close to 70° radial

eccentricity.

Scanning environment and MR imaging of the brain

An MR imaging session was conducted for each participant on each of two consecutive

days. During the scanning session, all sources of light within the scanner room were

extinguished and covered. Room lights were on when participants were positioned within the

scanner, but these were turned off at the start of data acquisition.

All MRI images were obtained with a 32-channel head coil on a 7T Siemens scanner.

The anatomical T1w images were acquired with voxel size=0.8 mm iso, resolution =

480x640x176, TR = 2800ms, TE=4.28ms, TI=1500ms, and flip angle=7°. For the BOLD images,

a multiband epfid2d1_130 sequence was used with multiband acceleration factor=5, voxel

size=1 mm iso, resolution=130x130x75, TR=1000ms, TE=26.8ms, flip angle=45°, and

average=1.

Experimental design

Each fMRI acquisition was 336 seconds in duration and presented modulations that

targeted one particular post-receptoral channel (LMS, L–M, or S). The acquisition was

composed of 28, 12 second trials. Each trial presented a spectral modulation that flickered at

one of 7, log-spaced frequencies (0, 2, 4, 8, 16, 32, 64 Hz). A pseudo-random, counterbalanced

ordering of the frequencies was defined (50 elements). This sequence was divided into two-

subsequences ("A" and "B"), each padded with three instances of the 0 Hz trial at the end. A set

of 6 acquisitions composed a scanning "block", and consisted of the ordered acquisitions:

LMS_A, L–M_A, S_A, S_B, L–M_B, LMS_B. A total of 6 blocks were collected for each

participant across the two sessions. The stimulus field was set to the half-on background before

and between each acquisition.

The sinusoidal spectral modulation presented in each block was windowed by a 1.5

second, half-cosine ramp at trial onset and offset. An intermittent attention event occurred with a

33% probability during each trial. The attention event could occur at any point during the trial,

except constrained not to take place during first or last 2 seconds of the trial. The attention

event was a 250 msec darkening of the stimulus field. The participant was asked to make a

bilateral button press on a fiber-optic pad in response to each attention event. Performance on

this attention task was uniformly high for the two participants.

Pre-processing of fMRI data

Three-dimensional surface reconstruction of the anatomical images for both participants

was performed using Freesurfer (Dale et al., 1999). For the preprocessing of the BOLD images,

we used the *fmriprep* pipeline with custom options (Esteban et al., 2019). The pipeline included

bias field correction (Tustison et al., 2010), brain extraction and normalization (Avants et al.,

2008), tissue segmentation (Zhang et al., 2001), slice timing correction (Cox, 1996), and motion

correction (Jenkinson et al., 2002). As no B0 field maps were collected, we performed

susceptibility correction with symmetric group-wise diffeomorphic normalization (*SyN*). This

method aims to correct distortion by inverting the intensity of BOLD images (Huntenburg, 2014;

Wang et al., 2017) and registering these inverted images to subject T1w image using an

average fieldmap template as the constraint (Treiber et al., 2016). Noise regressors were

obtained with AROMA (Pruim et al., 2015). The set of regressors was inspected and any

component with a spatial extent that was largely confined to the occipital cortex was excluded.

The effect of the remaining set of regressors was removed from the data using the "non-

aggressive" denoising strategy. Finally, we mapped the preprocessed volumetric BOLD data to

HCP CIFTI FSLR_32k template space with *ciftify* which uses the algorithms and templates from

the HCP minimal processing pipeline (Glasser et al., 2013) to map volumetric timeseries data to

CIFTI surfaces. Because we did not expect, and did not observe, response differences between

the hemispheres, we averaged the data from the left and right hemisphere into a single

"pseudo" hemisphere.

Modeling of fMRI time-series data

The time-series data in each voxel was fit with a non-linear model that simultaneously

estimated the amplitudes of response to the stimuli and the shape of the hemodynamic

response function. First, the time-series data in each voxel in each acquisition were converted

to percentage change units. The data from all acquisitions across both sessions were then

concatenated, for a total of 12,096 time points. The amplitude of response to each stimulus

frequency was modeled as a 12 second step function. Attention events were modeled as delta

functions. All trials of a given stimulus frequency in a single acquisition were modeled with a

single covariate, but different covariates modeled the response in each acquisition (for a total of

8×36 = 288 covariates). The covariates were subject to convolution by a model of the

hemodynamic response function, which itself was under the control of three parameters that

specify the weighted combination of an orthogonal basis (Woolrich et al., 2004). The form of the

HRF was held in common across all acquisitions for a given voxel for a given participant. Within

a non-linear model fitting routine the shape of the HRF and the amplitudes of the covariates

were adjusted to minimize the L2 norm of the difference between the observed and modeled

fMRI time series, after subjecting the model and data to a 0.0387 Hz high-pass filter.

This analysis yielded estimates of the amplitude of neural response to each stimulus

frequency for each acquisition, providing 12 measures in total for each crossing of post-

receptoral channel x stimulus frequency for each participant. For display of the fit of the model

to the time-series data, and for voxel selection, we first regressed out the effect of the attention

task, and then averaged across the six acquisitions for a given post-receptoral direction with

stimulus ordering "A", and the six acquisitions with stimulus ordering "B", and concatenated

these. The R^2 value of the model fit to these average data (which excludes variance attributable

to the attention task) was retained.

Anatomical regions of interest

An anatomical model of the retinotopic organization of V1 cortex was used to define the

location and eccentricity mapping of this cortical area for each participant (Benson et al., 2014).

For the analysis across eccentricity, the V1 ROI was divided into six, log-spaced divisions of

eccentricity (0, 2.8, 5.6, 11.25, 22.5, 45, and 90°). The LGN ROI was defined in CIFTI

FSLR_32k space. We segmented the LGN using the Freesurfer thalamic segmentation

algorithm (Iglesias et al., 2018). A warp was calculated between participant anatomy and the

Montreal Neurological Institute (MNI) space, and the segmentation masks were moved to the

MNI coordinates with nearest neighbor interpolation. The reason for this operation is because

the CIFTI FSLR atlas uses MNI for the subcortical voxel maps. Within a given eccentricity band

(or region mask), voxels with a time-series model fit of $R^2 > 0.1$ were selected and averaged to

obtain temporal sensitivity measurements.

Modeling of the temporal sensitivity function

TSFs were fit with a difference-of-exponentials model (Hawken et al., 1996) developed

from Watson's temporal sensitivity model (Watson, 1986):

264
$$H_{center}(\omega) = (i2\pi\omega\tau_{center} + 1)^{-nc},$$

265
$$H_{surround}(\omega) = G_{surround}(i2\pi\omega\tau_{surround} + 1)^{-ns},$$

266
$$H(\omega) = G(H_{center} - H_{surround})$$

where *nc* and *ns* represent the order of the center (fast) and surround (slow) filters, fixed at 9

and 10, respectively (Watson, 1986). Variables that were free to vary included τ_{center}, the

multiplier of the $\tau_{surround}$ relative to the τ_{center}, and the multiplier of the $H_{surround}$ relative to the

H_{center}. Linear constraints were placed on the model parameters, and an additional non-linear

constraint was added to ensure the model produced a unimodal fit. The amplitudes of BOLD

fMRI response across temporal frequency were fit with this model using the Matlab *fmincon*

routine. While the Watson model provided a good description of the temporal sensitivity

functions in our data, we found the model to be over-parameterized, in that model fits did not

provide unique parameter solutions. Consequently, the values of the parameters were not

especially informative. We instead summarized the model fits by noting the amplitude and

temporal frequency at which the maximum interpolated response was found.

Statistical analysis

Goodness-of-fit for fMRI timeseries modeling was determined by R^2. TSFs are shown with the

median and interquartile range (IQR). Median and IQR amplitude and peak temporal frequency

were determined by fitting 1,000 TSFs bootstrapped across acquisitions under each stimulus

condition. A map of temporal sensitivity was also created. To do so, we fit the Watson temporal

sensitivity function to the response data from each vertex, separately for each stimulus

direction. At each vertex, we found the interpolated temporal frequency corresponding to the

maximum response (expressed as log Hz). The set of frequency values across vertices was

then z-transformed by subtracting the across-voxel mean and then dividing by the standard

deviation. The three z-maps obtained from the three stimulus directions were then averaged to

provide an overall map of relative peak temporal sensitivity. This analysis was conducted only

for the posterior 1/3rd of the cortical surface, as minimal responses were found anterior to this

point.

Model of retinal signals

We created a model of predicted retinal output to our stimuli, based upon previous *in vivo*

measurements in the macaque from three classes of retinal ganglion cells (parasol, midget,

small bistratified). These data were extracted from Figures 1a, b of Solomon and colleagues

(2002) and Figures 9a, b and 13a of Solomon(2005). The electrophysiologic data were recorded

in response to stimuli similar to those employed in our study (spatially uniform, 4.7° diameter

modulations that targeted the DKL cardinal axes around a 2000 Troland background).

Measurements were made for log-spaced temporal frequencies and contrasts from cells in three

different retinal eccentricity bands (0-10°, 20-30°, >30°; only the first two eccentricity bands

available for the bistratified responses). We averaged the "on" and "off" cell responses when

both were provided. Amplitude was given in units of "gain", defined as spikes per second per

percent contrast of the stimulus relative to the maximum possible cone contrast available for a

given stimulus modulation in cone contrast space (measured at the initial slope of the contrast

response function).

To obtain the predicted response to our stimuli, we calculated the spikes / sec response that

would be expected for the contrast level used for our stimuli, with the assumption that parasol

cells saturate at 25% contrast (Lee et al., 1990). The response to the LMS modulation was

taken as the sum of the midget and parasol achromatic responses. We assigned the RGC

measurements to three eccentricity locations within the cell recoding bins (5°, 25°, and 40°). To

produce the predicted retinal response integrated across cell populations, we scaled responses

at each eccentricity by the total quantity of RGC receptive fields (Watson, 2014), the relative

proportions of the different RGC classes (Dacey, 1993; Drasdo et al., 2007), and the change in

annular retinal area with eccentricity. We characterized the model output by the amplitude and

temporal frequency at the peak response.

Code Accessibility

All software used for data processing and data analysis is publicly accessible. The software

used for overall data analysis and figure generation is found here:

https://github.com/gkaguirrelab/Patterson_2023_JNeurosci. The routine used for non-linear

fitting of the BOLD fMRI data may be found here: https://github.com/gkaguirrelab/forwardModel.

The model of retinal response may be found here:

https://github.com/gkaguirrelab/rgcTemporalSensitivity.

Results
328

We obtained BOLD fMRI data from two participants, P1 and P2, while they viewed a

spatially uniform, wide field of flickering light. Our stimuli separately targeted the three post-

receptoral channels (Figure 1a). The design of these stimuli accounted for variation in receptoral

spectral sensitivity across retinal eccentricity (Figure 1b), and were designed to produce

spatially uniform, high contrast on the targeted post-receptoral channel across the visual field

(Figure 1c). Below we refer to these stimuli as being oriented in different "directions" along the

cardinal axes of Derrington-Krauskopf-Lennie (DKL) space (Derrington et al., 1984). Across

many fMRI acquisitions, these spectral modulations were flickered at log-spaced temporal

frequencies ranging from 2 to 64 Hz (Figure 1d). An additional, "0 Hz" condition presented the

steady stimulus background. Participants viewed the stimulus field within an MRI-compatible

eye piece through a +36 or +32 D contact lens, increasing the effective stimulus radius to ~70°

eccentricity.

Figure 1. Targeting the post-receptoral channels with spectral flicker
(a) L-, M-, and S-cones interact at the level of the retina to form the achromatic (L+M), chromatic red-green (L–M),
and chromatic blue-yellow (S–[L+M]) post-receptoral channels. Midget RGCs contribute substantially to both the
achromatic and red-green channels. (b) The spectral sensitivities of the cones vary slightly between central (2
degrees, dotted line) and more peripheral (10 degrees, solid line) eccentricity locations, largely due to the filtering
effects of macular pigment. (c) Spectral modulations were generated to target combinations of cone classes while
silencing others, corresponding to the three cardinal axes of DKL space. Stimuli modulated between a suppressing
(black) and stimulating (red) spectrum, around a common background (dotted gray). Different modulations produced
(*left*) equal nominal stimulation of the L, M, and S cones, thus producing isolated stimulation of the L+M channel,
(*center*) differential stimulation of the L and M cones, while silencing S, and (*right*) isolated stimulation of the S cones,
producing targeted stimulation of the S-[L+M] channel. Nominal Michelson contrast upon the targeted post-receptoral
channels is given in brackets above each plot. Spectral targeting accounted for the 2° and 10° variants of spectral

sensitivity for the cone classes, in an attempt to equate stimulus contrast across the visual field. (d) Temporal flicker
was presented at logarithmically spaced frequencies between 2 Hz and 64 Hz (as well as 0 Hz) in each of many 12
second trials. The modulation was subjected to a half-cosine ramp at onset and offset. The ordering of the stimulus
frequencies was specified by a pseudo-random, counterbalanced sequence. During intermittent attention events the
entire stimulus field dimmed for 250 msecs, in response to which the participant was asked to make a bilateral button
press. Different fMRI acquisitions presented modulations that targeted a different post-receptoral channel.

Variation in flicker frequency induces consistent variation in BOLD fMRI responses

First, we fit the BOLD fMRI data from each participant with a time-series model to

recover the amplitude of response to each stimulus type at each voxel or vertex. The average

amplitude of BOLD fMRI signal was estimated separately for each stimulus direction (LMS, L–

M, S) for each stimulus frequency across the 12 second periods of stimulation; the amplitude of

response to the attention events was also estimated. These estimates were made for each

acquisition to allow assessment of measurement error. Figure 2a presents maps of the R^2 fit of

the time-series model for each participant across the cortical surface. A similar distribution of

high-quality model fits was seen in both participants, largely confined to the early visual areas

on the mesial cortical surface bilaterally. The highest model fit values were found within the

borders of area V1.

Figure 2. Fit to fMRI time series data

The time series data at each voxel and vertex were analyzed for participant P1 (top) and P2 (bottom) across all
acquisitions. (a) The partially inflated, pseudo-hemisphere, cortical surface map for each participant, with the R^2 value
of the model fit to the time series data at each vertex. A black line indicates the border of area V1, within which the
highest model fit values were found. (b) The average BOLD fMRI time series (gray dots) across area V1, and across
the acquisitions for each stimulus type (LMS, L–M, and S). The continuous line shows the fit of the time-series model.
An inset, vertical scale bar at center-left indicates a 2% BOLD fMRI signal response. An inset scale bar at center-right
indicates a 12 second interval. The plot at center provides the ordering of stimulus frequency. Inspection of the time
series reveals both a systematic relationship with the stimulus sequence, and a similar form of response between the
two participants.
383

We obtained the V1 time-series data and model fit (averaged across vertices and

acquisitions) for each participant (after removing the effect of the attention events) for each of

the three stimulus directions (Figure 2b). For both participants, the analysis provided a good

description of the variation in the BOLD fMRI signal. Further, the time-series data were similar

between the two participants, and systematically related to the pattern of stimulus temporal

frequencies illustrated at center. There was clear variation in the amplitude of the cortical

response as a function of the modulation frequency of the stimulus for each post-receptoral

direction.

Temporal filtering and amplification of chromatic signals between LGN and V1

The first level analysis provided the amplitude of BOLD fMRI response to each stimulus

frequency. We next created temporal sensitivity functions (TSFs) which describe the average

response to each stimulation frequency (relative to the 0 Hz condition). Figure 3a presents the

TSFs measured for each participant with the LGN and area V1. The data were fit with the

Watson (1986) difference-of-exponentials temporal model, which provided a reasonable

description. Visual inspection of the TSFs reveals two notable differences between LGN and V1

responses: (1) there was a relative increase in the amplitude of response to chromatic stimuli

between LGN and V1; and (2) the peak temporal sensitivity shifted to lower frequencies

between LGN and V1 across all post-receptoral channels. We quantified these impressions by

obtaining the median (across bootstrapped acquisitions) peak amplitude and peak temporal

sensitivity provided by the fit of the Watson model (Figure 3b). Larger responses were seen in

V1 as compared to the LGN for all three post-receptoral channels, but this was more

pronounced for chromatic stimuli, with an increase that was 2-3 times larger between these

regions for chromatic as compared to achromatic stimuli. There was also a shift of peak

temporal sensitivity towards lower frequencies between LGN and V1 for all three post-receptoral

channels. This was largest for the achromatic stimuli, for which peak sensitivity dropped by 5-10

Hz between the thalamic and cortical sites.

Figure 3. Temporal sensitivity to flicker in LGN and V1. (a) LGN and V1 TSFs for stimuli that target the 3
post-receptoral channels are shown for P1 (*top*) and P2 (*bottom*) fit with the Watson difference-of-exponentials
model. Data points represent the boot-strapped median across acquisitions, and shading provides the interquartile
range. A dashed reference line is positioned at 16 Hz in each plot, facilitating a comparison of the point of peak
temporal sensitivity across plots. (b) Peak amplitude (top) and peak temporal sensitivity (bottom) for each subject
across region. Peak amplitude and peak temporal sensitivity were determined by Watson model fits.

Peak temporal sensitivity in V1 slows towards the periphery

The retinotopic organization of V1 allowed us to examine temporal sensitivity across

eccentricity. BOLD fMRI responses at each vertex within V1 were fit with the Watson model to

obtain the peak amplitude and peak temporal frequency of the TSF. We then examined how

these properties varied across vertices as a function of eccentricity (Figure 4a). The

interpretation of the absolute amplitude of response is complicated by the effect of non-neural

influences upon the BOLD fMRI response; we nonetheless can examine the relative response

between the stimulus directions. The response to the chromatic L–M modulation is comparable

to the LMS-driven response out to ~5° eccentricity, and then drops steadily into the visual

periphery. In contrast, the response to S-directed stimuli across eccentricity follows the same

general form as the response to LMS, although is scaled down and reaches zero close to the

fovea.

We examined peak temporal sensitivity across post-receptoral channel and eccentricity.

Across eccentricity, higher peak temporal frequency values were generally found for the

achromatic as compared to the chromatic stimuli. For the chromatic stimuli, there was only a

small change in peak temporal frequency with eccentricity. In contrast, the peak response to

LMS stimuli steadily shifted to lower frequencies starting at 10 – 20° eccentricity, eventually

reaching the temporal sensitivity seen in response to L–M stimuli. Across the three post-

receptoral directions, we found no evidence of a shift to higher temporal sensitivity with greater

eccentricity.

Next, we divided area V1 into 6 concentric bands to visualize TSFs as a function of

visual field eccentricity. To account for cortical magnification of the visual field, the borders of

these bands were logarithmically spaced, yielding patches of cortex roughly matched in surface

area. Within each band, we obtained the TSF for each stimulus modulation (Figure 4b).

Generally, reliable TSFs were measured for both participants, for all three stimulus types,

across eccentricity. This includes in the most peripheral eccentricity band (centered at 64°).

Examination of these TSFs confirms the impression that the peak response to LMS stimuli shifts

to lower temporal frequencies at greater eccentricities within area V1. This is evident both in the

peak temporal frequency, as well as in a relatively larger low-pass component to the response.

Figure 4. Temporal flicker sensitivity within area V1 across eccentricity. (a) Peak amplitude (right) and peak temporal frequency (left) within area V1 for P1 and P2 for stimuli that targeted each of the 3 post-receptoral directions. Dots represent values for each vertex and lines represent the fit of a smoothing spline to the median value at each of 30 eccentricity bins. (a) V1 TSFs at logarithmically spaced eccentricities across the 3 post-receptoral channels are shown for P1 (*left*) and P2 (*right*) with Watson temporal model fits. Data points represent the boot-strapped median across acquisitions, and shading provides the interquartile range.

Peak temporal sensitivity increases beyond area V1

We created maps of peak temporal frequency across the cortex for each stimulus direction

(Supplementary Figure 2). Although these maps differed in their absolute values, we found that

the patterns of higher and lower temporal sensitivity were quite similar across the three stimulus

directions. Therefore, we standardized the maps across vertices for each stimulus type and

combined them (Figure 5a). A similar pattern of response was seen for both participants. Within

area V1, there was a relatively uniform peak sensitivity to mid-range temporal frequencies.

Within posterior extra-striate cortex, peak sensitivity shifted to higher temporal frequencies; this

was the case for both dorsal and ventral visual areas. Finally, at more anterior points within

visually responsive cortex, peak sensitivity was found at lower temporal frequencies.

We measured the peak temporal sensitivity of specific extra-striate regions (Figure 5b). For both

participants across all three stimulus directions, peak temporal sensitivity increased across the

visual hierarchy, rising between V1 and V2/V3, and again into both MT and hV4. For achromatic

stimuli, MT demonstrated the highest peak temporal sensitivity; this pattern was not reliably

seen in response to the chromatic stimuli.

Figure 5. Peak temporal frequency beyond area V1. (a) Relative peak temporal sensitivity map across
visual cortex, combined across stimulus directions for the two participants. Lateral, medial, and ventral views are
provided. Map values reflect the relative peak frequency across the cortex within each of the stimulus directions. (b)
Median temporal sensitivity (boot-strapped across acquisitions) for each of several visual cortex regions of interest.
Error bars provide the interquartile range.
479

Comparison of V1 responses to predicted retinal signals

To better understand the transformation of signals from the retina to the cortex, we generated a

prediction of integrated retinal output to our stimuli, based upon prior measurements of

macaque RGC responses (Solomon et al., 2002; Solomon, 2005). In Figure 6, the filled circles

and lines correspond to the predicted retinal response, while the squares and triangles are the

data measured from our two participants within area V1 across eccentricity.

Figure 6. Predicted retinal signals compared to V1 response. Peak amplitude (left) and peak temporal
frequency (right) of the V1 fMRI data from this study (squares and triangles) compared to predicted retinal
output (filled circles) based upon physiologic recordings of macaque RGCs, integrated across the total
cell population present at a given eccentricity location. Peak amplitude is expressed relative to responses
to LMS-directed stimuli. For the retina model, normalization is to the average response evoked by LMS
stimuli across eccentricity; for the V1 data, normalization is performed at each of 30 eccentricity bins.

The predicted amplitude of retinal response to LMS stimulation varies only a small amount

among the three eccentricity locations for which measurements are available. This is in contrast

to the response to L–M chromatic stimulation, which is initially greater than the response to

LMS, but drops by an order of magnitude into the visual periphery, as the combined

consequence of both a relative reduction in the midget RGC class and the reduced response

per cell, perhaps due to the effects of random cone wiring (Wool et al., 2018). Finally, the

predicted response to S-directed stimulation is markedly smaller due to the sparse population of

small bistratified cells that are presumed to carry this signal.

We may compare these model values to the fMRI data, normalized at each eccentricity location

by the response to LMS stimulation. Similar to the retinal prediction, the relative response to L–

M stimulation drops steadily beyond 5-10° eccentricity, although the V1 response is smaller than

might be predicted in central vision. The amplitude of response to S-directed stimuli within V1 is

markedly larger than predicted by the retinal model, rising futher towards 5-10° eccentricity, then

falling in concert with the response to L–M.

The retinal model predicts the temporal frequency that would elicit the peak response for each

post-receptoral direction. At 5° eccentricity, there was rough agreement between the retinal

model and the fMRI data. At greater eccentricities, however, retinal responses to LMS and L–M

stimuli were predicted to shift to ever higher peak temporal frequencies, in marked contrast to

the properties of the fMRI data.

Discussion

Our study provides detailed temporal sensitivity measurements across subcortical and

cortical regions in response to supra-threshold stimulation of the three canonical, post-

receptoral visual channels. These results also characterize a fundamental aspect of human

cortical visual function across eccentricity. Many of our findings replicate known properties of

the visual system, including a relative amplification of chromatic signals between the LGN and

area V1 (Cottaris and De Valois, 1998; Mullen et al., 2008), filtering of high temporal

frequencies between these neural stages (Horwitz, 2021), attenuated responses to S-cone

stimulation at the fovea (Williams et al., 1981), a systematic difference between chromatic and

achromatic channels in peak temporal sensitivity (Swanson et al., 1987; Spitschan et al., 2016),

and faster peak temporal sensitivity in extra-striate cortex (Spitschan et al., 2016). Contrary to

what might be expected from the properties of the retina and threshold psychophysical

measurements, however, we do not observe an increase in peak temporal sensitivity to

achromatic stimuli in the visual periphery. Unexpectedly, we find a shift in these cortical

responses to lower temporal frequencies.

Comparison to prior neuro-imaging studies

There have been prior studies of the temporal sensitivity for achromatic and chromatic

modulations within primary visual cortex (Engel et al., 1997; D'Souza et al., 2011) and extra-

striate cortex (Spitschan et al., 2016), but these experiments did not examine responses across

eccentricity. Chai and colleagues (2019) measured BOLD fMRI responses to a spatially

structured, 15° radial eccentricity, achromatic checkerboard pattern that flickered between 1 to

40 Hz. Their parcellation of cortex by temporal sensitivity is similar in appearance to our

temporal sensitivity map (Figure 5a). Kim and colleagues (Kim et al., 2023) developed a spatio-

temporal receptive field model and analyzed fMRI data collected in response to drifting bars that

contained multi-colored images. They find that a temporal integration parameter increases

across the first 12 degrees of V1 eccentricity, although there is not a straightforward relationship

between their model and our measurements.

Himmelberg and Wade (2019) reported that V1 response contrast sensitivity for a 20 Hz

achromatic grating was greater at 6-10° eccentricity as compared to the fovea. As we did not

measure a neural contrast response function for our stimuli, we are not able to directly compare

this prior result to our own. Additionally, temporal sensitivity varies substantially with luminance

and spatial frequency (Watson, 1986), and these stimulus properties differ between the current

study and that of Himmelberg and Wade. We measured responses to a spatially uniform field at

high retinal illuminance and took advantage of tailored spectral design to equate stimulus

contrast across eccentricity. An advantage of this approach is that we do not need to contend

with the effect of variation of acuity across the visual field (Virsu et al., 1982).

Normalization of retinal signals in the visual cortex

Comparison of our measurements from area V1 to a predicted retinal response suggests

that there is extensive post-retinal modification of visual signals across eccentricity. Variation in

response amplitude in V1 across stimulus directions is a log unit less than predicted to arise in

the retina. There is also a notable difference between predicted and observed temporal

sensitivity with eccentricity: while the visual cortex data at 5° eccentricity is consistent with the

retinal prediction, at further eccentricities cortical responses peak at far lower stimulation

frequencies than predicted. It appears that post-retinal modification between the retina and V1

normalizes responses across eccentricity and post-receptoral channel, modulating the relative

amplitude of chromatic responses and slowing peak temporal sensitivity towards the periphery.

Overall, visual cortex responses substantially reduce the striking differences across post-

receptoral channel and eccentricity observed in the retina.

We might consider alternative explanations for the relative stability of temporal frequency

sensitivity that we observe across eccentricity in area V1. Our stimuli were wide-field and

spatially uniform. We therefore derived temporal sensitivity measures from simultaneously

stimulated eccentricities. It is possible that lateral or hierarchical influences between neurons

across eccentricity locations cause homogenization of responses (Nassi et al., 2014).

Alternatively, our stimulus might have been ineffective at driving neurons that have both

narrower spatial tuning, and which vary in temporal tuning across eccentricity (Reinhard and

Münch, 2021). In either case, measurements made with spatially restricted stimuli that produce

isolated stimulation at different eccentricities might produce different findings.

We designed our stimulus to be spatially uniform, including accounting for variation in

spectral sensitivity of the cone fundamentals with eccentricity (Supplementary Figure 1).

Nonetheless, we may be concerned that stimulus contrast or background luminance declined

into the periphery, causing a decrease in temporal sensitivity (Snowden et al., 1995). The

absolute amplitude of BOLD fMRI response measured for all three stimulus directions

decreased beyond ~10° eccentricity, potentially supporting this concern. We are reassured,

however, that this effect is unlikely to account for the slowing of temporal sensitivity for the

luminance channel, as the amplitude of response in the far periphery is equal or greater than

found at the fovea, despite a slowing of the TSF. In future studies, measurement of temporal

sensitivity at varying contrast levels could be used to better address this concern, and

potentially disambiguate the effects of non-neural influences upon the BOLD fMRI response

(Kurzawski et al., 2021).

Relation to psychophysical measures

The cortical temporal sensitivity functions we have obtained here are in good agreement

with known properties of human perception. This includes the both the general form of the

responses—which are well-fit by the Watson difference-of-exponentials model—and the

variation in peak temporal sensitivity between the chromatic and achromatic channels.

Perceptual temporal sensitivity across visual field eccentricity has been well studied for

the detection of high-frequency achromatic flicker. Specifically, the critical flicker fusion (CFF)

frequency is a measure of highest frequency stimulus (in the presence of noise) that can be

distinguished from a static background. There is evidence that people are able to perceive

higher frequency achromatic flicker in the periphery of their visual field (Hartmann et al., 1979;

Tyler and Hamer, 1993), although it is unclear if this reflects a change in the shape of the entire

sensitivity function or variation in the signal:noise ratio with eccentricity (Rider et al., 2021).

While we might consider measuring an analog of the CFF in our BOLD fMRI data, this

analysis faces practical limitations. For almost all the measurements across eccentricity, the

BOLD fMRI response to 32 Hz LMS flicker is far above threshold, while the response at 64 Hz is

zero; the cortical CFF lies somewhere between these two frequencies. Therefore, we regard our

data as not able address this question. Anecdotally, however, we note that one observer (P1)

perceived a faint, shimmering flicker in an annular zone of their peripheral vision during

presentation of the 64 Hz LMS stimulus. This participant was also found to have a non-zero

cortical response to this same stimulus, only in the 16° eccentricity band. The second observer

did not report this percept and did not have measurable BOLD fMRI responses to 64 Hz flicker

within the V1 eccentricity bands.

Our study measured the supra-threshold cortical temporal sensitivity function. A study of

supra-threshold temporal frequency discrimination using achromatic flicker found that

performance at the fovea and at 30° eccentricity was quite similar (Waugh and Hess, 1994),

perhaps consistent with our finding here that the supra-threshold temporal sensitivity of visual

cortex changes minimally across eccentricity.

Conclusions

This study provides the eccentricity dependence of the temporal sensitivity functions for

the three cardinal post-receptoral channels within V1, and the spatially averaged sensitivity

functions across multiple levels of the visual hierarchy. After the LGN, peak temporal sensitivity

increases across the visual cortical hierarchy. We find that responses in primary visual cortex

across post-receptoral channel and temporal frequency are normalized relative to predicted

signals from the retina. This includes an unexpected slowing of cortical temporal sensitivity for

achromatic flicker in the visual periphery.

Author Contributions:

Conceptualization: Manuel Spitschan, Andrew S. Bock, and Geoffrey K. Aguirre. Data curation:

Huseyin O. Taskin. Formal analysis: Carlyn Patterson Gentile, Manuel Spitschan, and Geoffrey

K. Aguirre. Funding acquisition: Geoffrey K. Aguirre. Investigation: Andrew S. Bock and

Geoffrey K. Aguirre. Methodology: Geoffrey K. Aguirre. Project administration: Geoffrey K.

Aguirre. Resources: Geoffrey K. Aguirre. Software: Carlyn Patterson Gentile, Manuel Spitschan,

Huseyin O. Taskin, and Geoffrey K. Aguirre. Supervision: Geoffrey K. Aguirre. Validation:

Manuel Spitschan and Geoffrey K. Aguirre. Visualization: Carlyn Patterson Gentile, Manuel

Spitschan, and Geoffrey K. Aguirre. Writing - original draft: Carlyn Patterson Gentile and

Geoffrey K. Aguirre. Writing - review & editing: Carlyn Patterson Gentile, Manuel Spitschan,

Huseyin O. Taskin, Andrew S. Bock, and Geoffrey K. Aguirre.

Study Funding: Supported by the Research to Prevent Blindness, Lions Club International

Foundation Low Vision Award to GKA; the US-Israel Binational Science Foundation to GKA; the

National Institute of Neurological Disorders and Stroke NIH K23NS124986 to CPG; and P30

EY001583: Core Grant for Vision Research.

References:

(CIE) CI de L (2006) Fundamental chromaticity diagram with physiological axes, Part 1. Vienna.

Avants B, Epstein C, Grossman M, Gee J (2008) Symmetric diffeomorphic image registration with cross-correlation: Evaluating automated labeling of elderly and neurodegenerative brain. Med Image Anal 12:26–41.

Barnett MA, Aguirre GK, Brainard D (2021) A quadratic model captures the human V1 response to variations in chromatic direction and contrast. Elife 10:e65590.

Benson NC, Butt OH, Brainard DH, Aguirre GK (2014) Correction of Distortion in Flattened Representations of the Cortical Surface Allows Prediction of V1-V3 Functional Organization from Anatomy. PLoS Comput Biol 10:e1003538.

Chai Y, Handwerker DA, Marrett S, Gonzalez-Castillo J, Merriam EP, Hall A, Molfese PJ, Bandettini PA (2019) Visual temporal frequency preference shows a distinct cortical architecture using fMRI. Neuroimage 197:13–23.

Cottaris NP, De Valois RL (1998) Temporal dynamics of chromatic tuning in macaque primary visual cortex. Nature 395:896–900.

Cox RW (1996) AFNI: Software for Analysis and Visualization of Functional Magnetic Resonance Neuroimages. Comput Biomed Res 29:162–173.

D'Souza D V., Auer T, Strasburger H, Frahm J, Lee BB (2011) Temporal frequency and chromatic processing in humans: An fMRI study of the cortical visual areas. J Vis 11:8–8.

Dacey DM (1993) Morphology of a small-field bistratified ganglion cell type in the macaque and human retina. Vis Neurosci 10:1081–1098.

Dale AM, Fischl B, Sereno MI (1999) Cortical Surface-Based Analysis. Neuroimage 9:179–194.

Derrington AM, Krauskopf J, Lennie P (1984) Chromatic mechanisms in lateral geniculate nucleus of macaque. J Physiol 357:241–265.

Drasdo N, Millican CL, Katholi CR, Curcio CA (2007) The length of Henle fibers in the human retina and a model of ganglion receptive field density in the visual field. Vision Res 47:2901–2911.

Esteban O, Markiewicz CJ, Blair RW, Moodie CA, Isik AI, Erramuzpe A, Kent JD, Goncalves M, DuPre E, Snyder M, Oya H, Ghosh SS, Wright J, Durnez J, Poldrack RA, Gorgolewski KJ (2019) fMRIPrep: a robust preprocessing pipeline for functional MRI. Nat Methods 16:111–116.

Estévez O, Spekreijse H (1982) The "silent substitution" method in visual research. Vision Res 22:681–691.

Gelfand EC, Horwitz GD (2018) Model of parafoveal chromatic and luminance temporal contrast sensitivity of humans and monkeys. J Vis 18:1.

Glasser MF, Sotiropoulos SN, Wilson JA, Coalson TS, Fischl B, Andersson JL, Xu J, Jbabdi S, Webster M, Polimeni JR, Van Essen DC, Jenkinson M (2013) The minimal preprocessing pipelines for the Human Connectome Project. Neuroimage 80:105–124.

Hartmann E, Lachenmayr B, Brettel H (1979) The peripheral critical flicker frequency. Vision Res 19:1019–1023.

Hawken MJ, Shapley RM, Grosof DH (1996) Temporal-frequency selectivity in monkey visual cortex. Vis Neurosci 13:477–492.

Himmelberg MM, Wade AR (2019) Eccentricity-dependent temporal contrast tuning in human visual cortex measured with fMRI. Neuroimage 184:462–474.

Horwitz GD (2020) Temporal information loss in the macaque early visual system Demb J, ed. PLOS Biol 18:e3000570.

Horwitz GD (2021) Temporal filtering of luminance and chromaticity in macaque visual cortex. iScience 24:102536.

Huntenburg J (2014) Evaluating nonlinear coregistration of BOLD EPI and T1w images. Freie Univ Berlin.

Jenkinson M, Bannister P, Brady M, Smith S (2002) Improved Optimization for the Robust and

Accurate Linear Registration and Motion Correction of Brain Images. Neuroimage 17:825–
841.
Kim I, Kupers ER, Lerma-Usabiaga G, Grill-Spector K (2023) Characterizing spatiotemporal
population receptive fields in human visual cortex with fMRI. bioRxiv:585539.
Kurzawski JW, Gulban OF, Jamison K, Winawer J, Kay K (2021) The influence of non-neural
factors on BOLD signal magnitude. bioRxiv.
Lee BB, Pokorny J, Martin PR, Valbergt A, Smith VC (1990) Luminance and chromatic
modulation sensitivity of macaque ganglion cells and human observers. J Opt Soc Am A
7:2223.
Mullen KT, Dumoulin SO, Hess RF (2008) Color responses of the human lateral geniculate
nucleus: selective amplification of S-cone signals between the lateral geniculate nucleno
and primary visual cortex measured with high-field fMRI. Eur J Neurosci 28:1911–1923.
Nassi JJ, Gómez-Laberge C, Kreiman G, Born RT (2014) Corticocortical feedback increases
the spatial extent of normalization. Front Syst Neurosci 8.
Pruim RHR, Mennes M, van Rooij D, Llera A, Buitelaar JK, Beckmann CF (2015) ICA-AROMA:
A robust ICA-based strategy for removing motion artifacts from fMRI data. Neuroimage
112:267–277.
Reinhard K, Münch TA (2021) Visual properties of human retinal ganglion cells Barnes S, ed.
PLoS One 16:e0246952.
Rider AT, Henning GB, Stockman A (2021) A reinterpretation of critical flicker-frequency (CFF)
data reveals key details about light adaptation and normal and abnormal visual processing.
Prog Retin Eye Res:101001.
Seiple W, Holopigian K (1996) Outer-retina locus of increased flicker sensitivity of the peripheral
retina. J Opt Soc Am A 13:658.
Snowden RJ, Hess RF (1992) Temporal frequency filters in the human peripheral visual field.
Vision Res 32:61–72.
Snowden RJ, Hess RF, Waugh SJ (1995) The processing of temporal modulation at different
levels of retinal illuminance. Vision Res 35:775–789.
Solomon SG (2005) Chromatic Organization of Ganglion Cell Receptive Fields in the Peripheral
Retina. J Neurosci 25:4527–4539.
Solomon SG, Martin PR, White AJ., Rüttiger L, Lee BB (2002) Modulation sensitivity of ganglion
cells in peripheral retina of macaque. Vision Res 42:2893–2898.
Spitschan M, Aguirre GK, Brainard DH (2015) Selective stimulation of penumbral cones reveals
perception in the shadow of retinal blood vessels. PLoS One 10:e0124328.
Spitschan M, Datta R, Stern AM, Brainard DH, Aguirre GK (2016) Human Visual Cortex
Responses to Rapid Cone and Melanopsin-Directed Flicker. J Neurosci 36:1471–1482.
Spitschan M, Woelders T (2018) The Method of Silent Substitution for Examining Melanopsin
Contributions to Pupil Control. Front Neurol 9:941.
Swanson WH, Ueno T, Smith VC, Pokorny J (1987) Temporal modulation sensitivity and pulse-
detection thresholds for chromatic and luminance perturbations. J Opt Soc Am A 4:1992.
Thibos LN, Still DL, Bradley A (1996) Characterization of spatial aliasing and contrast sensitivity
in peripheral vision. Vision Res 36:249–258.
Treiber JM, White NS, Steed TC, Bartsch H, Holland D, Farid N, McDonald CR, Carter BS, Dale
AM, Chen CC (2016) Characterization and Correction of Geometric Distortions in 814
Diffusion Weighted Images Najbauer J, ed. PLoS One 11:e0152472.
Tustison NJ, Avants BB, Cook PA, Yuanjie Zheng, Egan A, Yushkevich PA, Gee JC (2010)
N4ITK: Improved N3 Bias Correction. IEEE Trans Med Imaging 29:1310–1320.
Tyler CW, Hamer RD (1993) Eccentricity and the Ferry-Porter law. J Opt Soc Am A 10:2084–
2087.
Virsu V, Rovamo J, Laurinen P, Näsänen R (1982) Temporal contrast sensitivity and cortical
magnification. Vision Res 22:1211–1217.

Wang S, Peterson DJ, Gatenby JC, Li W, Grabowski TJ, Madhyastha TM (2017) Evaluation of
Field Map and Nonlinear Registration Methods for Correction of Susceptibility Artifacts in
Diffusion MRI. Front Neuroinform 11.
Watson AB (1986) Temporal Sensitivity. In: Handbook of Perception and Human Performance,
Vol. 1: Sensory Processes and Perception.
Watson AB (2014) A formula for human retinal ganglion cell receptive field density as a function
of visual field location. J Vis 14:15.
Waugh SJ, Hess RF (1994) Suprathreshold temporal-frequency discrimination in the fovea and
the periphery. J Opt Soc Am A 11:1199.
Williams DR, MacLeod DIA, Hayhoe MM (1981) Punctate sensitivity of the blue-sensitive
mechanism. Vision Res 21:1357–1375.
Wool LE, Crook JD, Troy JB, Packer OS, Zaidi Q, Dacey DM (2018) Nonselective Wiring
Accounts for Red-Green Opponency in Midget Ganglion Cells of the Primate Retina. J
Neurosci 38:1520–1540.
Woolrich MW, Behrens TEJ, Smith SM (2004) Constrained linear basis sets for HRF modelling
using Variational Bayes. Neuroimage 21:1748–1761.
Zhang Y, Brady M, Smith S (2001) Segmentation of brain MR images through a hidden Markov
random field model and the expectation-maximization algorithm. IEEE Trans Med Imaging
20:45–57.
760
761