Identifying novel links between cardiovascular disease and insomnia by

***Drosophila* modeling of genes from a pleiotropic GWAS locus**

Farah Abou Daya[1], Torrey Mandigo[2,3,4], Lily Ober[1], Dev Patel[1], Matthew Maher[2], Suraj

Math[2,3], Cynthia Tchio[2], James A. Walker[2,3,4], Richa Saxena[2,4,5], Girish C. Melkani[1*]

[1]Department of Pathology, Division of Molecular and Cellular Pathology, Heersink

School of Medicine, The University of Alabama at Birmingham, AL 35294, USA

[2]Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA

[3]Department of Neurology, Massachusetts General Hospital, Harvard Medical School,

Boston, USA

[4]Cancer Program, Broad Institute of MIT and Harvard, Cambridge, USA

[5]Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General

Hospital and Harvard Medical School, Boston, MA, USA

[*]Correspondence: Department of Pathology, Division of Molecular and Cellular

Pathology, School of Medicine, University of Alabama at Birmingham, AL 35294, USA.

Tel.: 1-205-996-0591; Fax: 1-205-934-7447; E-mail: girishmelkani@uabmc.edu

Abstract:

Insomnia symptoms have been associated with cardiovascular disease (CVD), doubling

the risk of incident CVD, but specific shared pathways remain poorly understood.

Recently, genome-wide association studies (GWAS) identified genetic loci significantly

associated with insomnia symptoms, including one locus

(near *ATP5G1*, *UBE2Z*, *SNF8*, *IGF2BP1*, and *GIP*) that was previously linked with CVD

in an independent GWAS. To evaluate the cell-autonomous role of genes within the

17q21 insomnia and CVD locus, we used *Drosophila melanogaster* models to perform

tissue-specific RNAi knockdown of four conserved orthologues (*ATPSynC, Lsn, Bruce*,

and *Imp*) in neurons and in the heart. To identify non-cell-autonomous mechanisms, we

also assessed heart function in flies with neuronal-specific knockdown and sleep in flies

with heart-specific knockdown. Neuronal and cardiac-specific RNAi knockdown of

several of the genes conserved in *Drosophila* led to compromised sleep quality and

impaired cardiac performance. Neuronal-specific knockdown of *ATPSynC, Imp, and*

Lsn led to disruptions in sleep quantity and quality. Knockdown of *ATPSynC and Lsn in*

the heart led to significantly reduced cardiac performance without and with cardiac

dilation, respectively. Furthermore, *Lsn* and *ATPSynC*-suppressed hearts showed

disruption in the actin-containing myofibrillar organization and led to a significantly

shortened lifespan. Non-cell-autonomous effects were seen both from neurons to heart

(*Imp*), and heart to neurons (*ATPSynC and Lsn*). Specifically, *Imp* neuronal knockdown

led to a significantly compromised cardiac function, whereas knockdown

of *ATPSynC* and *Lsn* in the heart led to compromised sleep characterized by increased

sleep fragmentation, both accompanied by an increase in inflammation through Upd3,

an inflammatory cytokine, in the heart or head, respectively. We also demonstrate

disrupted cardiac function or sleep upon cardiac-specific or neuronal-specific

overexpression of Upd3, respectively, showing a direct link between cardiac dysfunction

and sleep disruption through inflammation. Our study reveals tissue-specific and cross-

tissue consequences of Drosophila knockdown of multiple genes at this locus, providing

novel insights into potential genetic mechanisms linking CVD and insomnia. Our study

also highlights the key role of these four conserved genes in both sleep and cardiac

function.

Key Words: Cardiovascular Disease, Insomnia, GWAS, Genetic Loci, inflammation,

rs4643373 locus, Drosophila Model

Nonstandard Abbreviations and Acronyms:

CVD: Cardiovascular disease

GWAS: Genome-Wide Association Study

SNP: Single-Nucleotide Polymorphism

ESCRT: Endosomal Sorting Complexes Required for Transport

HP: Heart Period

AI: Arrhythmia Index

DI: Diastolic Interval

SI: Systolic Interval

DD: Diastolic Diameter

SD: Systolic Diameter

FS: Fractional Shortening

Introduction:

Cardiovascular disease (CVD), one of the leading causes of death worldwide,

encompasses several conditions that affect heart structure and function.[1-3] The

incidence of CVD continues to rise, with approximately 18.2 million deaths worldwide in

2019, which contributes to rising healthcare costs and creates a significant

socioeconomic burden.[3,4] Many factors that increase the risk of CVD include genetic

factors, smoking, and lack of physical activity. One important risk factor for CVD that

has recently emerged is sleep dysfunction including insomnia.[4-7] Insomnia is the most

common sleep disorder, affecting 10 to 30% of the population, and is defined as

persistent difficulty in falling and/or staying asleep or non-restorative sleep resulting in

daytime sleepiness, fatigue, or dysfunction.[8-10] Studies suggest that insomnia has a

genetic component with heritability estimates ranging between 22 and 25% in

adults[11,12], and multiple GWASs have identified genetic loci with links to insomnia.[12-15]

Although genetic factors have been identified as contributors to CVD and insomnia, the

genetic mechanisms underlying these two diseases remain poorly understood.

Observational studies have demonstrated that insomnia increases the risk of several disorders, especially CVD.[16-19] Moreover, mendelian randomization analyses show that insomnia symptoms double the risk for incident CVD.[12,13] Similarly, cardiac dysfunction has been associated with sleep disruptions.[20,21] Furthermore, a recent study found that sleep modifies atherosclerosis through hematopoiesis in mice.[22] Together these findings establish a clear link between cardiovascular traits and sleep disruptions. However, the specific underlying causal genetic pathways and mechanisms connecting CVD and insomnia are unknown. Recent genome-wide association studies (GWASs) identified multiple significant loci for self-reported insomnia symptoms in UK Biobank and 23andMe participants.[12,13] From these loci, we identified a single locus, represented by lead SNP rs4643373, that has also been previously associated with coronary artery disease (CAD), and other cardiac disorders, including myocardial infarction.[23-27] This locus provides a valuable opportunity to identify genes important for CVD and/or insomnia, and dissect potential genetic mechanisms underlying the link between cardiovascular function and sleep. Near this locus, we identified five candidate genes, *ATP5G1*, *UBE2Z*, *SNF8*, *IGF2BP1*, and *GIP*. The known functions of these genes are very diverse, including energy metabolism (ATP5G1), protein ubiquitination (UBE2Z), multivesicular body biogenesis (SNF8), post-transcriptional regulation (IGF2BP1), and lipid metabolism (GIP).[28-32] However, it remains unclear which of these genes, if any, contribute to CVD or insomnia.

To elucidate the impact of these candidate genes on the regulation of cardiac function and sleep, we identified conserved *Drosophila melanogaster* orthologs for the

insomnia and CVD-related candidate genes near rs4643373. *Drosophila* genes, namely

ATPSynC (CG1746), Bruce (CG6303), Lsn (CG6637) and Imp (CG1691), were

identified as orthologs of *ATP5G1*, *UBE2Z*, *SNF8*, and *IGF2BP1,* respectively, however,

GIP doesn't have an ortholog. *Drosophila* has become a well-established model system

for studying both CVD and sleep disturbances.[33-38] The fly heart displays many

developmental and functional similarities to the mammalian heart.[33,39] Moreover, several

genes causing heart disease in humans are present in *Drosophila* and play similar

pathophysiological roles, and the manipulation of these genes in *Drosophila* leads to

disease phenotypes similar to humans.[33,39,40] Sleep in flies has also been demonstrated

to share many characteristics with human sleep, such as consolidation during the night

and similar responses to sleep altering drugs.[34,41-43] Therefore, studies investigating the

role of human-relevant *Drosophila* orthologs in the regulation of cardiovascular function

and sleep provide an efficient means to identify new causal genes related to CVD

and/or insomnia, and understand mechanisms relating both diseases to identify

potential future therapeutic targets.

Here, we have evaluated the cardiac and sleep roles of *Drosophila* genes

ATPSynC, Bruce, Lsn and Imp, in both, cell-autonomous and non-cell-autonomous

manners. To assess the role of these genes in cardiac and sleep physiology, we

performed tissue specific knockdown (KD) in the heart and nervous system,

respectively. Cardiac- and neuronal-specific KD of these genes led to cardiac and sleep

dysfunction, suggesting tissue-specific functions related to each disease. After

characterizing the cell-autonomous role of *ATPSynC*, *Bruce*, *Lsn*, and *Imp* in cardiac

function and sleep, we also identified non-cell-autonomous effects of these genes on

cardiac and sleep phenotypes, upon KD of these genes neuronally or in cardiac tissue,

respectively. Neuronal KD of *Imp* compromised cardiac function, while cardiac KD of

ATPSynC, Lsn and Bruce led to sleep dysfunction in a non-cell-autonomous manner.

We further identified inflammation as an underlying mechanism involved in the non-cell-

autonomous effects of cardiac dysfunction on sleep disruption and vice versa. In

conclusion, we were able to uncover novel genetic mechanisms with cell-autonomous

effects on the regulation of cardiac function and sleep as well as non-cell autonomous

genetic mechanisms linking cardiac function in the regulation of sleep and sleep on

cardiac function. Taken together, our data are among the first functional genetic proofs,

to our knowledge, that link CVD with sleep disorders and provide mechanistic insight

into potential therapeutic targets to prevent or attenuate both diseases.

Results

Shared GWAS locus at 17q21 is associated with CAD/CVD and insomnia.

A GWAS signal at 17q21.32 for CAD including myocardial infarction, percutaneous

transluminal coronary angioplasty, coronary artery bypass grafting, angina or chronic

ischemic heart disease from the CARDIoGRAMplusC4D Consortium, represented by

lead SNP rs4643373 (Fig. 1A, n=181,522 cases and 984,079 controls; OR=1.04 (95%

CI=1.028-1.050))[44] was found to co-localize with an association signal for insomnia

symptoms (n=593,724 cases and 1,771,286 controls; OR=1.04 (95% CI=1.025-1.046);

Fig. 1B; pp = 0.95 that both traits share the same causal SNP)[45]. The genomic region

around the co-localized signal encompasses multiple genes including *ATP5MC1,*

UBE2Z, SNF8, GIP and IGF2BP1. Multi-tissue eQTL analyses shows an effect of lead

SNP rs4643373 on the expression of these genes in multiple tissues including the left

ventricle of the heart and the hypothalamus (Table S1). Moreover, to check for other

CVD-related signals in the region of rs4643373, we catalogued cardiovascular and

cardiometabolic trait associations at genome-wide significance ($p<5\times10^{-8}$) of this region

from literature and the Cardiovascular Disease Knowledge Portal (see methods). We

found a total of 9 significant associations with cardiovascular and cardiometabolic traits,

including myocardial infarction, diastolic blood pressure, and triglycerides (Fig. 1C)

reinforcing the importance of this genomic region in CVD. The causal genes and

variants at this locus are unknown. Furthermore, it is unclear if the association signals

reflect independent contribution of effector genes to sleep and cardiovascular disease,

or if effector genes influence sleep through cardiovascular dysfunction or vice versa.

Thus, we developed a *Drosophila* screening approach to identify the role of fly orthologs

of these genes in sleep and cardiovascular function in both cell-autonomous and non-

cell-autonomous manners (Fig. 1D).

Neuronal-specific suppression of CVD- and insomnia- related genes leads to

altered sleep phenotypes along with enhanced sleep fragmentation.

Fly orthologs of four of the five human genes near rs4643373 with their % similarity are

shown in Table 1. The fifth gene, GIP, lacks a fly ortholog. In order to assess whether

any of the orthologs of the genes within the CVD- and insomnia-associated locus were

essential in *Drosophila*, we performed ubiquitous KD of these genes driven by *Act5C-*

Gal4 driver*.* Ubiquitous KD of *ATPSynC* and *Lsn* led to lethality, while that of *Bruce* and

Imp did not affect viability. We then performed a pan-muscle *KD* using the *24b-Gal4*

driver to determine if the function of these genes was essential in all muscle tissues. As

with ubiquitous KD, we found that pan-muscle KD of *ATPSynC* and *Lsn* resulted in

lethality, whereas *Bruce* and *Imp* were viable. We also tested viability with pan-neuronal

KD using the *Elav-Gal4* driver. Only *ATPSynC* Line 1 (see methods) led to lethality

when suppressed pan-neuronally, suggesting an essential role for *ATPSynC* in

neuronal function.

To test the impact of these 4 genes near the CVD- and insomnia- associated locus on

sleep, we used the neuronal-specific *Elav-Gal4* driver to KD gene expression. Level of

KD of each gene in the head is shown in Fig. S1A-D. We quantified total, day and night

measures of sleep, locomotor activity, sleep fragmentation, sleep bout length and sleep

bout number of three-to-four-day-old male flies, averaged over a 5-day period, with pan-

neuronal *Elav* KD of each of the four *Drosophila* orthologs.

Since KD of *ATPSynC* using RNAi Line 1 caused lethality, we used another available

line with efficient KD when driven by *Elav-Gal4* (Fig. S1A, see methods). Compared to

driver and UAS control flies, RNAi-mediated inhibition of *ATPSynC* significantly

increased overall sleep duration which primarily resulted from an increase in nighttime

sleep (Fig. 2A-D, Table S2). This increased sleep corresponded to a decrease in overall

locomotor activity (Fig. 2E, Table S2). KD of *Imp* also showed increased sleep which

was primarily due to increased daytime sleep (Fig. 2A-D, Table S2) with decreased

activity (Fig. 2E, Table S2). Even though the suppression of *Lsn* showed a significant

increase in nighttime sleep (Fig. 2A-D), it was not significantly different than its UAS

control (Table S2). However, neuronal KD of *Bruce* decreased daytime sleep but had no

effect on overall sleep or activity (Fig. 2A-E, Table S2). We observed similar sleep and

activity trends in *ATPSynC*, *Lsn* and *Bruce* females, but no change in *Imp* females

compared to controls (Fig. S3A). Since we observed a decrease in activity

accompanying an increase in sleep, we evaluated fly locomotion speed using

MARGO.[46,47] In males, we observed similar trends in locomotion speed as those

measured using the DAM system in *ATPSynC* and *Imp* males (Fig. S4A). However,

ATPSynC females, which also showed an increase in sleep, did not significantly affect

locomotion speed measured using MARGO (Fig. S4B). This suggests that the increase

in sleep in *ATPSynC* flies is not only due to a decrease in locomotor activity, however,

the increase in sleep in *Imp* flies could be due to an overall decrease in fly activity.

To further characterize these sleep alterations observed upon KD of each gene, we

measured the number and length of sleep bouts to assess sleep fragmentation.

Compared to driver and UAS controls, KD of only *Imp* resulted in increased sleep bout

number along with a decreased bout length which are indicative of sleep fragmentation

(Fig. 2F, G, Table S2). We also observed similar sleep and activity trends in 3-week-old

male flies (Fig. S5A). Moreover, to assess the effects of adult-specific KD of those

genes, we used the GeneSwitch system, and observed a consolidated increased sleep

phenotype in *ATPSynC* flies similar to that observed when the gene was KD throughout

development indicating the importance of this gene for sleep regulation, while sleep

dysfunction observed upon *Imp* KD was lost with adult-specific KD of this gene (Fig.

S6A). To determine the effect of KD of these genes on circadian rhythms, we measured

rhythmicity of flies under constant darkness. We found that neuronal-specific KD of *Imp*

led to circadian defects as all flies were arrhythmic (Table S3). KD of *ATPSynC*

decreased the number of rhythmic flies (41.18%) but it did not significantly affect the

period length or rhythm strength (FFT), while KD of *Lsn* or *Bruce* did not affect

rhythmicity (Table S3). Therefore, the neuronal suppression of genes within the CVD-

and insomnia-related locus led to a significantly altered sleep phenotype characterized

by an increase in overall sleep in two genes with only *Imp* affecting sleep fragmentation

and rhythmicity.

Cardiac-specific suppression of CVD- and insomnia-related genes leads to

cardiac dysfunction, myofibrillar disorganization, cardiac fibrosis, and shortened

lifespan.

To evaluate the effect of suppressing these genes on cardiac performance, KD of

ATPSynC, *Bruce*, *Lsn, or Imp,* was carried out using the cardiac-specific *Hand-*

Gal4 driver. Levels of KD of each gene in the heart is shown in Fig S1E-H. One-week-

old male and female flies were dissected and imaged for assessment of cardiac

physiological parameters. Interestingly, suppressing *Lsn* led to a non-beating, heart

failure-like phenotype where only 62.75% of hearts beat at 1 week of age which

decreased to 15.91% by 3 weeks of age in males (Fig. 3A). *ATPSynC* flies also

exhibited a decrease in the number of beating hearts with age, from 91.67% at 1-week

to 60% beating hearts at 3-weeks of age (Fig. 3A). This suggests an important role for

Lsn and *ATPSynC* in cardiac function. Upon analyzing beating hearts, cardiac-specific

KD of *ATPSynC* showed a bradycardic phenotype with cardiac dysfunction

characterized by significantly increased heart period (HP), arrhythmia index (AI),

diastolic interval (DI) and systolic diameter (SD) and reduced diastolic diameter (DD)

and fractional shortening (FS), a measure of cardiac performance, in both male and

female flies (Fig. 3B-H, Fig. S3B, Table S2). Suppression of *Lsn* led to cardiac

disfunction characterized by significantly increased DD and SD and reduced FS in both

sexes, in addition to a tachycardic phenotype in males characterized by decreased HP

(Fig. 3B-H, Fig. S3B, Table S2). Suppression of *Imp* led to a tachycardia characterized

by decreased HP without affecting cardiac function in males only (Fig. 3B-H, Fig. S3B,

Table S2). However, suppressing *Bruce* in males and females did not significantly affect

heart function in one-week-old flies (Fig. 3B-H, Fig. S3B, Table S2). To assess whether

or not the effects of these genes on the heart persist with age, we assessed cardiac

function in 3-week-old male flies with cardiac-specific suppression of these genes and

observed similar overall trends for cardiac parameters as observed in 1-week-old

ATPSynC and *Lsn* flies (Fig. S5B). Interestingly, in 3-week-old flies with KD of *Bruce*,

we observed cardiac dysfunction characterized by increased DD and SD and decreased

FS (Fig. S5B), suggesting an age-related component involved in the cardiac phenotype

observed with *Bruce* KD. To determine the effect of adult-specific KD of these genes,

we used the GeneSwitch system. Upon adult-specific KD of *ATPSynC* or *Lsn*, we

observe significantly decreased fractional shortening similar to that observed upon KD

throughout development; thus, confirming that *ATPSynC* or *Lsn* are essential genes for

cardiac function (Fig. S6B).

To characterize the morphology of hearts with KD of these genes, 1-week-old male

hearts were stained with phalloidin. KD of *ATPSynC* severely disrupted actin-containing

myofibrillar organization and led to almost complete loss of contractile circumferential

muscles (CF) and mostly non-contractile longitudinal muscles (LF) are seen (Fig. 4A,

Fig. S7A). KD of *Lsn* showed a dilated heart with more evident CF aggregations along

with myofibrillar disarray, while that of *Bruce* and *Imp* showed a less severe phenotype,

with visible CFs and LFs (Fig. 4A, Fig. S7A). Moreover, only suppression of *Lsn* led to

significantly increased pericardin deposition, which is a collagen-like protein and a

component of the extra cellular matrix (ECM) indicative of a fibrotic phenotype (Fig. 4B,

C). We also assessed the effects of cardiac KD of these genes on lifespan. Cardiac

suppression of *ATPSynC, Lsn,* and *Imp* led to a significantly shortened lifespan in both

sexes (p< 0.0001), while flies with suppressed *Bruce* showed an increased lifespan in

males (p=0.0057) (Fig. 4D). Our findings indicate that suppression of CVD- and

insomnia-related genes *Lsn* and *ATPSynC* in the heart led to significantly compromised

cardiac function with myofibril disorganization and shortened lifespan. Moreover, *Bruce*

and *Imp* showed less severe phenotypes with *Bruce* leading to cardiac dysfunction with

increased age.

Non-cell-autonomous mechanisms linking CVD with insomnia.

Mendelian randomization analyses in recent genetic studies confirm a causal role for

insomnia on CVD.[12] Moreover, cardiac dysfunction has also been associated with sleep

disruptions.[20,21,48] Therefore, to assess non-cell-autonomous roles of these genes in

influencing cardiac and sleep dysfunction, we suppressed genetic expression in the

neurons and measured cardiac function, or we suppressed it in the heart and assessed

sleep phenotypes in 3-week-old male flies (Fig. 5A). Unlike cardiac-specific KD,

neuronal suppression of *ATPSynC*, *Lsn* or *Bruce* resulted in no cardiac phenotype,

while neuronal suppression of *Imp* in 3-week-old male flies significantly decreased HP,

DI and significantly reduced FS (Fig. 5B-D). Neuronal-specific KD of only *Imp* led to

increased sleep fragmentation (Fig. 2F, G), which may be underlying the cardiac

dysfunction observed in those flies. This compromised cardiac performance indicates a

non-cell-autonomous effect of *Imp* in the causal role of sleep disruption on cardiac

function.

Upon the cardiac-specific KD of *Bruce* in 3-week-old flies, overall sleep significantly

increased (Fig. 5E). However, KD of ATPSynC or *Lsn* did not affect overall sleep

(Fig.5E). We then measured the number and length of sleep bouts to assess sleep

fragmentation. Only *ATPSynC* and *Lsn* showed a significant increase in total sleep bout

numbers and decrease in total bout length (Fig. 5F, G) indicative of significant sleep

fragmentation in these flies. In support of our data, cardiac dysfunction has been

associated with sleep disruptions in observational studies.[21,48]

Since inflammation was reported as a potential mechanism underlying the connection

between CVD and sleep dysfunction[18], we measured *Upd3* levels, an IL-6-like

proinflammatory cytokine in flies[49], in heads and hearts of flies with neuronal-specific

KD (Fig. 6A, B) or cardiac-specific KD (Fig. 6C, D). We first evaluated the effect of

neuronal-specific KD of these genes on *Upd3* levels. *ATPSynC* KD led to increased

Upd3 levels in heads of flies while *Imp, Lsn, and Bruce* KD led to a trend towards

increased Upd3 levels, but it was not significant (Fig. 6A). This may suggest that Upd3

is a mechanism involved in the sleep effects observed upon KD of these genes. Upon

the KD of *Imp* neuronally, *Upd3* levels in the heart significantly increased (Fig.6B). This

suggests a non-cell-autonomous effect of *Imp* in the causal role of sleep disruption on

cardiac dysfunction through an increased inflammatory state. We then measured the

levels of Upd3 in the heart upon cardiac-specific KD of these genes. KD of *Lsn* led to

significant increase in Upd3 levels, while suppression of *ATPSynC* led to a trend

towards increased Upd3 levels (p=0.0719) (Fig. 6C). This may suggest that Upd3 is the

specific mechanism underlying CVD-related inflammation observed upon KD of these

genes. As another measure of inflammatory-like state, we performed hemocyte counts.

Cardiac-specific suppression of *Lsn* led to a significantly increased number of

hemocytes in the hemolymph (Fig. S7B). Interestingly, the KD of *Lsn* in the heart led to

a significant increase in *Upd3* levels in the head, while that of *ATPSynC* showed an

increase that was not significant (Fig. 6D, p=0.33), both, Lsn and ATPSynC, being the

only 2 genes that led to an increased sleep fragmentation phenotype non-cell-

autonomously upon cardiac-specific KD (Fig. 5F,G).

To assess whether Upd3 elevation is a mechanism involved in the non-cell-autonomous

effects between cardiac and sleep dysfunction, we overexpressed Upd3 in the neurons

or heart using Elav- or Hand-Gal4, respectively. When we overexpressed it in neurons,

it increased overall sleep amount (Fig. S8A), similar to sleep dysfunction observed in

ATPSynC and Imp KD (Fig. 2B). We then assessed cardiac function upon neuronal-

specific Upd3 overexpression, and we observed a tachycardiac phenotype

characterized by decreased HP, DI and decreased cardiac performance reflected by

decreased FS (Fig. 7A), which is similar to that observed upon the neuronal KD of Imp

(Fig. 5B-D). When we overexpressed Upd3 in the heart, we observed a cardiac

dysfunction phenotype similar to Lsn, characterized by increased AI, DD, and SD, and

decreased FS (Fig. S8B). Interestingly, we also observed a non-cell-autonomous effect

on sleep where flies with cardiac-specific overexpression of Upd3 also had significantly

increased sleep fragmentation characterized by increase in bout numbers and decrease

in bout lengths without effecting sleep amount (Fig. 7B) similar to the non-cell-

autonomous effect of cardiac-specific KD of ATPSynC and Lsn on sleep (Fig. 5E-G).

These novel findings suggest inflammation through Upd3 as an underlying mechanism

linking cardiac and sleep dysfunction, bidirectionally.

Discussion

This study is the first to identify four genes at a single locus that link CVD and insomnia,

ATPSynC, Bruce, *Lsn,* and *Imp* (ATP5G1, UBE2Z, SNF8, and IGF2BP1 in humans).

Genetic screens have been previously applied in different model systems including

zebrafish, *Drosophila* and mice to identify genes involved in different CVDs and sleep

regulation.[50-56] Despite the numerous advantages of these models for functional and

behavioral screening, only few studies have utilized them to test genes identified after

human GWASs. A recent study[56] used *Drosophila* to identify causal variants reported in

an insomnia GWAS[13], including our insomnia-related locus, and screen candidate

genes to pinpoint those involved in sleep regulation. However, there are no studies to

date that identify genes related to both diseases or investigate functional genetic

mechanisms underlying a connection between CVD and sleep dysfunction.

Here, we used an innovative approach integrating the use of human genetics in

conjunction with fly genetics to identify genes related to each disease and advance the

understanding of the association between CVD and insomnia. We focused on a genetic

locus identified in both CVD and insomnia GWASs[13,15,23] and identified *Drosophila*

orthologs of potential nearby causal genes (Fig. 1). The locus we identified presented

as a colocalization signal for both diseases. Interestingly, of 554 risk loci for insomnia

identified thus far, this locus is among only three loci that colocalize with CVD (pp >0.90;

others include the ApoE region, *and LINGO4/RORC*)[45]. Functional dissection of other

association signals for cardiovascular/cardiometabolic traits in this same genomic

region suggests an importance of this region in CVD and complex contributions of

multiple genes at the locus to CVD pathogenesis.[57]

The first objective of our study was to functionally identify a novel role of these genes in

CVD and/or insomnia. Thus, we performed tissue-specific, neuronal or cardiac, KD of

each gene. KD of *Imp* significantly increased sleep which was fragmented and disrupted

rhythmicity (Fig. 2, Table S3), thus implicating this gene with sleep regulation and

maintenance of the circadian rhythm. However, KD of *Imp* also decreased the activity

and speed of flies (Fig.2, Fig. S4) which could influence the increase in sleep observed

in these flies. Therefore, future studies testing different levels of *Imp* KD and their

effects on sleep, activity and circadian rhythms are required for a better understanding

of the specific role of *Imp* on sleep physiology. KD of *Imp* did not have a severe cell-

autonomous effect on the heart. Similarly, KD of *Bruce* did not affect overall sleep but

started showing a decrease in cardiac function at 3 weeks of age in a cell-autonomous

manner, which in turn led to a non-cell-autonomous effect on sleep duration (Fig. S5,

Fig. 5).

Neuronal suppression of *ATPSynC* only increased overall sleep duration (Fig. 2). This

increased sleep phenotype is supported by published findings in another study,

screening insomnia-related genes identified from GWAS.[56] This increased sleep

corresponded with decreased locomotor activity, which has been recently reported in

humans and flies with mutations in *ATP5G1/ATPsynC*[58]. However, video tracking of

flies to probe for locomotion speed showed that speed was decreased only in males

even though both male and female flies showed an increase in sleep amount (Fig. 2,

Fig. S4) suggesting an effect of *ATPSynC* on sleep independent of locomotion defects.

Neuronal KD of ATPSynC also increased inflammation (Fig. 6A), which has been

associated with sleep dysfunction.[59] Cardiac suppression of *ATPSynC* significantly

compromised cardiac function characterized by severely increased arrythmia (Fig. 3)

and disrupted structure and fibrosis (Fig. 4). It also increased *Upd3*-specific

inflammation (Fig. 4), which is an important indicator of cardiac injury. These findings

revealed a novel role of *ATPSynC* in cardiac and sleep regulation in a cell-autonomous

manner. Both the neurons and heart require large amounts of ATP to perform their

functions. In both organs, ATP is essential for electrophysiological activities in resting

and active states[60,61], and reduction of ATP levels impairs neural and cardiac

functions[60,62,63]. ATP is produced by ATP Synthase, and impairing the function of ATP

Synthase is known to be associated with cardiovascular and neurological diseases.[64,65]

ATPSynC is an important component of ATP Synthase. Therefore, KD of *ATPSynC*

may disrupt the function of ATP Synthase thus contributing to CVD and sleep

disruptions we observed.

Cardiac-specific KD of *Lsn* also resulted in a severe cardiac phenotype with significant

compromised cardiac performance, characterized by significant dilation (Fig. 3), and

evident myofibril disorganization and fibrosis (Fig. 4). It also showed inflammation (Fig.

6C), indicating cardiac injury, along with a unique non-beating heart phenotype that

worsened with age (Fig. 3). These novel findings establish a cell-autonomous role of

Lsn in cardiac dysfunction. *Lsn* is part of the ESCRT pathway, which is a key

mechanism of multivesicular body (MVB) biogenesis.[30,66] MVBs form exosomes which

are crucial for intercellular communication and have been implicated in the

pathophysiology of CVD and other diseases[30,66,67], which supports the importance of

Lsn for cardiac function we observed.

Our next objective was to assess associations between CVD and insomnia and assess

the effects of one disease on the other through these genes (Fig. 5). First, we

suppressed these genes neuronally and assessed cardiac function. Unlike cardiac KD,

neuronal suppression of only *Imp* significantly reduced cardiac function which goes

along the strong sleep phenotypes observed upon neuronal *Imp* KD. One possible

mechanism reported to underly the effects of sleep dysfunction on cardiac function is

inflammation.[18] Notably, neuronal KD of *Imp* increased inflammation in the heart (Fig.

6B). Furthermore, the overexpression of Upd3 in neurons phenocopied those non-cell-

autonomous effects on cardiac function (Fig. 7A); thus, further suggesting the influence

of sleep dysfunction on cardiovascular performance through inflammation, supporting

mendelian randomization reports that show an effect of insomnia on CVD.

Although previous observational and genetic studies more commonly report an effect of

sleep on CVD, some human studies show an effect of heart failure on sleep

interruption.[20] Therefore, we were interested in assessing whether there is an influence

in the opposing direction, from heart on the neurons. Therefore, we suppressed these

genes in the heart and assessed sleep physiology (Fig. 5). We observed non-cell-

autonomous effects on sleep in genes with cardiac dysfunction upon cardiac-specific

KD, where flies with cardiac KD of Lsn and ATPSynC had increased sleep

fragmentation. Moreover, although cardiac KD of *Bruce* did not affect heart function in

1-week-old flies, it significantly reduced cardiac function in 3-week-old flies (Fig. S5)

which in turn increased sleep amount non-cell-autonomously. This shows an evident

effect of cardiac dysfunction on sleep. These findings suggest a non-cell-autonomous

influence of *ATPSynC*, *Lsn* and *Bruce* on sleep regulation.

While the influence of heart function on the nervous system has been reported,

underlying mechanisms remain poorly understood.[68,69] We hypothesized again that

inflammation is a mechanism underlying the effects of cardiac dysfunction on sleep

disruption. KD of *Lsn* and *ATPSynC* increased inflammation in the head after cardiac

suppression (Fig. 6D). Importantly, we show that the overexpression of Upd3 in the

heart leads to cardiac dysfunction cell-autonomously and sleep disruption non-cell-

autonomously (Fig. S8B, Fig. 7B); recapitulating non-cell-autonomous effects observed

by cardiac KD *ATPSynC* and *Lsn* on sleep. Overall, our results suggest inflammation as

an important underlying factor in the bidirectional connection between cardiovascular

function and sleep physiology.

One limitation of our study is the inability to characterize vascular and atherosclerotic

phenotypes in the fly as the locus was linked to coronary artery disease in pervious

GWASs. However, to address this, we measured inflammation, which has a crucial role

in CAD[70], to further characterize cardiac phenotypes observed upon KD of genes near

the CVD- and insomnia-related locus. Our study provides an important basis for future

studies in more complex model systems to further characterize the roles of these genes

on cardiovascular function. Moreover, our study focused on the effects of suppression

of these genes on sleep and cardiac physiology. Future work will evaluate the effects of

overexpressing these genes on each tissue and assess cell-autonomous and non-cell-

autonomous effects that will have on sleep and cardiac function.

In summary, we report four important findings: first, *ATPSynC* is the only gene that is

important cell-autonomously for both sleep and cardiac function and affects sleep non-

cell-autonomously through the heart; second, *Lsn* is important for cardiac function cell-

autonomously and only affects sleep non-cell-autonomously through the heart; third,

Imp is important cell-autonomously for sleep and affects the heart only non-cell-

autonomously through sleep; and fourth, we functionally identified inflammation as a

mechanism connecting cardiovascular and sleep dysfunction, bidirectionally (Fig. 8).

These findings advance our understanding of the association between CVD and sleep

disorders and provide basis for future studies to help develop therapeutic strategies that

prevent or attenuate insomnia and coincident CVD.

Materials and Methods

LocusZoom plots and phenotypic associations for the rs4643373 region and eQTL analyses.

The LocusZoom plots for CVD and insomnia SNP, rs4643373, were generated using LocusZoom v1.3 (06/20/2014). Nearby associations to rs4643373 were obtained from the Cardiovascular Disease Knowledge Portal (broadcvdi.org); 20230927; https://cvd.hugeamp.org/variant.html?variant=rs4643373 (RRID:SCR_016536). Results were plotted using LDassoc in LDLink. The GTEx V8 database was used for multi-tissue eQTL analyses. The GTEx V8 data contains a total of 17,382 RNA-seq samples from 948 post-mortem donors. The GTEx data used for multi-tissue eQTL analyses described in this manuscript were obtained from the GTEx Portal on 03/08/2023.

Drosophila Stocks

Drosophila stocks were cultured at 25°C on standard agar media.[71] UAS-RNAi transgenic stocks of CVD- and insomnia-related genes were obtained from Vienna Drosophila Resource Center (VDRC) and Bloomington Drosophila Stock Center (BDSC): *ATPSynC*-RNAi (VDRC: <u>106834</u>; BDSC: **35464**, 57705), *Lsn*-RNAi (VDRC: <u>110350</u>, **21658**; BDSC: 38289), *Bruce*-RNAi (VDRC: <u>107620</u>, 48309; BDSC: **51814**), *Imp*-RNAi (VDRC: 20321, <u>20322</u>; BDSC: 38219, **55645**, 34977), control lines (w^{1118}, VDRC: 60000, 60100, BDSC: 36303, 36304, 41552, 41553), *Act5C-Gal4* (BDSC: 4414); *24b-Gal4* (BDSC: 1767), *Elav-Gal4* (BDSC: X: 458 II: 8765). *Hand-Gal4* was obtained from Dr. Olson's lab. Genotypes are listed in Table S4. Data from RNAi lines were not combined if more than one RNAi line was used. Line 1 is underlined and is the main line used for all experiments. Line 2 is bolded and used only in Fig. S2. *ATPSynC*

Line 2 is used for neuronal-specific suppression as Line 1 was lethal when crossed with

Elav-Gal4. Elav-Gal4 (II) was used in overexpression experiments to allow for observing

more subtle phenotypes induced by lower expression of constructs, as overexpressing

with *Elav-Gal4 (X) was lethal*.

Ubiquitous- and Tissue-specific knock-down and genetic modulation

The GAL4-UAS system[72] was used to drive the knockdown of CVD- and insomnia-

related genes ubiquitously or tissue-specifically. Adult flies possessing UAS RNAi CVD-

and insomnia-related genes were crossed to *Hand-Gal4, Elav-Gal4, Act5C-Gal4, Ubi-*

Gal4, or *24b-Gal4* flies and incubated at 25°C throughout development. Adult male and

female F1 progeny were separated according to sex and allowed to age, with a new

food source supplied every three days prior to assays of cardiac function. Age-matched

adults from w^{1118} (wild-type), V60100 or BL36303 (VDRC and BDSC RNAi controls)

were crossed with each of the *Gal4* drivers as controls. Male and female flies were

screened at 1 and 3 weeks of age for cell-autonomous assays and 3 weeks for non-cell-

autonomous assays in at least 2 independent experiments. All flies were kept at 25 °C,

50% humidity in a 12 h light/12 h dark (LD) cycle.

Sleep-wake behavioral and rhythmicity analysis

Three-to-four-day old male and female progeny of *Elav-Gal4* (cell-autonomous) and 2.5-

week-old male progeny of *Hand-Gal4* (non-cell-autonomous) with control and RNAi

lines of each of the four genes were collected and individual flies were loaded into glass

tubes containing standard fly food (n>16). Sleep-wake behavior was recorded using

the Drosophila Activity Monitor (DAM, TriKinetics inc MA, USA) system in a 12L:12D

cycle at 25°C. Drosophila activity (or wake) is measured by infra-red beam crosses in

the DAM system.[73] To evaluate the role of CVD- and insomnia-related genes in

maintaining adult rhythms, 3-4-day old male flies were loaded into glass tubes and

entrained for at least 3 days to a 12h:12h LD cycle, followed by 3 days in constant

darkness. Rhythmicity analysis was performed for each 3-day period in constant

darkness. Data was analyzed using ClockLab and RStudio. Custom R scripts and

methodology used with RStudio can be found on

https://github.com/jameswalkerlab/Gill_et.al. One-way ANOVA with Dunnet's multiple

comparisons test for DAM system data was performed using GraphPad Prism.

Drosophila sleep was defined by a period of at least 5 minutes of inactivity,

demonstrated by zero beam breaks recorded.[74] Average sleep per 24 hours (ZT0-ZT24)

of each genotype was calculated. Five days were used for analysis of 1-week-old flies,

and 3 days were used for 3-week-old fly experiments to overcome decreased viability in

older flies. Sleep bouts were quantified by counting the number of periods of sleep as

defined above. Sleep fragmentation was defined by either the number of 1-minute

wakes or sleep bouts during a 24-hour period. Data for daytime sleep is from ZT0 to

ZT12 and that for nighttime sleep is from ZT12-ZT24.

MARGO locomotion monitoring

Drosophila locomotion was video monitored using the MARGO (Massively Automated

Real-time GUI for Object-tracking) system for automated tracking.[46,47] Behavior boards

were prepared by the addition of standard fly food at one end. Individual 3-5-day old

male and mated female flies were loaded into the behavior boards such that there was

one fly per channel. Flies were video recorded for 3 days at 20°C. Collected data from

days 2 and 3 were analyzed using a custom script in MatLab. Speed data was

calculated by the change in location of the centroid of the tracked fly between frames.

This data collected at a frame rate of 4Hz and an experimental setup such that 3.4667

pixels equal 1 millimeter. Using these values, pixels per frame were converted to

millimeters per second values. Frames in which flies were immobile were excluded from

speed calculations. Therefore, average speed measurements are representative of the

average speed while flies are in motion. Average speed values are an average of

speeds from days 2 and 3 and normalized to controls set up within the same behavior

board.

Cardiac physiological analyses of semi-intact Drosophila hearts

1-week-old male and female progeny of *Hand-Gal4* (cell-autonomous) and 2.5-week-old

male progeny of *Elav-Gal4* (non-cell-autonomous) with control and RNAi lines of each

of the four genes were collected, and semi-intact hearts were prepared as described

(n>30).[75,76] Direct immersion optics was used in conjunction with a digital high-speed

camera (at 200 frames/sec, Hamamatsu Flash 4 camera) to record 30 second movies

of beating hearts; images were captured using HC Image (Hamamatsu Corp.). Cardiac

function was analyzed from the high-speed movies using semi-automatic optical

heartbeat analysis (SOHA) software that quantifies heart rate, heart period, diastolic

and systolic diameters, diastolic and systolic intervals, cardiac rhythmicity, fractional

shortening and produced the Mechanical-mode records [75,76].

Cytological Studies of adult hearts

Dissected hearts from one-week old adults were relaxed by a one-minute treatment with 5 mM EGTA in hemolymph and then fixed with 4% paraformaldehyde in PBS for 30 min as previously described [76]. Fixed hearts were stained with anti-Pericardin antibody overnight (5ug/ml, 1:10; Developmental Biology Hybridoma Bank, University of Iowa) followed by Alexa488-phalloidin for 30 min (1:1000, U0281, Abnova), which stains F-actin containing myofibrils. Samples were then mounted with Diamond Antifade Mountant with DAPI. Confocal images were taken from a Nikon A1R HD microscope (UAB) at 10X for pericardin quantification and 20X for representative images for phalloidin staining. Quantification of pericardin area in the confocal images from three to five independent male hearts per genotype was performed by thresholding images in ImageJ, then percent area was measured.[77]

Adult-specific KD of genes

To induce adult-specific KD, we crossed flies from each genotype with either Hand- or Elav-GeneSwitch-Gal4 and put them on RU486-containing food three-to-four days after eclosion. For crosses with *Hand-GeneSwitch-Gal4*, we used food containing 200uM of RU486, while for crosses with *Elav-GeneSwitch-Gal4*, we used food containing 500uM of RU486 to allow activation of the GeneSwitch system. We changed the food twice a week for three weeks, then , we measured cardiac and sleep physiology of three-week-old male flies.

Viability

Adult flies (n>100, males and females) with suppression of CVD- and insomnia-related genes and controls were collected on the day of eclosion from the pupal case, designated as day zero. Approximately 30 flies were placed in each vial and transferred to a new vial every three to four days. The numbers of surviving adults were counted twice a week. The numbers of surviving adults were compared to the original number of adults collected on day zero and the percentage for each day was graphed.[78]

Hemocyte Counts

To evaluate inflammation, fly hemolymph was collected from n>100 (per replicate, 3 biological replicates) one-week-old adult male flies with cardiac-specific suppression using *Hand-Gal4* by making an incision in the thorax of flies and centrifuging them.[79] Hemocytes were then counted by staining the hemolymph with 1:1 Trypan blue dilution and using a hemocytometer.[79]

Real-time quantitative PCR

Dissected male hearts (n=10-12 per biological replicate, 3 biological replicates) and heads (n=10, per biological replicate, 3 biological replicates) from 1-week-old flies was placed in the RNA lysis buffer, and flash frozen. RNA from heads was extracted using Zymo Research Quick-RNA Microprep Kit with on column DNase I digestion. RNA from hearts was extracted using the RNeasy kit (QIAGEN). Quantitative PCR was performed using SsoAdvanced Universal SYBR Green supermix (Bio Rad) in a BIO-RAD CFX Opus Real-Time PCR System. Expression was normalized with 60S ribosomal protein

(RPL11). Primers for qPCR are listed below: *ATPSynC-F:*
GCAACAGTCGGTGTCGCT; ATPSynC -R: AGGCGAACAGCAGCAGGAA; Lsn-F:
TCACCAAGGAGGACATCCTAATGG; Lsn-R: TCCGGGAATGGACTGAACTATGTA;
Bruce-F: AATAGCGCTCCATCTCGACCAT; Bruce-R: ATCGACCATGCACAATGCTGT;
Imp-F: AATTCGCCGACCTGGAACTCT; Imp-R: ACTCGACACCGTTCAGACCAA;
Upd3-F: AGCCGGAGCGGTAACAAAA; Upd3-R: CGAGTAAGATCAGTGACCAGTTC;
Rpl11-F: CGATCTGGGCATCAAGTACGA; Rpl11-R: TTGCGCTTCCTGTGGTTCAC.
Results are presented as $2^{-\Delta\Delta Ct}$ values normalized to the expression of Rpl11 and
control samples. All reactions were performed using biological triplicates. The means
and standard error of the mean were calculated in GraphPad Prism 9 software.
619
620

Statistical Analysis
For all quantitation except transcript levels and lifespan analyses, statistical significance
was determined using one-way analysis of variance (ANOVA) followed by Dunnett's
post-hoc test to determine significance between groups for sleep and cardiac
physiological parameters. Comparisons to UAS controls were calculated by 1-way
ANOVA followed by Šidák's post-hoc test to assess off-target effects of RNAi lines.
Additional comparisons of sleep and cardiac parameters of flies to their respective RNAi
controls in Table S5 were calculated by 1-way ANOVA followed by Šidák's post-hoc
test. For expression of transcript levels in heads, statistics were calculated by 1-way
ANOVA. For expression of transcript levels in hearts, statistics were calculated by
Kruskal-Wallis test and without correcting for multiple comparisons to account for

variability. For overexpression experiments, statistical significance was determined

using an unpaired t-test. Bar graphs show mean $\pm$ SEM. For lifespan studies, data

were analyzed using the Kaplan-Meier test followed by multiple comparisons between

control and experimental groups. Significance was presented using p-values on figures.

All statistical analysis were performed with GraphPad Prism 9.

Acknowledgments

We would like to thank Dr. Philip R. Jansen for the insomnia LocusZoom plot, Dr.

Harrison for supplying the UAS-Upd3 overexpression stock and Dr. Olson for supplying

the Hand-Gal4 driver stock. We thank Dr. Louis Dell'Italia and Dr. Jonathan Roth for

their editorial comments on the manuscript. We would also like to thank Dr. Ruan

Moraes for his technical support. The Genotype-Tissue Expression (GTEx) Project was

supported by the Common Fund of the Office of the Director of the National Institutes of

Health, and by NCI, NHGRI, NHLBI, NIDA, NIMH, and NINDS. Stocks obtained from

the Bloomington Drosophila Stock Center, and Vienna Drosophila Resource Center

were used in this study. Illustrations throughout the manuscript were created with

BioRender.com.

Figures

650
Fig. 1. CVD- and insomnia-related locus, other phenotypic associations and

nearby genes. Manhattan plots (LocusZoom) showing CVD (A) and insomnia (B) SNP

association peaks with 5 nearby candidate genes, *ATP5MC1 (ATP5G1), UBE2Z, SNF8,*

GIP, IGF2BP1. GWAS associations meeting genome-wide significance (p<$^{5x10-8}$, blue

line) for cardiovascular/cardiometabolic traits in the region near rs4643373 obtained

from the Cardiovascular Disease Knowledge Portal (see methods) and literature. Each

point corresponds to a single trait in a single study. Blue dot represents rs4643373 (C).

Graphical Scheme showing workflow (D).

Human Symbol	Drosophila Symbol	% Similarity
ATP5G1	ATPSynC	83
UBE2Z	Bruce	54

SNF8	Lsn	71
IGF2BP1	Imp	58
GIP	-	-

Table 1. Human and fly symbols of CVD- and insomnia-related genes with %similarity

at a single locus.

Fig. 2. Neuronal-specific suppression of CVD- and insomnia- related genes leads

to compromised sleep phenotypes. Sleep profiles showing sleep minutes per hour for

24 hours (A). Violon plots for quantitative sleep parameters, total sleep amount (B),

daytime sleep amount (C), nighttime sleep amount (D), total locomotor activity (E), total

bout number (F), and total bout length (G) from 1-week-old male *Drosophila* with

neuronal RNAi knockdown of CVD- and insomnia-related genes with the pan-neuronal

Elav-Gal4 driver. N=16-24 for each group. For *ATPSynC*, Line 1 was lethal; Line 2 was

used (refer to methods section). Data was collected from at least 2 independent

experiments from one RNAi line (second line data shown in Fig. S2). Each data point

represents a fly. Statistics were calculated by one-way ANOVA for comparisons to

controls.

Fig. 3. Cardiac-specific suppression of CVD- and insomnia-related genes leads to

cardiac dysfunction. Representative 5-second mechanical-modes (A) from 1-week-old

male flies with cardiac RNAi knockdown of CVD- and insomnia-related genes with

cardiac-specific *Hand-Gal4* driver. Percentage of beating hearts at 1 versus 3 weeks of

age shows significant effect of Lsn KD with age (p<0.0001) (B). Violin plots for cardiac

physiological parameters, heart period (C), arrythmia index (D), diastolic interval (E),

diastolic diameter (F), systolic diameter (G), fractional shortening (H), and N= 29-33 for

each group for C-H, N=32-51 for each group for B. Each data point represents one fly.

Data was collected from at least 2 independent experiments from one RNAi line

(second line data shown in Fig. S2). Statistics were calculated by 1-way ANOVA for C-

H. Fisher's exact test was performed for A.

Fig. 4. Cardiac-specific suppression of CVD- and insomnia-related genes leads to

myofibrillar disorganization, cardiac fibrosis, and shortened lifespan.

Representative images showing actin-containing myofibrils (A) and pericardin (B) from

1-week-old male flies with cardiac RNAi knockdown of CVD- and insomnia-related

genes with *Hand-Gal4.* Quantification of pericardin signal (C). Each data point is a fly.

Lifespan assay (D) for male flies with cardiac RNAi knockdown of CVD- and insomnia-

related genes with cardiac-specific *Hand-Gal4* driver resulted in significant decrease in

lifespan (p<0.0001) of *ATPSynC, Lsn* and *Imp*, and a significant increase in lifespan of

Bruce (p=0.0057). Graph plots % survival (n☐>100 for each group) vs. time post-

eclosion. Statistics were calculated by 1-way ANOVA for C-D and a Kaplan-Meier test

was performed for E.

701
702 **Figure 5. Non-cell-autonomous mechanisms linking CVD with insomnia.** Graphical

703 scheme showing experimental layout (A). Violin plots for cardiac physiological

parameters, heart period (B), diastolic interval (C) and fractional shortening (D) from 3-week-old male flies with neuronal-specific knockdown of CVD- and insomnia-related genes (N= 19-32 per group). Violin plots for quantitative sleep parameters; total sleep amount (E), total bout number (F), and total bout length (G) from 3-week-old male *Drosophila* with cardiac-specific knockdown of CVD- and insomnia-related genes. N= 12-30 per group. Each data point represents one fly. Statistics were calculated by 1-way ANOVA.

Figure 6. Knockdown of CVD- and insomnia-related genes increases inflammation cell-autonomously and non-cell-autonomously. Transcript levels of Upd3 in heads of flies with neuronal-specific knockdown of these genes (N= 10-12 heads per data point per group) (A). Transcript levels of Upd3 in hearts of flies with neuronal-specific

knockdown of these genes (N= 10-12 hearts per data point per group) (B). Transcript

levels of Upd3 in hearts of flies with cardiac-specific knockdown of these genes (N= 10-

12 hearts per data point per group) (C). Transcript levels of Upd3 in heads of flies with

cardiac-specific knockdown of these genes (N= 10-12 heads per data point per group)

(D). Statistics were calculated by 1-way ANOVA.

Figure 7. Overexpressing Upd3 in neurons or the heart leads to cardiac or sleep

dysfunction, respectively. Bar graphs showing cardiac physiological parameters,

heart period, diastolic interval and fractional shortening from 3-week-old male flies with

neuronal-specific overexpression of Upd3 (N= 16-26 per group) (B). Bar graphs

showing quantitative sleep parameters; total sleep amount, total bout number, and total

bout length from 3-week-old male flies with cardiac-specific overexpression of Upd3. N=

13-14 per group. Each data point represents one fly. Statistics were calculated by

unpaired t-test.

Fig. 8. Graphical summary displaying main findings. Independent GWASs identified

one genetic locus associated with CVD and insomnia. Near this locus, we identified five

candidate genes, four of which have fly orthologs. To investigate the roles of genes near

this locus in both neurons and the heart, we suppressed their expression in each tissue

and assessed their effects on sleep and cardiac function in a cell-autonomous or non-

cell-autonomous manner. Our main findings include: one gene (*ATPSynC*) involved cell-

autonomously in sleep and cardiac function and affects sleep non-cell-autonomously

through the heart; one gene (*Imp*) involved cell-autonomously in sleep and affects

cardiac function non-cell-autonomously through sleep; and one gene (*Lsn*) involved cell-

autonomously in cardiac function and affects sleep non-cell-autonomously through the

heart. We also found that cell-autonomous and non-cell-autonomous effects were

accompanied with inflammation through elevation of *Upd3* levels.

Author Contributions: G.C.M. and F.A. designed the experiments in consultation with

R.S and J.W. F.A. performed all sleep and cardiac experiments. F.A. performed

analysis of cardiac physiological parameters, acquisition and analysis of cytological

imaging, and qPCR experiments and analysis. T.M. performed analysis of sleep data.

L.O and D.P. aided F.A. in experiments. M.M. and C.C. performed LocusZoom plots

with the help of R.S. S.M. performed MARGO experiments. F.A. prepared the paper

with G.C.M.'s input. All authors provided feedback on the manuscript.

References

Heart disease facts, <https://www.cdc.gov/heartdisease/facts.htm > ((2022, February 7)).

The top 10 causes of death. World Health Organization., <https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death> (

Virani, S. S. *et al.* Heart Disease and Stroke Statistics—2021 Update. *Circulation* **143**, e254-e743, doi:doi:10.1161/CIR.0000000000000950 (2021).

Larsson, S. C. & Markus, H. S. Genetic Liability to Insomnia and Cardiovascular Disease Risk. *Circulation* **140**, 796-798, doi:10.1161/circulationaha.119.041830 (2019).

Grandner, M. A. *et al.* Sleep: important considerations for the prevention of cardiovascular disease. *Curr Opin Cardiol* **31**, 551-565, doi:10.1097/hco.0000000000000324 (2016).

Zheng, B. *et al.* Insomnia symptoms and risk of cardiovascular diseases among 0.5 million adults. *A 10-year cohort* **93**, e2110-e2120, doi:10.1212/wnl.0000000000008581 (2019).

Quan, S. F. Sleep Disturbances and their Relationship to Cardiovascular Disease. *Am J Lifestyle Med* **3**, 55s-59s, doi:10.1177/1559827609331709 (2009).

Sleep Disorders, <https://medlineplus.gov/sleepdisorders.html> (

Suni, E. *Sleep Statistics*, <https://www.sleepfoundation.org/how-sleep-works/sleep-facts-statistics> (

Winkelman, J. W. Insomnia Disorder. *New England Journal of Medicine* **373**, 1437-1444, doi:10.1056/NEJMcp1412740 (2015).

Lind, M. J. & Gehrman, P. R. Genetic Pathways to Insomnia. *Brain Sci* **6**, doi:10.3390/brainsci6040064 (2016).

Lane, J. M. *et al.* Biological and clinical insights from genetics of insomnia symptoms. *Nat Genet* **51**, 387-393, doi:10.1038/s41588-019-0361-7 (2019).

Jansen, P. R. *et al.* Genome-wide analysis of insomnia in 1,331,010 individuals identifies new risk loci and functional pathways. *Nature Genetics* **51**, 394-403, doi:10.1038/s41588-018-0333-3 (2019).

Song, W. *et al.* Genome-wide association analysis of insomnia using data from Partners Biobank. *Scientific Reports* **10**, 6928, doi:10.1038/s41598-020-63792-0 (2020).

15   Watanabe, K. *et al.* Genome-wide meta-analysis of insomnia in over 2.3 million
individuals implicates involvement of specific biological pathways through gene-
prioritization.   *medRxiv*,   2020.2012.2007.20245209,
doi:10.1101/2020.12.07.20245209 (2020).
16   Hoevenaar-Blom, M. P., Spijkerman, A. M., Kromhout, D., van den Berg, J. F. &
Verschuren, W. M. Sleep duration and sleep quality in relation to 12-year
cardiovascular disease incidence: the MORGEN study. *Sleep* **34**, 1487-1492,
doi:10.5665/sleep.1382 (2011).
17   Hsu, C. Y. *et al.* The Association Between Insomnia and Increased Future
Cardiovascular Events: A Nationwide Population-Based Study. *Psychosom Med*
77, 743-751, doi:10.1097/psy.0000000000000199 (2015).
18   Javaheri, S. & Redline, S. Insomnia and Risk of Cardiovascular Disease. *Chest*
152, 435-444, doi:10.1016/j.chest.2017.01.026 (2017).
19   Bertisch, S. M. *et al.* Insomnia with objective short sleep duration and risk of
incident cardiovascular disease and all-cause mortality: Sleep Heart Health
Study. *Sleep* **41**, doi:10.1093/sleep/zsy047 (2018).
20   Zheng, T. Sleep disturbance in heart failure: A concept analysis. *Nurs Forum* **56**,
710-716, doi:10.1111/nuf.12566 (2021).
21   Parati, G. *et al.* Heart failure and sleep disorders. *Nature Reviews Cardiology* **13**,
389-403, doi:10.1038/nrcardio.2016.71 (2016).
22   McAlpine, C. S. *et al.* Sleep modulates haematopoiesis and protects against
atherosclerosis. *Nature* **566**, 383-387, doi:10.1038/s41586-019-0948-2 (2019).
23   Harst, P. v. d. & Verweij, N. Identification of 64 Novel Genetic Loci Provides an
Expanded View on the Genetic Architecture of Coronary Artery Disease.
Circulation Research **122**, 433-443, doi:doi:10.1161/CIRCRESAHA.117.312086
(2018).
24   Khera, A. V. & Kathiresan, S. Genetics of coronary artery disease: discovery,
biology and clinical translation. *Nature Reviews Genetics* **18**, 331-344,
doi:10.1038/nrg.2016.160 (2017).
25   Hartiala, J. A. *et al.* Genome-wide analysis identifies novel susceptibility loci for
myocardial infarction. *Eur Heart J* **42**, 919-933, doi:10.1093/eurheartj/ehaa1040
(2021).
26   Zhu, Z. *et al.* Genetic overlap of chronic obstructive pulmonary disease and
cardiovascular disease-related traits: a large-scale genome-wide cross-trait
analysis. *Respiratory Research* **20**, 64, doi:10.1186/s12931-019-1036-8 (2019).
27   Said, S. *et al.* Genetic analysis of over half a million people characterises C-
reactive protein loci. *Nat Commun* **13**, 2198, doi:10.1038/s41467-022-29650-5
(2022).
28   Saleh, A. A. *et al.* Evaluation of mRNA expression level of the ATP synthase
membrane subunit c locus 1 (ATP5G1) gene in patients with schizophrenia.
Biochem Biophys Rep **30**, 101234, doi:10.1016/j.bbrep.2022.101234 (2022).
29   Shi, X. *et al.* Upregulation of ubiquitin-conjugating enzyme E2Z is associated with
human hepatocellular carcinoma. *Biochem Biophys Res Commun* **523**, 25-32,
doi:10.1016/j.bbrc.2019.11.170 (2020).

Xu, B. *et al.* The ESCRT-III pathway facilitates cardiomyocyte release of cBIN1-containing microparticles. *PLOS Biology* **15**, e2002354, doi:10.1371/journal.pbio.2002354 (2017).

Huang, X. *et al.* Insulin-like growth factor 2 mRNA-binding protein 1 (IGF2BP1) in cancer. *J Hematol Oncol* **11**, 88, doi:10.1186/s13045-018-0628-y (2018).

Campbell, J. E. & Drucker, D. J. Pharmacology, physiology, and mechanisms of incretin hormone action. *Cell Metab* **17**, 819-837, doi:10.1016/j.cmet.2013.04.008 (2013).

Piazza, N. & Wessells, R. J. Drosophila models of cardiac disease. *Prog Mol Biol Transl Sci* **100**, 155-210, doi:10.1016/b978-0-12-384878-9.00005-4 (2011).

Bushey, D. & Cirelli, C. From genetics to structure to function: exploring sleep in Drosophila. *Int Rev Neurobiol* **99**, 213-244, doi:10.1016/b978-0-12-387003-2.00009-4 (2011).

Feng, G. *et al.* Control of Sleep Onset by Shal/K$_v$4 Channels in Drosophila Circadian Neurons. *The Journal of Neuroscience* **38**, 9059-9071, doi:10.1523/jneurosci.0777-18.2018 (2018).

Bhide, S. *et al.* Increasing autophagy and blocking Nrf2 suppress laminopathy-induced age-dependent cardiac dysfunction and shortened lifespan. *Aging Cell* **17**, e12747, doi:10.1111/acel.12747 (2018).

Gill, S., Le, H. D., Melkani, G. C. & Panda, S. Time-restricted feeding attenuates age-related cardiac decline in Drosophila. *Science* **347**, 1265-1269, doi:10.1126/science.1256682 (2015).

Wolf, M. J. & Rockman, H. A. Drosophila, Genetic Screens, and Cardiac Function. *Circulation Research* **109**, 794-806, doi:doi:10.1161/CIRCRESAHA.111.244897 (2011).

Souidi, A. & Jagla, K. Drosophila Heart as a Model for Cardiac Development and Diseases. *Cells* **10**, doi:10.3390/cells10113078 (2021).

Wolf, M. J. *et al. Drosophila* as a model for the identification of genes causing adult human heart disease. *Proceedings of the National Academy of Sciences* **103**, 1394-1399, doi:doi:10.1073/pnas.0507359103 (2006).

Cirelli, C. & Bushey, D. Sleep and wakefulness in Drosophila melanogaster. *Ann N Y Acad Sci* **1129**, 323-329, doi:10.1196/annals.1417.017 (2008).

Beckwith, E. J. & French, A. S. Sleep in Drosophila and Its Context. *Frontiers in Physiology* **10**, doi:10.3389/fphys.2019.01167 (2019).

Nall, A. H. & Sehgal, A. Small-molecule screen in adult Drosophila identifies VMAT as a regulator of sleep. *J Neurosci* **33**, 8534-8540, doi:10.1523/jneurosci.0253-13.2013 (2013).

Aragam, K. G. *et al.* Discovery and systematic characterization of risk variants and genes for coronary artery disease in over a million participants. *Nat Genet* **54**, 1803-1815, doi:10.1038/s41588-022-01233-6 (2022).

Watanabe, K. *et al.* Genome-wide meta-analysis of insomnia prioritizes genes associated with metabolic and psychiatric pathways. *Nat Genet* **54**, 1125-1132, doi:10.1038/s41588-022-01124-w (2022).

Werkhoven, Z., Rohrsen, C., Qin, C., Brembs, B. & de Bivort, B. MARGO (Massively Automated Real-time GUI for Object-tracking), a platform for high-

throughput ethology. *PLOS ONE* **14**, e0224243,
doi:10.1371/journal.pone.0224243 (2019).

47   Elya, C. *et al.* Neural mechanisms of parasite-induced summing behavior in
'zombie' Drosophila. *Elife* **12**, doi:10.7554/eLife.85410 (2023).

48   Sharma, B., Owens, R. & Malhotra, A. Sleep in congestive heart failure. *Med Clin*
North Am **94**, 447-464, doi:10.1016/j.mcna.2010.02.009 (2010).

49   Romão, D., Muzzopappa, M., Barrio, L. & Milán, M. The Upd3 cytokine couples
inflammation to maturation defects in Drosophila. *Current Biology* **31**, 1780-
1787.e1786, doi:https://doi.org/10.1016/j.cub.2021.01.080 (2021).

50   Koh, K. *et al.* Identification of SLEEPLESS, a sleep-promoting factor. *Science*
321, 372-376, doi:10.1126/science.1155942 (2008).

51   Neely, G. G. *et al.* A global in vivo Drosophila RNAi screen identifies NOT3 as a
conserved regulator of heart function. *Cell* **141**, 142-153,
doi:10.1016/j.cell.2010.02.023 (2010).

52   Kamp, A. *et al.* Genome-wide identification of mouse congenital heart disease
loci. *Hum Mol Genet* **19**, 3105-3113, doi:10.1093/hmg/ddq211 (2010).

53   Snider, P. & Conway, S. J. Probing human cardiovascular congenital disease
using transgenic mouse models. *Prog Mol Biol Transl Sci* **100**, 83-110,
doi:10.1016/b978-0-12-384878-9.00003-0 (2011).

54   Spielmann, N. *et al.* Extensive identification of genes involved in congenital and
structural heart disorders and cardiomyopathy. *Nature Cardiovascular Research*
1, 157-173, doi:10.1038/s44161-022-00018-8 (2022).

55   Lenz, O., Xiong, J., Nelson, M. D., Raizen, D. M. & Williams, J. A. FMRFamide
signaling promotes stress-induced sleep in Drosophila. *Brain Behav Immun* **47**,
141-148, doi:10.1016/j.bbi.2014.12.028 (2015).

56   Palermo, J. *et al.* Variant-to-gene mapping followed by cross-species genetic
screening identifies GPI-anchor biosynthesis as a regulator of sleep. *Science*
Advances **9**, eabq0844, doi:doi:10.1126/sciadv.abq0844 (2023).

57   Erbilgin, A. *et al.* Identification of CAD candidate genes in GWAS loci and their
expression in vascular cells[S]. *Journal of Lipid Research* **54**, 1894-1905,
doi:https://doi.org/10.1194/jlr.M037085 (2013).

58   Neilson, D. E. *et al.* A Novel Variant of ATP5MC3 Associated with Both Dystonia
and Spastic Paraplegia. *Mov Disord* **37**, 375-383, doi:10.1002/mds.28821 (2022).

59   Dzierzewski, J. M., Donovan, E. K., Kay, D. B., Sannes, T. S. & Bradbrook, K. E.
Sleep Inconsistency and Markers of Inflammation. *Frontiers in Neurology* **11**,
doi:10.3389/fneur.2020.01042 (2020).

60   Zhu, X. H., Lu, M. & Chen, W. Quantitative imaging of brain energy metabolisms
and neuroenergetics using in vivo X-nuclear (2)H, (17)O and (31)P MRS at ultra-
high field. *J Magn Reson* **292**, 155-170, doi:10.1016/j.jmr.2018.05.005 (2018).

61   Weiss, R. G., Gerstenblith, G. & Bottomley, P. A. ATP flux through creatine
kinase in the normal, stressed, and failing human heart. *Proceedings of the*
National Academy of Sciences **102**, 808-813, doi:doi:10.1073/pnas.0408962102
(2005).

62   Owen, L. & Sunram-Lea, S. I. Metabolic agents that enhance ATP can improve
cognitive functioning: a review of the evidence for glucose, oxygen, pyruvate,
creatine, and L-carnitine. *Nutrients* **3**, 735-755, doi:10.3390/nu3080735 (2011).

63 Beard, D. A. *et al.* Reduced cardiac muscle power with low ATP simulating heart
failure. *Biophysical Journal* **121**, 3213-3223,
doi:https://doi.org/10.1016/j.bpj.2022.07.029 (2022).
64 Galber, C., Carissimi, S., Baracca, A. & Giorgio, V. The ATP Synthase Deficiency
in Human Diseases. *Life (Basel)* **11**, doi:10.3390/life11040325 (2021).
65 Long, Q., Yang, K. & Yang, Q. Regulation of mitochondrial ATP synthase in
cardiac pathophysiology. *Am J Cardiovasc Dis* **5**, 19-32 (2015).
66 Burtenshaw, D. *et al.* Exosomal Composition, Biogenesis and Profiling Using
Point-of-Care Diagnostics—Implications for Cardiovascular Disease. *Frontiers in*
Cell and Developmental Biology **10**, doi:10.3389/fcell.2022.853451 (2022).
67 Neves, K. B., Rios, F. J., Sevilla-Montero, J., Montezano, A. C. & Touyz, R. M.
Exosomes and the cardiovascular system: role in cardiovascular health and
disease. *The Journal of Physiology* **n/a**, doi:https://doi.org/10.1113/JP282054.
68 Xie, W., Zheng, F., Yan, L. & Zhong, B. Cognitive Decline Before and After
Incident Coronary Events. *Journal of the American College of Cardiology* **73**,
3041-3050, doi:https://doi.org/10.1016/j.jacc.2019.04.019 (2019).
69 Marebwa, B. K. *et al.* Cardiovascular Risk Factors and Brain Health: Impact on
Long-Range Cortical Connections and Cognitive Performance. *Journal of the*
American Heart Association **7**, e010054, doi:doi:10.1161/JAHA.118.010054
(2018).
70 Christodoulidis, G., Vittorio, T. J., Fudim, M., Lerakis, S. & Kosmas, C. E.
Inflammation in coronary artery disease. *Cardiol Rev* **22**, 279-288,
doi:10.1097/crd.0000000000000006 (2014).
71 Livelo, C. *et al.* Time-restricted feeding promotes muscle function through purine
cycle and AMPK signaling in Drosophila obesity models. *Nature Communications*
14, 949, doi:10.1038/s41467-023-36474-4 (2023).
72 Duffy, J. B. GAL4 system in Drosophila: a fly geneticist's Swiss army knife.
Genesis **34**, 1-15, doi:10.1002/gene.10150 (2002).
73 Feng, G. *et al.* Control of Sleep Onset by Shal/K(v)4 Channels in Drosophila
Circadian Neurons. *J Neurosci* **38**, 9059-9071, doi:10.1523/jneurosci.0777-
18.2018 (2018).
74 Hendricks, J. C. *et al.* Rest in Drosophila is a sleep-like state. *Neuron* **25**, 129-
138, doi:10.1016/s0896-6273(00)80877-6 (2000).
75 Fink, M. *et al.* A new method for detection and quantification of heartbeat
parameters in Drosophila, zebrafish, and embryonic mouse hearts.
Biotechniques **46**, 101-113, doi:10.2144/000113078 (2009).
76 Melkani, G. C. *et al.* Huntington's disease induced cardiac amyloidosis is
reversed by modulating protein folding and oxidative stress pathways in the
Drosophila heart. *PLoS Genet* **9**, e1004024, doi:10.1371/journal.pgen.1004024
(2013).
77 Voskobiynyk, Y. *et al.* Alzheimer's disease risk gene BIN1 induces Tau-
dependent network hyperexcitability. *eLife* **9**, e57354, doi:10.7554/eLife.57354
(2020).
78 Melkani, G. C. *et al.* Huntington's Disease Induced Cardiac Amyloidosis Is
Reversed by Modulating Protein Folding and Oxidative Stress Pathways in the

Drosophila Heart. *PLOS Genetics* **9**, e1004024,
doi:10.1371/journal.pgen.1004024 (2013).
79 Au - Hiroyasu, A., Au - DeWitt, D. C. & Au - Goodman, A. G. Extraction of
Hemocytes from Drosophila melanogaster Larvae for Microbial Infection and
Analysis. *JoVE*, e57077, doi:doi:10.3791/57077 (2018).
980