Modulation of prefrontal couplings by prior belief-related responses in
ventromedial prefrontal cortex

Bin A. Wang[1,2*], Sabrina Drammis[3,4,5], Ali Hummos[4,5], Michael M. Halassa[6,7] and Burkhard
Pleger[1,2]

[1]Department of Neurology, BG University Hospital Bergmannsheil, Ruhr-University Bochum,
Bochum, Germany

[2]Collaborative Research Centre 874 "Integration and Representation of Sensory Processes",
Ruhr-University Bochum, Bochum, Germany

[3]Computer Science & Artificial Intelligence Laboratory, Massachusetts Institute of Technology,
Cambridge, Massachusetts, USA

[4]McGovern Institute for Brain Research, Massachusetts Institute of Technology, Cambridge,
Massachusetts, USA.

[5]Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology,
Cambridge, Massachusetts, USA.

[6]Department of Neuroscience, Tufts University School of Medicine, Boston, MA, USA

[7]Neuroscience Center, Helsinki Institute of Life Sciences, University of Helsinki, Helsinki,
Finland

[*]Correspondence to B.A.W: bin.a.wang@outlook.com

Short title: Modulation of prefrontal coupling by activity in vmPFC

Abstract

Humans and animals can maintain constant payoffs in an uncertain environment by steadily re-

evaluating and flexibly adjusting current strategy, which largely depends on the interactions

between the prefrontal cortex (PFC) and mediodorsal thalamus (MD). While the ventromedial PFC

(vmPFC) represents the level of uncertainty (i.e., prior belief about external states), it remains

unclear how the brain recruits the PFC-MD network to re-evaluate decision strategy based on the

uncertainty. Here, we leverage nonlinear dynamic causal modeling on fMRI data to test how prior

belief-dependent activity in vmPFC gates the information flow in the PFC-MD network when

individuals switch their decision strategy. We show that the prior belief-related responses in

vmPFC had a modulatory influence on the connections from dorsolateral PFC (dlPFC) to both,

lateral orbitofrontal (lOFC) and MD. Bayesian parameter averaging revealed that only the

connection from the dlPFC to lOFC surpassed the significant threshold, which indicates that the

weaker the prior belief, the less was the inhibitory influence of the vmPFC on the strength of

effective connections from dlPFC to lOFC. These findings suggest that the vmPFC acts as a

gatekeeper for the recruitment of processing resources to re-evaluate the decision strategy in

situations of high uncertainty.

Keywords: Cognitive flexibility, prefrontal cortex (PFC), mediodorsal thalamus (MD), nonlinear

Dynamic Causal Modeling, uncertainty, prior belief

Author Summary

Prefrontal cortex (PFC) together with the mediodorsal thalamus (MD) jointly establish

computations critical for behavioral adaptations. While the task uncertainty (i.e., prior belief) was

represented by the ventromedial PFC (vmPFC), it remains unclear how the PFC-MD network

reallocates the processing resources to re-evaluate decision strategy under uncertainty. Here we

filled this gap by leveraging the Bayesian hierarchical modelling and nonlinear dynamic causal

modelling in an associative learning task. We found that in situations of high uncertainty, the prior

belief-related responses in vmPFC significantly strengthened effective connectivity from the

dorsolateral PFC to the orbitofrontal cortex, but not to the MD. The findings provide evidence for

the role of vmPFC in driving the re-evaluation of the decision strategy during behavioral

adaptations in situations of uncertainty.

Introduction

The prefrontal cortex (PFC) consists of several regions that are thought to play an important role in flexible decision-making. The dorsolateral PFC (dlPFC) is assumed to support executive functions (Jones & Graff-Radford, 2021), whereas the orbitofrontal cortex (OFC) appears to be involved in the flexible adaptation of behavior (Schoenbaum et al., 2021; Wang et al., 2023). The ventromedial PFC (vmPFC) was shown to be associated with the estimation of the value and saliency of sensory events and thereby guides value-based decision-making (Dundon et al., 2021). Interactions between these regions are thought to implement a variety of functions relevant to the flexibility by which cognitive resources are deployed. Interaction between medial PFC and the anterior cingulate cortex, for instance, are thought to contribute to the updating of beliefs about higher-order contextual associations (M. Botvinick et al., 1999; M. M. Botvinick et al., 2001; Sarafyazd & Jazayeri, 2019), whereas the interplay between OFC and vmPFC appears important for the prediction of value-based behavioral changes (Howard et al., 2016).

Recent studies in animals (Halassa & Kastner, 2017; Mukherjee et al., 2021; Saal et al., 2017; Schmitt et al., 2017) and humans (Hwang et al., 2017; Wen et al., 2021) revealed compelling evidence that cognitive flexibility also depends on interactions between distinct PFC subregions and the mediodorsal nucleus of the thalamus (MD). These studies have provided a complementary perspective on thalamic function challenging the classical notion of the thalamus as a sensory relay (Sherman, 2016). Combining hierarchical Bayesian modeling with fMRI in humans, we recently unveiled distinct prefrontal connections targeting the MD in relation to the participant's prior belief during associative learning (Wang & Pleger, 2020). The surprise about an unexpected outcome lowered the prior belief about the cue-target association and hence triggered a switch of the decision strategy through modulations of connections between MD and lateral OFC (Wang &

Pleger, 2020). These findings are supported by neuronal recordings obtained from mice. When

mice decided between different sets of learned cues that directed attention to either visual or

auditory targets, responses from the medial PFC reflected both, the individual cue as well as its

importance as a task-rule (Rikhye et al., 2018). The MD, on the other hand, appeared to facilitate

switching between cueing contexts by supporting or suppressing task-associated representations

in the PFC (Mukherjee et al., 2021; Rikhye et al., 2018; Schmitt et al., 2017). Together, these

findings from mice and humans emphasize crucial prefrontal-MD computations necessary for

learning stimulus-incentive associations.

We recently refined a previously developed computational prefrontal-MD model, inspired by cell

recordings obtained from mice (Rikhye et al., 2018), and trained it on human empirical data to test

whether the re-evaluation and adjustment of the decision strategy in both species follow the same

computational principles (Hummos et al., 2022). We found that the MD learned abstract

representations of its cortical inputs through biologically plausible Hebbian learning rules. Direct

feedback from MD to prefrontal cortex supported switching between behavioral strategies, while

lateral OFC (lOFC) constantly accumulated evidence for a strategy switch based on rapid Bayesian

estimation. Following these computational rules, our human fMRI results revealed that lOFC

directed its outputs to MD, rendering MD as the brain site which dynamically integrates crucial

inputs relevant for forming the behavioral strategy. These abstract MD representations and their

ability to reorganize prefrontal computations describes an efficient way how the brain utilizes the

MD to integrate inputs from other brain regions and to dynamically select between competing

behavioral strategies (Hummos et al., 2022).

Another prefrontal region, the vmPFC, was shown to encode the beliefs about outcome values,

which represents an intermediate signal required for efficient prefrontal-MD computations

underpinning the re-evaluation of decision strategy. In our prefrontal-MD model (Hummos et al.,

2022), we directed the vmPFC output on the executive dlPFC, as the most relevant receiver of

value-related information, but this dlPFC-vmPFC interaction was not directly supported by

empirical evidence. Using bilinear Dynamic Causal Modelling (DCM), we could not directly

model the vmPFC as an additional hub in the prefrontal-MD network since the vmPFC was not

among the regions involved in decision switches. In the present study, we therefore applied non-

linear DCM, which allowed us to capture the non-linear history of prior synaptic activity (Stephan

et al., 2008), and hence the modulatory (second-order) effects of prior belief-dependent responses

in vmPFC on the gain of activity within the prefrontal-MD network. We re-analyzed our

previously collected fMRI dataset and combined Bayesian hierarchical modeling with nonlinear

DCM on fMRI data to test how prior belief-related activity in the vmPFC tunes functional

couplings in the dlPFC-lOFC-MD network during the adjustment of the decision strategy.

Results

The participants' behavioral performance during the associative learning task has been presented in more detail in our previous studies (Hummos et al., 2022; Wang et al., 2020; Wang & Pleger, 2020). In brief, we found a significant effect of learning (one-way ANOVA: $F_{(4,27)} = 256$, $p < 0.001$) and post-hoc paired t-tests revealed that participants made significantly more correct predictions in learning blocks with high predictability (i.e., 90%/10%) than in blocks with low predictability (i.e., 70%/30%, $t(1,27) = 20.75$, $P < 0.001$, Bonferroni- corrected) or unpredictability (i.e., 50%, $t(1,27) = 24.70$, $p < 0.001$, Bonferroni-corrected). In addition, we found that the participants had a lower prior belief and required more time to respond ($p < 0.001$) in trials when they changed their decision strategy (*Switching*), compared to the trials where the decision strategy did not change (*Staying*).

To assess the parametric modulation by the prior beliefs, both $\hat{\mu}_2^{(k)}$ and $\hat{\mu}_3^{(k)}$ derived from HGF model were included as modulatory parameters in the GLM of fMRI data. We found that responses from the vmPFC reflected the prior belief about the sample-target contingency ($\hat{\mu}_2^{(k)}$, Fig. 1C, $x = -6$, $y = 62$, $z = -4$, $t(31) = 6.69$, $p < 0.05$, FWE whole-brain corrected). Besides the vmPFC, also the left precuneus ($x = -10$, $y = -56$, $z = 26$) and middle cingulate cortex ($x = 2$, $y = -12$, $z = 38$) represented significant prior belief-related activity ($p < 0.05$, FWE whole-brain corrected). The analysis of parametric effects related to prior belief about volatility ($\hat{\mu}_3^{(k)}$) did not reveal any significant brain regions ($p > 0.05$, FWE whole-brain corrected).

Fig. 1. The fMRI activity and dynamic causal modeling (DCM). **A.** Strategy switches (*Switching>Staying*) entailed significant BOLD activity in right dlPFC, right lOFC and right MD (Hummos et al., 2022). **B.** The bilinear DCM revealed that four connections (orange), i.e., from lOFC to MD, from dlPFC to lOFC, as well as between dlPFC and MD in both directions were significantly strengthened by strategy switches (*Switching > Staying*) (Hummos et al., 2022). The black connections indicate endogenous connections between brain areas. **C**. Upper panel: The prior beliefs over the course of the experiment. The red shaded area indicates the standard error of the mean (SEM) of prior beliefs over time. Lower panel: Prior belief-related vmPFC activity is projected on sagittal and axial MRI brain slices (p < 0.05, whole brain-FWE corrected). **D**. The non-linear DCM, which we used to test how prior belief-related activity in the vmPFC (blue) gated the information flow in the dlPFC-lOFC-MD network. (dlPFC–dorsolateral prefrontal cortex; lOFC–lateral orbitofrontal cortex; MD–thalamic mediodorsal nucleus; vmPFC-ventromedial prefrontal cortex)

As shown in our recent study (Hummos et al., 2022), the Bayesian model comparison across different plausible bilinear dlPFC-lOFC-MD models revealed that the adjustment of the decision strategy (*Switching*) significantly modulated connections from lOFC to MD, from dlPFC to lOFC, as well as between dlPFC and MD in both directions (Fig. 1B). In the present study, using nonlinear

DCM, we tested how prior belief related response from vmPFC modulated the connection strength

in the dlPFC-lOFC-MD network (Fig. 1D). More specifically, we tested which two of the four

projections were directly modulated by vmPFC activity. This resulted in six competing models

that were further evaluated with Bayesian model selection (BMS) (Supplementary Fig. S1A). BMS

revealed that the prior belief associated activity in vmPFC had a modulatory influence on the

projections originating in the dlPFC and targeting both, lOFC and MD (Supplementary Fig. S1B

and Fig. 2A). The posterior probability for the winning model was 0.48, surpassing the posterior

probabilities of the other tested models which ranged from 0.01 to 0.25 (Supplementary Fig. S1B).

This winning nonlinear model assumed a direct effect of prior belief on vmPFC whose activity

then mediated the gain of the dlPFC→lOFC and dlPFC→MD connections. We next applied

Bayesian parameter averaging (BPA), which computes a joint posterior density for the entire

sample and found that the prior belief-dependent vmPFC activity significantly influenced the

dlPFC→lOFC connection (posterior probability = 0.94, Fig. 2B). The modulations of the

dlPFC→MD connection showed a trend into the same direction, but the posterior probabilities

failed to surpass the 90% threshold (posterior probability = 0.81, Fig. 2B). The modulatory effect

was generally inhibitory, suggesting that the weaker the activity in vmPFC, and the lower the prior

belief (i.e., high task uncertainty), the stronger was the connection strength from dlPFC to lOFC.

Fig. 2. The modulatory effect on the dlPFC-lOFC-MD network by prior belief-associated vmPFC activity. A.
The Bayesian Model Selection across the possible six models revealed the winning model in which prior belief-
dependent vmPFC activity specifically influenced the connections originating in the dlPFC and targeting both, lOFC
and MD. The black connections indicate endogenous connections between brain areas. **B.** Bayesian parameter
averaging revealed that the prior belief-dependent vmPFC activity significantly inhibited the dlPFC→lOFC
connection (posterior probability = 0.94, solid red line) suggesting that the higher the activity in vmPFC, and the
stronger prior belief, the weaker was the connection strength from dlPFC to lOFC. The negative numbers shown next
to the projections and above the posterior probabilities index the inhibitory inputs per second (Hz). The dlPFC→MD
connection showed a trend into the same direction, but posterior probabilities failed to surpass the 90% threshold
(posterior probability = 0.81, dotted red line).

Discussion

In this study, we investigated how prior belief-related activity in the vmPFC gates the information flow in the dlPFC-lOFC-MD network underpinning the re-evaluation of decision strategy during associative learning. We showed that the strength of connections from dlPFC to lOFC fluctuated from trial to trial in relation to the level of task uncertainty, i.e., the prior beliefs about the cue-target contingency, signaled by inhibitory inputs from the vmPFC. The weaker the activity in vmPFC, and the stronger the task uncertainty (i.e., low prior belief), the stronger was the strength of effective connection from executive dlPFC to lOFC. These findings suggest that in situations with high task uncertainty, there is an increase in lOFC responses to inputs from dlPFC, which provides direct empirical evidence for the key role of prior belief-dependent synaptic plasticity in driving the re-evaluation of the decision strategy during flexible decision-making.

Representations of sensory stimuli are modulated by internal states and beliefs about the world, which are crucial for flexibly adjusting perception and behavior in humans and animals (Akrami et al., 2018; Snyder et al., 2015). The prior beliefs have been shown to warp neural representations in the frontal cortex, which allows the mapping of sensory inputs to motor outputs and to incorporate prior statistics in accordance with Bayesian inference (Sohn et al., 2019). These results uncover a simple and general principle whereby prior beliefs exert their influence on behavior by sculpting cortical latent dynamics. The failure of beliefs about the environmental state forms the prediction error and updates expectations for upcoming stimuli and associated rewards. In a previous study, we investigated how humans apply probabilistic computations, following Bayesian rules, to infer on joint brain representations of prior belief and decision switches (Wang et al., 2020). We found, that during such switches prior belief specifically modulated connectivity among the anterior insular cortex (AIC), the premotor cortex (PMd), and the inferior parietal lobule (IPL).

On a trial-by-trial basis, prior belief weakened connectivity between AIC and IPL when the

sensory stimulus was expected, whereas it strengthened connectivity between AIC and PMd when

the stimulus was unexpected (Wang et al., 2020). AIC has been shown to act as a core hub

modulating the interaction of bodily, attentional, and anticipatory sensory signals (Allen et al.,

2016; Craig, 2009; Sridharan et al., 2008). Our results furnish a picture in which AIC, in

conjunction with other brain regions, contributes not only to the coordination of expectation and

sensory inputs, but also to the integration of priors and prediction outcomes for updating beliefs

specifically supporting strategy switches during associative learning. The present findings extend

the scope of belief-related brain functions by the vmPFC. According to our non-linear DCM,

vmPFC plays a key role in exerting modulatory influences on prefrontal interactions, which does

not directly reflect prior belief such as the aforementioned AIC-network (Allen et al., 2016), but

which critically depends on prior belief-dependent information from the vmPFC to flexibly guide

strategy switches.

It has been shown that vmPFC plays a major role in sensory integration to achieve abstract and

conceptual interpretation of the environment (Petrides & Pandya, 2007). The vmPFC represents

signals more suited for subject-centered, internally driven motivational processes, whereas lOFC

encodes signals for evaluating environment-centered, externally driven motivational processes

(Bouret & Richmond, 2010). Functional interactions between vmPFC and lOFC in humans are

thought to implement important functions relevant to cognitive flexibility and the prediction of

value-based behavioral changes (Howard et al., 2016). In primates, the dlPFC has been shown to

be anatomically (Mackey & Petrides, 2010; Saleem et al., 2014) and functionally (Kahnt et al.,

2012) connected to the lOFC, and also in rodents, mPFC terminals - i.e., homologue of the primate

dlPFC - were detected in the lateral and ventrolateral OFC (Dalley et al., 2004). Rodent mPFC

projection neurons furthermore target the thalamus MD to regulate adaptive control to flexibly

optimize behavioral responses in goal-directed behavior and receive an MD innervation (Carlén,

2017; Hoover & Vertes, 2007). Together, these anatomical evidence from rodents and non-human

primates provide a solid anatomical basis for our winning nonlinear DCM model, in which the

responses of lOFC to the presynaptic input from dlPFC depend on the history of inputs that they

receive from vmPFC. Inconsistencies in the nomenclature and anatomical boundaries of PFC areas

have made it difficult to compare data and interpret findings across species, especially between

primates and rodents. In addition, we cannot exclude that the modulatory effect on the PFC

interactions is implemented through an intermediate region, for example the amygdala or the

striatum, given the fact that both receive many cortical inputs from prefrontal areas (Chang &

Grace, 2018; Middleton & Strick, 2002).

The weaker the prior belief related activity in vmPFC, the stronger is the connection strength of

the dlPFC projections, suggesting that the re-evaluation of decision strategy under high task

uncertainty requires further processing resources controlled by vmPFC. This gating mechanism

agrees well with the theoretical accounts of the free-energy principle (Friston, 2009). Accordingly,

perception optimizes predictions by minimizing free energy with respect to perceptual inference,

memory, attention and salience (Friston, 2010). Based on these principles, the high prior belief

about the environmental state in our study may have weakened the strength of the dlPFC-output

projections to minimize the engagement of additional cognitive resources and hence to lower the

energy costs for the decision process. The process that alters synaptic strengths with time constants

in the range of milliseconds to minutes, i.e., the so-called "short-term synaptic plasticity" (STP),

is proposed to be the underlying neurobiological mechanisms for this nonlinear modulatory effect,

including NMDA-controlled rapid trafficking of AMPA receptors (Diering & Huganir, 2018),

synaptic depression/facilitation (Zucker & Regehr, 2002) or "early long-term potentiation (LTP)"

(Frey & Morris, 1998).

There are limitations that should be considered when interpreting the current findings. First, single

unit recording in mice are well suited to assess how distinct neuronal populations in subregions of

the PFC and MD nuclei respond to specific task-related cues and contexts (Rikhye et al., 2018).

FMRI recordings instead represent comparably crude measures of neural activity capturing

hemodynamic responses only from large neuronal populations. These topographic inaccuracies,

especially for signals originating in small cortical and subcortical structures, represent a general

limitation of fMRI. Second, we did not split the MD into its subnuclei when we constructed the

network models. A recently developed MRI-based atlas provides masks for four of these

subregions (the medial (MDm), central (MDc), dorsal (MDd), and lateral (MDl)), created with

connectivity-based   methods   applied   to   the   high-resolution   data   from   the

Human Connectome Project (Li et al., 2022). In the current study with lower functional MRI

resolution and signal-to-noise ratio, it remained difficult to reliably capture these subregions.

Therefore, we considered the MD as a single region of interest. Considering these topographic

inaccuracies, the trend level effect we found for the modulatory influence on the dlPFC-MD

connection should be interpreted with caution. Third, although we focused on thalamocortical

connections, other regions of the broader PFC-MD network, including the basal ganglia or the

amygdala, could have also been involved in guiding cognitive flexibility. For example, the

cortical-basal ganglia-thalamocortical subnetworks have been shown to be involved in impaired

cognitive flexibility in patients with obsessive-compulsive disorder (Kim et al., 2022). Given the

above limitations, it will be one of the major challenges for future decision-making research to

extend the here proposed PFC-MD network by other key regions in the PFC, basal ganglia, and

other subcortical structures, through combining various imaging techniques and cross-species

approaches.

Materials and Methods

Participants and associative learning task

The analyses in this study were based on the previously collected dataset from twenty-eight healthy human participants (mean age ± SD: 25.3 ± 3.9 years, only male participants). The study was approved by the local ethics committee of the Ruhr-University Bochum. All participants gave written informed consent prior to participation. Demographics and the experimental design were described in more detail elsewhere (Hummos et al., 2022; Wang et al., 2020; Wang & Pleger, 2020).

In each trial, participants first received one out of two tactile cues for 500ms to the tip of their right index finger using an MRI-compatible Braille piezo stimulator with 8-pins (2x4 array) (Metec, Stuttgart, Germany). Subsequently, they had to predict (within 1300ms) whether the following tactile stimulus (i.e., target) will show the same pattern (e.g., 4 upper pins lifted) or the alternative pattern (4 lower pins lifted) by pressing one of two buttons (LumiTouch keypads, Photon Control) with the index or middle finger of the left hand. After an interval of 500-1500ms (jitter), the target was presented for 500ms which indicated whether the preceding prediction was correct or incorrect. A variable delay of 1500-3000ms separated trials. The predictability of the target stimulus was modulated by the strength of the cue–target contingency over time (i.e., strongly predictive blocks (90% and 10%), moderately predictive (70% and 30%), and nonpredictive (50%) blocks; either 30 or 40 trials per block to avoid predictability of the block onsets). The order of blocks was pseudorandomized and fixed across participants to ensure inter-subject comparability of the learning process. The fMRI experiment consisted of 350 trials in total, which were split into three runs, each lasting about 10 min.

In order to examine the flexibility in decision-making, we tested the adjustment of the decision

strategy, i.e., whether the statistical property of the environment (sample stimulus matches or

mismatches target stimulus) has been changed or not. To this end, we separated the trials across

all sample–target associations into two conditions: 1) *Switching* condition (122 ± 10 trials): trials

with decision switches across two successive trials, and 2) Staying condition (211 ± 17 trials):

trials without decision switches across two trials.

Estimating the prior belief with the Hierarchical Gaussian Filter

Human behavioral data were applied to a three-level Hierarchical Gaussian Filter (HGF) model

using the HGF toolbox (v5.2) as implemented in TAPAS (Translational Algorithms for Psychiatry-

Advancing Science, https://www.tnu.ethz.ch/en/software/tapas, Frässle et al., 2021), to calculate

the individual trial-wise prior belief about external states at different levels. The first level of the

HGF represents a sequence of inputs about the environmental states (i.e., whether the sample

stimulus matches the target stimulus or not), the second level represents the sample-target

contingency (i.e., the conditional probability, in logit space, of the target stimulus given the sample

cue), and the third level represents the log-volatility of the environment. Each of these hidden

states is assumed to evolve as a Gaussian random walk, such that its variance depends on the state

at the next higher level (Mathys et al., 2011).

In the HGF, at any level *i* of the hierarchy, the prior belief about the external state on trial k ($\hat{\mu}_i^{(k)}$)

is evolved from the posterior belief of the previous trial (k-1)

$$\hat{\mu}_i^{(k)} = \mu_i^{(k-1)} \tag{1}$$

313 The posterior belief on trial t ($\mu_i^{(k)}$) is updated based on the prediction error from the level below

314 ($\delta_{i-1}^{(k)}$) weighted by the precision of prediction ($\varphi_i^{(k)}$):

$$315 \qquad \mu_i^{(k)} = \mu_i^{(k-1)} + \varphi_i^{(k)} * \delta_{i-1}^{(k)} \qquad (2)$$

316 The precision of prediction $\varphi_i^{(k)}$ is updated with every trial and can be regarded as equivalent to a

317 dynamic learning rate in reward learning models, as follows:

$$318 \qquad \varphi_i^{(k)} = \frac{1}{1/\hat{\varphi}_i^{(k)} + \hat{\varphi}_{i-1}^{(k)}} \qquad (3)$$

$$319 \qquad \hat{\varphi}_{i-1}^{(k)} = \hat{\mu}_{i-1}^{(k-1)}\left(1 - \hat{\mu}_{i-1}^{(k-1)}\right) \qquad (4)$$

$$320 \qquad \hat{\varphi}_i^{(k)} = \varphi_i^{(k-1)} + \exp\left(\mu_{i+1}^{(k-1)} + \omega\right) \qquad (5)$$

321 where ω is a free parameter of the perceptual model in HGF, which determines the step size

322 between consecutive time steps. The prediction error $\delta_{i-1}^{(k)}$, which drives learning at the second

323 level of our HGF model, is defined as the difference between the actual outcome and its

324 estimated probability before the outcome:

$$325 \qquad \delta_{i-1}^{(k)} = \mu_{i-1}^{(k)} - \hat{\mu}_{i-1}^{(k-1)} \qquad (6)$$

FMRI Data Processing

FMRI data were collected on a Philips Achieva 3.0 T X-series scanner using a 32-channel head

coil. For functional imaging, we used a T2*- weighted echoplanar imaging (EPI) sequence (voxel

size, 2x2x3 mm³; field of view, 224x224 mm²; interslice gap, 0.6 mm; TR= 2800 ms; TE = 36 ms)

to acquire 36 transaxial slices covering the whole brain. Pre- and post-processing of the fMRI data

was done using the Statistical Parametric Mapping software SPM12 (Wellcome Department of

Imaging Neuroscience, University College London, London, UK;

http://www.fil.ion.ucl.ac.uk/spm) implemented in MATLAB R2022a (MathWorks). All fMRI

images were first applied to slice time correction, spatial realignment, and normalization to the

MNI template using the unified segmentation approach (Ashburner & Friston, 2005). Finally,

normalized images were spatially smoothed using a Gaussian filter with a full-width half-

maximum kernel of 6 mm. Data were high pass filtered at 1/128 Hz. For each participant, we

conducted a first level general linear model (GLM). Events were time-locked to the onset of the

presentation of the cue stimulus using stick functions and split into two regressors, one for *Staying*

(no strategy switches) and the other one for *Switching* trials. For each of these two regressors, two

parametric modulators of prior belief were defined. The first parametric modulator was prior belief

about the sample-target contingency ($\hat{\mu}_2^{(k)}$). The second modulator was prior belief about the

volatility ($\hat{\mu}_3^{(k)}$), orthogonalized with respect to $\hat{\mu}_2^{(k)}$. Please note that the absolute value of prior

belief ($\left|\hat{\mu}_i^{(k)}\right|$) was used to enter the GLM as parametric modulators. Invalid trials (i.e., missing or

late responses) were modeled separately. Furthermore, six head motion parameters, as estimated

during the realignment procedure, were added as regressors of no interest to minimize false-

positive activations due to task-correlated motion.

Using the GLM, we investigated prefrontal and thalamic responses when switching the decision

strategy using the contrast '*Switching> Staying*'. The detailed data analysis for this comparison

and corresponding results have been reported in our recent paper (Hummos et al., 2022). In current

study, we primarily analyzed the main effect of parametric modulation by prior belief (both $\hat{\mu}_2^{(k)}$

and $\hat{\mu}_3^{(k)}$) for all trials using the GLM. The respective t-contrast images of the modulatory effect

for each subject were applied to the group-level one-sample *t* test (p < 0.05, family-wise error

(FWE) corrected for the whole brain).

Nonlinear Dynamic Causal Models (DCM)

In our previous study (Hummos et al., 2022), the fMRI GLM analyses revealed two prefrontal

regions, i.e., right dlPFC and lOFC, together with the MD thalamus, which were all significantly

modulated by decision switches (Fig. 1A). The Bayesian parameter averaging (BPA) across

participants revealed that connections from lOFC to MD, from dlPFC to lOFC, as well as between

dlPFC and MD in both directions were all significantly strengthened by strategy switches (Fig.

1B). In the current study, we show that the vmPFC reflects the value of the prediction, i.e., the

prior belief about the sample-target contingency (see results section). Based on these findings, we

next questioned how increased BOLD responses in the vmPFC, which were positively related to

stronger prior belief, modulated the connection strength of the prefrontal-MD network, thus

enhancing its computational efficiency and facilitating strategy switches. To test this, we

constructed a nonlinear DCM including right dlPFC, right lOFC, right MD and vmPFC, and

compared several alternative vmPFC modulations. For each brain region, subject-specific time

series were extracted from the nearest local maximum within a sphere with a radius of 8 mm

centered on each node's group maximum. The first Eigenvariate was extracted across all voxels

surviving $p = 0.05$, uncorrected, within a 4 mm sphere centered on the individual peak voxel. The

resulting BOLD time series were adjusted for effects of no interest (e.g., invalid trials, and

movement parameters).

The basic architecture of the model, shown in Fig. 1B and in Hummos et al., 2022, included the

driving sensory (i.e., tactile) input directed to the dlPFC, as well as the four connections that were

all significantly strengthened by strategy switches (i.e., dlPFC to lOFC, lOFC to MD, as well as

between dlPFC and MD in both directions). With the nonlinear DCM we extended this model by

the modulation of these connections through the activity in the vmPFC, which was driven by the

trial-by-trial prior belief derived from the HGF model (Fig. 1D). We specified the model space

with the modulatory influence of the vmPFC on different connections. More specifically, we tested

the modulatory influence of prior belief related vmPFC on two out of the four significant

connections within the dlPFC-lOFC-MD network, resulting in six models that were compared to

each other (Supplementary Fig. S1A). The fixed-effects Bayesian model selection (BMS) was used

to assess the most likely model among the six competing models. Parameters of the winning model

were then summarized by Bayesian parameter averaging (BPA), which computes a joint posterior

density for the entire group by combining the individual posterior densities. A posterior probability

criterion of 90% was considered to reflect significant modulatory effect on the connections.

Data and code availability

The data and code that were applied to assess the findings of this study are available from the

corresponding author upon reasonable request.

Acknowledgements

This work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research

Foundation): Project number 122679504 - SFB 874 'Integration and Representation of Sensory

Processes' (to B.A.W. and B.P.), the DFG grant with the project number PL602/6-1' Prefrontal-

thalamic control of cognitive flexibility – from mice to humans' (to B.P.) and National Natural

Science Foundation of China: Project number 32200867 (to B.A.W.).

Conflict of Interest

The authors declare no competing financial interests.

References

Akrami, A., Kopec, C. D., Diamond, M. E., & Brody, C. D. (2018). Posterior parietal cortex represents sensory history and mediates its effects on behaviour. *Nature*, *554*, 368–372. https://doi.org/10.1038/nature25510

Allen, M., Fardo, F., Dietz, M. J., Hillebrandt, H., Friston, K. J., Rees, G., & Roepstorff, A. (2016). Anterior insula coordinates hierarchical processing of tactile mismatch responses. *NeuroImage*, *127*, 34–43. https://doi.org/10.1016/j.neuroimage.2015.11.030

Ashburner, J., & Friston, K. J. (2005). Unified segmentation. *NeuroImage*, *26(3)*, 839–851. https://doi.org/10.1016/j.neuroimage.2005.02.018

Botvinick, M. M., Carter, C. S., Braver, T. S., Barch, D. M., & Cohen, J. D. (2001). Conflict monitoring and cognitive control. *Psychological Review*, *108(3)*, 624–652. https://doi.org/10.1037/0033-295X.108.3.624

Botvinick, M., Nystrom, L. E., Fissell, K., Carter, C. S., & Cohen, J. D. (1999). Conflict monitoring versus selection-for-action in anterior cingulate cortex. *Nature*, *402*(6758), 179–181. https://doi.org/10.1038/46035

Bouret, S., & Richmond, B. J. (2010). Ventromedial and orbital prefrontal neurons differentially encode internally and externally driven motivational values in monkeys. *The Journal of Neuroscience*, *30*(25), 8591–8601. https://doi.org/10.1523/JNEUROSCI.0049-10.2010

Carlén, M. (2017). What constitutes the prefrontal cortex? *Science (New York, N.Y.)*, *358*(6362), 478–482. https://doi.org/10.1126/science.aan8868

Chang, C.-H., & Grace, A. A. (2018). Inhibitory Modulation of Orbitofrontal Cortex on Medial Prefrontal Cortex-Amygdala Information Flow. *Cerebral Cortex (New York, N.Y. : 1991)*, *28*(1), 1–8. https://doi.org/10.1093/cercor/bhw342

Craig, A. D. (2009). How do you feel - now? The anterior insula and human awareness. *Nature Reviews Neuroscience*, *10*(1), 59–70. https://doi.org/10.1038/nrn2555

Dalley, J. W., Cardinal, R. N., & Robbins, T. W. (2004). Prefrontal executive and cognitive functions in rodents: neural and neurochemical substrates. *Neuroscience and Biobehavioral Reviews*, *28*(7), 771–784. https://doi.org/10.1016/j.neubiorev.2004.09.006

Diering, G. H., & Huganir, R. L. (2018). The AMPA Receptor Code of Synaptic Plasticity. *Neuron*, *100*(2), 314–329. https://doi.org/10.1016/j.neuron.2018.10.018

Dundon, N. M., Shapiro, A. D., Babenko, V., Okafor, G. N., & Grafton, S. T. (2021). Ventromedial Prefrontal Cortex Activity and Sympathetic Allostasis During Value-Based Ambivalence. *Frontiers in Behavioral Neuroscience*, *15*, 615796. https://doi.org/10.3389/fnbeh.2021.615796

Frässle, S., Aponte, E. A., Bollmann, S., Brodersen, K. H., Do, C. T., Harrison, O. K., Harrison, S. J., Heinzle, J., Iglesias, S., Kasper, L., Lomakina, E. I., Mathys, C., Müller-Schrader, M., Pereira, I., Petzschner, F. H., Raman, S., Schöbi, D., Toussaint, B., Weber, L. A., … Stephan, K. E. (2021). TAPAS: An Open-Source Software Package for Translational Neuromodeling and Computational Psychiatry. *Frontiers in Psychiatry*, *12*, 680811. https://doi.org/10.3389/fpsyt.2021.680811

Frey, U., & Morris, R. G. (1998). Synaptic tagging: implications for late maintenance of hippocampal long-term potentiation. *Trends in Neurosciences*, *21*(5), 181–188. https://doi.org/10.1016/s0166-2236(97)01189-2

Friston, K. (2009). The free-energy principle: a rough guide to the brain? *Trends in Cognitive Sciences*, *13*(7), 293–301. https://doi.org/10.1016/j.tics.2009.04.005

Friston, K. (2010). The free-energy principle: A unified brain theory? *Nature Reviews*
Neuroscience, *11*, 127–138. https://doi.org/10.1038/nrn2787
Halassa, M. M., & Kastner, S. (2017). Thalamic functions in distributed cognitive control.
Nature Neuroscience, *20*(12), 1669–1679. https://doi.org/10.1038/s41593-017-0020-1
Hoover, W. B., & Vertes, R. P. (2007). Anatomical analysis of afferent projections to the medial
prefrontal cortex in the rat. *Brain Structure & Function*, *212*(2), 149–179.
https://doi.org/10.1007/s00429-007-0150-4
Howard, J. D., Kahnt, T., & Gottfried, J. A. (2016). Converging prefrontal pathways support
associative and perceptual features of conditioned stimuli. *Nature Communications*, *7*,
11546. https://doi.org/10.1038/ncomms11546
Hummos, A., Wang, B. A., Drammis, S., Halassa, M. M., & Pleger, B. (2022). Thalamic
regulation of frontal interactions in human cognitive flexibility. *PLoS Computational*
Biology, *18*(9), e1010500. https://doi.org/10.1371/journal.pcbi.1010500
Hwang, K., Bertolero, M. A., Liu, W. B., & D'Esposito, M. (2017). The Human Thalamus Is an
Integrative Hub for Functional Brain Networks. *The Journal of Neuroscience*, *37*(23),
5594–5607. https://doi.org/10.1523/JNEUROSCI.0067-17.2017
Kahnt, T., Chang, L. J., Park, S. Q., Heinzle, J., & Haynes, J.-D. (2012). Connectivity-based
parcellation of the human orbitofrontal cortex. *The Journal of Neuroscience : The Official*
Journal of the Society for Neuroscience, *32*(18), 6240–6250.
https://doi.org/10.1523/JNEUROSCI.0257-12.2012
Kim, T., Kim, M., Jung, W. H., Kwak, Y. bin, Moon, S.-Y., Kyungjin Lho, S., Lee, J., & Kwon,
J. S. (2022). Unbalanced fronto-pallidal neurocircuit underlying set shifting in obsessive-
compulsive disorder. *Brain : A Journal of Neurology*, *145*(3), 979–990.
https://doi.org/10.1093/brain/awab483
Li, K., Fan, L., Cui, Y., Wei, X., He, Y., Yang, J., Lu, Y., Li, W., Shi, W., Cao, L., Cheng, L., Li,
A., You, B., & Jiang, T. (2022). The human mediodorsal thalamus: Organization,
connectivity, and function. *NeuroImage*, *249*, 118876.
https://doi.org/10.1016/j.neuroimage.2022.118876
Mackey, S., & Petrides, M. (2010). Quantitative demonstration of comparable architectonic areas
within the ventromedial and lateral orbital frontal cortex in the human and the macaque
monkey brains. *The European Journal of Neuroscience*, *32*(11), 1940–1950.
https://doi.org/10.1111/j.1460-9568.2010.07465.x
Mathys, C., Daunizeau, J., Friston, K., & Stephan, K. (2011). A Bayesian Foundation for
Individual Learning Under Uncertainty. *Frontiers in Human Neuroscience*, *5*, 1–20.
https://www.frontiersin.org/article/10.3389/fnhum.2011.00039
Middleton, F. A., & Strick, P. L. (2002). Basal-ganglia "projections" to the prefrontal cortex of
the primate. *Cerebral Cortex (New York, N.Y. : 1991)*, *12*(9), 926–935.
https://doi.org/10.1093/cercor/12.9.926
Mukherjee, A., Lam, N. H., Wimmer, R. D., & Halassa, M. M. (2021). Thalamic circuits for
independent control of prefrontal signal and noise. *Nature*, *600*(7887), 100–104.
https://doi.org/10.1038/s41586-021-04056-3
Petrides, M., & Pandya, D. N. (2007). Efferent association pathways from the rostral prefrontal
cortex in the macaque monkey. *The Journal of Neuroscience : The Official Journal of the*
Society for Neuroscience, *27*(43), 11573–11586.
https://doi.org/10.1523/JNEUROSCI.2419-07.2007

Rikhye, R. v., Gilra, A., & Halassa, M. M. (2018). Thalamic regulation of switching between
cortical representations enables cognitive flexibility. *Nature Neuroscience*, *21*, 1753–1763.
https://doi.org/10.1038/s41593-018-0269-z
Saal, H. P., Delhaye, B. P., Rayhaun, B. C., & Bensmaia, S. J. (2017). Simulating tactile signals
from the whole hand with millisecond precision. *Proceedings of the National Academy of*
Sciences, *114*(28), E5693–E5702. https://doi.org/10.1073/pnas.1704856114
Saleem, K. S., Miller, B., & Price, J. L. (2014). Subdivisions and connectional networks of the
lateral prefrontal cortex in the macaque monkey. *The Journal of Comparative Neurology*,
522(7), 1641–1690. https://doi.org/10.1002/cne.23498
Sarafyazd, M., & Jazayeri, M. (2019). Hierarchical reasoning by neural circuits in the frontal
cortex. *Science*, *364(6441):* https://doi.org/10.1126/science.aav8911
Schmitt, L. I., Wimmer, R. D., Nakajima, M., Happ, M., Mofakham, S., & Halassa, M. M.
(2017). Thalamic amplification of cortical connectivity sustains attentional control. *Nature*,
545(7653), 219–223. https://doi.org/10.1038/nature22073
Schoenbaum, G., Khamassi, M., Pessiglione, M., Gottfried, J. A., & Murray, E. A. (2021). The
magical orbitofrontal cortex. *Behavioral Neuroscience*, *135*(2), 108.
https://doi.org/10.1037/bne0000470
Sherman, S. M. (2016). Thalamus plays a central role in ongoing cortical functioning. *Nature*
Neuroscience, *19(4)*, 533–541. https://doi.org/10.1038/nn.4269
Snyder, J. S., Schwiedrzik, C. M., Vitela, A. D., & Melloni, L. (2015). How previous experience
shapes perception in different sensory modalities. *Frontiers in Human Neuroscience*, *9*,
594. https://doi.org/10.3389/fnhum.2015.00594
Sohn, H., Narain, D., Meirhaeghe, N., & Jazayeri, M. (2019). Bayesian Computation through
Cortical Latent Dynamics. *Neuron*, *103(5)*, 934–947.
https://doi.org/10.1016/j.neuron.2019.06.012
Sridharan, D., Levitin, D. J., & Menon, V. (2008). A critical role for the right fronto-insular
cortex in switching between central-executive and default-mode networks. *Proceedings of*
the National Academy of Sciences of the United States of America, *105*(34), 12569–12574.
https://doi.org/10.1073/pnas.0800005105
Stephan, K. E., Kasper, L., Harrison, L. M., Daunizeau, J., den Ouden, H. E. M., Breakspear, M.,
& Friston, K. J. (2008). Nonlinear dynamic causal models for fMRI. *NeuroImage*, *42(2)*,
649–662. https://doi.org/10.1016/j.neuroimage.2008.04.262
Wang, B. A., & Pleger, B. (2020). Confidence in Decision-Making during Probabilistic Tactile
Learning Related to Distinct Thalamo-Prefrontal Pathways. *Cerebral Cortex*, *30*, 4677–
4688,. https://doi.org/10.1093/cercor/bhaa073
Wang, B. A., Schlaffke, L., & Pleger, B. (2020). Modulations of Insular Projections by Prior
Belief Mediate the Precision of Prediction Error during Tactile Learning. *The Journal of*
Neuroscience : The Official Journal of the Society for Neuroscience, *40*, 3827–3837.
https://doi.org/10.1523/JNEUROSCI.2904-19.2020
Wang, B. A., Veismann, M., Banerjee, A., & Pleger, B. (2023). Human orbitofrontal cortex
signals decision outcomes to sensory cortex during behavioral adaptations. *Nature*
Communications, *14*(1), 3552. https://doi.org/10.1038/s41467-023-38671-7
Wen, X., Li, W., Liu, Y., Liu, Z., Zhao, P., Zhu, Z., & Wu, X. (2021). Exploring communication
between the thalamus and cognitive control-related functional networks in the cerebral
cortex. *Cognitive, Affective, & Behavioral Neuroscience*.

Zucker, R. S., & Regehr, W. G. (2002). Short-term synaptic plasticity. *Annual Review of*
Physiology, *64*, 355–405. https://doi.org/10.1146/annurev.physiol.64.092501.114547
537
538