Title: Responses of the hyper-diverse community of canopy-dwelling

Hymenoptera to oak decline

Running title: Canopy Hymenoptera and oak decline

Authors: E. Le Souchu[1], J. Cours[2,3,4], T. Cochenille[1], C. Bouget[4], S. Bankhead-Dronnet[1], Y. Braet[5], P.

Burguet[6], C. Gabard[1], C. Galkowski[7], B. Gereys[8], F. Herbrecht[9], B. Joncour[1], E. Marhic[10], D. Michez[11], P.

Neerup Buhl[12], T. Noblecourt[13], D. G. Notton[14], W. Penigot[15], J.-Y. Rasplus[16], T. Robert[17], A.

Staverlokk[18], C. Vincent-Barbaroux[1], A. Sallé[1]

Corresponding author: elodie.lesouchu2@gmail.com

Author details:

[1] Laboratoire de Biologie des Ligneux et des Grandes Cultures, INRAE, University of Orléans, Orléans,

France

[2] Department of Biological and Environmental Science, University of Jyväskylä, Jyväskylä, Finland

[3] School of Resource Wisdom, University of Jyväskylä, Jyväskylä, Finland

[4] INRAE, UR EFNO, Domaine des Barres, Nogent-sur-Vernisson, France

[5] Royal Belgian Institute of Natural Sciences, O.D. Phylogeny and Taxonomy, Entomology, Vautier

street 29, B-1000 Brussels, Belgium; Unité d'Entomologie fonctionnelle et évolutive, Gembloux Agro-

Bio Tech, Université de Liège, Passage des Déportés 2, B-5030 Gembloux, Belgium

[6] 18 rue du Massaud, 63118 Cébazat, France

[7] Société Linnéenne de Bordeaux, Bordeaux, France

[8] 4 Chemin des Escaranches, 04700 Oraison, France

[9] 2, le Bois Barré, 35390 Sainte-Anne-sur-Vilaine, France

[10] 25 rue du Maréchal de Lattre de Tassigny, 78780 Maurecourt, France

[11] University of Mons, Laboratory of Zoology, Research Institute for Biosciences, Place du parc, 20,

7000 Mons, Belgium

[12] Zoological Museum, Department of Entomology, University of Copenhagen, Universitet sparken 15,

DK-2100 Copenhagen, Denmark

[13] Réseau entomologie de l'Office National des Forêts, 14 rue de la Garenne, F-11190 Antugnac,
France

[14] Invertebrates Section, Natural Sciences Department, National Museums Collection Centre, 242
West Granton Road, Granton, Edinburgh EH5 1JA, Scotland

[15] 24 rue du Puech de la Borie, 71240 Saint-Juéry, France

[16] CBGP, INRAE, CIRAD, IRD, Montpellier SupAgro, Université de Montpellier, 34988 Montpellier,
France

[17] 12 rue du 20^e bataillon 54120 Baccarat, France

[18] Norwegian Institute for Nature Research, P.O.Box 5685 Torgard, NO-7485 Trondheim, Norway

ABSTRACT

1. Forest decline and dieback are growing phenomena worldwide, resulting in severe, large-
scale degradation of the canopy. This can profoundly alter forest microclimates and the
provision of trophic resources and microhabitats for forest arthropods, especially those
dwelling in the canopy.

2. In 2019, we assessed the effect of oak decline on the community of canopy-dwelling
Hymenoptera. We selected 21 oak stands, and 42 plots, located in three forests in France,
presenting contrasting levels of decline. Insects were sampled at the canopy level with green
multi-funnel and flight-interception traps.

3. We collected a particularly diverse community of 19,289 insect individuals belonging to
918 taxa, eight larval trophic guilds and five nesting guilds.

4. Oak decline had no effect on the abundance or richness of the overall community, but
significantly reshaped the community assemblages. Decline had contrasting effects
depending on the taxa and guilds considered. Parasitoids were more abundant at
intermediate levels of decline severity. Taxa depending on ground-related resources and
microhabitats were promoted. Saproxylic taxa were more diverse while xylophagous insects
were negatively impacted.

5. Reduced leaf area index promoted several guilds, and the diversity of the overall
community. While an increasing tree mortality rate enhanced the abundance and diversity of
deadwood resources, it had negative impacts on Hymenoptera guilds. Our results suggest
that micro-environmental changes at the ground-level due to canopy decline have major
cascading effects on the communities of canopy-dwelling Hymenoptera.

6. Our study highlights the relevance of studying Hymenoptera communities to investigate
the outcomes of disturbances on forest biodiversity.

Keywords: trophic guild, habitat guild, deadwood, canopy openness, temperate forest

INTRODUCTION

Global change is currently increasing the frequency, severity and spatial extent of major forest

disturbances in Europe such as droughts, windstorms and wildfires (Seidl et al., 2017; Samaniego et

al., 2018; Spinoni et al., 2018). Disturbances can markedly affect the amount, diversity and

distribution of key trophic resources and microhabitats for forest arthropods (Cours et al., 2023). One

of the main drivers of these changes is the degradation of the forest canopy (Sallé et al., 2021; Cours

et al., 2023). A reduction in canopy cover can result from direct disturbance impacts on the trees

themselves, but also from subsequent forest dieback or decline, as a progressive loss of tree vigour

generally translates into crown dieback (Sallé et al., 2021). Decline and dieback can consequently

affect the quality and quantity of foliage, and reduce the production of flowers, fruits and seeds in

the crown (Ishii et al., 2004; Günthardt-Goerg et al., 2013; Hu et al., 2013). On the other hand, they

may also increase the amount and diversity of deadwood resources and tree-related microhabitats

such as fruiting bodies of opportunistic fungi, trunk cavities or perched deadwood (Sallé et al., 2021;

Larrieu et al., 2022; Zemlerová et al., 2023). Increased canopy openness also considerably affects

microclimates and trophic resources for arthropods in the lower strata of forest ecosystems for

instance, by promoting the accumulation and diversification of floral resources in the herbaceous

layer, or of deadwood resources on the ground (Cours et al., 2023). Consequently, forest dieback and

decline can promote forest biodiversity to a certain extent by increasing structural complexity at

multiple scales, as predicted by the pulse dynamics theory (Jentsch and White, 2019; Cours et al.,

2022). These modifications can profoundly reshape communities of forest arthropods (Viljur et al.,

2022; Cours et al., 2023). The impact of canopy dieback on forest arthropods has been investigated

for several taxonomic groups and functional guilds, including some specifically dwelling in the canopy

(Martel & Mauffette, 1997; Stone et al., 2010; Sallé et al., 2020; Vincent et al., 2020). While the

responses of canopy arthropods is largely mediated by their functional and/or trophic guild (e.g.,

Sallé et al., 2020), idiosyncratic responses still occur among the taxa of a same guild or taxonomic

group (e.g., Vincent et al., 2020). Consequently, the impacts of disturbance-driven changes in canopy

structure on arthropod communities are still difficult to predict, especially in the largely under-

studied temperate forest canopy (Sallé et al., 2021; Cours et al., 2023). Considering that canopy

mortality rate has been on the rise for several decades in Europe (Senf et al., 2018), further studies

on how decline and dieback affect canopy-dwelling arthropods are urgently needed, especially for

the taxonomic groups that have received limited attention to date, like the Hymenoptera.

Hymenoptera is a hyper-diverse group of insects, sometimes regarded as the most speciose animal

order (Forbes et al., 2018). This group plays a wide variety of functional roles in temperate forests.

Ants and predatory and parasitic wasps help to regulate forest pests (Hilszczański, 2018). Bees and

wasps pollinate forest plants (Motten, 1986; Yumoto, 1987), while ants can disperse seeds (Rico-Gray
and Oliveira, 2008). Hymenoptera nests, galleries and galls also provide microhabitats for many
species (Stone et al., 2002). Wood-nesting bees and ants, as well as the larvae of xylophagous
species, contribute to the decomposition process of wood (Ulyshen, 2016). Phyllophagous and
xylophagous species can be major pests for trees (Lyytikäinen-Saarenmaa and Tomppo, 2002;
Slippers et al., 2015). The Hymenoptera, especially parasitic wasps, can be used as bioindicators in
forest ecosystems (Maleque et al., 2009). Their communities can be largely influenced by stand-
related variables such as stand composition (Fraser et al., 2007; Ulyshen et al., 2011a), deadwood
resources (Hilszczański et al., 2005; Ulyshen et al., 2011a; Jonsell et al., 2023) and silvicultural
practices (Lewis and Whitfield, 1999; Hilszczański et al., 2005; Rappa et al., 2023). Pollinators in
particular are sensitive to canopy opening and the subsequent positive impacts on the floral
resources in the herbaceous layer and on ground-nesting sites (Wermelinger et al., 2017; Burkle et
al., 2019; Viljur et al., 2022; Cours et al., 2023).

Many Hymenoptera use the canopy as a hunting ground, a nesting site or a breeding site (Ulyshen,
2011). The canopy can also act as a dispersal corridor for species with a preference for open habitats
(Pucci, 2008; Di Giovanni et al., 2017). As for other forest insects, the Hymenoptera community
exhibits a conspicuous vertical stratification in temperate forests (Smith et al., 2012). Certain
ecological guilds like the predatory and parasitic wasps, and the pollinators, differ significantly
between the ground level and the canopy (Sobek et al., 2009; Ulyshen et al., 2011b; Urban-Mead et
al., 2021), and several taxa are specific to the canopy layer (Ulyshen et al., 2010; Di Giovanni et al.,
2017). While some studies have investigated the vertical stratification of certain taxonomic groups or
ecological guilds of bees and wasps in temperate forests, few have taken into account the whole
Hymenoptera community (e.g., Vance et al., 2007), and none have considered this community's
response to forest decline.

In the present study, we investigated the impact of oak decline on the community of canopy-dwelling
Hymenoptera. We chose to focus on oak forests because they are the dominant forest ecosystem in
western Europe; they shelter a diverse entomofauna (Kennedy and Southwood, 1984); finally, they
have suffered from historic decline in France (Nageleisen, 2008), and have recently undergone severe
summer droughts (in 2018 and 2019) (Saintonge and Goudet, 2020). Our first objective was to
characterise the little-known community of canopy-dwelling Hymenoptera in temperate oak forests.
Our second objective was to assess how the severity of forest decline affected the richness and
taxonomic composition of this community. Our third objective was to evaluate the effects of decline
on certain ecological guilds of Hymenoptera, and how changes in stand-related variables drive those
effects. We focused on the trophic guilds of larvae and nesting guilds. For these guilds, we

formulated three hypotheses. First (H1), we expected to find positive effects of increasing decline
severity on pollinivorous/nectarivorous and phyllophagous species associated with host plants other
than oak due to the increased diversity and amount of plant-resources in the herbaceous and shrub
layers, commonly observed when canopy closure decreases (Romey et al., 2007; Lu et al., 2019;
Cacciatori et al., 2022). Likewise, we expected that soil-nesters would be promoted by more open
conditions resulting in warmer soil temperatures. We also hypothesised that xylophagous, saproxylic
and wood-nester taxa would benefit from oak dieback and the subsequent increased amount and
diversity of deadwood. Conversely (H2), we expected to find negative effects on the guilds that feed
on oak foliage, for example, gall-inducing and phyllophagous taxa, due to the decrease in the quality
and quantity of leaves. Finally (H3), we hypothesised that decline severity would have contrasted
effects on parasitoids and polyphagous taxa, depending on their hosts and trophic resources.

MATERIAL AND METHODS

Study sites

Three oak-dominated (*Quercus petraea* (Matt.) Liebl. and *Quercus robur* L.) French forests were
sampled in 2019 (Fig. 1, Tab. S1): one near Orléans (Forêt domaniale d'Orléans, 47°98′97″81N,
1°95′44″E), one near Vierzon (Forêt domaniale de Vierzon, 47°26′89″N, 02°10′74″E) and one near
Marcenat (Forêt domaniale de l'Abbaye, 46°21′12″N, 3°36′13″E). All the forests were managed by
the French National Forest Service. The Orléans forest covers 35,000 ha and is dominated by oaks
(47%) and pines (*Pinus sylvestris* L., 51%). The Vierzon forest covers 7,500 ha and is also dominated
by oaks (61%) and pines (*P. sylvestris* and *P. pinaster* Ait., 31%). The Marcenat forest covers 2,100 ha
and is dominated by oaks (71%), pines (11%) and Douglas fir (8%). The Vierzon forest suffered the
most from decline. Several decline events had been recorded there since 1920, and massive oak
mortality had occurred in several stands (Douzon, 2006). In the Marcenat forest, two summer
droughts (2018-2019) had led to severe decline in several stands (Saintonge and Goudet, 2020).
Conversely, no major decline event had been reported in the Orléans forest.

A total of 21 oak-dominated stands with a minimum surface area of 3 ha were selected in these
forests: nine in Orléans, nine in Vierzon and three in Marcenat (Fig. 1). In each forest, we selected
healthy, moderately declining or severely declining stands (3 for each category in Orléans and
Vierzon, 1 in Marcenat). In each of these stands, we selected one declining and one healthy tree, at
least 50 m from each other, in which to hang our traps. In each stand, we recorded average tree
height, density and diameter, and the stand tree species composition (Tab. S1). We also evaluated
the level of decline according to the DEPERIS protocol (Goudet and Nageleisen, 2019). This protocol

makes it possible to quantify decline at the tree level by evaluating the percentage of dead branches
and ramification loss in the crown. Based on these criteria, each tree was assigned to a decline class
ranging from A (no decline) to F (severe decline). Trees in the A–C classes were considered healthy.
Trees in the D–F classes were considered in decline. We used this protocol to describe tree decline at
two embedded spatial scales: (i) the ten closest oaks surrounding each trap-bearing tree, hereafter
referred as a *"plot"*; and (ii) 30 oaks in the stand where the two plots were located (i.e., the 20 trees
in the two monitored plots and ten trees in an additional plot located between the two monitored
plots), hereafter referred as a *"stand"*. Decline level was evaluated in January or February in 2019
and 2020, i.e., before and after Hymenoptera sampling, and we averaged the 2019 and 2020 values
for our analyses. We classified our plots and stands into three decline categories depending on the
proportion of declining oak trees. When the proportion of declining trees was < 30%, we considered
the plot/stand *"healthy"*; when the proportion was between 30% and 60%, the plot/stand was
considered *"moderately declining"*; and when the proportion was > 60%, the plot/stand was
considered *"severely declining"*.

In the early summer of 2019, we used a plant canopy analyzer LAI-2200C with a half view cap
attached to the optical sensor to quantify the leaf area index (LAI) in the 12 stands in the Orléans and
Vierzon forests. More than 50 measurements were carried out at each plot at twilight below the
canopy along two diagonal transects; these values were then compared with above-canopy readings
taken outside the plot. We applied an ellipsoidal post-processing correction to the first three zenith
angles and used the FV2200 software to compute LAI by the gap fraction method. Then, in October
2020, we used a Bitterlich relascope with an opening angle corresponding to counting factor n° 1
(ratio 1/50) with a mean plot area of about 0.3 ha (Bouget et al., submitted) to characterise the
woody resources in each of the 18 plots in the Orléans and Vierzon forests. For each tree, we
recorded its status (i.e., living tree, standing dead tree (snag) or downed dead tree (log)), decay
stage, tree species and diameter at breast height (DBH). We defined the mortality rate as the basal
area of standing and lying deadwood over the total basal area of all the recorded trees (i.e., the sum
of both living and dead trees). We calculated the overall volume of living trees and deadwood. For
deadwood, we calculated the volume for each status (standing or downed deadwood), decay stage
(slightly vs. highly decayed) and diameter class (deadwood DBH < 40 cm vs. ≥ 40 cm). We compiled all
this information (i.e., status, species, decay stage and DBH) to calculate deadwood diversity. Finally,
we used the dbFD function from the FD R-package to calculate the mean status value of the woody
resources and the dispersion value for each plot, based on an ordinal scale (i.e., living trees = 3, snags
= 2, logs = 1). For more information, see https://doi.org/10.5281/zenodo.7816954.

Insect sampling

We hung two types of traps in each selected tree: one green multi-funnel trap (ChemTica
Internacional, San José, Costa Rica) with 12 fluon-coated funnels, and one cross-vane flight
interception trap (Polytrap™). The first type is used to sample borers like *Agrilus* (Rassati et al.,
2019); the experiment was initially designed to sample the community of borers along the decline
gradient in oak stands (Le Souchu et al., submitted). This type of trap is also efficient for sampling
guilds of foliage-associated insects (Sallé et al., 2020). The second type of trap is commonly used in
forests to sample saproxylic beetles (Bouget et al., 2008).

Both types of traps were suspended among the lower branches in the canopy (i.e., approximately 10
- 15m above the ground). The collectors were filled with a 50% (v/v) monopropylene glycol - water
solution with a drop of detergent. The traps were in place from the end of March to the beginning of
September 2019, during the insects' activite period, and they were emptied every month.

Identification and ecological trait list

All sampled Hymenoptera were sorted and identified to the family or superfamily level, according to
the identification key by Goulet and Huber (1993). Then the specimens were sent to taxonomical
experts who identified them to the genus, species or morphospecies level. Only the Cynipidae and
Figitidae were not identified at a lower taxonomic level.

We arranged the taxa into eight larval trophic guilds: gall-inducers, parasitoids, oak-associated
phyllophagous larvae, other phyllophagous larvae, phytophagous larvae (i.e. feeding on plant parts
other than leaves, for example, seeds), social polyphagous larvae, xylophagous larvae and
pollinivorous/nectarivorous larvae. Preferred larval habitat (larval nesting guild) was also considered,
when known. Taxa were divided into gall-nesters (gall-inducers and their inquiline parasitoids), soil-
nesters, stem-nesters (generally in hollow plant stems) and wood-nesters. For the latter three guilds,
we separated the specialists (spe.) from the generalists (gen.), nesting in wood and/or soil and/or
stems. Specialist wood-nesters were considered analogous to saproxylic taxa (Jonsell et al., 2023).

Data analysis

All analyses and graphs were performed in R, version 4.1.1 (R Core Team, 2021). We used the ggplot2
R-package (Wickham, 2016) to produce the figures. Unless otherwise stated, analyses were realised
with the Hymenoptera community collected at the plot level, and the proportion of declining trees at
both plot and stand scales. The figures with stand decline levels are included in the body of the
present article since they are more informative than the figures with plot decline levels, which are
available in the Supplementary Data.

Some specimens of Cynipidae, Figitidae, Halictidae, Andrenidae, Pteromalidae, Encyrtidae and
Platygastridae were not identified, either because of a lack of expertise or because the specimens
were too damaged for identification. These unidentified specimens were removed from the
community analyses.

To extrapolate the species richness for the whole community (i.e., γ-diversity) and for each decline
category, we first plotted species rarefaction curves (functions *iNEXT* and *ggiNEXT*, iNEXT R-package;
Hsieh et al., 2016) and calculated Chao, Jackknife, Jackknife 2 and Bootstrap diversity indices
(function *specpool*, vegan R-package; Oksanen et al., 2022). We used the additive diversity
partitioning method (Lande, 1996; Crist et al., 2003) to evaluate the contribution of α- and β-diversity
to the γ-diversity of canopy-dwelling Hymenoptera over the entire sampled area (function *adipart*,
index = "richness", 1,000 permutations, vegan R-package). We tested whether observed species
richness differed from the one we expected by chance (null hypothesis). At the plot scale, we used
three levels of differentiation: plot, plot decline category and the whole sampled area. α plot
corresponds to the average species diversity per plot, β plot to the diversity among plots and β
cat_plot to the diversity among plot decline categories. γ-diversity is the sum of α plot, β plot and β
cat_plot (γ = α plot + β plot + β cat_plot). At the stand scale, we used a similar approach except that
four levels of differentiation were included: plot, stand, stand decline category and the whole
sampled area. In addition, we separated β-diversity (Sorensen dissimilarity index) between plots,
stands, decline categories and pairs of decline categories into β-turnover, i.e., species replacement
(Simpson dissimilarity index) and β-nestedness (nestedness-resultant fraction of Sorensen
dissimilarity) (function *beta.multi*, Sorensen family, betapart R-package; Baselga, 2010; Baselga and
Orme, 2012).

We assessed the effect of oak decline on the abundance and species richness of the overall
community, of each taxonomic family (when n > 30 ind.), and on the abundance of each taxa (when n
> 30 ind.). We also assessed the oak-decline effect on each guild (i.e., larval trophic guilds and nesting
guilds). To do this, we used generalised linear mixed-effects models (GLMMs; function *glmer,*
glmer.nb or *lmer*, lme4 R-package; Bates et al., 2022) fitted for the Poisson family, the negative
binomial family or the log-normal distribution (i.e., log-transformed response). We first selected the
best suited family distribution with the fitdistrplus R-package (function *fitdist* and *gofstat*; Delignette-
Muller and Dutang, 2015). We then performed GLMMs with either abundance or species richness at
both scales (plot and stand). To adjust the decline variable to our dataset, we used linear and
equations. We added forest and stand as nested random effects on the intercept in the mixed
models to account for repeated measurements and the spatial configuration of the sampling design.
Since some traps were not continuously functional, for example, because they fell from the tree, we

also added an offset of the log-number of effective traps across the entrapment season, weighted by

the efficacy of each type of trap. To assess model quality, we used marginal R^2 (function *R2*,

performance R-package; Lüdecke et al., 2021). We used the differences in AICc scores (function *AICc*,

AICCmodavg R-package; Mazerolle, 2023) to compare the fit among models. When the linear and

quadratic models were undifferentiated, we show only the results for the simplest one, i.e., the

linear model. We also used the indicspecies R-package (function *multipatt*, func = "IndVal.g", 1,000

permutations) to identify indicator species for healthy, moderately-declining and severely-declining

stands and plots (Dufrene and Legendre, 1997; De Cáceres and Legendre, 2009; De Cáceres et al.,

2010).

In a second step, we implemented Structural Equation Modelling (SEM, piecewiseSEM R-package;

Lefcheck et al., 2022) to evaluate the cascading effects of tree decline (i.e., the proportion of

declining trees) and tree mortality (i.e., the proportion of dead trees) on taxonomic families and

ecological guilds through changes in canopy closure (LAI) and deadwood resources. SEM was used

only for the plots located in the Orléans and Vierzon forests. LAI was originally measured in 12 plots

in the Orléans and Vierzon forests, and correlated strongly with tree decline rate at the stand scale

(Fig. S1). Consequently, in order to avoid excluding additional plots in the SEM analysis, we

extrapolated LAI from the tree decline rate in the six remaining plots using the equation of the

regression curve (Fig. S1). We studied Hymenoptera responses through (i) the abundance of each

taxonomic family, and (ii) the abundance and species richness of each ecological guild (see above).

Response variables that did not respect assumptions of residual normality and homoscedasticity for

linear regressions were log-transformed. In addition, in a first set of mixed models, we added forest

identity (i.e., Orléans or Vierzon) as a random effect in the *lmer* function from lme4 R-package. We

finally deleted this random effect since it was redundant with the fixed effects and caused model

non-convergence. Finally, we used a linear model with the *lm* function. Several plots were installed

close to wetlands, so we added this information to the models as a fixed effect since the

characteristic might affect soil-nesters and phyllophagous species associated with plants in the

herbaceous and shrub layers. We also added an offset term of the log-number of effective traps

across the entrapment season to our models. Finally, since we tested 14 predictors in SEMs to assess

the cascading effects of oak decline on each canopy-dwelling Hymenoptera community metric (i.e.,

multiple tests), we adjusted the p.value to control for an increased risk of type I errors and only

considered effects for which p.values were lower than 0.0036 (i.e., 0.05/14).

RESULTS

The community of canopy-dwelling Hymenoptera

We collected 19,289 individuals, belonging to 54 families and 918 taxa (Fig. 2; Tab. S2). The most

abundant families were the Ichneumonidae (22% of the specimens), the Cynipidae (22%) and the

Perilampidae (14%), while the most diverse were the Ichneumonidae (321 taxa), the Braconidae

(121) and the Tenthredinidae (88) (Fig. 2). The dominant species was the chalcid *Perilampus*

ruficornis (F.) (2,614 individuals), followed by the ant *Dolichoderus quadripunctatus* (L.), the common

wasp *Vespula vulgaris* (L.) (647 ind.), and the halictid *Lasioglossum pallens* (Brullé) (605 ind.). Most of

the community was composed of singletons (344 taxa) and doubletons (129), and only 198 taxa

occurred with more than ten individuals (Tab. S2). Several species cited for the first time in France

were sampled, especially members of the Ichneumonidae family (Tab. S2). We also collected 35 bee

species registered on Worldwide or European Red Lists (Tab. S2). Among them, three species were

near-threatened: *Andrena fulvida* Schenck, *Lasioglossum monstrificum* (Morawitz) and *Lasioglossum*

sexnotatum (Kirby). The other species were classified as either "data deficient" (13 spp.) or "of least

concern" (19 spp.).

The activity of most families peaked in the spring (Fig. S2), and the community was the most

abundant (64% of total specimens) in April. Species richness was more evenly distributed throughout

the sampling period, but also peaked in April (48% of all taxa). Overall, the green multi-funnel traps

clearly outperformed the flight interception traps (93% vs. 7% of total abundance, 94% vs. 28% of all

taxa) (Fig. S3); for each guild and family, both abundance and species richness were higher in the

multi-funnel traps.

According to our species-diversity estimators (Tab. 1), there should have been between 1,074 and

1,505 species in the entire sampled area. We collected between 61% and 85% of these estimations,

and consequently, the rarefaction curve for the whole community barely approached an asymptote

(Fig. 3). Both moderately declining and severely declining plots and stands were predicted to shelter

slightly more taxa than healthy ones (Tab. 1, Fig. 3, Tab. S3, Fig. S4). The communities in healthy and

severely declining stands were clearly distinct, but both were partially similar to those in moderately

declining stands (Fig. 4). At the plot scale, a similar pattern was observed, although the divergence

was less marked (Fig. S5). Consequently, the differences between decline categories were significant

at both the plot (PERMANOVA, $F = 2.49$, df = 2, $R^2 = 0.11$, $p = 0.001$) and stand scales (PERMANOVA, F

$= 3.76$, df = 2, $R^2 = 0.16$, $p = 0.001$).

Contribution of α- and β-diversity to γ-diversity

The additive partitioning of species richness showed that, at the plot scale, *β plot* did not present any

significant difference between observed and expected richness (Fig. S6). At the stand scale, the

observed richness of *β plot* was significantly lower than expected (Fig. 5), while observed *β stand* was

similar to predicted *β stand*. Conversely, decline categories contributed largely to γ-diversity.

Observed *β cat_plot* and *β cat_stand* were significantly higher than expected and respectively

represented 39% and 40% of the γ-diversity.

Overall *β*-diversity was mainly due to species turnover, among all plots and stands (Tab. 2), and

between each pair of plots (Tab. S4) and stand decline categories (Tab. 2). At the plot scale, *β*

nestedness was very low. At the stand scale, however, *β* nestedness was a significant part of the *β*-

diversity between moderately declining stands and healthy ones (22%), as well as between

moderately and severely declining stands (16%).

Indicator species

One species was associated with healthy plots, 11 with moderately declining plots and 11 with

severely declining plots (Tab. S5). At the stand scale, five species were associated with healthy

stands, five with moderately declining stands and seventeen with severely declining stands (Tab. S6).

Four species were indicators at both plot and stand scales. The majority of the indicators were

parasitoids. Several sawflies that feed on non-oak host plants were also indicators of severely

declining stands, and a few species with pollinivorous/nectarivorous larvae were indicators of

moderately declining plots.

Effect of oak decline on ecological guilds and taxonomic families and taxa

Neither the abundance nor the species richness of the community of canopy-dwelling Hymenoptera

was influenced by the extent of the decline. However, contrasting effects were observed among

guilds, taxonomic families and taxa (Tabs. 3 & 4, Fig. S7). For larval trophic guilds, the abundance of

pollinivorous/nectarivorous and polyphagous species increased with decline severity, while those of

xylophagous species decreased (Tab. 3). Parasitoid species were more abundant at intermediate

levels of decline (40% of declining trees, Fig. S7). Carnivorous and polyphagous taxa also became

more diverse as decline severity increased (Tab. 4). For larval habitat guilds, oak decline only slightly

affected the abundance of gall-nesting taxa, while the abundance of both specialist wood-nesters

and specialist soil-nesters was promoted (Tab. 3). Decline also promoted the diversity of generalist

soil-, wood- and stem-nesters (Tab. 4).

For taxonomic groups, as some guilds were composed of a single family, or sometimes a single

species, the effect of oak decline on these families concurred with the effect on the guilds (e.g.,

Xiphydriidae, *Xiphydria longicollis* (Geoffroy) for xylophagous taxa and Cynipidae for gall-inducers)

(Tabs. 3 & 4). Consistent results were also observed for the families making up the majority of a guild,

like the Formicidae (polyphagous), the Halictidae (pollinivorous/nectarivorous) and the Vespidae (carnivorous). In addition, the abundance of Evaniidae (*Brachygaster minuta* (Olivier)), Pamphilidae and Tiphiidae (*Tiphia femorata* F.) was promoted with decline severity. This was also true for Pamphilidae diversity. Conversely, the Perilampidae (*P. ruficornis*) were negatively affected by decline, and the abundance of Ichneumonidae peaked at intermediate levels of decline (approx. 50% of declining trees, Tab. 3). At the species level, the abundance of 18 species out of the 83 tested was influenced by oak decline, either positively (5 species), negatively (4) or non-linearly (9). Eleven of them were parasitoids / hyperparasitoids (Tab. 3, Fig. S7). Two species of ants (*D. quadripunctatus*, *Myrmecina graminicola* (Latreille)) were also promoted by intermediate or high levels of decline.

The SEM showed that tree mortality promoted the abundance and diversity of deadwood habitats and that tree decline markedly reduced LAI (Figs. 6 & 7). Tree mortality rate negatively affected overall species richness, the abundance of the Eulophidae and, to a lesser extent, that of the Tiphiidae (Fig. 6). Tree decline rate also had direct impacts and promoted both the Halictidae and the Tiphiidae, but reduced Xiphydriidae abundance. Few effects on taxonomic guilds were mediated by changes in forest structure (Fig. 6). The most notable effect was the positive effect of tree decline rate on the overall species richness of Hymenoptera mediated by its negative impact on LAI (Fig. 6).

Tree mortality and decline rates also had direct effects on ecological guilds (Fig. 7). An increasing tree mortality rate reduced the abundance and species richness of nectarivorous/pollinivorous species, and the species richness of non-oak phyllophagous, polyphagous and soil-nesting species. As mentioned above, tree decline rate negatively affected the abundance of xylophagous Hymenoptera (Xiphydriidae), but also promoted the species richness of polyphagous species. Nonetheless, the most important impact of tree decline rate was mediated by the reduction in LAI which promoted the species richness of non-oak phyllophagous species, and both the species richness and abundance of nectarivorous/pollinivorous and specialist soil-nesting taxa. LAI impacts on guilds were consistently similar to those of tree mortality, leading to opposite effects of tree decline and tree mortality on ecological guilds (Fig. 7). For the overall community, as well as for nectarivorous/pollinivorous and non-oak phyllophagous taxa, species richness consistently decreased when LAI increased, and was always low when tree mortality rate was high (Fig. S8). Finally, changes in deadwood resources, mediated by tree mortality, only marginally influenced xylophagous taxa and had no impact on the other ecological guilds (Fig. 7).

DISCUSSION

The community of canopy-dwelling Hymenoptera in temperate oak forests

With more than 150,000 species described worldwide, Hymenoptera are one of the most diverse

orders of insects (Huber, 2017; Aberlenc, 2020). In our study, we identified 918 taxa, indicating that

the community of Hymenoptera dwelling in the canopy of the oak stands was particularly diverse. As

a comparison, with the same sampling design, in the same plots and at the same time, we collected

approximately 97,000 Coleoptera and 10,000 Hemiptera, and identified 562 and 169 taxa

respectively (unpubl. data). In agreement with earlier predictions (Forbes et al., 2018), these results

suggest that the community of Hymenoptera could be one of the most diverse, if not the most

diverse, community of arthropods in temperate oak forests.

We collected numerous phyllophagous and gall-inducing species (Tab. S2), which supports using

green multi-funnel traps to collect foliage-feeding arthropods (Sallé et al., 2020). In the same traps,

we also collected a wide diversity of parasitoids and predators, which might have been attracted by

the green colour, indicative of potential hunting grounds for their hosts and prey. For example, the

most common species in our sampling, *P. ruficornis*, is a hyperparasitoid that frequently parasitises

leaf-chewing caterpillars and cynipids (Mitroiu and Koutsoukos, 2023). We also collected several

phytophagous species that feed on oak foliage (e.g., *Periclista* spp.) and oak pollen (e.g., *Andrena*

ferox Smith, *L. pallens*), and saproxylic taxa known to develop in oak branches (i.e., *X. longicollis*) or

to nest in standing deadwood and/or in branches (e.g., *D. quadripunctatus*, *Deuteragenia*

subintermedia (Magretti)). We therefore expected to find all of these taxa in the canopy (e.g.,

Kampichler and Teschner, 2002 for gall-inducers). Conversely, we found several unexpected taxa in

the canopy layer. For instance, we collected species that feed on plants in the herbaceous layer (e.g.,

Strongylogaster multifasciata (Geoffroy) a fern-feeder), as well as parasitoids of soil-dwelling grubs

(i.e., *T. femorata*) or spiders (i.e., *Aporus unicolor* Spinola). We also sampled several uncommon or

even rare species (e.g., *Chrysis equestris* Dahlbom, *Pamphilius balteatus* (Fallén), *Parascleroderma*

sulcatifrons (Kieffer), Marhic & Noblecourt, pers. obs.), including three species of bees that are near-

threatened at the European level (i.e., *A. fulvida*, *L. monstrificum*, and *L. sexnotatum;* Nieto et al.,

2014), and several species new for France (e.g., *Gastrancistrus laticeps* Graham, *Spathius polonicus*

Niezabitowski), or even Europe (i.e., *Plastanoxus evansi* Gorbatovsky). This highlights how little we

still know about the composition and ecology of canopy-dwelling arthropods in general, and

Hymenoptera in particular (Hilszczański, 2018; Sallé et al., 2021), in temperate forests.

Effects of forest decline on the community, guilds and taxa of canopy-dwelling Hymenoptera

Oak decline did not directly affect the abundance and richness of the community of canopy-dwelling
Hymenoptera. However, it did markedly reshape its composition, which differed significantly along
the decline gradient. Species turnover was the main component of these changes, as has been
commonly observed (Soininen et al., 2018), while difference among decline categories was the main
contributor to global diversity. The significant contribution of nestedness to the β-diversity between
moderately declining stands and both healthy and severely declining stands is congruent with the
NMDS results, and indicates that the community characteristic of an intermediate level of decline is
composed of taxa from the other two levels rather than of specific taxa new to the community.
Similarly, in a previous study, the species richness of canopy-dwelling beetles was not influenced by
oak decline, although both their abundance and their biomass increased (Sallé et al., 2020). Our
result is nonetheless surprising since a recent meta-analysis indicates that forest Hymenoptera
diversity (excluding ants) is generally promoted by disturbances like wildfires, windstorms and pest
outbreaks (Viljur et al., 2022). However, the meta-analysis also found that the magnitude of increase
in species richness was quite variable among studies (Viljur et al., 2022). In our case study, the
decline effect was generally consistent between the two spatial scales considered (plot vs. stand),
but the magnitude of the effects was sometimes different. This might reflect differences in foraging
behaviour and habitat use among taxa and/or guilds.

Changes in community composition were driven by contrasting responses of taxa and ecological
guilds to oak decline, supporting the importance of multi-taxonomic and multitrophic approaches in
research on how environmental changes affect community structure (Seibold et al., 2018). As
expected from the literature (Gaston, 1991), the parasitoid guild was both the most diverse and the
most abundant guild. Since parasitoids, and especially koinobionts, require a diversity of habitats to
support the full assemblage of their hosts, they are highly sensitive to changes in habitat (Fraser et
al., 2007; Hilszczański, 2018; Jonsell et al., 2023). Overall, in our study, they exhibited quadratic
relationships with decline levels, and were more abundant at intermediate levels of decline severity
(i.e., 40% of declining oaks). At the taxa level, a majority of species whose abundance was affected by
decline also exhibited a similar pattern, with a peak of abundance at low or moderate levels of
decline (i.e., 30% to 60% of declining oaks). This suggests that a moderate decline may provide a
greater diversity of habitats and trophic resources for parasitoids, as predicted, for instance, by the
intermediate disturbance hypothesis (Grime, 1973; Horn, 1975). This overall trend might be further
modulated by the host type of the parasitoid. We observed that both Tiphiidae (e.g., *T. femorata*)
and Evanidae (e.g., *B. minuta*), which parasitize chafers and cockroaches respectively, benefited from
tree decline, while the parasitoids of defoliating or mining moths (e.g., *Agrypon flaveolatum*
(Gravenhorst), *Bassus* sp. 1, *Earinus gloriatorius* (Panzer), *Macrocentrus nitidus* (Wesmael), *P.*

ruficornis (Mitroiu and Koutsoukos, 2023)), were somewhat disadvantaged by high levels of decline.
Further studies would be necessary to assess whether these effects depend on host type or are
idiosyncratic species responses. It might also be interesting to investigate whether koinobionts are
more responsive to forest decline than idiobionts. The other guild belonging to a higher trophic level,
the carnivorous taxa, was promoted by oak decline. Disturbances can have highly contrasted effects
on predators (Cours et al., 2023). In our study, this guild included several taxonomic families such as
Vespidae, Formicidae and Pemphredonidae that prey on a wide range of taxa. The positive decline
effect might result either from a higher abundance or accessibility of prey (Cours et al., 2023), or
from better habitat conditions. Further studies would be required to identify the factors
underpinning this positive effect.

As expected, the abundance of pollinivorous/nectarivorous taxa increased with severity of decline.
This is in line with previous studies showing that disturbances generally promote pollinators by
providing more floral resources and nesting sites (Wermelinger et al., 2017; Viljur et al., 2022; Perlík
et al., 2023). Interestingly, however, the dominant family of pollinivorous/nectarivorous bees,
Halictidae, was mostly represented by *L. pallens*, a soil-nesting bee which mainly feeds on oak pollen
and consequently, does not rely on other floral resources (Hermann et al., 2003). Since the overall
abundance of specialist soil-nesters was also promoted by stand decline, this suggests that soil
conditions were potentially a more important driver of community change for the guild of
pollinivorous/nectarivorous taxa than were floral resources in the herbaceous layer. The abundance
and the diversity of polyphagous species were also enhanced by oak decline. This is congruent with
the fact that ants, which dominated the guild of polyphagous species, are known to benefit from
silvicultural practices leading to more open conditions (Tausan et al., 2017; Grevé et al., 2018; Cours
et al., 2023).

Oak decline promoted the abundance, but not the diversity, of specialist wood-nesters, i.e. of
saproxylic taxa. Yet, the accumulation and diversification of deadwood resources and weakened
hosts during forest dieback and decline events generally favour saproxylic taxa abundance and
diversity (Beudert et al., 2015; Kozák et al., 2020; Cours et al., 2022, 2023), including those dwelling
in the canopy (Stone et al., 2010; Sallé et al., 2020; Le Souchu et al. *subm.*). This overall trend for the
guild might be contradicted at the species level, since some saproxylic species can be negatively
affected by forest dieback or decline (Vincent et al., 2020). In line with this, the only xylophagous
species in our survey i.e., *X. longicollis*, was negatively affected by decline severity. The larvae of this
species bore into the branches of weakened broadleaved trees and are sometimes associated with
oak decline (Dominik and Starzyk, 1988). Nonetheless, severe decline conditions may have reduced
the amount of living branches in the canopy, thereby reducing the amount of breeding substrates for

the species, as suggested by both the direct effect of tree decline rate, and the indirect effect of tree
mortality on *X. longicollis* abundance.

The diversity of generalist stem-, wood- and soil-nesters was promoted by oak decline, while the
species richness of the specialist guilds remained unaffected. The decline process may lead to
resources pulses, which may be more readily exploited by generalist species (Cours et al., 2023).

Contrary to our hypothesis, oak decline did not influence the phyllophagous species typically
associated with oaks, and only marginally affected gall-inducers and gall-nesters. Responses of leaf-
feeding insects to tree decline can be highly variable, ranging from negative to positive effects
(Martel and Mauffette, 1997; Stone et al., 2010; Sallé et al., 2020). Host specificity might modulate
the outcome of tree decline on leaf-feeding species, but we did not observe any such tendency in our
survey. Interestingly, oak decline had an overall negative effect on the parasitoids of defoliating or
mining caterpillars. This suggests a negative impact of decline on leaf-feeding moths, with potential
cascading impacts on higher trophic levels. Conversely, the diversity of leaf-feeders associated with
other host plants increased with decline severity, and several of them were indicators of high levels
of decline (i.e., *S. multifasciata* and *Aneugmenus padi* (L.), both fern-feeders; *Tenthredopsis nassata*
(L.) and *Dolerus gonager* (F.), bentgrass-feeders; *Eutomostethus luteiventris* (Klug), a rush-feeder; and
Pachyprotasis simulans (Klug), a goldenrod-feeder (Lacourt, 2020)). All of the above were
"Symphyta" and their increase might reflect the preference of these wasps for open forests (Lehnert
et al., 2013), but it might also result from a cascading effect of an increase in plant diversity and
biomass along with increased canopy openness (Lehnert et al., 2013; Lu et al., 2019; Cacciatori et al.,
2022).

Key parameters of decline-driven changes in Hymenoptera guilds and taxa

The results of the SEMs were consistent with those of the GLMMs, indicating that the effects
highlighted by the SEM approach were probably also involved in hymenopteran responses to oak
decline over the whole sampling design. Changes in community composition and structure resulted
either from the direct effects of tree mortality and tree decline, or from indirect effects mediated by
alterations of the forest structure. As dead branches accumulated in the crown of declining trees, an
increase in tree decline rate markedly reduced the LAI. Most of the tree decline effects on ecological
guilds were mediated by changes in LAI, which was therefore a major driver of community change. In
parallel, an increase in tree mortality rate increased the volume and diversity of deadwood
resources.

An increasing tree decline rate directly promoted the abundance of the Halictidae and Tiphiidae.
Both families rely on ground-related trophic resources (floral resources for the Halictidae and soil-

dwelling larvae for the Tiphiidae) and ground habitats since they are soil-nesters (Michez et al.,
2019). In declining stands, soil-nesters may benefit from better or warmer micro-environmental
conditions but also from better underground resources due to LAI reduction (Cours et al., 2023). In
line with this, generalist leaf-feeding weevils, with root-feeding larvae, have been shown to benefit
from oak decline (Sallé et al., 2020). The positive effect of a reduction in LAI on
nectarivorous/pollinivorous taxa is also congruent with its positive effect on the diversity of non-oak
phyllophagous taxa, supporting the above-mentioned hypothesis of a pulse in plant diversity and
floral resources when canopy openness increases (Romey et al., 2007; Lu et al., 2019; Cacciatori et
al., 2022). In our study, canopy opening promoted the overall diversity of the Hymenoptera taxa,
which supports previous observations (Eckerter et al., 2022; Perlík et al., 2023; Rappa et al., 2023).
This result also indicates that micro-environmental modifications at the ground level can have critical
impacts on the communities dwelling in higher forest strata, especially when species have
developmental instars that rely on ground resources or ground micro-habitats.

Surprisingly, while tree mortality rate altered a large array of deadwood resources, we did not
observe any effect of these alterations on the taxonomic families and guilds of Hymenoptera, aside
from the negative effect of a reduction in the number of living trees on the xylophagous
Xiphydriidae. This contradicts several studies where the accumulation and/or diversification of
deadwood resources promoted the abundance, species richness or community composition of
different Hymenoptera guilds and taxa like bees and wasps (Bogusch and Horák, 2018; Urban-Mead
et al., 2021; Rappa et al., 2023), and parasitoid wasps (Hilszczański et al., 2005; Ulyshen et al., 2011a).
However, our results are congruent with recent observations where changes in deadwood amounts
did not influence the species richness or community composition of either cavity- or non-cavity-
nesting Hymenoptera, at least at a local scale (Perlík et al., 2023). Our study plots were located in
managed oak forests, and active forest management likely restricted the magnitude of accumulation
and diversification of deadwood resources (Bouget et al., *subm.*). This may have limited the
potentially beneficial effects of tree mortality and decline on saproxylic Hymenoptera, thus only
weakly promoting saproxylic taxa and wood-nesters. It may also be a matter of spatial scale;
considering the amount and diversity of deadwood resources over larger spatial scales might be
more relevant, especially for taxa with a high dispersal capacity (Cours et al., 2022). In addition,
considering more precise parasitoid subguilds, especially those with saproxylic hosts, might also help
to unravel the impacts of changes in deadwood on the forest Hymenoptera community (but see
below).

In our study, the effects of tree mortality rate on the ecological guilds were consistently opposed to
those of the decline rate, and were, overall, negative. The stands with a high tree mortality rate
might very well have experienced more severe and frequent droughts in the years before the study,
leading to pervasive cascading effects on Hymenoptera through alterations in plant-pollinator or
host-parasitoid interactions, for instance (Rouault et al., 2006; Endres et al., 2021). The negative
effects could also indicate that canopies with dead trees are avoided by Hymenoptera, thus leading
to an overall reduction in abundance and species richness. Stands with high levels of tree mortality
might also have experienced critical changes in micro-environmental conditions and trophic
resources, but further studies are required to better understand the processes underpinning these
deleterious effects.

Forest Hymenoptera and conservation

We have shown that the β-diversity among decline categories at the stand level significantly
contributed to the overall γ-diversity, mostly through species turnover. This suggests that, when
possible, maintaining a landscape mosaic including stands at different levels of decline or dieback
may serve a conservation purpose (Socolar et al., 2016), especially since maintaining high habitat
heterogeneity is generally a key goal for the conservation of insect communities (Samways, 2015).
There are probably underlying mechanisms linked to resource and habitat supplementation and
complementation in the heterogeneous forest landscape (Dunning et al., 1992).

Our results highlight the extreme diversity of the forest Hymenoptera community, and the relevance
of Hymenoptera in studying changes in global community structure and forest functional processes.
Our results also support their value as forest bioindicators, as previously proposed by Maleque *et al.*
(2009). Nonetheless, our study also underlines the relative lack of knowledge of this group in
temperate forests (Hilszczański, 2018; Jonsell et al., 2023). Except for a few well-known species, it
was difficult to obtain information on the biology and ecology of the taxa we collected. We therefore
had to limit the number of ecological traits in our analyses, and our current list of traits is still
partially incomplete. This is unfortunate since databases of ecological traits have proven to be useful
tools for ecological studies and conservation purposes (e.g., Bouget et al., 2013; 2019).
Consequently, considering the growing threats to biodiversity in general, and to insect communities
in particular (Wagner, 2020), we call for thorough investigations on the diversity and ecology of
Hymenoptera to make better use of their potential as bioindicators (Maleque et al., 2009),
understand their ecological services, and better evaluate their conservation value.

CONCLUSION

Oak decline had mixed effects on canopy-dwelling Hymenoptera in our study. We found contrasting
responses for some taxa and ecological guilds, which profoundly reshaped the species assemblage
but did not affect the overall abundance or diversity of the community. Although tree mortality rate
influenced several guilds and taxa, local changes in deadwood resources only had marginal effects on
forest Hymenoptera. However, we confirmed the critical role of canopy openness on the ecological
guilds and taxa dependant on ground-related resources and microhabitats (floral resources,
herbaceous layer and soil conditions). This suggests that micro-environmental changes at the ground
level strongly influence changes in canopy communities. We call for a thorough investigation of the
changes in ecological processes and community structure at the ground level following forest decline
and dieback.

ACKNOWLEDGEMENTS

We thank C. Moliard (INRAE), G. Parmain (INRAE), X. Pineau (LBLGC) and O. Denux (INRAE) for their
technical assistance. We also thank T. Wood, S. Flaminio and P. Rosa for their help in determining the
Andrenidae, Halictidae and Chrysididae. We are grateful to the National Forest Office (Office National
des Forêts) for their field assistance. This work was supported by Région Centre-Val de Loire Project
no. 2018-00124136 (CANOPEE) coordinated by A. Sallé.

SUPPLEMENTARY INFORMATION

Representative specimens of Braconidae, Ichneumonidae, Halictidae, Andrenidae, Crabronidae and Pemphredonidae are housed at the Orleans Museum for Biodiversity and the Environment (MOBE, Orléans, France). R scripts, the list of species traits and the list of plot and stand characteristics can be found at: https://doi.org/10.57745/H8CBG1

REFERENCES

Aberlenc, H.-P. (Ed.), 2020. Les insectes du monde. Museo Éditions, Plaissan.

Baselga, A., 2010. Partitioning the turnover and nestedness components of beta diversity: Partitioning beta diversity. Glob. Ecol. Biogeogr. 19, 134–143. https://doi.org/10.1111/j.1466-8238.2009.00490.x

Baselga, A., Orme, C.D.L., 2012. betapart: an R package for the study of beta diversity: *Betapart package*. Methods Ecol. Evol. 3, 808–812. https://doi.org/10.1111/j.2041-210X.2012.00224.x

Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R.H.B., Singmann, H., Dai, B., Scheipl, F., Grothendieck, G., Green, P., Fox, J., Bauer, A., simulate.formula), P.N.K. (shared copyright on, 2022. lme4: Linear Mixed-Effects Models using "Eigen" and S4.

Beudert, B., Bässler, C., Thorn, S., Noss, R., Schröder, B., Dieffenbach-Fries, H., Foullois, N., Müller, J., 2015. Bark Beetles Increase Biodiversity While Maintaining Drinking Water Quality: Bark beetles, biodiversity and drinking water. Conserv. Lett. 8, 272–281. https://doi.org/10.1111/conl.12153

Bogusch, P., Horák, J., 2018. Saproxylic Bees and Wasps, in: Ulyshen, M.D. (Ed.), Saproxylic Insects, Zoological Monographs. Springer International Publishing, Cham, pp. 217–235. https://doi.org/10.1007/978-3-319-75937-1_7

Bouget, C., Brustel, H., Brin, A., Noblecourt, T., 2008. Sampling saproxylic beetles with window flight traps: methodological insights. Rev. Ecol. Terre Vie 21–32.

Bouget, C., Brustel, H., Noblecourt, T., Zagatti, P., 2019. Les coléoptères saproxyliques de France: catalogue écologique illustré, Patrimoines naturels. Publications scientifiques du Muséum national d'histoire naturelle, Paris.

Bouget, C., Larrieu, L., Nusillard, B., Parmain, G., 2013. In search of the best local habitat drivers for saproxylic beetle diversity in temperate deciduous forests. Biodivers. Conserv. 22, 2111–2130. https://doi.org/10.1007/s10531-013-0531-3

Burkle, L.A., Simanonok, M.P., Durney, J.S., Myers, J.A., Belote, R.T., 2019. Wildfires Influence Abundance, Diversity, and Intraspecific and Interspecific Trait Variation of Native Bees and

647   Flowering Plants Across Burned and Unburned Landscapes. Front. Ecol. Evol. 7, 252.
648   https://doi.org/10.3389/fevo.2019.00252

649 Cacciatori, C., Bacaro, G., Chećko, E., Zaremba, J., Szwagrzyk, J., 2022. Windstorm effects on
650   herbaceous vegetation in temperate forest ecosystems: Changes in plant functional diversity
651   and species trait values along a disturbance severity gradient. For. Ecol. Manag. 505, 119799.
652   https://doi.org/10.1016/j.foreco.2021.119799

653 Cours, J., Bouget, C., Barsoum, N., Horák, J., Le Souchu, E., Leverkus, A.B., Pincebourde, S., Thorn, S.,
654   Sallé, A., 2023. Surviving in Changing Forests: Abiotic Disturbance Legacy Effects on
655   Arthropod   Communities   of   Temperate   Forests.   Curr.   For.   Rep.
656   https://doi.org/10.1007/s40725-023-00187-0

657 Cours, J., Sire, L., Ladet, S., Martin, H., Parmain, G., Larrieu, L., Moliard, C., Lopez-Vaamonde, C.,
658   Bouget, C., 2022. Drought-induced forest dieback increases taxonomic, functional, and
659   phylogenetic diversity of saproxylic beetles at both local and landscape scales. Landsc. Ecol.
660   37, 2025–2043. https://doi.org/10.1007/s10980-022-01453-5

661 Crist, T.O., Veech, J.A., Gering, J.C., Summerville, K.S., 2003. Partitioning Species Diversity across
662   Landscapes and Regions: A Hierarchical Analysis of α, β, and γ Diversity. Am. Nat. 162, 734–
663   743. https://doi.org/10.1086/378901

664 De Cáceres, M., Legendre, P., 2009. Associations between species and groups of sites: indices and
665   statistical inference. Ecology 90, 3566–3574.

666 De Cáceres, M., Legendre, P., Moretti, M., 2010. Improving indicator species analysis by combining
667   groups of sites. Oikos 119, 1674–1684. https://doi.org/10.1111/j.1600-0706.2010.18334.x

668 Delignette-Muller, M.L., Dutang, C., 2015. fitdistrplus: An R Package for Fitting Distributions. J. Stat.
669   Softw. 64. https://doi.org/10.18637/jss.v064.i04

670 Di Giovanni, F., Mei, M., Cerretti, P., 2017. Vertical stratification of selected Hymenoptera in a
671   remnant forest of the Po Plain (Italy, Lombardy) (Hymenoptera: Ampulicidae, Crabronidae,
672   Sphecidae). Fragm. Entomol. 49, 71–77.

673 Dominik, J., Starzyk, J.R., 1988. Owady niszczące drewno, Ochrona drewna. Państwowe
674   Wydawnictwo Rolnicze i Leśne, Poland.

675 Douzon, G., 2006. Le point sur les dépérissements des chênes pédonculés en forêt de Vierzon. Dép.
676   Santé For. Bilan Santé For. En 2005 4.

677 Dufrene, M., Legendre, P., 1997. Species Assemblages and Indicator Species: The Need for a Flexible
678   Asymmetrical Approach. Ecol. Monogr. 67, 345. https://doi.org/10.2307/2963459

679 Dunning, J.B., Danielson, B.J., Pulliam, H.R., 1992. Ecological processes that affect populations in
680   complex landscapes. Oikos 2, 165-177.

Eckerter, T., Braunisch, V., Buse, J., Klein, A.M., 2022. Open forest successional stages and landscape

heterogeneity promote wild bee diversity in temperate forests. Conserv. Sci. Pract. 4.
https://doi.org/10.1111/csp2.12843

Endres, K.L., Morozumi, C.N., Loy, X., Briggs, H.M., CaraDonna, P.J., Iler, A.M., Picklum, D.A., Barr,
W.A., Brosi, B.J., 2021. Plant–pollinator interaction niche broadens in response to severe
drought perturbations. Oecologia 197, 577–588. https://doi.org/10.1007/s00442-021-05036-
0

Forbes, A.A., Bagley, R.K., Beer, M.A., Hippee, A.C., Widmayer, H.A., 2018. Quantifying the
unquantifiable: why Hymenoptera, not Coleoptera, is the most speciose animal order. BMC
Ecol. 18, 21. https://doi.org/10.1186/s12898-018-0176-x

Fraser, S.E.M., Dytham, C., Mayhew, P.J., 2007. Determinants of parasitoid abundance and diversity
in woodland habitats: Determinants of parasitoid abundance and diversity. J. Appl. Ecol. 44,
352–361. https://doi.org/10.1111/j.1365-2664.2006.01266.x

Gaston, K.J., 1991. The Magnitude of Global Insect Species Richness. Conserv. Biol. 5, 283–296.
https://doi.org/10.1111/j.1523-1739.1991.tb00140.x

Goudet, M., Nageleisen, L.M., 2019. Protocole Dépéris: Méthode de notation simplifiée de l'aspect
du houppier des arbres forestiers dans un contexte de dépérissement. For. Entrep. 246, 36–
40.

Goulet, H., Huber, J.T., 1993. Hymenoptera of the world: an identification guide to families,
Agriculture Canada Publication. ed, Publication. Centre for Land and Biological Resources
Research, Ottawa, Ontario, USA.

Grevé, M.E., Hager, J., Weisser, W.W., Schall, P., Gossner, M.M., Feldhaar, H., 2018. Effect of forest
management   on   temperate   ant   communities.   Ecosphere   9,   e02303.
https://doi.org/10.1002/ecs2.2303

Grime, J.P., 1973. Competitive Exclusion in Herbaceous Vegetation. Nature 242, 344–347.
https://doi.org/10.1038/242344a0

Günthardt-Goerg, M.S., Kuster, T.M., Arend, M., Vollenweider, P., 2013. Foliage response of young
central European oaks to air warming, drought and soil type: Oak foliage response to air
warming, drought and soil pH. Plant Biol. 15, 185–197. https://doi.org/10.1111/j.1438-
8677.2012.00665.x

Herrmann, M., Burger, F., Müller, A., & ischendorf, S., 2003. Verbreitung, Lebensraum und Biologie
der Furchenbiene Lasioglossum pallens (Brullé 1832) und ihrer Kuckucksbiene Sphecodes
majalis Pérez 1903 in Deutschland (Hymenoptera, Apidae, Halictinae). Carolinea, 61, 133-
144.

Hilszczański, J., 2018. Ecology, Diversity and Conservation of Saproxylic Hymenopteran Parasitoids, in:
Ulyshen, M.D. (Ed.), Saproxylic Insects, Zoological Monographs. Springer International

Publishing, Cham, pp. 193–216. https://doi.org/10.1007/978-3-319-75937-1_6

Hilszczański, J., Gibb, H., Hjältén, J., Atlegrim, O., Johansson, T., Pettersson, R.B., Ball, J.P., Danell, K.,

2005. Parasitoids (Hymenoptera, Ichneumonoidea) of Saproxylic beetles are affected by

forest successional stage and dead wood characteristics in boreal spruce forest. Biol.

Conserv. 126, 456–464. https://doi.org/10.1016/j.biocon.2005.06.026

Horn, H.S., 1975. Markovian properties of forest succession, in: Cody, M., Diamond, J.M. (Eds.),

Ecology and Evolution of Communities. Belknap Press, Cambridge, MA, pp. 196–211.

Hsieh, T.C., Ma, K.H., Chao, A., 2016. iNEXT: an R package for rarefaction and extrapolation of species

diversity (H ill numbers). Methods Ecol. Evol. 7, 1451–1456. https://doi.org/10.1111/2041-

210X.12613

Hu, B., Simon, J., Rennenberg, H., 2013. Drought and air warming affect the species-specific levels of

stress-related foliar metabolites of three oak species on acidic and calcareous soil. Tree

Physiol. 33, 489–504. https://doi.org/10.1093/treephys/tpt025

Huber, J.T., 2017. Biodiversity of Hymenoptera, in: Foottit, R.G., Adler, P.H. (Eds.), Insect Biodiversity.

John   Wiley   &   Sons,   Ltd,   Chichester,   UK,   pp.   419–461.

https://doi.org/10.1002/9781118945568.ch12

Ishii, H., Tanabe, S., Hiura, T., 2004. Exploring the Relationships Among Canopy Structure, Stand

Productivity, and Biodiversity of Temperate Forest Ecosystems. For. Sci. 50, 342–355.

Jentsch, A., White, P., 2019. A theory of pulse dynamics and disturbance in ecology. Ecology 100.

https://doi.org/10.1002/ecy.2734

Jonsell, M., Vårdal, H., Forshage, M., Stigenberg, J., 2023. Saproxylic Hymenoptera in dead wood

retained on clear cuts, relation to wood parameters and their degree of specialisation. J.

Insect Conserv. 27, 347–359. https://doi.org/10.1007/s10841-023-00468-w

Kampichler, C., Teschner, M., 2002. The spatial distribution of leaf galls of Mikiola fagi (Diptera:

Cecidomyiidae) and Neuroterus quercusbaccarum (Hymenoptera: Cynipidae) in the canopy

of   a   Central   European   mixed   forest.   Eur.   J.   Entomol.   99,   79–84.

https://doi.org/10.14411/eje.2002.014

Kennedy, C.E.J., Southwood, T.R.E., 1984. The Number of Species of Insects Associated with British

Trees: A Re-Analysis. J. Anim. Ecol. 53, 455. https://doi.org/10.2307/4528

Kozák, D., Svitok, M., Wiezik, M., Mikoláš, M., Thorn, S., Buechling, A., Hofmeister, J., Matula, R.,

Trotsiuk, V., Bače, R., Begovič, K., Čada, V., Dušátko, M., Frankovič, M., Horák, J., Janda, P.,

Kameniar, O., Nagel, T.A., Pettit, J.L., Pettit, J.M., Synek, M., Wieziková, A., Svoboda, M.,

2020. Historical Disturbances Determine Current Taxonomic, Functional and Phylogenetic

Diversity of Saproxylic Beetle Communities in Temperate Primary Forests. Ecosystems.

https://doi.org/10.1007/s10021-020-00502-x

Lacourt, J., 2020. Symphytes d'Europe, Hyménoptères d'Europe. NAP éditions, Verrières-le-Buisson.

Lande, R., 1996. Statistics and Partitioning of Species Diversity, and Similarity among Multiple
Communities. Oikos 76, 5. https://doi.org/10.2307/3545743

Larrieu, L., Courbaud, B., Drénou, C., Goulard, M., Bütler, R., Kozák, D., Kraus, D., Krumm, F., Lachat,
T., Müller, J., Paillet, Y., Schuck, A., Stillhard, J., Svoboda, M., Vandekerkhove, K., 2022.
Perspectives: Key factors determining the presence of Tree-related Microhabitats: A
synthesis of potential factors at site, stand and tree scales, with perspectives for further
research. For. Ecol. Manag. 515, 120235. https://doi.org/10.1016/j.foreco.2022.120235

Lehnert, L.W., Bässler, C., Brandl, R., Burton, P.J., Müller, J., 2013. Conservation value of forests
attacked by bark beetles: Highest number of indicator species is found in early successional
stages. J. Nat. Conserv. 21, 97–104. https://doi.org/10.1016/j.jnc.2012.11.003

Lewis, C.N., Whitfield, J.B., 1999. Braconid Wasp (Hymenoptera: Braconidae) Diversity in Forest Plots
Under   Different   Silvicultural   Methods.   Environ.   Entomol.   28,   986–997.
https://doi.org/10.1093/ee/28.6.986

Lu, D., Zhang, G., Zhu, J., Wang, G.G., Zhu, C., Yan, Q., Zhang, J., 2019. Early natural regeneration
patterns of woody species within gaps in a temperate secondary forest. Eur. J. For. Res. 138,
991–1003. https://doi.org/10.1007/s10342-019-01219-w

Lüdecke, D., Ben-Shachar, M., Patil, I., Waggoner, P., Makowski, D., 2021. performance: An R Package
for Assessment, Comparison and Testing of Statistical Models. J. Open Source Softw. 6, 3139.
https://doi.org/10.21105/joss.03139

Lyytikäinen-Saarenmaa, P., Tomppo, E., 2002. Impact of sawfly defoliation on growth of Scots pine
Pinus sylvestris (Pinaceae) and associated economic losses. Bull. Entomol. Res. 92, 137–140.
https://doi.org/10.1079/BER2002154

Maleque, M.A., Maeto, K., Ishii, H.T., 2009. Arthropods as bioindicators of sustainable forest
management, with a focus on plantation forests. Appl. Entomol. Zool. 44, 1–11.
https://doi.org/10.1303/aez.2009.1

Martel, J., Mauffette, Y., 1997. Lepidopteran Communities in Temperate Deciduous Forests Affected
by Forest Decline. Oikos 78, 48. https://doi.org/10.2307/3545799

Mazerolle, M.J., 2023. AICcmodavg: Model Selection and Multimodel Inference Based on (Q)AIC(c).

Michez, D., Rasmont, P., Terzo, M., Vereecken, N., 2019. Hymenoptera of Europe. NAP Editions,
Verrières-le-Buisson.

Mitroiu, M. D., Koutsoukos, E., 2023. *Perilampus neglectus* and other neglected species: new records
of Palaearctic Perilampidae (Hymenoptera, Chalcidoidea), with a key to European species of
Perilampus. Journal of Hymenoptera Research, 96, 57-99.

Motten, A.F., 1986. Pollination Ecology of the Spring Wildflower Community of a Temperate

Deciduous Forest. Ecol. Monogr. 56, 21–42. https://doi.org/10.2307/2937269

Nageleisen, L.M., 2008. Actualité sur les dépérissements du "chêne." Min. Agr. Pêche.

Nieto, A., Roberts, S.P.M., Kemp, J., Rasmont, P., Kuhlmann, M., García Criado, M., Biesmeijer, J.C.,
Bogusch, P., Dathe, H.H., De la Rúa, P., 2014. European red list of bees. Publications Office,
Luxembourg.

Oksanen, J., Simpson, G., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P., hara, R., Solymos, P.,
STEVENS, H., Szöcs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B., Borcard, D.,
Carvalho, G., Chirico, M., De Cáceres, M., Durand, S., Weedon, J., 2022. vegan community
ecology package version 2.6-2 April 2022.

Perlík, M., Kraus, D., Bußler, H., Neudam, L., Pietsch, S., Mergner, U., Seidel, D., Sebek, P., Thorn, S.,
2023. Canopy openness as the main driver of aculeate Hymenoptera and saproxylic beetle
diversity following natural disturbances and salvage logging. For. Ecol. Manag. 540, 121033.
https://doi.org/10.1016/j.foreco.2023.121033

Pucci, T., 2008. A Comparison of the Parasitic Wasps (Hymenoptera) at Elevated Versus Ground
Yellow Pan Traps in a Beech-Maple Forest. J. Hymenopt. Res. 17, 116–123.

R Core Team, 2021. R: A Language and Environment for Statistical Computing. R Foundation for
Statistical Computing, Vienna, Austria.

Rappa, N.J., Staab, M., Ruppert, L.-S., Frey, J., Bauhus, J., Klein, A.-M., 2023. Structural elements
enhanced by retention forestry promote forest and non-forest specialist bees and wasps.
For. Ecol. Manag. 529, 120709. https://doi.org/10.1016/j.foreco.2022.120709

Rassati, D., Marini, L., Marchioro, M., Rapuzzi, P., Magnani, G., Poloni, R., Di Giovanni, F., Mayo, P.,
Sweeney, J., 2019. Developing trapping protocols for wood-boring beetles associated with
broadleaf trees. J. Pest Sci. 92, 267–279. https://doi.org/10.1007/s10340-018-0984-y

Rico-Gray, V., Oliveira, P.S., 2008. The ecology and evolution of ant-plant interactions. University of
Chicago Press.

Romey, W.L., Ascher, J.S., Powell, D.A., Yanek, M., 2007. Impacts of Logging on Midsummer Diversity
of Native Bees (Apoidea) in a Northern Hardwood Forest. J. Kans. Entomol. Soc. 80, 327–338.
https://doi.org/10.2317/0022-8567(2007)80[327:IOLOMD]2.0.CO;2

Rouault, G., Candau, J.-N., Lieutier, F., Nageleisen, L.-M., Martin, J.-C., Warzée, N., 2006. Effects of
drought and heat on forest insect populations in relation to the 2003 drought in Western
Europe. Ann. For. Sci. 63, 613–624. https://doi.org/10.1051/forest:2006044

Saintonge, F.-X., Goudet, M., 2020. Situation sanitaire début 2020 de 52 massifs de chênes en forêt
publique. Dép. Santé For. 5.

Sallé, A., Cours, J., Le Souchu, E., Lopez-Vaamonde, C., Pincebourde, S., Bouget, C., 2021. Climate
Change Alters Temperate Forest Canopies and Indirectly Reshapes Arthropod Communities.

Front. For. Glob. Change 4, 8. https://doi.org/10.3389/ffgc.2021.710854

Sallé, A., Parmain, G., Nusillard, B., Pineau, X., Brousse, R., Fontaine-Guenel, T., Ledet, R., Vincent-
Barbaroux, C., Bouget, C., 2020. Forest decline differentially affects trophic guilds of canopy-
dwelling beetles. Ann. For. Sci. 77, 21. https://doi.org/10.1007/s13595-020-00990-w

Samaniego, L., Thober, S., Kumar, R., Wanders, N., Rakovec, O., Pan, M., Zink, M., Sheffield, J., Wood,
E.F., Marx, A., 2018. Anthropogenic warming exacerbates European soil moisture droughts.
Nat. Clim. Change 8, 421–426. https://doi.org/10.1038/s41558-018-0138-5

Samways, M.J., 2015. Future-proofing insect diversity. Curr. Opin. Insect Sci. 12, 71–78.
https://doi.org/10.1016/j.cois.2015.09.008

Seibold, S., Cadotte, M.W., MacIvor, J.S., Thorn, S., Müller, J., 2018. The Necessity of Multitrophic
Approaches in Community Ecology. Trends Ecol. Evol. 33, 754–764.
https://doi.org/10.1016/j.tree.2018.07.001

Seidl, R., Thom, D., Kautz, M., Martin-Benito, D., Peltoniemi, M., Vacchiano, G., Wild, J., Ascoli, D.,
Petr, M., Honkaniemi, J., Lexer, M.J., Trotsiuk, V., Mairota, P., Svoboda, M., Fabrika, M.,
Nagel, T.A., Reyer, C.P.O., 2017. Forest disturbances under climate change. Nat. Clim. Change
7, 395–402. https://doi.org/10.1038/nclimate3303

Senf, C., Pflugmacher, D., Zhiqiang, Y., Sebald, J., Knorn, J., Neumann, M., Hostert, P., Seidl, R., 2018.
Canopy mortality has doubled in Europe's temperate forests over the last three decades.
Nat. Commun. 9, 4978. https://doi.org/10.1038/s41467-018-07539-6

Slippers, B., Hurley, B.P., Wingfield, M.J., 2015. *Sirex* Woodwasp: A Model for Evolving Management
Paradigms of Invasive Forest Pests. Annu. Rev. Entomol. 60, 601–619.
https://doi.org/10.1146/annurev-ento-010814-021118

Smith, S.M., Islam, N., Bellocq, M.I., 2012. Effects of single-tree selection harvesting on
hymenopteran and saproxylic insect assemblages in the canopy and understory of northern
temperate forests. J. For. Res. 23, 275–284. https://doi.org/10.1007/s11676-012-0253-5

Sobek, S., Tscharntke, T., Scherber, C., Schiele, S., Steffan-Dewenter, I., 2009. Canopy vs. understory:
Does tree diversity affect bee and wasp communities and their natural enemies across forest
strata? For. Ecol. Manag. 258, 609–615. https://doi.org/10.1016/j.foreco.2009.04.026

Socolar, J.B., Gilroy, J.J., Kunin, W.E., Edwards, D.P., 2016. How Should Beta-Diversity Inform
Biodiversity Conservation? Trends Ecol. Evol. 31, 67–80.
https://doi.org/10.1016/j.tree.2015.11.005

Soininen, J., Heino, J., Wang, J., 2018. A meta-analysis of nestedness and turnover components of
beta diversity across organisms and ecosystems. Glob. Ecol. Biogeogr. 27, 96–109.
https://doi.org/10.1111/geb.12660

Spinoni, J., Vogt, J.V., Naumann, G., Barbosa, P., Dosio, A., 2018. Will drought events become more

frequent   and   severe   in   Europe?   Int.   J.   Climatol.   38,   1718–1736.
https://doi.org/10.1002/joc.5291

Stone, A.C., Gehring, C.A., Whitham, T.G., 2010. Drought negatively affects communities on a
foundation   tree: growth   rings   predict   diversity.   Oecologia   164,   751–761.
https://doi.org/10.1007/s00442-010-1684-3

Stone, G.N., Schönrogge, K., Atkinson, R.J., Bellido, D., Pujade-Villar, J., 2002. The population biology
of oak gall wasps (Hymenoptera: Cynipidae). Annu. Rev. Entomol. 47, 633–668.

Tausan, I., Dauber, J., Trica, M.R., Marko, B., 2017. Succession in ant communities (Hymenoptera:
Formicidae) in deciduous forest clear-cuts - an Eastern European case study. Eur. J. Entomol.
114, 92–100. https://doi.org/10.14411/eje.2017.013

Ulyshen, M.D., 2016. Wood decomposition as influenced by invertebrates: Invertebrates and wood
decomposition. Biol. Rev. 91, 70–85. https://doi.org/10.1111/brv.12158

Ulyshen, M.D., 2011. Arthropod vertical stratification in temperate deciduous forests: Implications
for   conservation-oriented   management.   For.   Ecol.   Manag.   261,   1479–1489.
https://doi.org/10.1016/j.foreco.2011.01.033

Ulyshen, M.D., Pucci, T.M., Hanula, J.L., 2011a. The importance of forest type, tree species and wood
posture to saproxylic wasp (Hymenoptera) communities in the southeastern United States. J.
Insect Conserv. 15, 539–546. https://doi.org/10.1007/s10841-010-9348-5

Ulyshen, M.D., Soon, V., Hanula, J.L., 2011b. Vertical Distribution and Seasonality of Predatory Wasps
(Hymenoptera: Vespidae) in a Temperate Deciduous Forest. Fla. Entomol. 94, 1068–1070.
https://doi.org/10.1653/024.094.0450

Urban-Mead, K.R., Muñiz, P., Gillung, J., Espinoza, A., Fordyce, R., Dyke, M. van, McArt, S.H.,
Danforth, B.N., 2021. Bees in the trees: Diverse spring fauna in temperate forest edge
canopies. For. Ecol. Manag. 482, 118903. https://doi.org/10.1016/j.foreco.2020.118903

Vance, C.C., Smith, S.M., Malcolm, J.R., Huber, J., Bellocq, M.I., 2007. Differences Between Forest
Type and Vertical Strata in the Diversity and Composition of Hymenpteran Families and
Mymarid Genera in Northeastern Temperate Forests. Environ. Entomol. 36, 1073–1083.
https://doi.org/10.1603/0046-225X(2007)36[1073:DBFTAV]2.0.CO;2

Viljur, M., Abella, S.R., Adámek, M., Alencar, J.B.R., Barber, N.A., Beudert, B., Burkle, L.A., Cagnolo, L.,
Campos, B.R., Chao, A., Chergui, B., Choi, C., Cleary, D.F.R., Davis, T.S., Dechnik-Vázquez, Y.A.,
Downing, W.M., Fuentes-Ramirez, A., Gandhi, K.J.K., Gehring, C., Georgiev, K.B., Gimbutas,
M., Gongalsky, K.B., Gorbunova, A.Y., Greenberg, C.H., Hylander, K., Jules, E.S., Korobushkin,
D.I., Köster, K., Kurth, V., Lanham, J.D., Lazarina, M., Leverkus, A.B., Lindenmayer, D., Marra,
D.M., Martín-Pinto, P., Meave, J.A., Moretti, M., Nam, H., Obrist, M.K., Petanidou, T., Pons,
P., Potts, S.G., Rapoport, I.B., Rhoades, P.R., Richter, C., Saifutdinov, R.A., Sanders, N.J.,

Santos, X., Steel, Z., Tavella, J., Wendenburg, C., Wermelinger, B., Zaitsev, A.S., Thorn, S.,
2022. The effect of natural disturbances on forest biodiversity: an ecological synthesis. Biol.
Rev. 97, 1930–1947. https://doi.org/10.1111/brv.12876

Vincent, A., Tillier, P., Vincent-Barbaroux, C., Bouget, C., Salle, A., 2020. Influence of forest decline on
the abundance and diversity of Raphidioptera and Mecoptera species dwelling in oak
canopies. Eur. J. Entomol. 117, 372–379. https://doi.org/10.14411/eje.2020.041

Wagner, D.L., 2020. Insect Declines in the Anthropocene. Annu. Rev. Entomol. 65, 457–480.
https://doi.org/10.1146/annurev-ento-011019-025151

Wermelinger, B., Moretti, M., Duelli, P., Lachat, T., Pezzatti, G.B., Obrist, M.K., 2017. Impact of
windthrow and salvage-logging on taxonomic and functional diversity of forest arthropods.
For. Ecol. Manag. 391, 9–18. https://doi.org/10.1016/j.foreco.2017.01.033

Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

Yumoto, T., 1987. Pollination systems in a warm temperate evergreen broad-leaved forest on Yaku
Island. Ecol. Res. 2, 133–145. https://doi.org/10.1007/BF02346922

Zemlerová, V., Kozák, D., Mikoláš, M., Svitok, M., Bače, R., Smyčková, M., Buechling, A., Martin, M.,
Larrieu, L., Paillet, Y., Roibu, C.-C., Petritan, I.C., Čada, V., Ferenčík, M., Frankovič, M., Gloor,
R., Hofmeister, J., Janda, P., Kameniar, O., Majdanová, L., Markuljaková, K., Matula, R.,
Mejstřík, M., Rydval, M., Vostarek, O., Svoboda, M., 2023. Natural Disturbances are Essential
Determinants of Tree-Related Microhabitat Availability in Temperate Forests. Ecosystems.
https://doi.org/10.1007/s10021-023-00830-8

Tables

Table 1. Species diversity estimators for the overall community and for each category of stand
decline in three temperate oak forests (21 stands and 42 plots).

Stand status	Observed species richness	Number of plots	Chao	Jackknife 1	Jackknife 2	Bootstrap	Range
All stands	918	42	1,407 ± 70	1,279 ± 65	1,505	1,074 ± 32	1,074 - 1,505
Healthy	501	14	943 ± 79	742 ± 76	909	604 ± 37	604 – 943
Moderately declining	629	16	943 ± 51	891 ± 76	1,041	745 ± 39	745 - 1,041
Severely declining	532	12	898 ± 61	783 ± 78	940	641 ± 35	641 - 940

919 Table 2. Overall β-diversity, β turnover and β nestedness between pairs of stand-decline category.
920 Overall β-diversity corresponds to Sorensen dissimilarity, β turnover to Simpson dissimilarity and β
921 nestedness to the difference between Sorensen and Simpson dissimilarity. H = healthy stands, MD =
922 moderately declining stands, SD = severely declining stands.

	Overall β-diversity	β turnover	β nestedness
Among plots	0.95	0.93 (97.89%)	0.02 (2.11%)
Among stands	0.89	0.85 (95.51%)	0.04 (4.49%)
Among categories of stand decline	0.47	0.41 (87.23%)	0.06 (12.77%)
Between H and MD stands	0.37	0.29 (78.38%)	0.08 (21.62%)
Between MD and SD stands	0.38	0.32 (84.21%)	0.06 (15.79%)
Between H and SD stands	0.42	0.40 (95.24%)	0.02 (4.76%)

926 Table 3. Effect of the proportion of declining oak trees (plot or stand decline) on the abundance of oak-dwelling Hymenoptera, and on the abundance of
927 guilds (larval trophic guilds, larval nesting guild, saproxylic guild), families (abundance > 30 ind.) and taxa (abundance > 30 ind.). GLMMs were fitted for the
928 negative binomial family (NB), the Poisson family (P) or the log-normal family (Log), with forest and stand as random effects. Models with either linear or
929 quadratic decline variables were tested. ΔAICc = AICc (decline model) − AICc (null model). The best model between plot and stand scales is highlighted in
930 bold. Only significant relationships are shown.

Community / taxa	Abund.	Dist. fam.	Plot decline					Stand decline				
			Delta AICc	Estimate	S. E.	z or t value	Marginal R^2	Delta AICc	Estimate	S. E.	z or t value	Marginal R^2
Overall community	19,282	Log	2.72	-	-	-	-	2.09	-	-	-	-
Larval trophic guilds												
Gall-inducer	4,199	Log	-4.29	d1. 1.74 ns	1.7	1.01	0.20	-5.15	d1. 2.44 ns	2.0	1.22	0.24
				d2. -3.06 ns	1.9	-1.65			d2. -4.06 .	2.2	-1.85	
Parasitoid	9,904	Log	1.75	-	-	-	-	-6.35	d1. 1.49 *	0.7	2.23	0.12
									d2. -2.25 **	0.7	-3.12	
Pollinivorous/Nectarivorous	1,246	Log	-3.05	d1. 3.19 *	1.5	2.13	0.07	-4.30	1.56 *	0.6	2.63	0.15
				d2. - 2.74 .	1.6	-1.72						
Polyphagous	1,281	NB	-9.35	1.29 ***	0.1	113.57	0.19	-3.99	1.22 ***	0.1	112.82	0.14
Xylophagous	149	NB	-3.51	-3.47 *	1.5	-2.33	0.35	2.57	-	-	-	-
Larval nesting guilds												
Gall nest	5,090	Log	0.72	-	-	-	-	-5.24	d1. 3.05 ns	2.0	1.55	0.25
									d2.-4.55 *	2.2	-2.12	
Spe. Soil nesters	1,384	NB	0.16					-4.60	0.75 ***	0.01	101.35	0.04
Spe. Wood nesters	1,866	Log	-3.07	0.74 *	0.3	2.59	0.10	-2.67	0.85 *	0.3	2.49	0.11
Families												
Cynipidae	4,199	Log	-4.29	d1. 1.74 ns	1.7	1.01	0.20	-5.15	d1. 2.44 ns	2.0	1.22	0.24
				d2. -3.06 ns	1.9	-1.65			d2. -4.06 .	2.2	-1.85	
Evanidae	33	P	-3.13	2.87 **	1.1	2.74	0.15	-11.88	5.03 ***	1.3	3.97	0.35
Formicidae	1,339	NB	-8.98	1.23 ***	0.1	113.00	0.18	-3.35	1.14 **	0.1	109.721	0.13

931   Tab. 3 continued

Community / taxa	Abund.	Dist. fam.	Plot decline					Stand decline				
			Delta AICc	Estimate	S. E.	z or t value	Marginal R²	Delta AICc	Estimate	S. E.	z or t value	Marginal R²
Halictidae	924	Log	-3.94	d1. 3.81 *	0.6	2.17	0.06	-4.90	1.82 *	0.7	2.70	0.12
				d2. -3.25 .	0.6	-1.75						
Ichneumonidae	4,242	Log	3.89	-	-	-	-	-4.46	d1. 2.24 **	0.7	3.40	0.11
				-	-	-	-		d2. -2.37 **	0.7	-3.33	
Pamphiliidae	38	NB	-1.55	-	-	-	-	-4.77	3 *	1.1	2.66	0.21
Perilampidae	2,614	NB	-6.30	-1.72 **	0.5	-3.24	0.38	-9.87	-2.67 *	0.8	-3.45	0.42
Tiphiidae	92	P	-2.44	2.39 *	1.1	2.21	0.05	-2.17	4.24 *	1.9	2.29	0.12
Xiphydriidae	149	NB	-3.51	-3.47 *	1.5	-2.33	0.35	2.57	-	-	-	-
Taxa												
Agrypon flaveolatum	32	P	-2.63	-2.18 *	1.1	-2.00	0.11	2.32	-	-	-	-
Bassus sp.1	38	NB	-4.54	d1. 9.06 *	3.5	2.58	0.36	0.07	-	-	-	-
				d2. -12.59 **	4.7	-2.67						
Brachygaster minuta	33	P	-3.13	2.87 *	1.1	2.74	0.15	-11.88	5.03 ***	1.3	3.97	0.35
Ctenopelmatinae sp.1	88	P	-86.47	d1. 41.56 ***	8.3	4.99	0.53	1.88	-	-	-	-
				d2. -51.48 ***	11.3	-4.55						
Ctenopelmatinae sp.7	73	P	-20.70	2.92 ***	0.7	4.45	0.18	2.41	-	-	-	-
Deuteragenia vechti	44	P	-6.11	-3.05 *	1.2	-2.62	0.14	0.11	-	-	-	-
Dolichoderus quadripunctatus	718	NB	-6.30	1.49 *	0.5	3.06	0.18	-3.30	1.34 *	0.6	2.38	0.12
Earinus elator	180	NB	2.53	-	-	-	-	2.56	-	-	-	-
Earinus gloriatorius	43	NB	-4.19	d1. 7.77 ***	1.7	4.50	0.21	2.57	-	-	-	-
				d2. -9.29 ***	2.0	-4.73						
Empria tridens	30	P	-2.72	-2.18 *	0.9	-2.33	0.11	1.07	-	-	-	-
Macrocentrus nitidus	42	NB	-3.08	d1. 5.12 ***	1.2	4.15	0.24	-0.01	-	-	-	-
				d2. -7.78 ***	1.6	-4.98						

934   Tab. 3 continued

Community / taxa	Abund.	Dist. fam.	Plot decline					Stand decline				
			Delta AICc	Estimate	S. E.	z or t value	Marginal R²	Delta AICc	Estimate	S. E.	z or t value	Marginal R²
Myrmecina graminicola	39	P	-10.86	d1. 12.48 **	3.8	3.29	0.30	-3.03	d1. 9.87 **	3.7	2.64	0.21
				d2. -12.13 **	4.0	-3.04			d2. -9.68 *	4.1	-2.39	
Perilampus ruficornis	2,614	NB	-8.50	d1. 2.00 ***	0.1	131.40	0.38	-22.79	d1. 3.53 ***	0.1	124.97	0.67
				d2. -4.36 ***	0.1	-287.19			d2. -7.43 ***	0.1	-262.53	
Priocnemis perturbator	38	P	-4.30	d1. 9.73 *	3.9	2.47	0.17	1.33	-	-	-	-
				d2. -8.37 *	3.9	-2.16						
Strongylogaster multifasciata	52	P	-12.98	d1. 15.97 **	4.7	3.40	0.31	-1.43	-	-	-	-
				d2. -11.87 **	3.5	-3.39						
Tiphia femorata	92	P	-2.44	2.39 *	1.1	2.21	0.04	-2.17	4.24 *	1.9	2.29	0.12
Vespa crabro	240	NB	-3.70	1.61 *	0.7	2.43	0.10	-0.27	-	-	-	-
Xiphydria longicollis	149	NB	-3.51	-3.47 *	1.5	-2.33	0.35	2.57	-	-	-	-
Xorides praecatorius	30	P	1.37	-	-	-	-	-3.07	d1. 9.88 *	3.9	2.53	0.22
									d2. -9.82 *	4.0	-2.46	

937 Table 4. Effect of the proportion of declining oak trees (plot or stand decline) on the species richness of oak-dwelling Hymenoptera, and on the species

938 richness of guilds (larval trophic guilds, larval nesting guild, saproxylic guild) and families (abundance > 30 ind.). Only significant relationships are shown. See

939 the caption for Table 3 for more information.

Community / taxa	Species richness	Dist. fam.	Plot decline					Stand decline				
			Delta AICc	Estimate	S. E.	z or t value	Marginal R²	Delta AICc	Estimate	S. E.	z or t value	Marginal R²
Overall community	918	Log	0.75	-	-	-	-	-0.97	-	-	-	-
Larval trophic guilds												
Carnivorous	70	P	-3.60	0.57 *	0.23	2.54	0.05	-3.23	0.63 *	0.25	2.56	0.05
Non-oak phyllophagous	86	Log	1.62	-	-	-	-	-3.16	0.89 *	0.64	2.6	0.11
Polyphagous	25	P	-2.66	0.56 *	0.25	2.79	0.04	-3.24	0.66 *	0.28	2.40	0.05
Larval nesting guilds												
Gen. Soil nesters	28	P	-2.13	0.53*	0.24	2.19	0.03	-0.68				
Gen. Stem nesters	32	P	-2.92	0.58 *	0.24	2.39	0.04	-3.65	0.67 *	0.27	2.53	0.04
Gen. Wood nesters	40	P	-3,89	0.55 *	0.22	2.57	0.04	-2,72	0.55 *	0.24	2.32	0.03
Families												
Formicidae	29	P	-2.08	0.49 *	0.23	2.15	0.03	-1.54	-	-	-	-
Pamphiliidae	9	P	-4.51	2.00 **	0.69	2.89	0.12	-7.45	2.58 **	0.79	3.29	0.17

Figures

Figure 1

Figure 1. Map of France showing the location of the three oak forests studied, and the 21 stands

sampled. Details on the stands can be found in Tab. S1.

Figure 2

Figure 2. Overview of the abundances (log-transformed) of the Hymenoptera families sampled in oak

canopies. Families are arranged in three taxonomic groups: "Symphyta", Apocrita Aculeata and

Apocrita parasitica. Numbers between brackets indicate the number of individuals and the number

of taxa identified for each family.

Figure 3

Figure 3. Rarefaction curves for canopy-dwelling Hymenoptera from the three forests, 21 stands and

42 plots. H = healthy stands, MD = moderately declining stands, SD = severely declining stands,

Overall = all stands. See main text for details on stand decline categories.

Figure 4

Figure 4. NMDS ordination (k=3, stress=0.18) of species composition per plot, grouped by stand

decline category. H = healthy stands, MD = moderately declining stands, SD = severely declining

stands. Species with fewer than 10 individuals were removed from the analysis. See main text for

details on stand decline categories.

Figure 5

Figure 5. Global additive partitioning of species richness of the canopy-dwelling Hymenoptera in the

three oak forests, at plot and stand scales. Four levels are represented: α plot (within plot), β plot

(among plots), β stand (among stands) and β cat_stand (among levels of stand decline). Significance

levels correspond to the difference between expected and observed values, with ***: $p < 0.001$; . : p

< 0.1.

Figure 6

Figure 6. Results of the Structural Equation Modelling (SEM) for the effects of oak decline on the

abundance of the main Hymenoptera taxonomic families through changes in forest structure. Since

we tested 14 predictors, we used 0.0036 as the p.value (0.05/14). The transparent lines had a p.value

below 0.05 but above 0.0036 and help to explain the response variables in the multiple regression

models. LAI: leaf area index, vol.: volume, DW: deadwood, LW: living wood, CWM: community-

weighted mean trait value, FDis: functional diversity, Med: medium-sized, LarVer: large and very

large, dec12: low level of decay, dec34: high level of decay.

Figure 7

Figure 7. Results of the Structural Equation Modelling (SEM) for the effects of oak decline on

Hymenoptera ecological guilds through changes in forest structure. Since we tested 14 predictors, we

used 0.0036 as the p.value (0.05/14). See the caption for Figure 6 for complementary information.

997