Enhancer plasticity in endometrial tumorigenesis demarcates non-coding driver mutations and alterations in 3D genome organization to stimulate oncogene expression

Sebastian Gregoricchio[1,#], Aleksandar Kojic[1,#], Marlous Hoogstraat[1,2], Karianne Schuurman[1], Suzan Stelloo[3], Tesa M. Severson[1,2], Tracy A. O'Mara[4], Marjolein Droog[1], Abhishek A. Singh[1], Dylan M. Glubb[4], Lodewyk F.A. Wessels[2], Michiel Vermeulen[3,5], Flora van Leeuwen[6], Wilbert Zwart[1]

[1] Division of Oncogenomics, Oncode Institute, The Netherlands Cancer Institute, Amsterdam, The Netherlands
[2] Division of Molecular Carcinogenesis, The Netherlands Cancer Institute, Amsterdam, The Netherlands
[3] Department of Molecular Biology, Faculty of Science, Radboud Institute for Molecular Life Sciences, Oncode Institute, Radboud University Nijmegen, Nijmegen, The Netherlands
[4] Cancer Research Program, QIMR Berghofer Medical Research Institute, Brisbane, Australia
[5] Division of Molecular Genetics, Oncode Institute, The Netherlands Cancer Institute, Amsterdam, The Netherlands
[6] Department of Epidemiology, The Netherlands Cancer Institute, Amsterdam, The Netherlands

[#] Authors equally contributed to this work

ABSTRACT

The incidence and mortality of Endometrial Cancer (EC) is on the rise. 85% of ECs depend on Estrogen Receptor alpha (ERα) for proliferation, but little is known about its transcriptional regulation in these tumors.

We generated epigenomics and Hi-C data streams in healthy and tumor endometrial tissues, identifying robust ERα reprogramming and profound alterations in 3D genome organization that lead to a gain of tumor-specific enhancer activity during EC development. Integration with WGS data from metastatic samples revealed a striking enrichment of non-coding somatic mutations at tumor-enriched ERα sites. Through machine learning-based predictions and interaction proteomics analyses, we identified an enhancer mutation which alters 3D genome organization, impairing recruitment of the transcriptional repressor EHMT2/G9a/KMT1C, thereby alleviating transcriptional repression of *ESR1* in EC.

In summary, we identified a complex genomic-epigenomic interplay in EC development and progression, altering 3D genome organization to enhance expression of the critical driver ERα.

INTRODUCTION

Endometrial cancer (EC) is the second most-common gynecological cancer with over 417,000 new cases and 97,000 deaths in 2020, with global incidence rates increasing every year[1]. Surprisingly, and in stark contrast to most other cancers, survival rates for endometrial cancer are decreasing[1,2]. This gradual deterioration of EC survival is likely related to the relatively understudied nature of the disease, with many molecular mechanisms driving tumor development and progression remaining largely elusive.

Endometrial tissue is under tight endocrine control, and this remains the case upon tumorigenesis. The majority of endometrial cancers (85%) are classified as low grade endometrioid tumors, expressing Estrogen Receptor alpha (ERα) and Progesterone Receptor (PR)[3–5]. PR agonists are used in the treatment of endometrial tumors as alternative to chemotherapy and hysterectomy to retain ovarian function in young patients. Furthermore, the estrogen competitive ERα antagonist tamoxifen is also prescribed to patients with endometrial cancer, often alternated with PR agonists (reviewed in ref [6]). Thus, while PR acts in a tumor suppressive manner, ERα serves as driver of tumor progression which is therapeutically blocked in endometrial cancer care.

ERα is an enhancer-acting transcription factor, regulating expression of its responsive genes through long-range 3D genome interactions[7,8]. To date, most literature on promoter-enhancer communication and ERα activity has been focused on breast cancer[8], and is far less understood in endometrial tumors. The observation that only around 30% of ERα sites are shared between breast cancer and endometrial cancer[9–11] highlights highly context dependent chromatin binding of ERα. Tamoxifen – prescribed in the treatment of breast cancer – is a risk factor for endometrial tumorigenesis[12]. We showed that tamoxifen treatment in endometrial tissue reprograms the ERα chromatin interaction landscape, phenocopying profiles found in breast cancer cells, driving endometrial tumor growth[13]. These observations highlighted that the endometrial cancer epigenome is not fixed, but rather dynamically affected by oncogenic factors.

Both ERα in breast tissue[14], and Androgen Receptor (AR) in prostate tissue[15–17], undergo extensive reprogramming throughout the genome during tumorigenesis. In case of prostate cancer, both somatic mutations and risk single-nucleotide polymorphisms (SNPs) are enriched at tumor-gained AR sites, with only a small fraction of which causally affecting transcriptional output[16]. In endometrial cancer, such studies thus far not reported functional interplay between somatic mutations, epigenetic alterations and the impact on 3D genome organization remain fully elusive.

Here, we investigated the plasticity of ERα genomic action upon endometrial tumor development, by comparing the epigenome of primary human healthy and tumor

endometrial tissues (ERα and H3K27ac ChIP-seq) and the effects on 3D genome

organization (Hi-C, 4C-seq and, H3K27ac HiChIP). Secondly, we studied the crosstalk

between epigenetic alterations and somatic variant events (WGS) that leads to EC

progression. We identified substantial reprogramming of ERα on the genome upon

tumorigenesis, which coincided with the occurrence of somatic variants in metastatic

samples. In particular, we discovered an EC-specific *ESR1* enhancer which is target of

somatic variants in metastatic EC, loosing capacity to recruit lysine methyl-transferase

G9a/EHMT2/KMT1C when mutated, to enhance ERα expression in metastatic disease.

Cumulatively, we show that non-coding mutations in endometrial cancer can have direct pro-

tumorigenic potential, through a tight interplay with epigenetic alterations and changes in the

3D genome structure.

RESULTS

ERα enhancer plasticity in endometrial tumorigenesis

To investigate the role of ERα signaling in endometrial tumor development, we first sought to assess how the ERα cistrome differs between endometrial tumors and healthy endometrial tissue. We collected five fresh-frozen endometrial tumors from post-menopausal patients that did not receive any previous systemic therapy, as well as four fresh-frozen samples from women with pathologically normal endometrial tissue (**Fig. 1a** and **Extended Data Fig. 1a**, for clinicopathological characteristics see **Extended Data Table 1**). All samples were subjected to ERα and H3K27 acetylation (H3K27ac) ChIP-seq in order to identify active regulatory elements that are under ERα control. On average, we identified 20,815 ERα peaks (range 1,399-51,888) and 44,824 H3K27ac peaks (range 8,791-71,898) in our samples. The ChIP-seq library size was comparable among the different samples for both targets (**Extended Data Fig. 1b**). However, the number of ERα peaks was higher in tumor samples compared to healthy tissue (Mann-Whitney P = 0.063, **Fig. 1b**) while the global number of H3K27ac peaks for these samples was comparable between both groups (Mann-Whitney P = 0.29, **Fig. 1c**). To asses data quality and reproducibility, we performed Principal Component Analysis (PCA) on the peak intensities. These analyses revealed a clustering of the samples by ChIP target (first component) and tissue type (second component) (**Extended Data Fig. 1c**).

Differential binding analysis (*DiffBind*[18]) for ERα between tumor and normal samples revealed over 10,292 differentially-bound genomic locations (FDR ≤ 0.05 & $|\log_2(\text{FoldChange})| \geq 1$) (**Fig. 1d-e** and **Extended Data Fig. 1d**), of which 6,488 ERα sites were lost and 3,804 ERα sites gained in tumor samples, hereafter referred to as "tumor-depleted" (blue) and "tumor-enriched" (orange) ERα binding sites, respectively. Tornado plots depicting normalized ERα ChIP-seq signal for all 10,292 differential-enriched ERα sites confirmed the selective enrichment of signals between samples, with high reproducibility (**Fig. 1f**). As ERα acts as enhancer-selective transcription factor, we next analyzed the differential ERα sites for presence of H3K27ac, inferring regulatory element activity. Interestingly, persistent activity in both tissue states was observed at tumor-depleted ERα sites, while tumor-enriched ERα sites showed a marked increase of activity upon tumorigenesis (**Fig. 1g**).

Collectively, these data suggest a selective gain of regulatory element potential upon tumorigenesis, marked by dynamically altered ERα sites.

Genomic distributions and transcription complex composition at differential ERα sites

To further investigate the nature of the ERα regions differentially bound upon tumorigenesis, we first inspected their genome wide distribution with respect to different genomic elements. In agreement with previous reports from us (tumors[9,13]) and others (cell lines[10]), tumor-enriched ERα sites were mostly found at distal intergenic and intronic regions (**Fig. 1h**). However, tumor-depleted ERα sites, selectively occupied by ERα in healthy endometrial tissue, display a strong enrichment for promoter regions of genes involved in cell cycle regulation and DNA damage processes (**Extended Data Fig. 1e**). Motifs for ERα/β (ESR1/2) and AP-1 (JUN, FOS) binding sites were specifically enriched at tumor-depleted ERα regions, whereas tumor-enriched regions demonstrated an overabundance of SOX and FOX transcription factor family member motifs (**Fig 1i**).

Differential motif enrichment suggests selective enrichment of ERα-interaction partners for different regions. To test this hypothesis, we integrated publicly available ChIP-seq data from the Ishikawa endometrial cancer cell line (**Fig. 1j** and **Extended Data Table 1**), revealing striking differences between transcription factors on chromatin selectivity for the two ERα binding site categories. Interestingly, ERα chromatin binding in hormone-deprived cells showed a significant overlap with the tumor-depleted ERα peaks (adjusted Fisher's exact $P = 3.15 \times 10^{-21}$), while ERα peaks from estradiol-stimulated cells showed significant overlap with the tumor-enriched ERα peaks (adjusted Fisher's exact $P = 4.60 \times 10^{-59}$, **Fig. 1j**). In addition, both FOXM1 and FOXA1 binding sites in Ishikawa showed significant overlaps with our tumor enriched regions (adjusted Fisher's exact $P = 3.37 \times 10^{-21}$ and $P = 6.25 \times 10^{-57}$, respectively, **Fig. 1k**).

Altogether, these data suggest prominent enhancer plasticity in endometrial tumorigenesis, with ERα relocating to alternatively engaged and activated non-canonical enhancers in endometrial cancer.

3D high-order chromatin structures are altered in tumors

Enhancer regions can modulate gene transcription through interactions with promoters in 3D genomic space[19]. With the observed plasticity of ERα chromatin profiles in tumorigenesis, we hypothesized that these epigenetic alterations were accompanied by reorganization of the 3D chromatin structure. To test this hypothesis, we performed high-throughput chromosome conformation capture (Hi-C) analyses on healthy endometrial tissue (n=3) and primary tumors (n=3) derived from post-menopausal women that did not receive any previous systemic therapy (**Fig. 2a,** for clinicopathological features see **Extended Data Table 1**). Hi-C is particularly suitable to study translocations in cancer[20], which were not

observed in any of the tumor samples we analyzed (**Extended Data Fig. 2a**). We did
however observe a distinct clustering of the healthy tissues from the tumor specimens,
based on 3D genome organization (**Extended Data Fig. 2b**). In addition, tumor samples
displayed a higher Hi-C contact heterogeneity in comparison to the more correlated healthy
tissues (**Extended Data Fig. 2b**). Next, we quantified the Relative Hi-C Contact Probability
(RCP) as function of the contact distances. Interestingly, tumors displayed an increased
probability of short-range interactions (<2Mb) concomitant with a loss of long-range
chromatin contacts (**Fig. 2b-d** and **Extended Data Fig. 2c-d**). Furthermore, analyzing the
distribution of the loop distances (**Fig. 2e**), we found that loops whose anchors include
tumor-enriched ERα binding sites tend to be shorter in tumor tissues than in healthy tissues.
On the other hand, tumor-depleted and shared ERα binding sites do not show differences in
loop length between two tissues stages. We next wondered whether this loop shortening in
tumor samples was associated with changes in chromatin compartmentalization. Analysis of
compartment polarization (**Fig. 2b,f-g** and **Extended Data Fig. 2e-f**) revealed decreased
compartment strength in tumor samples compared to healthy tissues. In particular, we
observed that A compartments (euchromatin) were more robust in healthy samples, while B
compartments (heterochromatin) where slightly stronger in tumor tissues (**Fig. 2g** and
Extended Data Fig. 2f). Overall, around one third of compartments (A: 565/1431 (39.5%);
B: 440/1306 (33.6%)) underwent class switching in tumor samples (**Fig. 2h**). To investigate
whether this compartment class switch was occurring specifically in compartments where
ERα was bound in a tissue type-specific manner, we overlapped the compartments with the
differential ERα occupied regions upon tumorigenesis (**Fig. 2i**). We found that compartments
switching from A to B class were enriched for tumor-depleted ERα binding sites, relative to
tumor-enriched ones (A-to-B depleted/enriched ratio 2.65 (135/51)). In contrast,
compartments switching from B-to-A were clearly enriched for tumor-gained ERα binding
sites (B-to-A depleted/enriched ratio 0.33 (35/105)). These findings were further confirmed
studying the behavior of smaller genomic regions (compartment bins) relative to
differentially-occupied ERα binding sites, excluding regions that harbor multiple ERα binding
site categories (Fisher's exact test: B-to-A depleted-vs-enriched: $P = 1.317 \times 10^{-56}$, OR 0.20;
A-to-B depleted-vs-enriched: $P = 6.013 \times 10^{-11}$, OR 1.87) (**Fig. 2j** and **Extended Data Fig.**
2g).
Taken together, our findings highlight a widespread reorganization of the 3D genome
upon tumorigenesis that leads to a loss of chromatin compartment polarization. This was
accompanied by an extensive class-switching, with tumor-enriched ERα binding sites being
enriched in B-to-A compartments. Moreover, we identified a gain of short-range chromatin
interactions upon tumor development, at the expense of long-range contacts.
189

Somatic mutations are enriched at tumor-gained ERα sites

Somatic mutations in cancer are often found enriched at non-coding regulatory

elements[21]. To determine whether that is also the case in endometrial cancer, we analyzed

Whole Genome Sequencing (WGS) data of 26 fresh frozen biopsy samples, derived from 24

cases of advanced endometrial cancer. Biopsies were collected from tumors that

metastasized to lung, liver, and other abdominal regions (**Fig. 3a** and **Extended Data Table**

3). In these tumors, the vast majority (>96%) of variants were located at non-coding regions,

with 85% of the identified variants found to populate intronic and intergenic regions.

Considerably, less variants (mean 9%, range 8.1-10.3) were located proximally upstream of

a coding gene, potentially representing promoter regions (**Fig. 3b**). To assess WGS data

quality and explore additional genomic features of our samples, we investigated mutational

signatures, counts, and frequently-mutated genes (**Fig. 3c-d**). Four samples were

hypermutated and displayed 20-67% of variants associated with defective DNA mismatch

repair (SBS15). Six samples from tumors pre-treated with carboplatin showed traces of

signatures 31 and 35, both associated with platinum treatment[22]. Cumulatively, these results

confirm previously-described enrichment of tumor-intrinsic[23,24] and treatment-induced[22]

mutational features of endometrial cancer. The most-frequently mutated genes with protein

sequence altering mutations included *ARID1A* and various members of the PIK3CA

pathway, such as *PTEN*, *PIK3CA* and *PIK3R1*, recapitulating previous observations for

endometrial tumor[23].

As somatic mutations in metastatic samples were enriched in non-coding regions, we

integrated these somatic mutation data with tissues-type enriched ERα ChIP-seq data (**Fig.**

3e). Interestingly, differentially bound ERα regions were significantly enriched for somatic

mutations (11.9%, 1,221/10,292), relative to non-differential ERα-bound sites (8%,

5,881/73,312; Fisher's exact P = 5.64×10^{-36}, OR 1.54, 95% CI 1.44-1.65). All samples

contributed to this enrichment, with a median somatic variant count of 18 (range 1-358) in

differentially ERα-bound sites. Moreover, comparing tumor-enriched with tumor-depleted

regions, we observed a significant enrichment of somatic variants in the tumor-enriched ERα

sites compared to tumor-depleted ones (14.6% vs 10.1%, Fisher's exact P = 2.98×10^{-11}, OR

1.51, 95% CI 1.34-1.71, **Fig. 3e**). In most instances (n=1,002), regions overlapped with only

one somatic variant, while some ERα-bound regions were observed to harbor 2 (183

regions), 3 (34 regions) or 4 (2 regions) somatic mutations.

To investigate genes whose expression may be affected by enhancer mutations, we

intersected tumor-depleted and tumor-enriched ERα binding sites (FDR ≤ 0.01) bearing

mutations with H3K27ac HiChIP chromatin looping data in Ishikawa cells (**Extended Data**

Table 4), aimed to couple putative enhancrs with the transcription start sites (TSS) they

control, yielding a total of 331 genes. Out of these, 13 genes were differentially expressed in

Ishikawa endometrial cancer cells upon 8h of estradiol-stimulation[25] (**Fig. 3f**). Interestingly,

ESR1 (encoding ERα) was represented as well, being downregulated at both RNA and

protein level (**Fig. 3f-g** and **Extended Data Fig. 3a**) and harboring 3 metastatic somatic

mutations in two *cis* regulatory elements upstream of the *ESR1* locus, hereafter referred as

'*Enhancer 1*' and '*Enhancer 2*' (**Fig. 4a** and **Extended Data Fig. 3b**). Of note, high *ESR1*

expression is associated with favorable outcome in primary endometrial cancer patients

(TCGA[26]) (**Fig. 3h**). Of note, these two enhancers are specific to the endometrium cancer

samples, relative to healthy endometrial tissues and breast cancer specimens, based on

ATAC-seq (TCGA[27]) and ERα ChIP-seq signal (our previous work[28]) (**Extended Data Fig.**

3b-c) at the *ESR1* locus.

ESR1 enhancer mutation alters 3D genome contacts and decreases

EHMT2/G9a enhancer binding to boost ERα expression

Since *ESR1* expression is strongly associated with favorable outcome in endometrial

cancer (**Fig. 3h**), alterations in its transcriptional regulation may have direct implications on

tumor development and progression. Since both *Enhancer 1* and *Enhancer 2* – targets of

somatic mutations – showed induction of ERα binding and H3K27ac signals in tumors (**Fig.**

4a), we investigated whether these loci may undergo alternative regulation in different

stages of tumor progression leading to *ESR1* expression deregulation.

Hi-C analyses of endometrial tumors (**Fig. 4b**) illustrated increased short-range and

decreased long-range chromatin interactions at this locus, when compared to healthy

endometrial tissue. These data were confirmed with 4C-seq analyses using the *ESR1*

promoter as view-point (**Fig. 4c**), showing tumor-specific gained interactions of newly ERα-

engaged enhancers with the *ESR1* promoter (**Extended Data Fig. 2h**).

To identify which *ESR1* enhancer mutation had an higher probability to impact *ESR1*

gene regulation and potentially drive tumor progression, we employed the *Akita*[29] machine

learning tool to predict the effect of the somatic mutations on the 3D genome organization

(**Fig. 4d**). Our analyses predicted the mutation occurring at position chr6:152,002,679 (TTC-

to-T) of *Enhancer 1* to affect chromatin architecture surrounding the *ESR1* locus. Based on

these results, this specific mutation was selected for further analyses.

To determine whether enhancer mutations can alter the composition of the

transcription complex recruitment, we performed DNA affinity purification experiments

coupled to quantitative mass spectrometry. To this end, we incubated nuclear extracts from

Ishikawa endometrial cancer cells with immobilized DNA-oligonucleotides encompassing the

center of the ERα-bound site at the *Enhancer 1* – containing either the reference or mutant

sequence – with a ±25bp window around it (**Fig. 5a-b**). Quantitative proteomics identified 13

gained and 11 lost proteins upon mutation of the regulatory element, respectively (**Fig. 5b**).

To identify differentially bound proteins that may directly regulate *ESR1* expression, we used

publicly available RNA-seq (TCGA[26]) data of 589 endometrial cancer tissues to correlate the

expression of *ESR1* with the expression of genes corresponding to the 24 differentially-

bound proteins (**Fig. 5c**). We then compared these values with the distribution of the

correlation of *ESR1* expression against the full transcriptome, after which the genes most-

strongly correlating positively (*POLK*, *ZBTB21*, *XPA*, *SIN3A*) or negatively (*GTF2IRD1*,

EHMT2, *ZNF768*) with *ESR1* expression, were selected (**Extended Data Fig. 4a**). For these

genes, expression was analyzed in relation to the progression-free survival probability of

endometrial cancer patients (TCGA[26]) (**Fig. 5d** and **Extended Data Fig. 4b**). Interestingly,

our analyses identified only one of the 7 genes analyzed – the lysine methyl-transferase

EHMT2 (also known as G9a or KMT1C) – to significantly associate with a poor outcome of

endometrial cancer patients (**Fig. 5d** and **Extended Data Fig. 4b**).

EHMT2/G9a interacts with ERα (**Fig. 5e** and **Extended Data Fig. 4c**), based on

immunoprecipitation experiments. In line with this, EHMT2/G9a was recruited to the *ESR1*

Enhancer 1 and promoter upon estradiol treatment, as assessed by ChIP-qPCR in Ishikawa

cells (**Fig. 5f**). Furthermore, siRNA-mediated knock-down of EHMT2/G9a in Ishikawa cell

lines, resulted in an increase of ERα protein expression (**Fig. 5g** and **Extended Data Fig.**

4d), confirming the reciprocal relationship between these proteins.

Altogether, these data show that EHMT2/G9a DNA binding is impaired upon

mutation of the *ESR1 Enhancer 1* locus, thereby enhancing expression of tumor driver ERα

in metastatic endometrial cancer and that its activity is required to repress *ESR1* expression

in endometrial cancer in a ERα-dependent manner.

DISCUSSION

Despite the central role of ERα in the endometrial cancer development and progression[5], the molecular actions of ERα in this cancer type remain largely unknown.

Though multi-omics analyses (**Fig. 1a**), we investigated how ERα activity is altered in endometrial cancer and how this feature may be leveraged by tumor cells to progress into the metastatic stage. Cumulatively, our findings allowed us to propose a molecular model describing the dynamic nature of ERα action in endometrial cancer development and progression (**Fig. 5h**): i) in healthy endometrial tissue, ERα regulates genes involved in cell cycle and DNA damage regulation; ii) during tumorigenesis, ERα re-localizes to the tumor-specific *ESR1* enhancer where recruits and interacts with EHMT2/G9a which acts as a co-repressor to reduce *ESR1* expression; iii) during tumor progression, the *ESR1* enhancer acquires a somatic mutation that may impair EHMT2/G9a chromatin occupancy to allow for enhanced *ESR1* expression.

Notably, our analysis of the ERα cistrome in both normal and neoplastic endometrial tissues revealed limited overlap of chromatin occupancy between the two tissue states. Interestingly, tumor-depleted ERα binding sites did not display differences in H3K27ac, in contrast to the tumor-enriched ones which instead showed a consistent increase in this active histone mark. These results highlighted a specific gain-of-function for newly-engaged ERα-bound *cis*-regulatory elements in tumors, while the regulatory elements for which ERα binding is lost remain active through action of other transcription factors. Surprisingly, studying the genomic distribution of the ERα binding sites, we observed an unexpected enrichment of promoter regions for the tumor-depleted binding sites over the tumor-enriched ones. Indeed, ERα has been reported to primarily occupy promoter-distal *cis*-regulatory elements rather than promoters[9,13,30,31]. However, the molecular biology of this nuclear receptor has been studied mainly using cancer models, introducing therefore a potential cancer-related bias. Indeed, our results highlight an extended re-programming of the ERα regulation in tumors involving a promoter-to-enhancer shift in ERα chromatin occupancy.

Our data, combined with previous findings showing that cell-specific ERα sites lack high-affinity estrogen responsive elements (EREs)[10], suggest that tumor-depleted binding sites (enriched for ERα/β motifs) are essential for the cell maintenance in normal tissues. However, during tumorigenesis these sites progressively lose ERα binding in favor of sites enriched in tumors (lacking EREs), likely driving cell transformation. Interestingly, at these tumor-enriched binding sites we found an overrepresentation of SOX and FOX transcription factor family motifs. These finding are particularly relevant for the endometrial cancer biology since one of the most recurrently mutated genes in this cancer type is *SOX17*[23], which acts as tumor suppressor[32]. Therefore, we can hypothesize that *SOX17* loss-of-function may

leave low-affinity ERα binding sites more accessible for ERα, increasing receptor occupancy
at these regions. On the other hand, enrichment of motifs for forkhead pioneer transcription
factor class (FOX family) at tumor-enriched ERα binding sites is consistent with the
increased H3K27ac signal at these low-affinity ERα binding sites. Indeed, FOXA1 is known
to facilitate ERα binding through its pioneer activity in breast cancer[33], and we previously
proposed FOXA1 to serve a comparable role in endometrial cancer[9]. In support of these
hypothesis, tumor-enriched ERα binding sites display an enrichment in FOXA1 ChIP-seq
peaks detected in Ishikawa endometrial cancer cells (**Fig. 1j-k**).

Enhancers interact with promoter regions to regulate gene expression, facilitated
through spatial rearrangement of the chromatin structure[19]. Performing Hi-C analyses in
healthy and tumor tissues, we observed a global enrichment of short-range (<2Mb)
chromatin interactions, at the expense of long-range (>2Mb) contacts, and compartment
depolarization in tumor samples. Chromatin compartment destabilization is observed in
many tumor types[34] (e.g., colorectal cancer[35] and glioblastoma[36]). Interestingly, as tumor-
enriched ERα binding sites were enriched at B-to-A compartment switched sub-regions,
these data suggest that alterations in 3D genome organization directly contribute to
enhancer plasticity in tumorigenesis, concordant with increased H3K27ac signal at these
sites.

As somatic mutations preferentially occur at non-coding regions of the genome[21], we
investigated whether tumor-enriched ERα sites could drive tumor progression through
acquisition of mutations. Through integration of somatic mutation data with RNA-seq in
Ishikawa cells, we concluded that most genes with proximal ERα-bound enhancer mutations
are not under direct ERα control. This is in agreement with our previous observation in
prostate cancer, in which only a minor fraction of SNVs at regulatory elements functionally
affected their activity[16].

While the interaction between ERα and EHMT2/G9a has already been reported in
breast[37,38], EHMT2/G9a is considered to be an ERα co-activator in that cellular context.
However, studies in eryhtroleukemia[39,40] describe a dual role of EHMT2/G9a as both gene
activator and repressor depending on its interaction with Mediator rather than JARID1,
respectively. As we observed ERα expression levels being upregulated following knockdown
of EHMT2/G9a (**Fig. 5g**), an ERα co-repressive role for EHMT2/G9a in the endometrial
cancer cell context is supported by our data, but further investigation would be required to
define what determines the repressive transcriptional effect of EHMT2/G9a in the interplay
with ERα in this tumor type.

In this study we shed light on the molecular mechanisms underlying endometrial
cancer development and progression, connecting alterations in 3D genome organization
with epigenetic plasticity and non-coding somatic mutations. This study may serve as

blueprint for studies in other hormone-driven cancer types, in which such complex
connections are yet to be identified.

METHODS

Human tissue collection and quality assessment

Fresh-frozen healthy and tumor samples were obtained from post-operative tissue at the Netherlands Cancer Institute (Amsterdam, the Netherlands) from patients who did not receive neoadjuvant endocrine treatment. Tumor content was assessed by haematoxylin and eosin (H&E) staining on slides taken throughout the tissue sample. For tumors, only samples displaying a tumor percentage greater or equal to 70% were deemed eligible for further analysis.

The study was approved by the institutional review board of the Netherlands Cancer Institute, written informed consent was signed by all participants enrolled in the study, and all research was carried out in accordance with relevant guidelines and regulations.

Cell culture and chemicals

Ishikawa endometrial cancer cell lines (Merck Sigma Aldrich) have been cultured in Dulbecco's Modified Eagle Medium (DMEM, Gibco) supplemented with 10% Fetal Bovine Serum (FBS-12A, Capricorn Scientific) and penicillin/streptomycin (100µg/mL, Gibco). Cell lines were subjected to regular *Mycoplasma* testing, and underwent authentication by short tandem repeat profiling (Eurofins Genomics).

For hormone stimulation, cells were pre-cultured for 3 days in phenol red-free DMEM (Gibco) supplied with 5% dextran-coated charcoal (DCC) stripped FBS, 2mM L-glutamine (Gibco), and penicillin/streptomycin (100µg/mL, Gibco), then stimulated with 10nM DMSO-solubilized 17β-estradiol (MedChemExpress, #HY-B0141) for the indicated amount of time. Inhibition of ERα activity was performed by treatment of the cells with 100nM DMSO-solubilized Fulvestrant (MedChemExpress, #HY-13636).

ChIP-qPCR and ChIP-seq library preparation

Chromatin Immunoprecipitation of ERα (SantaCruz #sc-543, 5µg/IP) and H3K27ac (ActiveMotif #39133, 5µg/IP) were performed as previously described[28] employing the combination of DSG and Formaldehyde to perform the crosslinking. The same protocol, but using only Formaldehyde for the crosslinking, have been used to perform EHMT2/G9a (Abcam #ab133482, 5µg/IP) and Normal-IgG (Merck Millipore #12-370, same amount than IP) ChIPs in Ishikawa cells.

Immunoprecipitated DNA is compared to input sample by the comparative C_t method, and signal enrichment is expressed as percentage of input (% Input). Real-Time quantitative PCR (RT-qPCR) was performed on the QuantStudio™ 5 Real-Time PCR System 384-well (Applied Biosystems, #A28140) using SenSMix™ SYBR® No-ROX buffer (Meridian

397 Bioscience). The following genomic locations have been tested: *ESR1_Enhancer1*-125kb

398 (Fw: 5'-TGGTAGGTGCTCAGGAGATAA-3', Rv: 5'-CAGCGACTCGAACAGGATTT-3'),

399 *ESR1_*promoter-0.9kb (Fw: 5'-CCACTCCTGGCATTGTGATTA-3', Rv: 5'-

400 CAGGACACATGACACCCAAT-3'), *CDK12_*TSS (Fw: 5'-GGACCTGATCTCGCGTTGTT-3',

401 Rv: 5'-TAGCCTCTCGCGATGTTTCG-3').

ChIP-seq data processing, motif and Gene Ontology enrichment analyses

Reads were aligned to the human genome build GRCh37 using *BWA* 0.5.9-r26-dev[41].

Reads with a mapping quality (MAPQ) < 20 were removed from further analysis. Enrichment

over input control was determined using both *MACS2*[42] and *DFilter*[43]. Only peaks identified

by both methods, and not overlapping with the ENCODE GRCh37/Hg19 blacklisted

regions[44], were retained.

Differentially bound regions between tumor and healthy samples were identified using

DiffBind[18] with *EdgeR*[45] mode at an FDR < 0.05. RPGC-normalized (Reads Per Genomic

Content; *bamCoverage*[46]) ChIP-seq signal at peaks was visualized using *deepTools*[46], while

genomic tracks and average density profiles were generated with *Rseb*[47] R-package

(https://github.com/sebastian-gregoricchio/Rseb) or *pyGenomeTracks*[48,49].

Peak genomic annotation was performed using *Homer*[50], while for motif enrichment in peak

regions we used *AME* v5.5.05[51] algorithm from the *MEME* suite[52] using the JASPAR

database[53,54]. Visualization of the enrichment motifs was made using *ggwordcloud* R-

package using the *AME* computed E-value.

Area proportional venn diagrams were created using the *Vennerable* R package.

Analyses for Gene Ontology biological process (GO-BP) enrichments for tumor-depleted

ERα promoter-bound genes were performed using DAVID[55] (v6.8) and employing tumor-

enriched ERα promoter-bound genes as background data set.

Hi-C library preparation and data processing

Flash-frozen primary tissues have been processed as described in **Fig. 2a**. Briefly, 30µm

slices are cross-linked, washed, electronically homogenized and filtered using a 75µm cell

strainer and collected in a 1.5mL microcentrifuge loBind tube. Then, Hi-C single-index library

preparation was performed as previously described[56] using MboI (New England Biolabs)

restriction enzyme.

Quality and quantification of the Hi-C libraries was assessed using the 2100 Bioanalyzer

(Agilent, DNA 7500 kit). An equimolar pool of the different samples was sequenced on the

Illumina NextSeq 550 System in a 75bp paired-end setup. De-multiplexed fastq data were

analyzed at 2kb, 10kb, 40kb, 100kb and 500kb resolution using the *snHiC*[57] v0.1.1 pipeline

(https://github.com/sebastian-gregoricchio/snHiC) using default parameters and the

Hg19/GRCh37 genome assembly. Downstream analyses have been performed using
GENOVA[58] v1.0.1 (Hi-C contact matrices and Relative Contact Probabilities),
HiCExplorer[59,60] v3.7.2 (compartment polarization, loops distance, Hi-C correlations, TAD
detection, long/short contacts ratio), *intervene*[61] (upset plot and compartment-ChIP peaks
overlaps), *dcHiC*[62] (compartments detection and differential compartment analyses),
pyGenomeTracks[48,49] (data visualization).
440

Whole Genome Sequencing (WGS) analyses
Sampling, sequencing and initial computational steps have been previously described
extensively by Priestley *et al.*[63] and performed at the central sequencing facility at the
Hartwig Medical Foundation.
Mutational signatures were determined using the *deconstructSigs*[64] R package, using the
COSMIC mutational signatures v3 database[65]. Frequently mutated genes were determined
by overlapping genes with somatic mutations affecting protein coding regions in more than
one sample in our dataset, with the mutated genes described in the endometrial carcinoma
publication from TCGA[26] and the OncoKB Cancer Gene List[66]. Overlap between somatic
variants and ChIP-seq regions was determined using *bedtools* v2.25.0 intersect and
multiinter[67].
452

RNA-seq and differential gene expression analyses
454   Ishikawa gene expression data was obtained from GEO, accession numbers GSE109892[25].
455   Differential gene expression between estradiol and DMSO treated cells was computed using
456   *DESeq2*[68].
457

Circularized Chromosome Conformation Capture sequencing (4C-seq)
For each sample, 10-15 30µM-thick slices of flash frozen tissue were cross-linked under
rotation at room temperature for 20 minutes in 5mL of 1× PBS supplemented with 2% FBS
and 2% Methanol-free Formaldehyde. The cross-linked slices have been washed twice in
10mL of ice-cold 1× PBS before to be resupended in 0.5-1mL of ice-cold 1× PBS in 1.5mL
DNA-loBind microcentrifuge tubes. Tissues have been electronically homogenized and
lysates filtered using a 75µm cell strainer. Then, experiments of 4C-seq were performed as
previously described[69]. Briefly, nuclei were isolated and permeabilized to allow digestion of
the chromatin by the primary RE (DpnII, New England Biolabs). Upon dilution, chromatin
fragments have been ligated and then de-crosslinked. DNA was purified and digested with
the secondary RE (NlaIII, New England Biolabs) and circularized by ligation. PCR
amplification of the re-purified circular fragments have been performed using View-Point
specific primers (Reading Primer: 5'-tacacgacgctcttccgatctAACTCGATTTGGAGCGATC-3';

Non-reading Primer: 5'-actggagttcagacgtgtgctcttccgatctCTGGGACTGCACTTGCTC-3')

using the Expand Long Template PCR System (Roche). Amplicons were purified with

AMPure XP beads (Beckman Coulter) in a 0.8× ratio and then amplified using standard

indexed Illumina primers as previously described[69] using the Expand Long Template PCR

System (Roche). Second-round PCR products were purified with PCR purification columns

(Qiagen) and quantified by 2100 Bioanalyzer (Agilent, DNA 7500 kit).

4C libraries have been pooled equimolarly and sequenced using Illumina MiSeq with a 75bp

Single-End reads setup. Fastq files have been demultiplexed by Cutadapt[70] and mapped on

Hg19/GRCh37 genome assembly and signal normalized by *pipe4C*[69] v1.1 R-package in "cis"

mode and with default parameters. The signal of different technical replicates was averaged

to obtain a unique signal for each sample. The, each tissue sample have been considered

as biological replicate. Downstream analyses of 4C-seq data have been performed in an R

v4.0.3 environment by using *get.single.base.score.bw* and *genomic.track* functions from

Rseb[47] (v0.3.0) (https://github.com/sebastian-gregoricchio/Rseb) R-package in combination

with *ggplot2* (v3.3.5) and *ggforce* (v0.3.3) R-packages.

Protein extraction and immunoblotting

For immunoprecipiated proteins, elution buffer (1% SDS, 0.1M NaHCO$_3$) and Laemmli

buffer, supplied with 100mM DTT (Dithiothreitol, Merck Sigma Aldrich), were added to input

samples and beads, and then boiled at 95°C for 30min. For whole-cell extracts, cells have

been lysed using 2× Laemmli buffer supplied with 100mM DTT (Dithiothreitol, Merck Sigma

Aldrich), boiled at 95°C for 10min and then DNA was sheared by sonication (Bioruptor®

Pico, Diagenode; 4 cycles of 30s ON + 90s OFF). In both cases, proteins were then resolved

by SDS-PAGE and transferred to a 0.22µm nitrocellulose membrane (BioRad). Membranes

were incubated for 2h with blocking solution (BS: 1× PBS, 0.1% Tween20, 5% non-fat milk

powder) and then incubated with primary antibody against ERα (ThermoFisher #MA5-14104,

1:1000 in BS), EHMT2/G9a (Abcam #ab133482, 1:1000 in BS) or GAPDH (ThermoFisher

#PA1-987, 1:5000 in BS) overnight at +4°C. After washing with washing solution (1× PBS,

0.1% Tween20), membranes were incubated with secondary antibodies donkey-α-mouse

IRDye® 680RD (926-68073, LI-COR Biosciences, 1:10,000 in BS), donkey-α-mouse

IRDye® 800CW (926-32212, LI-COR Biosciences, 1:10,000 in BS), donkey-α-rabbit IRDye®

800CW (926-32213, LI-COR Biosciences, 1:10,000 in BS) or donkey-α-rabbit IRDye®

680RD (926-68073, LI-COR Biosciences, 1:10,000 in BS) for 1h. Membranes were washed

again with washing solution, scanned and analyzed using an Odyssey® CLx Imaging

System (LI-COR Biosciences) and *ImageStudio™ Lite* v.5.2.5 software (LI-COR

Biosciences).

Sequence-based Machine Learning 3D genome folding predictions

The 3D genome organization perturbation predictions have been performed using a window of 2^{20}bp around the *ESR1* locus employing the Convolutional Neural Networks (CNN) model from *Akita*[29] tool. We used the pre-existing model based on Hi-C data originated from human foreskin fibroblast (HFF). Heatmaps have been generated using *matplotlib.pyplot.matshow*.

DNA-oligo protein pull-down

Ishikawa cells were harvested and washed twice with ice-cold PBS and nuclear extracts were prepared as described previously[71]. The ~50bp oligonucleotide probes encompassing the SNP were ordered with the forward strand containing a 5'-biotin moiety (Integrated DNA Technologies) (Fw_WT: 5'-/5Biosg/ACTGTGGAAACTGGAAGCTGTTCTTGGACTATTTCGCAACACTTTTCTCC-3'; Rv_WT: 5'-GGAGAAAAGTGTTGCGAAATAGTCCAAGAACAGCTTCCAGTTTCCACAGT-3'; Fw_SNP: 5'-/5Biosg/ACTGTGGAAACTGGAAGCTGTTTGGACTATTTCGCAACACTTTTCTCCTC-3', Rv_SNP: (5'-GAGGAGAAAAGTGTTGCGAAATAGTCCAAACAGCTTCCAGTTTCCACAGT-3'; mutated site is underlined). DNA affinity purifications, on-bead trypsin digestion and dimethyl labeling were performed as described[72]. Matching light and medium labelled samples were then combined and analyzed using a gradient from 7 to 30% buffer B in Buffer A over 44min, followed by a further increase to 95% in the next 16min at flow rate of 250nL/min using an Easy-nLC 1000 (Thermo Fisher Scientific) coupled online to an Orbitrap Exploris 480 (Thermo Fisher Scientific). MS1 spectra were acquired at 120,000 resolution with a scan range from 350 to 1300 m/z, normalized AGC target of 300% and maximum injection time of 20 ms. The top 20 most intense ions with a charge state 2-6 from each MS1 scan were selected for fragmentation by HCD. MS2 resolution was set at 15,000 with a normalized AGC target of 75%. Raw MS spectra were analyzed using MaxQuant software (version 1.6.0.1) with standard settings[72,73]. Data was searched against the human UniProt database (downloaded 2017) using the integrated search engine. N-terminal and lysine modification for dimethyl labeling was specified under "labels". Carbamidomethylation was specified as a fixed modification. N-terminal acetylation and methionine oxidation were selected as variable modifications.

Transient small interfering RNA (siRNA) cell transfection

Transient transfections of Ishikawa cell lines was performed according to the manufacturer's instructions using Lipofectamine™ RNAiMAX (Invitrogen) for siRNA knockdown experiments. The ON-TARGETplus™ siRNA SMARTpool targeting human EHMT2/G9a (L-

006937-00-00050) and the siGENOME™ Non-Targeting control siRNA #5 (D-001210-05-20)

were purchased from Dharmacon.

Statistical analysis

All statistical analyses were performed in R versions 3.4.4, 3.5.1 or 4.0.2 (R Core Team

2020, https://www.R-project.org). Enrichment of Ishikawa peaks and somatic mutations in

differentially bound regions was calculated using Fisher's exact test, and, where appropriate,

corrected for multiple testing using the FDR.

DATA AVAILABILITY

Endometrial healthy and tumor tissues ERα and H3K27ac ChIP-seq, Hi-C and, 4C-seq

sequencing data has been deposited at the European Genome-phenome Archive (EGA),

which is hosted by the EBI and the CRG, under accession number EGAS00001007240.

Raw sequencing data of the RNA-seq performed in Ishikawa cells can be found at the Gene

Expression Omnibus (GEO) database under accession number GSE109892[25]. ERα ChIP-

seq data in breast cancer tissues from Singh *et al.*[28] are accessible in GEO under accession

number GSE114737. Ishikawa ChIP-seq data in the form of optimal Irreproducibility

Discovery Rate (IDR) thresholded peaks (GRCh37/Hg19 genome build) were obtained from

the ENCODE data portal[74] (**Extended Data Table 1**). The H3K27ac HiChIP data in Ishikawa

cells have been obtained from O'Mara *et al.*[75] accessible in GEO under accession number

GSE137936. Mass spectrometry data have been deposited at the ProteomeXchange

Consortium through the PRIDE[76] partner repository with the identifier PXD029822. The

breast and endometrial cancer ATAC-seq, gene expression and progression-free survival

used in this this work are in whole or part based upon data generated by the TCGA

Research Network[26,27] (https://www.cancer.gov/tcga).

ACKNOWLEDGEMENTS

We thank members of the Zwart and Bergman labs for valuable feedback, suggestions and

input. We would like to acknowledge the Research High Performance Computing (RHPC)

facility of the Netherlands Cancer Institute (NKI) to have enabled us to perform all the

computations required to analyze the data generated, the NKI Genomics Core Facility for

next-generation sequencing and bioinformatics support, and the NKI-AVL Core Facility

Molecular Pathology and Biobanking for patient samples handling.

This publication and the underlying study have been made possible partly based on data
that Hartwig Medical Foundation has made available to the study through the Hartwig
Medical Database. Further, we would like to thank Hartwig Medical Foundation for facilitating
and supporting whole-genome sequencing data generation, data analyses and data
interpretation.

583
584

AUTHOR CONTRIBUTIONS

Conceptualization: S.G., A.K and W.Z. W.Z. was responsible for project funding. S.G. and
A.K. performed Hi-C and 4C-seq experiments; S.G., K.S. and M.D. performed ChIP/ChIP-
seq and immunoblotting experiments; S.S. performed and analyzed the protein DNA-oligo
pulldowns experiments; S.G., M.H, T.M.S. and A.A.S. analyzed the omics data; T.A.O. and
D.M.G. provided support in the HiChIP data analyses; M.V. provided support for the
proteomics experiments; L.F.A.W. provided bioinformatics support; F.v.L. designed and
performed study for patient sample collection. S.G., A.K. and W.Z. wrote the manuscript,
with input from all co-authors.

594
595

596 ## FUNDING

597 This work was supported by Oncode Institute and *Saxum Volutum*. The Vermeulen lab is
598 part of the Oncode Institute, which is partly funded by the Dutch Cancer Society. T.O. is
599 supported by a National Health and Medical Research Council of Australia Emerging Leader
600 Investigator Fellowship (GNT1173170).

601
602

603 ## COMPETING INTERESTS

604 The authors declare no competing interests.

REFERENCES

1. Sung, H. *et al.* Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA. Cancer J. Clin.* **71**, 209–249 (2021).

2. Gu, B. *et al.* Variations in incidence and mortality rates of endometrial cancer at the global, regional, and national levels, 1990-2019. *Gynecol. Oncol.* **161**, 573–580 (2021).

3. Morice, P., Leary, A., Creutzberg, C., Abu-Rustum, N. & Darai, E. Endometrial cancer. *Lancet Lond. Engl.* **387**, 1094–1108 (2016).

4. Carlson, M. J., Thiel, K. W. & Leslie, K. K. Past, present, and future of hormonal therapy in recurrent endometrial cancer. *Int. J. Womens Health* **6**, 429–435 (2014).

5. Rodriguez, A. C., Blanchard, Z., Maurer, K. A. & Gertz, J. Estrogen Signaling in Endometrial Cancer: a Key Oncogenic Pathway with Several Open Questions. *Horm. Cancer* **10**, 51–63 (2019).

6. Crosbie, E. J. *et al.* Endometrial cancer. *Lancet Lond. Engl.* **399**, 1412–1428 (2022).

7. Heldring, N. *et al.* Estrogen receptors: how do they signal and what are their targets. *Physiol. Rev.* **87**, 905–931 (2007).

8. Flach, K. D. & Zwart, W. The first decade of estrogen receptor cistromics in breast cancer. *J. Endocrinol.* **229**, R43-56 (2016).

9. Droog, M. *et al.* Comparative Cistromics Reveals Genomic Cross-talk between FOXA1 and ERα in Tamoxifen-Associated Endometrial Carcinomas. *Cancer Res.* **76**, 3773–3784 (2016).

10. Gertz, J. *et al.* Distinct Properties of Cell-Type-Specific and Shared Transcription Factor Binding Sites. *Mol. Cell* **52**, 25–36 (2013).

11. Gertz, J., Reddy, T. E., Varley, K. E., Garabedian, M. J. & Myers, R. M. Genistein and bisphenol A exposure cause estrogen receptor 1 to bind thousands of sites in a cell type-specific manner. *Genome Res.* **22**, 2153–2162 (2012).

12. Leeuwen, F. E. van *et al.* Risk of endometrial cancer after tamoxifen treatment of breast cancer. *The Lancet* **343**, 448–452 (1994).

13. Droog, M. *et al.* Estrogen receptor α wields treatment-specific enhancers between morphologically similar endometrial tumors. *Proc. Natl. Acad. Sci.* **114**, E1316–E1325 (2017).

14. Chi, D. *et al.* Estrogen receptor signaling is reprogrammed during breast tumorigenesis. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 11437–11443 (2019).

15. Pomerantz, M. M. *et al.* The androgen receptor cistrome is extensively reprogrammed in human prostate tumorigenesis. *Nat. Genet.* **47**, 1346–1351 (2015).

16. Mazrooei, P. *et al.* Cistrome Partitioning Reveals Convergence of Somatic Mutations and Risk Variants on Master Transcription Regulators in Primary Prostate Tumors. *Cancer Cell* **36**, 674-689.e6 (2019).

17. Pomerantz, M. M. *et al.* Prostate cancer reactivates developmental epigenomic programs during metastatic progression. *Nat. Genet.* **52**, 790–799 (2020).

18. Ross-Innes, C. S. *et al.* Differential oestrogen receptor binding is associated with clinical outcome in breast cancer. *Nature* **481**, 389–393 (2012).

19. Robson, M. I., Ringel, A. R. & Mundlos, S. Regulatory Landscaping: How Enhancer-Promoter Communication Is Sculpted in 3D. *Mol. Cell* **74**, 1110–1122 (2019).

20. Harewood, L. *et al.* Hi-C as a tool for precise detection and characterisation of chromosomal rearrangements and copy number variation in human tumours. *Genome Biol.* **18**, 125 (2017).

21. Hindorff, L. A. *et al.* Potential etiologic and functional implications of genome-wide association loci for human diseases and traits. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 9362–9367 (2009).

22. Boot, A. *et al.* Characterization of colibactin-associated mutational signature in an Asian oral squamous cell carcinoma and in other mucosal tumor types. *Genome Res.* **30**, 803–813 (2020).

23. Cancer Genome Atlas Research Network *et al.* Integrated genomic characterization of endometrial carcinoma. *Nature* **497**, 67–73 (2013).

24. O'Hara, A. J. & Bell, D. W. The genomics and genetics of endometrial cancer. *Adv. Genomics Genet.* **2012**, 33–47 (2012).

25. Vahrenkamp, J. M. *et al.* Clinical and Genomic Crosstalk between Glucocorticoid Receptor and Estrogen Receptor α In Endometrial Cancer. *Cell Rep.* **22**, 2995–3005 (2018).

26. Cancer Genome Atlas Research Network *et al.* The Cancer Genome Atlas Pan-Cancer analysis project. *Nat.*
Genet. **45**, 1113–1120 (2013).

27. Corces, M. R. *et al.* The chromatin accessibility landscape of primary human cancers. *Science* **362**, eaav1898
(2018).

28. Singh, A. A. *et al.* Optimized ChIP-seq method facilitates transcription factor profiling in human tumors. *Life*
Sci. Alliance **2**, (2019).

29. Fudenberg, G., Kelley, D. R. & Pollard, K. S. Predicting 3D genome folding from DNA sequence with Akita.
Nat. Methods **17**, 1111–1117 (2020).

30. Carroll, J. S. *et al.* Genome-wide analysis of estrogen receptor binding sites. *Nat. Genet.* **38**, 1289–1297
(2006).

31. Lin, C.-Y. *et al.* Whole-Genome Cartography of Estrogen Receptor α Binding Sites. *PLOS Genet.* **3**, e87
(2007).

32. Zhang, Y. *et al.* SOX17 is a tumor suppressor in endometrial cancer. *Oncotarget* **7**, 76036–76046 (2016).

33. Hurtado, A., Holmes, K. A., Ross-Innes, C. S., Schmidt, D. & Carroll, J. S. FOXA1 is a critical determinant of
Estrogen Receptor function and endocrine response. *Nat. Genet.* **43**, 27–33 (2011).

34. Yamaguchi, K., Chen, X., Oji, A., Hiratani, I. & Defossez, P.-A. Large-Scale Chromatin Rearrangements in
Cancer. *Cancers* **14**, 2384 (2022).

35. Johnstone, S. E. *et al.* Large-Scale Topological Changes Restrain Malignant Progression in Colorectal
Cancer. *Cell* **182**, 1474-1489.e23 (2020).

36. Yang, Q. *et al.* Alterations in 3D chromatin organization contribute to tumorigenesis of EGFR-amplified
glioblastoma. *Comput. Struct. Biotechnol. J.* **20**, 1967–1978 (2022).

37. Purcell, D. J., Jeong, K. W., Bittencourt, D., Gerke, D. S. & Stallcup, M. R. A distinct mechanism for
coactivator versus corepressor function by histone methyltransferase G9a in transcriptional regulation. *J. Biol.*
Chem. **286**, 41963–41971 (2011).

38. Zhang, X. *et al.* G9a-mediated methylation of ERα links the PHF20/MOF histone acetyltransferase complex to
hormonal gene expression. *Nat. Commun.* **7**, 10810 (2016).

39. Chaturvedi, C.-P. *et al.* Dual role for the methyltransferase G9a in the maintenance of beta-globin gene
transcription in adult erythroid cells. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 18303–18308 (2009).

40. Chaturvedi, C.-P. *et al.* Maintenance of gene silencing by the coordinate action of the H3K9 methyltransferase
G9a/KMT1C and the H3K4 demethylase Jarid1a/KDM5A. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 18845–18850
(2012).

41. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinforma. Oxf.*
Engl. **25**, 1754–1760 (2009).

42. Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137 (2008).

43. Kumar, V. *et al.* Uniform, optimal signal processing of mapped deep-sequencing data. *Nat. Biotechnol.* **31**,
615–622 (2013).

44. Amemiya, H. M., Kundaje, A. & Boyle, A. P. The ENCODE Blacklist: Identification of Problematic Regions of
the Genome. *Sci. Rep.* **9**, 9354 (2019).

45. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential expression
analysis of digital gene expression data. *Bioinforma. Oxf. Engl.* **26**, 139–140 (2010).

46. Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic*
Acids Res. **44**, W160-165 (2016).

47. Gregoricchio, S. *et al.* HDAC1 and PRC2 mediate combinatorial control in SPI1/PU.1-dependent gene
repression in murine erythroleukaemia. *Nucleic Acids Res.* **50**, 7938–7958 (2022).

48. Lopez-Delisle, L. *et al.* pyGenomeTracks: reproducible plots for multivariate genomic datasets. *Bioinformatics*
37, 422–423 (2021).

49. Ramírez, F. *et al.* High-resolution TADs reveal DNA sequences underlying genome organization in flies. *Nat.*
Commun. **9**, 189 (2018).

50. Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-regulatory elements
required for macrophage and B cell identities. *Mol. Cell* **38**, 576–589 (2010).

51. McLeay, R. C. & Bailey, T. L. Motif Enrichment Analysis: a unified framework and an evaluation on ChIP data.
BMC Bioinformatics **11**, 165 (2010).

52. Bailey, T. L., Johnson, J., Grant, C. E. & Noble, W. S. The MEME Suite. *Nucleic Acids Res.* **43**, W39-49
(2015).

53. Castro-Mondragon, J. A. *et al.* JASPAR 2022: the 9th release of the open-access database of transcription
factor binding profiles. *Nucleic Acids Res.* **50**, D165–D173 (2022).

54. Sandelin, A., Alkema, W., Engström, P., Wasserman, W. W. & Lenhard, B. JASPAR: an open-access
database for eukaryotic transcription factor binding profiles. *Nucleic Acids Res.* **32**, D91-94 (2004).

55. Sherman, B. T. *et al.* DAVID: a web server for functional enrichment analysis and functional annotation of
gene lists (2021 update). *Nucleic Acids Res.* **50**, W216–W221 (2022).

56. Donaldson-Collier, M. C. *et al.* EZH2 oncogenic mutations drive epigenetic, transcriptional, and structural
changes within chromatin domains. *Nat. Genet.* **51**, 517–528 (2019).

57. Gregoricchio, S. & Zwart, W. snHiC: a complete and simplified snakemake pipeline for grouped Hi-C data
analysis. *Bioinforma. Adv.* **3**, vbad080 (2023).

58. van der Weide, R. H. *et al.* Hi-C analyses with GENOVA: a case study with cohesin variants. *NAR Genomics*
Bioinforma. **3**, lqab040 (2021).

59. Wolff, J. *et al.* Galaxy HiCExplorer 3: a web server for reproducible Hi-C, capture Hi-C and single-cell Hi-C
data analysis, quality control and visualization. *Nucleic Acids Res.* **48**, W177–W184 (2020).

60. Wolff, J. *et al.* Galaxy HiCExplorer: a web server for reproducible Hi-C data analysis, quality control and
visualization. *Nucleic Acids Res.* **46**, W11–W16 (2018).

61. Khan, A. & Mathelier, A. Intervene: a tool for intersection and visualization of multiple gene or genomic region
sets. *BMC Bioinformatics* **18**, 287 (2017).

62. Chakraborty, A., Wang, J. G. & Ay, F. dcHiC detects differential compartments across multiple Hi-C datasets.
Nat. Commun. **13**, 6827 (2022).

63. Priestley, P. *et al.* Pan-cancer whole-genome analyses of metastatic solid tumours. *Nature* **575**, 210–216
(2019).

64. Rosenthal, R., McGranahan, N., Herrero, J., Taylor, B. S. & Swanton, C. deconstructSigs: delineating
mutational processes in single tumors distinguishes DNA repair deficiencies and patterns of carcinoma
evolution. *Genome Biol.* **17**, 31 (2016).

65. Tate, J. G. *et al.* COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids Res.* **47**, D941–
D947 (2019).

66. Chakravarty, D. *et al.* OncoKB: A Precision Oncology Knowledge Base. *JCO Precis. Oncol.* **2017**,
PO.17.00011 (2017).

67. Quinlan, A. R. BEDTools: The Swiss-Army Tool for Genome Feature Analysis. *Curr. Protoc. Bioinforma.* **47**,
11.12.1-34 (2014).

68. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data
with DESeq2. *Genome Biol.* **15**, 550 (2014).

69. Krijger, P. H. L., Geeven, G., Bianchi, V., Hilvering, C. R. E. & de Laat, W. 4C-seq from beginning to end: A
detailed protocol for sample preparation and data analysis. *Methods San Diego Calif* **170**, 17–32 (2020).

70. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17**,
10–12 (2011).

71. Vermeulen, M. Identifying chromatin readers using a SILAC-based histone peptide pull-down approach.
Methods Enzymol. **512**, 137–160 (2012).

72. Makowski, M. M. *et al.* An interaction proteomics survey of transcription factor binding at recurrent TERT
promoter mutations. *Proteomics* **16**, 417–426 (2016).

73. Cox, J. & Mann, M. MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass
accuracies and proteome-wide protein quantification. *Nat. Biotechnol.* **26**, 1367–1372 (2008).

74. Korch, C. *et al.* DNA profiling analysis of endometrial and ovarian cell lines reveals misidentification,
redundancy and contamination. *Gynecol. Oncol.* **127**, 241–248 (2012).

75. O'Mara, T. A., Spurdle, A. B., Glubb, D. M., & Endometrial Cancer Association Consortium. Analysis of
Promoter-Associated Chromatin Interactions Reveals Biologically Relevant Candidate Target Genes at
Endometrial Cancer Risk Loci. *Cancers* **11**, 1440 (2019).

76. Perez-Riverol, Y. *et al.* The PRIDE database resources in 2022: a hub for mass spectrometry-based
proteomics evidences. *Nucleic Acids Res.* **50**, D543–D552 (2022).

Fig. 1 | ERα cistrome changes upon tumorigenesis.

A. Schematic workflow of the multi-omics approach applied in this work.

B/C. Boxplot depicting the distribution of peak number detected by ERα (**B**) or H3K27ac (**C**)

ChIP-

seq in healthy (blue) or tumor (orange) endometrial tissues.

D. Venn diagram of the overlap between consensus ERα binding sites detected in healthy

(blue)

and tumor (orange) tissues.

E. MA-plot of differential binding analyses performed on the consensus ERα ChIP-seq peaks

(Healthy vs Tumor). Differential binding sites (FDR ≤ 0.05 & $|\log_2(\text{FoldChange})| \geq 1$) are

highlighted in pink.

F/G. Tornado plot of the ERα (**F**) or H3K27ac (**G**) ChIP-seq signal at tumor-depleted (blue,

upper

blocks) or tumor-enriched (orange, lower blocks) ERα consensus peaks for all the 4

healthy

(blue, left heatmaps) and 5 tumor (orange, right heatmaps) endometrial tissues. On the

top of each heatmap is plotted the average density signal for each region group.

H. Bar plot with counts per genomic localization of tumor-depleted (blue) and tumor-enriched

(orange) ERα consensus peaks.

I. Wordcloud of the top-50 enriched motifs at tumor-depleted (blues) and tumor-enriched

(oranges) ERα consensus peaks. Size and color intensity are proportional to the -

$\log_{10}(\text{E-value})$.

J. The upper part of the heatmap shows the individual tumor-depleted (blues) and tumor-

enriched (oranges) ERα consensus peaks colored by signal intensity. On the lower part,

each

black bar represents an overlapping peak of ChIP-seq data of several targets publicly

available for the Ishikawa endometrial cancer cell line.

K. Heatmap of the percentage of tumor-depleted or tumor-enriched ERα consensus peaks

overlapping with each ChIP-seq target in Ishikawa cells from **j**. Ranking is performed by

descending number of overlaps in tumor-depleted peaks.

Fig. 2 | 3D genome landscape is remodeled during tumorigenesis.

A. Schematic of the Hi-C library preparation starting from fresh frozen tissues. Ten 30µm-thick slices of flash-frozen tissue are cross-linked 25min in 2% Formaldehyde. Then, electronically homogenized tissues are filtered using a 75µm cell strainer and subjected to Hi-C library preparation. Hi-C libraries are then sequenced and resulting reads are analyzed by the *snHiC* pipeline.

B. On the left, 40kb-resolution matrices of the average Hi-C score in Tumor (n = 3) and Healthy endometrial tissues (n = 3) at chromosome 6. On the right, differences of the scores shown on the left side (Tumor – healthy).

C. Relative Hi-C Contact Probability (RCP) as function of the distance for each individual sample (40kb resolution).

D. Violin plot of the distribution of the short-range (<2Mb) over long-rage (>2Mb) Hi-C contacts ratio in each individual sample (40kb resolution). For each comparison the Wilcoxon's test *P*-value is indicated.

E. Distribution of the Hi-C loop distance (10kb resolution) for loops overlapping with tumor-depleted, tumor-enriched or shared ERα consensus peaks in combined healthy (blue) or tumor (orange) endometrial tissues.

F. Compartment polarization ratio (100kb resolution), defined as (AA + BB) / (AB + BA), for each individual sample.

G. Top: saddle plot of A/B compartments interactions as computed in **F** for each individual sample. Bottom: difference of saddle-score with the reference H.005.A2 (healthy tissue), where orange indicates an higher score in tumor samples while purple an higher score in healthy tissues.

H. Sankey plot depicting the proportion of compartment state transition (100kb resolution) of healthy compartments (left) upon tumorigenesis (right).

I. Upset plot showing the overlaps between different compartment transition states (100kb resolution) in healthy compared to tumor as in **H** (A-to-A, B-to-B, A-to-B, B-to-A) and different categories of ERα consensus peaks.

J. Stacked bar plot of the proportion of compartment transition status as in **H** for individual genomic bins overlapping with only one of the different categories of ERα consensus peaks.

Fig. 3 | Tumor-enriched ERα binding sites represent non-coding regions target of somatic mutation in metastatic endometrial cancer tumor.

A. Scheme of the number metastases and metastatic site, for all metastatic samples used for the WGS analyses.

B/C. Genomic localization (**b**) and mutational signature (**c**) distribution of the somatic mutations

detected by WGS in the metastatic samples described in **a**.

D. In the outer plot, it is shown the number of different types of somatic variants detected in each metastatic sample. In the inner plot, stacked bar plot of the most-frequently protein sequence mutated genes and relative number and type of somatic variants identified in metastatic samples.

E. ERα consensus peaks in primary tumors were ranked by differential signal Fold Change (Tumor vs Healthy). On the top is plotted the number of somatic variants overlapping with each individual ERα peak: higher the amount of overlapping somatic variants darker the bar color.

F. MA-plot of differential expression analyses in Ishikawa cell lines upon 8h of 10nM estradiol (E2) stimulation. Black dots indicate genes which promoter has been linked, by H3K27ac Hi-C analyses, to a tumor-depleted or tumor-enriched ERα-bound regulatory element bearing somatic mutations in metastatic samples (n=311). If one these genes was differentially expressed ($|\log_2$(Fold Change expression)$| \geq 1$ & p-value adjusted $<$ 0.05) upon estradiol stimulation, it has been then labeled and highlighted by a red dot (n=12).

G. Ishikawa endometrial cancer cells were treated or not for 6 or 12 hours with 10nM estradiol. Whole-cell extracts were analyzed by immunoblotting with antibodies against ERα and GAPDH (loading control).

H. Progression-Free Kaplan-Meier curve of endometrial cancer patients (TCGA data) divided into two groups using the median of *ESR1* expression (FPKM) as cut-off.

Fig. 4 | Short-range chromatin contacts at the *ESR1* locus are stronger in tumors.

A. ChIP-seq genomic tracks for ERα (upper panel) and H3K27ac (lower panel) in healthy
(blue) and tumor (orange) endometrial primary tissues at the *ESR1* locus. *ESR1*
Enhancer 1 and *Enhancer 2* are indicated, as well as the mutations found by WGS
analyses in metastatic samples.

B. Representation of the averaged 40kb resolution Hi-C matrix at the *ESR1* locus for the 3
healthy (top) and 3 tumor (middle) tissues or the score difference (bottom) tumor –
healthy, where orange indicates higher scores in the tumors while purple higher scores
in healthy. The matrices scores are dived by the sum of the matrix. Black lines indicate
the Topologically Associated Domains (TADs) identified.

C. 4C-seq genomic tracks at the *ESR1* locus using the *ESR1* TSS as View-Point (VP) for 2
healthy (top, blue) and 3 tumor (middle, orange) endometrial tissues, and the average
difference of score tumor – healthy (bottom). With green arcs are depicted loops
detected by H3K27ac HiChIP in Ishikawa endometrial cancer cells. The ribbon around
the 4C-seq signal lines indicates the Standard Error Mean (SEM) among biological
replicates.

D. In the top row, Observed/Expected matrices of sequence-based machine learning
prediction in a ±500kb window surrounding the *ESR1* locus. In order from left to right can
be found: wild-type sequence, point mutation in a genomic desert (negative control),
deletion of the full *Enhancer 1* sequence (positive control), introduction of SNPs found by
WGS analyses of metastatic endometrial cancer samples. On the bottom row, is showed
the difference of Observed/Expected score over the wild-type sequence, where orange
indicates higher scores in the altered sequence and purple a higher score in the wild-
type one.

Fig. 5 | EHMT2/G9a is a negative regulator of *ESR1* expression in endometrial cancer.

A.   Schematic workflow used to perform DNA-oligo protein pull-downs. Biotin-conjugated

wild-type or mutated DNA-oligos are immobilized on streptavidin magnetic beads and

mixed with Ishikawa nuclear lysates. Captured proteins are then dimethyl labeled and

analyzed by mass spectrometry.

B.   A DNA oligo with the sequence of the ±25bp surrounding the ERα binding site in the

ESR1 Enhancer 1, in wild-type or chr6:152,002,679–TCT-to-T form, was used to perform

DNA-oligo protein pull downs in Ishikawa cells as described in **a**. The scatter plot shows

the $\log_2$ ratios of all identified and quantified proteins in both experiments plotted against

each other. Proteins significantly enriched at the wild type sequence are highlighted in

red, and proteins significantly enriched at the mutant sequence are highlighted in blue. .

C.   Gene expression correlation heatmap for all corresponding 24 differentially bound

proteins identified in **b** in 589 endometrial cancer patients (TCGA). Dashed lines indicate

the separation between positive and negative correlation scores. Genes are ranked by

correlation with *ESR1* gene expression.

D.   Progression-Free Kaplan-Meier curve of endometrial cancer patients (TCGA data)

divided into two groups using the median of *EHMT2/G9a* expression (FPKM) as cut-off.

E.   Ishikawa endometrial cancer cells were cross-linked and DNA sheared by sonication.

Protein lysates were then immunoprecipitated (IP) with antibodies against EHMT2/G9a

and IgG (negative control) to be then immunoblotted with antibodies against EHMT2/G9a

and ERα. For EHMT2/G9a, two different exposure images were used for input and IP, as

indicated by the vertical dashed bar.

F.   Ishikawa cells were stimulated for 6h with 10nM estradiol before performing EHMT2/G9a

ChIP. Quantitative PCR (qPCR) analyses were performed to determine the percentage

of enrichment over the input (% Input) at the *ESR1 Enhancer 1*, *ESR1* promoter and

CDK12 promoter (negative control). Mean of 2 independent experiments is shown.

G.   Ishikawa endometrial cancer cells were treated or not for 72 hours with 10nM estradiol in

combination or not with 100nM ICI (Fulvestrant). In all conditions, cells where incubated

either with a Non-Targeting (NT) siRNA or with an siRNA against EHMT2/G9a. Then,

whole-cell extracts were analyzed by immunoblotting with antibodies against ERα,

EHMT2/G9a and GAPDH (loading control).

H.   Schematic representation of the *ESR1* gene regulation working model. Upon

tumorigenesis, ERα is re-located to an endometrial cancer specific *ESR1* enhancer

where interacts with EHMT2/G9a and comes in contact with the *ESR1* promoter. These

changes lead to a decrease in *ESR1* gene expression. In more advanced endometrial

cancer stages, a somatic mutation is acquired at the tumor-specific *ESR1* enhancer

leading to a loss of EHMT2/G9a chromatin occupancy which allows re-expression of

ESR1.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5