Enhancer plasticity in endometrial tumorigenesis demarcates non-coding somatic mutations and 3D-genome alterations boosting the oncogenic driver *ESR1*

Sebastian Gregoricchio[1,*], Aleksandar Kojic[1], Marlous Hoogstraat[1,2], Karianne Schuurman[1], Suzan Stelloo[3], Tesa M. Severson[1,2], Tracy A. O'Mara[4], Marjolein Droog[1], Abhishek A. Singh[1], Dylan M. Glubb[4], Lodewyk F.A. Wessels[2], Michiel Vermeulen[3,5], Flora E. van Leeuwen[6], Wilbert Zwart[1,*]

[1] Division of Oncogenomics, Oncode Institute, The Netherlands Cancer Institute, Plesmanlaan 121, 1066CX, Amsterdam, The Netherlands

[2] Division of Molecular Carcinogenesis, The Netherlands Cancer Institute, Plesmanlaan 121, 1066CX, Amsterdam, The Netherlands

[3] Department of Molecular Biology, Faculty of Science, Radboud Institute for Molecular Life Sciences, Oncode Institute, Radboud University Nijmegen, Geert Grooteplein Zuid 28, 6525GA, Nijmegen, The Netherlands

[4] Cancer Research Program, QIMR Berghofer Medical Research Institute, Locked Bag 2000, QLD 4029, Brisbane, Australia

[5] Division of Molecular Genetics, Oncode Institute, The Netherlands Cancer Institute, Plesmanlaan 121, 1066CX, Amsterdam, The Netherlands

[6] Department of Epidemiology, The Netherlands Cancer Institute, Plesmanlaan 121, 1066CX, Amsterdam, The Netherlands

* To whom correspondence should be addressed. e-mail: s.gregoricchio@nki.nl and w.zwart@nki.nl

ABSTRACT

The incidence and mortality of Endometrial Cancer (EC) is on the rise. 85% of ECs depend on Estrogen Receptor alpha (ERα) for proliferation, but little is known about its transcriptional regulation in these tumors.

We generated epigenomics, transcriptomics and Hi-C datastreams in healthy and tumor endometrial tissues, identifying robust ERα reprogramming and profound alterations in 3D genome organization that lead to a gain of tumor-specific enhancer activity during EC development. Integration with endometrial cancer risk single-nucleotide polymorphisms, as well as WGS data from primary tumors and metastatic samples revealed a striking enrichment of risk variants and non-coding somatic mutations at tumor-enriched ERα sites. Through machine learning-based predictions and interaction proteomics analyses, we identified an enhancer mutation which alters 3D genome conformation, impairing recruitment of the transcriptional repressor EHMT2/G9a/KMT1C, thereby alleviating transcriptional repression of *ESR1* in EC.

39   In summary, we identified a complex genomic-epigenomic interplay in EC
40   development and progression, altering 3D genome organization to enhance expression of
41   the critical driver ERα.

INTRODUCTION

Endometrial cancer (EC) is the second most-common gynecological cancer with over

417,000 new cases and 97,000 deaths in 2020, with global incidence rates increasing every

year[1]. Surprisingly, and in stark contrast to most other cancers, survival rates for endometrial

cancer are decreasing[1,2]. This gradual deterioration of EC survival is likely related to the

relatively understudied nature of the disease, with many molecular mechanisms driving

tumor development and progression remaining largely elusive.

Endometrial tissue is under tight endocrine control, and this remains the case upon

tumorigenesis. The majority of endometrial cancers (85%) are classified as low grade

endometrioid tumors, expressing Estrogen Receptor alpha (ERα) and Progesterone

Receptor (PR)[3–5]. PR agonists are used in the treatment of endometrial tumors as alternative

to chemotherapy and hysterectomy to retain uterine function in young patients[6].

Furthermore, the estrogen competitive ERα antagonist tamoxifen is also prescribed to

patients with endometrial cancer, often alternated with PR agonists (reviewed in ref [7]). Thus,

while PR acts in a tumor suppressive manner, ERα serves as driver of tumor progression

which is therapeutically blocked in endometrial cancer care.

ERα is an enhancer-acting transcription factor, regulating expression of its responsive genes

through long-range 3D genome interactions[8,9]. To date, most literature on promoter-

enhancer communication and ERα activity has been focused on breast cancer[9], and is far

less understood in endometrial tumors. Between breast cancer and endometrial cancer,

substantial differences are observed in ERα chromatin binding profiles[10–12], highlighting the

highly context-dependent nature of ERα genomic action. Paradoxically, while tamoxifen can

be prescribed in the treatment of endometrial cancer, its use in the treatment of breast

cancer is also reported as risk factor for endometrial tumorigenesis[13]. Previously, we showed

that tamoxifen treatment in endometrial tissue reprograms the ERα chromatin interaction

landscape, phenocopying profiles found in breast cancer cells, driving endometrial tumor

growth[14]. These observations highlighted that the endometrial cancer epigenome is not

fixed, but rather dynamically affected by oncogenic factors.

Both ERα in breast tissue[15], and Androgen Receptor (AR) in prostate tissue[16–18],

undergo extensive reprogramming throughout the genome during tumorigenesis. In case of

prostate cancer, both somatic mutations and risk single-nucleotide polymorphisms (SNPs)

are enriched at tumor-gained AR sites, with only a small fraction of which causally affecting

transcriptional output[17]. In endometrial cancer, such studies have thus far not been reported,

and functional interplay between somatic mutations, epigenetic alterations and the impact on

3D genome organization remain fully elusive.

Here, we investigated the plasticity of ERα genomic action upon endometrial tumor
development, by comparing the epigenome of primary human healthy and tumor
endometrial tissues (ERα and H3K27ac ChIP-seq) and the effects on 3D genome
organization (Hi-C, 4C-seq and, H3K27ac HiChIP). Secondly, we studied the crosstalk
between epigenetic alterations and somatic variant events (WGS) that are associated with
EC progression. We identified substantial epigenetic reprogramming upon tumorigenesis
that results in the ERα re-localization at tumor-specific regions throughout the genome,
which coincided with the occurrence of somatic variants in metastatic samples. In particular,
we discovered an EC-specific *ESR1* enhancer which is selectively found mutated in
metastatic EC. *In vitro* analyses show diminished capacity to bind lysine methyl-transferase
G9a/EHMT2/KMT1C to this region when mutated, and perturbation of G9a expression
enhanced ERα expression in cell line studies. Cumulatively, we show that non-coding
mutations in endometrial cancer may have direct pro-tumorigenic potential, through a tight
interplay with epigenetic alterations and changes in the 3D genome structure.

RESULTS

ERα enhancer plasticity in endometrial tumorigenesis

To investigate the role of ERα signaling in endometrial tumor development, we first sought to assess how the ERα cistrome differs between endometrial tumors and healthy endometrial tissue. We collected five fresh-frozen endometrial tumors from post-menopausal patients that did not receive any estrogen receptor targeting for cancer therapy, as well as four fresh-frozen samples from post-menopausal women with pathologically normal endometrial tissue (**Fig. 1a** and **Supplementary Fig. 1a**, for clinicopathological characteristics see **Supplementary Table 1**). All samples were subjected to ERα and H3K27 acetylation (H3K27ac) ChIP-seq in order to identify active regulatory elements that are under ERα control. On average, we identified 20,815 ERα peaks (range 1,399-51,888) and 44,824 H3K27ac peaks (range 8,791-71,898) in our samples. The ChIP-seq library size was comparable among the different samples for both targets (**Supplementary Fig. 1b**). Interestingly, the number of ERα peaks was higher in tumor samples compared to healthy tissue (Mann-Whitney P = 0.063, **Fig. 1b**) while the global number of H3K27ac peaks for these samples was comparable between both groups (Mann-Whitney P = 0.29, **Fig. 1c**). To assess data quality and reproducibility, we performed Principal Component Analysis (PCA) on the peak intensities which revealed a clustering of the samples by ChIP target (first component) and tissue type (second component) (**Supplementary Fig. 1c**). Furthermore, given the relatively small sample size, we tested for over-training and performed unsupervised permutation clustering analyses that showed a clear separation of the two tissues types, except for the sample T33 showing mixed features, between normal and tumor tissues (**Supplementary Fig. 1d**), possibly due to a lower tumor cell representation in those sections that were used for ChIP-seq studies.

Differential binding analysis (*DiffBind*[19]) for ERα between tumor and normal samples revealed 10,292 differentially-bound genomic locations (FDR ≤ 0.05 & |log$_2$(FoldChange)| ≥ 1) (**Fig. 1d-e** and **Supplementary Fig. 1f**), of which 6,488 ERα sites were lost and 3,804 ERα sites gained in tumor samples, hereafter referred to as "tumor-depleted" (blue) and "tumor-enriched" (orange) ERα binding sites, respectively. Tornado plots depicting normalized ERα ChIP-seq signal for all 10,292 differential-enriched ERα sites (**Fig. 1f**), and label-swap permutation differential binding analyses (**Supplementary Fig. 1e**), confirmed the selective enrichment of signals between samples, with high reproducibility. As ERα is a transcription factor that acts predominantly by occupying enhancers[8,9], we next analyzed the differential ERα sites for presence of H3K27ac, inferring regulatory element activity. Interestingly, persistent activity in both tissue states was observed at tumor-depleted ERα

126 sites, while tumor-enriched ERα sites showed a marked increase of activity upon
127 tumorigenesis (**Fig. 1g**).

128   Collectively, these data suggest a selective gain of regulatory element potential upon
129 tumorigenesis, marked by dynamically altered ERα sites.

Genomic distributions and transcription complex composition at differential ERα sites

To further investigate the nature of the ERα regions differentially bound upon
tumorigenesis, we first inspected their genome-wide distribution with respect to different
genomic elements. In agreement with previous reports from us (tumors[10,14]) and others (cell
lines[11]), tumor-enriched ERα sites were mostly found at distal intergenic and intronic regions
(**Fig. 1h**) and at larger distance from the associated Transcription start site (TSS) (**Fig. 1i**).
However, tumor-depleted ERα sites, selectively occupied by ERα in healthy endometrial
tissue, display a strong enrichment for promoter regions of genes involved in cell cycle
regulation and DNA damage processes (**Supplementary Fig. 1g**). To determine whether
loss of ERα binding at these promoters leads to transcriptional deregulation of the
associated genes in tumor samples, we performed RNA sequencing (RNA-seq) in 3 healthy
and 3 tumor endometrial tissues from post-menopausal patients that did not receive any
prior estrogen receptor targeting for cancer therapy (**Supplementary Table 1**). Principle
component analyses display a clear separation of normal and tumor tissues with a higher
variability for tumor samples than healthy ones (**Supplementary Fig. 2a**). Differential
expression analyses identified 1945 genes to be down-regulated and 2392 to be up-
regulated in tumors, compared to healthy tissue (**Supplementary Fig. 2b**). Next, we
analyzed the expression of the tumor-depleted ERα promoter-bound genes included in the
previously identified over-represented biological processes (**Supplementary Fig. 1g** and
Supplementary Fig. 2c). This investigation confirmed our former results highlighting a
significant deregulation of cell division and DNA-repair related genes (**Supplementary Fig.
2c**). Furthermore, results from Gene Set Enrichment Analyses (GSEA) for the "hallmarks"
(H) dataset also corroborated with these findings in a more global manner (**Supplementary
Fig. 2d**). Of note, we observed that differentially expressed genes associated with tumor-
enriched ERα binding sites were mostly up-regulated in tumor samples compared to genes
associated with tumor-depleted ERα binding sites (Fisher's exact $P < 2.2 \times 10^{-16}$, OR 6.8,
95% CI 4.35-10.84; **Supplementary Fig. 2e-f**).

Consensus motifs for ERα/β (ESR1/2) and AP-1 (JUN, FOS) binding sites were
specifically enriched at tumor-depleted ERα regions, whereas tumor-enriched regions
demonstrated an overabundance of SOX and FOX transcription factor family member motifs

(**Fig. 1j**). Interestingly, tumor-enriched ERα binding sites, despite the consensus ERα/ESR1

motif being poorly detected, do harbor this motif in around 85% of the cases in a measure

comparable to the tumor-depleted sites (**Supplementary Fig. 1h**), albeit highly degenerated

as compared to tumor-depleted or common ones (**Supplementary Fig. 1i**). Differential motif

enrichment analyses led us to hypothesize that the ERα transcription complex may differ

between tumor-enriched and tumor-depleted sites. To test this hypothesis, we next

integrated our tumor-enriched and tumor-depleted sites with publicly available ChIP-seq

data from the Ishikawa endometrial cancer cell line (**Fig. 1k** and **Supplementary Table 2**),

revealing striking differences between transcription and epigenetic factors on chromatin

occupancy for the two ERα binding site categories. Interestingly, ERα chromatin binding in

hormone-deprived cells showed a significant overlap with the tumor-depleted ERα peaks

(adjusted Fisher's exact $P = 3.15 \times 10^{-21}$), while ERα peaks from estradiol-stimulated cells

showed significant overlap with the tumor-enriched ERα peaks (adjusted Fisher's exact $P =$

4.60×10^{-59}, **Fig. 1l**). In addition, both FOXM1 and FOXA1 binding sites in Ishikawa showed

significant overlaps with our tumor enriched regions (adjusted Fisher's exact $P = 3.37 \times 10^{-21}$

and $P = 6.25 \times 10^{-57}$, respectively, **Fig. 1l**). These findings were successfully confirmed by

GIGGLE[20] analyses (testing for overlap in a large repository of thousands of publicly

available ChIP-seq datasets) at differential ERα binding sites (**Supplementary Fig. 1j**)

implicating a strongly divergent transcription complex composition between tumor-enriched

and tumor-depleted ERα sites.

Altogether, these data suggest prominent enhancer plasticity in endometrial

tumorigenesis, with ERα relocating to alternatively engaged and activated non-canonical

enhancers in endometrial cancer.

187 3D high-order chromatin structures are altered in tumors

Enhancer regions can modulate gene transcription through interactions with

promoters in 3D genomic space[21]. With the observed plasticity of ERα chromatin profiles in

tumorigenesis, we hypothesized that these epigenetic alterations were accompanied by

reorganization of the 3D chromatin structure. To test this hypothesis, we performed high-

throughput chromosome conformation capture (Hi-C) analyses on healthy endometrial tissue

(n=3) and primary tumors (n=3) derived from post-menopausal women that did not receive

any previous systemic therapy (**Fig. 2a,** for clinicopathological features see **Supplementary**

Table 1). Hi-C is particularly suitable to study translocations in cancer[22], which were not

observed in any of the tumor samples we analyzed (**Supplementary Fig. 3a**). We did

however observe a distinct clustering of the healthy tissues from the tumor specimens,

based on 3D genome organization (**Supplementary Fig. 3b**) and independent of Hi-C
contact quality bias, intrinsic to the tissue type (**Supplementary Fig. 3c**). In addition, tumor
samples displayed a higher Hi-C contact heterogeneity in comparison to the more correlated
healthy tissues (**Supplementary Fig. 3c**). Next, we quantified the Relative Hi-C Contact
Probability (RCP) as function of the contact distances. Interestingly, tumors displayed an
increased probability of shorter-range interactions (<2Mb) relative to healthy tissue, which is
concomitant with a loss of longer-range chromatin contacts (**Fig. 2b-d** and **Supplementary**
Fig. 3d-e), highlighting a major 3D genome reorganization in tumor samples independently
of the inter-individual variability. Furthermore, analyzing the distribution of the loop distances
(**Fig. 2e** and **Supplementary Fig. 4**), we found that loops whose anchors include tumor-
enriched ERα binding sites tend to be shorter in tumor tissues than in healthy tissues. On
the other hand, tumor-depleted and shared ERα binding sites do not show differences in
loop length between two tissues stages. We therefore investigate the distribution of the Hi-C
contacts as function of the distance from different categories of ERα binding sites, by
performing paired-end spatial chromatin analysis (PE-SCAn[23]) between healthy and tumor
tissues (**Fig. 2f**). We observed that shared and tumor-depleted ERα binding sites lose
chromatin interaction in their surrounding regions in tumor tissues, whereas tumor-enriched
binding sites show an increase in relatively shorter-range interactions (**Fig. 2f**). We next
wondered whether the loop shortening, and increased chromatin contacts, at tumor-enriched
binding sites in tumor samples was associated with changes in chromatin
compartmentalization. Analysis of compartment polarization (**Fig. 2b,g-i** and
Supplementary Fig. 3f-g) revealed decreased compartment strength in tumor samples
compared to healthy tissues. In particular, we observed that A compartments (euchromatin)
were more robust in healthy samples, while B compartments (heterochromatin) were slightly
stronger in tumor tissues (**Fig. 2h** and **Supplementary Fig. 3g**). Overall, around one third of
compartments (A: 565/1431 (39.5%); B: 440/1306 (33.6%)) underwent class switching in
tumor samples (**Fig. 2i**). To investigate whether this compartment class switch was
occurring specifically in compartments where ERα was bound in a tissue type-specific
manner, we overlapped the compartments with the differential ERα occupied regions upon
tumorigenesis (**Fig. 2j**). We found that compartments switching from A to B class were
enriched for tumor-depleted ERα binding sites, relative to tumor-enriched ones (A-to-B
depleted/enriched ratio 2.65 (135/51)). In contrast, compartments switching from B-to-A
were clearly enriched for tumor-gained ERα binding sites (B-to-A depleted/enriched ratio
0.33 (35/105)). These findings were further confirmed studying the behavior of smaller
genomic regions (compartment bins) relative to differentially-occupied ERα binding sites,
excluding regions that harbor multiple ERα binding site categories (Fisher's exact test: B-to-

A depleted-vs-enriched: $P = 1.317×10^{-56}$, OR 0.20; A-to-B depleted-vs-enriched: $P =$
$6.013×10^{-11}$, OR 1.87) (**Fig. 2k** and **Supplementary Fig. 3h**).

Taken together, our findings highlight a widespread reorganization of the 3D genome
upon tumorigenesis that leads to a loss of chromatin compartment polarization. This was
accompanied by an extensive class-switching, with tumor-enriched ERα binding sites being
enriched in B-to-A compartments. Moreover, we identified a gain of relatively shorter-range
chromatin interactions upon tumor development, at the expense of longer-range contacts.

241
242

Somatic mutations are selectively found at a higher frequency at tumor-
enriched ERα binding sites

Recently, pan-cancer whole-genome analysis highlighted the presence of driver
mutations in non-coding regulatory elements[24]. To determine whether that might also be the
case in endometrial cancer, we re-analyzed Whole-Genome Sequencing (WGS) data of 41
primary endometrial cancer samples from the TCGA Uterine Corpus Endometrial Carcinoma
(UCEC) cohort[25] (see Methods for details on the cases selection). Most of the mutations
were located in intronic and intergenic non-coding regions (**Fig. 3a**). Mutational signatures
(**Fig. 3b**) and variant counts (**Fig. 3c**) well correlated with the Micro-Satellite Stability (MSS)
status of the source samples. These analyses show that most of the Micro-Satellite Instable
(MSI) samples are characterized by UV light exposure (SBS7c) and defective DNA
mismatch repair (SBS15/44) signatures (**Fig. 3b**), accompanied by a higher number of
mutations per sample (**Fig. 3c**). On the other hand, Micro-Satellite Stable (MSS) samples,
as expected, display a lower number of somatic mutation counts (**Fig. 3c**) and prevalently
mutational signatures associated to defective DNA (polymerase ε exonuclease domain
mutations, SBS10b) (**Fig. 3b**).

To investigate whether somatic mutations in primary tumors are associated with
selective ERα activity, we compared the number of somatic mutations that are overlapping
with the different categories of ERα binding sites. We found that tumor-enriched binding
sites have a higher representation of somatic mutations over the tumor-depleted ones
(Fisher's exact $P = 1.143×10^{-5}$, OR 1.89, 95% CI 1.41-2.53; **Fig. 3d**). Notably, the somatic
variants overlapping with the ERα binding sites are carried almost exclusively by hyper-
mutated samples (**Supplementary Fig. 6a**). Interestingly, analyzing the enrichment of
Genome-Wide Association Study (GWAS) loci associated with endometrial cancer risk at
ERα-binding sites[26], we observed that tumor-enriched binding sites are significantly enriched
($P_{adj} = 0.006$) for endometrial cancer risk single-nucleotide polymorphisms (rSNP) in
comparison to tumor-depleted regions ($P_{adj} = 0.995$) (**Fig. 3e**). These data imply that both

germline as well as somatic variants are not equally distributed over the genome, but rather
show a selective occurrence at tumor-gained ERα sites.

To investigate whether somatic variant enrichment for particular ERα sites also
persisted after endometrial cancer progression, we next analyzed WGS data of 26 fresh
frozen biopsy samples, derived from 24 cases of advanced endometrial cancer. Biopsies
were collected from tumors that metastasized to lung, liver, and other abdominal regions
(**Fig. 3f** and **Supplementary Table 3**). In these tumors, the vast majority (>96%) of variants
were located at non-coding regions, with 85% of the identified variants found to populate
intronic and intergenic regions. Considerably, less variants (mean 9%, range 8.1-10.3) were
located proximally upstream of a coding gene, potentially representing promoter regions
(**Fig. 3g**). To assess WGS data quality and explore additional genomic features of our
samples, we investigated mutational signatures, counts, and frequently-mutated genes (**Fig.**
3h-i). Four samples were hypermutated and displayed 20-67% of variants associated with
defective DNA mismatch repair (SBS15). Six samples from tumors pre-treated with
carboplatin showed traces of signatures 31 and 35, both associated with platinum
treatment[27]. Cumulatively, these results confirm previously-described enrichment of tumor-
intrinsic[28,29] and treatment-induced[27] mutational features of endometrial cancer. The most-
frequently mutated genes with protein sequence altering mutations included *ARID1A* and
various members of the PIK3CA pathway, such as *PTEN*, *PIK3CA* and *PIK3R1*,
recapitulating previous observations for endometrial tumor[28].

As expected, analogously to the primary tumor analyses (**Fig. 3a**), somatic mutations
in metastatic samples were also predominantly occurring in non-coding regions. Next, we
integrated these somatic mutation data with tissues-type enriched ERα ChIP-seq data (**Fig.**
3j), and again found differentially bound ERα regions were significantly enriched for somatic
mutations (11.9%, 1,221/10,292), relative to the total number of ERα-bound sites from our
entire cohort, not found to differ between normal tissue and tumors (8%, 5,881/73,312;
Fisher's exact $P = 5.64 \times 10^{-36}$, OR 1.54, 95% CI 1.44-1.65). All samples contributed to this
enrichment, with a median somatic variant count of 18 (range 1-358) in differentially ERα-
bound sites. Moreover, comparing tumor-enriched with tumor-depleted regions, we observed
a significant enrichment of somatic variants in the tumor-enriched ERα sites compared to
tumor-depleted ones (14.5% vs 10.1%, Fisher's exact $P = 2.98 \times 10^{-11}$, OR 1.51, 95% CI
1.34-1.71, **Fig. 3j**). In contrast to primary tumors, these observations are independent of the
mutational frequency status (**Supplementary Fig. 6b**). In most instances (n=1,002), regions
overlapped with only one somatic variant, while some ERα-bound regions were observed to
harbor 2 (183 regions), 3 (34 regions) or 4 (2 regions) somatic mutations.

To investigate genes whose expression may be affected by enhancer mutations, we
intersected tumor-depleted and tumor-enriched ERα binding sites (FDR ≤ 0.01) bearing

mutations with H3K27ac HiChIP chromatin looping data in Ishikawa cells (**Supplementary**
Table 4), aimed to couple putative enhancers with the transcription start sites (TSS) they
control, yielding a total of 331 genes. Out of these, 13 genes were differentially expressed in
Ishikawa endometrial cancer cells upon 8h of estradiol-stimulation[30] (**Fig. 3k**). Interestingly,
ESR1 (encoding ERα) was represented as well, being modestly downregulated at both RNA
and protein level (**Fig. 3k-l** and **Supplementary Fig. 7c**), and harboring 3 metastatic
somatic mutations in two *cis* regulatory elements upstream of the *ESR1* locus, hereafter
referred as '*Enhancer 1*' (P17 and P22; non-hypermutated) and '*Enhancer 2*' (P03b;
hypermutated) (**Fig. 4a** and **Supplementary Fig. 7b**). Of note, high *ESR1* expression is
associated with favorable outcome in primary endometrial cancer patients (TCGA[25]) (**Fig.**
3m)**.** Of note, these two enhancers are specific to the endometrium cancer samples, relative
to healthy endometrial tissues and breast cancer specimens, based on ATAC-seq (TCGA[31])
and ERα ChIP-seq signal (our previous work[32]) (**Supplementary Fig. 7b-c**) at the *ESR1*
locus.

ESR1 enhancer mutation alters 3D genome contacts and decreases
EHMT2/G9a enhancer binding to boost ERα expression

Since *ESR1* expression is strongly associated with favorable outcome in endometrial
cancer (**Fig. 3h**), alterations in its transcriptional regulation may have direct implications on
tumor development and progression. Since both *Enhancer 1* and *Enhancer 2* – found
mutated in metastatic, but not primary endometrial cancer – showed induction of ERα
binding and H3K27ac signals in tumors (**Fig. 4a**), we investigated whether these loci may
undergo alternative regulation in different stages of tumor progression leading to *ESR1*
expression deregulation. Of note, in both our (**Supplementary Fig. 5a**) and TNM-plot[33]
(**Supplementary Fig. 5b**) RNA-seq expression data, *ESR1* expression in primary tumors
was comparable to the levels found in normal tissue, suggesting that altered regulatory
potential alone at these enhancers was not sufficient to affect *ESR1* expression.

Hi-C analyses of endometrial tumors (**Fig. 4b**) illustrated increased short-range and
decreased long-range chromatin interactions at this locus, when compared to healthy
endometrial tissue. These data were confirmed with 4C-seq analyses using the *ESR1*
promoter as view-point (**Fig. 4c**), showing tumor-specific gained interactions of newly ERα-
engaged enhancers with the *ESR1* promoter (**Supplementary Fig. 3i**).

To identify which *ESR1* enhancer mutation had an higher probability to impact *ESR1*
gene regulation and potentially drive tumor progression, we employed the *Akita*[34] machine
learning tool to predict the effect of the somatic mutations on the 3D genome organization

(**Fig. 4d**). Our analyses predicted the mutation occurring at position chr6:152,002,679 (TTC-
to-T, P22) of *Enhancer 1* to affect chromatin architecture surrounding the *ESR1* locus.
Based on these results, this specific mutation was selected for further analyses.

To determine whether enhancer mutations can alter the composition of the
transcription complex recruitment, we performed DNA affinity purification experiments
coupled to quantitative mass spectrometry. To this end, we incubated nuclear extracts from
Ishikawa endometrial cancer cells with immobilized DNA-oligonucleotides encompassing the
center of the ERα-bound site at the *Enhancer 1* – containing either the reference or mutant
sequence – with a ±25bp window around it (**Fig. 5a-b**). Quantitative proteomics identified 13
gained and 11 lost proteins upon mutation of the regulatory element, respectively (**Fig. 5b**).
To identify differentially bound proteins that may directly regulate *ESR1* expression, we used
publicly available RNA-seq (TCGA[25]) data of 589 endometrial cancer tissues to correlate the
expression of *ESR1* with the expression of genes corresponding to the 24 differentially-
bound proteins (**Fig. 5c**). Then, we selected the genes that most-strongly positively (*POLK*,
ZBTB21, *XPA*, *SIN3A*) or negatively (*GTF2IRD1*, *EHMT2*, *ZNF768*) correlated with *ESR1*
expression (**Supplementary Fig. 8a**) For these genes, expression was analyzed in relation
to the progression-free survival probability of endometrial cancer patients (TCGA[25]) (**Fig. 5d**
and **Supplementary Fig. 8b**). Interestingly, our analyses identified only one of the 7 genes
analyzed – the lysine methyl-transferase EHMT2 (also known as G9a or KMT1C) – to
significantly associate with a poor outcome of endometrial cancer patients (**Fig. 5d** and
Supplementary Fig. 8b).

EHMT2/G9a interacts with ERα (**Fig. 5e** and **Supplementary Fig. 8c**), based on co-
immunoprecipitation experiments. In line with this, ChIP-qPCR revealed that EHMT2/G9a
was recruited to both *Enhancer 1* and promoter of the *ESR1* gene in Ishikawa cells,
following estradiol treatment (**Fig. 5f**). As negative control, these experiments were
reperformed in the ERα-negative endometrial cancer cell line AN3CA (**Supplementary Fig.**
9a-b). In line with the absence of ERα expression, estradiol treatment did not alter tumor cell
growth (**Supplementary Fig. 9c-d**). In absence of ERα, there is little recruitment of
EHMT2/G9a to the *ESR1 Enhancer1* and promoter when compared to the negative control
region (**Supplementary Fig. 9e**). Furthermore, siRNA-mediated knock-down of EHMT2/G9a
in Ishikawa cells, resulted in an increase of ERα protein expression (**Fig. 5g** and
Supplementary Fig. 8d), confirming the reciprocal relationship between these proteins.
Interestingly, we observed a slight decrease in EHTM2/G9a protein levels upon estradiol
stimulation, an effect that was abolished when ERα was targeted for degradation[35] by ICI
182,780 (Fulvestrant) treatment (**Fig. 5g** and **Supplementary Fig. 8d**). As negative control,
comparable experiments performed in ERα-negative AN3CA cells, which did not show any
gain of ERα protein levels upon EHMT2/G9a knock-down (**Supplementary Fig. 9f-g**).

Finally, we wondered whether the ERα-dependent chromatin recruitment of
EHMT2/G9a is a specific feature of *ESR1* locus, or rather represents a more general feature
of EHMT2/G9a action. To address this question, we studied the EHMT2/G9a ChIP-seq
signal around the different categories of ERα binding sites in the presence or absence of
estradiol stimulation in Ishikawa cells (**Fig. 5h**). These analyses revealed an absence of
EHMT2/G9a signal at tumor-depleted and common regions, with no differences in its
recruitment upon estradiol stimulation. In contrast, EHMT2/G9a co-localizes with ERα at
tumor-enriched binding sites with a significantly increased signal upon estradiol stimulation
(**Fig. 5h**, *right*), suggesting a more general behavior of EHMT2/G9a for tumor-gained ERα
sites.
Altogether, these data show that EHMT2/G9a DNA binding is impaired upon
mutation of the *ESR1 Enhancer 1* locus, thereby enhancing expression of tumor driver ERα
in metastatic endometrial cancer and that its activity is required to repress *ESR1* expression
in endometrial cancer in a ERα-dependent manner.

DISCUSSION

High expression levels of *ESR1* (encoding ERα) in endometrial cancer are associated with a favorable outcome (**Fig. 3m**), since *ESR1*-positive tumors typically proliferate slower as compared to *ESR1*-negative tumors. Nonetheless, for these tumors, ERα plays a critical role in endometrial cancer development and progression[5], the molecular actions of ERα in this cancer type remain largely unknown.

Though multi-omics analyses (**Fig. 1a**), we investigated how ERα activity is altered in endometrial cancer and how this feature may be leveraged by tumor cells to progress into the metastatic stage. Cumulatively, our findings allowed us to propose a molecular model describing the dynamic nature of ERα action in endometrial cancer development and progression (**Fig. 5i**): i) in healthy endometrial tissue, ERα regulates genes involved in cell cycle and DNA damage response (**Supplementary Fig. 1g** and **Supplementary Fig. 2c**); ii) during tumorigenesis, ERα re-localizes to the tumor-specific enhancers (**Fig. 1f-g**), including the *ESR1* enhancer itself (**Fig. 4c**). At this enhancer ERα interacts with EHMT2/G9a (**Fig. 5e**) in estradiol depend manner (**Fig. 5f**); iii) during tumor progression, the *ESR1* enhancer acquires a somatic mutation for a subset of tumors (**Fig. 4a**). *In vitro* analyses revealed that this mutation impairs EHMT2/G9a DNA binding (**Fig. 5b**), and perturbation of EHMT2/G9a in Ishikawa endometrial cancer cells enhanced *ESR1* expression (**Fig. 5g**). Given the low frequency of SNVs at ERα binding sites, many other paths to metastasis may exist, next to the mechanism we propose in this study.

Our analysis of the ERα cistrome in both normal and neoplastic endometrial tissues revealed limited overlap of chromatin occupancy between the two tissue states. Importantly, despite the healthy and tumor tissues were not matched for the same individual patient, large scale programmatic changes were consistently observed on chromatin binding of ERα, in comparing the two tissue types. Performing these analyses in matched healthy and tumor samples might lead to the identification of a higher number of significantly differentially bound sites, as noise and inter-sample variation is likely lower. This notion implies that our findings may reflect the strongest effects induced by ERα reprogramming, that are to be confirmed in larger future studies. Interestingly, tumor-depleted ERα binding sites did not display differences in H3K27ac, in contrast to the tumor-enriched ones which instead showed a consistent increase in this active histone mark. These results highlighted a specific gain-of-function for newly-engaged ERα-bound *cis*-regulatory elements in tumors, while the regulatory elements for which ERα binding is lost remain active through action of other transcription factors. Surprisingly, studying the genomic distribution of the ERα binding sites, we observed an unexpected enrichment of promoter regions for the tumor-depleted binding sites over the tumor-enriched ones. This promoter-enrichment is in contrast to prior

studies, that reported ERα to primarily occupy promoter-distal *cis*-regulatory elements rather
than promoters[10,14,36,37]. Importantly, as the molecular biology of this nuclear receptor has
been studied mainly using cancer models, the non-tumor context of ERα action remains less
well known, and may involve a higher degree of promoter involvement.

Our data, combined with previous findings showing that cell-specific ERα sites lack
high-affinity estrogen responsive elements (EREs)[11], suggest that tumor-depleted binding
sites (enriched for ERα/β motifs) are essential for the cell maintenance in normal tissues.
However, during tumorigenesis these sites progressively lose ERα binding in favor of sites
enriched in tumors (low-affinity EREs), likely driving cell transformation. Interestingly, at
these tumor-enriched binding sites we found an overrepresentation of SOX and FOX
transcription factor family motifs. These finding are particularly relevant for the endometrial
cancer biology since one of the most recurrently mutated genes in this cancer type is
SOX17[28], which acts as tumor suppressor[38]. Therefore, we can hypothesize that *SOX17*
loss-of-function may leave low-affinity ERα binding sites more accessible for ERα,
increasing receptor occupancy at these regions. On the other hand, enrichment of motifs for
forkhead pioneer transcription factor class (FOX family) at tumor-enriched ERα binding sites
is consistent with the increased H3K27ac signal at these low-affinity ERα binding sites.
FOXA1 is known to facilitate ERα binding through its pioneer activity in breast cancer[39], and
we previously proposed FOXA1 to serve a comparable role in endometrial cancer[10]. In
support of these hypothesis, tumor-enriched ERα binding sites display an enrichment in
FOXA1 ChIP-seq peaks detected in Ishikawa endometrial cancer cells (**Fig. 1k-l** and
Supplementary Fig. 1j). Furthermore, one can speculate on this mechanism to represent a
more general feature of hormone-dependent cancers. Indeed, comparison between healthy
prostate epithelium and prostate adenocarcinoma – being strongly dependent on Androgen
Receptor (AR) action – showed tumor-gained AR binding sites to be selectively enriched for
FOXA1 motifs, as opposed to AR sites found exclusively in healthy tissues[16].

Enhancers interact with promoter regions to regulate gene expression, facilitated
through spatial rearrangement of the chromatin structure[21]. Performing Hi-C analyses in
healthy and tumor tissues, we observed a global enrichment of short-range (<2Mb)
chromatin interactions, at the expense of long-range (>2Mb) contacts. Moreover, chromatin
loop distance in tumor tissues is reduced at tumor-enriched ERα binding sites when
compared to normal samples, while loop distance at shared and tumor-depleted sites
remains unchanged. This corroborates with the observed newly acquired H3K27ac signal at
tumor-enriched ERα binding sites in tumor samples in contrast to common and tumor-
depleted sites, where this active enhancer/promoter histone mark is persistent in both tissue
types. Combined, these findings imply a gain of enhancer activity at tumor-enriched binding

sites that may lead to an increase in enhancer-promoter proximal contacts, explaining the
loop distance shortening.

Next to the short/long range interaction alterations between heathy and tumor tissues, we
also observed a compartment depolarization in tumor samples. Chromatin compartment
destabilization is observed in many tumor types[40] (e.g., colorectal cancer[41] and
glioblastoma[42]). Interestingly, as tumor-enriched ERα binding sites were enriched at B-to-A
compartment switched sub-regions, these data suggest that alterations in 3D genome
organization directly contribute to enhancer plasticity in tumorigenesis, concordant with
increased H3K27ac signal at these sites.

As the vast majority of mutations are found at non-coding regions of the genome, we
investigated whether tumor-enriched ERα sites could drive tumor progression through
acquisition of mutations. Through integration of somatic mutation data with RNA-seq in
Ishikawa cells, we concluded that most genes with proximal ERα-bound enhancer mutations
are not under direct ERα control. This is in agreement with our previous observation in
prostate cancer, in which only a minor fraction of SNVs at regulatory elements functionally
affected their activity[17].

We also observed a depletion of somatic mutations in tumor-depleted ERα binding sites,
relative to the common and tumor-enriched ERα sites (**Fig. 3d,j**). We could speculate that
two mechanisms may underlie this observation. The first is based on previous reports
showing that, since DNA-bound proteins prevent efficient DNA repair[43], somatic mutations
preferentially accumulate at active TF-binding sites. On the other hand, accumulation of
functional mutations may occur at tumor-enriched sites by other mechanisms. In our
previous work in prostate cancer[17], we analogously observed that tumor-gained AR binding
sites were more often somatically mutated, but were also more-often positive for cancer risk
SNPs. These results put our current study in perspective, suggesting that the observed
higher frequency of SNVs in tumor-gained regulatory elements for hormone receptors may
be a feature that is common to hormone-dependent cancers.

While the interaction between ERα and EHMT2/G9a has already been reported in
breast[44,45], EHMT2/G9a is considered to be an ERα co-activator in that cellular context.
However, studies in eryhtroleukemia[46,47] describe a dual role of EHMT2/G9a as both gene
activator and repressor depending on its interaction with Mediator rather than JARID1,
respectively. As we observed ERα expression levels being upregulated following knockdown
of EHMT2/G9a (**Fig. 5g**), an ERα co-repressive role for EHMT2/G9a in the endometrial
cancer cell context is supported by our data. Due to the lack of other ERα-positive
endometrial cancer cell line availability[48], only Ishikawa cells were used in this study.
Therefore, despite experiments carried out in AN3CA ERα-negative cells support our
findings, further investigation would be required to further substantiate what determines the

repressive transcriptional effect of EHMT2/G9a in the interplay with ERα in this tumor type.

Moreover, despite many data streams were generated in tumor tissues, EHMT2/G9a

functional experiments performed in Ishikawa cells – deriving from endometrial epithelial

cells – places our conclusion in an intrinsically tumor cell-centric context that ignores any

potential influence of the microenvironment (e.g., stromal and immunity cells). Further

studies would be required to determine whether the endometrial cancer stromal

compartment may influence ERα regulation in the tumor itself.

In this study we shed light on the molecular mechanisms underlying endometrial

cancer development and progression, connecting alterations in 3D genome organization

with epigenetic plasticity and non-coding somatic mutations, using ESR1 enhancer mutation

as an example. This study may serve as blueprint for studies in other hormone-driven

cancer types, in which such complex connections are yet to be identified.

METHODS

Human tissue collection and quality assessment

Fresh-frozen healthy and tumor samples were obtained from post-operative tissue at the Netherlands Cancer Institute (Amsterdam, the Netherlands) from patients who did not receive neoadjuvant endocrine treatment. Tumor content was assessed by haematoxylin and eosin (H&E) staining on slides taken throughout the tissue sample. For tumors, only samples displaying a tumor percentage greater or equal to 70% were deemed eligible for further analysis. Of note, non-cancerous healthy tissues were obtained from patients whose uterus was removed because of: (a) surgery for cervical carcinoma or ovarian cancer; (b) endometrial cancer diagnoses, but the area that was cryo-preserved was devoid of tumor cells.

The study was approved by the institutional review board of the Netherlands Cancer Institute, written informed consent was signed by all participants enrolled in the study, and all research was carried out in accordance with relevant guidelines and regulations.

Cell culture and chemicals

Ishikawa (Merck Sigma Aldrich) cells were cultured in Dulbecco's Modified Eagle Medium (DMEM, Gibco) and DMEM Mixture F-12 (DMEM/F-12, Gibco), respectively, supplemented with 10% Fetal Bovine Serum (FBS-12A, Capricorn Scientific) and penicillin/streptomycin (100µg/mL, Gibco). AN3CA endometrial cancer cells (ATCC) were cultured in Dulbecco's Modified Eagle Medium (DMEM, Gibco) supplemented with 10% Fetal Bovine Serum (FBS-12A, Capricorn Scientific) and penicillin/streptomycin (100µg/mL, Gibco). Cell lines were subjected to regular *Mycoplasma* testing, and underwent authentication by short tandem repeat profiling (Eurofins Genomics).

For hormone stimulation, cells were pre-cultured for 3 days in phenol red-free DMEM (Gibco, Ishikawa) or DMEM/F-12 (Gibco, AN3CA) supplied with 5% dextran-coated charcoal (DCC) stripped FBS, 2mM L-glutamine (Gibco), and penicillin/streptomycin (100µg/mL, Gibco), then stimulated with 10nM DMSO-solubilized 17β-estradiol (MedChemExpress, #HY-B0141) for the indicated amount of time. Inhibition of ERα activity was performed by treatment of the cells with 100nM DMSO-solubilized Fulvestrant/ICI-182,780 (MedChemExpress, #HY-13636).

Cell proliferation analyses

Cells were plated in a 96-well plate at a density of 10,000 cells/well. Cells were treated with 100nM DMSO-solubilized Fulvestrant/ICI-182,780 (MedChemExpress, #HY-13636). Cells were imaged every 4 hours by using an IncuCyte ZOOM Live-Cell Analysis System (Essen

551 BioScience, Sartorious), and cell confluency percentage was calculated using the
552 IncucyteZoom (v2018A) software. Normalization, by subtraction of the first point (T_0), and
553 visualization were performed using *Rseb*[49] *(v0.3.2)* R-package (https://github.com/sebastian-
554 gregoricchio/Rseb).

ChIP-qPCR and ChIP-seq library preparation
Chromatin Immunoprecipitation of ERα (SantaCruz #sc-543, 5μg/IP) and H3K27ac
(ActiveMotif #39133, 5μg/IP) were performed as previously described[32] employing the
combination of DSG and Formaldehyde to perform the crosslinking. The same protocol, but
using 1% formaldehyde for the crosslinking, have been used to perform EHMT2/G9a
(Abcam #ab133482, 5μg/IP) and Normal-IgG (Merck Millipore #12-370, same amount than
IP) ChIPs in Ishikawa cells.
Immunoprecipitated DNA is compared to input sample by the comparative C_t method, and
signal enrichment is expressed as percentage of input (% Input). Real-Time quantitative
PCR (RT-qPCR) was performed on the QuantStudio™ 5 Real-Time PCR System 384-well
(Applied Biosystems, #A28140) using SenSMix™ SYBR® No-ROX buffer (Meridian
Bioscience). The following genomic locations have been tested: *ESR1_Enhancer1*-125kb
(Fw: 5'-TGGTAGGTGCTCAGGAGATAA-3', Rv: 5'-CAGCGACTCGAACAGGATTT-3'),
*ESR1_*promoter-0.9kb (Fw: 5'-CCACTCCTGGCATTGTGATTA-3', Rv: 5'-
CAGGACACATGACACCCAAT-3'), *CDK12*_TSS (Fw: 5'-GGACCTGATCTCGCGTTGTT-3',
Rv: 5'-TAGCCTCTCGCGATGTTTCG-3').

ChIP-seq data processing, motif and Gene Ontology enrichment analyses
Reads were aligned to the human genome build GRCh37 using *BWA* 0.5.9-r26-dev[50].
Reads with a mapping quality (MAPQ) < 20 were removed from further analysis, and
duplicates were marked using GATK markDuplicates[51]. Enrichment over input control was
determined using both *MACS2*[52] (*q* < 0.01) and *DFilter*[53]. Only peaks identified by both
methods, and not overlapping with the ENCODE GRCh37/Hg19 blacklisted regions[54], were
retained. ERα consensus peaks for a given condition (healthy or normal tissues) have been
defined as peaks found in at least 75% of the samples.
Differentially bound regions between tumor and healthy samples were identified using
DiffBind[19] with *EdgeR*[55] mode at an FDR < 0.05. RPGC-normalized (Reads Per Genomic
Content-1× coverage: (mapped reads × fragment length) / effective genome size;
bamCoverage[56]) ChIP-seq signal at peaks was visualized using *deepTools*[56], while genomic
tracks and average density profiles were generated with *Rseb*[49] *(v0.3.2)* R-package
(https://github.com/sebastian-gregoricchio/Rseb) or *pyGenomeTracks*[57,58].

Genomic feature annotations and distance to TSS were performed using *ChIPseeker*[59]

(promoter: -2kb:TSS:+1kb, flanking distance: 2kb), while for motif enrichment in peak

regions we used *AME* v5.5.05[60] algorithm from the *MEME* suite[61] (v.5.5.5) using the

JASPAR database[62,63] as reference. Visualization of the enrichment motifs was made using

ggwordcloud R-package using the *AME* computed *E*-value.

Area proportional Venn diagrams were created using the *Vennerable* R package.

Analyses for Gene Ontology biological process (GO-BP) enrichments for tumor-depleted

ERα promoter-bound genes were performed using DAVID[64] (v6.8) and employing tumor-

enriched ERα promoter-bound genes as background data set. Unsupervised clustering

permutation test was performed using *ConsensusClusterPlus*[65] (v1.54.0) on the *DiffBind*

normalized counts using the following parameters: reps = 100 (number of permutations),

pItem = 0.8 (80% of sample shuffling), pFeature = 1 (100% of features to sample), clusterAlg

= "hc" (hclust, hierarchical clustering), distance = "spearman". To determine the ESR1/ERα

motif strength at each individual ERα binding site, fasta-formatted sequences of the peaks

was obtained using BED2FASTA from the *MEME* suite[61] (v.5.5.5). Then, *PWMScore*, from

PWMTools[66], was used to compute the ESR1/ERα motif (JASPAR id: MA0112.3) sum

occupancy score for each site. *GIGGLE*[20] analyses were performed using the Cistrome Data

Browser Toolkit (http://dbtoolkit.cistrome.org/).

Hi-C library preparation and data processing

Flash-frozen primary tissues have been processed as described in **Fig. 2a**. Briefly, 30µm

slices are cross-linked, washed, electronically homogenized and filtered using a 75µm cell

strainer and collected in a 1.5mL microcentrifuge loBind tube. Then, Hi-C single-index library

preparation was performed as previously described[67] using MboI (New England Biolabs)

restriction enzyme.

Quality and quantification of the Hi-C libraries was assessed using the 2100 Bioanalyzer

(Agilent, DNA 7500 kit). An equimolar pool of the different samples was sequenced on the

Illumina NextSeq 550 System in a 75bp paired-end setup. De-multiplexed fastq data were

analyzed at 2kb, 10kb, 40kb, 100kb and 500kb resolution using the *snHiC*[68] (v0.1.1) pipeline

(https://github.com/sebastian-gregoricchio/snHiC) using default parameters and the

Hg19/GRCh37 genome assembly. This pipeline relies on *HiCExplorer*[69,70] (v3.7.2) for the

matrix generation, normalization and correction, TAD and loop detection, and long/short

contacts ratio and quality controls. Compartments are identified by *dcHiC*[71] *(v2.1).*

Downstream analyses have been performed using *GENOVA*[72] (v1.0.1), which was used to

plot Hi-C contact matrices heatmaps, compute Relative Contact Probabilities, and perform

paired-end spatial chromatin analyses (*PE-SCAn*[23]) at ERα binding sites. *HiCExplorer*[69,70]

(v3.7.2) *hicCompartmentalization* was used to quantify the compartment polarization. For the

compartment switching computation, A and B compartments locations in the reference
condition were obtained merging adjacent bins displaying positive or negative, respectively,
compartmentalization scores. Then, average compartmentalization score in the two
conditions was computed for each reference compartment. A compartment in the reference
condition was defined as "switching" when the average compartmentalization score in the
reference condition switched sign in the test condition. Compartment-ChIP peaks overlaps
and relative upset plot have been computed using *intervene*[73], while Hi-C tracks were
generated by *pyGenomeTracks*[57,58].

Whole Genome Sequencing (WGS) analyses
Primary samples. From the full Uterine Corpus Endometrial Carcinoma (UCEC) cohort
available in the TCGA[25] database, 41 samples have been selected in order to match the
following criteria: gender = "FEMALE", histological_type = "Endometrioid endometrial
adenocarcinoma", history_of_neoadjuvant_treatment = "No", horm_ther = "No, I have never
taken menopausal hormone therapy.", menopause_status = "Post (prior bilateral
ovariectomy OR >12 mo since LMP with no prior hysterectomy)",
prior_tamoxifen_administered_usage_category = "Never Used", radiation_therapy = "NO",
sample_type = "Primary Tumor", targeted_molecular_therapy = "NO", tumor_tissue_site =
"Endometrial", *ESR1* TPM-normalized gene counts ≥ 5. Somatic mutation calling was
performed on base-quality recalibrated aligned Whole Genome Sequencing data available
under restricted access on the TCGA portal (data access authorization, project ID: 36269),
using blood-derived matched normal samples as germline reference, by *GATK* (v 4.3.0.0)
Mutec2[51]. Only mutations passing the GATK *FilterMutectCalls* filter and displaying a
sequencing depth greater than 20 (DP>0) were retained.
Metastatic samples. Sampling, sequencing and initial computational steps have been
previously described extensively by Priestley *et al.*[74] and performed at the central
sequencing facility at the Hartwig Medical Foundation.
Frequently mutated genes were determined by overlapping genes with somatic mutations
affecting protein coding regions in more than one sample in our dataset, with the mutated
genes described in the endometrial carcinoma publication from TCGA[25] and the OncoKB
Cancer Gene List[75]. Overlap between somatic variants and ChIP-seq regions was
determined using *bedtools intersect* (v2.25.0)[76].
For both datasets, mutational signatures were determined using the *deconstructSigs*[77] R-
package, using the COSMIC mutational signatures v3 database[78]. Genomic feature
annotation was performed using *ChIPseeker*[59] (promoter: -2kb:TSS:+1kb, flanking distance:
2kb).
660

Risk Single Nucleotide Polymorphism (rSNP) enrichment analyses

We assessed the enrichment of endometrial cancer germline risk variants among differentially bound ERα sites using the *VSE*[79] (v0.99) R-package. 27 endometrial cancer risk loci (lead variant $P < 1 \times 10^{-5}$) were identified from a genome-wide association study of endometrial cancer[26]. Variants in strong LD ($r^2 > 0.8$) with the lead SNP at each locus were selected using 1000 Genomes Project phase 3[80] to generate an associated variant set (AVS). A null-distribution was built on the basis of 500 matched random variant sets. We assessed enrichment within tumor-enriched and tumor-depleted ERα binding sites, with and without including a 500bp window surrounding each binding site. A Bonferroni-corrected *P*-value < 0.05 (adjusting for four tests) was considered statistically significant.

Endometrial human tissue derived RNA-seq library preparation and differential expression analyses

RNA from ~30mg of endometrial healthy or tumor tissue was extracted using the RNeasy mini kit (Qiagen) following manufacture instructions. Quality of the RNA extraction was assessed using the 2100 Bioanalyzer (Agilent, RNA 6000 Nano Kit) and selecting samples with a RNA Integrity Number (RIN) above 9. PolyA+ stranded RNA library was prepared using the Illumina Stranded mRNA Prep kit and quality was assessed by using the 2100 Bioanalyzer (Agilent, DNA 7500 kit). RNA-seq libraries have been pooled equimolarly and sequenced using NovaSeq6000 (Illumina) sequencer with a 51bp Paired-End reads setup. Fastq files have been demultiplexed by Cutadapt[81] and mapped on Hg38/GRCh38 genome assembly using HISAT2[82] (v2.1.0) using the following parameters: --wrapper basic-0 --min-intronlen 20 --max-intronlen 500000 --rna-strandness FR -k 5 --minins 0 --maxins 500 --fr --new-summary. HISAT2[82] (v2.1.0) was used to generate raw gene counts, gene counts normalization and differential expression analyses were performed using *DESeq2*[83] (v1.30.1). Differentially expressed genes were defined by |Fold Change Expression| ≥ 2 & $P_{adj} < 0.05$. Data were visualized using *Rseb*[49] (v0.3.2) (https://github.com/sebastian-gregoricchio/Rseb).

Gene Set Enrichment Analyses (GSEA)

Differentially expressed genes were ranked by the $\log_2$(Fold Change expression) calculated using *DESeq2*[83] (v1.30.1) and used to perform GSEA analyses by *clusterProfiler*[84] (v3.18.1) on the "Hallmarks" (H) datasets retrieved through *msigdbr* (v7.5.1). Visualization of the results was performed using *Rseb*[49] (v0.3.2) (https://github.com/sebastian-gregoricchio/Rseb).

Public RNA-seq and differential gene expression analyses on Ishikawa cells

698 Ishikawa gene expression data was obtained from GEO, accession numbers GSE109892[30].

699 Differential gene expression between estradiol and DMSO treated cells was computed using

700 *DESeq2*[83] (v1.30.1). Differentially expressed genes were defined by |Fold Change

701 Expression| ≥ 1.5 & P_{adj} < 0.05.

Circularized Chromosome Conformation Capture sequencing (4C-seq)

For each sample, 10-15 30µM-thick slices of flash frozen tissue were cross-linked under

rotation at room temperature for 20 minutes in 5mL of 1× PBS supplemented with 2% FBS

and 2% Methanol-free Formaldehyde. The cross-linked slices have been washed twice in

10mL of ice-cold 1× PBS before to be resuspended in 0.5-1mL of ice-cold 1× PBS in 1.5mL

DNA-loBind microcentrifuge tubes. Tissues have been electronically homogenized and

lysates filtered using a 75µm cell strainer. Then, experiments of 4C-seq were performed as

previously described[85]. Briefly, nuclei were isolated and permeabilized to allow digestion of

the chromatin by the primary RE (DpnII, New England Biolabs). Upon dilution, chromatin

fragments have been ligated and then de-crosslinked. DNA was purified and digested with

the secondary RE (NlaIII, New England Biolabs) and circularized by ligation. PCR

amplification of the re-purified circular fragments have been performed using View-Point

specific primers (Reading Primer: 5'-tacacgacgctcttccgatctAACTCGATTTGGAGCGATC-3';

Non-reading Primer: 5'-actggagttcagacgtgtgctcttccgatctCTGGGACTGCACTTGCTC-3')

using the Expand Long Template PCR System (Roche). Amplicons were purified with

AMPure XP beads (Beckman Coulter) in a 0.8× ratio and then amplified using standard

indexed Illumina primers as previously described[85] using the Expand Long Template PCR

System (Roche). Second-round PCR products were purified with PCR purification columns

(Qiagen) and quantified by 2100 Bioanalyzer (Agilent, DNA 7500 kit).

4C libraries have been pooled equimolarly and sequenced using Illumina MiSeq with a 75bp

Single-End reads setup. Fastq files have been demultiplexed by Cutadapt[81] and mapped on

Hg19/GRCh37 genome assembly and signal normalized by *pipe4C*[85] v1.1 R-package in "cis"

mode and with default parameters. The signal of different technical replicates was averaged

to obtain a unique signal for each sample. The, each tissue sample have been considered

as biological replicate. Downstream analyses of 4C-seq data have been performed in an R

v4.0.3 environment by using *get.single.base.score.bw* and *genomic.track* functions from

Rseb[49] (v0.3.2) (https://github.com/sebastian-gregoricchio/Rseb) R-package in combination

with *ggplot2* (v3.3.5) and *ggforce* (v0.3.3) R-packages.

Protein extraction and immunoblotting

For immunoprecipitated proteins, elution buffer (1% SDS, 0.1M NaHCO$_3$) and Laemmli buffer, supplied with 100mM DTT (Dithiothreitol, Merck Sigma Aldrich), were added to input samples and beads, and then boiled at 95°C for 30min. For whole-cell extracts, cells have been lysed using 2× Laemmli buffer supplied with 100mM DTT (Dithiothreitol, Merck Sigma Aldrich), boiled at 95°C for 10min and then DNA was sheared by sonication (Bioruptor® Pico, Diagenode; 4 cycles of 30s ON + 90s OFF). In both cases, proteins were then resolved by SDS-PAGE and transferred to a 0.22μm nitrocellulose membrane (BioRad). Membranes were incubated for 2h with blocking solution (BS: 1× PBS, 0.1% Tween20, 5% non-fat milk powder) and then incubated with primary antibody against ERα (ThermoFisher #MA5-14104, 1:1000 in BS), EHMT2/G9a (Abcam #ab133482, 1:1000 in BS) or GAPDH (ThermoFisher #PA1-987, 1:5000 in BS) overnight at +4°C. After washing with washing solution (1× PBS, 0.1% Tween20), membranes were incubated with secondary antibodies donkey-α-mouse IRDye® 680RD (926-68073, LI-COR Biosciences, 1:10,000 in BS), donkey-α-mouse IRDye® 800CW (926-32212, LI-COR Biosciences, 1:10,000 in BS), donkey-α-rabbit IRDye® 800CW (926-32213, LI-COR Biosciences, 1:10,000 in BS) or donkey-α-rabbit IRDye® 680RD (926-68073, LI-COR Biosciences, 1:10,000 in BS) for 1h. Membranes were washed again with washing solution, scanned and analyzed using an Odyssey® CLx Imaging System (LI-COR Biosciences) and *ImageStudio™ Lite* v.5.2.5 software (LI-COR Biosciences).

Sequence-based Machine Learning 3D genome folding predictions

The 3D genome organization perturbation predictions have been performed using a window of 2^{20}bp around the *ESR1* locus employing the Convolutional Neural Networks (CNN) model from *Akita*[34] tool. We used the pre-existing model based on Hi-C data originated from human foreskin fibroblast (HFF). Heatmaps have been generated using *matplotlib.pyplot.matshow*.

DNA-oligo protein pull-down

Ishikawa cells were harvested and washed twice with ice-cold PBS and nuclear extracts were prepared as described previously[86]. Briefly, cells are washed with 1× PBS and trypsinized. Trypsin is neutralized by adding the appropriate SILAC medium, and "light"/"heavy" cells are collected separately at 4°C. Cells are washed in 1× ice-cold PBS and resuspended in five volumes of ice-cold buffer A (10 mM KCl, 20 mM Hepes KOH pH 7.9, 1.5 mM MgCl$_2$). Cells are then pelleted and resuspended in two volumes of buffer A supplemented with protease inhibitors and 0.15% Igepal NP40 (v/v) before to be dounce homogenized. Nuclei are then collected and resuspended in two volumes of buffer C (420 mM NaCl, 20 mM Hepes KOH pH 7.9, 20% glycerol (v/v), 2 mM MgCl$_2$, 0.2 mM EDTA, and

0.1% Igepal CA630 (NP40; Sigma-Aldrich, I8896), EDTA-free complete protease inhibitors
(Roche), and 0.5 mM DTT) and incubated for 1 hour. The suspension is centrifuged, and the
supernatant containing the nuclear extracts is collected.
The ~50bp oligonucleotide probes encompassing the SNP were ordered with the forward
strand containing a 5'-biotin moiety (Integrated DNA Technologies) (Fw_WT: 5'-
/5Biosg/ACTGTGGAAACTGGAAGCTGTTCTTGGACTATTTCGCAACACTTTTCTCC-3';
Rv_WT: 5'-GGAGAAAAGTGTTGCGAAATAGTCCAAGAACAGCTTCCAGTTTCCACAGT-
3'; Fw_SNP: 5'-
/5Biosg/ACTGTGGAAACTGGAAGCTGTTTGGACTATTTCGCAACACTTTTCTCCTC-3',
Rv_SNP: (5'-GAGGAGAAAAGTGTTGCGAAATAGTCCAAACAGCTTCCAGTTTCCACAGT-
3'; mutated site is underlined). DNA affinity purifications, on-bead trypsin digestion and
dimethyl labeling were performed as described[87]. Matching light and medium labelled
samples were then combined and analyzed using a gradient from 7 to 30% buffer B in Buffer
A over 44min, followed by a further increase to 95% in the next 16min at flow rate of
250nL/min using an Easy-nLC 1000 (Thermo Fisher Scientific) coupled online to an Orbitrap
Exploris 480 (Thermo Fisher Scientific). MS1 spectra were acquired at 120,000 resolution
with a scan range from 350 to 1300 m/z, normalized AGC target of 300% and maximum
injection time of 20 ms. The top 20 most intense ions with a charge state 2-6 from each
MS1 scan were selected for fragmentation by HCD. MS2 resolution was set at 15,000 with a
normalized AGC target of 75%. Raw MS spectra were analyzed using MaxQuant software
(version 1.6.0.1) with standard settings[87,88]. Data was searched against the human UniProt
database (downloaded 2017) using the integrated search engine. N-terminal and lysine
modification for dimethyl labeling was specified under "labels". Carbamidomethylation was
specified as a fixed modification. N-terminal acetylation and methionine oxidation were
selected as variable modifications.
794

Transient small interfering RNA (siRNA) cell transfection
Transient transfections of Ishikawa and AN3CA cell lines was performed according to the
manufacturer's instructions using Lipofectamine™ RNAiMAX (Invitrogen) for siRNA
knockdown experiments. The ON-TARGETplus™ siRNA SMARTpool targeting human
EHMT2/G9a (L-006937-00-00050) and the siGENOME™ Non-Targeting control siRNA #5
(D-001210-05-20) were purchased from Dharmacon.
801

802 **Statistical analysis**
803 All statistical analyses were performed in R versions 3.4.4, 3.5.1 or 4.0.2 (R Core Team
804 2020, https://www.R-project.org). Enrichment of Ishikawa peaks and somatic mutations in

805   differentially bound regions was calculated using Fisher's exact test, and, where appropriate,

806   corrected for multiple testing using the FDR.

DATA AVAILABILITY

Endometrial healthy and tumor tissues ERα and H3K27ac ChIP-seq, RNA-seq, Hi-C and,

4C-seq raw sequencing data (GRCh37/Hg19 genome build) have been deposited at the

European Genome-phenome Archive (EGA), which is hosted by the EBI and the CRG,

under accession number EGAS00001007240, while endometrial tissue processed data and

Ishikawa ChIP-seq data are deposited at the Gene Expression Omnibus (GEO) database

under accession number GSE235241.

Raw sequencing data of the RNA-seq performed in Ishikawa cells can be found at GEO

database under accession number GSE109892[30]. ERα ChIP-seq data in breast cancer

tissues from Singh *et al.*[32] are accessible in GEO under accession number GSE114737.

Ishikawa ChIP-seq data in the form of optimal Irreproducibility Discovery Rate (IDR)

thresholded peaks (GRCh37/Hg19 genome build) were obtained from the ENCODE data

portal[48] (**Supplementary Table 2**). The H3K27ac HiChIP data in Ishikawa cells have been

obtained from O'Mara *et al.*[89] accessible in GEO under accession number GSE137936.

Mass spectrometry data have been deposited at the ProteomeXchange Consortium through

the PRIDE[90] partner repository with the identifier PXD029822. The breast and endometrial

cancer ATAC-seq, gene expression, progression-free survival and Whole Genome

Sequencing (data access authorization for project ID: 36269) used in this this work are in

whole or part based upon data generated by the TCGA Research Network[25,31]

(https://www.cancer.gov/tcga).

ACKNOWLEDGEMENTS

We thank members of the Zwart and Bergman labs for valuable feedback, suggestions and

input. We would like to acknowledge the Research High Performance Computing (RHPC)

facility of the Netherlands Cancer Institute (NKI) to have enabled us to perform all the

computations required to analyze the data generated, the NKI Genomics Core Facility for

next-generation sequencing and bioinformatics support, and the NKI-AVL Core Facility

Molecular Pathology and Biobanking for patient samples handling.

This publication and the underlying study have been made possible partly based on data

that Hartwig Medical Foundation has made available to the study through the Hartwig

Medical Database. Further, we would like to thank Hartwig Medical Foundation for facilitating
and supporting whole-genome sequencing data generation, data analyses and data
interpretation.
843
844

AUTHOR CONTRIBUTIONS

Conceptualization: S.G., A.K and W.Z. W.Z. was responsible for project funding. S.G. and
A.K. performed Hi-C and 4C-seq experiments; S.G., K.S. and M.D. performed ChIP/ChIP-
seq and immunoblotting experiments; S.S. performed and analyzed the protein DNA-oligo
pulldowns experiments; S.G., M.H, T.M.S. and A.A.S. analyzed the omics data; T.A.O. and
D.M.G. provided support in the HiChIP and rSNP data analyses; M.V. provided support for
the proteomics experiments; L.F.A.W. provided bioinformatics support; F.v.L. designed and
performed study for patient sample collection. S.G., A.K. and W.Z. wrote the manuscript,
with input from all co-authors.

854
855

856   ## FUNDING

857   This work was supported by Oncode Institute and *Saxum Volutum.* The Zwart, Wessels and
858   Vermeulen labs are part of the Oncode Institute, which is partly funded by the Dutch Cancer
859   Society. T.O. is supported by a National Health and Medical Research Council of Australia
860   Emerging Leader Investigator Fellowship (GNT1173170).

861
862

863   ## COMPETING INTERESTS

864   The authors declare no competing interests.

REFERENCES

1. Sung, H. *et al.* Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA. Cancer J. Clin.* **71**, 209–249 (2021).

2. Gu, B. *et al.* Variations in incidence and mortality rates of endometrial cancer at the global, regional, and national levels, 1990-2019. *Gynecol. Oncol.* **161**, 573–580 (2021).

3. Morice, P., Leary, A., Creutzberg, C., Abu-Rustum, N. & Darai, E. Endometrial cancer. *Lancet Lond. Engl.* **387**, 1094–1108 (2016).

4. Carlson, M. J., Thiel, K. W. & Leslie, K. K. Past, present, and future of hormonal therapy in recurrent endometrial cancer. *Int. J. Womens Health* **6**, 429–435 (2014).

5. Rodriguez, A. C., Blanchard, Z., Maurer, K. A. & Gertz, J. Estrogen Signaling in Endometrial Cancer: a Key Oncogenic Pathway with Several Open Questions. *Horm. Cancer* **10**, 51–63 (2019).

6. Gunderson, C. C., Fader, A. N., Carson, K. A. & Bristow, R. E. Oncologic and reproductive outcomes with progestin therapy in women with endometrial hyperplasia and grade 1 adenocarcinoma: a systematic review. *Gynecol. Oncol.* **125**, 477–482 (2012).

7. Crosbie, E. J. *et al.* Endometrial cancer. *Lancet Lond. Engl.* **399**, 1412–1428 (2022).

8. Heldring, N. *et al.* Estrogen receptors: how do they signal and what are their targets. *Physiol. Rev.* **87**, 905–931 (2007).

9. Flach, K. D. & Zwart, W. The first decade of estrogen receptor cistromics in breast cancer. *J. Endocrinol.* **229**, R43-56 (2016).

10. Droog, M. *et al.* Comparative Cistromics Reveals Genomic Cross-talk between FOXA1 and ERα in Tamoxifen-Associated Endometrial Carcinomas. *Cancer Res.* **76**, 3773–3784 (2016).

11. Gertz, J. *et al.* Distinct Properties of Cell-Type-Specific and Shared Transcription Factor Binding Sites. *Mol. Cell* **52**, 25–36 (2013).

12. Gertz, J., Reddy, T. E., Varley, K. E., Garabedian, M. J. & Myers, R. M. Genistein and bisphenol A exposure cause estrogen receptor 1 to bind thousands of sites in a cell type-specific manner. *Genome Res.* **22**, 2153–2162 (2012).

13. Leeuwen, F. E. van *et al.* Risk of endometrial cancer after tamoxifen treatment of breast cancer. *The Lancet* **343**, 448–452 (1994).

14. Droog, M. *et al.* Estrogen receptor α wields treatment-specific enhancers between morphologically similar endometrial tumors. *Proc. Natl. Acad. Sci.* **114**, E1316–E1325 (2017).

15. Chi, D. *et al.* Estrogen receptor signaling is reprogrammed during breast tumorigenesis. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 11437–11443 (2019).

16. Pomerantz, M. M. *et al.* The androgen receptor cistrome is extensively reprogrammed in human prostate tumorigenesis. *Nat. Genet.* **47**, 1346–1351 (2015).

17. Mazrooei, P. *et al.* Cistrome Partitioning Reveals Convergence of Somatic Mutations and Risk Variants on Master Transcription Regulators in Primary Prostate Tumors. *Cancer Cell* **36**, 674-689.e6 (2019).

18. Pomerantz, M. M. *et al.* Prostate cancer reactivates developmental epigenomic programs during metastatic progression. *Nat. Genet.* **52**, 790–799 (2020).

19. Ross-Innes, C. S. *et al.* Differential oestrogen receptor binding is associated with clinical outcome in breast cancer. *Nature* **481**, 389–393 (2012).

20. Layer, R. M. *et al.* GIGGLE: a search engine for large-scale integrated genome analysis. *Nat. Methods* **15**, 123–126 (2018).

21. Robson, M. I., Ringel, A. R. & Mundlos, S. Regulatory Landscaping: How Enhancer-Promoter Communication Is Sculpted in 3D. *Mol. Cell* **74**, 1110–1122 (2019).

22. Harewood, L. *et al.* Hi-C as a tool for precise detection and characterisation of chromosomal rearrangements and copy number variation in human tumours. *Genome Biol.* **18**, 125 (2017).

23. de Wit, E. *et al.* The pluripotent genome in three dimensions is shaped around pluripotency factors. *Nature* **501**, 227–231 (2013).

24. Rheinbay, E. *et al.* Analyses of non-coding somatic drivers in 2,658 cancer whole genomes. *Nature* **578**, 102–111 (2020).

25. Cancer Genome Atlas Research Network *et al.* The Cancer Genome Atlas Pan-Cancer analysis project. *Nat.*
Genet. **45**, 1113–1120 (2013).

26. O'Mara, T. A. *et al.* Identification of nine new susceptibility loci for endometrial cancer. *Nat. Commun.* **9**, 3166
(2018).

27. Boot, A. *et al.* Characterization of colibactin-associated mutational signature in an Asian oral squamous cell
carcinoma and in other mucosal tumor types. *Genome Res.* **30**, 803–813 (2020).

28. Cancer Genome Atlas Research Network *et al.* Integrated genomic characterization of endometrial
carcinoma. *Nature* **497**, 67–73 (2013).

29. O'Hara, A. J. & Bell, D. W. The genomics and genetics of endometrial cancer. *Adv. Genomics Genet.* **2012**,
33–47 (2012).

30. Vahrenkamp, J. M. *et al.* Clinical and Genomic Crosstalk between Glucocorticoid Receptor and Estrogen
Receptor α In Endometrial Cancer. *Cell Rep.* **22**, 2995–3005 (2018).

31. Corces, M. R. *et al.* The chromatin accessibility landscape of primary human cancers. *Science* **362**, eaav1898
(2018).

32. Singh, A. A. *et al.* Optimized ChIP-seq method facilitates transcription factor profiling in human tumors. *Life*
Sci. Alliance **2**, (2019).

33. Bartha, Á. & Győrffy, B. TNMplot.com: A Web Tool for the Comparison of Gene Expression in Normal, Tumor
and Metastatic Tissues. *Int. J. Mol. Sci.* **22**, 2622 (2021).

34. Fudenberg, G., Kelley, D. R. & Pollard, K. S. Predicting 3D genome folding from DNA sequence with Akita.
Nat. Methods **17**, 1111–1117 (2020).

35. Zwart, W. *et al.* Resistance to antiestrogen arzoxifene is mediated by overexpression of cyclin D1. *Mol.*
Endocrinol. Baltim. Md **23**, 1335–1345 (2009).

36. Carroll, J. S. *et al.* Genome-wide analysis of estrogen receptor binding sites. *Nat. Genet.* **38**, 1289–1297
(2006).

37. Lin, C.-Y. *et al.* Whole-Genome Cartography of Estrogen Receptor α Binding Sites. *PLOS Genet.* **3**, e87
(2007).

38. Zhang, Y. *et al.* SOX17 is a tumor suppressor in endometrial cancer. *Oncotarget* **7**, 76036–76046 (2016).

39. Hurtado, A., Holmes, K. A., Ross-Innes, C. S., Schmidt, D. & Carroll, J. S. FOXA1 is a critical determinant of
Estrogen Receptor function and endocrine response. *Nat. Genet.* **43**, 27–33 (2011).

40. Yamaguchi, K., Chen, X., Oji, A., Hiratani, I. & Defossez, P.-A. Large-Scale Chromatin Rearrangements in
Cancer. *Cancers* **14**, 2384 (2022).

41. Johnstone, S. E. *et al.* Large-Scale Topological Changes Restrain Malignant Progression in Colorectal
Cancer. *Cell* **182**, 1474-1489.e23 (2020).

42. Yang, Q. *et al.* Alterations in 3D chromatin organization contribute to tumorigenesis of EGFR-amplified
glioblastoma. *Comput. Struct. Biotechnol. J.* **20**, 1967–1978 (2022).

43. Sabarinathan, R., Mularoni, L., Deu-Pons, J., Gonzalez-Perez, A. & López-Bigas, N. Nucleotide excision
repair is impaired by binding of transcription factors to DNA. *Nature* **532**, 264–267 (2016).

44. Purcell, D. J., Jeong, K. W., Bittencourt, D., Gerke, D. S. & Stallcup, M. R. A distinct mechanism for
coactivator versus corepressor function by histone methyltransferase G9a in transcriptional regulation. *J. Biol.*
Chem. **286**, 41963–41971 (2011).

45. Zhang, X. *et al.* G9a-mediated methylation of ERα links the PHF20/MOF histone acetyltransferase complex to
hormonal gene expression. *Nat. Commun.* **7**, 10810 (2016).

46. Chaturvedi, C.-P. *et al.* Dual role for the methyltransferase G9a in the maintenance of beta-globin gene
transcription in adult erythroid cells. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 18303–18308 (2009).

47. Chaturvedi, C.-P. *et al.* Maintenance of gene silencing by the coordinate action of the H3K9 methyltransferase
G9a/KMT1C and the H3K4 demethylase Jarid1a/KDM5A. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 18845–18850
(2012).

48. Korch, C. *et al.* DNA profiling analysis of endometrial and ovarian cell lines reveals misidentification,
redundancy and contamination. *Gynecol. Oncol.* **127**, 241–248 (2012).

49. Gregoricchio, S. *et al.* HDAC1 and PRC2 mediate combinatorial control in SPI1/PU.1-dependent gene
repression in murine erythroleukaemia. *Nucleic Acids Res.* **50**, 7938–7958 (2022).

50. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinforma. Oxf.*
Engl. **25**, 1754–1760 (2009).

51. Cibulskis, K. *et al.* Sensitive detection of somatic point mutations in impure and heterogeneous cancer
samples. *Nat. Biotechnol.* **31**, 213–219 (2013).

52. Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137 (2008).

53. Kumar, V. *et al.* Uniform, optimal signal processing of mapped deep-sequencing data. *Nat. Biotechnol.* **31**,
615–622 (2013).

54. Amemiya, H. M., Kundaje, A. & Boyle, A. P. The ENCODE Blacklist: Identification of Problematic Regions of
the Genome. *Sci. Rep.* **9**, 9354 (2019).

55. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential expression
analysis of digital gene expression data. *Bioinforma. Oxf. Engl.* **26**, 139–140 (2010).

56. Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic*
Acids Res. **44**, W160-165 (2016).

57. Lopez-Delisle, L. *et al.* pyGenomeTracks: reproducible plots for multivariate genomic datasets. *Bioinformatics*
37, 422–423 (2021).

58. Ramírez, F. *et al.* High-resolution TADs reveal DNA sequences underlying genome organization in flies. *Nat.*
Commun. **9**, 189 (2018).

59. Yu, G., Wang, L.-G. & He, Q.-Y. ChIPseeker: an R/Bioconductor package for ChIP peak annotation,
comparison and visualization. *Bioinformatics* **31**, 2382–2383 (2015).

60. McLeay, R. C. & Bailey, T. L. Motif Enrichment Analysis: a unified framework and an evaluation on ChIP data.
BMC Bioinformatics **11**, 165 (2010).

61. Bailey, T. L., Johnson, J., Grant, C. E. & Noble, W. S. The MEME Suite. *Nucleic Acids Res.* **43**, W39-49
(2015).

62. Castro-Mondragon, J. A. *et al.* JASPAR 2022: the 9th release of the open-access database of transcription
factor binding profiles. *Nucleic Acids Res.* **50**, D165–D173 (2022).

63. Sandelin, A., Alkema, W., Engström, P., Wasserman, W. W. & Lenhard, B. JASPAR: an open-access
database for eukaryotic transcription factor binding profiles. *Nucleic Acids Res.* **32**, D91-94 (2004).

64. Sherman, B. T. *et al.* DAVID: a web server for functional enrichment analysis and functional annotation of
gene lists (2021 update). *Nucleic Acids Res.* **50**, W216–W221 (2022).

65. Wilkerson, M. D. & Hayes, D. N. ConsensusClusterPlus: a class discovery tool with confidence assessments
and item tracking. *Bioinformatics* **26**, 1572–1573 (2010).

66. Ambrosini, G., Groux, R. & Bucher, P. PWMScan: a fast tool for scanning entire genomes with a position-
specific weight matrix. *Bioinformatics* **34**, 2483–2484 (2018).

67. Donaldson-Collier, M. C. *et al.* EZH2 oncogenic mutations drive epigenetic, transcriptional, and structural
changes within chromatin domains. *Nat. Genet.* **51**, 517–528 (2019).

68. Gregoricchio, S. & Zwart, W. snHiC: a complete and simplified snakemake pipeline for grouped Hi-C data
analysis. *Bioinforma. Adv.* **3**, vbad080 (2023).

69. Wolff, J. *et al.* Galaxy HiCExplorer 3: a web server for reproducible Hi-C, capture Hi-C and single-cell Hi-C
data analysis, quality control and visualization. *Nucleic Acids Res.* **48**, W177–W184 (2020).

70. Wolff, J. *et al.* Galaxy HiCExplorer: a web server for reproducible Hi-C data analysis, quality control and
visualization. *Nucleic Acids Res.* **46**, W11–W16 (2018).

71. Chakraborty, A., Wang, J. G. & Ay, F. dcHiC detects differential compartments across multiple Hi-C datasets.
Nat. Commun. **13**, 6827 (2022).

72. van der Weide, R. H. *et al.* Hi-C analyses with GENOVA: a case study with cohesin variants. *NAR Genomics*
Bioinforma. **3**, lqab040 (2021).

73. Khan, A. & Mathelier, A. Intervene: a tool for intersection and visualization of multiple gene or genomic region
sets. *BMC Bioinformatics* **18**, 287 (2017).

74. Priestley, P. *et al.* Pan-cancer whole-genome analyses of metastatic solid tumours. *Nature* **575**, 210–216
(2019).

75. Chakravarty, D. *et al.* OncoKB: A Precision Oncology Knowledge Base. *JCO Precis. Oncol.* **2017**,
PO.17.00011 (2017).

76. Quinlan, A. R. BEDTools: The Swiss-Army Tool for Genome Feature Analysis. *Curr. Protoc. Bioinforma.* **47**,
11.12.1-34 (2014).

77. Rosenthal, R., McGranahan, N., Herrero, J., Taylor, B. S. & Swanton, C. deconstructSigs: delineating
mutational processes in single tumors distinguishes DNA repair deficiencies and patterns of carcinoma
evolution. *Genome Biol.* **17**, 31 (2016).

78. Tate, J. G. *et al.* COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids Res.* **47**, D941–
D947 (2019).

79. Ahmed, M. *et al.* Variant Set Enrichment: an R package to identify disease-associated functional genomic
regions. *BioData Min.* **10**, 9 (2017).

80. The 1000 Genomes Project Consortium *et al.* A global reference for human genetic variation. *Nature* **526**, 68–
74 (2015).

81. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17**,
10–12 (2011).

82. Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome alignment and genotyping
with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* **37**, 907–915 (2019).

83. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data
with DESeq2. *Genome Biol.* **15**, 550 (2014).

84. Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *The Innovation* **2**,
100141 (2021).

85. Krijger, P. H. L., Geeven, G., Bianchi, V., Hilvering, C. R. E. & de Laat, W. 4C-seq from beginning to end: A
detailed protocol for sample preparation and data analysis. *Methods San Diego Calif* **170**, 17–32 (2020).

86. Vermeulen, M. Identifying chromatin readers using a SILAC-based histone peptide pull-down approach.
Methods Enzymol. **512**, 137–160 (2012).

87. Makowski, M. M. *et al.* An interaction proteomics survey of transcription factor binding at recurrent TERT
promoter mutations. *Proteomics* **16**, 417–426 (2016).

88. Cox, J. & Mann, M. MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass
accuracies and proteome-wide protein quantification. *Nat. Biotechnol.* **26**, 1367–1372 (2008).

89. O'Mara, T. A., Spurdle, A. B., Glubb, D. M., & Endometrial Cancer Association Consortium. Analysis of
Promoter-Associated Chromatin Interactions Reveals Biologically Relevant Candidate Target Genes at
Endometrial Cancer Risk Loci. *Cancers* **11**, 1440 (2019).

90. Perez-Riverol, Y. *et al.* The PRIDE database resources in 2022: a hub for mass spectrometry-based
proteomics evidences. *Nucleic Acids Res.* **50**, D543–D552 (2022).

1049

 FIGURE LEGENDS

Main figures

Fig. 1: ERα cistrome changes upon tumorigenesis.

a Schematic workflow of the multi-omics approach applied in this work. **b-c** Boxplot depicting

the distribution of peak number detected by ERα (**b**) or H3K27ac (**c**) ChIP-seq in healthy

(blue) or tumor (orange) endometrial tissues. **d** Venn diagram of the overlap between

consensus ERα binding sites detected in healthy (blue) and tumor (orange) tissues. **e** MA-

plot of differential binding analyses performed on the consensus ERα ChIP-seq peaks

(Healthy vs Tumor). Differential binding sites (FDR ≤ 0.05 & |log$_2$(FoldChange)| ≥ 1) are

highlighted in pink. **f-g** Tornado plot of the ERα (**f**) or H3K27ac (**g**) ChIP-seq signal at tumor-

depleted (blue, upper blocks) or tumor-enriched (orange, lower blocks) ERα consensus

peaks for all the 4 healthy (blue, left heatmaps) and 5 tumor (orange, right heatmaps)

endometrial tissues. On the top of each heatmap is plotted the average density signal for

each region group. **h** Stacked bar plot depicting the genomic localization frequency of tumor-

depleted and tumor-enriched ERα ChIP-seq consensus peaks. **i** Stacked bar plot displaying

the distance to associated gene TSS frequency distribution of tumor-depleted and tumor-

enriched ERα ChIP-seq consensus peaks. **j** Wordcloud of the top-50 enriched motifs at

tumor-depleted (blues) and tumor-enriched (oranges) ERα consensus peaks. Size and color

intensity are proportional to the -log$_{10}$(E-value). **k** The upper part of the heatmap shows the

individual tumor-depleted (blues) and tumor-enriched (oranges) ERα consensus peaks

colored by signal intensity. On the lower part, each black bar represents an overlapping

peak of ChIP-seq data of several targets publicly available for the Ishikawa endometrial

cancer cell line. **l** Heatmap of the percentage of tumor-depleted or tumor-enriched ERα

consensus peaks overlapping with each ChIP-seq target in Ishikawa cells from (**k**). Ranking

is performed by descending number of overlaps in tumor-depleted peaks.

Fig. 2: 3D genome landscape is remodeled during tumorigenesis.

a Schematic of the Hi-C library preparation starting from fresh frozen tissues. Ten 30μm-thick slices of flash-frozen tissue are cross-linked 25min in 2% formaldehyde. Then, electronically homogenized tissues are filtered using a 75μm cell strainer and subjected to Hi-C library preparation. Hi-C libraries are then sequenced and resulting reads are analyzed by the *snHiC*[68] pipeline. **b** On the left, 40kb-resolution matrices of the average Hi-C score in Tumor (n = 3) and Healthy endometrial tissues (n = 3) at chromosome 6. On the right, differences of the scores shown on the left side (Tumor – healthy). **c** Relative Hi-C Contact Probability (RCP) as function of the distance for each individual sample (40kb resolution). **d** Violin plot of the distribution of the short-range (<2Mb) over long-rage (>2Mb) Hi-C contacts ratio in each individual sample (40kb resolution). For each comparison the Wilcoxon's test *P*-value is indicated. **e** Distribution of the Hi-C loop distance (10kb resolution) for loops overlapping with tumor-depleted, tumor-enriched or shared ERα consensus peaks in combined healthy (blue) or tumor (orange) endometrial tissues. **f** PE-SCAn results in healthy and tumor tissues at common, tumor-depleted and tumor-enriched ERα binding sites. For each panel, top and middle rows depict the 3D and 2D, respectively, representation of the Hi-C contact frequency distribution in healthy (left) and tumor (right) tissues; lower row shows the difference of Hi-C contact frequencies between Tumor (orange) and Healthy (blue) tissue scores. **g** Compartment polarization ratio (100kb resolution), defined as (AA + BB) / (AB + BA), for each individual sample. **h** *Top*: saddle plot of A/B compartments interactions as computed in (**g**) for each individual sample. *Bottom*: difference of saddle-score with the reference H.005.A2 (healthy tissue), where orange indicates a higher score in tumor samples while purple indicates a higher score in healthy tissues. **i** Sankey plot depicting the proportion of compartment state transition (100kb resolution) of healthy compartments (left) upon tumorigenesis (right). **j** Upset plot showing the overlaps between different compartment transition states (100kb resolution) in healthy compared to tumor as in (**g**) (A-to-A, B-to-B, A-to-B, B-to-A) and different categories of ERα consensus peaks. **j** Stacked bar plot of the proportion of compartment transition status as in (**g**) for individual genomic bins overlapping with only one of the different categories of ERα consensus peaks.

Fig. 3: Tumor-enriched ERα binding sites represent non-coding regions target of somatic mutation in metastatic endometrial cancer tumor.

a-c Genomic localization (**a**), mutational signature (**b**) and counts (**c**) distribution of the somatic mutations detected by WGS in the primary endometrial cancer samples form a selection of samples from the TCGA-UCEC cohort (n=41). **d** Stacked bar blot depicting the fraction of ERα peaks overlapping with somatic mutations identified in the primary cohort. **e** Variant Set Enrichment (VSE) analysis depicting the enrichment of differentially bound ERα sites over endometrial cancer risk loci identified by genome-wide association study (GWAS, $P < 1×10^{-5}$)[26]. Density plot represents distribution for the Z-score from matched controls defining the null distribution. Blue (tumor-depleted) or orange (tumor-enriched) vertical lines represent observed enrichments, with tumor-enriched enrichment being statistically significant (P_{adj} = 0.0060). Gray vertical lines define 0, 25, 75, 100 percentiles of the distribution, while dotted black line indicates the median. **f** Scheme of the number metastases and metastatic site, for all metastatic samples used for the WGS analyses. **g-h** Genomic localization (**g**) and mutational signature (**h**) distribution of the somatic mutations detected by WGS in the metastatic samples described in (**f**). **i** In the outer plot, it is shown the number of different types of somatic variants detected in each metastatic sample. In the inner plot, stacked bar plot of the most-frequently protein sequence mutated genes and relative number and type of somatic variants identified in metastatic samples. **j** Stacked bar blot depicting the fraction of ERα peaks overlapping with somatic mutations identified in the metastatic cohort. **k** MA-plot of differential expression analyses in Ishikawa cell lines upon 8h of 10nM β-estradiol (E2) stimulation. Black dots indicate genes which promoter has been linked, by H3K27ac Hi-C analyses, to a tumor-depleted or tumor-enriched ERα-bound regulatory element bearing somatic mutations in metastatic samples (n=311). Differentially expressed genes (|Fold Change expression| ≥ 1.5 & $P_{adj} < 0.05$) upon estradiol stimulation have been labeled and highlighted by a red dot (n=12). **l** Ishikawa endometrial cancer cells were treated or not for 6 or 12 hours with 10nM estradiol. Whole-cell extracts were analyzed by immunoblotting with antibodies against ERα and GAPDH (loading control). **m** Progression-Free Kaplan-Meier curve of endometrial cancer patients (TCGA data) divided into two groups using the median of *ESR1* expression (FPKM) as cut-off.

Fig. 4: Short-range chromatin contacts at the *ESR1* locus are stronger in tumors.

a ChIP-seq genomic tracks for ERα (upper panel) and H3K27ac (lower panel) in healthy (blue) and tumor (orange) endometrial primary tissues at the *ESR1* locus. *ESR1 Enhancer 1* and *Enhancer 2* are indicated, as well as the mutations found by WGS analyses in metastatic samples. **b** Representation of the averaged 40kb resolution Hi-C matrix at the *ESR1* locus for the 3 healthy (top) and 3 tumor (middle) tissues or the score difference (bottom) tumor – healthy, where orange indicates higher scores in the tumors while purple higher scores in healthy. The matrices scores are dived by the sum of the matrix. Black lines indicate the Topologically Associated Domains (TADs) identified. **c** 4C-seq genomic tracks at the *ESR1* locus using the *ESR1* TSS as View-Point (VP) for 2 healthy (top, blue) and 3 tumor (middle, orange) endometrial tissues, and the average difference of score tumor – healthy (bottom). With green arcs are depicted loops detected by H3K27ac HiChIP in Ishikawa endometrial cancer cells. The ribbon around the 4C-seq signal lines indicates the Standard Error Mean (SEM) among biological replicates. **d** In the top row, Observed/Expected matrices of sequence-based machine learning prediction in a ±500kb window surrounding the *ESR1* locus. In order from left to right can be found: wild-type sequence, point mutation in a genomic desert (negative control), deletion of the full *Enhancer 1* sequence (positive control), introduction of SNVs found by WGS analyses of metastatic endometrial cancer samples. On the bottom row, is showed the difference of Observed/Expected score over the wild-type sequence, where orange indicates higher scores in the altered sequence and purple a higher score in the wild-type one.

Fig. 5: EHMT2/G9a is a negative regulator of *ESR1* expression in endometrial cancer.

a Schematic workflow used to perform DNA-oligo protein pull-downs. Biotin-conjugated wild-type or mutated DNA-oligos are immobilized on streptavidin magnetic beads and mixed with Ishikawa nuclear lysates. Captured proteins are then dimethyl labeled and analyzed by mass spectrometry. **b** A DNA oligo with the sequence of the ±25bp surrounding the ERα binding site in the *ESR1 Enhancer 1*, in wild-type or chr6:152,002,679–TCT-to-T form, was used to perform DNA-oligo protein pull downs in Ishikawa cells as described in (**a**). The scatter plot shows the $\log_2$ ratios of all identified and quantified proteins in both experiments plotted against each other. Proteins significantly enriched at the wild type sequence are highlighted in red, and proteins significantly enriched at the mutant sequence are highlighted in blue. **c** Gene expression correlation heatmap for all corresponding 24 differentially bound proteins identified in (**b**) in 589 endometrial cancer patients (TCGA). Dashed lines indicate the separation between positive and negative correlation scores. Genes are ranked by correlation with *ESR1* gene expression. **d** Progression-Free Kaplan-Meier curve of endometrial cancer patients (TCGA data) divided into two groups using the median of *EHMT2/G9a* expression (FPKM) as cut-off. **e** EHMT2/G9a and IgG (negative control) ChIP followed by Western blot in Ishikawa endometrial cancer cells. Antibodies against EHMT2/G9a and ERα have been used. For EHMT2/G9a, two different exposure images were used for input and IP, as indicated by the vertical dashed bar. **f** EHMT2/G9a ChIP in Ishikawa cells stimulated for 6h with 10nM β-estradiol. Barplot shows percentage of enrichment over the input (% Input) at the *ESR1 Enhancer 1*, *ESR1* promoter and *CDK12* promoter (negative control) analyzed by quantitative PCR (qPCR). Mean of 2 independent experiments is shown. **g** Ishikawa endometrial cancer cells were stimulated for 72h with 10nM β-estradiol in combination or not with 100nM ICI-182,780 (Fulvestrant, negative control). In all conditions, cells where incubated either with a Non-Targeting (NT) siRNA or with an siRNA against *EHMT2/G9a*. Then, whole-cell extracts were analyzed by immunoblotting with antibodies against ERα, EHMT2/G9a and GAPDH (loading control). **h** Differential ERα peak centered EHMT2/G9a average ChIP-seq density profiles in Ishikawa cells upon 6h treatment with 10nM β-estradiol (purple) or DMSO (blue, control). Ribbon indicates SEM. Paired Wilcoxon test was performed on the average score of the highlighted area for each individual region analyzed; *P* values are indicated. **i** Schematic representation of the *ESR1* gene regulation working model. Upon tumorigenesis, ERα is re-located to an endometrial cancer specific *ESR1* enhancer (**Fig. 1f-g**). ERα interacts with EHMT2/G9a (**Fig. 5e**) and binds both enhancer and promoter regions of ESR1, in a hormone-dependent fashion (**Fig. 5f**). In a subset of metastatic endometrial cancers, a somatic mutation is acquired at the tumor-specific *ESR1* enhancer (**Fig. 4a**). *In vitro* analyses probing this mutation revealed a loss of EHMT2/G9a DNA binding capacity (**Fig. 5b**), and EHMT2/G9a

1193   knockdown in endometrial cancer cell lines leads to an increase of *ESR1* expression levels

1194   (**Fig. 5g**).

Supplementary Figures

Supplementary Fig. 1: Endometrial cancer tissues and ChIP-seq quality controls.

a Hematoxylin and eosin stain of healthy (top) and tumor (bottom) representative endometrial tissues. These tissues correspond to tissues used for Hi-C and RNA-seq analyses. **b** Bar plot (top) and box plot (bottom) showing the total number of reads of the ERα (black) and H3K27ac (red) ChIP-seq libraries among all the healthy (blue) and tumor (orange) tissue samples. Wilcoxon test: $P_{ERα} = 0.659$, $P_{H3K27ac} = 0.21$ (ns = not significant, $P \geq 0.05$). **c** Principal Component Analyses (PCA) of ERα (circles) and H3K27ac (triangles) ChIP-seq data in 4 healthy (blues) and 5 tumor (orange) endometrial primary tissues. **d** Unsupervised permutation clustering test results. Gradient indicates the clustering probability between samples. **e** Distribution of the number of ERα differential peaks obtained by 100 iterations of label reshuffling. **f** Boxplot of the normalized read count distribution at ERα bindings sites detected in healthy (blue) or tumor (orange) tissues, as well as at differential consensus ERα bindings sites with lower (–) or higher (+) affinity in tumor tissues. **g** Bubble plot showing the relative enrichment of DAVID Gene Ontology biological processes (GO-BP) enrichments for tumor-depleted ERα promoter-bound genes. Tumor-enriched ERα promoter-bound genes were used as background data set. **h** Stacked bar plot showing the presence (PWMScore > 0) frequency of ESR1/ERα motif in different ERα binding categories. **i** Bar plot depicting the average PWMScore ± SEM at different ERα binding categories. P value of Wilcoxon test is indicated. **j** Top 20 ranked factors identified to be enriched at different ERα binding categories as identified by GIGGLE[20] analyses.

Supplementary Fig. 2: Differential gene expression between normal and tumor endometrial tissues.

a Principal Components analyses of RNA-seq data for 3 healthy (blue shades) and 3 tumor (orange shades) samples. Scatter plot between PC1 and PC2 is shown. **b** Volcano plot of the differential gene expression analyses between 3 normal (blue) and 3 tumor (orange) endometrial tissues. Vertical lines indicate a Fold Change Expression equal to 2 (positive side)/0.5 (negative side), while the horizontal line indicates a $P_{adj} = 0.05$. **c** Violin plot showing the *DESeq2* normalized counts in healthy (blue) vs tumor (orange) tissues for the genes included in the over-represented biological processes identified in **Supplementary Fig. 1g**. *P*-value of a paired Wilcoxon test is indicated above each category, in gray when above 0.05, and in black when below 0.05 (statistically significant). **d** GSEA analyses on full transcriptome based on genes ranked by fold change differential expression between tumor and normal tissues. Bar plot of the Normalized Enrichment Score (NES) is shown for statistically significant gene sets ($P < 0.05$). **e** Stacked plot of the differential expression

1231 frequency of genes associated to different classes of ERα binding sites. **f** Volcano plot of the
1232 differential gene expression analyses between 3 normal (blue) and 3 tumor (orange)
1233 endometrial tissues of genes associated with different classes of ERα binding sites. Vertical
1234 lines indicate a Fold Change Expression equal to 2 (positive side)/0.5 (negative side), while
1235 the horizontal line indicates a P_{adj} = 0.05.

1236

Supplementary Fig. 3: 3D genome organization analyses in endometrial primary
tissues.
a Chromosome matrix for each individual sample showing the chromosome translocation
score per each chromosome based on the Hi-C contact probability (500kb resolution). **b**
Pearson Hi-C contacts correlation matrix among different healthy (blue) and tumor (orange)
endometrial primary tissues. **c** Stacked bar plot showing the Hi-C contact type frequency per
each sample, as computed by *HiCExplorer*. Numbers indicate the millions of Hi-C contacts
(paired-end fragments) detected. **d** Average Relative Hi-C Contact Probability as function of
the distance for healthy and tumor samples (40kb resolution). **e** Violin plot of the average
distribution of the short-range (<2Mb) over long-rage (>2Mb) Hi-C contacts ratio in healthy
and tumor samples (40kb resolution). Wilcoxon's test *P*-value is indicated. **f** Average
compartment polarization ratio (100kb resolution), defined as (AA + BB) / (AB + BA), for
healthy and tumor samples. **g** On the upper part, average saddle plot of A/B compartments
interactions as computed in (**f**) for each tissue type. On the lower part, Tumor-Healthy
difference of saddle-score, where orange indicates a higher score in tumor samples while
purple a higher score in healthy tissues. **h** On the left, scatter plot showing the correlation of
average individual compartment bin scores in tumor vs healthy endometrial tissues and
overlapping with the different categories of ERα consensus peaks. On the right, bin count
fold change for each compartment transition subcategory (A-to-A, B-to-B, A-to-B, B-to-A)
relative to the count of bins not overlapping with any ERα binding site. **i** Quantification of the
average 4C-seq signal from **Fig. 4c** at the "*Enhancer 1*" and "*Enhancer 2*" regions per each
tissue sample. Each point represents the replicate mean value of each sample. Genomic
location (GRCh37/Hg19) of the two regions are indicated in parenthesis.

1260

Supplementary Fig. 4: Example of chromatin loops in healthy and tumor tissues
Heatmaps depict the averaged 10kb resolution Hi-C matrix for the 3 healthy (upper) and 3
tumor (lower) tissues. Loops specific to healthy (blue) or tumor (orange) tissues are
indicated by squares on the heatmaps, while the dashed lines represent the diagonal
projection of the loop anchors. Loops are indicated by arcs that connect the loop anchors.
Genomic tracks are showing the normalized ChIP-seq signal for ERα (top tracks) and
H3K27ac (bottom tracks) in healthy (blue) and tumor (orange) endometrial tissues.

1268

Supplementary Fig. 5: *ESR1* gene expression in normal and tumor endometrial human

samples

a RNA-seq *ESR1* gene expressed as *DESeq2* normalized counts between healthy and

tumor endometrial tissues from data generated in this work. Each individual sample is

represented by a dot. *P* value of Wilcoxon test is indicated. **b** Violin and box plot showing the

RNA-seq *ESR1* gene expression from endometrial carcinoma TNMplot data using normal

samples from non-cancerous patients as control. Average of 146 healthy and 547 tumor

samples is shown. *P* value of Wilcoxon test is indicated. **c** Violin and box plot showing the

RNA-seq *ESR1* gene expression from endometrial carcinoma TNMplot data using paired

tumor and adjacent normal tissues samples. Average of 23 patients is shown. *P* value of

paired Wilcoxon test is indicated.

1280

1281 **Supplementary Fig. 6: Somatic mutation overlaps with ERα binding sites in**

1282 **hypermutated and non-hypermutated primary and metastatic samples**

1283 **a-b** Stacked bar blot depicting the fraction of ERα peaks overlapping with somatic mutations

1284 identified in the primary (**a**) or metastatic (**b**) patient cohort. Overlaps of mutation occurring in

1285 non-hypermutated (*left*) and hypermutated (*right*) are shown independently.

1286

Supplementary Fig. 7: Genomic and epigenomic landscape comparison of the *ESR1*

locus in endometrial and breast cancer.

a Source images and Ponceau Red staining of the western blot shown in **Fig. 3l**. Cropped

areas in the main figure are indicated by a red rectangle. **b** Average density profile of ATAC-

seq signal from publicly available data (TCGA) for endometrial and breast cancer at the

ESR1 locus (*Enhancer 1*, *Enhancer 2* and promoter). Signal was smoothed using the loess

regression method. **c** ChIP-seq genomic tracks for ERα in healthy endometrial tissues (top,

blue), endometrial tumors (middle, orange) and breast cancer (bottom, pink) at the *ESR1*

locus. Focused average density signal at *ESR1 Enhancer 1*, *Enhancer 2* and promoter are

plotted on the lower panel. Average density signal was smoothed using the loess regression

method.

1298

Supplementary Fig. 8: Validation of the TCGA gene expression analyses.

a Density distribution of the *ESR1* gene expression correlation with the whole transcriptome

in TCGA RNA-seq data. Vertical blue and red lines indicate the gene expression correlation

score between *ESR1* and the top 3 anti-correlated and 3 correlated genes showed in **Fig.**

5c. Black dotted vertical lines indicate respectively the 5^{th}, 50^{th} (median) and 95^{th} percentile

of the distribution. **b** Progression-Free Kaplan-Meier curve of endometrial cancer patients

(TCGA data) divided into two groups using the median of expression gene (FPKM) as cut-

off: blue high expression, red low expression. On the upper part are shown the curves for

GTF2IRD1, *ZNF768*, *SIN3A*, while on the lower part *XPA*, *ZBTB21*, *POLK*. **c-d** Source

images and Ponceau Red staining of the western blot shown in **Fig. 5e** (**c**) and **Fig. 5g** (**d**).

Cropped areas in the main figure are indicated by red rectangles.

1310

Supplementary Fig. 9: EHMT2/G9a functional analyses in AN3CA ERα-deficient

endometrial cancer cell lines

a Whole protein extracts from Ishikawa and AN3CA endometrial cancer cell lines were

immunoblotted using antibodies against EHMT2/G9a, ERα and GAPDH (loading control). **b**

Source images and Ponceau Red staining of the western blot shown in (**a**). Cropped areas

in the main figure are indicated by red rectangles. **c-d** Normalized cell confluency of

Ishikawa (**c**) and AN3CA (**d**) endometrial cancer cells upon DMSO (black) or 100nM ICI-

182,780/Fulvestrant (green) treatment over time. Average ± SEM of 6 replicates is shown. **e**

EHMT2/G9a ChIP in AN3CA cells stimulated for 6h with 10nM β-estradiol. Bar plot shows

percentage of enrichment over the input (% Input) at the *ESR1 Enhancer 1*, *ESR1* promoter

and *CDK12* promoter (negative control) analyzed by quantitative PCR (qPCR). Mean of 2

independent experiments is shown. **f** AN3CA endometrial cancer cells were stimulated for

72h with 10nM β-estradiol in combination or not with 100nM ICI-182,780 (Fulvestrant). In all

conditions, cells where incubated either with a Non-Targeting (NT) siRNA or with an siRNA

against *EHMT2/G9a*. Then, whole-cell extracts were analyzed by immunoblotting with

antibodies against ERα, EHMT2/G9a and GAPDH (loading control). **g** Source images and

Ponceau Red staining of the western blot shown in (**f**). Cropped areas in the main figure are

indicated by red rectangles.

 TABLE LEGENDS

Supplementary Table 1: Clinicopatholigical features of patient samples and omics data streams generated.
Description of the collection site, histological status, site of collection, age at diagnosis, tumor percentage, grade and FIGO classification for each patient sample. The last column indicates the omics data generated.

Supplementary Table 2: List of publicly available ChIP-seq data sets in Ishikawa cells.
List of transcription and epigenetic factors ChIP-seq data in Ishikawa cells and corresponding ENCODE accession number.

Supplementary Table 3: Metadata relative to metastatic endometrial cancer samples.
In this tables are reported the fraction of tumor cells, year of birth, biopsy site and eventual pre-treatments of the 24 metastatic endometrial cancer samples.

Supplementary Table 4: List of ERα binding sites and H3K27ac HiChIP linked target genes
List of candidate target genes for the tumor-gained mutated ERα sites based on H3K27ac HiChIP analysis of Ishikawa cells[89]. The structure of the table is the following: *chr_enh/start_enh/end_enh*: genomic position of the ERα binding sites; *count_enh*: enhancer identifier; *target_gene/ensemble_ID*: candidate target gene of enhancer from endometrial cancer H3K27ac HiChIP data; *enhancer_in_HiChIP_target_gene_promoter_anchor?*: *yes*, enhancer is located within the HiChIP anchor that encompasses the gene TSS – *no*, enhancer is located in the HiChIP anchor distal to the gene; *enhancer_ within_3kb_TSS_of_target_gene?*: *yes*, enhancer is located within ±3kb of the TSS of the gene – *no*, enhancer is not located within 3kb of the TSS of the gene.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5