1 **Population genetics of *Glossina palpalis gambiensis* in the sleeping sickness focus**
2 **of Boffa (Guinea) before and after eight years of vector control: no effect of control**
3 **despite a significant decrease of human exposure to the disease**
4

5 Moise S. Kagbadouno[1], Modou Séré[2], Adeline Ségard[3,4], Abdoulaye Dansy Camara[1],
6 Mamadou Camara[1], Bruno Bucheton[1,3,4], Jean-Mathieu Bart[1,3,4], Fabrice Courtin[3,4,5],
7 Thierry de Meeûs[3,4$*] Sophie Ravel[3,4$]
8
1 Programme National de Lutte contre la THA, Ministère de la Santé, Conakry, Guinée.
2 University of Dédougou, Dédougou B.P. 176, Burkina Faso.
3 Univ Montpellier, IRD, Cirad, Intertryp, Baillarguet, 34398 Montpellier Cedex 5, France.
4 IRD, Intertryp, UMR 177, Montpellier, France.
5 Représentation de l'IRD au Burkina Faso, 668, avenue, Pr Joseph Ki-Zerbo-Koulouba,
01 BP 182 Ouagadougou 01, Burkina Faso.
15
Moise Kagbadouno: moise.kagbadouno@gmail.com; Modou Séré:
modousere11@gmail.com; Adeline Ségard: adeline.segard@ird.fr; Abdoulaye Dansy
Camara: camaradansy@gmail.com; Mamadou Camara: mamadycamarafr@yahoo.fr;
Bruno Bucheton: bruno.bucheton@ird.fr; Jean-Mathieu Bart: jean-mathieu.bart@ird.fr;
Fabrice Courtin: fabrice.courtin@ird.fr; Thierry de Meeûs: thierry.demeeus@ird.fr; Sophie
Ravel: sophie.ravel@ird.fr
22
23 $ equal supervision
24
25 *Corresponding author: thierry.demeeus@ird.fr

26
27 Index keywords: Tsetse flies, Human African Trypanosomiasis, Population structure,
28   Effective population size, Vector control, Neglected Tropical Diseases.
29

Abstract

Human African trypanosomosis (HAT), aka sleeping sickness, is still a major concern in endemic countries. Its cyclical vector are biting insects of the genus *Glossina* or tsetse flies. In Guinea, the mangrove ecosystem contains the main HAT foci of Western Africa. There, the cyclical vector is *Glossina palpalis gambiensis*. A still ongoing vector control campaign (VCC) started in 2011 in the focus of Boffa, using tiny targets, with a significant impact on the prevalence of the disease. To assess the sustainability of these results, we have studied the impact of this VCC on the population biology of *G. p. gambiensis* in Boffa. We used the genotyping at several microsatellite markers and population genetics tools of tsetse flies from different sites and at different dates before and after the beginning of the VCC. In variance with a significant impact of VCC on the apparent densities of flies captured in the traps deployed, the global population of *G. p. gambiensis* displayed no genetic signature of control, and behaved as a very large population occupying the entire zone. This implies that targets deployment efficiently protects the human populations locally, but not where targets cannot be deployed and where the main tsetse population exploits available resources. This also entails the pursuit of vector control measures until HAT can be considered as entirely eradicated from the focus.

Introduction

Human African trypanosomosis (HAT), aka sleeping sickness, is still a major concern for the WHO (Holmes, 2014; Simarro et al., 2015). Its agent is an euglenozoan kinetoplastid parasite, *Trypanosoma brucei gambiense* 1 (Jamonneau et al., 2019) (Tbg1), with a clonal propagation (Mathurin Koffi et al., 2009; M. Koffi et al., 2015), which recently emerged 10,000 years ago and spread from West Africa (W. Weir et al., 2016). It encompasses a complex life cycle, with a phase in blood sucking insects of the genus *Glossina* (Diptera, Hipposcoidea), its cyclical vector (Bouyer et al., 2015).

In Guinea, the mangrove ecosystem still contains the main HAT foci of Western Africa (Simarro et al., 2015). There, *Glossina palpalis gambiensis* is the only known vector of the disease (Courtin et al., 2015). Boffa, is one of the three active HAT foci of the country (Kagbadouno et al., 2012). An active vector control campaign (VCC), aiming at reducing the human/tsetse contact, and subsequently at interrupting the transmission of Tbg1, was implemented in 2011 (Courtin et al., 2015).This VCC produced a significant decrease in exposure risks and prevalence of the disease (Courtin et al., 2015). As regard to the vertebrate host, the exact role of animal or asymptomatic human reservoirs is still uncertain. Nevertheless, the genetic diversity of this parasite maintained by the Guinean populations suggests the existence of a significant amount of individuals clones outside the patients involved in surveys (Mathurin Koffi et al., 2009), in human and/or animal reservoirs. Several other studies appeared to confirm this hypothesis (Bucheton et al., 2002; Büscher et al., 2018).

A recent study of *G. palpalis palpalis* in the focus of Bonon (Côte d'Ivoire), revealed that a special allele at a given trinucleotide locus (GPCAG), experienced a brutal increase in frequency after control (Berté et al., 2019). This suggested the selection for resistance against control measure.

Given these observations, deciding what strategies to use in the future will depend on the most precise knowledge we can get on the population impact of medical campaigns and VCC.

In this paper, we investigated the impact of VCC over 11 years on the HAT focus of Boffa (Figure 1). We report here the impact of VCC on capture densities and sex-ratio, as well as on *G. palpalis gambiensis* population genetics, in particular on locus GPCAG. We then discuss the best strategy to be used to optimize the protection of human populations from this neglected tropical disease.

Material and Methods

Data collection

All sites used for the entomological survey, number of trapped flies and their gender can be seen in Kagbadouno et al's paper (Kagbadouno et al., 2012) and Supplementary file S1.

Localization, date of sampling and number of genotyped flies can be found in Figure 1 and Table 1. A total of 290 flies were genotyped at 11 loci. Vector control was implemented with tiny targets from February 2012 in the sites of the left bank of river Rio Pongo and then on both sides after 2016. Then, according to Table 1, flies from 2009 and 2011 samples correspond to T0 (before control) on both sides of River Rio Pongo, while samples from 2019 and 2020 correspond to TX (after control has begun), on both sides of the river.

Figure 1: Localization of traps for the capture of *Glossina palpalis gambiensis* in the HAT focus of Boffa in Guinea. Dates of capture, number of flies genotyped, and other geographical information can be found in the Table 1.

Table 1: Number of genotyped *Glossina palpalis gambiensis* (*N*) in the different traps of different sites and date of capture in the HAT Focus of Boffa (Guinea). The cohort (considering six generations of tsetse flies per year) and the time (T) after the beginning of control in months (0=before), the bank of the Rio Pongo River (see Figure 1) and the land type are also indicated.

Site	Trap	Date	Cohort	T	Bank	Land	*N*	N C-T-Bank
Guiapet	4	10/2009	0	0	Left	Island	23	C0-T0-LeftBank
Yangoya	1	10/2009	0	0	Left	Island	30	53
Bakyia	1	05/2011	10	0	Left	Continent	11	
Bakyia	2	05/2011	10	0	Left	Continent	8	
Bakyia	3	05/2011	10	0	Left	Continent	5	
Guiapet	2	05/2011	10	0	Left	Island	7	
Guiapet	3	05/2011	10	0	Left	Island	4	C10-T0-LeftBank
Yangoya	1	05/2011	10	0	Left	Island	7	65
Yangoya	5	05/2011	10	0	Left	Island	3	
Yangoya	6	05/2011	10	0	Left	Island	2	
Yangoya	7	05/2011	10	0	Left	Island	16	
Yangoya	8	05/2011	10	0	Left	Island	2	
Dobire	6	05/2011	10	0	Right	Island	20	
Dobire	7	05/2011	10	0	Right	Island	12	
Santani	2	05/2011	10	0	Right	Continent	10	C10-T0-RightBank
Santani	4	05/2011	10	0	Right	Continent	4	66
Santani	5	05/2011	10	0	Right	Continent	20	
Bakyia	2	10/2019	61	92	Left	Continent	4	
Guiapet	2	10/2019	61	92	Left	Island	2	
Guiapet	3	10/2019	61	92	Left	Island	11	
Yangoya	1	11/2019	61	93	Left	Island	2	C61-TX-LeftBank
Yangoya	5	11/2019	61	93	Left	Island	4	35
Yangoya	6	11/2019	61	93	Left	Island	1	
Yangoya	7	11/2019	61	93	Left	Island	11	
Dobire	6	11/2019	67	33	Right	Island	9	
Dobire	7	10/2019	67	32	Right	Island	1	
Santani	2	11/2019	67	33	Right	Continent	8	C61-TX-RightBank
Santani	4	11/2019	67	33	Right	Continent	1	27
Santani	5	11/2019	67	33	Right	Continent	8	
Bakyia	2	10/2020	67	104	Left	Continent	3	
Guiapet	1	10/2020	67	104	Left	Island	15	C67-TX-LeftBank
Guiapet	2	10/2020	67	104	Left	Island	1	24
Yangoya	7	10/2020	67	104	Left	Island	5	
Dobire	1	10/2020	67	44	Right	Island	6	
Dobire	5	10/2020	67	44	Right	Island	2	C67-TX-RightBank
Santani	1	10/2020	67	44	Right	Continent	10	20
Santani	3	10/2020	67	44	Right	Continent	2	

N C-T-Bank: Names and numbers of riverbank subsamples, with the cohort (C) and the time (T) before (0) or after (X) the beginning of control.

Active surveillance began in 2011, where 2730 individuals (1474 females and 1256

males) were captured. In 2019, 143 flies (80 females and 63 males) and in 2020, 215 (106

females and 109 males) were caught. The details on number of flies in the different dates

and locations, and all genotypes, are available in the supplementary file S1.

Assuming a two months generation time (Williams, 1990), as can be seen from

Table 1, the genotyped subsamples extended from cohort 0 (2009), 10 (2011), 61 (2019)

and 67 (2020). Because individuals belonging to one of these cohorts have no chance of

interacting with any individual from another of these cohorts, we then always considered

these four cohorts into separate entities.

Genotyping

Genotyping was undertaken for 11 loci: X55-3 (Solano et al., 1997), XpGp13,

pGp24 (Luna et al., 2001), GPCAG (Baker & Krafsur, 2001), pGp15, pGp16, and pGp27

(Ravel et al., 2020). Loci A10, B3, XB104, C102 were kindly provided by A. Robinson

(Insect Pest Control Sub-program, Joint Food and Agriculture Organization of the United

Nations/International Atomic Energy Agency Program of Nuclear Techniques in Food and

Agriculture). Despite used at several occasions (Camara et al., 2006; Bouyer et al., 2007;

Solano et al., 2009; Melachio et al., 2011), information about these loci were never

published. Readers will find this information in Table A1 in the Appendix. Locus name with

an X means that the locus is located on the X chromosome, and thus haploid in males.

Cells with a "0" correspond to individuals with no amplification at the concerned locus,

while uninterpretable profiles were coded "NA" (see Supplementary file S1).

Data analyzed

Genotyping was undertaken in different ways and under different conditions for the

different dates of sampling (see Table S1). For samples collected in 2009 and 2011, allele

bands were resolved on a 4300 DNA Analysis System (LICOR, Lincoln, NE) using two

infrared dyes (700 and 800nm) as described in (Solano et al., 2009). For samples

collected in 2019 and 2020, an ABI 3500xl sequencer (Applied Biosystem, Waltham,

Massachusetts) was used with four different dyes (FAM, NED, VIC and PET) as described

in (Berté et al., 2019). Consequently, the 11 loci were not analyzed with the same dyes

between samples from 2009-2011 and samples from 2019-2020. Because it is generally

admitted that the dye used can modify the apparent size of the amplified alleles, we

genotyped again some samples from 2009-2011 at the different loci using the ABI 3500xl

sequencer to recode all the genotypes from the samples from 2009-2011 In some subsets,

too many missing data were observed at some loci or subsamples. Consequently, to avoid

multiple analyses, keep a reasonable number of loci and for the sake of consistency,

population genetics data analyses were undertaken with females only, at the eight loci that

were successfully amplified over all the dataset (see Table S1): X55-3, XpGp13, pGp24,

A10, B3, XB104, C102, and GPCAG.

Unless specified otherwise, the global dataset (File S1) was processed with Create

(Coombs et al., 2008) to produce datasets in different formats, depending on the kind of

analyses to be undertaken.

Selection of the relevant subsample units

The reproductive system of a population from a genotyped sample can be assessed

through Wright's F_{IS} (Wright, 1965), which is a measure of the relative inbreeding of

individuals as compared to the subpopulation they belong to. This parameter was

estimated with Weir and Cockerham's unbiased estimator f (B. S. Weir & Cockerham,

1984), that we kept labelling F_{IS} for the sake of simplicity. In case of random mating in a

dioecious population, a negative value is expected (Robertson, 1965; Pudovkin et al.,

1996; Balloux, 2004; De Meeûs & Noûs, 2023). In tsetse flies, the trap can be the unit at

which a significant subdivision may occur (Ravel et al., 2023), including *G. palpalis*

gambiensis (Bouyer et al., 2009). Here, traps, and even sites, in combination with the

cohort, often contained very few flies. To determine if traps, sites or river banks (right or

left of the Pongo River, see Figure 1) really mattered, we used the Wahlund effect

approach (Goudet et al., 1994; De Meeûs et al., 2006). We compared the F_{IS_T} in traps,

the F_{IS_S} in sites (ignoring traps), the F_{IS_B} within the two riverbanks (ignoring sites), and

the F_{IS_C} within the four cohorts (0, 10, 61 and 67) (ignoring river banks). For this, we

undertook a Friedman two-way analysis of variance by ranks for data paired by locus with

the package R-commander (J. Fox, 2005; J. Fox, 2007) (rcmdr) for R (R-Core-Team,

2022). We then undertook planned one sided Wilcoxon signed rank test for paired data. If

traps and/or sites and/or river banks matter, we expect a Wahlund effect (De Meeûs et al.,

2007; De Meeûs, 2018) when pooling individuals from different traps, and/or from different

sites and or different river banks of the same cohort, meaning that the alternative

hypothesis was $F_{IS\_T} < F_{IS\_S} < F_{IS\_B} < F_{IS\_C}$. When necessary (significant *p*-values), we

corrected the *p*-values with the Benjamini and Yekutieli correction (Benjamini & Yekutieli,

2001) with R (command "p.adjust"), for test series with dependency.

Quality testing of loci and subsamples

Quality testing of the subsamples and genotyping was assessed with linkage disequilibrium (LD), Wright's F_{IS} and F_{ST} (Wright, 1965) and their variance across loci.

Linkage disequilibrium was tested with the *G*-based randomization test between each locus pair across subsamples as described in (De Meeûs et al., 2009), with 10,000 random shuffling of genotypes of the two loci of each pair. Because there are as many non-independent tests as locus pairs, we adjusted the *p*-values with the Benjamini and Yekutieli (BY) procedure with R (command "p.adjust").

Wright *F*-statistics were estimated with Weir and Cockerham's unbiased estimators (B. S. Weir & Cockerham, 1984): *f* for F_{IS}, *θ* for F_{ST} and *F* for F_{IT}. Their significant deviation from the expected value under the null hypothesis (H0: F_{IS}=0 or F_{ST}=0) was tested with 10,000 randomizations of alleles between individuals within subsamples (for F_{IS}) and of individuals between subsamples (for F_{ST}). For F_{IS}, the statistic used was Weir and Cockerham's *f* for each locus and overall. For testing subdivision we used the *G* based test (Goudet et al., 1996) for each locus and overall. For F_{IS}, tests are one-sided (F_{IS}>0), and two-sided *p*-values were obtained by doubling the one-sided *p*-value if <0.5 or we doubled 1-*p*-value otherwise. Confidence intervals (95%CI) were computed with 5000 bootstraps over loci for the average over loci, and over individuals for the confidence intervals around each locus. The standard error of jackknives over loci (SE) was computed for F_{IS} and F_{ST} (SE_F_{IS} and SE_F_{ST} respectively) for null allele diagnostics (see below).

All this computations and randomizations were undertaken with Fstat 2.9.4 (Goudet, 2003), updated from Fstat 1.2 (Goudet, 1995), except the bootstraps over individuals, for which we used Genetix (Belkhir et al., 2004). Since Genetix computes bootstraps for each locus in each subsamples, we averaged these values across subsamples to obtain 95%CI of F_{IS} for each locus.

Null allele signatures were looked for with several criterions: the ratio of SE of F_{IS} over F_{ST} (r_{SE}), as computed with jackknives over loci; the correlation between F_{IS} and F_{ST}, and between F_{IS} and the number of missing genotypes (N_b) (De Meeûs, 2018). These correlations were tested with a one-sided (positive correlation) Spearman's signed rank correlation test. The regression F_{IS}~N_b was also undertaken to determine the proportion of the variance of F_{IS} that was explained by missing genotypes (putative null homozygotes) with the determination coefficient R^2, and to extrapolate the possible basic F_{IS} (F_{IS_0}) in absence of null alleles through the intercept of this regression. Null allele frequencies were also estimated with the EM algorithm (Dempster et al., 1977) implemented by FreeNA (Chapuis & Estoup, 2007). For these analyses, missing genotypes (true blanks) were

coded as null homozygotes for allele 999, as recommended (Chapuis & Estoup, 2007).

The goodness of fit of observed blank genotypes, with the expected number of missing

data, was assessed with a one-sided exact binomial test with R (command "binom.test"). If

null alleles and random mating fully explain the data, there should be at least as many

blank genotypes as expected. For this test we computed the square of null allele

frequency of each locus in each subsample and multiplied it by the subsample size. To

increase statistical power, we summed these expected number of null homozygotes over

all subsamples ($N_{\text{Exp-b}}$) and compared those to the observed number of missing genotypes

for each locus, over all the considered dataset. We then also undertook the regression

$F_{\text{IS}} \sim p_{\text{n}}$, where p_{n} is the global null allele frequency.

Stuttering detection and correction followed the procedure described elsewhere (De

Meeûs et al., 2021; De Meeûs & Noûs, 2022) over all subsamples and for each locus,

using the associated template "TestStutterDioecious-n1000N100-1-10%Stuttering.xlsx"

available at https://zenodo.org/record/7029324, exact binomial tests and adjustment of *p*-

values with Benjamini and Hochberg (BH) procedure with R (command "p.adjust").

Short allele dominance (SAD) was looked for with the correlation method between

F_{IT} and the size of the alleles (Manangwa et al., 2019) using a one-sided (negative

relationship) Spearmans rank correlation test. In case of doubt, we also undertook the

regression $F_{\text{IS}} \sim$ Allele size, weighted with $p_i(1-p_i)$, and where p_i was the frequency of the

allele (De Meeûs et al., 2004). Correlation and regression tests were undertaken with

rcmdr. For SAD, the regression allows to minimize spurious correlations due to rare

alleles. In rcmdr, regression tests are two-sided. To obtain a one-sided *p*-value, when the

slope was negative, we simply halved the two-sided *p*-value, or computed 1-*p*-value/2

otherwise. We labelled the correlation *p*-value p_{cor} and the regression one p_{reg}.

Subdivision

Comparisons of *F*-statistics between different subsamples or between F_{IS} and F_{IT}

was undertaken with Wilcoxon signed rank tests for paired data with rcmdr, using the loci

as pairing factor. Tests were one-sided in the appropriate direction (e.g. $F_{\text{IT}} > F_{\text{IS}}$).

Subdivision was measured and tested with Wright's F_{ST} and the *G* randomization test, as

described above. Genetic differentiation was measured tested between each pair of

cohorts in the same way, except that systematic paired test required BY corrections for

significant *p*-values.

Effective population sizes and effective population densities

We estimated effective population sizes (N_e) with several algorithms and softwares.

The first method is the heterozygote excess method from De Meeûs and Noûs (De Meeûs

& Noûs, 2023), computed for each locus in each cohort, ignoring values below or equal to

0, and averaged across loci, for each subsample (e.g. cohort). The second method was

the linkage disequilibrium based method (Waples & Do, 2010), corrected for missing data

(Peel et al., 2013), implemented by NeEstimator (Do et al., 2014), and assuming random

mating and selecting results for alleles with frequency above or equal to 5%. The third

method was the co-ancestry method (Nomura, 2008), also implemented by NeEstimator.

The fourth method used the averages obtained with four temporal methods. Three were

implemented in NeEstimator: Pollak, Nei and Tajima, and Jorde and Ryman (Nei & Tajima,

1981; Pollak, 1983; Jorde & Ryman, 2007) assuming random mating and selecting results

for alleles with frequency above or equal to 5%. For these three temporal method, we

averaged N_e across the different comparisons (different cohort pairs), ignoring "Infinite"

results The last temporal method was the maximum likelihood temporal method

implemented in MLNe (Wang & Whitlock, 2003). For MLNe, maximum and minimum

values corresponded to the confidence interval outputted by the procedure. We then

computed the average across all temporal methods, weighted by the number of usable

values and averaged minimum and maximum values in the same way. For MLNe, the

weight was the number of cohort pairs (6). The fifth method was the within and between

loci identity probabilities (Vitalis & Couvet, 2001b) with Estim (Vitalis & Couvet, 2001a).

The sixth method was the sibship frequency method (Wang, 2009) with Colony (Jones &

Wang, 2010), assuming female and male polygamy, and inbreeding. For each method, we

computed the average, the minimum and maximum across usable values, and assigned a

weight (number of usable values). For temporal methods, the weight was set to 4 (number

of cohorts). We then obtained the grand average for N_e and minimum and maximum

(minimax) values through an average across methods, weighted by the aforementioned

weights. This methodology obviously provided a rough approximation of N_e and the space

of its possible values, and finding the most accurate way will need specific multi-scenarios

simulation approaches, as discussed elsewhere (De Meeûs & Noûs, 2023).

To compute effective population densities, we considered three different surfaces,

computed with the function "insert a polygon" of Google Earth Pro. The smallest area was

defined by the traps containing flies that were indeed genotyped, as in Figure 1 and Table

1, and labelled S_{Genet}=132 km². The second surface was defined by all traps that captured

at least one tsetse fly in 2011, during the most extensive survey campaign (Kagbadouno et

al., 2012) (see File S1), S_C=224 km². The third area was defined by the limits of the survey

defined before 2011 (see File S1); S_L=629 km². The largest surface was drawn

considering all "mangrove-like" environments in Google Earth Pro, S_{max}=1301 km².

Effective population densities were then computed by dividing the effective population size

by these surfaces. For the sake of comparison, we also computed the densities of

captured flies in 2011 (initial survey, T0), in 20019 and 2020 (see Files S1 for raw data).

We also compute the sex-ratios of captured (c) flies, $SR=N_{c\text{-}males}/N_{c\text{-}females}$ with these data.

The homogeneity of the proportion of females across cohorts was tested with a Fisher

exact test with rcmdr. The evenness of sex-ratios was then tested with a two-sided exact

binomial test with R (command "binom.test").

Finally, maximum distances between the most remote sites defined by each

aforementioned surface was computed with Google Earth Pro with the menu "Add a

trajectory".

Bottleneck signatures

We tried to find bottleneck signatures with the diversity excess method and the

Wilcoxon signed rank test method with Bottleneck (Cornuet & Luikart, 1996), with a

particular interest to subsamples genotyped after the beginning of control (cohorts 61 and

67). We used the three mutation models: IAM, TPM with default options, and SMM.

Significant signatures of a bottleneck can be suspected if the test is highly significant with

the IAM and significant with the TPM at least. Weaker signals tend to be produced in

population with very small effective population sizes (TdM, unpublished simulation results).

Results

Defining the relevant geographic scales for subsample units

There was no significant difference between F_{IS_T}, F_{IS_S} and F_{IS_B} (all p-values>0.4)

(Figure 2). We thus ignored traps, sites and river banks for further analyses, and kept only

the different cohorts (cohorts 0, 10, 61 and 67) as subsample units.

Figure 2: Comparisons of F_{IS} within traps (F_{IS_T}), within sites (ignoring traps) (F_{IS_S}), within
river banks (ignoring sites) (F_{IS_B}), and within each cohort (ignoring river banks) (F_{IS_C}) in
female subsamples of *Glossina palpalis gambiensis* across 11 years in the mangroves of
Boffa, Guinea, for eight loci (crosses of different colors), average (empty circles) and
95%CI (black dashes). The global comparison outputted a *p*-value=0.9663 and none of
paired tests provided a *p*-value<0.4.

Quality testing of the subsamples and loci

Only one locus pair out of 28 appeared in significant LD (less than 4%) (*p*-

value=0.039). It did not remain significant after BY correction. We thus considered all loci

to be statistically independent from each other's.

There was a substantial but variable heterozygote deficit: F_{IS}= 0.143 in

95%CI=[0.076, 0.215] (*p*-value<0.0002) (Figure 3). With r_{SE}=19, and the positive

correlation between F_{IS} and F_{ST} (ρ=0.5988, *p*-value=0.0584), we could suspect that most of

this heterozygote deficit could be explained by null alleles. Nevertheless, the correlation

between F_{IS} and N_b did not seem to confirm this (ρ=0.0952, *p*-value=0.4201). Nonetheless,

this lack of signal resulted from three outlier loci (pGp24, B3 and C102), which displayed

too many missing genotypes as regard to their relatively low F_{IS} (Figure 3). This means

that many of these missing data, at these three loci, probably did not correspond to true

null homozygotes.

Figure 3: Variation of F_{IS} across loci in the four cohorts of *Glossina palpalis gambiensis* from Boffa (Guinea). Averages (black dots), and 95% confidence intervals (black dashes) were computed by 5000 bootstraps over individuals (for each locus) and over loci (All). The two sided *p*-values for significant deviation from panmixia and the number of observed missing genotypes are also provided.

We thus undertook the correlation and the regression $F_{IS} \sim N_b$ without these three loci. The results can be observed in the Figure 4. The correlation became significant and the regression suggested that missing genotypes (null alleles) explains almost all (96%) of F_{IS} variation. The intercept was negative, as expected in small, pangamic dioecious populations ($F_{IS\_0}=-0.0412$ in 95%CI=[-0.1872, 0.1171]).

For null allele frequency estimates, we recoded as homozygous for the null allele (allele coded 999) all missing genotypes but for the three outlier loci. Indeed, for these loci, most missing data (coded "0", and kept as such) probably corresponded to another problem. With these data, estimate of null allele frequencies, and the resulting expected number of missing data over all subsamples fitted very well with the observed ones at each locus (all *p*-values>0.2). Moreover, the regression $F_{IS} \sim p_n$ displayed very good results ($R^2=0.8566$, $F_{IS\_0}=0.012$ in 95%CI=[-0.1951, 0.2143]).

We also undertook the stuttering and SAD analyses. We found a significant stuttering signature for locus A10 only (*p*-value=0.0286). Nevertheless, null alleles explained this locus well enough (Figures 3 and 4) so that we could consider this result as coincidental. Three loci displayed marginally significant signatures of SAD: X55-3 ($p_{cor}=0.0423$, $p_{reg}=0.1551$), pGp24 ($p_{cor}=0.0963$, $p_{reg}=0.0489$), and B3 ($p_{cor}=0.288$, $p_{reg}=0.4841$). Given the weakness of such results, and that these three loci are very well

358 explained by null alleles, we did not consider that any locus was affected by SAD, and that
359 a few null alleles randomly associated with small alleles provided these weak signatures.
360   We detected no Wahlund effect and the significant heterozygote deficits observed
361 were entirely caused by null alleles of frequencies 0.02 to 0.28.
362

363
Figure 4: Regression of F_{IS} by the number of missing genotypes (black crosses and
continuous line) without loci displaying an excess of missing genotypes (pGp24, B3 and
C102) (empty circles) and of the corresponding 95%CI of bootstraps over individuals for
Glossina palpalis gambiensis from Boffa (Guinea) (black dashes and broken lines).
Regression equations, the determination coefficient (R^2) and the result of the correlation
test are also given.
370

Average subdivision was globally not significant with a weak variance across the
different loci: F_{ST}=-0.001 in 95%CI=[-0.004, 0.004] (*p*-value=0.6938). In particular, GPCAG
displayed a negative value: F_{ST}=-0.007 (-0.006 with FreeNA correction) (*p*-value=0.1709),
and allele 219 (here labelled 220, due to probable lag produced by the change in size
calibration on this species), did not increase in frequency after control had begun: 0.094,
0.093, 0.065, and 0.022 for cohorts 0, 10, 61 and 67, respectively.

Subdivision

As demonstrated above, we could not detect any geographic subdivision signature.
Subdivision was globally weak across the different cohorts: average F_{ST}=-0.001 in
95%CI=[-0.004, 0.004] and not significant (*p*-value=0.6938). Moreover, F_{IT}=0.142 in

95%CI=[0.073, 0.218] was not significantly bigger than F_{IS}=0.143 in 95%CI=[0.076, 0.215]

(*p*-value=0.7695).

We also measured and tested subdivision between each pair of cohorts. In the

Figure 5, we can confirm the absence of any subdivision. Moreover, subdivision between

cohorts before control (T0) and after control had begun (TX) did not appear bigger than for

other cohort comparisons.

Figure 5: Subdivision measures (F_{ST}) and test between each pair of cohorts (C0, C10, C61
and C67). The result of the test (*p*-values) and the comparison type between cohorts
392   before control (T0) or after the beginning of control (TX) are also indicated.
393

Effective population sizes and effective population densities

For this section, Estim provided no usable values. For the remaining methods,

results are presented in the Table 2. These values were quite variable, but did not display

particularly smaller values after control (26 and 18 for cohorts 61 and 67 respectively) than

before control (22 and 42 for cohorts 0 and 10 respectively). Nevertheless, such

comparisons could only be made for the heterozygote excess, co-ancestries and sibships

methods.

Table 2: Effective population size, minimum and maximum values estimated with different methods, and over all method (average weighted with "Weight"). Codes for methods are HzEs: heterozygote excess; LD: linkage disequilibrium; CoAn: coancestries.

| | Methods | | | | | |
	HzEs	LD	CoAn	Temporal	Sibship	Over all
Average	14	461	24	15130	44.5	3830
Minimum	5	461	21	404	32	143
Maximum	24	461	28	18957	73	4798
Weight	4	1	3	4	4	

Taking the surface occupied by genotyped flies, the resulting effective population density was $D_{e\text{-Genet}}=8$ in minimax=[1, 17] flies per km². However, given the absence of any population subdivision signature, we considered that this quantity was a very strong overestimate. In the Figure 6, we can see the effective population densities computed with S_C, S_L and S_{max}.

Figure 6: Densities of captured flies (Dc) and corresponding effective population densities (De) while considering the whole surface of Boffa (as defined in the text) occupied by tsetse flies in the HAT focus of Boffa (Guinea), for different years and different effective population size averages: with all methods, or without temporal methods.

There was no significant variations of the sex ratio across years (0.85, 0.79 and 1.03 for 2011, 2019 and 2020 respectively) (p-value=0.3253). Combining all years, captured flies displayed a highly significant female biased sex ratio $SR=0.86$ (p-value<0.0001).

Capturing flies after control had begun, in the same trapping sites, obviously appeared more difficult than before control (Figure 6).

The maximum distances between the most remote points were 19 km, 40 km, 44

km and 62 km for the areas S_{Genet}, S_C, S_L and S_{max} respectively. Depending on the true

area occupied by the population of tsetse flies in Boffa, these may represent the average

dispersal distances from one generation to the next.

Bottleneck signatures

The results of these analyses are given in Table 3, where an absence of any

bottleneck signature can be noticed, in particular after the beginning of control (cohorts 61

and 67).

Table 3: One sided *p*-values for the detection of a genetic signature of a bottleneck with
the three models of mutation (IAM, TPM and SMM) for the different subsamples before the
beginning of vector control (Cohorts 0 and 10) and after the beginning of control (Cohorts
61 and 67).

Cohort	IAM	TPM	SMM
Cohort 0	0.3711	0.9629	0.9981
Cohort 10	0.6289	0.9980	1
Cohort 61	0.5781	0.9863	0.9961
Cohort 67	0.6289	0.6797	1

Discussion

Despite the presence of null alleles, the eight loci studied here in female *G. palpalis*

gambiensis converged with the existence of a large pangamic population of this species

occupying the HAT focus of Boffa. This was confirmed by the absence of any Wahlund

effect when pooling all individuals from all the sampling area into one subsample in each

cohort. The regression of the F_{IS} with null alleles or number of missing data, with a

determination coefficient close to unity and a negative intercept confirmed this perception.

Subdivision tests also confirmed a free migration of tsetse flies in the area defined by this

tsetse population. Taking the total surface where flies could be captured and the total

surface of the survey, as the most realistic values, the corresponding dispersal distance

would be between 40 and 44 km per generation. The constant shading, high humidity and

wind conditions of the particular continuously favorable mangrove ecosystem probably

explain such important values (Courtin et al., 2010; Courtin & Kagbadouno, 2011; Courtin

et al., 2015). Within these surfaces, the effective population density could be considered to

vary between 3 and 6 individuals per km², in minimax≈[0.1, 8]. These range within the

smallest values as compared to what can be found elsewhere (De Meeûs et al., 2019).

This is even worse if we consider that, here, we could use temporal methods for estimating

effective population sizes. Temporal methods are almost never available in the literature,

but provided very high values in the present study. Without temporal methods, densities

would have fallen to very small values: between 0.05 and 0.1 individuals per km² only, for

the two surfaces respectively.

We observed a significant female biased *SR* in this population with small effective

population densities. In a recent study of *G. fuscipes fuscipes* from Chad (Ravel et al.,

2023), authors found a negative correlation between *SR* and densities. Scattered

distribution of hosts, leading to smaller effective population densities of the flies, would

impose to these insects to travel more to find blood meals. Because of larval feeding,

female tsetse would need to feed more often than males and be captured more often than

those in such kind of ecological frameworks. On the contrary, abundance of hosts would

maintain dense populations of tsetse flies, with no need to spend time seeking for blood

meals, especially so for females, while males still need looking for mates.

The absence of any significant genetic differentiation between subsamples before

and after the beginning of control, especially at locus GPCAG, of any strong (if any) effect

on effective population sizes, and the total absence of any genetic signature of a

bottleneck, testify of an absence of consequences of the VCC on the genetic structure of

this tsetse population, at the scale of the focus. Nevertheless, the dramatic drop in

captured flies after the VCC had begun, indicated an efficient protection of the human

population against tsetse bites around the targeted zones. The drop in tsetse bites and

HAT prevalence and incidence in selected controlled zones has already proved such a

protection (Courtin et al., 2015). It also indicates that in the areas targeted by vector

control tsetse densities would probably regain initial levels rapidly in case of scaling back.

The fact that a significant decrease in HAT prevalence was observed

(https://apps.who.int/neglected_diseases/ntddata/hat/hat.html) also advocates for the

benefic effects of the VCC. Finally, the absence of impact on the targeted population at the

scale of the whole mangrove of Boffa, suggests that VCC as it was undertaken, is efficient

at protecting human populations, but does not significantly affect the biodiversity of this

particular ecosystem.

In conclusion, due to the difficulties to access to all tsetse-infested zones, where

multiple hosts are available for these blood feeding insects, maintaining control measures

in zones frequented by human populations seems a reasonable measure, in order to

protect these populations from tsetse bites and subsequent transmission of HAT. Given

that the existence of hidden human and/or animal reservoirs can be suspected, such a

measure appears mandatory for now and a close future, before more research results

shed more light on the eco-epidemiology of this deadly disease in mangrove ecosystems.

Acknowledgements

Funding

This work was funded by the Bill & Melinda Gates Foundation

(http://www.gatesfoundation.org, grant agreement INV-001785)" and the JEAI RECIT

(#400982/00) of the IRD.

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial

conflicts of interest in relation to the content of the article

Author contributions

Moise S. Kagbadouno: data collection, data analyses and manuscript correction.

Modou Séré: genotyping, data analyses and manuscript correction.

Adeline Ségard: genotyping and manuscript correction.

Abdoulaye Dansy Camara: data collection and manuscript correction.

Mamadou Camara: supervision and manuscript correction.

Bruno Bucheton: supervision, data collection and manuscript correction.

Jean-Mathieu Bart: supervision, data collection and manuscript correction.

Fabrice Courtin: supervision, data collection and manuscript correction.

Thierry de Meeûs: supervision, data analyses, writing of the manuscript and design of

figures.

Sophie Ravel: supervision, genotyping and manuscript correction.

Data, scripts, code, and supplementary information availability

Raw data are available at https://zenodo.org/record/8181166. For data analyses, all

R scripts used are inserted in the main text. For other analyses we used facilities without

the need of scripts (e.g. click and point programs or packages).

References

Baker MD, Krafsur ES (2001) Identification and properties of microsatellite markers in tsetse flies *Glossina morsitans* sensu lato (Diptera: Glossinidae). *Molecular Ecology Notes,* **1**, 234-236. https://doi.org/https://doi.org/10.1046/j.1471-8278.2001.00087.x

Balloux F (2004) Heterozygote excess in small populations and the heterozygote-excess effective population size. *Evolution,* **58**, 1891-1900. https://doi.org/https://doi.org/10.1554/03-692

Belkhir K, Borsa P, Chikhi L, Raufaste N, Bonhomme F (2004) GENETIX 4.05, logiciel sous Windows TM pour la génétique des populations. Laboratoire Génome, Populations, Interactions, CNRS UMR 5000, Université de Montpellier II, Montpellier (France). https://doi.org/https://kimura.univ-montp2.fr/genetix/

Benjamini Y, Yekutieli D (2001) The control of the false discovery rate in multiple testing under dependency. *The Annals of Statistics,* **29**, 1165–1188. https://doi.org/ https://doi.org/10.1214/aos/1013699998

Berté D, De Meeus T, Kaba D, Séré M, Djohan V, Courtin F, N'Djetchi KM, Koffi M, Jamonneau V, Ta BTD, Solano P, N'Goran EK, Ravel S (2019) Population genetics of *Glossina palpalis palpalis* in sleeping sickness foci of Côte d'Ivoire before and after vector control. *Infection Genetics and Evolution,* **75**, 103963. https://doi.org/https://doi.org/10.1016/j.meegid.2019.103963

Bouyer J, Balenghien T, Ravel S, Vial L, Sidibé I, Thévenon S, Solano P, De Meeûs T (2009) Population sizes and dispersal pattern of tsetse flies: rolling on the river? *Molecular Ecology,* **18**, 2787-2797. https://doi.org/ https://doi.org/10.1111/j.1365-294X.2009.04233.x

Bouyer J, Dicko AH, Cecchi G, Ravel S, Guerrini L, Solano P, Vreysen MJB, De Meeûs T, Lancelot R (2015) Mapping landscape friction to locate isolated tsetse populations candidate for elimination. *Proceedings of the National Academy of Sciences of the*

United States of America, **112**, 14575–14580.

https://doi.org/https://doi.org/10.1073/pnas.1516778112

Bouyer J, Ravel S, Dujardin JP, De Meeûs T, Vial L, Thévenon S, Guerrini L, Sidibe I,

Solano P (2007) Population structuring of *Glossina palpalis gambiensis* (Diptera:

Glossinidae) according to landscape fragmentation in the Mouhoun river, Burkina

Faso. *Journal of Medical Entomology,* **44**, 788-795.

https://doi.org/https://doi.org/10.1603/0022-2585(2007)44[788:psogpg]2.0.co;2

Bucheton B, Kheir MM, El-Safi SH, Hammad A, Mergani A, Mary C, Abel L, Dessein A

(2002) The interplay between environmental and host factors during an outbreak of

visceral leishmaniasis in eastern Sudan. *Microbes and Infection,* **4**, 1449-1457.

https://doi.org/https://doi.org/10.1016/s1286-4579(02)00027-8

Büscher P, Bart JM, Boelaert M, Bucheton B, Cecchi G, Chitnis N, Courtin D, Figueiredo

LM, Franco JR, Grébaut P, Hasker E, Ilboudo H, Jamonneau V, Koffi M, Lejon V,

MacLeod A, Masumu J, Matovu E, Mattioli R, Noyes H, Picado A, Rock KS,

Rotureau B, Simo G, Thévenon S, Trindade S, Truc P, Van Reet N (2018) Do

cryptic reservoirs threaten gambiense-sleeping sickness elimination? *Trends in*

Parasitology, **34**, 197-207. https://doi.org/10.1016/j.pt.2017.11.008

Camara M, Caro-Riano H, Ravel S, Dujardin JP, Hervouet JP, De Meeûs T, Kagbadouno

MS, Bouyer J, Solano P (2006) Genetic and morphometric evidence for population

isolation of *Glossina palpalis gambiensis* (Diptera : Glossinidae) on the Loos

islands, Guinea. *Journal of Medical Entomology,* **43**, 853-860.

https://doi.org/https://doi.org/10.1603/0022-2585(2006)43[853:gamefp]2.0.co;2

Chapuis MP, Estoup A (2007) Microsatellite null alleles and estimation of population

differentiation. *Molecular Biology and Evolution,* **24**, 621-631.

https://doi.org/10.1093/molbev/msl191

Coombs JA, Letcher BH, Nislow KH (2008) CREATE: a software to create input files from

diploid genotypic data for 52 genetic software programs. *Molecular Ecology*

Resources, **8**, 578–580. https://doi.org/https://doi.org/10.1111/j.1471-

8286.2007.02036.x

Cornuet JM, Luikart G (1996) Description and power analysis of two tests for detecting

recent population bottlenecks from allele frequency data. *Genetics,* **144**, 2001-2014.

https://doi.org/https://doi.org/10.1093/genetics/144.4.2001

Courtin F, Camara M, Rayaisse JB, Kagbadouno M, Dama E, Camara O, Traore IS,

Rouamba J, Peylhard M, Somda MB, Leno M, Lehane MJ, Torr SJ, Solano P,

Jamonneau V, Bucheton B (2015) Reducing human-tsetse contact significantly

enhances the efficacy of sleeping sickness active screening campaigns: a

promising result in the context of elimination. *PLoS Neglected Tropical Diseases,* **9**.

https://doi.org/10.1371/journal.pntd.0003727

Courtin F, Jamonneau V, Camara M, Camara O, Coulibaly B, Diarra A, Solano P,

Bucheton B (2010) A geographical approach to identify sleeping sickness risk

factors in a mangrove ecosystem. *Tropical Medicine & International Health,* **15**,

881-889. https://doi.org/10.1111/j.1365-3156.2010.02559.x

Courtin F, Kagbadouno M (2011) Peuplements, mobilités et paysages en zone de

mangrove guinéenne : le cas de la baie de Sangaréah (Guinée). *Les Cahiers*

d'Outre-Mer, **256**, 453-466. https://doi.org/10.4000/com.6358

De Meeûs T (2018) Revisiting F_{IS}, F_{ST}, Wahlund effects, and Null alleles. *Journal of*

Heredity, **109**, 446-456. https://doi.org/10.1093/jhered/esx106

De Meeûs T, Chan CT, Ludwig JM, Tsao JI, Patel J, Bhagatwala J, Beati L (2021)

Deceptive combined effects of short allele dominance and stuttering: an example

with *Ixodes scapularis*, the main vector of Lyme disease in the U.S.A. *Peer*

Community Journal, **1**, e40. https://doi.org/https://doi.org/10.24072/pcjournal.34

De Meeûs T, Guégan JF, Teriokhin AT (2009) MultiTest V.1.2, a program to binomially

combine independent tests and performance comparison with other related

methods on proportional data. *BMC Bioinformatics,* **10**, 443.

https://doi.org/https://doi.org/10.1186/1471-2105-10-443

De Meeûs T, Humair PF, Grunau C, Delaye C, Renaud F (2004) Non-Mendelian

transmission of alleles at microsatellite loci: an example in *Ixodes ricinus*, the vector

of Lyme disease. *International Journal for Parasitology,* **34**, 943-950.

https://doi.org/https://doi.org/10.1016/j.ijpara.2004.04.006

De Meeûs T, Lehmann La, Balloux F (2006) Molecular epidemiology of clonal diploids: A

quick overview and a short DIY (do it yourself) notice. *Infection Genetics and*

Evolution, **6**, 163-170. https://doi.org/https://doi.org/10.1016/j.meegid.2005.02.004

De Meeûs T, McCoy KD, Prugnolle F, Chevillon C, Durand P, Hurtrez-Boussès S, Renaud

F (2007) Population genetics and molecular epidemiology or how to "débusquer la

bête". *Infection Genetics and Evolution,* **7**, 308-332.

https://doi.org/https://doi.org/10.1016/j.meegid.2006.07.003

De Meeûs T, Noûs C (2022) A simple procedure to detect, test for the presence of

stuttering, and cure stuttered data with spreadsheet programs. *Peer Community*

Journal, **2**, e52. https://doi.org/https://doi.org/10.24072/pcjournal.165

De Meeûs T, Noûs C (2023) A new and almost perfectly accurate approximation of the

eigenvalue effective population size of a dioecious population: comparisons with

other estimates and detailed proofs *Peer Community Journal,* **3**, e51.

https://doi.org/https://doi.org/10.24072/pcjournal.280

De Meeûs T, Ravel S, Solano P, Bouyer J (2019) Negative density dependent dispersal in

tsetse flies: a risk for control campaigns? *Trends in Parasitology,* **35**, 615-621.

https://doi.org/https://doi.org/10.1016/j.pt.2019.05.007

Dempster AP, Laird NM, Rubin DB (1977) Maximum likelihood from incomplete data via

the EM algorithm. *Journal of the Royal Statistical Society Series B,* **39**, 1–38.

https://doi.org/http://links.jstor.org/sici?sici=0035-

9246%281977%2939%3A1%3C1%3AMLFIDV%3E2.0.CO%3B2-Z

Do C, Waples RS, Peel D, Macbeth GM, Tillett BJ, Ovenden JR (2014) NeEstimator v2:

re-implementation of software for the estimation of contemporary effective

population size (N_e) from genetic data. *Molecular Ecology Resources,* **14**, 209-214.

https://doi.org/https://doi.org/10.1111/1755-0998.12157

Fox J (2005) The R commander: a basic statistics graphical user interface to R. *Journal of*

Statistical Software, **14**, 1–42. https://doi.org/https://doi.org/10.18637/jss.v014.i09

Fox J (2007) Extending the R commander by "plug in" packages. *R News,* **7**, 46–52.

https://doi.org/https://stat.ethz.ch/pipermail/r-

help/attachments/20071101/3603125e/attachment.pdf

Goudet J (1995) FSTAT (Version 1.2): A computer program to calculate F-statistics.

Journal of Heredity, **86**, 485-486.

https://doi.org/https://doi.org/10.1093/oxfordjournals.jhered.a111627

Goudet J (2003) Fstat (ver. 2.9.4), a program to estimate and test population genetics

parameters. Available at http://www.t-de-meeus.fr/Programs/Fstat294.zip, Updated

from Goudet (1995).

Goudet J, De Meeûs T, Day AJ, Gliddon CJ (1994) The different levels of population

structuring of dogwhelks, *Nucella lapillus*, along the south Devon coast. In: *Genetics*

and Evolution of Aquatic Organisms (ed. Beaumont A), pp. 81-95. Chapman and

Hall, London.

Goudet J, Raymond M, De Meeûs T, Rousset F (1996) Testing differentiation in diploid

populations. *Genetics,* **144**, 1933-1940.

https://doi.org/https://doi.org/10.1093/genetics/144.4.1933

Holmes P (2014) First WHO meeting of stakeholders on elimination of gambiense human

african trypanosomiasis. *PLoS Neglected Tropical Diseases,* **8**, e3244.

https://doi.org/https://doi.org/10.1371/journal.pntd.0003244

Jamonneau V, Truc P, Grébaut P, Herder S, Ravel S, Solano P, De Meeus T (2019)

Trypanosoma brucei gambiense group 2: the unusual suspect. *Trends in*

Parasitology, **35**, 983-995. https://doi.org/https://doi.org/10.1016/j.pt.2019.09.002

Jones OR, Wang JL (2010) COLONY: a program for parentage and sibship inference from

multilocus genotype data. *Molecular Ecology Resources,* **10**, 551-555.

https://doi.org/10.1111/j.1755-0998.2009.02787.x

Jorde PE, Ryman N (2007) Unbiased estimator for genetic drift and effective population

size. *Genetics,* **177**, 927-935.

https://doi.org/https://doi.org/10.1534/genetics.107.075481

Kagbadouno MS, Camara M, Rouamba J, Rayaisse JB, Traoré IS, Camara O,

Onikoyamou MF, Courtin F, Ravel S, De Meeûs T, Bucheton B, Jamonneau V,

Solano P (2012) Epidemiology of sleeping sickness in boffa (Guinea): where are the

trypanosomes? *PLoS Neglected Tropical Diseases,* **6**, e1949.

https://doi.org/10.1371/journal.pntd.0001949

Koffi M, De Meeûs T, Bucheton B, Solano P, Camara M, Kaba D, Cuny G, Ayala FJ,

Jamonneau V (2009) Population genetics of *Trypanosoma brucei gambiense*, the

agent of sleeping sickness in Western Africa. *Proceedings of the National Academy*

of Sciences of the United States of America, **106**, 209-214.

https://doi.org/https://doi.org/10.1073/pnas.0811080106

Koffi M, De Meeûs T, Séré M, Bucheton B, Simo G, Njiokou F, Salim B, Kaboré J,

MacLeod A, Camara M, Solano P, Belem AMG, Jamonneau V (2015) Population

genetics and reproductive strategies of african trypanosomes : revisiting available

published data. *PLoS Neglected Tropical Diseases,* **9**, e0003985.

https://doi.org/https://doi.org/10.1371/journal.pntd.0003985

Luna C, Bonizzoni MB, Cheng Q, Aksoy S, Zheng L (2001) Microsatellite polymorphism in

the tsetse fies (Diptera: Glossinidae). *Journal of Medical Entomology,* **38**, 376–381

Manangwa O, De Meeûs T, Grébaut P, Segard A, Byamungu M, Ravel S (2019) Detecting

Wahlund effects together with amplification problems : cryptic species, null alleles

and short allele dominance in Glossina pallidipes populations from Tanzania.

Molecular Ecology Resources, **19**, 757-772.

https://doi.org/https://doi.org/10.1111/1755-0998.12989

Melachio T, T. T., Simo G, Ravel S, De Meeûs T, Causse S, Solano P, Lutumba P,

Asonganyi T, Njiokou F (2011) Population genetics of Glossina palpalis palpalis

from central African sleeping sickness foci. *Parasites and Vectors,* **4**, 140.

https://doi.org/10.1186/1756-3305-4-140

Nei M, Tajima F (1981) Genetic drift and estimation of effective population size. *Genetics,*

98, 625-640. https://doi.org/https://doi.org/10.1093/genetics/98.3.625

Nomura T (2008) Estimation of effective number of breeders from molecular coancestry of

single cohort sample. *Evolutionary Applications,* **1**, 462-474.

https://doi.org/https://doi.org/10.1111%2Fj.1752-4571.2008.00015.x

Peel D, Waples RS, Macbeth GM, Do C, Ovenden JR (2013) Accounting for missing data

in the estimation of contemporary genetic effective population size (N_e). *Molecular*

Ecology Resources, **13**, 243-253. https://doi.org/https://doi.org/10.1111/1755-

0998.12049

Pollak E (1983) A new method for estimating the effective population size from allele

frequency changes. *Genetics,* **104**, 531-548.

https://doi.org/https://doi.org/10.1093/genetics/104.3.531

Pudovkin AI, Zaykin DV, Hedgecock D (1996) On the potential for estimating the effective

number of breeders from heterozygote excess in progeny. *Genetics,* **144**, 383-387.

https://doi.org/https://doi.org/10.1093/genetics/144.1.383

R-Core-Team (2022) R: A Language and Environment for Statistical Computing (Version

4.2.2 (2022-10-31 ucrt)).

Ravel S, Mahamat MH, Ségard A, Argiles-Herrero R, Bouyer J, Rayaisse J-B, Solano P,

Mollo BG, Pèka M, Darnas J, Belem AMG, Yoni W, Noûs C, De Meûs T (2023)

Population genetics of *Glossina fuscipes fuscipes* from southern Chad. *Peer*

Community Journal, **3**, e31. https://doi.org/https://doi.org/10.24072/pcjournal.257

Ravel S, Sere M, Manangwa O, Kagbadouno M, Mahamat MH, Shereni W, Okeyo WA,

Argiles-Herrero R, De Meeûs T (2020) Developing and quality testing of

microsatellite loci for four species of *Glossina*. *Infection Genetics and Evolution,* **85**,

104515. https://doi.org/https://doi.org/10.1016/j.meegid.2020.104515

Robertson A (1965) The interpretation of genotypic ratios in domestic animal populations.

Animal Production, **7**, 319-324.

https://doi.org/https://doi.org/10.1017/S0003356100025770

Ségard A, Romero A, Ravel S, Truc P, Dobigny G, Gauthier P, Etougbetche J, Dossou H-

J, Badou S, Houéménou G, Morand S, Chaisiri K, Noûs C, De Meeûs T (2022)

Development of nine microsatellite loci for *Trypanosoma lewisi*, a potential human

pathogen in Western Africa and South-East Asia, and preliminary population

genetics analyses. *Peer Community Journal,* **2**, e69.

https://doi.org/https://doi.org/10.24072/pcjournal.188

Simarro PP, Cecchi G, Franco JR, Paone M, Diarra A, Priotto G, Mattioli RC, Jannin JG

(2015) Monitoring the Progress towards the Elimination of Gambiense Human

African Trypanosomiasis. *PLoS Neglected Tropical Diseases,* **9**, e0003785.

https://doi.org/10.1371/journal.pntd.0003785

Solano P, Duvallet G, Dumas V, Cuisance D, Cuny G (1997) Microsatellite markers for

genetic population studies in *Glossina palpalis* (Diptera: Glossinidae). *Acta Tropica,*

65, 175-180

Solano P, Ravel S, Bouyer J, Camara M, Kagbadouno MS, Dyer N, Gardes L, Herault D,

Donnelly MJ, De Meeûs T (2009) The population structure of *Glossina palpalis*

gambiensis from island and continental locations in coastal Guinea. *PLoS*

Neglected Tropical Diseases, **3**, e392. https://doi.org/10.1371/journal.pntd.0000392

Vitalis R, Couvet D (2001a) ESTIM 1.0: a computer program to infer population

parameters from one- and two-locus gene identity probabilities. *Molecular Ecology*

Notes, **1**, 354-356. https://doi.org/https://doi.org/10.1046/j.1471-8278.2001.00086.x

Vitalis R, Couvet D (2001b) Estimation of effective population size and migration rate from

one- and two-locus identity measures. *Genetics,* **157**, 911-925.

https://doi.org/https://doi.org/10.1093/genetics/157.2.911

Wang JL (2009) A new method for estimating effective population sizes from a single

sample of multilocus genotypes. *Molecular Ecology,* **18**, 2148-2164.

https://doi.org/https://doi.org/10.1111/j.1365-294X.2009.04175.x

Wang JL, Whitlock MC (2003) Estimating effective population size and migration rates

from genetic samples over space and time. *Genetics,* **163**, 429-446.

https://doi.org/https://doi.org/10.1093/genetics/163.1.429

Waples RS, Do C (2010) Linkage disequilibrium estimates of contemporary N_e using

highly variable genetic markers: a largely untapped resource for applied

conservation and evolution. *Evolutionary Applications,* **3**, 244-262.

https://doi.org/https://doi.org/10.1111/j.1752-4571.2009.00104.x

Weir BS, Cockerham CC (1984) Estimating F-statistics for the analysis of population

structure. *Evolution,* **38**, 1358-1370. https://doi.org/https://doi.org/10.1111/j.1558-

5646.1984.tb05657.x

Weir W, Capewell P, Foth B, Clucas C, Pountain A, Steketee P, Veitch N, Koffi M, De

Meeûs T, Kaboré J, Camara M, Cooper A, Tait A, Jamonneau V, Bucheton B,

Berriman M, MacLeod A (2016) Population genomics reveals the origin and asexual

evolution of human infective trypanosomes. *eLife,* **5**, e11473.

https://doi.org/https://doi.org/10.7554%2FeLife.11473

Williams BG (1990) Tsetse fly (Diptera: Glossinidae) population dynamics and the

estimation of mortality rates from life-table data. *Bulletin of Entomological Research,*

80, 479-485. https://doi.org/http://dx.doi.org/10.1017/S0007485300050756

Wright S (1965) The interpretation of population structure by F-statistics with special

regard to system of mating. *Evolution,* **19**, 395-420.

https://doi.org/https://doi.org/10.1111/j.1558-5646.1965.tb01731.x

766
767

Appendix

Appendix 1: Information on unpublished microsatellite loci

Table A1: Information on microsatellite loci provided by A. Robinson (Insect Pest Control
Sub-program, Joint Food and Agriculture Organization of the United Nations/International
Atomic Energy Agency Program of Nuclear Techniques in Food and Agriculture) used in
the present study. Name of primers motif, annealing temperature (T_ann in °C), the range
of sizes of amplified fragments, and primers' sequences are indicated.

Locus	Primers (5'-3')		Motif	T_ann	Size range
			(CA)	54.5	190
A10	A10F	GCAACGCCAAGTGAAATAAAG			
	A10R	TACTGGGCTCGCGTACATAAT			
			(GA)	55	181-195
B3	B3F	TTCGCTTTTGTCAGAAGTG			
	B3R	TCCCAGACGGTTCATATTAC			
			(GA)	52	171-227
XB104	B104F	ACGATTGTTCTTCTTCATTCC			
	B104R	AAGAAGCCAAACTTTCAACGCTCGC			
			(TGA)	52	257-278
C102	C102F	CGTTAAATTGTCACGATGAAG			
	C102R	AGCACATTTTTGGTACATTTATACC			

778 Touch-down PCR reactions were the same as in (Ségard et al., 2022):

Santani traps 1, 5
Santani traps 2, 4
Bakyia trap 3
Bakyia traps 1, 2
Dobire traps 6, 7
Guiapet trap 1
Guiapet traps 2, 3, 4
Yangoya traps 1, 5, 6, 7

0   4   8 km

TBG
PATALA