Early-life stress triggers long-lasting organismal resilience and longevity via tetraspanin

Wei I. Jiang[1], Henry De Belly[1,2], Bingying Wang[1], Andrew Wong[1], Minseo Kim[1], Fiona Oh[1],

Jason DeGeorge[1], Xinya Huang[3], Shouhong Guang[3], Orion D. Weiner[1,2], Dengke K. Ma[1,4,5,*]

[1]Cardiovascular Research Institute, University of California, San Francisco, San Francisco, USA

[2]Department of Biochemistry and Biophysics, University of California, San Francisco, San

Francisco, CA, USA

[3]The USTC RNA Institute, Division of Life Sciences and Medicine, Department of Obstetrics and

Gynecology, The First Affiliated Hospital of USTC, Biomedical Sciences and Health Laboratory

of Anhui Province, University of Science and Technology of China, Hefei, Anhui, China

[4]Department of Physiology, University of California, San Francisco, San Francisco, CA, USA

[5]Innovative Genomics Institute, University of California, Berkeley, Berkeley, CA, USA.

* Correspondence: Dengke.Ma@ucsf.edu (D.K.M.)

Abstract

Early-life stress experiences can produce lasting impacts on organismal adaptation and fitness.

How transient stress elicits memory-like physiological effects is largely unknown. Here we show

that early-life thermal stress strongly up-regulates *tsp-1*, a gene encoding the conserved

transmembrane tetraspanin in *C. elegans*. TSP-1 forms prominent multimers and stable web-

like structures critical for membrane barrier functions in adults and during aging. Up-regulation

of TSP-1 is long lasting even after transient early-life stress. Such regulation requires CBP-1, a

histone acetyl-transference that facilitates initial *tsp-1* transcription. Tetraspanin webs form

regular membrane structures and mediate resilience-promoting effects of early-life thermal

stress. Gain-of-function TSP-1 confers striking *C. elegans* longevity extension and thermal

resilience in human cells. Together, our results reveal a cellular mechanism by which early-life

thermal stress produces long-lasting memory-like impact on organismal resilience and longevity.

Epidemiological and clinical evidence in humans shows that life stress of various forms

can exert profound lasting impacts on mental and physical health outcomes and lifespans (*1–4*).

For example, severe nutritional stress early in life can incur lifespan costs while psychosocial

early-life stress increases vulnerability to psychiatric disorders (*5–7*). By contrast, stresses in

milder physiological forms, such as fasting with adequate nutrition or thermal stimuli via sauna

exposure, are associated with long-lasting health benefits (*8, 9*). Transient periods of stress can

induce persistent changes in the endocrine response, epigenetic regulation of gene expression

and plasticity changes in various organs (*10–12*). However, the underlying molecular and

cellular mechanisms by which transient early-life stress can produce memory-like physiological

effects remain poorly understood. Additionally, it is challenging to establish causal relationship

between putative cellular mechanisms and long-term health outcomes in human studies.

In this work, we use a robust thermal stress paradigm in *C. elegans* to identify causal

mechanisms of how transient stress may exert lasting impacts on organismal resilience and

longevity. We show that transient heat exposure during late larval development activates the

gene *tsp-1*, which encodes a *C. elegans* homolog of the evolutionarily conserved tetraspanin

protein family. TSP-1 proteins form tetraspanin web-like structures and are essential for

maintaining membrane permeability, barrier functions and heat-induced organismal resilience

and longevity. Initial *tsp-1* induction by heat requires a histone acetyltransferance CBP-1, yet

surprisingly, leads to sustained up-regulation of TSP-1 protein but not mRNA abundance. We

propose that a tetraspanin web-like structure through stabilized TSP-1 multimerization mediates

long-lasting organismal impact triggered by transient mild thermal stress in early life.

Using RNA-seq, we previously identified *tsp-1* as one of the genes highly up-regulated

by cold-warming (CW) stress (transient exposure to 4 or -20 ^{o}C followed by recovery at 20 ^{o}C) in

C. elegans (*13–15*). We generated translational reporters by fusing GFP with endogenous

regulatory DNA sequences (promoter and coding regions) for many of the CW stress-inducible

genes to monitor their induction kinetics and protein localization in live *C. elegans* during

development and adult responses to stresses. We focus on *tsp-1* in this study as the

constructed *tsp-1*p::*tsp-1*::GFP translational reporter shows robust and striking induction at L4

stages by not only CW stresses, but also mild heat stress at 25 °C, more so at 28 °C, less so at

20 °C or above 32 °C (**Fig. 1A, B; fig. S1**). Induction by 28 °C is heat-duration dependent (**fig. S1**) and does not require HSF-1 (heat shock factor) (**fig. S2**) that mediates the canonical heat

shock response in *C. elegans* (*16–20*). By contrast, the heat-shock chaperon gene *hsp-16* is

also induced by 28 °C and requires HSF-1 for heat induction (**fig. S2**). These results identify a

previously unknown, HSF-independent, thermal heat induction of TSP-1 in *C. elegans*.

TSP-1 is a *C. elegans* homolog of the evolutionarily conserved tetraspanin protein family

in eukaryotes (*21–23*) predicted to form 4-transmembrane α-helical segments and two

extracellular loop domains (**Fig. 1C**). We fused GFP to the C-terminus of TSP-1, leaving the N-

terminus of TSP-1 intact, which contains the first transmembrane helix targeting TSP-1 to

plasma membranes (**Fig. 1C**). Tetraspanins are versatile scaffolding transmembrane proteins,

specific members of which can interact with themselves to form multimers and control the

spatial organization of membrane lipids and proteins in networks called tetraspanin webs or

tetraspanin-enriched microdomains (*21–23*). We used sodium dodecyl-sulfate polyacrylamide

gel electrophoresis (SDS-PAGE) and native PAGE to analyze the thermal induction and

multimerization of TSP-1::GFP (**Fig. 1D**). Under baseline condition at 20 °C, TSP-1::GFP

formed protein species corresponding to predicted monomers, SDS-resistant dimers and trimers

(in molecular weights). Heat at 28 °C for 24 hrs drastically increased all TSP-1::GFP species in

SDS-PAGE, and native PAGE further revealed formation of high molecular-weight species

larger than dimers and trimers (**Fig. 1D**). Confocal microscopy revealed that TSP-1::GFP was

highly enriched along the apical membrane of intestinal cells (**Fig. 1E**). Heat at 28 °C for 72 hrs

or a transient period of 24 hrs starting from L4 markedly increased the abundance of TSP-

1::GFP, forming striking tetraspanin web-like structures discernable at high resolution (**Fig. 1E**).

Such structures were not caused by temperature effect on the tag GFP *per se* since intestinal

apical membrane-tethered GFP alone did not show such a pattern (**fig. S3**). In addition, *tsp-1*

endogenously tagged with a worm-codon optimized fluorescent protein wrmScarlet by

CRISPR/Cas showed similar up-regulation by heat at 28 ^{o}C and formation of tetraspanin web-

like structures (**fig. S4**). Compared with multi-copy transgenic TSP-1::GFP, endogenously

tagged TSP-1::wrmScarlet exhibited overall weaker fluorescent intensity and enabled us better

to resolve finer tetraspanin webs with regular lattice-like structures (**fig. S4F-I**).

Tetraspanin has been implicated in numerous biological processes, yet its precise

cellular functions remain largely unknown. We next sought to determine normal TSP-1

expression pattern under non-stress conditions and its adult physiological function during aging

in *C. elegans*. At day 1 (young adult), 5 (adult), and 9 (old adult) post L4 stages, TSP-1::GFP

exhibited a progressive increase in abundance (**Fig. 2A-E**). Cultivation at 25 ^{o}C accelerated the

time-dependent increase of TSP-1::GFP abundance, albeit to a lesser degree than a transient

24 hrs period at 28 ^{o}C (**Fig. 2A-E, Fig. 1E**) (28 ^{o}C represents a harsh stress precluding chronic

analyses of TSP-1::GFP at day 5 and 9 because of organismal death by prolonged exposure,

see below). The striking tetraspanin web-like structure formed by TSP-1::GFP (**Fig. 1E**)

prompted us to test the role of TSP-1 in maintaining membrane integrity. We developed a

fluorescein-based assay modified from our previous studies (*13*) to measure the acute barrier

function or permeability of intestinal membranes in *C. elegans* (**Fig. 2F**). In wild-type animals,

short (10 min) incubation with fluorescein did not result in detectable fluorescence signals in the

intestine (**Fig. 2G**). By contrast, the same procedure led to dramatic accumulation of fluorescein

in the intestine of *tsp-1* deficient animals (**Fig. 2G-2J**). Such a difference was notable at the L4

stage and particularly prominent in adults at the day 5 post L4 (**Fig. 2J**). We observed similar

defects in membrane barrier functions using RNA interference (RNAi) against two different

coding regions of *tsp-1* (**fig. S5**). These results indicate that TSP-1 up-regulation during aging or

by thermal stress may physiologically maintain intestinal barrier functions.

We next characterized how early-life thermal stress (ELTS, 28 $^{\circ}$C for 24 hrs starting from

L4 stages) may trigger lasting TSP-1::GFP expression. In control animals at 20 $^{\circ}$C, TSP-1::GFP

remained at low baseline levels largely unaltered for the first 48 hrs and became slightly

elevated at 72 hrs (**Fig. 3A**). By contrast, ELTS (28 $^{\circ}$C for 24 hrs at L4) triggered robust up-

regulation of TSP-1::GFP abundance at 24 hrs post L4, which remained still high at 20 $^{\circ}$C even

for another 24 and 48 hrs after the initial ELTS (**Fig. 3A**). Chronic exposure to 28 $^{\circ}$C starting at

L4 induced stronger expression of TSP-1::GFP for 48 hrs than 24 hrs, but reached peak levels

at 72 hrs (**Figs. 3B, C**). We took advantage of the robust TSP-1::GFP activation by ELTS and

performed RNAi screens for genes that are required for TSP-1::GFP up-regulation. Given the

importance of transcription factors and histone modifying enzymes in epigenetic gene regulation

(*24–26*), we assembled a customized library of RNAi clones targeting >100 genes with

adequate expression in the intestine (transcript per million >2) and that encode proteins

including stress-responding transcription factors and chromatin/epigenetic regulators (**Fig. 3D,

table S1**). From such candidate screens, using RNAi vector only as negative control and *tsp-1*

RNAi as a positive control, we found that RNAi against only one hit, *cbp-1*, robustly prevented

TSP-1::GFP up-regulation by ELTS (**Fig. 3E, 3F**). *cbp-1* encodes the *C. elegans* ortholog of

histone acetyltransferase p300/CBP that promotes gene transcription (*27–35*). RNAi against

many of the known genes encoding heat-responding transcription factor, including *hsf-1*, *hsf-2*,

hif-1 and *nhr-49*, did not block TSP-1::GFP up-regulation (**table S1**). Although TSP-1::GFP

remained high for 24 and 48 hrs at 20 $^{\circ}$C post initial ELTS (**Fig. 3C**), qRT-PCR measurements

revealed that the up-regulation of *tsp-1* mRNA transcripts by 28 $^{\circ}$C was transient but not

sustained, and required CBP-1 (**Fig. 3F**). These results show that transient ELTS can trigger

lasting up-regulation of TSP-1 abundance, which requires initial CBP-1-dependent *tsp-1*

transcription but remains stable even after *tsp-1* expression has normalized after ELTS.

We next sought to determine the subcellular localization and stability of the heat-induced

TSP-1 tetraspanin web formation. We generated strains with CRISPR-mediated wrmScarlet

knock-in at the endogenous *tsp-1* locus that was genetically crossed with previously validated

GFP reporters for intestinal subcellular structures, including Akt-PH::GFP that binds to

PIP2/PIP3 at the inner leaflet of plasma membranes, or GFP::C34B2.10(SP12) that labels the

endoplasmic reticulum membrane (ERm) (*13, 36, 37*). Confocal microscopic imaging revealed

the striking proximity of Akt-PH::GFP and TSP-1::wrmScarlet throughout intestinal apical

membranes (**Fig. 4A**), corroborated by quantitative fluorescent intensity correlation analysis

(**Fig. 4B**). By contrast, ERm::GFP and TSP-1::wrmScarlet did not exhibit apparent co-

localization (**Fig. 4C, 4D**). We further assessed the temporal dynamics of tetraspanin webs by

imaging endogenous TSP-1::wrmScarlet (**Fig. 4E**) and found that the tetraspanin web structure

exhibited markedly consistent stability across the entire field and time scale of imaging (**Fig. 4F,**

Movie S1). These results indicate that TSP-1-dependent tetraspanin webs, once induced by

heat, form rather stable structures closely associated with the intestinal apical membrane.

What are the consequences of TSP-1 up-regulation triggered by ELTS at the organismal

level? To address this question, we measured the population lifespan and survival rates of wild

type and *tsp-1* deletion mutants under various heat stress conditions. Under constant 28 $^{\circ}$C

starting from L4, deficiency of *tsp-1* caused shortened lifespan and premature population death

compared with wild type animals (**Fig. 5A**). Transient ELTS (28 $^{\circ}$C for 24 hrs at L4) markedly

extended the longevity of wild-type animals at subsequent normal cultivation temperature (20

$^{\circ}$C). By contrast, the ELTS effect on longevity extension was abolished in *tsp-1* mutants (**Fig.**

5B). We applied 28 $^{\circ}$C exposure to animals growing constantly for 48 hrs at 20 $^{\circ}$C post L4, and

found that transient ELTS (28 $^{\circ}$C for 24 hrs at L4) enhanced survival rates in wild type but not

tsp-1-deficient animals (**Fig. 5C**). We used RNAi against *cbp-1* (null mutations are lethal) and

observed that reducing *cbp-1* expression caused even more shortened population lifespans (**Fig.**

5D). Given the more severe phenotype of *cbp-1* deficiency, CBP-1 likely regulates additional

genes beyond *tsp-1*. Indeed, we found that the heat shock chaperon-encoding gene *hsp-16* was

transiently induced by ELTS, and its induction required both HSF-1 and CBP-1 (**fig. S6**). Unlike

hsp-16, ELTS induction of *tsp-1* is HSF-1 independent (**fig. S2**), yet its similarly transient

induction leads to stable tetraspanin multimerization and web-like structural formation (**Fig. 3**).

Finally, we used a CRISPR-mediated GFP knock-in allele of *cbp-1* to show that ELTS markedly

increased the probability of nuclear entry by endogenous CBP-1 in the intestine, suggesting that

CBP-1 plays an instructive rather than permissive role in *tsp-1* up-regulation by heat (**Fig. 5E, F**).

Like induction of *tsp-1* mRNAs but not TSP-1 proteins, increased nuclear entry of CBP-1 also

followed a similarly transient pattern upon ELTS (**fig. S7**, **Fig. 3**). Taken together, these results

demonstrate that TSP-1 promotes survival under 28 ^{o}C heat stress and its CBP-1-dependent

induction mediates effects of transient ELTS on lasting benefits to longevity and organismal

resilience to subsequent heat stress conditions (**Fig. 5G**).

To assess the sufficiency of TSP-1 in conferring stress resilience, we next determined

gain-of-function effects of TSP-1 in *C. elegans* and ectopically in human cells. In *C. elegans*, we

generated two transgenic strains carrying genomic *tsp-1* regions as extrachromosomal arrays at

either low or high copy numbers (over-expression or OE, line 1 and 2, respectively) (**Fig. 6A**).

qRT-PCR confirmed that *tsp-1* mRNA expression levels are differentially up-regulated in the two

strains with array-positive animals (**Fig. 6A**). In thermal resilience assays, we found that OE line

2 with high-copy *tsp-1* arrays exhibited markedly enhanced survival rates upon continuous 28

^{o}C heat stress, compared with array-negative control or OE line 1 with low-copy *tsp-1* arrays

(**Fig. 6B**). In lifespan assays, OE line 2 exhibited similarly extended survival and longevity under

20 ^{o}C cultivation conditions, with a marked increase in both median and maximal lifespans in

animals carrying high-copy expression of *tsp-1* (**Fig. 6C**). Furthermore, we over-expressed *C.*

elegans tsp-1 cDNA with a tagged *GFP* driven by the CMV promoter in human embryonic

kidney cell lines (HEK293) (**Fig. 6D**). Such ectopic expression of *tsp-1* resulted in prominent

tetraspanin web-like structures (**Fig. 6E**) and functionally protected HEK293 cells against

thermal stress conditions (42 $^{\circ}$C). These results demonstrate striking gain-of-function effects of

tetraspanin-encoding *tsp-1* in both *C. elegans* and heterologous mammalian cells.

Previous studies have shown that transient metabolic or environmental stress can be

established in early life and elicit beneficial effects to extend longevity in *C. elegans* (*30*, *38–41*).

While heat exposure activates numerous genes involved in proteostasis and defense responses,

specific HSF-independent target genes with causal effects on longevity remain unidentified. In

this work, we identify *tsp-1* as a heat-inducible gene that mediates effects of ELTS on

organismal resilience and longevity. Regulatory mechanisms of how heat stress activates CBP-

1 in coordination with other unidentified factors await further studies. Our data suggest that *tsp-1*

expression leads to TSP-1 protein multimerization and formation of stable tetraspanin web

structures that are long lasting even in the absence of initial stimuli and *tsp-1* transcript up-

regulation. Such tetraspanin web-based stable protein structure formation represents a novel

mechanism of cellular memory distinct from previously known modes of epigenetic regulation

that occurs primarily in the nucleus, including DNA and histone modifications. We show that

TSP-1 is critical for maintaining intestinal membrane barrier functions and promotes animal

adaptation and survival under heat stress conditions and during aging. Several mammalian

tetraspanins have been shown to play important roles in maintaining blood-brain or retina-blood

barriers in endothelial cells and attenuate inflammation, cell senescence and organismal aging

(*42–44*). Thus, we propose that functional roles of tetraspanins are evolutionarily conserved and

likely mediate long-lasting physiological effects of transient cellular and organismal stresses in

mammals, including humans.

Main Figures

Fig. 1. Thermal stress induces TSP-1 expression and tetraspanin web-like structure

formation. (A) Representative bright field and epifluorescence images showing expression of

an integrated transgene *tsp-1p::tsp-1::GFP* under temperatures and durations indicated. Scale

bars: 100 µm. (B) Quantification of TSP-1::GFP fluorescence intensities under conditions

indicated. *** indicates $P < 0.001$ (n > 40 animals per condition). (C) Schematic of the *tsp-1*

gene structure, predicted membrane topology (by DeepTMHMM) and structure (by AlphaFold2)

of TSP-1. (D) SDS-PAGE and native-PAGE western blots showing heat-induced increase in

abundance and formation of dimers and multimers (asterisk) by TSP-1::GFP. (E)

Representative confocal fluorescence images showing high-resolution Z-stack views of

tetraspanin web structure formed by TSP-1::GFP, with enlarged view at right bottom. Scale bars:

10 µm.

2 **Fig. 2. Expression and roles of TSP-1 in membrane barrier integrity in adults and during**

3 **aging.** (A-D) Representative epifluorescence images showing TSP-1::GFP up-regulation from

L4 (A) to young adults (B) and during aging (C-D). (E) Quantification of fluorescence intensities

of TSP-1::GFP under conditions indicated. *** indicates $P < 0.001$ (n > 30 animals per condition).

(F) Schematic of the assay for acute membrane barrier functions. (G) Quantification of

intensities of fluorescein accumulated by live animals under conditions indicated. *** indicates P

< 0.001 (n > 20 animals per condition). (H-J) Representative epifluorescence images showing

fluorescein accumulation in animals of indicated stages, temperature and genotypes (wild type

and *tsp-1* deletion mutant allele *ok3594*). Scale bars: 100 µm.

Fig. 3. Early-life thermal stress-triggered TSP-1::GFP is long-lasting and requires CBP-1.

(A) Representative epifluorescence images showing long-lasting up-regulation of TSP-1::GFP

upon transient early-life thermal stress (ELTS, T28 for 24 hrs at L4). (B) Representative

epifluorescence images showing up-regulation of TSP-1::GFP upon sustained thermal stress

(T28 for 24, 48 or 72 hrs starting at L4). (C) Quantification of fluorescence intensities of TSP-

1::GFP under conditions indicated. *** indicates $P < 0.001$ (n > 30 animals per condition). (D)

Schematic of experimental flow using RNAi screen to identify genes required for ELTS induction

of TSP-1::GFP. (E) Representative bright-field and epifluorescence images showing that RNAi

against *tsp-1* or *cbp-1* blocks up-regulation of TSP-1::GFP by ELTS. (F) Quantitative RT-PCR

1   measurements of *tsp-1* expression levels under conditions indicated, showing transient *tsp-1*

2   up-regulation by ELTS in a CBP-1-dependent manner. *** indicates $P < 0.001$ (three

3   independent biological replicates). Scale bars: 100 μm.

Fig. 4. Subcellular localization and stability of tetraspanin web formed by endogenous

TSP-1::wrmScarlet. (A) Representative confocal images showing co-localization of the marker

Akt-PH::GFP, which binds to intestinal apical plasma membrane inner leaflet PIP2/3, with

endogenous TSP-1 tagged with wrmScarlet by CRISPR. (B) Fluorescent intensity correlation

analysis showing close proximity of Akt-PH::GFP and TSP-1::wrmScarlet. (C) Representative

confocal images showing non-co-localization of the marker ERm::GFP, which labels intestinal

ER membrane, with endogenous TSP-1 tagged with wrmScarlet by CRISPR. Scale bars: 10 µm.

(D) Fluorescent intensity correlation analysis for ERm::GFP and TSP-1::wrmScarlet. (E)

Representative confocal time-series images showing stability of tetraspanin webs formed by

endogenous TSP-1::wrmScarlet, with enlarged views in (F). Scale bars: 10 µm.

Fig. 5. Early-life thermal stress-triggered thermal resilience and longevity requires TSP-1.

(A) Lifespan curves of wild type and *tsp-1* mutants at 28 ^{o}C starting at L4. (B) Lifespan curves of

wild type and *tsp-1* mutants at constant 20 ^{o}C or after transient 28 ^{o}C for 24 hrs at L4 (ELTS). (C)

Lifespan curves of wild type and *tsp-1* mutants at 28 ^{o}C after transient 28 ^{o}C for 24 hrs at L4

(ELTS). (D) Lifespan curves of wild type animals with RNAi against *cbp-1* or control at 28 ^{o}C

starting at L4. (E) Representative confocal fluorescence images showing ELTS-induced

increase in numbers of nuclei with GFP::CBP-1 in the posterior intestine. (F) Quantification of

numbers of nuclei with GFP::CBP-1 in the posterior intestine under conditions indicated. (G)

Model of how ELTS induces lasting organismal thermal resilience through HSF-independent

epigenetic regulation of TSP-1 and stable tetraspanin web structure formation.

Fig. 6. Gain-of-function of TSP-1 confers longevity extension in *C. elegans* and thermal

resilience in human cells. (A) Schematic of the transgene that produces *tsp-1* gain-of-function

over-expression lines (OE line 1 at low level, OE line 2 at high level as measured by qRT-PCR).

(B) Lifespan curves of wild type and two representative *tsp-1* OE lines at 28 ^{o}C starting at L4. (C)

Lifespan curves of wild type and *tsp-1* OE lines at constant 20 ^{o}C. (D) Serial confocal

fluorescence images showing tetraspanin web-like structures formed in the human cell line

HEK293 by CMV promoter-driven expression of *C. elegans tsp-1::GFP* (bottom) but not *GFP*

alone (top). Scale bars: 10 μm. (E) Quantification of survival rates after heat shock in HEK293

cells showing enhanced thermal resilience conferred by ectopic gain-of-function of *C. elegans*

TSP-1::GFP. *** indicates *P* < 0.001 (three independent biological replicates).

1 Supplemental Figures

1　**fig. S1. Duration and temperature-dependent induction of TSP-1::GFP.** (A-D)

2　Representative bright-field and epifluorescence images showing expression of *tsp-1p::tsp-*

3　*1::GFP* under temperatures and durations indicated. Scale bars: 100 μm.

2 **fig. S2. TSP-1 induction by T28 is independent of HSF.** (A) Representative bright-field and

3 epifluorescence images showing expression of *tsp-1p::tsp-1::GFP* or *hsp-16p::GFP* at 28 $^{\circ}$C for

4 24 hrs, with control and RNAi against *hsf-1*. (B) Representative bright-field and epifluorescence

5 images showing expression of *tsp-1p::tsp-1::GFP* at 28 $^{\circ}$C for 24 hrs in wild type, *hsf-1(sy441)*

6 reduction-of-function heterozygous or homozygous mutants. Scale bars: 100 μm.

3 **fig. S3. Tetraspanin web regulation by heat is specific for TSP-1.** (A) Representative

4 confocal fluorescence images showing T28-induced tetraspanin web structure formation by *tsp-*

5 *1p::tsp-1::GFP* transgenes. (B), Representative confocal fluorescence images showing intestinal

6 membrane GFP from *glo-1p::GFP::CAAX* under identical conditions (T28 48 hrs).

fig. S4. Heat-induced *tsp-1* endogenously tagged with wrmScarlet. (A) Schematic of tsp-1

gene structure showing CRISPR-mediated knock-in of wrmScarlet at the C-terminus of TSP-1.

(B) Representative confocal fluorescence images showing up-regulation of endogenous TSP-

1::wrmScarlet by 28 ^{o}C for 24 hrs. (C) Representative confocal fluorescence images showing

low and high-mag views of endogenous TSP-1::wrmScarlet induced by 28 ^{o}C for 72 hrs.

(D)Representative epifluorescence images showing up-regulation of endogenous TSP-

1::wrmScarlet by 28 ^{o}C for 72 hrs. (E) Representative epifluorescence images showing up-

regulation of endogenous TSP-1::wrmScarlet by RNAi against *sptl-1*, loss of which disrupts the

biosynthesis of sphingolipids mimicking heat-induced membrane effects. (F-I) Representative

spinning disc confocal images showing high-resolution views of tetraspanin web structures.

2 **fig. S5. RNAi against *tsp-1* recapitulates mutant phenotype in membrane barrier**

3 **functions.** (A) Schematic of *tsp-1* gene structure showing two genomic regions used to

4 construct RNAi for expression in *E. Coli* and feeding to *C. elegans*. (B) Representative

5 epifluorescence images showing enhanced fluorescein uptake in *tsp-1* RNAi-treated animals, at

6 both L4 stage and Day 5-old animals. Scale bars: 100 μm.

fig. S6. Heat shock protein induction by T28 requires both HSF-1 and CBP-1. (A)

Quantitative RT-PCR measurements of *hsp-16* expression levels showing its transient induction

by ELTS (T28 for 24 hrs at L4). *** indicates $P < 0.001$ (three independent biological replicates).

(B) Representative epifluorescence images of animals with RNAi against *hsf-1* or *cbp-1* showing

up-regulation of *hsp-16*p::GFP by ELTS that depends on both HSF-1 and CBP-1, but not HSF-2.

Scale bars: 100 μm.

fig. S7. Heat-induced nuclear entry of CBP-1 endogenously tagged with GFP. (A)

Representative confocal fluorescence images showing specific nuclear signals of GFP::CBP-1

(by CRISPR knock-in at the endogenous *cbp-1* locus) that were diminished by RNAi against

cbp-1. (B) Representative confocal fluorescence images showing increased nuclear entry of

endogenous GFP::CBP-1 by 28 °C for 24 hrs. (C) Representative confocal fluorescence images

showing unaltered GFP::CBP-1 by 28 °C for 24 hrs and 20 °C for 24 hrs. (D) Representative

1   confocal fluorescence images showing unaltered GFP::CBP-1 by 28 ^{o}C for 24 hrs and 20 ^{o}C for

2   48 hrs. Shown are both high and low magnification views. Scale bars are indicated.

Methods and materials

C. elegans **strains**

C. elegans strains were grown on nematode growth media (NGM) plates seeded with

Escherichia coli OP50 at 20°C with laboratory standard procedures unless otherwise specified.

The N2 Bristol strain was used as the reference wild type (*45*). Mutants and integrated

transgenes were back-crossed at least 5 times. Genotypes of strains used are as follows:

dmaIs125 [tsp-1p::tsp-1::GFP; unc-54p::mCherry], dmaIs8 [hsp-16p::GFP; unc-54p::mCherry];

him-5(e1490), hjIs14 [vha-6p::GFP::C34B2.10(SP12 ER membrane) + unc-119(+)], etEx2 [glo-

1p::GFP::ras-2 CAAX + rol-6], tsp-1(ok3594), hsf-1(sy441), tsp-1(syb7456[tsp-1::wrmScarlet]),

cbp-1(ust475[GFP::cbp-1]).

For CRISPR knock-in of *cbp-1* with *3xFLAG::GFP::cbp-1*, a *cbp-1* promoter region was

PCR-amplified with the primers 5′- gggtaacgccagCACGTGtgggctgactcgtgctg -3′ and 5′-

ccgtcatggtctttgtagtctggtggttcatccatcaattagta -3′ from N2 genomic DNA. A *cbp-1* coding

sequence region was PCR amplified with the primers 5′-

AAGGAGGTGGAGGTGGAGCTatggatgaaccaccatcaa -3′ and 5′-

cagcggataacaatttcacaatgcatcattggatatccacc -3′ from N2 genomic DNA. A 3xFLAG::GFP region

was PCR-amplified with the primers 5′- gactacaaagaccatgacgg -3′ and 5′-

AGCTCCACCTCCACC -3′ from SHG1890 genomic DNA. ClonExpress MultiS One Step

Cloning Kit (Vazyme C113-02, Nanjing) was used to connect these fragments with vector

amplified with 5′- tgtgaaattgttatccgctgg -3′ and 5′- caCACGTGctggcgttacc -3′ from L4440. The

injection mix contained pDD162 (50 ng/ul), *cbp-1* repair plasmid (50 ng/ul), pCFJ90 (5 ng/ul),

and three sgRNAs (30 ng/ul). The mix was injected into young adult N2 animals, and the coding

sequence of 3xFLAG::GFP was inserted after the start codon using the CRISPR/Cas9 system.

The *tsp-1::wrmScarlet* knock-in allele was generated similarly (Sunybiotech), with *wrmScarlet*

sequences inserted before the termination codon of *tsp-1* in the N2 background.

For overexpression of *tsp-1* (3.26 kb), the *tsp-1* promotor (1.98 kb) and 1,283 bp

genomic DNA fragment of *tsp-1* coding region was PCR-amplified with the primers 5′ -

aaatataatttcagggatgtggtctcaaat -3′ and 5′ - tgacaagggtactgtagttcgtct -3′ from N2 genomic

DNA. The injection mix contained *tsp-1p::tsp-1* (10-50 ng/ul) and *unc-54*p::GFP (25 ng/ul) to

establish transgenic strains carrying extrachromosomal arrays of varying copy numbers.

Compound and confocal imaging

Epifluorescence compound microscopes (Leica DM5000 B Automated Upright Microscope

System) were used to capture fluorescence images (with a 10× objective lens). Animals of

different genotypes and different stages (L4, Day1, Day5, Day9) and different heat treatment

were randomly picked and treated with 10 mM sodium azide solution (71290-100MG, Sigma-

Aldrich) in M9, aligned on an 2% agarose pad on slides for imaging. The same settings (for

bright-field: exposure time 1 second, for GFP: exposure time 10 seconds) were maintained for

the images of all samples. The integrated density (IntDen) of TSP-1::GFP was measured by NIH

image program (Fiji image J) , average of mean gray value (three background area of each

image randomly selected) were employed to normalized the TSP-1::GFP. For confocal images,

the worms were randomly chosen and treated with 10 mM sodium azide in M9 solution and

aligned on an 2% agarose pad on slides and images acquired using a confocal microscope

(Leica TCS SPE) with a 63× objective lens, the same settings were maintained for the images of

all samples. Imaging in Figure 4 was conducted under room temperature conditions using a

Nikon Eclipse Ti inverted microscope equipped with a Borealis beam conditioning unit (Andor),

a CSU-W1 Yokogawa spinning disk (Andor; Belfast, Northern Ireland), a 100X PlanApo TIRF

1.49 numerical aperture (NA) objective (Nikon; Toyko, Japan), an iXon Ultra EMCCD camera

(Andor), and a laser merge module (LMM5, Spectral Applied Research; Exton, PA) containing

405, 440, 488, and 561-nm laser lines. Micro-Manager (UCSF) was used to control all the

hardware. Fiji (NIH) (*46*) and Prism (Graphpad software, Inc) were used for image display and

co-localization analysis.

SDS-page and native-page western blotting

For SDS-PAGE samples, stage-synchronized animals for control and experiment groups were

picked (n = 50) in 60 µl M9 buffer and lysed directly by adding 20 µl of 4x Laemmli sample

buffer (1610747, Bio-Rad) contain 10% of 2-Mercaptoethanol (M6250-100ML, Sigma(v/v)).

Protein extracts were denatured at 95 °C for 10 min and separated on 10% SDS-PAGE gels

(1610156, Bio-Rad) at 80 V for ~40 min followed by 110 V for ~70 min. The proteins were

transferred to a nitrocellulose membrane (1620094, Bio-Rad,) at 25 V for 40 mins by Trans-

Blot® Turbo™ Transfer System (Bio-Rad). The NC membrane was initially blocked with 5%

nonfat milk and 2% BSA (A4503, Sigma (v/v)) in tris buffered saline with 0.1% Tween 20 (93773,

Sigma) (TBS-T) at room temperature for 1 h. Proteins of interest were detected using antibodies

against GFP (A6455, Invitrogen) and Histone H3 (ab1791, Abcam) in cold room for overnight.

After three washes of 5 min each with tris-buffered saline with 0.1% Tween 20, anti-rabbit IgG,

HRP-linked Antibody (7074S, Cell Signaling Technology) was added at a dilution of 1:5000.

For native-PAGE samples, stage-synchronized animals were washed down from NGM

plates using M9 solution and subjected to 850 *g* for 60 seconds, and the pellet animals were

resuspended in pre-cooled 300 µl lysis buffer (M9 buffer + Protein inhibitors cocktails, A32965,

Thermo Fisher), then lysed by TissueRuptor (Motor unit '6') for 10 seconds and taken out,

repeated 3-5 times, followed by diluting the samples with Native Sample Buffer (161-0738, Bio-

Rad). Proteins were resolved by 4–15% Mini-PROTEAN® TGX™ Precast Protein Gels

(4561086, Bio-Rad,) and transferred to a nitrocellulose membrane (1620094, Bio-Rad). The NC

membrane was initially blocked with 5% nonfat milk and 2% BSA (A4503, Sigma (v/v)) in Tris

buffered saline with 0.1% Tween 20 (93773, Sigma) (TBS-T) at room temperature for 1 h. native

TSP-1::GFP were detected using antibodies against GFP (1:1000, 66002-1-Ig, Proteintech) at

cold room overnight. After three washes of 5 min each with Tris-buffered saline with 0.1%

Tween 20, goat anti-Mouse IgG (H+L) Secondary Antibody [HRP] (NB7539, Novus) added at a

dilution of 1:5000 as the secondary antibody.

Fluorescein assay

Stage-synchronized animals (L4, Day1 and Day5) of different genotypes at 20°C were randomly

picked to 20 µl M9 solution in 1.5 mL tube, followed by treatment with 100 µl of Fluorescein

(F0095-25G, TCI America, 10 mg/ml in M9 buffer) at room temperature for 10 minutes. Worms

were transferred to a NGM plate (without OP50), collected to new NGM plate with 200 µl M9

buffer to wash for three times, followed by randomly picking and treatment with 10 mM sodium

azide in M9 buffer and aligned on an 2% agarose pad on slides for compound microscope

imaging. The integrated density (IntDen) of fluorescein was measured by NIH image program

(Fiji image J), average of mean gray value (three background areas of each image randomly

selected) was employed to normalize the fluorescein signals.

RNA interference (RNAi) cloning and screen

To clone the *tsp-1* RNAi sequences, *tsp-1* exon 1-3 (Chr. III, C02F5.8, 1124 bp size, primer

forward: GGCGGCCGCTCTAGAACTAGTACAGATTTCCTCCACCTCTTCC, primer reverse:

TCCACGCGTCACGTGGCTAGCATTCCTAATTTTTCAGAGCCCACC) and exon 1-5 (Chr. III,

C02F5.8, 1472 bp size, primer forward:

GGCGGCCGCTCTAGAACTAGTTCCACCTCTTCCACCTTCATTAC, primer reverse:

TCCACGCGTCACGTGGCTAGCTGACAAGGGTACTGTAGTTCGTCT) were PCR-amplified

from wild-type N2 gDNA and subcloned into the SpeI and NheI sites of a pL4440 expression

vector with NEBuilder® HiFi DNA Assembly Cloning Kit (E2621L, NEB). Briefly, PCR was

performed with the following protocol on a MyCyclerTM Thermal Cycler (Bio–Rad): 98 °C for 30

s, 98 °C for 10 s, 55 °C for 30 s, 72 °C for 2 min (35 cycles); 72 °C for 5 min, and a final hold at

12 °C. The PCR products were analyzed on a 1.5 % agarose gel. 25 ng of precut pL4440 vector

with 2-fold excess of *tsp-1* PCR fragment (50 ng) were used for assembly in a thermocycler at

50°C for 30 minutes, 25 ng of precut pL4440 vector with same water instead of PCR fragment

were employed as native assembly control. 5 µl of assembly preparations were transformed to

NEB 5-alpha Competent *E. coli* (C2987H, NEB) by heat shock at exactly 42°C for exactly 30

seconds. Positive clones were verified by bacteria PCR by 2x thermo scientific dream taq green

PCR master mix (pL4440 forward: CTTATCGAAATTAATACG, pL4440 reverse:

AGGGCGAATTGGGTACCG). Positive pL4440-*tsp-1* RNAi plasmids were transformed to

competent HT115 by electroporation and verified the positive HT115 clones by bacteria PCR.

RNAi and screen for hits blocking TSP-1::GFP were performed by feeding worms with *E.*

coli strain HT115 (DE3) expressing double-strand RNA (dsRNA) targeting endogenous genes.

Briefly, dsRNA–expressing bacteria were replicated from the Ahringer library to LB plates

containing 100 µg/ml ampicillin (BP1760-25, Fisher Scientific) at 37 °C for 16 hrs. Single clone

was picked to LB medium containing 100 µg/ml ampicillin at 37 °C for 16 hrs and positive clones

(verified by bacteria PCR with pL4440 forward and pL4440 reverse primers) were spread onto

NGM plates containing 100 µg/ml ampicillin and 1 mM isopropyl 1-thio-β-Dgalactopyranoside

(IPTG, 420322, Millopore) for 24 hrs (namely RNAi plates). Developmentally synchronized

embryos from bleaching of gravid adult hermaphrodites were plated on RNAi plates and grown

at 20 °C to L4 stage followed by transfer to 28 °C incubator for 24 hrs. Randomly selected

population was observed under the epifluorescence microscope (SMZ18, Nikon) with hits

considered to block TSP-1::GFP when observing GFP levels comparable to that by *tsp-1* RNAi

(positive control).

qRT-PCR

Animals were washed down from NGM plates using M9 solution and subjected to RNA

extraction using TissueDisruptor and RNA lysis buffer (Motor unit '6' for 10 seconds and take it

out, repeat 3-5 times on ice) and total RNA was extracted following the instructions of the Quick-

RNA MiniPrep kit (Zymo Research, R1055) and reverse transcription was performed by

SuperScript™ III (18080093, Thermo scientific). Real-time PCR was performed by using

ChamQ Universal SYBR qPCR Master Mix (Q711-02, Vazyme) on the Roche LightCycler96

(Roche, 05815916001) system. Ct values of *tsp-1* were normalized to measurements of *rps-23*

(*C. elegans*) levels. Primer for qRT-PCR: *tsp-1* (forward, CTTTGATTGCCGTTGGATTT; reverse,

CCCAAAGAAAGGCCGATAAT), *hsp-16.2* (forward: ACGTTCCGTTTTTGGTGATCTTAT;

reverse, TCTGGTTTAAACTGTGAGACGTTG) and *rps-23* (forward,

CGCAAGCTCAAGACTCATCG; reverse, AAGAACGATTCCCTTGGCGT).

Thermal resilience and lifespan assays

Animals were cultured under non-starved conditions for at least 2 generations before heat

stress assays. For treatment of "early-life thermal stress" (ELTS), bleach-synchronized eggs

were growth to the L4 stage at 20°C, and populations were kept at 28 °C for 24 hrs and then

recovered for 0-72 hrs at 20 °C. For thermal resilience assays, stage-synchronized L4 stage

worms (n > 30) were picked to new NGM plates seeded with OP50 and transferred to 28 °C

incubator. Animals were scored for survival per 24 hrs. Worms failing to respond to repeated

touch of a platinum wire were scored as dead. For lifespan assays, stage-synchronized L4

stage worms (n = 50) were picked to new NGM plates seeded with OP50 containing 50 µM 5-fluoro-2′-deoxyuridine (FUDR) to prevent embryo growth at 20 °C incubator. Animals were scored for survival per 24 hrs. Worms failing to respond to repeated touch of a platinum wire were scored as dead.

Mammalian expression constructs for *C. elegans* TSP-1

The *C. elegans tsp-1* open reading frame was PCR-amplified with the primers 5′-GCGGCCTTAATTAAACCTCTAGAATGGCAACTTGGAAATTTATCATACG -3′ and 5′-AGCTCGAGATCTGAGTCCGGcAAAACGAGTGTCTTCGGTGATG -3′ from cDNA prepared from heat-induced (T28 24 hrs) wild-type animals. GFP fragment was PCR-amplified with 5′-gCCGGACTCAGATCTCGAGCTATGGTGAGCAAGGGCGCCG -3′ and 5′-CGGATCTTACTTACTTAGCGGCCGCTTACTTGTACAGCTCATCCATGCC -3′ from the pHAGE2-gfp plasmid. The *tsp-1*-GFP fragment was PCR-amplified using overlapping PCR and sub-cloned by T4 DNA Ligase to the pHAGE2-gfp plasmid at the XbaI and NotI sites.

HEK293T cells and thermal resilience assay

Human embryonic kidney (HEK) 293T cells were maintained in Dulbecco's modified Eagle's medium with 10% inactive fetal bovine serum and penicillin-streptomycin (Gibco, Grand Island, 15140122) at 37 °C supplied with 5% CO_2 in an incubator (Thermo Fisher Scientific) with a humidified atmosphere. Cells were washed once using PBS and digested with 0.05% trypsin-EDTA (Gibco) at 37 °C for routine passage of the cells. All HEK 293T cells were transiently transfected with indicated constructions using the lipo2000 (1 mg/ml, LIFE TECHNOLOGIES) reagent. The lipo2000/DNA mixture with the ratio of lipo2000 to DNA at 3:1 was incubated for

30 min at room temperature before being added to the HEK 293T cell cultures dropwise. For

thermal resilience assay, mock control and transfected HEK293T cells (48 h) in 24 well plate

were placed in a culture incubator with an ambient temperature at 42°C and humidified 5% CO_2

for 8 h followed by cell death assay or imaging with 4% PFA treatment for 12 min at room

temperature. For cell death assay, the collected cells were resuspended with 100 µl buffer with

addition of 0.1 µl Sytox blue (Thermo Fisher Scientific) for an additional 15 min at room

temperature. 25 ul of incubated cells were loaded into ArthurTM cell analysis slide (Nanoentek,

AC0050). The fluorescence intensity was measured for individual cells using automated

cytometry (ArthureTM image based cytometer, Nanoentek, AT1000) by viability assay. The 226

RFU (Fluorescence) threshold and cell size min 5 to max 25 were used for cell death analysis

and quantification.

Statistics

Data were analyzed using GraphPad Prism 9.2.0 Software (Graphpad, San Diego, CA) 389 and

presented as means ± S.D. unless otherwise specified, with P values calculated by unpaired

two-tailed t-tests (comparisons between two groups), one-way ANOVA (comparisons across

more than two groups) and two-way ANOVA (interaction between genotype and treatment), with

post-hoc Tukey HSD and Bonferroni's corrections. The lifespan assay was quantified using

Kaplan–Meier lifespan analysis, and *P* values were calculated using the log-rank test.

Acknowledgments

Some strains were provided by *the Caenorhabditis* Genetics Center (CGC), which is funded by

the NIH Office of Research Infrastructure Programs (P40 OD010440). The work was supported

Author contributions

W.J. and D.K.M. designed, performed and analyzed the *C. elegans* experiments, contributed to

project conceptualization and wrote the manuscript. B.W. generated *tsp-1* transgenic strains

and contributed to manuscript writing. A.W., M.K., F.O., and J.G. contributed to the *C. elegans*

experiments. H.D.B. and O.D.W. contributed to the tetraspanin web imaging, project

conceptualization and manuscript editing. D.K.M. supervised the project.

Competing interests

The authors declare no competing interests.

Materials & Correspondences.

Correspondence and material requests should be addressed to Dengke K. Ma, Ph. D.

(dengke.ma@ucsf.edu)

References

1.   E. R. de Kloet, M. Joëls, F. Holsboer, Stress and the brain: from adaptation to disease. *Nat Rev Neurosci.* **6**, 463–475 (2005).

2.   S. E. Taylor, B. M. Way, T. E. Seeman, Early adversity and adult health outcomes. *Dev Psychopathol.* **23**, 939–954 (2011).

3.   N. Fogelman, T. Canli, Early Life Stress, Physiology, and Genetics: A Review. *Front Psychol.* **10**, 1668 (2019).

4.   V. J. Felitti, R. F. Anda, D. Nordenberg, D. F. Williamson, A. M. Spitz, V. Edwards, M. P. Koss, J. S. Marks, Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study. *Am J Prev Med.* **14**, 245–258 (1998).

5.   S. English, T. Uller, Does early-life diet affect longevity? A meta-analysis across experimental studies. *Biol Lett.* **12**, 20160291 (2016).

6.   K. A. McLaughlin, K. C. Koenen, E. J. Bromet, E. G. Karam, H. Liu, M. Petukhova, A. M. Ruscio, N. A. Sampson, D. J. Stein, S. Aguilar-Gaxiola, J. Alonso, G. Borges, K. Demyttenaere, R. V. Dinolova, F. Ferry, S. Florescu, G. de Girolamo, O. Gureje, N. Kawakami, S. Lee, F. Navarro-Mateu, M. Piazza, B.-E. Pennell, J. Posada-Villa, M. Ten Have, M. C. Viana, R. C. Kessler, Childhood adversities and post-traumatic stress disorder: evidence for stress sensitisation in the World Mental Health Surveys. *Br J Psychiatry.* **211**, 280–288 (2017).

7.   D. W. Brown, R. F. Anda, H. Tiemeier, V. J. Felitti, V. J. Edwards, J. B. Croft, W. H. Giles, Adverse childhood experiences and the risk of premature mortality. *Am J Prev Med.* **37**, 389–396 (2009).

8.   V. D. Longo, M. P. Mattson, Fasting: molecular mechanisms and clinical applications. *Cell Metab.* **19**, 181–192 (2014).

9.   T. Laukkanen, S. K. Kunutsor, H. Khan, P. Willeit, F. Zaccardi, J. A. Laukkanen, Sauna bathing is associated with reduced cardiovascular mortality and improves risk prediction in men and women: a prospective cohort study. *BMC Med.* **16**, 219 (2018).

10.   S. J. Lupien, B. S. McEwen, M. R. Gunnar, C. Heim, Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nat Rev Neurosci.* **10**, 434–445 (2009).

11.   B. S. McEwen, In pursuit of resilience: stress, epigenetics, and brain plasticity. *Ann N Y Acad Sci.* **1373**, 56–64 (2016).

12.   D. K. Ma, M.-H. Jang, J. U. Guo, Y. Kitabatake, M.-L. Chang, N. Pow-Anpongkul, R. A. Flavell, B. Lu, G.-L. Ming, H. Song, Neuronal activity-induced Gadd45b promotes epigenetic DNA demethylation and adult neurogenesis. *Science.* **323**, 1074–1077 (2009).

13.   C. Wang, B. Wang, T. Pandey, Y. Long, J. Zhang, F. Oh, J. Sima, R. Guo, Y. Liu, C. Zhang, S. Mukherjee, M. Bassik, W. Lin, H. Deng, G. Vale, J. G. McDonald, K. Shen, D. K. Ma, A conserved megaprotein-based molecular bridge critical for lipid trafficking and cold resilience. *Nat Commun.* **13**, 6805 (2022).

14.   C. Wang, Y. Long, B. Wang, C. Zhang, D. K. Ma, GPCR signaling regulates severe stress-induced organismic death in Caenorhabditis elegans. *Aging Cell.* **22**, e13735 (2023).

15.   W. Jiang, Y. Wei, Y. Long, A. Owen, B. Wang, X. Wu, S. Luo, Y. Dang, D. K. Ma, A genetic program mediates cold-warming response and promotes stress-induced phenoptosis in C. elegans. *Elife.* **7** (2018), doi:10.7554/eLife.35037.

16. J. Brunquell, S. Morris, Y. Lu, F. Cheng, S. D. Westerheide, The genome-wide role of HSF-1 in the regulation of gene expression in Caenorhabditis elegans. *BMC Genomics*. **17** (2016), doi:10.1186/s12864-016-2837-5.

17. N. A. Baird, P. M. Douglas, M. S. Simic, A. R. Grant, J. J. Moresco, S. C. Wolff, J. R. Yates, G. Manning, A. Dillin, HSF-1-mediated cytoskeletal integrity determines thermotolerance and life span. *Science*. **346**, 360–363 (2014).

18. R. Gomez-Pastor, E. T. Burchfiel, D. J. Thiele, Regulation of heat shock transcription factors and their roles in physiology and disease. *Nat. Rev. Mol. Cell Biol.* (2017), doi:10.1038/nrm.2017.73.

19. R. I. Morimoto, The heat shock response: systems biology of proteotoxic stress in aging and disease. *Cold Spring Harb Symp Quant Biol*. **76**, 91–99 (2011).

20. V. Prahlad, T. Cornelius, R. I. Morimoto, Regulation of the Cellular Heat Shock Response in Caenorhabditis elegans by Thermosensory Neurons. *Science*. **320**, 811–814 (2008).

21. S. Charrin, S. Jouannet, C. Boucheix, E. Rubinstein, Tetraspanins at a glance. *J Cell Sci*. **127**, 3641–3648 (2014).

22. M. E. Hemler, Tetraspanin functions and associated microdomains. *Nat Rev Mol Cell Biol*. **6**, 801–811 (2005).

23. S. J. van Deventer, V.-M. E. Dunlock, A. B. van Spriel, Molecular interactions shaping the tetraspanin web. *Biochem Soc Trans*. **45**, 741–750 (2017).

24. M. Cui, M. Han, Roles of chromatin factors in C. elegans development. *WormBook*, 1–16 (2007).

25. C. González-Aguilera, F. Palladino, P. Askjaer, C. elegans epigenetic regulation in development and aging. *Brief Funct Genomics*. **13**, 223–234 (2014).

26. R. M. Woodhouse, A. Ashe, How do histone modifications contribute to transgenerational epigenetic inheritance in C. elegans? *Biochem Soc Trans*. **48**, 1019–1034 (2020).

27. Y. Shi, C. Mello, A CBP/p300 homolog specifies multiple differentiation pathways in Caenorhabditis elegans. *Genes Dev*. **12**, 943–955 (1998).

28. D. J. Eastburn, M. Han, A gain-of-function allele of cbp-1, the Caenorhabditis elegans ortholog of the mammalian CBP/p300 gene, causes an increase in histone acetyltransferase activity and antagonism of activated Ras. *Mol Cell Biol*. **25**, 9427–9434 (2005).

29. M. Vora, M. Shah, S. Ostafi, B. Onken, J. Xue, J. Z. Ni, S. Gu, M. Driscoll, Deletion of microRNA-80 activates dietary restriction to extend C. elegans healthspan and lifespan. *PLoS Genet*. **9**, e1003737 (2013).

30. L. Zhou, B. He, J. Deng, S. Pang, H. Tang, Histone acetylation promotes long-lasting defense responses and longevity following early life heat stress. *PLoS Genet*. **15**, e1008122 (2019).

31. D. S. Cabianca, C. Muñoz-Jiménez, V. Kalck, D. Gaidatzis, J. Padeken, A. Seeber, P. Askjaer, S. M. Gasser, Active chromatin marks drive spatial sequestration of heterochromatin in C. elegans nuclei. *Nature.* **569**, 734–739 (2019).

32. A. Ganner, J. Gerber, A.-K. Ziegler, Y. Li, J. Kandzia, T. Matulenski, S. Kreis, G. Breves, M. Klein, G. Walz, E. Neumann-Haefelin, CBP-1/p300 acetyltransferase regulates SKN-1/Nrf cellular levels, nuclear localization, and activity in C. elegans. *Exp Gerontol.* **126**, 110690 (2019).

33. L. D. Mathies, J. H. Lindsay, A. P. Handal, G. G. Blackwell, A. G. Davies, J. C. Bettinger, SWI/SNF complexes act through CBP-1 histone acetyltransferase to regulate acute functional tolerance to alcohol. *BMC Genomics.* **21**, 646 (2020).

34. T. Y. Li, M. B. Sleiman, H. Li, A. W. Gao, A. Mottis, A. M. Bachmann, G. El Alam, X. Li, L. J. E. Goeminne, K. Schoonjans, J. Auwerx, The transcriptional coactivator CBP/p300 is an evolutionarily conserved node that promotes longevity in response to mitochondrial stress. *Nat Aging.* **1**, 165–178 (2021).

35. L. N. Barrett, S. D. Westerheide, The CBP-1/p300 Lysine Acetyltransferase Regulates the Heat Shock Response in C. elegans. *Front Aging.* **3**, 861761 (2022).

36. A. Shi, O. Liu, S. Koenig, R. Banerjee, C. C.-H. Chen, S. Eimer, B. D. Grant, RAB-10-GTPase-mediated regulation of endosomal phosphatidylinositol-4,5-bisphosphate. *Proc Natl Acad Sci U S A.* **109**, E2306-2315 (2012).

37. N. Xu, S. O. Zhang, R. A. Cole, S. A. McKinney, F. Guo, J. T. Haas, S. Bobba, R. V. Farese, H. Y. Mak, The FATP1-DGAT2 complex facilitates lipid droplet expansion at the ER-lipid droplet interface. *J Cell Biol.* **198**, 895–911 (2012).

38. R. Xiao, B. Zhang, Y. Dong, J. Gong, T. Xu, J. Liu, X. Z. S. Xu, A genetic program promotes C. elegans longevity at cold temperatures via a thermosensitive TRP channel. *Cell.* **152**, 806–817 (2013).

39. G. J. Lithgow, T. M. White, S. Melov, T. E. Johnson, Thermotolerance and extended life-span conferred by single-gene mutations and induced by thermal stress. *Proc Natl Acad Sci U S A.* **92**, 7540–7544 (1995).

40. D. Zhu, X. Wu, J. Zhou, X. Li, X. Huang, J. Li, J. Wu, Q. Bian, Y. Wang, Y. Tian, NuRD mediates mitochondrial stress-induced longevity via chromatin remodeling in response to acetyl-CoA level. *Sci Adv.* **6**, eabb2529 (2020).

41. Y. Tian, G. Garcia, Q. Bian, K. K. Steffen, L. Joe, S. Wolff, B. J. Meyer, A. Dillin, Mitochondrial Stress Induces Chromatin Reorganization to Promote Longevity and UPR(mt). *Cell.* **165**, 1197–1208 (2016).

42. Y. Jin, Y. Takeda, Y. Kondo, L. P. Tripathi, S. Kang, H. Takeshita, H. Kuhara, Y. Maeda, M. Higashiguchi, K. Miyake, O. Morimura, T. Koba, Y. Hayama, S. Koyama, K. Nakanishi, T. Iwasaki, S. Tetsumoto, K. Tsujino, M. Kuroyama, K. Iwahori, H. Hirata, T. Takimoto, M. Suzuki, I. Nagatomo, K. Sugimoto, Y. Fujii, H. Kida, K. Mizuguchi, M. Ito, T. Kijima, H. Rakugi, E. Mekada, I. Tachibana, A. Kumanogoh, Double deletion of tetraspanins CD9 and CD81 in mice leads to a syndrome resembling accelerated aging. *Sci Rep.* **8**, 5145 (2018).

43. C. Zhang, M. B. Lai, M. G. Pedler, V. Johnson, R. H. Adams, J. M. Petrash, Z. Chen, H. J. Junge, Endothelial Cell-Specific Inactivation of TSPAN12 (Tetraspanin 12) Reveals Pathological Consequences of Barrier Defects in an Otherwise Intact Vasculature. *Arterioscler Thromb Vasc Biol*. **38**, 2691–2705 (2018).

44. J. Franz, M. Tarantola, C. Riethmüller, How tetraspanins shape endothelial and leukocyte nano-architecture during inflammation. *Biochem Soc Trans*. **45**, 999–1006 (2017).

45. S. Brenner, The genetics of Caenorhabditis elegans. *Genetics*. **77**, 71–94 (1974).

46. J. Schindelin, I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.-Y. Tinevez, D. J. White, V. Hartenstein, K. Eliceiri, P. Tomancak, A. Cardona, *Nat Methods*, in press, doi:10.1038/nmeth.2019.