Improved chromosome-level genome assembly for marigold (*Tagetes

***erecta*)**

Fan Jiang[1]*, Lihua Yuan[1,2,3]*, Sen Wang[1], Hengchao Wang[1], Dong Xu[1], Anqi Wang[1],

and Wei Fan[1]†

Affiliations:

[1]Shenzhen Branch, Guangdong Laboratory for Lingnan Modern Agriculture; Genome Analysis

Laboratory of the Ministry of Agriculture and Rural Affairs; Agricultural Genomics Institute at

Shenzhen, Chinese Academy of Agricultural Sciences, Shenzhen, Guangdong, 518120, China.

[2]State Key Laboratory of Crop Stress Adaptation and Improvement, School of Life Sciences, Henan

University, Kaifeng 475004, China.

[3]Shenzhen Research Institute of Henan university, Shenzhen 518000, China.

*These authors contributed equally to this work.

†Corresponding authors: Agricultural Genomics Institute at Shenzhen, Chinese Academy of

Agricultural Sciences, Shenzhen 518120, China. Email: fanwei@caas.cn

Abstract

Marigold (*Tagetes erecta* L.) is a popular ornamental plant of the Asteraceae family, and its petals are considered the most abundant source of lutein. A low-continuity chromosome-level genome sequence of marigold was published recently, with poor annotation of the protein-coding genes, which hinders the studies of lutein biosynthesis. Here, we generated a near telomere-to-telomere level genome assembly of marigold based on highly accurate high-fidelity (HiFi) long reads and Hi-C sequencing data. Compared to the previously reported marigold genome, the current assembly had obviously higher contiguity and higher completeness of gene set. The current genome assembly has a 27-fold increase in contig N50 size, a 12.1% increase in chromosome anchoring rate, and a 9.0% increase in BUSCO complete rate for the gene set. Besides, the current assembly has much fewer assembly errors. Based on this high-quality genome assembly, we found that the 170-bp repeats are the most abundant centromeric unit and all centromeric regions are distributed along the whole chromosomes for all 12 centromeres, indicating the existence of the holocentromeres in marigold. In addition, we analyzed the structure and phylogenetic relationship of the four *PSY* genes, and revealed that these genes have diversified and possibly executed different functions in various tissues. Our near telomere-to-telomere level genome assembly and comprehensive gene annotation will greatly facilitate the breeding of marigold and researches aimed at improving lutein production.

Keywords: *Tagetes erecta*; marigold; Asteraceae family; near telomere-to-telomere assembly; PacBio HiFi sequencing

1. Introduction

Marigold (*Tagetes erecta* L.) is one of the most popular ornamental plants of the Asteraceae family,

and its flower petals has become the most important raw material for lutein production (Lin et al.,

2015; Zhang et al., 2020b; Zhang et al., 2020a). Marigold is an annual herbaceous plant with yellow

or orange flowers (Burlec et al., 2021; Moliner et al., 2018), and its flower petals are considered the

most abundant source of lutein (Nwachukwu et al., 2016; Park et al., 2017; Tanaka et al., 2008),

which is effective in the prevention and treatment of cancer, coronary artery disease, and many eye

diseases (Buscemi et al., 2018; Chung et al., 2017; Gansukh et al., 2019; Li et al., 2020).

The lutein biosynthetic pathway has been well studied in many high plants, and it starts with the

condensation of two geranylgeranyl diphosphate (GGPP) molecules to produce phytoene, which is

catalyzed by phytoene synthase (PSY) (Nisar et al., 2015; Sun et al., 2018). Subsequently, phytoene

is transformed to lycopene by a series of desaturation and isomerization reactions (Giuliano, 2014).

Then, lycopene is cyclized twice by lycopene ε-cyclase and lycopene β-cyclase to generate α-

carotene, which is hydroxylated to lutein by hydroxylase (Kim et al., 2009). In this pathway, PSY

catalyzes the first and rate-limiting step in lutein biosynthesis (Maass et al., 2009; Nisar et al., 2015;

Rodriguez-Villalon et al., 2009), and four *PSY* genes in marigold were identified on the basis of a

whole genome sequence (Xin et al., 2023). However, the functional differences of these genes

among roots, stems, leaves, and flowers in marigold was less understood (Feng et al., 2018).

A recently study by Xin et al has reported a low-continuity marigold genome assembly, which

contains 3,451 contigs with a contig N50 size of 1.4 Mb, and only 87.6% of contig sequences were

anchored to 12 pseudo-chromosomes (Xin et al., 2023). In addition, the ratio of transposable

elements (57.3%) and BUSCO complete rate (89.3%) for the genes were obviously lower than those

of the closely related species (Badouin et al., 2017; Fan et al., 2022; Liu et al., 2020; Xu et al., 2021).

In this study, we generated a near telomere-to-telomere level reference genome assembly for

marigold, explored the structure of centromeres, and investigated the expression patterns of the *PSY*

genes in different tissues.

2. Materials and Methods

2.1. Plant materials and genome sequencing

The seeds of marigold were provided by Suqian Sunrise Seed Industry Co., Ltd, in March 2022, and

then were grown in a greenhouse of Agricultural Genomics Institute at Shenzhen, Chinese Academy

of Agricultural Sciences, Guangdong province, China. Total genomic DNA was extracted from

fresh leaves using the Hi-DNAsecure Plant Kit (TIANGEN, China). Then, sequencing libraries

were constructed by SMRTbell Express Template Prep Kit 2.0 (PacBio, USA), and sequenced on a

PacBio Sequel II platform with Circular consensus sequence (CCS) mode (PacBio, USA). For Hi-

C sequencing, the cross-linked DNA sample was digested with the MboI at motifs GATC, and

paired-end sequenced (2 x 150 bp) on an Illumina NovaSeq 6000 platform.

2.2. Transcriptome sequencing

To assist the prediction of protein-coding genes, total RNA samples were extracted by RNeasy Plant

Mini Kit (QIAGEN, Germany) from the different tissues, and then were reverse-transcribed into

cDNA using NEBNext Single Cell/Low Input cDNA Synthesis & Amplification Module (NEB,

UK). Sequencing libraries were constructed by SMRTbell Express Template Prep Kit 2.0 (PacBio,

USA) and sequenced on PacBio Sequel II system (PacBio, USA).

To investigate the expression patterns of candidate genes involved in lutein biosynthesis, the

samples collected from the roots, stems, leaves and flowers of 80-day-old plants were used for tissue

specificity analysis. Each plant represents a biological repeat, and five biological replicates were

used in this study. All samples were immediately frozen in liquid nitrogen and stored at -80°C until

use. Total RNA for each sample was extracted using RNeasy Plant Mini Kit (QIAGEN, Germany)

according to the manufacturer's instructions, and the cDNA libraries were paired-end (2 x 150 bp)

sequenced on an Illumina NovaSeq 6000 system.

2.3 Genome assembly

Before genome assembly, the PacBio HiFi reads with read length < 2,000 bp or average quality <

99% were filtered out. Then, the genome size was estimated by GCE v1.02 (Liu et al., 2013) based

on k-mer frequencies (K = 17). The filtered PacBio HiFi reads were assembled using Hifiasm

v0.16.1 (Cheng et al., 2021) with parameter "-l 3". Purge_dups v1.2.5 (Guan et al., 2020) with

parameters "-a 83 -f 0.8 -e" was used to filter the duplicate contigs in the primary assembly. After

removing contaminant contigs, we obtained 47 contigs with a total length of 778.9 Mb and a contig

N50 size of 38.0 Mb (**Table 1**). The completeness of the assembled contigs were evaluated using

the BUSCO v5.2.2 (Simao et al., 2015) with embryophyta_odb10 database, and the contig

correctness was estimated based on yak QV (https://github.com/lh3/yak).

The pseudo-chromosomes were constructed using Hi-C reads. We first mapped the Hi-C reads

onto the assembled contigs by Bowtie2 v2.3.4.3 (Langmead and Salzberg, 2012), and then used

HiC-Pro v2.11.4 (Servant et al., 2015) pipeline to detect valid ligation pairs and generate the Hi-C

link matrixes among different contigs. At last, the contigs were clustered, ordered, and oriented into

pseudo-chromosomes using EndHiC v1.0 (Wang et al., 2022) with parameters "--rounds 3 --

binnumstep 6".

2.4. Transposable elements identification

Transposable elements (TEs) were identified through three steps: (1) we used EDTA v1.9.9 (Ou et

al., 2019) to produce a filtered TE library; (2), we performed homology-searching against the above

structural TE library, Repbase v26.05 (Bao et al., 2015), and protein-coding TE database using

RepeatMasker v4.1.2 (Smit et al.); (3) we masked the above identified TEs with Ns in the genome,

and used RepeatModeler v2.0.2 (Flynn et al., 2020) to construct an extra *de novo* TE library, which

was further classified by TERL (da Cruz et al., 2021). All classified TE sequences in the *de novo*

TE library were used by RepeatMasker to identify the remaining TEs in the genome.

2.5. Gene prediction and functional annotation

We used Augustus v3.4.0 (Stanke et al., 2006) to predict the protein-coding gene models. The

training parameters for Augustus were generated during the BUSCO (Simao et al., 2015)

completeness assessment of the genome assembly. The homology hints were generated by aligning

the protein-coding sequences from the published genomes of *Smallanthus sonchifolius* (Fan et al.,

2022), *Mikania micrantha* (Liu et al., 2020), *Helianthus annuus* (Badouin et al., 2017), and *Stevia*

rebaudiana (Xu et al., 2021) to marigold genome, using Exonerate v2.2.0 (Slater and Birney, 2005).

To produce transcription hints, the full-length transcripts and Illumina RNA-Seq reads were aligned

to the genome, and then were transformed to hints by blat2hints and bam2hints in Augustus,

respectively. The tRNA genes were predicted by tRNAscan-SE v2.0 (Chan et al., 2021), and the

rRNA genes were predicted by cmscan from infernal v1.1.4 (Nawrocki and Eddy, 2013). Functional

annotation was performed by searching the protein sequences against the NCBI-NR and KEGG

databases using DIAMOND v2 (Buchfink et al., 2021). The protein domain annotation was

performed using InterProScan v5.52-86 (Blum et al., 2021) against InterPro database.

2.6. Identification of telomeres and centromeres

To investigate the telomeres and centromeres, tandem repeats were explored using Tandem Repeats

Finder (TRF) v4.07 (Benson, 1999) with parameter "2 5 7 80 10 50 2000 -h -d". We identified

telomeres on the chromosome scaffolds based on TRF results and telomeric units of "TTTAGGG"

and "CCCTAAA". For centromere annotation, we investigated the tandem repeat units with length

of 100 - 200 bp, and counted the total length of each repeat units, revealing the 170-bp repeats were

the most abundant unit in the genome, followed by 167-bp, 178-bp and 169-bp repeats. The

distribution of the four repeat units in different pseudo-chromosomes were visualized by IGV

v2.11.4 (Thorvaldsdottir et al., 2013).

2.7. Analysis of genes involved in lutein biosynthesis

The lutein biosynthesis genes in marigold genome were identified by homology alignment. Firstly,

we downloaded the known genes involved in the lutein biosynthesis pathway (map00906) from

KEGG database for the closely related species in Asterales order, including *Helianthus annuus*,

Erigeron canadensis, *Lactuca sativa*, and *Cynara cardunculus* var. *scolymus*. Then, we aligned the

protein-coding genes of marigold to the downloaded lutein biosynthesis genes using DIAMOND

v2 (Buchfink et al., 2021) with parameters "blastp --more-sensitive --evalue 0.00001". The genes

in marigold with best alignment identity > 80% and coverage > 80% were taken as potential lutein

synthesis genes. Furthermore, the orthogroups constructed by OrthoFinder v2.5.2 (Emms and Kelly,

2019) that contained these genes were obtained and checked.

To construct the tree for *PSY* genes, we downloaded the protein sequences of *PSY* genes from the

KEGG database for the 4 Asterales species (*Helianthus annuus*, *Erigeron canadensis*, *Lactuca*

sativa, *Cynara cardunculus* var. *scolymus*, and *Solanum lycopersicum*) and 4 Solanales species

(*Solanum lycopersicum*, *Solanum pennellii*, *Solanum stenotomum*, and *Ipomoea triloba*). The tree

was constructed by fasttree v2.1.11 (Price et al., 2010) with parameters "-lg -gamma" based on

multiple sequences alignments from protein sequences. The expression level of these candidate

genes was evaluated by transcripts per million (TPM) calculated from the HISAT v2.2.1 (Kim et

al., 2019) mapping results of root, stem, leaf, and flower transcriptome data.

3. Results

3.1. Near telomere-to-telomere level genome assembly of marigold

A total of 49.7 Gb (64 ×) PacBio HiFi reads with a read N50 length of 18.3 Kb (**Supplementary**

Table S1) were assembled into contigs using Hifiasm (Cheng et al., 2021). The primary assembly

includes 381 contigs with a total length of 1,011 Mb, a contig N50 size of 36.5 Mb, and a contig

N90 size of 8.8 Mb (**Supplementary Table S2**). This assembly is obviously larger than the

estimated genome size (~780 Mb) that evaluated based on the distribution of k-mer frequencies

(**Supplementary Figure S1**), which may result from genome heterozygosity. After purging the

heterozygous contigs from the primary assembly using Purge_dups (Guan et al., 2020), we obtained

a genome assembly with a total length of 778.9 Mb (**Table 1**), which is comparable to the estimated

genome size. The genome assembly comprises 47 contigs with contig N50 and N90 size of 38.0 and

11.0 Mb, respectively (**Table 1 and Supplementary Table S2**), revealing a BUSCO complete rate

of 98.3% (**Table 1**). In addition, the contig correctness was evaluated by using yak, which revealed

an average base accuracy of 99.9998% (QV57), suggesting the high accuracy of our genome

assembly. Compared to recently published genome assembly of Marigold by Xin et al. (defined as

assembly v1) (Xin et al., 2023), the huge improvement of our genome assembly (defined as

assembly v2) was highlighted by the total contig number (~73-fold decrease), contig N50 size (27-

folds increase), and genome size (71.7 Mb increase) (**Table 1**). Based on the Hi-C data

(**Supplementary Table S1 and S3**), we successfully anchored 99.7% (776.8 Mb) of contig

sequences onto 12 pseudo-chromosomes (**Table 1, Figure 1a and b**), which was obviously higher

than that of v1 assembly (87.6%) (**Table 1**). Finally, we obtained a near telomere-to-telomere level

reference genome for marigold, with scaffold N50 size of 69.8 Mb and N90 size of 40.8 Mb.

We identified a total of 590.4 Mb (75.8%) of TEs in v2 assembly (**Table 1 and Supplementary**

Figure S2), including 84.1 Mb (14.2%) of structural intact TEs. The proportion of TEs in v2

assembly falls in the range of that for the closely related species (73.4% - 84.7%) (Badouin et al.,

2017; Fan et al., 2022; Liu et al., 2020; Xu et al., 2021), and showed a 18.1% increase compared to

v1 assembly (57.7%) (**Table 1 and Supplementary Table S4**). In addition, we identified a total of

242.8 Mb (31.2% of the genome) TRs in marigold (**Table 1**). Then, we compared the TEs and TRs

in v2 assembly, and found that 97.5% of TRs were overlapped with TEs (**Supplementary Figure**

S3), which means that most of the TRs were formed from tandem clusters of the same type of TEs.

We predicted 42,529 protein-coding genes in v2 assembly, similar to that of *Mikania micrantha*

(46,351), *Helianthus annuus* (57,126), and *Stevia rebaudiana* (44,143), and obviously fewer than

that of *Smallanthus sonchifolius* (89,959) (Badouin et al., 2017; Fan et al., 2022; Liu et al., 2020;

Xu et al., 2021), which experienced a recent whole genome duplication (Fan et al., 2022). The

genome has a mean coding sequence length of 1,158 bp and exon number of 5.1, which was similar

to that of the related species (**Supplementary Table S5**). Functional annotation showed that 86.7%

of genes could be annotated by one of the NCBI-NR, KEGG, InterPro or GO databases

(**Supplementary Table S6**). Compared to v1 assembly, v2 assembly identified more genes (8,705

genes), which resulted in a 9.0% increase in BUSCO complete rate (**Table 1 and Supplementary**

Table S5), indicating the higher completeness of the v2 gene set. In addition, we identified 1,098

tRNA genes, 2,126 rRNA genes, and 929 other non-coding RNA genes in v2 assembly

(**Supplementary Table S7**).

3.2. Comprehensive assembly of repetitive regions and correction of large-scale

assembly errors

In v2 assembly, 79.2% of the assembled chromosome-ends have telomeric repeats (**Table 1**),

including 7 pseudo-chromosomes have telomeric repeats at both ends and 5 pseudo-chromosomes

have telomeric repeats at a single end (**Figure 1b and Supplementary Figure S4**). Compared to

v1 assembly, the v2 assembly reconstructed obviously more telomeres (79.2% vs 8.3%) (**Table 1**),

indicating the improvement in assembly of the ends for pseudo-chromosomes.

Compared to v1 assembly, the v2 assembly has an obvious increase (155.6 Mb) in the total length

of the 12 pseudo-chromosomes, which caused by both the higher total contig size (778.9 Mb vs

707.2 Mb) and the higher anchor rate (99.7% vs 87.6%) in v2 assembly (**Table 1**). Comparisons of

the pseudo-chromosomes between the assembly v2 and v1 revealed an obviously one-to-one match

relationship for all 12 pseudo-chromosomes (**Figure 2**). Compared to v2 assembly, large blocks

were missed in v1 assembly for many pseudo-chromosomes, including Chr02 (33.1 Mb), Chr03

(27.5 Mb), Chr04 (9.1 Mb), Chr05 (32.4 Mb), Chr06 (10.0 Mb), Chr08 (20.1 Mb), and Chr10 (25.5

Mb) (**Figure 2 and Supplementary Table S8**). After annotating of these large blocks in v2

assembly, we found that most of these sequences were identified as TRs, which were barely

assembled in v1 assembly (**Figure 2**).

Moreover, the pseudo-chromosomes with large blocks that showed inconsistent ordering or

orientation in v1 assembly were analyzed. For Chr03, the whole pseudo-chromosome in v1

assembly was split into two large blocks by a huge gap according to the alignment result, and the

ordering of the two blocks was inconsistent with that in v2 assembly (**Figure 2**). Similar results

were observed for Chr04, Chr05, and Chr08 (**Figure 2**). Because the huge gap will greatly reduce

the Hi-C linkage signal between the two blocks (Zhang et al., 2019), it is easy to result in wrong

connections among different chromosome-ends in v1 assembly. In addition, no telomeric repeats

was identified from the ends of these pseudo-chromosomes in v1 assembly, but identified from the

single or both ends in v2 assembly (**Figure 1b and Supplementary Figure S4**), which suggested

the correctness of these pseudo-chromosomes in v2 assembly. For Chr01, Chr06 and Chr11, the

chromosomes in v1 assembly were split into many large blocks, which were inconsistent with that

in v2 assembly, though the length of these chromosomes are similar (**Figure 2**). Since v1 assembly

has telomeric repeats located in the sub-terminal of the three pseudo-chromosomes instead of the

chromosome-ends, while v2 assembly has telomeric repeats at single or both ends (**Figure 1b and**

Supplementary Figure S4), we inferred that the v2 assembly has corrected the order and

orientation mistakes in v1 assembly. In total, the v2 assembly corrected 7 pseudo-chromosomes that

contained wrong ordering or orientation blocks in v1 assembly.

3.3. High proportion of centromeric 170-bp unit reveals the existence of

holocentromeres

The Hi-C heatmap showed many large blocks in genome assembly with no or weak Hi-C linkage

signals (**Figure 1a**), revealing a high proportion of high similar repeating sequences existed in the

genome. We identified a total of 242.8 Mb (31.2% of the genome) TRs, and the distribution of the

TRs was highly consistent with the location of these large blocks in Hi-C heatmap (**Supplementary**

Figure S5). Noticeably, the proportion of TRs in marigold genome was obviously higher than that

of *Smallanthus sonchifolius* (101.3 Mb, 3.7% of the genome), *Mikania micrantha* (108.6 Mb, 6.1%

of the genome), *Helianthus annuus* (163.1 Mb, 5.4% of the genome), and *Stevia rebaudiana* (72.8

Mb, 5.1% of the genome). In addition, the TRs and TEs were highly overlapped in marigold genome

(**Supplementary Figure S3**), and more than 97.5% of TRs were classified as TEs.

Previous studies have reported that centromere consisted of highly repetitive DNA sequences with monomers of length 100 - 200 bp (Melters et al., 2013; Naish et al., 2021; Shi et al., 2023; Sundararajan and Straight, 2022). Here, we found that 170-bp repeats were the most abundant unit in the genome, which had 682,138 repetitions accounted for 14.8% of the total genome sequences, followed by 167-bp (2.7%), 178-bp (1.2%) and 169-bp (0.9%) repeats (**Supplementary Table S9**). These four repeat units localized on all 12 pseudo-chromosomes, with the total length ranged from 4.0 Kb (Chr09) to 29.0 Mb (Chr02) (**Supplementary Table S10**). In detail, the 170-bp was the main repeat unit in Chr01, Chr02, Chr03, Chr05, Chr06, and Chr07, and the 169-bp was the main repeat unit in Chr11 (**Figure 3a and Supplementary Figure S6**). In addition, the 167-bp and 170-bp repeats were the main units for Chr04, Chr08, and Chr10, the 167-bp and 178-bp repeats were the main units for Chr09, and 169-bp and 170-bp repeats were the main units for Chr12 (**Figure 3a and Supplementary Figure S6**). Noticeably, these repeat units distributed across the whole chromosome for all 12 pseudo-chromosomes, but not constrained in a single region (**Figure 3b**). Besides, these repeat regions have obviously lower gene density than that of other parts on all pseudo-chromosomes (**Figure 3b**). Moreover, more than 80.2% of sequences within these repeat regions were annotated as retrotransposons (**Figure 3b**), which was consistent with that centromeres are composed of long arrays of tandem repeats and/or retrotransposons in many species (Henikoff et al., 2001; Zhang et al., 2014). Based on these results, we inferred that these repeat regions in marigold genome were centromeric regions, and the chromosomes of marigold should be classified as holocentric chromosomes, similar to *Holhymenia histrio*, *Rhynchospora pubera*, and *Vitis vinifera* (Bardella et al., 2020; Hofstatter et al., 2022; Marques et al., 2015; Shi et al., 2023).

3.4. Differential expression patterns of the four *PSY* genes in different tissues of marigold

In marigold, lutein accounts for up to 90% of the total carotenoid contents in the flower petals, and all genes involved in lutein biosynthesis were identified by Xin *et al.* based on a whole genome sequence (Xin et al., 2023). Phytoene synthase (PSY) catalyzes the first and rate-limiting step in lutein biosynthesis (**Figure 4a**), and four *PSY* genes were identified from marigold genome (Xin et al., 2023). However, the structure and the phylogenetic relationship of the four genes in marigold were not investigated. Based on our high-quality genome assembly, we identified four *PSY* genes (*PSYa*, *PSYb*, *PSYc*, and *PSYd*) (**Supplementary Figure S7**), consistent with the previous study (Xin et al., 2023). The four *PSY* genes have a length of 2,841 bp, 2,760bp, 3,235 bp, and 1,917 bp, respectively, all of which contain 6 exons and 5 introns (**Supplementary Figure S7**). The predicted protein sequences have 425, 398, 427, and 375 amino acids for *PSYa*, *PSYb*, *PSYc*, and *PSYd* genes, respectively. To investigate the phylogenetic relationship of these four *PSY* genes, 28 PSY protein sequences from four Solanales species and five Asterales species were used (**Figure 4b**). The phylogenetic tree showed that *PSYa* and *PSYc* were tightly grouped together, which was then clustered with *PSYb* (**Figure 4b**). In addition, *PSYd* was clustered into another group, and showed a high divergence to other three *PSY* genes (**Figure 4b**), which was consistent with that *PSYd* showed a low amino acid sequence identity to *PSYa*, *PSYb* and *PSYc* (60.9 - 62.2%).

Based on the recently published marigold genome, Xin *et al.* investigated the expression patterns of lutein biosynthesis genes at four flowering development stages in marigold flowers (Xin et al.,

2023), but the differences among different tissues were not studied. In this study, we compared the

expression levels of these genes among different tissues, including roots, stems, leaves, and flowers.

For *PSY* genes, *PSYa* only expressed in flowers and showed a high expression level, *PSYb* showed

moderate expression levels in flowers and leaves, and low level in stems, and *PSYc* exhibited

moderate expression levels in leaves, and low level in stems and roots (**Figure 4c**). In addition,

PSYd only expressed in roots with very low expression level (**Figure 4c**), which was similar to its

homolog *SlPSY3* in tomato that showed low but detectable transcript levels in roots (Fantini et al.,

2013). These results suggested that the four *PSY* genes have diversified and possibly executed

different functions in roots, stems, leaves, and flowers. In addition, the expression levels of the other

genes were all obviously higher in flowers than those in leaves, stems, and roots (**Figure 4c**), which

was consistent with that the flowers accumulated with the highest level of lutein in marigold (Moehs

et al., 2001).

320　**4. Discussions**

In this study, we reported a near telomere-to-telomere level genome assembly of marigold based on

highly accurate HiFi long reads and Hi-C sequencing data. Compared to the previously reported

marigold genome, the current assembly had obviously higher contiguity and higher completeness

of gene set. Besides, the current assembly has much fewer assembly errors. Although all the

assembled pseudo-chromosomes contain less than 5 contigs in this study, a few gaps still exist in

the genome assembly. In addition, we have assembled telomere sequences for 19 chromosome-ends,

but 5 chromosome-ends still have no assembled telomere sequences. To generate a real telomere-

to-telomere genome assembly, further investigations based on ultra-long sequencing and other

technologies will be necessary.

Based on the high-quality genome assembly, we found that the 170-bp repeats are the most

abundant centromeric unit, which followed by 167-bp, 178-bp and 169-bp repeats. All centromeric

units are distributed along the whole chromosomes for all 12 centromeres, indicating the existence

of the holocentromeres in marigold. Here, the centromeres were identified mainly based on the

analysis of the tandem repeat units. To ascertain the centromere type of marigold, more studies are

needed to investigate the organization of spindle fibres at metaphase, since the previous studies

reported that the spindle fibres are attached along almost the entire poleward surface of the

chromatids in holocentromeres (Schubert et al., 2020). In addition, the chromosomal distribution of

the centromere-specific histone H3 (CENH3) protein and the cell cycle-dependent pericentromeric

phosphorylation of histone H3 and H2A need further investigations (Baez et al., 2020).

The reference genome sequence of marigold greatly facilitates lutein biosynthesis studies. Xin *et*

al. have identified all the genes involved in lutein biosynthesis based on a low-continuity reference

genome of marigold, and investigated the expression patterns of these genes at four flowering

development stages in flowers (Xin et al., 2023). However, in marigold, the relationship of the four

PSY genes, and the expression patterns of these genes among different tissues were not studied. In

this study, we analyzed the structure and phylogenetic relationship of the four *PSY* genes, revealing

that *PSYa* and *PSYc* were tightly grouped together, then clustered with *PSYb*, and *PSYd* formed a

separate group that with a high divergence to other three *PSY* genes. In addition, our results revealed

that the four *PSY* genes have diversified and possibly executed different functions in roots, stems,

leaves, and flowers. To verify the functions of these *PSY* genes, more investigations based on the

molecular genetic methods, such as gene editing technologies, are needed. Our near telomere-to-

telomere level genome assembly and comprehensive gene annotation will greatly facilitate the

breeding of marigold and researches aimed at improving lutein production.

Data Availability Statement

All raw sequencing data generated during the current study have been deposited at

DDBJ/ENA/GenBank under project accession PRJNA975275. Genomic sequence reads have been

deposited in the SRA database with accession SRR24689467 for PacBio sequencing. Hi-C

sequencing reads have been deposited in the SRA database with accession SRR24718810. Full-

length transcript sequence reads have been deposited in the SRA database with accession

SRR24733750. Illumina transcript sequences for roots, stems, leaves, and flowers have been

deposited in SRA database with accession SRR24733730-SRR24733749. The genome assembly

and annotation have deposited at NCBI with accession JAUHHV000000000, and also available at

Zenodo: https://doi.org/10.5281/zenodo.8107331.

CRediT authorship contribution statement

368 **Fan Jiang**: Investigation, Methodology, Formal analysis, Writing - original draft. **Lihua Yuan**:

369 Investigation, Methodology, Formal analysis. **Sen Wang**: Formal analysis, Validation. **Hengchao**

370 **Wang**: Formal analysis. **Dong Xu**: Formal analysis. **Anqi Wang**: Validation, Software. **Wei Fan**:

371 Supervision, Writing - review & editing, Funding acquisition.

Declaration of competing Interest

374 The authors declare no competing interest.

Acknowledgment

377 The work was funded by the Agricultural Science and Technology Innovation Program and the Elite

378 Young Scientists Program of CAAS, the fund of Key Laboratory of Shenzhen

379 (ZDSYS20141118170111640), and the National Natural Science Foundation of China (32172430).

References

Badouin H, Gouzy J, Grassa CJ, Murat F, Staton SE, Cottret L, Lelandais-Briere C, Owens GL,

Carrere S, Mayjonade B, Legrand L, Gill N, Kane NC, Bowers JE, Hubner S, Bellec A,

Berard A, Berges H, Blanchet N, Boniface MC, Brunel D, Catrice O, Chaidir N, Claudel

C, Donnadieu C, Faraut T, Fievet G, Helmstetter N, King M, Knapp SJ, Lai Z, Le Paslier

MC, Lippi Y, Lorenzon L, Mandel JR, Marage G, Marchand G, Marquand E, Bret-Mestries

E, Morien E, Nambeesan S, Nguyen T, Pegot-Espagnet P, Pouilly N, Raftis F, Sallet E,

Schiex T, Thomas J, Vandecasteele C, Vares D, Vear F, Vautrin S, Crespi M, Mangin B,

Burke JM, Salse J, Munos S, Vincourt P, Rieseberg LH, and Langlade NB, 2017. The

sunflower genome provides insights into oil metabolism, flowering and Asterid evolution.

Nature. 546 (7656), 148. https://doi.org/10.1038/nature22380.

Baez M, Kuo YT, Dias Y, Souza T, Boudichevskaia A, Fuchs J, Schubert V, Vanzela ALL, Pedrosa-

Harand A, and Houben A, 2020. Analysis of the small chromosomal Prionium serratum

(Cyperid) demonstrates the importance of reliable methods to differentiate between mono-

and holocentricity. Chromosoma. 129 (3-4), 285-97. https://doi.org/10.1007/s00412-020-

00745-6.

Bao W, Kojima KK, and Kohany O, 2015. Repbase Update, a database of repetitive elements in

eukaryotic genomes. Mob. DNA. 6, 11. https://doi.org/10.1186/s13100-015-0041-9.

Bardella VB, Milani D, and Cabral-de-Mello DC, 2020. Analysis of Holhymenia histrio genome

provides insight into the satDNA evolution in an insect with holocentric chromosomes.

Chromosome Research. 28 (3), 369-80. https://doi.org/10.1007/s10577-020-09642-1.

Benson G, 1999. Tandem repeats finder: a program to analyze DNA sequences. Nucleic Acids Res.

27 (2), 573-80. https://doi.org/10.1093/nar/27.2.573.

Blum M, Chang HY, Chuguransky S, Grego T, Kandasaamy S, Mitchell A, Nuka G, Paysan-Lafosse

T, Qureshi M, Raj S, Richardson L, Salazar GA, Williams L, Bork P, Bridge A, Gough J,

Haft DH, Letunic I, Marchler-Bauer A, Mi H, Natale DA, Necci M, Orengo CA,

Pandurangan AP, Rivoire C, Sigrist CJA, Sillitoe I, Thanki N, Thomas PD, Tosatto SCE,

Wu CH, Bateman A, and Finn RD, 2021. The InterPro protein families and domains

database: 20 years on. Nucleic Acids Res. 49 (D1), D344-D54.

https://doi.org/10.1093/nar/gkaa977.

Buchfink B, Reuter K, and Drost HG, 2021. Sensitive protein alignments at tree-of-life scale using

DIAMOND. Nat. Methods. 18 (4), 366. https://doi.org/10.1038/s41592-021-01101-x.

Burlec AF, Pecio L, Kozachok S, Mircea C, Corciova A, Verestiuc L, Cioanca O, Oleszek W, and

Hancianu M, 2021. Phytochemical profile, antioxidant activity, and cytotoxicity

assessment of Tagetes erecta L. flowers. Molecules. 26 (5), 1201.

https://doi.org/10.3390/molecules26051201.

Buscemi S, Corleo D, Di Pace F, Petroni ML, Satriano A, and Marchesini G, 2018. The effect

oflutein on eye and extra-eye health. Nutrients. 10 (9), 1321.

https://doi.org/10.3390/nu10091321.

Chan PP, Lin BY, Mak AJ, and Lowe TM, 2021. tRNAscan-SE 2.0: improved detection and

functional classification of transfer RNA genes. Nucleic Acids Res. 49 (16), 9077-96.

https://doi.org/10.1093/nar/gkab688.

Cheng HY, Concepcion GT, Feng XW, Zhang HW, and Li H, 2021. Haplotype-resolved de novo

assembly using phased assembly graphs with hifiasm. Nat. Methods. 18 (2), 170.

https://doi.org/10.1038/s41592-020-01056-5.

Chung RWS, Leanderson P, Lundberg AK, and Jonasson L, 2017. Lutein exerts anti-inflammatory

effects in patients with coronary artery disease. Atherosclerosis. 262, 87-93.

https://doi.org/10.1016/j.atherosclerosis.2017.05.008.

da Cruz MHP, Domingues DS, Saito PTM, Paschoal AR, and Bugatti PH, 2021. TERL:

classification of transposable elements by convolutional neural networks. Brief. Bioinform.

22 (3), 1-15. https://doi.org/10.1093/bib/bbaa185.

Emms DM and Kelly S, 2019. OrthoFinder: phylogenetic orthology inference for comparative

genomics. Genome Biol. 20 (1), 238. https://doi.org/10.1186/s13059-019-1832-y.

Fan W, Wang S, Wang HC, Wang AQ, Jiang F, Liu HW, Zhao HB, Xu D, and Zhang Y, 2022. The

genomes of chicory, endive, great burdock and yacon provide insights into Asteraceae

palaeo-polyploidization history and plant inulin production. Molecular Ecology Resources.

22 (8), 3124-40. https://doi.org/10.1111/1755-0998.13675.

Fantini E, Falcone G, Frusciante S, Giliberto L, and Giuliano G, 2013. Dissection of Tomato

Lycopene Biosynthesis through Virus-Induced Gene Silencing. Plant Physiology. 163 (2),

986-98. https://doi.org/10.1104/pp.113.224733.

Feng G, Huang S, Liu Y-R, Xiao F, Liu J, Zhang Z, Chen Q, Mao Y, Cao XR, Wang Y, Chen D,

Zhou Y, Yu F-M, Liu G, Liu Y, and Niu X, 2018. The transcriptome analyses of Tagetes

erecta provides novel insights into secondary metabolite biosynthesis during flower

development. Gene. 660, 18-27.

Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, and Smit AF, 2020.

RepeatModeler2 for automated genomic discovery of transposable element families. Proc.

Natl. Acad. Sci. U.S.A. 117 (17), 9451-57. https://doi.org/10.1073/pnas.1921046117.

Gansukh E, Mya KK, Jung M, Keum YS, Kim DH, and Saini RK, 2019. Lutein derived from

marigold (Tagetes erecta) petals triggers ROS generation and activates Bax and caspase-3

mediated apoptosis of human cervical carcinoma (HeLa) cells. Food Chem. Toxicol. 127,

11-18. https://doi.org/10.1016/j.fct.2019.02.037.

Giuliano G, 2014. Plant carotenoids: genomics meets multi-gene engineering. Curr Opin Plant Biol.

19, 111-7. https://doi.org/10.1016/j.pbi.2014.05.006.

Guan D, McCarthy SA, Wood J, Howe K, Wang Y, and Durbin R, 2020. Identifying and removing

haplotypic duplication in primary genome assemblies. Bioinformatics. 36 (9), 2896-98.

https://doi.org/10.1093/bioinformatics/btaa025.

Henikoff S, Ahmad K, and Malik HS, 2001. The centromere paradox: stable inheritance with rapidly

evolving DNA. Science. 293 (5532), 1098-102. https://doi.org/10.1126/science.1062939.

Hofstatter PG, Thangavel G, Lux T, Neumann P, Vondrak T, Novak P, Zhang M, Costa L, Castellani

M, Scott A, Toegelova H, Fuchs J, Mata-Sucre Y, Dias Y, Vanzela ALL, Huettel B, Almeida

CCS, Simkova H, Souza G, Pedrosa-Harand A, Macas J, Mayer KFX, Houben A, and

Marques A, 2022. Repeat-based holocentromeres influence genome architecture and

karyotype   evolution.   Cell.   185   (17),   3153-68   e18.

https://doi.org/10.1016/j.cell.2022.06.045.

Kim D, Paggi JM, Park C, Bennett C, and Salzberg SL, 2019. Graph-based genome alignment and

genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol. 37 (8), 907-15.

https://doi.org/10.1038/s41587-019-0201-4.

Kim J, Smith JJ, Tian L, and Dellapenna D, 2009. The evolution and function of carotenoid

487    hydroxylases  in  Arabidopsis.  Plant  Cell  Physiol.  50  (3),  463-79.

488    https://doi.org/10.1093/pcp/pcp005.

489 Langmead B and Salzberg SL, 2012. Fast gapped-read alignment with Bowtie 2. Nat. Methods. 9

490    (4), 357-9. https://doi.org/10.1038/nmeth.1923.

491 Li LH, Lee JC, Leung HH, Lam WC, Fu Z, and Lo ACY, 2020. Lutein supplementation for eye

492    diseases. Nutrients. 12 (6), 1721. https://doi.org/10.3390/nu12061721.

493 Lin JH, Lee DJ, and Chang JS, 2015. Lutein production from biomass: marigold flowers versus

494    microalgae.  Bioresour.  Technol.  184,  421-28.

495    https://doi.org/10.1016/j.biortech.2014.09.099.

496 Liu B, Shi Y, Yuan J, Hu X, Zhang H, Li N, Li Z, Chen Y, Mu D, and Fan W, 2013. Estimation of

497    genomic characteristics by analyzing k-mer frequency in de novo genome projects. arXiv:

498    Genomics.

499 Liu B, Yan J, Li WH, Yin LJ, Li P, Yu HX, Xing LS, Cai ML, Wang HC, Zhao MX, Zheng J, Sun

500    F, Wang ZZ, Jiang ZY, Ou QJ, Li SB, Qu L, Zhang QL, Zheng YP, Qiao X, Xi Y, Zhang Y,

501    Jiang F, Huang C, Liu CH, Ren YW, Wang S, Liu HW, Guo JY, Wang HH, Dong H, Peng

502    CL, Qian WQ, Fan W, and Wan FH, 2020. Mikania micrantha genome provides insights

503    into the molecular mechanism of rapid growth. Nat. Commun. 11 (1), 340.

504    https://doi.org/ARTN 340

505 10.1038/s41467-019-13926-4.

506 Maass D, Arango J, Wust F, Beyer P, and Welsch R, 2009. Carotenoid crystal formation in

507    Arabidopsis and carrot roots caused by increased phytoene synthase protein levels. PLoS

508    One. 4 (7), e6373. https://doi.org/10.1371/journal.pone.0006373.

Marques A, Ribeiro T, Neumann P, Macas J, Novak P, Schubert V, Pellino M, Fuchs J, Ma W,

Kuhlmann M, Brandt R, Vanzela ALL, Beseda T, Simkova H, Pedrosa-Harand A, and

Houben A, 2015. Holocentromeres in Rhynchospora are associated with genome-wide

centromere-specific repeat arrays interspersed among euchromatin. Proc. Natl. Acad. Sci.

USA. 112 (44), 13633-38. https://doi.org/10.1073/pnas.1512255112.

Melters DP, Bradnam KR, Young HA, Telis N, May MR, Ruby JG, Sebra R, Peluso P, Eid J, Rank

D, Garcia JF, DeRisi JL, Smith T, Tobias C, Ross-Ibarra J, Korf I, and Chan SWL, 2013.

Comparative analysis of tandem repeats from hundreds of species reveals unique insights

into centromere evolution. Genome Biol. 14 (1), R10. https://doi.org/ARTN R10

10.1186/gb-2013-14-1-r10.

Moehs CP, Tian L, Osteryoung KW, and DellaPenna D, 2001. Analysis of carotenoid biosynthetic

gene expression during marigold petal development. Plant Molecular Biology. 45 (3), 281-

93. https://doi.org/Doi 10.1023/A:1006417009203.

Moliner C, Barros L, Dias MI, Lopez V, Langa E, Ferreira ICFR, and Gomez-Rincon C, 2018.

Edible flowers of Tagetes erecta L. as functional ingredients: phenolic composition,

antioxidant and protective effects on Caenorhabditis elegans. Nutrients. 10 (12), 2002.

https://doi.org/ARTN 2002

10.3390/nu10122002.

Naish M, Alonge M, Wlodzimierz P, Tock AJ, Abramson BW, Schmucker A, Mandakova T, Jamge

B, Lambing C, Kuo P, Yelina N, Hartwick N, Colt K, Smith LM, Ton J, Kakutani T,

Martienssen RA, Schneeberger K, Lysak MA, Berger F, Bousios A, Michael TP, Schatz

MC, and Henderson IR, 2021. The genetic and epigenetic landscape of the Arabidopsis

531   centromeres. Science. 374 (6569), eabi7489. https://doi.org/10.1126/science.abi7489.

532   Nawrocki EP and Eddy SR, 2013. Infernal 1.1: 100-fold faster RNA homology searches.

533   Bioinformatics. 29 (22), 2933-5. https://doi.org/10.1093/bioinformatics/btt509.

534   Nisar N, Li L, Lu S, Khin NC, and Pogson BJ, 2015. Carotenoid metabolism in plants. Mol Plant.

535   8 (1), 68-82. https://doi.org/10.1016/j.molp.2014.12.007.

536   Nwachukwu ID, Udenigwe CC, and Aluko RE, 2016. Lutein and zeaxanthin: Production technology,

537   bioavailability, mechanisms of action, visual function, and health claim status. Trends in

538   Food Science & Technology. 49, 74-84. https://doi.org/10.1016/j.tifs.2015.12.005.

539   Ou S, Su W, Liao Y, Chougule K, Agda JRA, Hellinga AJ, Lugo CSB, Elliott TA, Ware D, Peterson

540   T, Jiang N, Hirsch CN, and Hufford MB, 2019. Benchmarking transposable element

541   annotation methods for creation of a streamlined, comprehensive pipeline. Genome Biol.

542   20 (1), 275. https://doi.org/10.1186/s13059-019-1905-y.

543   Park YJ, Park SY, Valan Arasu M, Al-Dhabi NA, Ahn HG, Kim JK, and Park SU, 2017.

544   Accumulation of carotenoids and metabolic profiling in different cultivars of Tagetes

545   flowers. Molecules. 22 (2), 313. https://doi.org/10.3390/molecules22020313.

546   Price MN, Dehal PS, and Arkin AP, 2010. FastTree 2--approximately maximum-likelihood trees for

547   large alignments. PLoS One. 5 (3), e9490. https://doi.org/10.1371/journal.pone.0009490.

548   Rodriguez-Villalon A, Gas E, and Rodriguez-Concepcion M, 2009. Phytoene synthase activity

549   controls the biosynthesis of carotenoids and the supply of their metabolic precursors in

550   dark-grown Arabidopsis seedlings. Plant Journal. 60 (3), 424-35.

551   https://doi.org/10.1111/j.1365-313X.2009.03966.x.

Schubert V, Neumann P, Marques A, Heckmann S, Macas J, Pedrosa-Harand A, Schubert I, Jang

TS, and Houben A, 2020. Super-resolution microscopy reveals diversity of plant

centromere architecture. International Journal of Molecular Sciences. 21 (10), 3488.

https://doi.org/ARTN 3488

10.3390/ijms21103488.

Servant N, Varoquaux N, Lajoie BR, Viara E, Chen CJ, Vert JP, Heard E, Dekker J, and Barillot E,

2015. HiC-Pro: an optimized and flexible pipeline for Hi-C data processing. Genome Biol.

16, 259. https://doi.org/10.1186/s13059-015-0831-x.

Shi X, Cao S, Wang X, Huang S, Wang Y, Liu Z, Liu W, Leng X, Peng Y, Wang N, Wang Y, Ma Z,

Xu X, Zhang F, Xue H, Zhong H, Wang Y, Zhang K, Velt A, Avia K, Holtgrawe D, Grimplet

J, Matus JT, Ware D, Wu X, Wang H, Liu C, Fang Y, Rustenholz C, Cheng Z, Xiao H, and

Zhou Y, 2023. The complete reference genome for grapevine (Vitis vinifera L.) genetics

and breeding. Hortic Res. 10 (5), uhad061. https://doi.org/10.1093/hr/uhad061.

Simao FA, Waterhouse RM, Ioannidis P, Kriventseva EV, and Zdobnov EM, 2015. BUSCO:

assessing genome assembly and annotation completeness with single-copy orthologs.

Bioinformatics. 31 (19), 3210-2. https://doi.org/10.1093/bioinformatics/btv351.

Slater GS and Birney E, 2005. Automated generation of heuristics for biological sequence

comparison. BMC Bioinformatics. 6, 31. https://doi.org/10.1186/1471-2105-6-31.

Smit A, Hubley R, and Green P, RepeatMasker Open-4.0. 2015. http://repeatmasker.org/. Accessed:

10 November, 2020.,

Stanke M, Schoffmann O, Morgenstern B, and Waack S, 2006. Gene prediction in eukaryotes with

a generalized hidden Markov model that uses hints from external sources. BMC

Bioinformatics. 7 (1), 62. https://doi.org/10.1186/1471-2105-7-62.

Sun TH, Yuan H, Cao HB, Yazdani M, Tadmor Y, and Li L, 2018. Carotenoid Metabolism in Plants:

The Role of Plastids. Molecular Plant. 11 (1), 58-74.

https://doi.org/10.1016/j.molp.2017.09.010.

Sundararajan K and Straight AF, 2022. Centromere Identity and the Regulation of Chromosome

Segregation. Front Cell Dev Biol. 10, 914249. https://doi.org/10.3389/fcell.2022.914249.

Tanaka Y, Sasaki N, and Ohmiya A, 2008. Biosynthesis of plant pigments: anthocyanins, betalains

and carotenoids. Plant J. 54 (4), 733-49. https://doi.org/10.1111/j.1365-313X.2008.03447.x.

Thorvaldsdottir H, Robinson JT, and Mesirov JP, 2013. Integrative Genomics Viewer (IGV): high-

performance genomics data visualization and exploration. Brief. Bioinform. 14 (2), 178-

92. https://doi.org/10.1093/bib/bbs017.

Wang S, Wang HC, Jiang F, Wang AQ, Liu HW, Zhao HB, Yang BY, Xu D, Zhang Y, and Fan W,

2022. EndHiC: assemble large contigs into chromosome-level scaffolds using the Hi-C

links from contig ends. BMC Bioinformatics. 23 (1), 528. https://doi.org/10.1186/s12859-

022-05087-x.

Xin H-b, Ji F, Wu J, Zhang S, Yi CJ, Zhao S, Cong R-C, Zhao L, Zhang H-l, and Zhang Z, 2023.

Chromosome-scale genome assembly of marigold (Tagetes erecta L.): an ornamental plant

and feedstock for industrial lutein production. Horticultural Plant Journal.

Xu XY, Yuan HY, Yu XQ, Huang SZ, Sun YM, Zhang T, Liu QQ, Tong HY, Zhang YX, Wang YJ,

Liu CX, Wu L, Hou ML, and Yang YH, 2021. The chromosome-level Stevia genome

provides insights into steviol glycoside biosynthesis. Horticulture Research. 8 (1), 129.

https://doi.org/ARTN 129

10.1038/s41438-021-00565-4.

Zhang C, Wei L, Wang W, Qi W, Cao Z, Li H, Bao M, and He Y, 2020a. Identification,

characterization and functional analysis of AGAMOUS subfamily genes associated with

floral organs and seed development in Marigold (Tagetes erecta). BMC Plant Biol. 20 (1),

439. https://doi.org/10.1186/s12870-020-02644-5.

Zhang H, Zhang S, Zhang H, Chen X, Liang F, Qin H, Zhang Y, Cong R, Xin H, and Zhang Z,

2020b. Carotenoid metabolite and transcriptome dynamics underlying flower color in

marigold (Tagetes erecta L.). Sci Rep. 10 (1), 16835. https://doi.org/10.1038/s41598-020-

73859-7.

Zhang H, Koblizkova A, Wang K, Gong Z, Oliveira L, Torres GA, Wu Y, Zhang W, Novak P, Buell

CR, Macas J, and Jiang J, 2014. Boom-Bust Turnovers of Megabase-Sized Centromeric

DNA in Solanum Species: Rapid Evolution of DNA Sequences Associated with

Centromeres. Plant Cell. 26 (4), 1436-47. https://doi.org/10.1105/tpc.114.123877.

Zhang X, Zhang S, Zhao Q, Ming R, and Tang H, 2019. Assembly of allele-aware, chromosomal-

scale autopolyploid genomes based on Hi-C data. Nat Plants. 5 (8), 833-45.

https://doi.org/10.1038/s41477-019-0487-8.

622 **Table and legends**

623 Table 1. Comparisons of the genomic assembly and annotation for marigold v2 and v1 assembly.

Characteristics	v2	v1
Genome assembly		
Contig number	47	3,451
Contig size (bp)	778,905,021	707,210,696
Contig N50 size (bp)	37,953,657	1,391,023
Scaffold number	26	1,651
Scaffold size (bp)	778,926,021	709,010,696
Scaffold N50 size (bp)	69,673,756	54,151,473
Pseudo-chromosomes length (bp)	776,792,443	621,195,486
Anchored to Pseudo-chromosomes (%)	99.7	87.6
Ratio of telomeres assembled	19/24	2/24
BUSCO complete rate of the genome (%)	98.3	98.2
Genome annotation		
Transposable element length (bp)	590,391,399	407,962,073
% transposable elements	75.8	57.5
Tandem repeat length (bp)	243,461,285	60,279,323
% tandem repeats	31.3	8.5
Protein-coding gene number	42,529	33,824
Total CDS size (bp)	49,266,489	38,796,777
BUSCO complete rate of the genes (%)	98.3	89.3

Figure legends

Figure 1. Assessment of the genome assembly for marigold. (a) Hi-C heatmap of the genome

assembly. We scanned the genome by 200-Kb non-overlapping window as a bin and calculated

valid interaction links of Hi-C data between any pair of bins, and color represents Log2(links

number); (b) View of the pseudo-chromosomes. The thick lines represent contigs, and the thin lines

represent the links between the two contigs. The chromosome ends assembled with telomere-

specific repeats (CCCTAAA) were highlighted with red solid circle.

Figure 2. Comparisons between the genomes of marigold v2 and v1 assembly. The above and

right parts showed the distributions of tandem repeats in v2 and v1 assembly, respectively. We

scanned the genome by 100-Kb non-overlapping window as a bin to calculate the length of tandem

repeats. The core part of the figure showed the pair-wise alignment of genome sequences between

the v2 and v1 assembly. The corresponding segments for highly enriched tandem repeats in v2

assembly were highlighted with red box.

Figure 3. Characterization of holocentromeres in marigold genome. (a) Views of genes and four

centromeric repeat units for Chr02 with IGV. We take Chr02 as an example to show the distributions

of genes and different centromeric units. The head number showed the total length of the

chromosome. (b) Distribution of genes, centromeric repeat units and the retrotransposons in

centromeric regions for all 12 pseudo-chromosomes. The left columns showed the distribution of

genes. The length represents Log2(length) in 100 Kb windows.

657 **Figure 4.** Analysis of lutein biosynthesis in marigold. (a) Primary steps for lutein

658 biosynthesis pathway. (b) Phylogenetic tree for *PSY* genes from some Solanales and

659 Asterales species. (c) Comparisons of the expression levels of all genes involved in

660 lutein biosynthesis among different tissues.

661