Calcineurin contributes to RNAi-mediated transgene silencing and small interfering RNA production in the human fungal pathogen *Cryptococcus neoformans*

Vikas Yadav, Riya Mohan, Sheng Sun, and Joseph Heitman*

Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, NC 27710, USA

*Corresponding author

Joseph Heitman
Box 3546, 322 CARL Building, Research Drive
Department of Molecular Genetics and Microbiology
Duke University Medical Center
Durham, NC 27710
Phone: 919-684-2824
Fax: 919-684-2790
Email: heitm001@duke.edu

Running title: Calcineurin governs RNAi

Keywords: Calcium signaling; RNA interference; P-body; gene silencing; epigenetics.

Abstract

Adaptation to external environmental challenges at the cellular level requires rapid responses and involves relay of information to the nucleus to drive key gene expression changes through downstream transcription factors. Here, we describe an alternative route of adaptation through a direct role for cellular signaling components in governing gene expression via RNA interference-mediated small RNA production. Calcium-calcineurin signaling is a highly conserved signaling cascade that plays central roles in stress adaptation and virulence of eukaryotic pathogens, including the human fungal pathogen *Cryptococcus neoformans*. Upon activation in *C. neoformans*, calcineurin localizes to P-bodies, membrane-less organelles that are also the site for RNA processing. Here, we studied the role of calcineurin and its substrates in RNAi-mediated transgene silencing. Our results reveal that calcineurin regulates both the onset and the reversion of transgene silencing. We found that some calcineurin substrates that localize to P-bodies also regulate transgene silencing but in opposing directions. Small RNA sequencing in mutants lacking calcineurin or its targets revealed a role for calcineurin in small RNA production. Interestingly, the impact of calcineurin and its substrates was found to be different in genome-wide analysis, suggesting that calcineurin may regulate small RNA production in *C. neoformans* through additional pathways. Overall, these findings define a mechanism by which signaling machinery induced by external stimuli can directly alter gene expression to accelerate adaptative responses and contribute to genome defense.

Article summary (80 words)

Signaling cascades primarily drive responses to external stimuli through gene expression

changes via transcription factors that localize to the nucleus and bind to DNA. Our study

identifies an alternative mechanism whereby calcineurin, a key and direct downstream effector

of calcium signaling, is involved in post-transcriptional regulation of gene expression through

RNAi-mediated small RNA production. We propose that such signaling allows cells to bypass

the requirement for communication to the nucleus and rapidly drive stress responses in a

reversible fashion.

Introduction

RNA interference (RNAi) is a mechanism by which cellular gene expression is suppressed post-transcriptionally through targeted RNA cleavage and degradation. RNAi has been proposed to have evolved as a genome defense mechanism against invading RNA viruses and transposons (BUCHON AND VAURY 2006; OBBARD *et al.* 2009). The basic process of RNAi involves either the identification of double-stranded RNA or the generation of double-stranded RNA molecules from single-stranded RNA via RNA-dependent RNA polymerases (SHABALINA AND KOONIN 2008; OBBARD *et al.* 2009; CASTEL AND MARTIENSSEN 2013). The double-stranded RNA moiety is recognized by Dicer and cleaved into smaller RNA fragments, which are then bound by Argonaute and serve as a template to identify longer target RNA molecules. As a result, the target RNA molecule is destroyed rendering it non-functional and resulting in suppression or silencing of gene expression. RNAi has also been utilized as an important tool to study gene functions by perturbing gene expression levels (BOUTROS AND AHRINGER 2008; CASTEL AND MARTIENSSEN 2013).

While there are differences in the accessory proteins that operate during RNAi, the core proteins and key steps are highly conserved across evolution from yeasts to humans (SHABALINA AND KOONIN 2008; LAX *et al.* 2020). In fungi, the RNAi machinery has been well characterized in both model and pathogenic species since its initial discovery in *Neurospora crassa* (COGONI AND MACINO 1999; NAKAYASHIKI 2005; NICOLAS AND GARRE 2016). In addition to its conventional genome defense role, RNAi contributes to genome integrity by establishing centromere identity, maintaining centromere structure, and driving antifungal drug resistance through epimutations as well as by controlling transposition (VOLPE *et al.* 2003; FOLCO *et al.* 2008; CALO *et al.* 2014; YADAV *et al.* 2018; CHANG *et al.* 2019; PRIEST *et al.* 2022).

Cryptococcus neoformans is a basidiomycetous yeast and a human fungal pathogen that

primarily causes infections in immune-compromised patients and accounts for ~20% of

HIV/AIDS-related deaths (RAJASINGHAM *et al.* 2017; ZHAO *et al.* 2019; RAJASINGHAM *et al.*

2022). RNAi is required for genome defense in *C. neoformans* by silencing both transposons

and transgene arrays (JANBON *et al.* 2010; WANG *et al.* 2010; DUMESIC *et al.* 2013). Transgene

silencing occurs at ~50% frequency during sexual reproduction and at a lower frequency (0.1%)

during mitotic growth and has been termed sex-induced silencing (SIS) or mitotic-induced

silencing (MIS), respectively (WANG *et al.* 2010; WANG *et al.* 2012). Studies on the RNAi-

mediated silencing mechanism identified several unconventional proteins as part of the RNAi

machinery in *C. neoformans* (FERETZAKI *et al.* 2016; BURKE *et al.* 2019).

Localization of core RNAi components by direct fluorescence of epitope-tagged proteins

revealed their localization at processing bodies (P-bodies), membrane-less organelles involved in

RNA processing (WANG *et al.* 2010). Interestingly, previous studies demonstrated that the heat-

stress-activated phosphatase calcineurin also localizes to P-bodies during 37°C heat stress in *C.*

neoformans (KOZUBOWSKI *et al.* 2011). Calcineurin is a heterodimeric phosphatase comprised

of the catalytic A and regulatory B subunits (RUSNAK AND MERTZ 2000; PARK *et al.* 2019;

ULENGIN-TALKISH AND CYERT 2023). Calcineurin is activated upon calcium influx in the cell by

calcium-bound calmodulin and calcium binding to calcineurin B (YADAV AND HEITMAN 2023).

Calcineurin is also the target of two immunosuppressive drugs, FK506 and cyclosporin A, and its

activity in *C. neoformans* is inhibited by these drugs. Previous studies have established that

calcineurin is essential for both growth at 37°C and sexual reproduction in *C. neoformans*

(ODOM *et al.* 1997; CRUZ *et al.* 2001; FOX *et al.* 2001). Phosphoproteome studies identified

several candidate calcineurin substrates localized in P-bodies and explored their roles in

thermotolerance and sexual reproduction (PARK *et al.* 2016). Specifically, these substrates

included Gwo1, which was previously identified to be associated with Argonaute, a core

component of RNAi machinery; Pbp1, a homolog of the *Saccharomyces cerevisiae* poly-A-

binding protein (Pab1) binding protein, a regulator of mRNA polyadenylation; Puf4, a member

of the pumilio-family of RNA binding proteins that was previously shown to be involved in cell

wall biosynthesis in *C. neoformans*; and Vts1, a member of the Smaug family of transcriptional

regulators that binds to RNA through the sterile alpha motif (SAM) (MANGUS *et al.* 1998; RENDL

et al. 2008; DUMESIC *et al.* 2013; KALEM *et al.* 2021). Interestingly, all of the targets of

calcineurin localized to P-bodies are involved in RNA binding or processing, with one of them,

Gwo1, being part of the RNAi machinery itself (DUMESIC *et al.* 2013; PARK *et al.* 2016). Based

on calcineurin localization in P-bodies and the role of its targets in RNA binding and processing,

we explored if calcineurin plays a role in RNAi given that RNAi is executed within P-bodies.

We hypothesized that calcineurin might regulate RNAi through its action on substrates in

P-bodies. This was studied by analyzing the possible roles of calcineurin and its P-body

localized substrates in MIS (WANG *et al.* 2010). Our results suggest that calcineurin and its

substrates contribute to transgene silencing although none are essential for silencing during

mitotic growth. We also found that calcineurin plays a role in the reactivation of silenced

transgenes. Based on genetic epistasis analysis, calcineurin, and its substrates regulate transgene

silencing in different fashions. Genome-wide small RNA sequencing revealed that calcineurin

mutations down-regulate small RNA production throughout the genome including from

transgenes. Interestingly, *GWO1* deletion abolished small RNA from specific loci in the genome

without impacting the majority of RNAi targets throughout the genome. Overall, our studies

identify a novel role for calcineurin in RNAi-mediated silencing as well as small RNA

production that may contribute to genome defense in *C. neoformans*.

Materials and Methods

Strains and media

C. neoformans strain JF289**a** served as the reference strain for mitotic-induced silencing

(MIS) assays and the congenic H99α strain served as the wild-type isogenic parental for sex-

induced silencing (SIS) assays (PERFECT *et al.* 1993; WANG *et al.* 2010). Strains were grown in

YPD media for all experiments at 30°C for non-selective growth conditions. G418 and/or NAT

were added at a final concentration of 200 and 100 µg/ml, respectively, for the selection of

transformants. MS media was used for all the mating assays, which were performed as described

previously (SUN *et al.* 2019). Random spore dissections were performed after two-three weeks of

mating, and the spore germination frequency was scored after five days of dissection. All strains

and primers used in this study are listed in Supplementary Table S1 and Supplementary Table

S2, respectively.

Mitotic-induced silencing and unsilencing assays

The frequency for sex-induced and mitotic-induced silencing for the wild-type and

various mutants was calculated as described previously (WANG *et al.* 2010). Briefly, for MIS

assays, single colonies were obtained for each strain on SD-URA media to ensure expression of

URA5 and incubated at 30°C for two days. For each strain, 3 to 5 single colonies were

inoculated in YPD and grown overnight at 30°C per experiment. From the overnight culture,

various dilutions were plated on YPD and 5-FOA-containing media. The colonies were counted

in the YPD media after 2 days and 5-FOA-containing media after 7 to 10 days and the frequency

for mitotic-induced silencing was calculated.

The colonies obtained on the 5-FOA media from the MIS assays were directly

resuspended in 100 µl dH$_2$O. From this resuspension, 5 µl was directly spotted on the SD-URA

media. The colonies exhibiting a robust growth i.e., uniform growth in at least 50% of the

spotted area were considered as reverted colonies and the frequency of reversion was calculated.

PacBio sequencing and genome assembly

Long-read PacBio sequencing was conducted to assemble the complete transgene array

for the strain JF289 along with its chromosome-level genome assembly. DNA isolation was

performed to enrich the large fragment as described previously (YADAV *et al.* 2020). After a

quality check, the DNA was submitted to Duke University Sequencing and Genomic

Technologies Core facility for PacBio sequencing. The sequencing reads obtained were then

subjected to a de novo genome assembly using Canu 2.0. This resulted in an assembly for all 14

chromosomes, including the full sequence for the transgene array. The genome assembly was

then error-corrected (5X) with Pilon using short-read Illumina sequencing reads. The following

genome assembly was then employed for the downstream analysis of small RNA reads. The

genome annotations were generated by transferring the *C. neoformans* H99 genome annotations

onto the JF289 genome based on sequence identity and have not been curated.

Small RNA preparation and analysis

Single colonies obtained on either 5-FOA or YPD media from the MIS assays were used

for small RNA sample preparation and analysis. Small RNA preparation and analysis were

performed as described previously (PRIEST *et al.* 2022). Briefly, colonies were inoculated in 8

ml YPD medium and grown overnight at standard laboratory conditions. Cells were harvested,

frozen, and then subjected to lyophilization overnight. 50 to 70 mg of lyophilized material was

used for sRNA isolation following the mirVana miRNA Isolation Kit manufacturer's

instructions. sRNA was quantified with a Qubit 3 Fluorometer and submitted for sequencing at

the Duke University Sequencing and Genomic Technologies Core facility. sRNA libraries were

prepared with a QiaPrep miRNA Library Prep Kit and 1 x 75 bp reads were sequenced on the

Illumina NextSeq 500 System.

Post-sequencing, the small RNA reads were processed with cutadapt to remove the

adapter sequences, and reads smaller than 14-bp were discarded. The reads obtained after

trimming were mapped to the reference JF289 genome using Bowtie2 and resulting SAM files

were analyzed for reads length distribution and 5'-nucleotide preference with tools described

previously (Priest et al. 2022). For read mapping, we used the default option in Bowtie2 that

allows mapping of each read to only one location in the genome based on the mapping quality in

order to avoid artificial accumulation of siRNA reads at transposons and repeats. GraphPad

Prism was used to calculate percent reads and plot the graphs presented in the figures.

Furthermore, SAM files were then converted to BAM files and then to TDF format.   The TDF

files were visualized in the IGV for their distribution pattern as maps that are presented in

figures. The TDF normalization, which multiplies each value by [1,000,000/total read count] for

each sample, was employed for visualization in IGV to account for different coverage between

samples. The maps were combined with the annotations from the Geneious Prime 2022.1.1

(https://www.geneious.com/) software to overlay the annotation tracks with the read mapping.

Differential expression analysis for the small RNAs was conducted using Geneious

Prime. For this purpose, the trimmed reads were mapped to the newly generated JF289 reference

genome using Bowtie2, and plug-in DEseq2 (using "parametric" Fit Type) was used to calculate

the differential expression of sRNA levels between samples (See details in Supplementary

Materials and Methods). The analysis was conducted in two different manners; 1) where only

two 5-FOA-specific samples were compared, and 2) where all three samples (two 5-FOA and

one YPD) were pooled together. The loci with differential expression are shown in

supplementary table S3.

Results

Calcineurin deletion results in defective transgene silencing and reactivation

To probe the possible role of calcineurin in RNAi, we employed a previously

characterized system to analyze transgene array silencing in *C. neoformans* (WANG *et al.* 2010).

The transgene array includes the *URA5* gene arranged in a tandem fashion as a repeat cluster

where the expression of the *URA5* gene can be suppressed by RNAi-mediated small RNA

generation. The silencing of *URA5* within the transgene was initially discovered during sexual

reproduction where it happens in 50% of the progeny that inherit the transgene and requires an

active RNAi machinery (WANG *et al.* 2010). This process was defined as sex-induced silencing

(SIS) and was shown to act as a defensive mechanism for the *C. neoformans* genome. An

analysis of silencing during mitotic growth found transgene silencing, albeit at a much lower

frequency of approximately 1/1000, and is defined as mitotic-induced silencing (MIS).

Because calcineurin mutants fail to undergo sexual reproduction, we measured the frequency of

MIS in the wild-type and isogenic mutant strains. We generated two independent calcineurin A

(Cna1) and calcineurin B (Cnb1) mutants in the reporter JF289 strain background that bears the

URA5 transgene array, and these mutant strains were then subjected to MIS RNAi silencing

assays (Figure 1A-B). Specifically, isolated single colonies were grown in rich media and then

plated to synthetic defined (SD) media containing 5-Fluoroorotic Acid (5-FOA). 5-FOA is

metabolized into a toxin in cells expressing *URA5*, and thus only colonies that do not express

URA5 or harbor a *ura5* mutation grow on 5-FOA-containing medium. With this assay,

calcineurin mutant strains exhibited an approximate two-fold reduction in transgene RNAi

silencing as compared to the wildtype parent strain JF289, suggesting a modest but

demonstratable role for calcineurin in this process (Figure 1C-D). To further assess this, we

performed MIS assays after treating JF289 cells with the calcineurin inhibitor FK506 in an

overnight culture and then plated cells on 5-FOA media containing FK506. This experiment

revealed that prolonged calcineurin inhibition had a similar impact on the MIS frequency (Figure

1D). As expected, a known RNAi mutant (*rdp1*Δ) did not produce any 5-FOA resistant colonies

in MIS assays. Calcineurin mutation or inhibition confers growth defect at temperatures higher

than 25°C, we directly tested whether such a growth defect could lead to MIS reduction in these

mutants. First, we measured the doubling time of these mutants at 30°C which showed that both

the *cna1*Δ and the *cnb1*Δ mutants exhibited slower growth growing 1.3-1.5 times slower than the

wild-type (Table S4). We also measured the growth rate in SD media (which is supplemented

with 5-FOA for assessing MIS frequency) and found a similar reduction in growth rate for the

mutants. Next, we measured the viability of these strains in these two media conditions and

found a similar loss in viability in the mutants in both YPD and SD media (Figure S1A).

Combined together, these results show that the mutants are similar in both nutrient-rich YPD and

nutrient-limiting SD media. During the MIS frequency calculation, we always factor in the

number of viable cells plated by counting colonies on the YPD plate (Figure 1B), which accounts

for the potential loss of viability in these mutants, negating any impact on MIS frequency as a

result of viability loss. Next, we measured MIS frequency in the wild-type and two calcineurin

mutants after growth for 18 hours and 40 hours from the same culture to determine the direct

impact of the number of cell divisions. We found that the MIS frequency was unchanged for all

three strains over time indicating the number of cell divisions had little to no impact on transgene

silencing in these strains (Figure S1B). Considering the epigenetic nature of this phenomenon,

where a given cell could switch between silenced and unsilenced states at any point of time

irrespective of the number of divisions, this might be a predictable finding. Based on these

results, we conclude that MIS reduction observed in calcineurin mutants is not a result of cell

growth defects.

During the MIS assays, we observed that the colony morphology for calcineurin mutants

on 5-FOA media as well as for the wild-type cells treated with FK506 differed from the wild-

type parent without any drug treatment (Figure 1C). Notably, the majority of colonies (>70%)

obtained in the calcineurin mutants or upon calcineurin inhibition were homogeneous with

smooth edges, in stark contrast to those from the wild-type strain where >80% of the colonies

were sectored with rough peripheries. We hypothesized that this might be attributable to a

difference in the rate at which cells within the colonies reverted to *URA5* expression and that

calcineurin mutation might interfere with the reversion of RNAi-mediated transgene silencing.

To test this, we scored >100 colonies from 5-FOA media for their ability to grow on a defined

medium lacking uracil (SD-URA) that only allows the growth of cells that express *URA5*.

Colonies that exhibited a uniform growth in more than 50% of the spot were counted as reverted

or unsilenced whereas the rest were counted as silenced. These experiments revealed that a

significantly higher number of JF289 colonies could directly grow on SD-URA, as compared to

calcineurin mutants or JF289 cells growing in the presence of FK506 (Figure 1E). This result

suggests that calcineurin plays a role in controlling the frequency at which a silenced transgene

reverts and thereby escapes RNAi silencing. We also tested the MIS reversion frequency for

calcineurin mutants from the time-point experiment and found that colonies from both 18 hours

and 40 hours time points reverted at a similar rate (Figure S1B). Taken together, we conclude

that calcineurin mutants are not only modestly defective in silencing the *URA5* gene, but also

exhibit a reduced rate of unsilencing, thus contributing to prolonging the state of RNAi-mediated

transgene silencing.

Calcineurin substrates in P-bodies play different roles in silencing

To understand how calcineurin regulates RNAi silencing, we hypothesized that

calcineurin targets/substrates in P-bodies might be involved in this process (PARK *et al.* 2016).

To test this, we generated and tested two independent mutants of known calcineurin substrates

that localize to P-bodies and bind to RNA (Gwo1, Pbp1, Puf4, and Vts1) in the transgene array

strain background (JF289) and determined the MIS frequency for these mutants. Interestingly,

gwo1Δ mutations led to an approximately five-fold reduction whereas *puf4Δ* mutations led to an

approximately three-fold increase in MIS (Figure 2A). While no significant difference in the

MIS frequency was observed for the *pbp1Δ* mutants, we did observe more heterogeneity in the

shape and sizes of the colonies, suggesting that silencing might be delayed in the *pbp1Δ* mutant

without an alteration in the overall rate of silencing (Figure 2B). Finally, the *vts1Δ* mutant did

not exhibit any difference in the MIS frequency. Unlike the calcineurin mutants, none of these

mutants exhibited a growth defect (Table S4) ruling this out as a possible cause of the MIS

defect in both the *gwo1Δ* and the *puf4Δ* mutants. Next, we tested the frequency of transgene

silencing reversion for the mutants of the four P-body components by directly transferring the

colonies from 5-FOA media to SD-URA media. These experiments showed that *puf4Δ*

mutations completely abolished the MIS reversion as none of the colonies exhibited uniform

growth on uracil lacking media. Deletion of the other three genes did not have a significant

impact on transgene reactivation with the frequency ranging between 30 to 39% (Figure 2C).

Among these targets, *gwo1Δ* and *vts1Δ* mutants can successfully undergo sexual

reproduction and produce spores whereas *puf4Δ* and *pbp1Δ* mutations impair sexual

reproduction. This allowed us to determine the RNAi silencing frequency during sexual

reproduction (SIS) for the *gwo1Δ* and v*ts1Δ* mutants. For this purpose, we generated mutants for

these genes in a congenic strain of opposite mating-type, H99α, and crossed them with respective

mutants in the transgene reporter strain JF289**a**. The basidiospores produced from these crosses

were randomly dissected by micromanipulation to obtain viable progeny on the non-selective

media (YPD). The germinated spores were then genotyped for the presence of the *URA5*

transgene in their genome and in parallel assayed for their growth on 5-FOA-containing media.

SIS frequency was then calculated as the number of progeny growing on 5-FOA-containing

media out of the total number of progeny that inherited the transgene array. Previous studies

have established that a wild-type cross results in *URA5* silencing in approximately 50% of the

progeny (WANG *et al.* 2010; FERETZAKI *et al.* 2016) and it occurred at ~45% in our experiments

(Table 1). Interestingly, *gwo1Δ* mutations led to a complete abolishment of SIS in the H99

gwo1Δ x JF289 *gwo1Δ* cross and none of the analyzed progeny grew on 5-FOA. By

comparison, a 30% reduction (SIS frequency ~33%) in SIS was observed in a cross between H99

vts1Δ between JF289 *vts1Δ* strains. These results indicate that the impact of Gwo1, and

probablyVts1, on RNAi-mediated silencing is considerably more evident during sexual

reproduction. Overall, these results suggest that calcineurin targets in P-bodies are involved in

RNAi-mediated silencing but fulfill different roles suggesting the presence of multiple pathways

controlling this process.

Calcineurin and its targets impact RNAi-mediated silencing through multiple pathways

Previously, studies showed that Gwo1, Pbp1, Puf4, and Vts1 localize more prominently

to P-bodies in response to heat stress (PARK *et al.* 2016). Here, we first explored if calcineurin

controls the localization of these targets and thus impacts RNAi-mediated silencing. To study

this, we performed co-localization of these calcineurin targets at 37°C in the presence of the

calcineurin inhibitor, FK506. Interestingly, these experiments revealed that the localization

pattern of all four proteins is independent of calcineurin activity and they co-localize with a P-

body marker, Dcp1, irrespective of calcineurin activity (Figure S2). Similar to a previous study

(KOZUBOWSKI *et al.* 2011), we also noted that calcineurin activity is not required for its own

localization to P-bodies (Figure S2 – top row). These results suggest that the localization of

calcineurin and its targets to P-bodies may be driven by other factors such as structural

determinants.

To further understand the connection between calcineurin and its targets for their roles in

RNAi-mediated silencing, we generated double deletion mutants lacking Cna1 in combination

with loss of Puf4 or Gwo1 because these two targets exhibited significant differences in MIS

frequency.   Surprisingly, both *cna1Δ gwo1Δ* and *cna1Δ puf4Δ* mutants exhibited MIS

frequencies similar to *gwo1Δ* and *puf4Δ* single mutants, respectively (Figure 2A). These results

indicate a more dominant role for both Puf4 and Gwo1 than Cna1 in MIS silencing. It is also

possible that calcineurin-mediated dephosphorylation may not be required for the role of Gwo1

or Puf4 in RNAi-mediated silencing. Furthermore, calcineurin might have additional substrates

in P-bodies that regulate MIS in different ways negating the effects of calcineurin in these double

mutants. Because Gwo1 and Puf4 regulate MIS in opposite directions, we also generated a

gwo1Δ puf4Δ double mutant and determined the MIS frequency. We attempted to obtain this

double mutant strain by crossing the strains lacking either one of the genes in the parents and

analyzing the progeny by PCR analysis. We failed to obtain a *gwo1Δ puf4Δ* double mutant from

H99 *gwo1Δ* x JF289 *puf4Δ* (0 out of 73 viable progeny) suggesting the double mutant might

have a fitness defect. *GWO1* is located on chromosome 1 whereas *PUF4* is present on

chromosome 3, ruling out linkage as an explanation for this observation. Interestingly, we

recovered 11 progeny that were *gwo1Δ puf4Δ* double mutants from a reciprocal cross H99 *puf4Δ*

x JF289 *gwo1Δ* (11 out of 80), albeit at a lower frequency than expected (25%). The MIS assays

with the *gwo1Δ puf4Δ* double mutant revealed the MIS frequency at the same rate as that of the

gwo1Δ single mutant suggesting that Gwo1 plays a more dominant role in transgene silencing

than Puf4 (Figure 2A). Overall, these studies show that calcineurin and its known targets impact

RNAi-mediated transgene silencing through multiple pathways that may or may not be

connected with each other.

Calcineurin mutants produce less RNAi-mediated small RNA from the transgene

A previous study showed that transgene silencing is mediated by RNAi-mediated small-

interfering RNAs (siRNAs) (WANG *et al.* 2010). We therefore assessed if calcineurin and its

targets impact siRNA production from the transgene array. We first conducted long-read PacBio

sequencing to assemble the genome of the transgene array containing reporter strain, JF289. The

genome assembly revealed 15 tandem copies of the transgene integrated next to the endogenous

URA5 gene.   After assembling the complete transgene array, we conducted small RNA

sequencing from the wild-type transgene array strain JF289 as well as mutant strains.

We selected two 5-FOA resistant colonies (transgene silenced) obtained from the MIS

assays and one YPD colony (transgene expressed) from the wild-type (JF289) as well as mutants

that exhibited differences in MIS silencing (*cna1*Δ, *cnb1*Δ, *gwo1*Δ, and *puf4*Δ strains).

Following sRNA-sequencing, the reads were processed to trim adapters followed by mapping to

the reference genome sequence. This mapped small RNA population was first analyzed the

small RNA population for RNAi-mediated signatures such as RNA length and 5' nucleotide

preference. This analysis revealed that all strains are proficient in RNAi-mediated siRNA

production and exhibited a peak at 21-23 nucleotides with a 5'-Uracil preference, both signatures

of canonical RNAi processing (Figure 3). Interestingly, the *cna1*Δ and *cnb1*Δ mutants exhibited

a significant 30-40% reduction in this population of siRNA compared to the wild-type and other

two mutants (Table 2). These results confirm the role of calcineurin in small RNA production

and provide additional evidence for its role in transgene silencing.

Next, we mapped the small RNA reads to the JF289 genome and first analyzed the reads

mapping specifically to the transgene array (Figure 4). In accordance with the previous report

(WANG *et al.* 2010), we identified siRNA only against the *URA5* component of the transgene,

and no siRNA peaks were identified against the rest of the transgene including the *SXI2***a** gene.

More specifically, this siRNA population against the *URA5* gene was present only in the cultures

grown from 5-FOA colonies (*URA5* silenced), but not in the YPD colonies cultures (*URA5*

expressed) (Figure 4). The presence of RNAi-mediated siRNA population in these 5-FOA

colonies is expected because they are required for suppression of *URA5* expression, whereas

YPD colonies have not been selected for *URA5* silencing in accord with the lack of siRNAs.

Both *cna1Δ* and *cnb1Δ* mutants exhibited fewer siRNAs against *URA5* as compared to the wild-

type (Figure 4), in accordance with the overall reduction in RNAi-mediated siRNA population in

these mutants as described above. In our analysis, we mapped each read to only one location in

the genome and as a result all reads generated from the transgene locus are uniformly distributed

across all copies of *URA5*. It is possible that some *URA5* copies might produce more siRNA

than others but if it occurs it cannot be inferred from this analysis.

Further analysis revealed mutant-specific behavior of the siRNA population where both

gwo1Δ and *puf4Δ* mutants exhibited a change in siRNA distribution. *gwo1Δ* mutant isolates

produced more siRNA against the 3' end of the *URA5* gene and less against the 5' end as

compared to the wild-type transgene array strain JF289. On the other hand, *puf4Δ* mutant

isolates generated similar levels of siRNA against *URA5* to that of wild-type but additionally

produced siRNAs from the upstream region of the *URA5* gene spreading into the neighboring

gene. A similar pattern of siRNA was also observed in one of the *cnb1Δ* mutant 5-FOA isolates,

albeit at a lower level. These results reveal that calcineurin, Gwo1, and Puf4 regulate the siRNA

population in different manners affecting either the level or the profile of siRNAs against the

URA5 gene.

Genome-wide siRNA production is altered in calcineurin and gwo1Δ mutants

After analyzing the transgene-specific small RNAs, we next assessed whether calcineurin

mutants also affect small RNA production on a genome-wide level. Analysis of the small RNA

levels throughout the genome revealed that both the *cna1Δ* and *cnb1Δ* mutants produced fewer

siRNAs (Figure 5, S3 and Table 2). To quantify this, a differential expression analysis was

performed which revealed that both the *cna1Δ* and *cnb1Δ* mutants exhibited a change in the

production of siRNAs against several loci with the *cnb1Δ* mutation affecting at slightly more loci

(Table S3). This comparative analysis also showed that the *gwo1Δ* mutant also impacted a large

number of loci whereas the *puf4Δ* mutant altered siRNA levels from only a few loci, suggesting

a more dominant role for Gwo1 in RNAi in accord with results from the MIS assays.

Most genomic regions exhibited mutant-specific differential expression where the siRNA

population from certain loci showed changes across all samples, two from 5-FOA, and one from

YPD. However, some loci produced different levels of siRNA depending on the condition for

the same mutant. For example, siRNA production from a region in centromere 3 and an

intergenic region on chromosome 6 was different between YPD and 5-FOA samples in the *puf4Δ*

mutant (Figure 5, top and bottom panels). A more striking example was observed for the *gwo1Δ*

mutant which showed no siRNA from a locus on chromosome 8 in the 5-FOA condition and

abundant siRNA production from the same loci in the YPD condition (Figure S3, left side,

middle panel). While these regions do not exhibit any similarity to the *URA5* gene, whose

expression 5-FOA counter-selects, it is important to note that 5-FOA-containing media provides

a stressful growth condition and these siRNA changes observed may be due to this additional

stress resulting in epi-alleles at these loci.

The genome-wide analysis also revealed that the *gwo1Δ* deletion affects siRNA

production from certain loci more dramatically than others. For several loci, *gwo1Δ* deletion

caused a complete abolishment of siRNA production (Figure 5 and S3), whereas for many others

this led to an increase in siRNA production (compare various loci shown in the upper two panels

of Figure 5). This particular pattern was observed for multiple loci throughout the genome

showing that the *gwo1Δ* mutation has a significant impact on siRNA production. Overall,

siRNA sequencing analysis revealed that calcineurin and Gwo1 are major factors in the global

regulation of small RNAs in *C. neoformans*.

Discussion

In this study, we showed that calcineurin contributes to RNAi-mediated silencing as well

as the reactivation of a transgene array. Our results suggest that this role of calcineurin may be

exerted both through and independent of its known targets in P-bodies, and requires interactions

with the RNAi machinery (Figure 6). Interestingly, one of the known substrates, Gwo1, has

been identified in an earlier study (DUMESIC *et al.* 2013) as a component of the RNAi machinery

via its binding to Argonaute (Ago1), a core component of the RNAi machinery, and notably

showed the largest impact on transgene silencing both during MIS and SIS in our assays.

Interestingly, Dumesic et al had concluded that Gwo1 is not required for siRNA biogenesis

based on the analysis of three genomic loci. Contrary to that, our analysis identified that loss of

Gwo1 has a locus-specific impact and is required for siRNA biogenesis from certain genomic

loci but not others, unlike conventional RNAi proteins. We also identified that deletion of Puf4

results in enhanced silencing by approximately three-fold and this is the first gene whose

mutation has been identified to enhance silencing. Puf4 belongs to the Pumilio and Fem-3 (PUF)

family of mRNA binding factors and has been shown to bind to several RNAs in *C. neoformans*

(KALEM *et al.* 2021; KALEM *et al.* 2022). RNA interactions of Puf4 involve genes required for

cell wall synthesis, drug resistance, as well as regulating gene expression. More importantly,

Puf4 immuno-precipitation also revealed a direct interaction with Ago1 (KALEM *et al.* 2022). It

is possible that both Gwo1 and Puf4 controls siRNA loading on Ago1 and future studies would

explore siRNA cargo of Ago1 in the absence of these two proteins. Our results also indicate a

more global role for calcineurin than either Gwo1 or Puf4 indicating that calcineurin might be

regulating siRNA production and transgene silencing independently of these two substrates.

Such a global impact might involve either a direct interaction of calcineurin with the core RNAi

machinery or an indirect effect through one of its uncharacterized substrates that plays a major

role in RNAi function. Delineating these direct and indirect effects of calcineurin on RNAi-

mediated transgene silencing and siRNA production will be crucial to better understanding the

impact of calcineurin signaling in genome defense mechanisms.

One of the main findings of this study is the demonstration of a role for calcineurin in

reducing both silencing and un-silencing of the transgene array, revealing that calcineurin

maintains the flux between the silenced and expressed states. Three of four calcineurin

substrates analyzed in this study (Gwo1, Pbp1, and Vts1) do not impact the un-silencing of the

transgene array whereas Puf4 deletion resulted in the complete abolishment of transgene

reactivation. This result suggests that calcineurin could impact the transcription status of cells in

response to stress conditions in a previously unknown manner. A recent study in plants

identified a link between calcium signaling and antiviral RNAi defense (WANG *et al.* 2021).

This study showed that calcium signaling activated calmodulin-binding transcription activator-3

(CAMTA-3) to drive expression changes of key RNAi machinery components in plants. In our

study, we show a direct link between RNAi and calcineurin, a phosphatase that is also activated

by calcium and calmodulin. Whether this might be conserved in animals is unknown.

Combined, these studies indicate that calcium signaling might have a broader role in genome

defense mechanisms such as RNAi and might act through different downstream components in

different organisms.

Previous studies revealed a large group of genes differently regulated at 37°C in the

absence of Cna1 independent of Crz1, a transcription factor that is activated by calcineurin

(CHOW *et al.* 2017). How calcineurin governs the mRNA abundance of these genes independent

of Crz1 is not well understood but might involve post-transcriptional regulation in P-bodies.

First, calcineurin activity may be essential for the processing of certain mRNAs, and in its

absence, those mRNAs might be either degraded or stored for longer times resulting in an

alteration in RNA levels. A second possible mechanism of calcineurin regulation might be

through its role in RNAi where calcineurin might be responsible for degrading certain mRNAs

via RNAi at high temperatures. The third possible role of calcineurin in regulating RNA

metabolism might be through a direct role in the nucleus. Calcineurin localizes to the nucleus in

mammalian cells, and this localization varies among different calcineurin isoforms (SHIBASAKI *et*

al. 1996; JABR *et al.* 2007). While calcineurin has not as yet been reported to localize to the

nucleus in *C. neoformans*, it cannot be excluded as a possible mechanism of action. Lastly, it is

also possible that calcineurin regulates other transcription factors in addition to Crz1 that drive

the expression of a different set of genes. Our experiments employing small RNA sequencing

revealed that calcineurin has a significant impact on siRNA production from hundreds of

genomic loci providing evidence for the second proposed hypothesis above. However, more

than 30% of these affected loci are non-coding RNAs, and siRNA changes could account for

only 29 genes whose expression is differentially regulated in the calcineurin mutant based on

RNA-sequencing, suggesting that calcineurin's role on RNA metabolism could be through

multiple independent regulatory processes. Additionally, it is possible that the impact of

calcineurin on siRNA production is more enhanced at 37°C and could account for a larger

number of genes whose expression is altered upon heat stress. No study has analyzed siRNA

levels at different temperatures making it a subject of investigation for future studies to better

understand the roles of the RNAi machinery in heat stress responses in *C. neoformans*.

RNAi-mediated silencing is known for its roles in genome integrity and defense

mechanisms (OBBARD *et al.* 2009; DUMESIC *et al.* 2013; YADAV *et al.* 2018; PRIEST *et al.* 2022).

A role for calcineurin in RNAi regulation suggests that calcineurin might also be contributing to

genome integrity in ways that have not been defined. This is further strengthened by recent

studies in humans where calcineurin roles at nuclear pore complexes and centrosomes were

discovered (WIGINGTON *et al.* 2020; TSEKITSIDOU *et al.* 2023). In *Schizosaccharomyces pombe*,

RNAi is required for centromere function and if calcineurin has a similar role in regulating RNAi

in this fission yeast, it might be directly contributing to genome maintenance and nuclear

division (VOLPE *et al.* 2003; FOLCO *et al.* 2008). Similarly, RNAi regulates centromere structure

in *C. neoformans* (YADAV *et al.* 2018), and whether calcineurin also plays a role will be of

interest to study. Whether calcineurin has broadly conserved or species-specific roles in genome

integrity in other organisms will be an interesting avenue to pursue, especially given that no clear

link has been established between calcineurin and genome defense mechanisms.

Our results also suggest that processes like RNAi can be affected by environmental

stimuli through signaling pathways such as calcineurin. This not only expands the repertoire of

factors that affect genome defense mechanisms but also provides important insights on the

mechanisms of RNAi-associated processes. For example, epimutations that drive drug resistance

in some fungi might have an origin in stress-induced signaling that is initiated in response to

infection (CALO *et al.* 2014; CHANG *et al.* 2019; CHANG AND HEITMAN 2019). In *C. neoformans*,

transgene silencing occurs at a much higher rate during meiosis than in mitosis (WANG *et al.*

2010). While it has been shown that RNAi is more active during meiosis, what triggers this

change remains to be identified. It is possible that environmental stimuli play an important role

in this switch and signaling pathways such as calcineurin may play a role. Calcineurin in *C.*

neoformans is not required for vegetative growth at 25°C and cells lacking calcineurin behave

like the wild-type at this temperature. However, when calcineurin mutant cells are subjected to

mating that also occurs at 25°C, they fail to produce filaments revealing the impact of calcineurin

in response to mating-inducing cues specifically (CRUZ *et al.* 2001). Previous studies have also

identified that the RNAi machinery is activated during the early stages of mating itself, possibly

in response to the same environmental stimuli that triggers the calcineurin pathway (WANG *et al.*

2010). Whether there is a direct connection between calcineurin and RNAi during mating and

later stages of sexual reproduction remains to be explored.

Overall, our study opens avenues for exploration with respect to the roles of calcineurin

in gene expression, maintaining genome integrity, as well as linking environmental stimuli with

processes such as RNAi. In this study, we provide a connection between calcineurin signaling

and RNAi-mediated silencing that cannot be explained entirely through known substrates of

calcineurin. Thus, a better understanding of calcineurin signaling is required to fully understand

its roles in *C. neoformans* biology, specifically in stress-response pathways.

Data Availability Statement

All of the sequencing data generated in this study including the genome assembly

generated for JF289 and small RNA sequencing data has been submitted to NCBI under the

BioProject PRJNA996625. The genome assembly and annotations are also available via the

figshare repository (DOI: 10.6084/m9.figshare.24975069).

Funding

This work was supported by NIH/NIAID R01 awards AI039115-26, AI050113-19, and

AI170543-02 awarded to JH, and R01 grant AI33654-06 awarded to JH and Paul Magwene. JH

is also a Co-Director and Fellow of the CIFAR program Fungal Kingdom: Threats &

Opportunities. This work was funded in part by a 2022 developmental grant to VY from the

Duke University Center for AIDS Research (CFAR), an NIH-funded program (5P30 AI064518).

References

Boutros, M., and J. Ahringer, 2008 The art and design of genetic screens: RNA interference. Nat
Rev Genet 9**:** 554-566.

Buchon, N., and C. Vaury, 2006 RNAi: A defensive RNA-silencing against viruses and
transposable elements. Heredity 96**:** 195-202.

Burke, J. E., A. D. Longhurst, P. Natarajan, B. Rao, J. Liu *et al.*, 2019 A non-dicer RNase III and
four other novel factors required for RNAi-mediated transposon suppression in the
human pathogenic yeast *Cryptococcus neoformans*. G3 9**:** 2235-2244.

Calo, S., C. Shertz-Wall, S. C. Lee, R. J. Bastidas, F. E. Nicolas *et al.*, 2014 Antifungal drug
resistance evoked via RNAi-dependent epimutations. Nature 513**:** 555-558.

Castel, S. E., and R. A. Martienssen, 2013 RNA interference in the nucleus: Roles for small
RNAs in transcription, epigenetics and beyond. Nat Rev Genet 14**:** 100-112.

Chang, Z., R. B. Billmyre, S. C. Lee and J. Heitman, 2019 Broad antifungal resistance mediated
by RNAi-dependent epimutation in the basal human fungal pathogen *Mucor*
circinelloides. PLoS Genetics 15**:** e1007957.

Chang, Z., and J. Heitman, 2019 Drug-resistant epimutants exhibit organ-specific stability and
induction during murine infections caused by the human fungal pathogen *Mucor*
circinelloides. mBio 10.

Chow, E. W., S. A. Clancey, R. B. Billmyre, A. F. Averette, J. A. Granek *et al.*, 2017
Elucidation of the calcineurin-Crz1 stress response transcriptional network in the human
fungal pathogen *Cryptococcus neoformans*. PLoS Genetics 13**:** e1006667.

Cogoni, C., and G. Macino, 1999 Gene silencing in *Neurospora crassa* requires a protein
homologous to RNA-dependent RNA polymerase. Nature 399**:** 166-169.

Cruz, M. C., D. S. Fox and J. Heitman, 2001 Calcineurin is required for hyphal elongation during
mating and haploid fruiting in *Cryptococcus neoformans*. EMBO J 20**:** 1020-1032.

Dumesic, P. A., P. Natarajan, C. Chen, I. A. Drinnenberg, B. J. Schiller *et al.*, 2013 Stalled
spliceosomes are a signal for RNAi-mediated genome defense. Cell 152**:** 957-968.

Feretzaki, M., R. B. Billmyre, S. A. Clancey, X. Wang and J. Heitman, 2016 Gene network
polymorphism illuminates loss and retention of novel RNAi silencing components in the
Cryptococcus pathogenic species complex. PLoS Genetics 12**:** e1005868.

Folco, H. D., A. L. Pidoux, T. Urano and R. C. Allshire, 2008 Heterochromatin and RNAi are
required to establish CENP-A chromatin at centromeres. Science 319**:** 94-97.

Fox, D. S., M. C. Cruz, R. A. Sia, H. Ke, G. M. Cox *et al.*, 2001 Calcineurin regulatory subunit
is essential for virulence and mediates interactions with FKBP12-FK506 in *Cryptococcus*
neoformans. Mol Microbiol 39**:** 835-849.

Jabr, R. I., A. J. Wilson, M. H. Riddervold, A. H. Jenkins, B. A. Perrino and L. H. Clapp, 2007
Nuclear translocation of calcineurin Aβ but not calcineurin Aα by platelet-derived growth
factor in rat aortic smooth muscle. Am J Physiol Cell Physiol 292**:** C2213-2225.

Janbon, G., S. Maeng, D. H. Yang, Y. J. Ko, K. W. Jung *et al.*, 2010 Characterizing the role of
RNA silencing components in *Cryptococcus neoformans*. Fungal Genet Biol 47**:** 1070-
1080.

Kalem, M. C., S. Duffy, S. Shen, J. N. Kaur, J. Qu and J. C. Panepinto, 2022 Arginine
methylation of Puf4 drives diverse protein functions. bioRxiv**:** 2022.2006.2021.497104.

Kalem, M. C., H. Subbiah, J. Leipheimer, V. E. Glazier and J. C. Panepinto, 2021 Puf4 mediates
post-transcriptional regulation of cell wall biosynthesis and caspofungin resistance in
Cryptococcus neoformans. mBio 12**:** e03225-03220.

Kozubowski, L., E. F. Aboobakar, M. E. Cardenas and J. Heitman, 2011 Calcineurin colocalizes
with P-bodies and stress granules during thermal stress in *Cryptococcus neoformans*.
Eukaryot Cell 10**:** 1396-1402.

Lax, C., G. Tahiri, J. A. Patino-Medina, J. T. Canovas-Marquez, J. A. Perez-Ruiz *et al.*, 2020
The evolutionary significance of RNAi in the fungal kingdom. Int J Mol Sci 21.

Mangus, D. A., N. Amrani and A. Jacobson, 1998 Pbp1p, a factor interacting with
Saccharomyces cerevisiae poly(A)-binding protein, regulates polyadenylation. Mol Cell
Biol 18**:** 7383-7396.

Nakayashiki, H., 2005 RNA silencing in fungi: mechanisms and applications. FEBS Lett 579**:**
5950-5957.

Nicolas, F. E., and V. Garre, 2016 RNA interference in fungi: Retention and loss. Microbiol
Spectr 4.

Obbard, D. J., K. H. Gordon, A. H. Buck and F. M. Jiggins, 2009 The evolution of RNAi as a
defence against viruses and transposable elements. Philos Trans R Soc Lond B Biol Sci
364**:** 99-115.

Odom, A., S. Muir, E. Lim, D. L. Toffaletti, J. Perfect and J. Heitman, 1997 Calcineurin is
required for virulence of *Cryptococcus neoformans*. EMBO J 16**:** 2576-2589.

Park, H. S., E. W. Chow, C. Fu, E. J. Soderblom, M. A. Moseley *et al.*, 2016 Calcineurin targets
involved in stress survival and fungal virulence. PLoS Pathogens 12**:** e1005873.

Park, H. S., S. C. Lee, M. E. Cardenas and J. Heitman, 2019 Calcium-calmodulin-calcineurin
signaling: A globally conserved virulence cascade in eukaryotic microbial pathogens.
Cell Host Microbe 26**:** 453-462.

Perfect, J. R., N. Ketabchi, G. M. Cox, C. W. Ingram and C. L. Beiser, 1993 Karyotyping of
Cryptococcus neoformans as an epidemiological tool. J Clin Microbiol 31**:** 3305-3309.

Priest, S. J., V. Yadav, C. Roth, T. A. Dahlmann, U. Kuck *et al.*, 2022 Uncontrolled transposition
following RNAi loss causes hypermutation and antifungal drug resistance in clinical
isolates of *Cryptococcus neoformans*. Nat Microbiol 7**:** 1239-1251.

Rajasingham, R., N. P. Govender, A. Jordan, A. Loyse, A. Shroufi *et al.*, 2022 The global burden
of HIV-associated cryptococcal infection in adults in 2020: A modelling analysis. Lancet
Infect Dis 22**:** 1748-1755.

Rajasingham, R., R. M. Smith, B. J. Park, J. N. Jarvis, N. P. Govender *et al.*, 2017 Global burden
of disease of HIV-associated cryptococcal meningitis: an updated analysis. Lancet Infect
Dis 17**:** 873-881.

Rendl, L. M., M. A. Bieman and C. A. Smibert, 2008 *S. cerevisiae* Vts1p induces deadenylation-
dependent transcript degradation and interacts with the Ccr4p-Pop2p-Not deadenylase
complex. RNA 14**:** 1328-1336.

Rusnak, F., and P. Mertz, 2000 Calcineurin: form and function. Physiol Rev 80**:** 1483-1521.

Shabalina, S. A., and E. V. Koonin, 2008 Origins and evolution of eukaryotic RNA interference.
Trends Ecol Evol 23**:** 578-587.

Shibasaki, F., E. R. Price, D. Milan and F. McKeon, 1996 Role of kinases and the phosphatase
calcineurin in the nuclear shuttling of transcription factor NF-AT4. Nature 382**:** 370-373.

Sun, S., S. J. Priest and J. Heitman, 2019 *Cryptococcus neoformans* mating and genetic crosses.
Curr Protoc Microbiol 53**:** e75.

Tsekitsidou, E., C. J. Wong, I. Ulengin-Talkish, A. I. M. Barth, T. Stearns *et al.*, 2023
Calcineurin associates with centrosomes and regulates cilia length maintenance. J Cell
Sci 136.

Ulengin-Talkish, I., and M. S. Cyert, 2023 A cellular atlas of calcineurin signaling. Biochim
Biophys Acta Mol Cell Res 1870**:** 119366.

Volpe, T., V. Schramke, G. L. Hamilton, S. A. White, G. Teng *et al.*, 2003 RNA interference is
required for normal centromere function in fission yeast. Chromosome Res 11**:** 137-146.

Wang, X., Y. P. Hsueh, W. Li, A. Floyd, R. Skalsky and J. Heitman, 2010 Sex-induced silencing
defends the genome of *Cryptococcus neoformans* via RNAi. Genes Dev 24**:** 2566-2582.

Wang, X., P. Wang, S. Sun, S. Darwiche, A. Idnurm and J. Heitman, 2012 Transgene induced
co-suppression during vegetative growth in *Cryptococcus neoformans*. PLoS Genetics 8**:**
e1002885.

Wang, Y., Q. Gong, Y. Wu, F. Huang, A. Ismayil *et al.*, 2021 A calmodulin-binding
transcription factor links calcium signaling to antiviral RNAi defense in plants. Cell Host
Microbe 29**:** 1393-1406 e1397.

Wigington, C. P., J. Roy, N. P. Damle, V. K. Yadav, C. Blikstad *et al.*, 2020 Systematic
discovery of short linear motifs decodes calcineurin phosphatase signaling. Mol Cell 79**:**
342-358 e312.

Yadav, V., and J. Heitman, 2023 Calcineurin: The Achilles' heel of fungal pathogens. PLoS
Pathog 19**:** e1011445.

Yadav, V., S. Sun, R. B. Billmyre, B. C. Thimmappa, T. Shea *et al.*, 2018 RNAi is a critical
determinant of centromere evolution in closely related fungi. Proc Natl Acad Sci U S A
115**:** 3108-3113.

Yadav, V., S. Sun, M. A. Coelho and J. Heitman, 2020 Centromere scission drives chromosome
shuffling and reproductive isolation. Proc Natl Acad Sci U S A 117**:** 7917-7928.

Zhao, Y., J. Lin, Y. Fan and X. Lin, 2019 Life cycle of *Cryptococcus neoformans*. Annu Rev
Microbiol 73**:** 17-42.

Figures legends

Figure 1.   Calcineurin regulates the onset and reversion of transgene silencing. (A) A cartoon depicting the *URA5* transgene array in the reporter strain JF289. **(B)** A schematic showing the workflow to determine the mitotic-induced silencing (MIS) frequency with the reporter strain. **(C)** Plate photos showing the difference in colony morphology for the wild-type JF289, *cna1Δ*, *cnb1Δ*, and JF289 plated on FK506 containing media when an approximately equal number of colonies were obtained in each case after 10 days of incubation. **(D)** A graph showing the MIS frequency of calcineurin mutants as compared to the JF289 strain. P-value (**) < 0.01. Error bars represent the standard error of the mean (SEM). **(E)** Images of petri-dishes showing the growth of colonies on media lacking uracil (SD-URA) directly from 5-FOA-containing media during MIS assays. The numbers below represent the average percent of colonies growing (Mean ± SEM) from three independent experiments.

Figure 2.   Calcineurin substrates play diverged roles during transgene silencing. (A) A graph showing the MIS frequency of mutants for known calcineurin targets (Gwo1, Pbp1, Puf4, and Vts1) as well as double deletion mutants of genes that showed an effect on MIS (p-values: ** < 0.01, **** < 0.0001). The double deletion mutants were not found to be significant as compared to their respective mutants (p-value >0.05). **(B)** Photographs of petri-dishes showing the colony number variation in different mutants, as compared to JF289 after 10 days of incubation when an estimated equal number of cells were plated in each case. **(C)** Images showing the growth of individual colonies on SD-URA media from 5-FOA containing media for MIS reversion assays. The numbers represent the reversion rates for each mutant.

Figure 3. Calcineurin and its substrates are not essential for RNAi-mediated siRNA

production. Graphs showing the (A) length distribution and (B) 5' nucleotide preference of

small RNAs in the wild-type (JF289) and mutant strains. The isolates analyzed for the small

RNA preparation were from both non-selective (YPD) and selective (5-FOA) conditions for the

MIS assays.

Figure 4. Calcineurin mutants produce less siRNA against the transgene array. A series of

genomic location maps showing the status of siRNA against the transgene array is presented with

the top panel showing the full chromosome 8 on which the transgene array is integrated. The

middle panel shows a zoomed view of the entire transgene array and mapping of siRNA from the

wild-type as well as various mutants. The lower panel depicts a single unit of transgene array

including the location of *URA5* and *SXI2***a** genes. *URA5* outside of the transgene array represents

the endogenous copy of *URA5* that was targeted for transgene insertion. The mapped siRNA

data is presented in the same order across all three panels and is labeled in the lower panel. The

siRNA mapping revealed a lower level of siRNA in both *cna1*Δ and *cnb1*Δ mutants whereas a

difference in siRNA distribution was observed in *gwo1*Δ and *puf4*Δ mutants.

Figure 5. Calcineurin and Gwo1 regulate genome-wide siRNA production differently.

Snapshots of various genomic regions showing small RNA reads mapping for various strains

including calcineurin mutants. The top panel depicts a centromere from chromosome 3, whereas

other panels present two other genomic loci. These regions show similarities and differences in

the siRNA distribution profiles for the wild-type and mutant strains. Also see Figure S3.

Figure 6. A model showing the role of calcineurin, and its known substrates, Gwo1 and

Puf4, in regulating RNAi-mediated small RNA production.

Supplementary figure legends

Figure S1. Growth rate in calcineurin mutants does not impact their MIS frequency. (A)

Plate pictures showing the growth for the wild-type (JF289) and calcineurin mutants in both the

nutrient-rich YPD media and nutrient limiting standard defined (SD) media. The viability

percent is listed for each strain in both media conditions. **(B)** A graph showing the MIS

frequency and a table presenting the MIS reversion rate in the wild-type (WT), *cna1Δ* and *cnb1Δ*

mutants at two different time points from the same culture (p-values: * < 0.05, *** < 0.001, ns >

0.05).

Figure S2. Calcineurin activity is not required for the P-body localization of its substrates.

(A) Microscopy images showing the co-localization of the mCherry-tagged version of

calcineurin and its substrates with GFP-Dcp1, a P-body marker, at 37°C in the absence and

presence of calcineurin inhibitor, FK506. Bar, 5 µm. Some of the co-localization events are

highlighted with white arrowheads in all cases. **(B)** A table describing the analysis of the

number of cells showing co-localization with the P-body marker, GFP-Dcp1, in the presence and

absence of FK506.

Figure S3. Loss of calcineurin leads to reduced siRNA production across multiple genomic

loci. Small RNA reads mapping revealed a lower level of siRNA against multiple different

regions of the genome in calcineurin mutants and a complete abolishment of siRNA against

some genomic regions in the *gwo1*Δ mutant.

Table 1. SIS frequency for JF289, *vts1*Δ, and *gwo1*Δ mutant crosses.

Cross	Viable progeny obtained	Progeny inheriting transgene array		SIS frequency
		Total progeny	Growth on 5-FOA	
H99α x JF289**a**	71	36	16	44.4%
H99α *vts1*Δ x JF289**a** *vts1*Δ	59	27	9	33.3%
H99α *gwo1*Δ x JF289**a** *gwo1*Δ	92	49	0	0%

739 **Table 2. siRNA population levels in the wild-type and various mutants.**

Condition	siRNA length	Percent of reads in siRNA population				
		Wild-type	*cna1*Δ	*cnb1*Δ	*gwo1*Δ	*puf4*Δ
YPD	21-nt	7.30	4.08	3.48	8.06	5.57
	22-nt	15.15	8.02	6.66	15.30	10.02
	23-nt	8.58	5.40	4.55	8.02	5.34
	Total	31.03	17.50	14.69	31.38	20.93
5-FOA-1	21-nt	8.68	6.01	5.78	7.33	8.56
	22-nt	16.57	11.39	10.62	13.89	15.54
	23-nt	8.54	6.59	6.13	6.33	7.82
	Total	33.79	23.99	22.53	27.55	31.92
5-FOA-2	21-nt	9.69	6.52	5.94	8.68	9.65
	22-nt	17.85	12.47	11.19	17.14	16.88
	23-nt	8.86	7.29	6.32	8.04	7.07
	Total	36.40	26.28	23.45	33.86	33.60
Mean ± SEM of total siRNA/condition		33.74 ±1.55	22.59 ±2.63	20.22 ±2.78	30.93 ±1.84	28.81 ±3.97
Significant; p-value (Compared to wild-type)			Yes; 0.011	Yes; 0.012	No; 0.278	No; 0.199

741 **Supplementary tables**

743 **Table S1. List of strains used in this study.**

745 **Table S2. List of primers used in this study.**

747 **Table S3. Differential expression analysis of small RNA reads.**

749 **Table S4. Growth rates of wild-type and mutant strains in different media.**

A Transgene array x15

Native *URA5* copy

B Isolate single colonies on SD-Uracil

Overnight cultures in YPD (x5)

Dilute and plate cells

5-FOA

YPD

$$\text{MIS frequency} = \frac{\text{\# colonies on 5-FOA plate}}{\text{\# colonies on YPD plate} \times 10^3}$$

C Wild-type (WT)   *cna1Δ*   *cnb1Δ*   WT on FK506

D

MIS frequency

1.5×10^{-3}

1×10^{-3}

5×10^{-4}

**

WT   *cna1Δ*   *cnb1Δ*   WT on FK506   *rdp1Δ*

E

WT   *cna1Δ*   *cnb1Δ*   WT on FK506

43 ± 5%   14 ± 3%   15 ± 2%   14 ± 5%

A

MIS frequency

B MIS assay

WT

gwo1Δ   puf4Δ

pbp1Δ   vts1Δ

C MIS Reversion

gwo1Δ = 39%   pbp1Δ = 32%   puf4Δ = 0%   vts1Δ = 30%

Chromosome 8

Transgene array

Transgene unit

	Genes/Exons
	ncRNA/Transgene

0-134	WT - 5-FOA-1
0-130	WT - 5-FOA-2
0-153	WT - YPD
0-201	*cna1Δ* - 5-FOA-1
0-139	*cna1Δ* - 5-FOA-2
0-141	*cna1Δ* - YPD
0-153	*cnb1Δ* - 5-FOA-1
0-167	*cnb1Δ* - 5-FOA-2
0-152	*cnb1Δ* - YPD
0-287	*gwo1Δ* - 5-FOA-1
0-142	*gwo1Δ* - 5-FOA-2
0-198	*gwo1Δ* - YPD
0-167	*puf4Δ* - 5-FOA-1
0-150	*puf4Δ* - 5-FOA-2
0-103	*puf4Δ* - YPD