Complementary opposing D2-MSNs and D1-MSNs dynamics during interval timing

R. Austin Bruce[1], Matthew A. Weber[1], Alexandra S. Bova[1], Rachael A. Volkman[1], Casey E. Jacobs[1], Kartik Sivakumar[1], Hannah R Stutt[1], Young-cho Kim[1], Rodica Curtu[2,3], Nandakumar S. Narayanan[1,3]

[1]Department of Neurology, University of Iowa, [2]Department of Mathematics, University of Iowa, and [3]The Iowa Neuroscience Institute

*Corresponding Author
Nandakumar Narayanan
nandakumar-narayanan@uiowa.edu
169 Newton Road
Pappajohn Biomedical Discovery Building—5336
University of Iowa, Iowa City, 52242
319-353-5698

Abstract

The role of striatal pathways in cognitive processing is unclear. We studied dorsomedial striatal cognitive processing during interval timing, an elementary cognitive task that requires mice to estimate intervals of several seconds and involves working memory for temporal rules as well as attention to the passage of time. We harnessed optogenetic tagging to record from striatal D2-dopamine receptor-expressing medium spiny neurons (D2-MSNs) in the indirect pathway and from D1-dopamine receptor-expressing MSNs (D1-MSNs) in the direct pathway. We found that D2-MSNs and D1-MSNs exhibited distinct dynamics over temporal intervals as quantified by principal component analyses and trial-by-trial generalized linear models. MSN recordings helped construct and constrain a four-parameter drift-diffusion computational model. This model predicted that disrupting either D2-MSNs or D1-MSNs would increase interval timing response times and alter MSN firing. In line with this prediction, we found that optogenetic inhibition or pharmacological disruption of either D2-MSNs or D1-MSNs increased interval timing response times. Pharmacologically disrupting D2-MSNs or D1-MSNs also changed MSN dynamics and degraded trial-by-trial temporal decoding. Together, our findings demonstrate that D2-MSNs and D1-MSNs make complementary contributions to interval timing despite opposing dynamics, implying that striatal direct and indirect pathways work together to shape temporal control of action. These data provide novel insight into basal ganglia cognitive operations beyond movement and have implications for human striatal diseases and therapies targeting striatal pathways.

Introduction

The basal ganglia have been heavily studied in motor control (Albin et al., 1989). These subcortical nuclei are composed of two pathways defined by medium spiny neurons (MSNs) in the striatum: the indirect pathway, defined by D2-dopamine receptor-expressing MSNs (D2-MSNs), and the direct pathway, defined by D1-dopamine receptor-expressing MSNs (D1-MSNs). It has been proposed that the indirect and direct pathways play opposing roles in movement (Alexander and Crutcher, 1990; Cruz et al., 2022; Kravitz et al., 2010), although recent work identified how these pathways can play more complex and complementary roles (Cui et al., 2013; Tecuapetla et al., 2016). MSNs receive dense input from both motor and cognitive cortical areas (Averbeck et al., 2014; Graybiel, 1997; Middleton and Strick, 2000), but the respective roles of D2-MSNs and D1-MSNs in cognitive processing are largely unknown. Understanding basal ganglia cognitive processing is critical for diseases affecting the striatum such as Huntington's disease, Parkinson's disease, and schizophrenia (Andreasen, 1999; Hinton et al., 2007; Narayanan and Albin, 2022). Furthermore, pharmacological and brain stimulation therapies directly modulate D2-MSNs, D1-MSNs, and downstream basal ganglia structures such as the globus pallidus or subthalamic nucleus. Determining basal ganglia pathway dynamics during cognitive processes will help avoid side effects of current treatments and inspire novel therapeutic directions.

We studied striatal D2-MSNs and D1-MSNs during an elementary cognitive task, interval timing, which requires estimating an interval of several seconds. Interval timing provides an ideal platform to study cognition in the striatum because timing 1) requires cognitive resources including working memory for temporal rules and attention to the passage of time

(Parker et al., 2013); 2) is reliably impaired in human striatal diseases such as Huntington's

disease, Parkinson's disease, and schizophrenia (Merchant and de Lafuente, 2014); 3) requires

nigrostriatal dopamine that modulates MSNs (Emmons et al., 2017; Gouvea et al., 2015; Matell

et al., 2003; Mello et al., 2015; Monteiro et al., 2023; Wang et al., 2018); and 4) can be

rigorously studied in animal models (Balci et al., 2008; Buhusi and Meck, 2005; Kim et al.,

2017a; Parker et al., 2017, 2015). We and others have found that striatal MSNs encode time

across multiple intervals by time-dependent ramping activity or monotonic changes in firing rate

across a temporal interval (Emmons et al., 2017; Gouvea et al., 2015; Mello et al., 2015; Wang et

al., 2018). However, the respective roles of D2-MSNs and D1-MSNs are unknown. Past work

has shown that disrupting either D2-dopamine receptors (D2) or D1-dopamine receptors (D1)

powerfully impairs interval timing by increasing estimates of elapsed time (Drew et al., 2007;

Meck, 2006). Similar behavioral effects were found with systemic (Stutt et al., 2024) or local

dorsomedial striatal D2 or D1 disruption (De Corte et al., 2019a). These data lead to the

hypothesis that D2 MSNs and D1 MSNs have similar patterns of ramping activity across a

temporal interval.

We tested this hypothesis with a combination of optogenetics, neuronal ensemble

recording, computational modeling, and behavioral pharmacology. We use a well-described

mouse-optimized interval timing task (Balci et al., 2008; Bruce et al., 2021; Larson et al., 2022;

Stutt et al., 2024; Tosun et al., 2016; Weber et al., 2023). Strikingly, optogenetic tagging of D2-

MSNs and D1-MSNs revealed distinct neuronal dynamics, with D2-MSNs tending to increase

firing over an interval and D1-MSNs tending to decrease firing over the same interval, similar to

opposing movement dynamics (Cruz et al., 2022; Kravitz et al., 2010; Tecuapetla et al., 2016).

MSN dynamics helped construct and constrain a four-parameter drift-diffusion computational

model of interval timing, which predicted that disrupting either D2-MSNs or D1-MSNs would

increase interval timing response times. Accordingly, we found that optogenetic inhibition of

either D2-MSNs or D1-MSNs increased interval timing response times. Furthermore,

pharmacological blockade of either D2- or D1-receptors also increased response times and

degraded trial-by-trial temporal decoding from MSN ensembles. Thus, D2-MSNs and D1-MSNs

have opposing temporal dynamics yet disrupting either MSN type produced similar effects on

behavior. These data demonstrate how striatal pathways play complementary roles in elementary

cognitive operations and are highly relevant for understanding the pathophysiology of human

diseases and therapies targeting the striatum.

RESULTS

Mouse-optimized interval timing

We investigated cognitive processing in the striatum using a well-described mouse-optimized interval timing task which requires mice to respond by switching between two nosepokes after a 6-second interval (Fig 1A; see Methods; (Balci et al., 2008; Bruce et al., 2021; Larson et al., 2022; Tosun et al., 2016; Weber et al., 2023)). In this task, mice initiate trials by responding at a back nosepoke, which triggers auditory and visual cues for the duration of the trial. On 50% of trials, mice were rewarded for nosepoking after 6 seconds at the designated 'first' front nosepoke; these trials were not analyzed. On the remaining 50% of trials, mice were rewarded for nosepoking at the 'first' nosepoke and then switching to the 'second' nosepoke; initial nosepokes at the second nosepoke after 18 seconds triggered reward when preceded by a first nosepoke. The first nosepokes occurred before switching responses and the second nosepokes occurred much later in the interval in anticipation of reward delivery at 18 seconds (Fig 1B-D). During the task, movement velocity peaked before 6 seconds as mice traveled to the front nosepoke (Fig 1E).

We focused on the *switch response time*, defined as the moment mice exited the first nosepoke before entering the second nosepoke. Switch responses are a time-based decision guided by temporal control of action because mice switch nosepokes only if nosepoking at the first nosepokes does not lead to a reward after 6 seconds (Fig 1B-E). Switch responses are guided by internal estimates of time because no external cue indicates when to switch from the first to the second nosepoke (Balci et al., 2008; Bruce et al., 2021; Tosun et al., 2016; Weber et al., 2023). We defined the first 6 seconds after trial start as the 'interval', because during this epoch mice are estimating whether 6 seconds have elapsed and if they need to switch responses.

139   In 30 mice, switch response times were 9.3 seconds (8.4 – 9.7; median (IQR)); see Table 1 for a

140   summary of mice, experiments, trials, and sessions). We studied dorsomedial striatal D2-MSNs

141   and D1-MSNs using a combination of optogenetics and neuronal ensemble recordings in 9

142   transgenic mice (4 D2-Cre mice switch response time 9.7 (7.0 – 10.3) seconds; 5 D1-Cre mice

143   switch response time 8.2 (7.7 – 8.7) seconds; rank sum $p = 0.73$; Table 1).

Figure 1: Mouse-optimized interval timing. A) We trained mice to perform an interval timing task in which they had to respond by switching nosepokes after an ~6 second interval (in gray shade in all panels). Mice start trials by making a back nosepoke, which triggers an auditory and visual cue. On 50% of trials, mice were rewarded for a nosepoke after 6 seconds at the designated 'first' front nosepoke; these trials were not analyzed. On the remaining 50% of trials, mice were rewarded for switching to the 'second' nosepoke.; initial nosepokes at the second nosepoke after 18 seconds triggered reward when preceded by a first nosepoke. *Switch response time* was defined as the moment mice depart the first nosepoke prior to second nosepoke arrival. Because cues are identical and on for the full trial on all trials, switch responses are a time-based decision guided explicitly by temporal control of action; indeed, mice switch nosepokes only if they don't receive a reward at the first nosepokes after the 6 second interval. Top row - screen captures from the operant chambers during a trial with switch response. B) Response probability distribution from 30 mice for first nosepokes (purple), switch responses (green), and second nosepokes (orange). Responses at the first nosepoke peaked at 6 seconds, and switch responses peaked after 6 seconds. Because nosepoking at the second nosepoke was only rewarded after 18 seconds, second nosepokes tended to be highly skewed. Shaded area is standard error. C) Cumulative switch response density for each of 30 mice. D) Average cumulative switch response density; shaded area is standard error. E) DeepLabCut tracking of position during interval timing from a single mouse behavioral session revealed increased velocity after trial start and then constant velocity throughout the trial. Shaded area is standard error. In B-E, the 6 second interval is indicated in gray.

Opposing D2-MSN and D1-MSN dynamics

Striatal neuronal populations are largely composed of MSNs expressing D2-dopamine or

D1-dopamine receptors. We optogenetically tagged D2-MSNs and D1-MSNs by implanting

optrodes in the dorsomedial striatum and conditionally expressing channelrhodopsin (ChR2; Fig

S1) in 4 D2-Cre (2 female) and 5 D1-Cre transgenic mice (2 female). This approach expressed

ChR2 in D2-MSNs or D1-MSNs, respectively (Fig 2A-B; Kim et al., 2017a). We identified D2-

MSNs or D1-MSNs by their response to brief pulses of 473 nm light; neurons that fired within 5

milliseconds were considered optically tagged putative D2-MSNs (Fig S1B-C). We tagged 32

putative D2-MSNs and 41 putative D1-MSNs in a single recording session during interval

timing. There were no consistent differences in overall firing rate between D2-MSNs and D1-

MSNs (D2-MSNs: 3.4 (1.4 – 7.2) Hz; D1-MSNs 5.2 (3.1 – 8.6) Hz; F = 2.7, p = 0.11 accounting

for variance between mice). Peri-event rasters and histograms from a tagged putative D2-MSN

(Fig 2C) and from a tagged putative D1-MSN (Fig 2D) demonstrate prominent modulations for

the first 6 seconds of the interval after trial start. Z-scores of average peri-event time histograms

(PETHs) from 0 to 6 seconds after trial start for each putative D2-MSN are shown in Fig 2E and

for each putative D1-MSN in Fig 2F. These PETHs revealed that for the 6-second interval

immediately after trial start, many putative D2-MSN neurons appeared to ramp up while many

putative D1-MSNs appeared to ramp down. For 32 putative D2-MSNs average PETH activity

increased over the 6-second interval immediately after trial start, whereas for 41 putative D1-

MSNs, average PETH activity decreased. These differences resulted in distinct activity early in

the interval (0-1 seconds; F = 6.0, p = 0.02 accounting for variance between mice), but not late in

the interval (5-6 seconds; F = 1.9, p = 0.17 accounting for variance between mice) between D2-

MSNs and D1-MSNs. Examination of a longer interval of 10 seconds before to 18 seconds after

trial start revealed the greatest separation in D2-MSN and D1-MSN dynamics during the 6-

second interval after trial start (Fig S2). Strikingly, these data suggest that D2-MSNs and D1-

MSNs might display opposite dynamics during interval timing.

Figure 2: D2-MSNs and D1-MSNs have opposing dynamics during interval timing. A) D2-MSNs in the indirect pathway, which project from the striatum to the globus pallidus external segment (GPe; sagittal section) and internal segment (GPi) and B) D1-MSNs, which project from the striatum to the GPe, GPi, and substantia nigra (SNr; sagittal section). Peri-event raster C) from an optogenetically tagged putative D2-MSN (red) and D) from an optogenetically tagged putative D1-MSN (blue). Shaded area is the bootstrapped 95% confidence interval. E) Peri-event time histograms (PETHS) from all D2-MSNs and F) from all D1-MSNs. were binned at 0.2 seconds, smoothed using kernel-density estimates using a bandwidth of 1, and z-scored. Average activity from PETHs revealed that G) D2-MSNs (red) tended to ramp up, whereas H) D1-MSNs (blue) tended to ramp down. Shaded area is standard error. Data from 32 tagged D2-MSNs in 4 D2-Cre mice and 41 tagged D1-MSNs in 5 D1-Cre mice.

Differences between D2-MSNs and D1-MSNs

To quantify differences between D2-MSNs vs D1-MSNs, we turned to principal

component analysis (PCA), a data-driven tool to capture the diversity of neuronal activity (Kim

et al., 2017a). Work by our group and others has uniformly identified PC1 as a linear component

among corticostriatal neuronal ensembles during interval timing (Bruce et al., 2021; Emmons et

al., 2020, 2019, 2017; Kim et al., 2017a; Narayanan et al., 2013; Narayanan and Laubach, 2009;

Parker et al., 2014; Wang et al., 2018). We analyzed PCA calculated from all D2-MSN and D1-

MSN PETHs over the 6-second interval immediately after trial start. PCA identified time-

dependent ramping activity as PC1 (Fig 3A), a key temporal signal that explained 54% of

variance among tagged MSNs (Fig 3B; variance for PC1 $p = 0.009$ vs 46 (44-49)% variance for

PC1 derived from random data; Narayanan, 2016). In line with population averages from Fig

2G&H, D2-MSNs and D1-MSNs had opposite patterns of activity with negative PC1 scores for

D2-MSNs and positive PC1 scores for D1-MSNs (Fig 3C; PC1 for D2-MSNs: -3.4 (-4.6 – 2.5);

PC1 for D1-MSNs: 2.8 (-2.8 – 4.9); F = 8.8, $p = 0.004$ accounting for variance between mice

(Fig S3A); Cohen's $d = 0.7$; power = 0.80; no reliable effect of sex (F = 0.44, $p = 0.51$) or

switching direction (F = 1.73, $p = 0.19$)). Thus, analysis of PC1, which exhibited time-dependent

ramping similar to D2-MSN/D1-MSN average activity (Fig 2G/H and Fig S2C), S2E suggested

that D2-MSNs and D1-MSNs had opposing ramping dynamics.

To interrogate these dynamics at a trial-by-trial level, we calculated the linear slope of

D2-MSN and D1-MSN activity over the first 6 seconds of each trial using generalized linear

modeling (GLM) of effects of time in the interval vs trial-by-trial firing rate (Latimer et al.,

2015). Nosepokes were included as a regressor for movement. GLM analysis also demonstrated

that D2-MSNs had significantly different slopes (-0.01 spikes/second (-0.10 – 0.10)), which were

distinct from D1-MSNs (-0.20 (-0.47– -0.06; Fig 3D; F = 8.9, p = 0.004 accounting for variance

between mice (Fig S3B); Cohen's d = 0.8; power = 0.98; no reliable effect of sex (F = 0.02, p =

0.88) or switching direction (F = 1.72, p = 0.19)). We found that D2-MSNs and D1-MSNs had

significantly different slopes even when excluding outliers (4 outliers excluded outside of 95%

confidence intervals; F = 7.51, p = 0.008 accounting for variance between mice) and when the

interval was defined as the time between trial start and the switch response on a trial-by-trial

basis for each neuron (F = 4.3, p = 0.04 accounting for variance between mice). Trial-by-trial

GLM slope was correlated with PC1 scores in Fig 3A-C (PC1 scores vs. GLM slope r = -0.60, p

= 10^{-8}). These data demonstrate that D2-MSNs and D1-MSNs had distinct slopes of firing rate

across the interval and were consistent with analyses of average activity and PC1, which

exhibited time-related ramping.

Our findings could not be easily explained by movement because 1) few responses

occurred before 6 seconds (23%) and 44% of tagged neurons were recorded in trials without a

single nosepoke before 6 seconds (Fig 1B-C), 2) our GLM included a regressor accounting for

the nosepokes when present, and 3) nosepoke GLM βs were not reliably different between D2-

MSNs and D1-MSNs (F = 1.5, p = 0.22 accounting for variance between mice).

In summary, we used optogenetic tagging to record from D2-MSNs and D1-MSNs

during interval timing. Analyses of average activity, PC1, and trial-by-trial firing-rate slopes

over the interval provide convergent evidence that D2-MSNs and D1-MSNs had distinct and

opposing dynamics during interval timing. These data provide insight into temporal processing

by striatal MSNs.

Figure 3: Quantification of opposing D2-MSN and D1-MSN dynamics A) Principal component analysis revealed that the first component (PC1) exhibited time-dependent ramping. B) The first principal component explained ~54% of variance across tagged MSN ensembles. C) Differences between D2-MSNs (red) and D1-MSNs (blue) were captured by PC1 which exhibited time-dependent ramping. D) These differences were also apparent in the linear slope of firing rate vs time in the interval, with D1-MSNs (blue) having a more negative slope than D2-MSNs (red). In C and D, each point represents data from a tagged MSN. * = $p < 0.05$ via linear mixed effects models accounting for variance between mice. Data from 32 tagged D2-MSNs in 4 D2-Cre mice and 41 tagged D1-MSNs in 5 D1-Cre mice.

Drift-diffusion models of opposing D2-MSN and D1-MSN dynamics

Our data demonstrate that D2-MSNs and D1-MSNs have opposite activity patterns.

However, past computational models of interval timing have relied on drift-diffusion dynamics

with a positive slope that accumulates evidence over time (Nguyen et al., 2020; Simen et al.,

2011). To reconcile how these MSNs might complement to effect temporal control of action, we

constructed a four-parameter drift-diffusion model (DDM). Our goal was to construct a DDM

inspired by average differences in D2-MSNs and D1-MSNs that predicted switch-response time

behavior. We constructed a DDM where x represents the neuronal firing rate of an 'output' unit

collecting evidence on the activity of striatal D2-MSNs or D1-MSNs, t represents time measured

in seconds, and dx and dt mean "change" in x and t respectively (equivalent to the derivative

dx/dt):

237            $$dx = (F - x)Ddt + \sigma d\xi(t)$$ (Equation 1)

238            $$x(0) = b$$ (Equation 2)

239   The model has four independent parameters $F, D, \sigma,$ and b (described below) and a threshold

240   value T defined by

241            $$T = T(F, b) = \left(1 - \frac{1}{4}b\right)F + \frac{1}{4}b(1 - F)$$ (Equation 3)

The firing rate is set initially at baseline b (see Equation 2), then driven by input F akin to the

dendritic current induced by the overlap of corticostriatal postsynaptic potentials (Shepherd,

2013). With each unit of time dt, we suppose there is corticostriatal stimulation that provides

incremental input proportional to the activity itself, $(F - x)D$, to reach a decision. D is *the drift*

rate for the event sequence and can be interpreted as a parameter inverse proportional to the

neural activity's integration time constant. The drift, together with noise $\xi(t)$ (of zero mean and

strength σ), leads to fluctuating accumulation which eventually crosses a threshold T (see

Equation 3; Fig 4A-B). The time t^* it takes the firing rate to reach the threshold, $x(t^*) = T$, is

the *switch response time*.

Our model aimed to fit statistical properties of mouse behavioral responses while

incorporating MSN network dynamics. However, the model does not attempt to fit individual

neurons' activity, because our model predicts a single behavioral parameter – switch time – that

can be caused by the aggregation of diverse neuronal activity. We first analyzed trial-based

aggregated activity of MSN recordings from each mouse $(x_j(t))$ where $j = 1, \dots, N$ neurons. For

D2-MSN or D1-MSN ensembles of $N > 11$, we found linear combinations of their neuronal

activities, with some β_j coefficients,

$$258 \qquad x(t) = \sum_{j=1}^{N} \beta_j \, x_j(t)$$

that could predict the trial-by-trial switch response times (accuracy > 90%, Fig S4; compared

with < 20% accuracy for Poisson-generated spikes of same trial-average firing rate). The

predicted switch time t^*_{pred} was defined by the time when the weighted ensemble activity $x(t)$

first reached the value $x(t^*_{pred}) = 0.5$. Finally, we built DDMs to account for this opposing trend

(increasing vs decreasing) of MSN dynamics and for ensemble threshold behavior defining

t^*_{pred}; see the resulting model (Equations 1-3) and its simulations (Figure 4A-B).

This instantiation of DDMs captured the complimentary D2-MSN and D1-MSN

dynamics that we discovered in our optogenetic tagging experiments (Fig 2G-H vs. 4A-B). The

model's parameters were chosen to fit the distribution of switch response times: $F = 1, b = 0.52$

(so $T = 0.87$), $D = 0.135, \sigma = 0.052$ for intact D2-MSNs (Fig 4A, in black); and $F = 0, b =$

0.48 (so $T = 0.13$), $D = 0.141, \sigma = 0.052$ for intact D1-MSNs (Fig 4B, in black). Interestingly,

we observed that two-parameter gamma distributions almost perfectly accounted for the model

dynamics (D2-MSNs: Gamma parameters $\alpha = 6.08$, $\beta = 0.69$, R^2 vs Model = 0.99; D1-MSNs $\alpha =$

5.89, $\beta = 0.69$; R^2 vs Model = 0.99; Fig 4C-D, dotted black lines). Gamma distributions provided

a surprisingly good approximation for the probability distribution and cumulative distribution

functions of mouse switch response times (Fig 1B-D). Thus, in the next sections, we used

gamma distributions as a proxy for both the distribution of times generated by the model and the

distribution of mice switch response times when comparing those for the goodness of fit. Our

model provided the opportunity to computationally explore the consequences of disrupting D2-

MSNs or D1-MSNs. Because both D2-MSNs and D1-MSNs accumulate temporal evidence,

disrupting either MSN type in the model changed the slope. The results were obtained by

simultaneously decreasing the drift rate D (equivalent to lengthening the neurons' integration

time constant) and lowering the level of network noise σ: $D = 0.129, \sigma = 0.043$ for D2-MSNs

in Fig 4A (in red; changes in noise had to accompany changes in drift rate to preserve switch

response time variance); and $D = 0.122, \sigma = 0.043$ for D1-MSNs in Fig 4B (in blue). The

model predicted that disrupting either D2-MSNs or D1-MSNs would increase switch response

times (Fig 4C and Fig 4D) and would shift MSN dynamics. In the next section, we interrogated

these ideas with a combination of optogenetics, behavioral pharmacology, and

electrophysiology.

Figure 4: Four-parameter drift-diffusion computational model of striatal activity during interval timing.
A) We modeled interval timing with a low parameter diffusion process with a drift rate *D*, noise $\xi(t)$, and a baseline firing rate *b* that drifts toward a threshold *T* indicated by dotted lines. With D2-MSNs disrupted (solid red curves), this drift process decreases in slope and takes longer to reach the threshold. B) The same model also accounted for D1-MSNs with an opposite drift slope. With D1-MSNs disrupted (solid blue curves), the drift process again takes longer to reach the threshold. Because both D2-MSNs and D1-MSNs contribute to the accumulation of temporal evidence, this model predicted that C) disrupting D2-MSNs would increase response times during interval timing (dotted red line) and D) disrupting D1-MSNs would also increase response times (dotted blue line). Threshold *T* depends on *b* and target firing *F*.

Disrupting D2-MSNs or D1-MSNs increases switch response times

DDMs captured opposing MSN dynamics and predicted that disrupting either D2-MSNs or D1-MSNs should slow temporal processing and increase switch response times (Fig 4). We tested this idea with optogenetics. We bilaterally implanted fiber optics and virally expressed the inhibitory opsin halorhodopsin in the dorsomedial striatum of 10 D2-Cre mice (5 female) to inhibit D2-MSNs (Fig S5; this group of mice was entirely separate from the optogenetic tagging mice). We found that D2-MSN inhibition reliably increased switch response times (Fig 5A-B; Laser Off: 8.6 seconds (8.3–9.3); Laser On: 10.2 seconds (9.4–10.2); signed rank $p = 0.002$, Cohen's $d = 1.7$). Importantly, we found no reliable effects for D2-MSNs with opsin-negative controls (Fig S6). Remarkably, DDM predictions were highly concordant with D2-MSN inhibition behavioral data ($R^2 = 0.55$; parameters chosen to fit D2-MSN inhibition trials) as well as with behavioral data from laser-off trials ($R^2 = 0.63$; parameters chosen to fit laser-off trials; Fig S7).

Next, we investigated D1-MSNs (Kravitz et al., 2010). In 6 D1-Cre mice (3 female), optogenetic inhibition of dorsomedial striatal D1-MSNs increased switch response times (Fig 5C-D; Laser Off: 8.7 (8.0–9.1) seconds; Laser On: 10.5 (9.6 – 11.1) seconds; signed rank $p = 0.03$, Cohen's $d = 1.4$). As with D2-MSNs, we found no reliable effects with opsin-negative controls in D1-MSNs (Fig S6). DDM predictions again were highly concordant with D1-MSN inhibition behavioral data ($R^2 = 0.56$; parameters chosen to fit D1-MSN inhibition trials) and with laser off trials ($R^2 = 0.64$; parameters chosen to fit laser off trials; Fig S7).

To test the generality of the effects observed in the optogenetic inhibition experiment, we turned to systemic pharmacology. We used sulpiride (12.5 mg/kg intraperitoneally (IP)) to block D2-dopamine receptors in 10 wild-type (WT) mice, which increased switch response times

relative to saline sessions (median (IQR); saline: 8.9 (8.4 – 9.8) seconds; D2 blockade: 10.1 (9.4

– 10.8) seconds; signed rank $p = 0.002$, Cohen's $d = 1.0$; Fig 5E-F). Of note, there was no

difference in switch response time between saline sessions and laser-off trials from D2-MSN

optogenetic inhibition in D2-Cre mice, implying that in optogenetic sessions, D2-MSN inhibition

did not disrupt behavior during laser-off trials (rank sum $p = 0.31$). These data are consistent

with past work demonstrating that disrupting dorsomedial D2-MSNs slows timing (De Corte et

al., 2019b; Drew et al., 2007; Stutt et al., 2024). In the same 10 wild-type mice, systemic drugs

blocking D1-dopamine receptors (D1 blockade; SCH23390 0.05 mg/kg IP) increased switch

response times (saline: 9.7 seconds (9.0 – 9.9); D1 blockade: 10.1 seconds (9.3 – 11.4); signed

rank $p = 0.04$, Cohen $d = 0.7$; Fig 5G-H). Once again, there was no difference between saline

sessions and D1-MSN inhibition laser-off trials in D1-Cre mice (rank sum $p = 0.18$). These

results are in line with our DDM predictions and demonstrate that disrupting either D2-MSNs or

D1-MSNs increases switch response time.

We found no evidence that inhibiting D2-MSNs in the dorsomedial striatum changed

task-specific movements such as nosepoke duration (i.e., time of nosepoke entry to exit; signed

rank $p = 0.63$; Fig S8 for the 10 D2-Cre mice in optogenetics trials) or switch traversal time

between the first and second nosepokes (signed rank $p = 0.49$; Fig S8). Similarly, we found no

evidence that D1-MSN inhibition changed nosepoke duration (signed rank $p = 0.31$; Fig S8 for

the 6 D1-Cre mice) or traversal time (signed rank $p = 0.22$; Fig S8). Furthermore, disrupting D2-

MSNs or D1-MSNs did not change switch response time standard deviations or the number of

rewards (Fig S9). These results demonstrate that disrupting dorsomedial striatal D2-MSNs and

D1-MSNs specifically slowed interval timing without consistently changing task-specific

movements.

Figure 5: Disrupting D2- or D1-MSNs increases response times. A) As predicted by our DDM in Fig 4, optogenetic inhibition of D2-MSNs (red) shifted cumulative distributions of response times to the right, and B) increased response times; data from 10 D2-Cre mice expressing halorhodopsin (Halo). Also as predicted by our DDM, C) optogenetic inhibition of D1-MSNs shifted cumulative distribution functions to the right, and D) increased response times; data from 6 D1-Cre mice expressing Halo. Similarly, E) pharmacologically disrupting D2-dopamine receptors (red) with the D2 antagonist sulpiride shifted cumulative distribution functions to the right, and F) increased response times; data from 10 wild-type mice. Also, G) pharmacologically disrupting D1-dopamine receptors (blue) with the D1 antagonist SCH23390 shifted cumulative distribution functions to the right, and H) increased response times; data from the same 10 wild-type mice as in E-F. In B, D, F, and H connected points represent the mean response time from each animal in each session, and horizontal black lines represent group medians. *$p = < 0.05$, signed rank test. See Figure S6 for data from opsin-negative controls.

D2 blockade and D1 blockade shift MSN dynamics

MSN ensembles strongly encode time (Bruce et al., 2021; Emmons et al., 2017; Gouvea

et al., 2015; Mello et al., 2015; Wang et al., 2018), but it is unknown how disruptions in D2-

dopamine or D1-dopamine receptors affect these ensembles (Yun et al., 2023). Although non-

specific, pharmacological experiments have two advantages over optogenetics for recording

experiments: 1) many clinically approved drugs target dopamine receptors, and 2) they are more

readily combined with recordings than optogenetic inhibition, which silences large populations

of MSNs. We recorded from dorsomedial striatal MSN ensembles in 11 separate mice during

sessions with saline, D2 blockade with sulpiride, or D1 blockade with SCH23390 (Fig 6A; Fig

S10; data from one recording session each for saline, D2 blockade, and D1 blockade during

interval timing).

We analyzed 99 MSNs in sessions with saline, D2 blockade, and D1 blockade. We

matched MSNs across sessions based on waveform and interspike intervals; waveforms were

highly similar across sessions (correlation coefficient between matched MSN waveforms: saline

vs D2 blockade r = 1.00 (0.99 – 1.00 rank sum vs correlations in unmatched waveforms p =

$3x10^{-44;}$ waveforms; saline vs D1 blockade r = 1.00 (1.00 – 1.00), rank sum vs correlations in

unmatched waveforms $p = 4x10^{-50}$). There were no consistent changes in MSN average firing

rate with D2 blockade or D1 blockade (F = 1.1, p = 0.30 accounting for variance between

MSNs; saline: 5.2 (3.3 – 8.6) Hz; D2 blockade 5.1 (2.7 – 8.0) Hz; F = 2.2, p = 0.14; D1 blockade

4.9 (2.4 – 7.8) Hz).

We noticed differences in MSN activity across the interval with D2 blockade and D1

blockade at the individual MSN level (Fig 6B-D) as well as at the population level (Fig 6E). We

used PCA to quantify effects of D2 blockade or D1 blockade (Bruce et al., 2021; Emmons et al.,

2017; Kim et al., 2017a). We constructed principal components (PC) from z-scored peri-event

time histograms of firing rate from saline, D2 blockade, and D1 blockade sessions for all mice

together. The first component (PC1), which explained 54% of neuronal variance, exhibited

"time-dependent ramping", or monotonic changes over the 6 second interval immediately after

trial start (Fig 6F-G; variance for PC1 $p = 0.001$ vs 46 (45-47)% variance in random data;

Narayanan, 2016). Interestingly, PC1 scores shifted with D2 blockade (Fig 6F; PC1 scores for

D2 blockade: -0.6 (-3.8 – 4.7) vs saline: -2.3 (-4.2 – 3.2), F = 5.1, $p = 0.03$ accounting for

variance between MSNs; no reliable effect of sex (F = 0.2, $p = 0.63$) or switching direction (F =

2.8, $p = 0.10$)). PC1 scores also shifted with D1 blockade (Fig 6F; PC1 scores for D1 blockade: -

0.0 (-3.9 – 4.5), F = 5.8, $p = 0.02$ accounting for variance between MSNs; no reliable effect of

sex (F = 0.0, $p = 0.93$) or switching direction (F = 0.9, $p = 0.34$)). There were no reliable

differences in PC1 scores between D2 and D1 blockade. Furthermore, PC1 was distinct even

when sessions were sorted independently and assumed to be fully statistically independent

(Figure S10; D2 blockade vs saline: F = 5.8, $p = 0.02$; D1 blockade vs saline: F = 4.9, $p = 0.03$;

all analyses accounting for variance between mice). Higher components explained less variance

and were not reliably different between saline and D2 blockade or D1 blockade. Taken together,

this data-driven analysis shows that D2 and D1 blockade produced similar shifts in MSN

population dynamics represented by PC1. When combined with the major contributions of

D1/D2 MSNs to PC1 (Fig 3C) these findings indicate that pharmacological D2 blockade and D1

blockade disrupt ramping-related activity in the striatum.

Figure 6: D2 and D1 blockade shift temporal dynamics. A) We recorded dorsomedial striatal medium spiny neuron (MSN) ensembles during interval timing in sessions with saline, D2 blockade with sulpiride, or D1 blockade with SCH23390. Made with BioRender.com. B-D) Example peri-event raster from MSNs in sessions with saline (black), D2-dopamine blockade (red), or D1-dopamine blockade (blue). Shaded area is the bootstrapped 95% confidence interval. E) MSNs from 99 neurons in 11 mice from saline, D2 blockade, or D1 blockade session; MSNs were matched across sessions based on waveforms and interspike interval. Each row represents a peri-event time histogram (PETH) binned at 0.2 seconds, smoothed using kernel-density estimates using a bandwidth of 1, and z-scored. Colors indicate z-scored firing rate. See Fig S10 for analyses that assume statistical independence. F) Principal component analysis (PCA) identified MSN ensemble patterns of activity. The first principal component (PC1) exhibited time-dependent ramping. G) PC1 explained 54% of population variance among MSN ensembles; higher components were not analyzed. H) PC1 scores were closer to zero and significantly different with D2 or D1 blockade; * = $p < 0.05$ via linear mixed effects; data from 99 MSNs in 11 mice.

D2 blockade and D1 blockade degrades MSN temporal encoding

Finally, we quantified striatal MSN temporal encoding via a naïve Bayesian classifier

that generates trial-by-trial predictions of time from MSN ensemble firing rates (Fig 7A-C;

(Bruce et al., 2021; Emmons et al., 2017). Our DDMs predict that disrupted temporal decoding

would be a consequence of an altered DDM drift rate. We used leave-one-out cross-validation to

predict objective time from the firing rate within a trial. Saline sessions generated strong

temporal predictions for the first 6 seconds of the interval immediately after trial start (0–6

seconds; $R^2 = 0.91$ (0.83 – 0.94)) with weaker predictions for later epochs (6–12 seconds: $R^2 =$

0.55 (0.34 – 0.70); rank sum $p = 0.00002$ vs 0–6 seconds, Cohen $d = 2.0$; 12–18 seconds: $R^2 =$

0.18 (0.10 – 0.62); rank sum $p = 0.00003$ vs 0–6 seconds, Cohen $d = 2.4$; all analyses considered

each epoch statistically independent; Fig 7D). We found that temporal encoding early in the

interval (0–6 seconds) was degraded with either D2 blockade ($R^2 = 0.69$ (0.58 – 0.84); rank sum

$p = 0.0002$ vs saline, Cohen $d = 1.4$) or D1 blockade ($R^2 = 0.71$ (0.47 – 0.87); rank sum $p = 0.004$

vs saline, Cohen $d = 1.1$; Fig 7D), consistent with predictions made from DDMs. Later in the

interval (6–12 and 12-18 seconds), there were no significant differences between saline sessions

and D2 blockade or D1 blockade (Fig 7D).

Taken together, these data demonstrate that disrupting either D2-MSNs or D1-MSNs

degrades temporal encoding by MSN ensembles. In combination with our optogenetic tagging,

computational modeling, and optogenetic inhibition experiments, these data provide insight into

cognitive computations by the striatum.

410
411

Figure 7: D2 and D1 blockade degrade MSN temporal encoding. We used naïve Bayesian classifiers to decode time from MSN ensembles in A) saline sessions, B) D2 blockade sessions, and C) D1 blockade sessions. Color represents the temporal prediction across 20 trials with red representing stronger predictions. D) Temporal encoding was strong early in the interval, and D2 or D1 blockade degraded classification accuracy. Temporal encoding was decreased later in the interval. Each point represents the R^2 for each trial of behavior for MSN ensembles from 11 mice. * $p < 0.05$ vs saline from 0-6 seconds. Horizontal black lines in (D) represent group medians.

DISCUSSION

We describe how striatal MSNs work together in complementary ways to encode an elementary cognitive process, interval timing. Strikingly, optogenetic tagging showed that D2-MSNs and D1-MSNs had distinct dynamics during interval timing. MSN dynamics helped construct and constrain a four-parameter drift-diffusion model in which D2- and D1-MSN spiking accumulated temporal evidence. This model predicted that disrupting either D2-MSNs or D1-MSNs would increase switch response times. Accordingly, we found that optogenetically or pharmacologically disrupting striatal D2-MSNs or D1-MSNs increased switch response times without affecting task-specific movements. Disrupting D2-MSNs or D1-MSNs shifted MSN temporal dynamics and degraded MSN temporal encoding. These data, when combined with our model predictions, demonstrate that D2-MSNs and D1-MSNs contribute temporal evidence to controlling actions in time. Our results provide new insight into how opposing patterns of striatal MSN activity control behavior in similar ways and show that they play a complementary role in elementary cognitive operations.

Striatal MSNs are critical for temporal control of action (Emmons et al., 2017; Gouvea et al., 2015; Mello et al., 2015). Three broad models have been proposed for how striatal MSN ensembles represent time: 1) the striatal beat frequency model, in which MSNs encode temporal information based on neuronal synchrony (Matell and Meck, 2004); 2) the distributed coding model, in which time is represented by the state of the network (Paton and Buonomano, 2018); and 3) the DDM, in which neuronal activity monotonically drifts toward a threshold after which responses are initiated (Emmons et al., 2017; Simen et al., 2011; Wang et al., 2018). While our data do not formally resolve these possibilities, our results show that D2-MSNs and D1-MSNs exhibit opposing linear changes in firing rate dynamics in PC1 over the interval. Past work by

our group and others has demonstrated that these linear dynamics can scale over multiple

intervals to represent time (Emmons et al., 2020, 2017; Gouvea et al., 2015; Mello et al., 2015;

Wang et al., 2018). We find that low-parameter DDMs account for interval timing behavior with

both intact and disrupted striatal D2- and D1-MSNs. While other models can capture interval

timing behavior and account for MSN neuronal activity, our model does so parsimoniously with

relatively few parameters (Matell and Meck, 2004; Paton and Buonomano, 2018; Simen et al.,

2011). We and others have shown previously that ramping activity scales to multiple intervals,

and DDMs can be readily adapted by changing the drift rate (Emmons et al., 2017; Gouvea et al.,

2015; Mello et al., 2015; Simen et al., 2011). Interestingly, decoding performance was high early

in the interval; indeed, animals may have been focused on this initial interval (Balci and

Gallistel, 2006) in making temporal comparisons and deciding whether to switch response

nosepokes.

D2-MSNs and D1-MSNs play complementary roles in movement. For instance,

stimulating D1-MSNs facilitates movement, whereas stimulating D2-MSNs impairs movement

(Kravitz et al., 2010). Both populations have been shown to have complementary patterns of

activity during movements (Tecuapetla et al., 2016), with MSNs firing at different phases of

action initiation and selection. Further dissection of action selection programs reveals that

opposing patterns of activation among D2-MSNs and D1-MSNs suppress and guide actions,

respectively, in the dorsolateral striatum (Cruz et al., 2022). A particular advantage of interval

timing is that it captures a cognitive behavior within a single dimension — time. When projected

along the temporal dimension, it was surprising that D2-MSNs and D1-MSNs had opposing

patterns of activity. Past pharmacological work from our group and others has shown that

disrupting D2- or D1-MSNs slows timing (De Corte et al., 2019b; Drew et al., 2007, 2003; Stutt

et al., 2024), in line with pharmacological and optogenetic results in this manuscript.

Computational modeling predicted that disrupting either D2-MSNs or D1-MSNs increased self-

reported estimates of time, which was supported by both optogenetic and pharmacological

experiments. Notably, these disruptions are distinct from increased timing variability reported

with administrations of amphetamine, ventral tegmental area dopamine neuron lesions, and

rodent models of aging (Balci et al., 2008; Gür et al., 2020; Weber et al., 2023). Furthermore, our

current data demonstrate that disrupting either D2-MSN or D1-MSN activity shifted MSN

dynamics and degraded temporal encoding, supporting prior work (De Corte et al., 2019b; Drew

et al., 2007, 2003; Stutt et al., 2024). Our recording experiments do not identify where a possible

response threshold T is instantiated, but downstream basal ganglia structures may have a key role

in setting response thresholds (Toda et al., 2017).

Because interval timing is reliably disrupted in human diseases of the striatum such as

Huntington's disease, Parkinson's disease, and schizophrenia (Hinton et al., 2007; Singh et al.,

2021; Ward et al., 2011), these results have relevance to human disease. Our task version has

been used extensively to study interval timing in mice and humans (Balci et al., 2008; Bruce et

al., 2021; Stutt et al., 2024; Tosun et al., 2016; Weber et al., 2023). However, temporal bisection

tasks, in which animals hold during a temporal cue and respond at different locations depending

on cue length, have advantages in studying how animals time an interval because animals are not

moving while estimating cue duration (Paton and Buonomano, 2018; Robbe, 2023; Soares et al.,

2016). Our interval timing task version – in which mice switch between two response nosepokes

to indicate their interval estimate has elapsed – has been used extensively in rodent models of

neurodegenerative disease (Larson et al., 2022; Weber et al., 2024, 2023; Zhang et al., 2021), as

well as in humans (Stutt et al., 2024). Furthermore, because many therapeutics targeting

dopamine receptors are used clinically, these findings help describe how dopaminergic drugs

might affect cognitive function and dysfunction. Future studies of D2-MSNs and D1-MSNs in

temporal bisection and other timing tasks may further clarify the relative roles of D2- and D1-

MSNs in interval timing and time estimation.

Our approach has several limitations. First, systemic drug injections block D2- and D1-

receptors in many different brain regions, including the frontal cortex, which is involved in

interval timing (Kim et al., 2017a). D2 blockade or D1 blockade may have complex effects,

including corticostriatal or network effects that contribute to changes in D2-MSN or D1-MSN

ensemble activity. We note that optogenetic inhibition of D2-MSNs and D1-MSNs produces

similar effects to pharmacology in Figure 5. Pharmacology is compatible with neuronal

ensemble recordings as optogenetic inhibition would silence a large fraction of neurons,

complicating interpretations. Future studies might extend our work by combining local

pharmacology with neuronal ensemble recording. Second, although we had adequate statistical

power and medium-to-large effect sizes, optogenetic tagging is low-yield, and it is possible that

recording more of these neurons would afford greater opportunity to identify more robust results

and alternative coding schemes, such as neuronal synchrony. Furthermore, we find that recording

more neurons simultaneously would also help further constrain DDMs to predict trial-by-trial

firing rate and generate a more sophisticated neuronal network model of time (Fig S4). Third, the

striatum includes diverse cell types, some of which express both D1 and D2 dopamine receptors;

these cell types may also contribute to cognitive processing. Fourth, MSNs can laterally inhibit

each other, which may profoundly affect striatal function. Regardless, we show that cell-type-

specific disruption of D2-MSNs and D1-MSNs both slow timing, implying that these cell types

play a key role in interval timing. Future experiments may record from non-MSN striatal cell

types, including fast-spiking interneurons that shape basal ganglia output. Fifth, we did not

deliver stimulation to the striatum because our pilot experiments triggered movement artifacts or

task-specific dyskinesias (Kravitz et al., 2010). Future stimulation approaches carefully titrated

to striatal physiology may affect interval timing without affecting movement. Finally, movement

and motivation contribute to MSN dynamics (Robbe, 2023). Four lines of evidence argue that

our findings cannot be directly explained by motor confounds: 1) D2-MSNs and D1-MSNs

diverge between 0-6 seconds after trial start well before the first nosepoke (Fig S2), 2) our GLM

accounted for nosepokes and nosepoke-related βs were similar between D2-MSNs and D1-

MSNs, 3) optogenetic disruption of dorsomedial D2-MSNs and D1-MSNs did not change task-

specific movements despite reliable changes in switch response time, and 4) ramping dynamics

were quite distinct from movement dynamics. Furthermore, disrupting D2-MSNs and D1-MSNs

did not change the number of rewards animals received, implying that these disruptions did not

grossly affect motivation. Still, future work combining motion tracking with neuronal ensemble

recording and optogenetics and including bisection tasks may further unravel timing vs.

movement in MSN dynamics (Robbe, 2023).

In summary, we examined the role of dorsomedial striatal D2-MSNs and D1-MSNs

during an elementary cognitive behavior, interval timing. Optogenetic tagging revealed that D2-

MSNs and D1-MSNs exhibited opposite and complementary patterns of neuronal activity. These

dynamics could be captured by computational drift-diffusion models, which predicted that

disrupting either D2-MSNs or D1-MSNs would slow the accumulation of temporal evidence and

increase switch response time. In line with this prediction, we found that optogenetic or

pharmacological disruption of either D2-MSNs or D1-MSNs increased switch response times,

with pharmacological D2 or D1 blockade shifting MSN dynamics and degrading temporal

decoding. Collectively, our data provide insight into how the striatum encodes cognitive

information, which could be highly relevant for human diseases that disrupt the striatum and for

next-generation neuromodulation that targets the basal ganglia to avoid cognitive side effects or

to treat cognitive dysfunction.

Methods and Materials

Rodents. All procedures were approved by the Institutional Animal Care and Use Committee

(IACUC) at the University of Iowa, and all experimental methods were performed in accordance

with applicable guidelines and regulations (Protocol #0062039). We used five cohorts of mice,

summarized in Table 1: 1) 30 wild-type C57BL/6J mice (17 female) for behavioral experiments

(Fig 1), 2) 4 *Drd2-cre+* mice derived from Gensat strain ER44 (2 female) and 5 *Drd1-cre+* mice

derived from Gensat strain EY262 (2 female) for optogenetic tagging and neuronal ensemble

recordings in Fig 2-3; 3) 10 *Drd2-cre+* (5 female), and 6 *Drd1-cre+* mice (3 female) for

optogenetic inhibition (Fig 5) with 5 *Drd2-cre+* and 5 *Drd1-cre+* controls and 4) 10 wild-type

mice for behavioral pharmacology (Fig 5), and 5) 11 mice (4 C57BL/6J (2 female), 5 *Drd2-cre+*

mice (2 female), and 2 *Drd1-cre+* mice (0 female)) for combined behavioral pharmacology and

neuronal ensemble recording (Fig 6-7) (Table 1). Our recent work shows that D2-blockade and

D1-blockade have similar effects in both sexes(Stutt et al., 2024).

Interval timing switch task. We used a mouse-optimized operant interval timing task described

in detail previously (Balci et al., 2008; Bruce et al., 2021; Tosun et al., 2016; Weber et al., 2023).

Briefly, mice were trained in sound-attenuating operant chambers, with two front nosepokes

flanking either side of a food hopper on the front wall, and a third nosepoke located at the center

of the back wall. The chamber was positioned below an 8-kHz, 72-dB speaker (Fig 1A;

MedAssociates, St. Albans, VT). Mice were 85% food restricted and motivated with 20 mg

sucrose pellets (Bio-Serv, Flemington, NJ). Mice were initially trained to receive rewards during

fixed ratio nosepoke response trials. Nosepoke entry and exit were captured by infrared beams.

After shaping, mice were trained in the "switch" interval timing task. Mice self-initiated trials at

the back nosepoke, after which tone and nosepoke lights were illuminated simultaneously. Cues

were identical on all trial types and lasted the entire duration of the trial (6 or 18 seconds). On

50% of trials, mice were rewarded for a nosepoke after 6 seconds at the designated first 'front'

nosepoke; these trials were not analyzed. On the remaining 50% of trials, mice were rewarded

for nosepoking first at the 'first' nosepoke location and then switching to the 'second' nosepoke

location; the reward was delivered for initial nosepokes at the second nosepoke location after 18

seconds when preceded by a nosepoke at the first nosepoke location. Multiple nosepokes at each

nosepokes were allowed. Early responses at the first or second nosepoke were not reinforced.

Initial responses at the second nosepoke rather than the first nosepoke, alternating between

nosepokes, going back to the first nosepoke after the second nosepoke were rare after initial

training. Error trials included trials where animals responded only at the first or second nosepoke

and were also not reinforced. We did not analyze error trials as they were often too few to

analyze; these were analyzed at length in our prior work (Bruce et al., 2021).

Switch response time was defined as the moment animals departed the first nosepoke

before arriving at the second nosepoke. Critically, switch responses are a time-based decision

guided by temporal control of action because mice switch nosepokes only if nosepokes at the

first location did not receive a reward after 6 seconds. That is, mice estimate if more than 6

seconds have elapsed without receiving a reward to decide to switch responses. Mice learn this

task quickly (3-4 weeks), and error trials in which an animal nosepokes in the wrong order or

does not nosepoke are relatively rare and discarded. Consequently, we focused on these switch

response times as the key metric for temporal control of action. *Traversal time* was defined as

the duration between first nosepoke exit and second nosepoke entry and is distinct from switch

response time when animals departed the first nosepoke. *Nosepoke duration* was defined as the

time between first nosepoke entry and exit for the switch response times only. Trials were self-

initiated, but there was an intertrial interval with a geometric mean of 30 seconds between trials.

Surgical and histological procedures. Surgical procedures were identical to methods described

previously (Bruce et al., 2021). Briefly, mice were anesthetized using inhaled 4% isoflurane and

surgical levels of anesthesia were maintained at 1-2% for the duration of the surgery.

Craniotomies were drilled above bilateral dorsal striatal anatomical targets, and optogenetic

viruses (AAV5-DIO-eNHPR2.0 (halorhodopsin), AAV5-DIO-ChR2(H134R)-mcherry (ChR2),

or AAV5-DIO-cherry (control) from the University of North Carolina Viral Vector Core) were

injected into the dorsal striatum (0.5 uL of virus, +0.9, ML +/-1.3, DV –2.7). Either fiber optics

(Doric Lenses, Montreal Quebec; AP +0.9, ML +/-1.3, DV –2.5) or 4 x 4 electrode or optrode

arrays (AP +0.4, ML -1.4, DV -2.7 on the left side only; Microprobes, Gaithersburg, MD) were

positioned in the dorsal striatum. Holes were drilled to insert skull screws to anchor headcap

assemblies and/or ground electrode arrays and then sealed with cyanoacrylate ("SloZap," Pacer

Technologies, Rancho Cucamonga, CA), accelerated by "ZipKicker" (Pacer Technologies) and

methyl methacrylate (AM Systems, Port Angeles, WA). Following postoperative recovery, mice

were trained on the switch task and acclimated to experimental procedures prior to undergoing

experimental sessions.

Optogenetics. We leveraged cell-type-specific optogenetics to manipulate D2-MSNs and D1-

MSNs in D2-cre or D1-cre mice. In animals injected with optogenetic viruses, optical inhibition

was delivered via bilateral patch cables for the entire trial duration of 18 seconds via 589-nm

laser light at 12 mW power on 50% of randomly assigned trials. We performed control

experiments in mice without opsins using identical laser parameters in D2-cre or D1-cre mice

(Fig S6). We did not stimulate for epochs less than the interval because we did not want to

introduce a cue during the interval. For optogenetic tagging, putative D1- and D2-MSNs were

optically identified via 473-nm photostimulation. Units with mean post-stimulation spike

latencies of ≤5 milliseconds and a stimulated-to-unstimulated waveform correlation ratio of >0.9

were classified as putative D2-MSNs or D1-MSNs (Ryan et al., 2018; Shin et al., 2018). Only

one recording session was performed for each animal per day, and one recording session was

included from each animal.

Behavioral pharmacology procedures. C57BL/6J mice were injected intraperitoneally (IP) 20-

40 minutes before interval timing trials with either SCH23390 ($C_{17}H_{18}ClNO$; D1 antagonist),

sulpiride ($C_{15}H_{23}N_3O_4S$; D2 antagonist), or isotonic saline. The sulpiride dosage was 12.5 mg/kg,

0.01 mL/g, and the SCH23390 was administered at a dosage of 0.05 mg/kg, 0.01 mL/g.

Behavioral performance was compared with interval timing behavior on the prior day when

isotonic saline was injected IP. Only one recording session was performed for each animal per

day, and one recording session was included from saline, D2 blockade, and D1 blockade

sessions.

Electrophysiology. Single-unit recordings were made using a multi-electrode recording system

(Open Ephys, Atlanta, GA). After the experiments, Plexon Offline Sorter (Plexon, Dallas, TX),

was used to remove artifacts. Principal component analysis (PCA) and waveform shape were

used for spike sorting. Single units were defined as those 1) having a consistent waveform shape,

2) being a separable cluster in PCA space, and 3) having a consistent refractory period of at least

2 milliseconds in interspike interval histograms. The same MSNs were sorted across saline, D2

blockade, and D1 blockade sessions by loading all sessions simultaneously in Offline Sorter and

sorted using the preceding criteria. MSNs had to have consistent firing in all sessions to be

included. Sorting integrity across sessions was quantified by comparing waveform similarity via

correlation coefficients between sessions.

Spike activity was analyzed for all cells that fired between 0.5 Hz and 20 Hz over the

entire behavioral session. Putative MSNs were further separated from striatal fast-spiking

interneurons (FSIs) based on hierarchical clustering of the waveform peak-to-trough ratio and the

half-peak width (*fitgmdist* and *cluster.m*; Fig S1; (Berke, 2011)). We calculated kernel density

estimates of firing rates across the interval (-4 seconds before trial start to 22 seconds after trial

start) binned at 0.2 seconds, with a bandwidth of 1. We used PCA to identify data-driven patterns

of z-scored neuronal activity, as in our past work (Bruce et al., 2021; Emmons et al., 2017; Kim

et al., 2017a). The variance of PC1 was empirically compared against data generated from 1000

iterations of data from random timestamps with identical bins and kernel density estimates.

Average plots were shown with Gaussian smoothing for plotting purposes only.

Immunohistochemistry. Following completion of experiments, mice were transcardially

perfused with ice-cold 1x phosphate-buffered saline (PBS) and 4% paraformaldehyde (PFA)

after anesthesia using ketamine (100 mg/kg IP) and xylazine (10 mg/kg IP). Brains were then

fixed in solutions of 4% PFA and 30% sucrose before being cryosectioned on a freezing

microtome. Sections were stained for tyrosine hydroxylase with primary antibodies for >12

hours (rabbit anti-TH; Millipore MAB152; 1:1000) at 4° C. Sections were subsequently

visualized with Alexa Fluor fluorescent secondary antibodies (goat anti-rabbit IgG Alexa 519;

Thermo Fisher Scientific; 1:1,000) matched to host primary by incubating 2 hours at room

temperature. Histological reconstruction was completed using postmortem analysis of electrode

placement by slide-scanning microscopy on an Olympus VS120 microscope (Olympus, Center

Valley, PA; Fig S2&S4).

Trial-by-trial GLMs. To measure time-related ramping over the first 6 seconds of the interval,

we used trial-by-trial generalized linear models (GLMs) at the individual neuron level in which

the response variable was firing rate and the predictor variable was time in the interval or

nosepoke rate (Shimazaki and Shinomoto, 2007). For each neuron, its time-related "ramping"

slope was derived from the GLM fit of firing rate vs time in the interval, for all trials per neuron.

GLMs accounted for nosepoke movements unless there were no nosepokes within the 0-6 second

interval. All GLMs were run at a trial-by-trial level to avoid the effects of trial averaging

(Latimer et al., 2015) as in our past work (Bruce et al., 2021; Emmons et al., 2017; Kim et al.,

2017b). We performed additional sensitivity analyses excluding outliers outside of 95%

confidence intervals and measuring the firing rate from the start of the interval to the time of the

switch response on a trial-by-trial level for each neuron.

Machine-learning analyses. To predict time, we used a naïve Bayesian classifier to evaluate

neuronal ensemble decoding as in our past work (Emmons et al., 2020, 2017; Gouvea et al.,

2015; Kim et al., 2017a; Mello et al., 2015). Data from neurons with more than 20 trials across

all mice with the goal of evaluating how the decoding of time is affected by the epoch in the

interval and by D2/D1 blockade. To preclude edge effects that might bias classifier performance,

we included data from 6 seconds before trial start and 6 seconds after the end of the interval. We

used leave-one-out cross-validation to predict an objective time from the firing rate within a trial.

Classifier performance was quantified by computing the R^2 of objective time vs predicted time,

only for bins during the interval (0-6, 6-12, and 12-18 seconds; see Fig 7). Classifier

performance was compared using time-shuffled firing rates via a Wilcoxon signed-rank test.

Drift-diffusion models (DDM). We constructed a four-parameter DDM (see Equations 1-3 in

Results) that accounted for the ensemble threshold behavior observed in the neural data, then

used it to simulate switch response times compatible with mice behavioral data. The model was

implemented in MATLAB, starting from initial value b and with discrete computer simulation

steps $x_{new} = x_{old} + (F - x_{old})D\Delta t + \sigma\sqrt{\Delta t}N(0,1)$. Here $N(0,1)$ represents random values

generated from the standardized Gaussian distribution (mean = 0 and standard deviation = 1).

The integration timestep was $\Delta t = 0.1$. For each numerical simulation, the "switch response

time" was defined as the time t^* when variable x first reached the threshold value $T =$

$F(1 - b/4) + (1 - F)b/4$. For each condition, we ran 500 simulations of the model (up to 25

seconds per trial) and recorded the switch response times. Examples of firing rate x dynamics are

shown in Fig 4A-B. We observed that the distributions of interval timing switch response times

could be fit by a gamma probability distribution function $PDF(x) = \frac{\beta^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\beta x}$ with shape

α and rate β (see Fig S7 and Table S2).

Analysis and modeling of mouse MSN-ensemble recordings. Our preliminary analysis found

that, for sufficiently large number of neurons ($N > 11$), each recorded ensemble of MSNs on a

trial-by-trial basis could predict when mice would respond. We took the following approach:

First, for each MSN, we convolved its trial-by-trial spike train $Spk(t)$ with a 1-second

exponential kernel $K(t) = w\,e^{-t/w}$ if $t > 0$ and $K(t) = 0$ if $t \leq 0$ (Zhou et al., 2018; here $w =$

$1\,s$). Therefore, the smoothed, convolved spiking activity of neuron j ($j = 1,2,...N$),

687
$$x_j(t) = (Spk_j * K)(t) = \int_{-\infty}^{\infty} K(s)Spk_j(t-s)ds = \int_0^{\infty} K(s)Spk_j(t-s)ds$$

688 tracks and accumulates the most recent (one second, in average) firing-rate history of the j-th

689 MSN, up to moment t. We hypothesized that the ensemble activity $(x_1(t), x_2(t), \ldots, x_N(t))$,

690 weighted with some weights β_j, could predict the trial switch time t^* by considering the sum

691
$$x(t) = \sum_{j=1}^{N} \beta_j x_j(t)$$

692 and the sigmoid

693
$$y(t) = \frac{1}{1 + \exp\left(-(x(t) - 0.5)/k\right)} = \frac{1}{1 + \exp\left[-\left(\hat{\beta}_0 + \sum_{j=1}^{N} \hat{\beta}_j x_j(t)\right)\right]}$$

that approximates the firing rate of an output unit. Here parameter k indicates how fast

$x(t)$ crosses the threshold 0.5 coming from below (if $k > 0$) or coming from above (if $k < 0$)

and relates the weights β_j to the unknowns $\hat{\beta}_j = \beta_j/k$ and $\hat{\beta}_0 = -0.5/k$. Next, we ran a logistic

fit for every trial for a given mouse over the spike count predictor matrix

$(x_1(t), x_2(t), \ldots, x_N(t))$ from the mouse MSN recorded ensemble, and observed value t^*,

estimating the coefficients $\hat{\beta}_0$ and $\hat{\beta}_j$, and so, implicitly, the weights β_j. From there, we compute

the predicted switch time t^*_{pred} by condition $x(t) = 0.5$. Accuracy was quantified comparing the

predicted accuracy within a 1 second window to switch time on a trial-by-trial basis (Fig S4).

Statistics. All data and statistical approaches were reviewed by the Biostatistics, Epidemiology,

and Research Design Core (BERD) at the Institute for Clinical and Translational Sciences

(ICTS) at the University of Iowa. All code and data are made available at

http://narayanan.lab.uiowa.edu/article/datasets. We used the median to measure central tendency

and the interquartile range to measure spread. We used Wilcoxon nonparametric tests to compare

behavior between experimental conditions and Cohen's *d* to calculate effect size. Analyses of

putative single-unit activity and basic physiological properties were carried out using custom

routines for MATLAB.

For all neuronal analyses, variability between animals was accounted for using

generalized linear-mixed effects models and incorporating a random effect for each mouse into

the model, which allows us to account for inherent between-mouse variability. We used *fitglme*

in MATLAB and verified main effects using *lmer* in R. We accounted for variability between

MSNs in pharmacological datasets in which we could match MSNs between saline, D2

blockade, and D1 blockade. P values < 0.05 were interpreted as significant.

Data Availability
All raw data are available at http://narayanan.lab.uiowa.edu/article/datasets.

Code Availability

All code is available at http://narayanan.lab.uiowa.edu/article/datasets.

Acknowledgements

This work was funded by NIMH R01MH116043 to NSN. We thank Bernardo Sabatini for

detailed feedback on our manuscript.

Author Contributions

RAB, MAW, and NN designed the experiments. RAB, MAW, ASB, RV, CJ, and HRS

performed all experiments and collected data. The data was analyzed by RAB, MAW, ASB,

HRS, YK, RC, and NN. RC developed the computational model. RAB, MAW, ASB, KS, RC,

and NN wrote the manuscript, and all authors reviewed and revised the manuscript.

Table 1: Summary of mice, sessions, # of switch trials, and MSNs

Experiment	Figure	Cohort	Mice	# Sessions median (Q1-Q3)	# of Switch Responses	Neurons
Interval timing behavior	Fig 1B-C	1	30 wild-type mice	2 (2-2)	34 (25-42)	~
Optogenetic Tagging of D2-MSNs	Fig 2-3	2	4 D2-Cre mice	1 (1-1)	25 (22-25)	32 D2-MSNs
Optogenetic Tagging of D1-MSNs	Fig 2-3		5 D1-Cre mice	1(1-1)	22 (10-28)	41 D1-MSNs
Drift-diffusion models (DDM)	Fig 4A-D, Fig S4&S7	N/A	N/A	N/A	N/A	N/A
Optogenetic Inhibition of D2-MSNs	Fig 5A-B	3	10 D2-Cre mice	6(4-8)	127(78-135)	~
Optogenetic Inhibition of D1-MSNs	Fig 5C-D		6 D1-Cre mice	6(4-8)	81(60-106)	~
Optogenetic D2-MSN controls	Fig S7		5 D2-Cre mice	6(4-7)	104(60-140)	~
Optogenetic D1-MSN controls	Fig S7		5 D1-Cre mice	7(4-7)	102(78-128)	~
Pharmacological D2 blockade	Fig 5E-F	4	10 wild-type mice	2(2-2)	28(17-36)	~
Pharmacological D1 blockade	Fig 5G-H			2(2-2)	29(25-30)	~
MSN ensemble recording - saline	Fig 6-7, Fig S10	5	4 wild-type mice	1(1-1)	23(21-30)	158 MSNs
MSN ensembles – D2 blockade	Fig 6-7, Fig S10		5 D2-cre mice, and	1(1-1)	15(11-22)	167 MSNs
MSN ensembles – D1 blockade	Fig 6-7, Fig S10		2 D1-cre mice	1(1-1)	16(13-26)	144 MSNs

Supplementary Information

Condition	Source	α	β	
D2-MSNs Laser Off	Empirical Data	6.03	0.67	Fig S7A
D2-MSNs Laser Off	Model	6.08	0.69	Fig S7E
D2-MSNs Inhibited	Empirical Data	7.62	0.78	Fig S7B
D2-MSNs Inhibited	Model	7.67	0.79	Fig S7F
D1-MSNs Laser Off	Empirical Data	5.87	0.68	Fig S7C
D1-MSNs Laser Off	Model	5.89	0.69	Fig S7G
D1-MSNs Inhibited	Empirical Data	7.13	0.70	Fig S7D
D1-MSNs Inhibited	Model	7.31	0.72	Fig S7H

Table S2: Gamma distribution parameters

748
749

751
752

Figure S1: A) Recording locations in the dorsomedial striatum (targeting AP +0.4, ML -1.4, DV

-2.7). Electrode reconstructions for D2-Cre (red), D1-Cre (blue), and wild-type mice (green).

Only the left striatum was implanted with electrodes in all animals. B) MSN classification by

waveform criteria for sessions with optogenetic tagging. C) Example of an optogenetically

tagged MSN. This neuron expresses ChR2 and fired action potentials within 5 milliseconds of

473 nm laser pulses (red line). Spikes from laser trials shown as red ticks; trials without laser

shown as blue ticks. Inset on bottom right – waveforms from laser trials (red) and trials without

laser (blue). Across 73 tagged neurons, waveform correlation coefficients for laser trials vs. trials

without laser was r = 0.97 (0.92-0.99), indicating that optogenetically triggered spikes were

similar to non-optogenetically triggered spikes. D) MSN classification by waveform criteria for

pharmacology sessions.

Figure S2: D2- and D1-MSN activity over a longer epoch from 10 seconds prior to trial start,

when mice initiated trials at the back nosepoke, to the end of 18 seconds, after which making a

second nosepoke led to reward. A) Tagged D2-MSN from Figure 2C shown over a longer

interval, and B) tagged D1-MSN from Figure 2D shown over a longer interval. C) Peri-event

time histograms from D2-MSNs and D) from D1-MSNs over a longer interval. E) We noticed

that on average, D2-MSNs and D1-MSNs had the biggest differences in dynamics, where they

tended to have distinct slopes (Fig 3C&D); slope analyses were less reliable for other epochs.

Data from 32 tagged D2-MSNs in 4 D2-Cre mice and 41 tagged D1-MSNs in 5 D1-Cre mice as

in Fig 2.

Figure S3: Effects in individual mice from optogenetic tagging experiments for A) PC1 and B)

trial-by-trial GLM slope of firing rate over the interval; red = D2-MSN, and blue = D1-MSN. C)

Effects in individual mice for PC1 for all neurons from combined pharmacology and neuronal

ensemble recording; black is saline, red is D2 Blockade, and blue is D1 Blockade. Black lines

indicate the median for each condition in each mouse. All statistics analyzing these data used

linear-mixed effects models as incorporating a random effect for each mouse into the model

allows to account for inherent between-mouse variability.

Figure S4: Trial-by-trial predictions of switch response time from D2-MSN and D1-MSN ensemble

dynamics. A) Exemplar integral of network activity $x(t) = \sum \beta_j x_j(t)$ generated from an ensemble of 13

D2-MSNs for a single animal on a single trial. The coefficients β_j were computed from the logistic

regression fit to the switch time t^* using the neurons firing rates $(x_j(t))_j$ as the predictor matrix (see

Methods). On this trial, this integral drifted towards the response threshold 0.5, and logistic regression

accurately predicted the switch response time. B) Exemplar integral network activity $x(t)$ generated from

an ensemble of 15 D1-MSNs in a single animal on a single trial. C) Across all D2-MSN and D1-MSN

ensembles per individual mouse, we found that the trial-by-trial accuracy increased with ensemble size,

exceeding >90% when D2-MSN and D1-MSN ensembles were >11 neurons. Neuronal data from 5 D1-

Cre (blue dots) and 4 D2-Cre (red dots) mice in Figures 2-3. Corresponding light blue/light red dots show

accuracy values computed in 100 simulations of respective D2-MSN/D1-MSN ensembles with Poisson

spikes matched to MSN firing rates.

800
Figure S5: Fiber optic locations from A) an opsin-expressing mouse with mCherry-tagged

halorhodopsin and bilateral fiber optics, and B) across 10 D2-Cre mice (red) and 6 D1-cre mice

(blue) with fiber optics (targeting AP +0.9, ML +/-1.3, DV –2.5).

Figure S6: Experiments in D2-Cre mice injected with virus without opsins did not reliably affect

A) cumulative density functions (CDFs) or B) switch response times (signed rank $p = 0.44$).

Experiments in D1-Cre mice expressing virus without opsins did not reliably affect C) CDFs or

D) switch response times (signed rank $p = 0.81$). Laser parameters (589 nm laser, 12 mW, 18

second duration) were identical to experimental animals in Fig 5.

Figure S7: Model details. Histograms of behavioral data from D2-Cre mice with A) Laser Off

and B) D2-MSN inhibition (red). Data from D1-Cre mice with C) Laser Off and with D) D1-

MSN inhibition (blue). E-H) model predictions. I-L) Comparisons of empirical data vs model.

All panels: fits for the gamma distribution with dotted circles; see Table S2 for the parameter

values defining each gamma distribution. Behavioral data: from 10 D2-Cre mice and 6 D1-mice

from Fig 5A-D. Model data: from numerical simulations of the DDM model shown in Fig 4. All

DDM parameters were chosen to fit behavioral data in each condition.

Figure S8: Optogenetically inhibiting D2-MSNs or D1-MSNs does not affect task-specific

motor control. We measured nosepoke duration (time of nosepoke entry to exit) on switch

responses. During interval timing there was no effect of optogenetic inhibition (red) of

dorsomedial striatal D2-MSNs on A-B) nosepoke duration or C) the traversal time between the

first and second nosepokes; note that traversal time is distinct from the switch response, which is

the moment animals depart the first nosepoke prior to arriving at the second nosepoke.

There was also no effect of optogenetic inhibition (blue) of dorsomedial striatal D1-MSNs on

nosepoke duration (D-E) or F) switch traversal time. Data from the same 10 D2-Cre mice and 6

D1-Cre mice, as in Fig 5. Horizontal black lines in B, C, E, and F represent group medians.

839
840

Figure S9: A) Standard deviation of switch response times for D2-MSN inhibition sessions

(signed rank test, $p = 0.19$) and B) D1-MSN inhibition sessions ($p = 0.84$), and the number of

total rewards for C) D2-MSN inhibition sessions ($p = 0.07$) and d) D1-MSN inhibition sessions

($p = 0.25$). Data from 10 D2-Cre mice and 6 D1-Cre mice as in Fig 5.

Figure S10: We analyzed MSN ensembles in sessions with saline (158 neurons), D2 blockade

(167 neurons), or D1 blockade (144 neurons) – unlike Figure 6; all sessions were sorted

independently and assumed to be fully statistically independent. A) Principal component analysis

(PCA) identified MSN ensemble patterns of activity. The first principal component (PC1)

exhibited time-dependent ramping. B) PC1 explained 55% of population variance among MSN

ensembles. C) PC1 scores were shifted and significantly different with D2 or D1 blockade as

when all sessions were sorted together in Fig 6. * = $p < 0.05$ via linear mixed effects models

accounting for variance between mice; all analyses assumed statistical independence.

REFERENCES

Albin RL, Young AB, Penney JB. 1989. The functional anatomy of basal ganglia disorders. *Trends Neurosci* **12**:366–375. doi:10.1016/0166-2236(89)90074-X

Alexander GE, Crutcher MD. 1990. Functional architecture of basal ganglia circuits: neural substrates of parallel processing. *Trends Neurosci* **13**:266–271. doi:10.1016/0166-2236(90)90107-L

Andreasen NC. 1999. Understanding the causes of schizophrenia. *N Engl J Med* **340**:645–647. doi:10.1056/NEJM199902253400811

Averbeck BB, Lehman J, Jacobson M, Haber SN. 2014. Estimates of projection overlap and zones of convergence within frontal-striatal circuits. *J Neurosci* **34**:9497–9505.

Balci F, Gallistel CR. 2006. Cross-domain transfer of quantitative discriminations: Is it all a matter of proportion? *Psychon Bull Rev* **13**:636–642. doi:10.3758/BF03193974

Balci F, Papachristos EB, Gallistel CR, Brunner D, Gibson J, Shumyatsky GP. 2008. Interval timing in genetically modified mice: a simple paradigm. *Genes Brain Behav* **7**:373–384. doi:10.1111/j.1601-183X.2007.00348.x

Berke JD. 2011. Functional properties of striatal fast-spiking interneurons. *Front Syst Neurosci* **5**:45.

Bruce RA, Weber MA, Volkman RA, Oya M, Emmons EB, Kim Y, Narayanan NS. 2021. Experience-related enhancements in striatal temporal encoding. *Eur J Neurosci* **54**:5063–5074. doi:10.1111/ejn.15344

Buhusi CV, Meck WH. 2005. What makes us tick? Functional and neural mechanisms of interval timing. *Nat Rev Neurosci* **6**:755–765. doi:10.1038/nrn1764

Cruz BF, Guiomar G, Soares S, Motiwala A, Machens CK, Paton JJ. 2022. Action suppression reveals opponent parallel control via striatal circuits. *Nature* **607**:521–526. doi:10.1038/s41586-022-04894-9

Cui G, Jun SB, Jin X, Pham MD, Vogel SS, Lovinger DM, Costa RM. 2013. Concurrent activation of striatal direct and indirect pathways during action initiation. *Nature* **494**:238–242. doi:10.1038/nature11846

De Corte BJ, Wagner LM, Matell MS, Narayanan NS. 2019a. Striatal dopamine and the temporal control of behavior. *Behav Brain Res* **356**:375–379. doi:10.1016/j.bbr.2018.08.030

De Corte BJ, Wagner LM, Matell MS, Narayanan NS. 2019b. Striatal dopamine and the temporal control of behavior. *Behav Brain Res* **356**:375–379. doi:10.1016/j.bbr.2018.08.030

Drew MR, Fairhurst S, Malapani C, Horvitz JC, Balsam PD. 2003. Effects of dopamine antagonists on the timing of two intervals. *Pharmacol Biochem Behav* **75**:9–15.

Drew MR, Simpson EH, Kellendonk C, Herzberg WG, Lipatova O, Fairhurst S, Kandel ER, Malapani C, Balsam PD. 2007. Transient Overexpression of Striatal D2 Receptors Impairs Operant Motivation and Interval Timing. *J Neurosci* **27**:7731–7739. doi:10.1523/JNEUROSCI.1736-07.2007

Emmons EB, De Corte BJ, Kim Y, Parker KL, Matell MS, Narayanan NS. 2017. Rodent medial frontal control of temporal processing in the dorsomedial striatum. *J Neurosci* **37**:8718–8733.

Emmons EB, Kennedy M, Kim Y, Narayanan NS. 2019. Corticostriatal stimulation compensates for medial frontal inactivation during interval timing. *Sci Rep* **9**:1–9. doi:10.1038/s41598-019-50975-7

Emmons EB, Tunes-Chiuffa G, Choi J, Bruce RA, Weber MA, Kim Y, Narayanan NS. 2020. Temporal Learning Among Prefrontal and Striatal Ensembles. *Cereb Cortex Commun* **1**. doi:10.1093/texcom/tgaa058

Gouvea TS, Monteiro T, Motiwala A, Soares S, Machens C, Paton JJ. 2015. Striatal dynamics explain duration judgments. *eLife* **4**. doi:10.7554/eLife.11386

Graybiel AM. 1997. The Basal Ganglia and Cognitive Pattern Generators. *Schizophr Bull* **23**:459–469. doi:10.1093/schbul/23.3.459

Gür E, Duyan YA, Arkan S, Karson A, Balcı F. 2020. Interval timing deficits and their neurobiological correlates in aging mice. *Neurobiol Aging* **90**:33–42. doi:10.1016/j.neurobiolaging.2020.02.021

Hinton SC, Paulsen JS, Hoffmann RG, Reynolds NC, Zimbelman JL, Rao SM. 2007. Motor timing variability increases in preclinical Huntington's disease patients as estimated onset of motor symptoms approaches. *J Int Neuropsychol Soc* **13**:539–543. doi:10.1017/S1355617707070671

Kim Y-C, Han S-W, Alberico SL, Ruggiero RN, De Corte B, Chen K-H, Narayanan NS. 2017a. Optogenetic Stimulation of Frontal D1 Neurons Compensates for Impaired Temporal Control of Action in Dopamine-Depleted Mice. *Curr Biol CB* **27**:39–47. doi:10.1016/j.cub.2016.11.029

Kim Y-C, Han S-W, Alberico SL, Ruggiero RN, De Corte B, Chen K-H, Narayanan NS. 2017b. Optogenetic stimulation of frontal D1 neurons compensates for impaired temporal control of action in dopamine-depleted mice. *Curr Biol* **27**:39–47.

Kravitz AV, Freeze BS, Parker PR, Kay K, Thwin MT, Deisseroth K, Kreitzer AC. 2010. Regulation of parkinsonian motor behaviours by optogenetic control of basal ganglia circuitry. *Nature* **466**:622–626.

Larson T, Khandelwal V, Weber MA, Leidinger MR, Meyerholz DK, Narayanan NS, Zhang Q. 2022. Mice expressing P301S mutant human tau have deficits in interval timing. *Behav Brain Res* **432**:113967. doi:10.1016/j.bbr.2022.113967

Latimer KW, Yates JL, Meister ML, Huk AC, Pillow JW. 2015. Single-trial spike trains in parietal cortex reveal discrete steps during decision-making. *Science* **349**:184–187.

Matell MS, Meck WH. 2004. Cortico-striatal circuits and interval timing: coincidence detection of oscillatory processes. *Cogn Brain Res*, Neuroimaging of Interval Timing **21**:139–170. doi:10.1016/j.cogbrainres.2004.06.012

Matell MS, Meck WH, Nicolelis MAL. 2003. Interval timing and the encoding of signal duration by ensembles of cortical and striatal neurons. *Behav Neurosci* **117**:760–773. doi:10.1037/0735-7044.117.4.760

Meck WH. 2006. Neuroanatomical localization of an internal clock: A functional link between mesolimbic, nigrostriatal, and mesocortical dopaminergic systems. *Brain Res* **1109**:93–107. doi:10.1016/j.brainres.2006.06.031

Mello GBM, Soares S, Paton JJ. 2015. A Scalable Population Code for Time in the Striatum. *Curr Biol* **25**:1113–1122. doi:10.1016/j.cub.2015.02.036

Merchant H, de Lafuente V. 2014. Introduction to the neurobiology of interval timing. *Adv Exp Med Biol* **829**:1–13. doi:10.1007/978-1-4939-1782-2_1

Middleton FA, Strick PL. 2000. Basal ganglia and cerebellar loops: motor and cognitive circuits. *Brain Res Rev* **31**:236–250. doi:10.1016/S0165-0173(99)00040-5

Monteiro T, Rodrigues FS, Pexirra M, Cruz BF, Gonçalves AI, Rueda-Orozco PE, Paton JJ. 2023. Using temperature to analyze the neural basis of a time-based decision. *Nat Neurosci*. doi:10.1038/s41593-023-01378-5

Narayanan NS. 2016. Ramping activity is a cortical mechanism of temporal control of action. *Curr Opin Behav Sci* **8**:226–230. doi:10.1016/j.cobeha.2016.02.017

Narayanan NS, Albin RL. 2022. Cognition in Parkinson's Disease, Progress in Brain Research. Elsevier, Acad. Press.

Narayanan NS, Cavanagh JF, Frank MJ, Laubach M. 2013. Common medial frontal mechanisms of adaptive control in humans and rodents. *Nat Neurosci* **16**:1888–1897. doi:10.1038/nn.3549

Narayanan NS, Laubach M. 2009. Delay activity in rodent frontal cortex during a simple reaction time task. *J Neurophysiol* **101**:2859–2871. doi:10.1152/jn.90615.2008

Nguyen Q-A, Rinzel J, Curtu R. 2020. Buildup and bistability in auditory streaming as an evidence accumulation process with saturation. *PLOS Comput Biol* **16**:e1008152. doi:10.1371/journal.pcbi.1008152

Parker KL, Chen K-H, Kingyon JR, Cavanagh JF, Narayanan NS. 2015. Medial frontal ∼4-Hz activity in humans and rodents is attenuated in PD patients and in rodents with cortical dopamine depletion. *J Neurophysiol* **114**:1310–1320. doi:10.1152/jn.00412.2015

Parker KL, Chen K-H, Kingyon JR, Cavanagh JF, Narayanan NS. 2014. D1-Dependent 4 Hz Oscillations and Ramping Activity in Rodent Medial Frontal Cortex during Interval Timing. *J Neurosci* **34**:16774–16783. doi:10.1523/JNEUROSCI.2772-14.2014

Parker KL, Kim YC, Kelley RM, Nessler AJ, Chen K-H, Muller-Ewald VA, Andreasen NC, Narayanan NS. 2017. Delta-frequency stimulation of cerebellar projections can compensate for schizophrenia-related medial frontal dysfunction. *Mol Psychiatry* **22**:647–655. doi:10.1038/mp.2017.50

Parker KL, Lamichhane D, Caetano MS, Narayanan NS. 2013. Executive dysfunction in Parkinson's disease and timing deficits. *Front Integr Neurosci* **7**:75. doi:10.3389/fnint.2013.00075

Paton JJ, Buonomano DV. 2018. The Neural Basis of Timing: Distributed Mechanisms for Diverse Functions. *Neuron* **98**:687–705. doi:10.1016/j.neuron.2018.03.045

Robbe D. 2023. Lost in time: Relocating the perception of duration outside the brain. *Neurosci Biobehav Rev* **153**:105312. doi:10.1016/j.neubiorev.2023.105312

Ryan MB, Bair-Marshall C, Nelson AB. 2018. Aberrant striatal activity in parkinsonism and levodopa-induced dyskinesia. *Cell Rep* **23**:3438–3446.

Shepherd GMG. 2013. Corticostriatal connectivity and its role in disease. *Nat Rev Neurosci* **14**:278–291. doi:10.1038/nrn3469

Shimazaki H, Shinomoto S. 2007. A Method for Selecting the Bin Size of a Time Histogram. *Neural Comput* **19**:1503–1527. doi:10.1162/neco.2007.19.6.1503

Shin JH, Kim D, Jung MW. 2018. Differential coding of reward and movement information in the dorsomedial striatal direct and indirect pathways. *Nat Commun* **9**:1–14.

Simen P, Balci F, de Souza L, Cohen JD, Holmes P. 2011. A model of interval timing by neural integration. *J Neurosci Off J Soc Neurosci* **31**:9238–9253. doi:10.1523/JNEUROSCI.3121-10.2011

Singh A, Cole RC, Espinoza AI, Evans A, Cao S, Cavanagh JF, Narayanan NS. 2021. Timing variability and midfrontal ∼4 Hz rhythms correlate with cognition in Parkinson's disease. *NPJ Park Dis* **7**:14. doi:10.1038/s41531-021-00158-x

Soares S, Atallah BV, Paton JJ. 2016. Midbrain dopamine neurons control judgment of time.
Science **354**:1273–1277. doi:10.1126/science.aah5234
Stutt HR, Weber MA, Cole RC, Bova AS, Ding X, McMurrin MS, Narayanan NS. 2024. Sex
similarities and dopaminergic differences in interval timing. *Behav Neurosci* **138**:85–93.
doi:10.1037/bne0000577
Tecuapetla F, Jin X, Lima SQ, Costa RM. 2016. Complementary Contributions of Striatal
Projection Pathways to Action Initiation and Execution. *Cell* **166**:703–715.
doi:10.1016/j.cell.2016.06.032
Toda K, Lusk NA, Watson GDR, Kim N, Lu D, Li HE, Meck WH, Yin HH. 2017. Nigrotectal
Stimulation Stops Interval Timing in Mice. *Curr Biol* **27**:3763-3770.e3.
doi:10.1016/j.cub.2017.11.003
Tosun T, Gur E, Balci F. 2016. Mice plan decision strategies based on previously learned time
intervals, locations, and probabilities. *Proc Natl Acad Sci U S A* **113**:787–792.
doi:10.1073/pnas.1518316113
Wang J, Narain D, Hosseini EA, Jazayeri M. 2018. Flexible timing by temporal scaling of
cortical responses. *Nat Neurosci* **21**:102. doi:10.1038/s41593-017-0028-6
Ward RD, Kellendonk C, Kandel ER, Balsam PD. 2011. Timing as a window on cognition in
schizophrenia. *Neuropharmacology*. doi:10.1016/j.neuropharm.2011.04.014
Weber MA, Kerr G, Thangavel R, Conlon MM, Gumusoglu SB, Gupta K, Abdelmotilib HA,
Halhouli O, Zhang Q, Geerling JC, Narayanan NS, Aldridge GM. 2024. Alpha-Synuclein
Pre-Formed Fibrils Injected into Prefrontal Cortex Primarily Spread to Cortical and
Subcortical Structures. *J Park Dis* **14**:81–94. doi:10.3233/JPD-230129
Weber MA, Sivakumar K, Tabakovic EE, Oya M, Aldridge GM, Zhang Q, Simmering JE,
Narayanan NS. 2023. Glycolysis-enhancing α1-adrenergic antagonists modify cognitive
symptoms related to Parkinson's disease. *Npj Park Dis* **9**:1–7. doi:10.1038/s41531-023-
00477-1
Yun S, Yang B, Anair JD, Martin MM, Fleps SW, Pamukcu A, Yeh N-H, Contractor A,
Kennedy A, Parker JG. 2023. Antipsychotic drug efficacy correlates with the modulation
of D1 rather than D2 receptor-expressing striatal projection neurons. *Nat Neurosci* 1–12.
doi:10.1038/s41593-023-01390-9
Zhang Q, Abdelmotilib H, Larson T, Keomanivong C, Conlon M, Aldridge GM, Narayanan NS.
2021. Cortical alpha-synuclein preformed fibrils do not affect interval timing in mice.
Neurosci Lett **765**:136273. doi:10.1016/j.neulet.2021.136273
1029