Complementary cognitive roles for D2-MSNs and D1-MSNs in interval timing

R. Austin Bruce[1], Matthew A. Weber[1], Alexandra S. Bova[1], Rachael A. Volkman[1], Casey E. Jacobs, Kartik Sivakumar[1], Young-cho Kim[1], Rodica Curtu[2,3], Nandakumar S. Narayanan[1,3]

[1]Department of Neurology, University of Iowa, [2]Department of Mathematics, University of Iowa, and [3]The Iowa Neuroscience Institute

*Corresponding Author
Nandakumar Narayanan
nandakumar-narayanan@uiowa.edu
169 Newton Road
Pappajohn Biomedical Discovery Building—5336
University of Iowa, Iowa City, 52242
319-353-5698

Abstract

Striatal pathways control motivation and movement; however, their role in cognition is unclear. We studied dorsomedial striatal cognitive processing during interval timing, an elementary cognitive task that requires mice to estimate intervals of several seconds. Interval timing involves working memory for temporal rules and attention to the passage of time. We discovered that both pharmacological and optogenetic disruptions of D2-dopamine receptor-expressing medium spiny neurons (D2-MSNs) or D1-dopamine receptor-expressing MSNs (D1-MSNs) delayed timing. Disrupting D2-MSNs or D1-MSNs did not affect task-specific movements. Furthermore, pharmacological blockade of D2-dopamine receptors or D1-dopamine receptors degraded dorsomedial MSN temporal encoding. We compared the timing-related firing of optogenetically-tagged D2-MSNs and D1-MSNs. Strikingly, we found that D2-MSNs and D1-MSNs exhibited opposite dynamics over temporal intervals despite similar effects of D2-MSN and D1-MSN disruptions. MSN dynamics helped construct and constrain a four-parameter drift-diffusion computational model that captured interval timing behavior and D2-MSN and D1-MSN perturbations. Despite opposing dynamics, disrupting either D2-MSNs or D1-MSNs slowed timing, implying that D2-MSNs and D1-MSNs make complementary contributions to interval timing. These data provide novel insight into basal ganglia cognitive operations beyond movement. Our findings carry implications for a broad range of human striatal diseases and for therapies targeting striatal pathways.

Introduction

The basal ganglia have been heavily studied in motor control[1–4]. These subcortical nuclei are composed of two pathways defined by medium spiny neurons (MSNs) in the striatum: the indirect pathway, defined by D2-dopamine receptor-expressing MSNs (D2-MSNs), and the direct pathway, defined by D1-dopamine receptor-expressing MSNs (D1-MSNs). It has been proposed that the indirect and direct pathways play opposing roles in movement[1,2,5–7], although recent work identified how these pathways can play more complex and complementary roles[3,4,8,9]. MSNs receive dense input from both motor and cognitive cortical areas[10–14], but the respective roles of D2-MSNs and D1-MSNs in cognitive processing are largely unknown. Understanding the cognitive basal ganglia is critical for diseases affecting the striatum such as Huntington's disease, Parkinson's disease, and schizophrenia[15–17]. Furthermore, pharmacological and brain stimulation therapies directly modulate D2-MSNs, D1-MSNs, or downstream basal ganglia structures such as the globus pallidus or subthalamic nucleus. Determining striatal pathway contributions to cognition may help avoid side effects of current treatments and inspire novel therapeutic directions.

We studied striatal D2-MSNs and D1-MSNs during an elementary cognitive task, interval timing. This task requires estimating an interval of several seconds and provides an ideal platform to study cognition in the striatum because 1) interval timing requires cognitive resources including working memory for temporal rules and attention to the passage of time[18,19]; 2) it is reliably impaired in human striatal diseases such as Huntington's disease, Parkinson's disease, and schizophrenia[16,19–22]; 3) interval timing requires nigrostriatal dopamine that modulates MSNs[23–32]; and 4) can be rigorously studied in animal models[33–35]. Past work has

shown that disrupting D2-dopamine receptors (D2s) in the dorsomedial striatum powerfully

impairs interval timing[25,36–38]. D1-dopamine receptors (D1s) can also contribute to interval

timing[37–39]. These observations suggest that D2-MSNs and D1-MSNs control the timing of

action; however, the respective roles remain poorly delineated.

We investigated this problem with a combination of behavioral pharmacology, cell-type-

specific optogenetics, neuronal ensemble recordings, and computational modeling in the

dorsomedial striatum of mice. Remarkably, we found that pharmacologic and optogenetic

disruption of either indirect pathway D2-MSNs or direct pathway D1-MSNs slowed timing

behavior without affecting task-specific movements. Furthermore, pharmacological blockade of

either D2- or D1-receptors degraded MSN ensemble temporal encoding, suggesting that both

D2- and D1-MSNs are intimately involved in the temporal control of action. We optogenetically

tagged D2-MSNs and D1-MSNs to compare their patterns of activity. Strikingly, despite similar

effects of D2-MSN and D1-MSN disruptions, we found opposing neuronal dynamics with D2-

MSNs increasing firing over an interval and D1-MSNs decreasing firing over an interval. These

data helped construct and constrain a four-parameter drift-diffusion model of MSN activity. Our

results provide insight into how striatal pathways encode elementary cognitive operations and

elucidate how opposing D2- and D1-MSN activity exerts similar roles in the temporal control of

action. These data demonstrate that D2-MSNs and D1-MSNs play complementary cognition

roles and are highly relevant for understanding the pathophysiology of human diseases and

therapies targeting the striatum.

RESULTS

Interval timing measures elementary cognitive processing in mice.

We investigated cognitive processing in the striatum using a mouse-optimized interval timing

task[33,40]. This task requires mice to *switch* from a short to a long nosepoke response port after

approximately 6 seconds. There is no external cue indicating when to switch response ports,

meaning the decision to switch response ports, or the *switch time*, is an explicit time-based

decision guided by the animals' internal estimate of time (Fig 1a). Interval timing is ideal for

studying elementary cognitive processing in mice using pharmacology, optogenetics, and

neuronal ensemble recordings (Fig 1b) of D2-MSNs and D1-MSNs (Fig 1c-d)[41], as it requires

both working memory and attention, has high relevance for human disease, and requires both

striatal MSNs and nigrostriatal dopamine[16,19–31,35].

Figure 1: Interrogating striatal cognitive contributions to interval timing. a) We examined cognitive behavior during a mouse-optimized interval timing task. In this task, mice are rewarded for switching from a short to a long nosepoke response port after approximately 6 seconds; switching response ports (in green) is an explicit time-based decision, as in other interval timing tasks. b) During surgery, we injected viruses, implanted fiber optics, and implanted electrode or optrode arrays. Then we trained mice to perform the interval timing task; and finally, we performed pharmacological, optogenetic, and neuronal ensemble recording experiments. We manipulated striatal pathways involving c) D2-MSNs (red pseudocolor of Cre-dependent yellow-fluorescent protein (YFP) in a D2-Cre mouse) or d) D1-MSNs (blue pseudocolor of Cre-dependent YFP in a D1-Cre mouse) which project from the striatum (Str) to downstream basal ganglia structures such as the globus pallidus external (GPe) and internal (GPi) segments, as well as the substantia nigra pars reticulata (SNr). Panels a and b were created with BioRender.com.

Disrupting D2-MSNs or D1-MSNs delay interval timing.

In 10 mice (5 female), we found that systemic drug injections targeting D2-dopamine

receptors (D2 blockade; sulpiride 12.5 mg/kg) reliably slowed timing by increasing switch times

relative to saline sessions (median (IQR); saline: 8.9 (8.4-9.8); D2 blockade: 10.1 (9.4-10.8); $p =$

0.002, Cohen's $d = 1.0$; Fig 2a–b)[37,38]. Because systemic D2 blockade can be non-specific, we

leveraged cell-type-specific optogenetics. We virally expressed the inhibitory opsin

halorhodopsin (Halo) in 10 D2-Cre mice (5 female) to inhibit dorsomedial striatal D2-MSNs

(Fig 1c; Fig S1). As with D2 blockade, we found that D2-MSN inhibition reliably increased

switch times (Fig 2c-d; Laser Off: 8.6 seconds (8.3–9.3); Laser On: 10.2 seconds (9.4–10.2);

signed rank $p = 0.002$, Cohen's $d = 1.7$; no effect in no-opsin controls; Fig S2). These data are

consistent with past work demonstrating that dorsomedial D2-MSNs reliably contribute to

timing[36–38].

Next, we investigated D1-MSNs[3,4,42]. In the same 10 mice as above with D2 blockade,

we found that systemic drugs blocking D1-dopamine receptors (D1 blockade; SCH23390 0.05

mg/kg) slowed interval timing by increasing switch times (saline: 9.7 (9.0-9.9); D1 blockade:

10.1 (9.3-11.4); $p = 0.04$, Cohen $d = 0.7$; Fig 2e-f). Furthermore, in 6 D1-Cre mice (3 female),

optogenetic inhibition of dorsomedial striatal D1-MSNs increased switch times (Fig 2g-h; Laser

Off: 8.7 seconds (8.0–9.1); Laser On: 10.5 seconds (9.6–11.1); signed rank $p = 0.03$, Cohen's $d =$

1.4; no effect in no-opsin controls; Fig S1-2). These data suggest that D1-MSNs and D2-MSNs

play similar roles in interval timing because disrupting either MSN type slowed timing by

increasing switch times[38,39].

Figure 2: Disrupting D2- or D1-MSNs increases switch times. Pharmacologically disrupting D2-dopamine receptors (red) with the D2 antagonist sulpiride slowed timing by shifting a**)** cumulative distribution functions to the right, and b) increased switch times; data from 10 wild-type (WT) mice. Optogenetic inhibition of D2-MSNs (red) also slowed timing by shifting c) cumulative distributions to the right, and d) increased switch times; data from 10 D2-Cre mice expressing halorhodopsin (Halo). Similarly, pharmacologically disrupting D1-dopamine receptors (blue) with SCH23390 slowed timing and shifted e) cumulative distribution functions to the right, and f) increased switch times; data from the same 10 WT mice as in a-b. In addition, optogenetic inhibition of D1-MSNs (blue) slowed timing and shifted g) cumulative distributions to the right and h) increased switch times; data from 6 D1-Cre mice expressing Halo. In c, d, f, and h, connected points represent the mean switch time from each animal in each session, and horizontal black lines represent group medians. *p = < 0.05, signed rank test.

Disrupting D2-MSNs or D1-MSNs does not affect task-specific movement.

MSNs can influence motor control, particularly in the dorsolateral striatum[2,3,5,8].

Critically, we found no evidence that inhibiting D2-MSNs in the dorsomedial striatum changed

task-specific movements such as response duration (signed rank $p = 0.63$; Fig 3a-b) or switch

traversal time between the short and long response ports (signed rank $p = 0.49$; Fig 3c).

Similarly, we found no evidence that D1-MSN inhibition changed response duration (signed

rank $p = 0.31$; Fig d-e) or traversal time (signed rank $p = 0.22$; Fig 3f). Furthermore, disrupting

D2-MSNs or D1-MSNs did not change switch time standard deviations or the number of rewards

(Fig S3). These results indicate that disrupting dorsomedial striatal D2-MSNs and D1-MSNs

specifically slowed interval timing without consistently changing task-specific movement.

Figure 3: Optogenetically inhibiting D2-MSNs or D1-MSNs does not affect task-specific motor control.
During interval timing there was no effect of optogenetic inhibition (red) of dorsomedial striatal D2-MSNs on a-
b) switch response duration (time of response port entry to exit) or c) the switch traversal time between the short-
and long-response ports. There was also no effect of optogenetic inhibition (blue) of dorsomedial striatal D1-
MSNs on switch response duration (d-e) or f) switch traversal time. Data from the same 10 D2-Cre mice and 6
D1-Cre mice, as in Figure 2. Horizontal black lines in b, c, e, and f represent group medians.

D2 blockade and D1 blockade degrade MSN temporal encoding.

MSNs strongly encode time[28–31,34,43], but it is unknown how striatal ensembles are

affected by drugs disrupting D2- or D1-dopamine receptors[9]. We investigated this idea by

recording from dorsomedial striatal MSN ensembles in 11 mice during sessions with saline, D2

blockade with sulpiride, or D1 blockade with SCH23390 (Fig 4a; Figs S4 and S5). We analyzed

MSN ensembles in sessions with saline (158 neurons), D2 blockade (167 neurons), or D1

blockade (144 neurons; Fig 4b-c).

We quantified striatal MSN temporal encoding via a naïve Bayesian classifier that

generates trial-by-trial predictions of time from MSN ensemble firing rates (Fig 4d)[28,34].

Inspection of prediction matrices from saline sessions revealed strong temporal predictions early

in the interval (0–6 seconds; $R^2 = 0.91$ (0.83–0.94)) with weaker predictions later in the interval

(6–12 seconds: $R^2 = 0.55$ (0.33–0.70); rank sum $p = 0.00002$ vs 0–6 seconds, Cohen $d = 2.1$; 12–

18 seconds: $R^2 = 0.16$ (0.08–0.64); rank sum $p = 0.00005$ vs 0–6 seconds, Cohen $d = 2.3$; all

sessions were considered statistically independent; Fig 4e). Remarkably, we found that temporal

encoding early in the interval (0–6 seconds) was degraded with either D2 blockade ($R^2 = 0.69$

(0.59–0.84); rank sum $p = 0.0003$ vs saline, Cohen $d = 1.4$) or D1 blockade ($R^2 = 0.70$ (0.47–

0.87); rank sum $p = 0.006$ vs saline, Cohen $d = 1.2$; Fig 4e). Later in the interval (6–12 and 12-18

seconds), there were no significant differences between saline sessions and D2 blockade or D1

blockade (Fig 4e). These data demonstrate that disrupting either D2-MSNs or D1-MSNs

degrades MSN temporal encoding.

Figure 4: D2 and D1 blockade degrade temporal encoding. a) We recorded dorsomedial striatal medium spiny neuron (MSN) ensembles during interval timing in sessions with saline, D2 blockade with sulpiride, or D1 blockade with SCH23390. Made with BioRender.com. b) Peri-event raster from a single MSN in sessions with saline (black), D2-dopamine blockade (red), or D1-dopamine blockade (blue). D2 blockade or D1 blockade changed this MSN's temporal dynamics. c) Neuronal ensemble recording from the same 11 animals with saline, D2 blockade, or D1 blockade; each row represents a peri-event time histogram. Colors indicate z-scored firing rate. d) We used naïve Bayesian classifiers to decode time from MSN ensembles (see Methods and Materials). Each color represents the temporal prediction across 20 trials and red represents stronger predictions. e) Temporal encoding was strong early in the interval, and D2 or D1 blockade degraded classification accuracy. Temporal encoding was decreased later in the interval. Each point represents the R^2 for each trial of behavior for MSN ensembles from 11 mice. * $p < 0.05$ vs saline from 0-6 seconds. Horizontal black lines in (e) represent group medians.

We harnessed principal component analyses (PCA) to identify patterns of MSN ensemble

activity across the interval[28,34,44]. We focused our analyses early in the interval (0–6 seconds)

because temporal encoding during this epoch was powerfully affected by D2 and D1 blockade

(Fig 4e), and this window captured MSN population dynamics before most switch responses (Fig

5a). Population analyses were performed by constructing principal components (PC) from z-

scored peri-event time histograms of firing rate on switch trials. Saline, D2 blockade, and D1

blockade sessions were considered independently (see Fig S6-7 for analyses of neurons across

sessions). PCs were computed from all sessions simultaneously. Examination of the individual

components revealed that the first component (PC1), which explained 55% of neuronal variance,

exhibited "time-dependent ramping", or monotonic changes in firing rate, over the first 6

seconds of the interval. The second component (PC2) explained 31% of the variance (Fig 5b).

Interestingly, PC1 was decreased with both D2 blockade (D2 blockade (0.73 (-6.38–5.08) vs

saline: 3.68 (-4.04–5.78); rank sum $p = 0.03$ vs saline, Cohen's $d = 0.2$) and in sessions with D1

blockade (0.71 (-6.69– 5.19); rank sum $p = 0.01$ vs saline, Cohen's $d = 0.3$; Fig 5c). Few reliable

differences were found in PC2 or higher components (Fig 5d). Together, these findings further

suggest that D2 and D1 blockade have similar effects on MSN population dynamics.

Figure 5: D2 and D1 blockade disrupt prominent patterns of MSN population activity. a) We leveraged principal component analysis to identify patterns of activity among MSN ensembles. The first principal component (PC1) exhibited time-dependent ramping. b) PC1 explained 55% of population variance among MSN ensembles; the second principal component (PC2) explained 31%; higher components explained <10% and were not analyzed. c) PC1 scores were significantly different with D2 or D1 blockade, but d) PC2 scores were not. MSN ensembles from 11 mice. *$p <$ 0.05; data from the same mice as in Figure 4.

D2-MSNs and D1-MSNs have opposite patterns of temporal encoding.

Striatal neuronal populations are composed of MSNs expressing D2-dopamine or D1-

dopamine receptors. We studied dorsomedial D2-MSN vs D1-MSN activity with optogenetic

tagging in transgenic mice in which we could identify each cell class[45–48]. To tag MSNs, we

expressed channelrhodopsin (ChR2) in 4 D2-Cre mice (2 female) and 5 D1-Cre mice (2 female)

and identified D2-MSNs or D1-MSNs neurons by their response to 5 millisecond pulses of 473-

nm light (Fig S8). We tagged 32 D2-MSNs and 41 D1-MSNs (Fig 6a–d). Strikingly, D2-MSN

populations tended to ramp up while D1-MSN populations tended to ramp down (Fig 6c–e).

We used PCA to analyze MSN populations calculated from peri-event matrices from D1-

MSNs and D2-MSNs[47,49]. PCA identified time-dependent ramping activity as PC1 (Fig 6f), a key

temporal signal that explained 56% of variance among tagged neurons[50]. We found that D2-

MSNs and D1-MSNs had opposite patterns of activity with negative PC1 scores for D2-MSNs

and positive PC1 scores for D1–MSNs (Fig 6f–g; PC1 for D2-MSNs: -5.05 (-6.40–3.35); PC1

for D1-MSNs: 4.21 (-4.52–6.92); rank sum p = 0.003, Cohen's d = 0.7). These data imply that

D2-MSN populations tended to ramp up, while D1-MSN populations tended to ramp down

opposite that of D2-MSNs.

To explore these dynamics in detail, we calculated the linear slope of D2-MSN and D1-

MSN activity using generalized linear modeling (GLM) of trial-by-trial firing rate[51,52]. GLM

analysis revealed that D2-MSNs tended to have a positive linear slope of firing rate vs time in

the interval (0.02 spikes/second (-0.07–0.21)), also implying that D2-MSNs tended to ramp up

(Fig 6h; compare with PCA in Fig 6f-g) compared to D1-MSNs, which tended to ramp down (-

0.14 spikes/second (-0.35–0.01); rank sum p = 0.002, Cohen's d = 0.7). Differences between D2-

MSNs and D1-MSNs were consistent over the entire 0–18 second interval (Fig S9). Together,

this line of evidence indicates that D2-MSNs and D1-MSNs have opposing patterns of neuronal

firing over the temporal interval defined by PC1 and by GLM-defined linear slope.

Figure 6: D2-MSNs and D1-MSNs have opposite dynamics. We used optogenetic tagging to record from D2-MSNs and D1-MSNs (Figure S8). Peri-event raster a) from a D2-MSN and b) from a D1-MSN. Histograms derived from the firing rate across trials. c) Peri-event time histograms from D2-MSNs and d) from D1-MSNs. e) D2-MSNs (red) tended to ramp up, whereas D1-MSNs (blue) tended to ramp down. f) Principal component analysis revealed that the first component (PC1) exhibited time-dependent ramping. This component explained ~56% of variance across tagged MSN ensembles. g) Differences between D2-MSNs (red) and D1-MSNs (blue) were captured in PC1, which captures time-dependent ramping. h) These differences were also apparent in the linear slope of firing rate vs time in the interval, with D1-MSNs (blue) having a more negative slope than D2-MSNs (red). Data from 32 tagged D2-MSNs in 4 D2-Cre mice and 41 tagged D1-MSNs in 5 D1-Cre mice. In g and h, each point represents data from each tagged MSN, and horizontal black lines represent group medians. * $p < 0.05$.

Drift-diffusion models capture complementary D2-MSN and D1-MSN dynamics.

Our data demonstrate that D2-MSNs and D1-MSNs have similar effects on interval timing behavior and MSN temporal encoding but opposite activity patterns. We turned to computational modeling to reconcile similar behavioral/encoding effects despite opposing dynamics. Past computational models of interval timing have relied on drift-diffusion dynamics with a positive slope that accumulates evidence over time[53–56]. However, our optogenetic tagging experiments indicated that D2-MSNs tend to ramp up while D1-MSNs tend to ramp down. To demonstrate how these MSNs might complement each other to control movement in time, we constructed a four-parameter drift-diffusion model, where x represents the neuronal firing rate, t represents time measured in seconds, and dx and dt mean "change" in x and t respectively (equivalent to the derivative dx/dt):

$$dx = (F - x)Ddt + \sigma d\xi(t) \qquad \text{(Equation 1)}$$
$$x(0) = b \qquad \text{(Equation 2)}$$

The model has four independent parameters $F, D, \sigma,$ and b (described below) and a threshold value T defined by

$$T = T(F, b) = \left(1 - \tfrac{1}{4}b\right)F + \tfrac{1}{4}b(1 - F). \qquad \text{(Equation 3)}$$

The firing rate is set initially at baseline b (see Equation 2), then driven by input F akin to the dendritic current induced by the overlap of cortico-striatal postsynaptic potentials. With each unit of time dt, we suppose there is corticostriatal input that provides incremental input proportional to the activity itself, $(F - x)D$, to reach a decision. D is *the drift rate* for the event sequence, which with noise $\xi(t)$ (of zero mean and strength σ) leads to fluctuating accumulation and eventually crosses a threshold T (see Equation 3; Fig 7A–B). The time t^* it takes the firing rate to reach the threshold, $x(t^*) = T,$ is the *switch time*.

We report that models like Equations 1–3 generate switch times consistent with gamma-

like response statistics observed in the behavioral data (one-sample Kolmogorov-Smirnov test

for goodness of fit: control $p = 0.27$, inhibition $p = 0.11$; Fig S10). Surprisingly, the same model

strongly and parsimoniously fit four conditions: 1) interval timing behavior for D2-Cre mice

(data from Fig 2d-f; Laser Off trials $R^2 = 0.63$ vs model; Fig 7C, in black); 2) interval timing

behavior for D1-Cre mice (data from Fig 2j-l; trials with Laser Off trials $R^2 = 0.64$ vs model; Fig

7d, in black); 3) interval timing behavioral data with D2-MSN optogenetic inhibition ($R^2 = 0.55$;

Fig 7c, in red); and 4) interval timing behavior with D1-MSN optogenetic inhibition ($R^2 = 0.57$;

Fig 7d, in blue). In Figs 7 and S10, the parameters were chosen as follows. 1) D2: $F = 1$; $b =$

0.52 (so $T = 0.87$); $D = 0.14, \sigma = 0.05$ for control; and $D = 0.14, \sigma = 0.04$ for inhibition. 2)

D1: $F = 0$; $b = 0.48$ (so $T = 0.13$); $D = 0.14, \sigma = 0.05$ for control; and $D = 0.12$, $\sigma = 0.04$

for inhibition. These data refine prior drift-diffusion models with dynamics inspired by

electrophysiology and, with relatively few parameters, can explain interval timing behavior as

well as disruptions in D2- and D1-MSNs[54,55]. This model describes how D2-MSNs and D1-

MSNs make complementary cognitive contributions to the temporal control of action.

In summary, we demonstrated that disrupting either D2-MSNs or D1-MSNs slowed

interval timing and degraded temporal encoding. Surprisingly, we found that D2-MSNs and D1-

MSNs have opposite dynamics during interval timing. Finally, we used these data to construct a

four-parameter drift-diffusion model of interval timing that explains baseline behavior as well as

D2/D1 perturbations. This line of work provides novel insight into cognitive computations by

the striatum.

Figure 7: Four-parameter drift-diffusion computational model of complementary interval timing dynamics. Drift-diffusion computational model of striatal activity during interval timing. a) We modeled interval timing with a low parameter diffusion process with a drift rate D, noise $\xi(t)$, a baseline firing rate b drift toward a threshold T indicated by dotted lines. With D2-MSNs disrupted, this drift process decreases in slope and takes longer to reach the threshold. b) The same model also accounted for D1-MSNs that had an opposite drift slope. Fit between empirical behavioral data (solid lines) and computational models (dashed lines) for both c) D2-Cre mice and d) D1-Cre mice expressing halorhodopsin. Data from inhibition trials are in red (D2-MSNs) or blue (D1-MSNs); data from trials with the Laser Off are in black. Threshold T depends on b and target firing F. This four-parameter model explained timing behavior as well as D2-MSN and D1-MSN disruptions.

DISCUSSION

We explored how striatal MSN ensembles encode an elementary cognitive process,

interval timing. We found that pharmacologically or optogenetically disrupting striatal D2-MSNs

or D1-MSNs slowed timing, increasing an animal's self-reported estimates of time without

affecting task-specific motor behavior. Furthermore, D2 blockade or D1 blockade degraded

MSN temporal encoding. To our surprise, optogenetic tagging revealed that D2-MSNs and D1-

MSNs had opposing patterns of activity with D2-MSNs ramping up over the interval and D1-

MSNs ramping down. We reconciled these opposing dynamics and similar behavioral and

temporal encoding effects of D2-MSN and D1-MSN perturbations with a four-parameter drift-

diffusion model. Taken together, these data provide new insight into how opposing patterns of

striatal MSN activity might influence behavior in similar ways, suggesting that they make

complementary contributions to elementary cognitive operations.

Striatal MSNs are critical for temporal control of action[29,57]. Three broad models have

been proposed for how striatal MSN ensembles represent time: 1) the striatal beat frequency

model, in which MSNs encode temporal information based on neuronal synchrony[58]; 2) the

distributed coding model, in which time is represented by the state of the network[29,59]; and 3) the

drift-diffusion model, in which neuronal activity monotonically drifts toward a threshold after

which responses are initiated[30,54,60]. While our data do not formally resolve these possibilities,

our results suggest that D2-MSN and D1-MSN activity exhibit opposing firing rate dynamics

with linear changes in firing rates over the interval. Furthermore, we find that low-parameter

drift-diffusion models account for interval timing behavior with both intact and disrupted striatal

D2- and D1-MSNs. While it is possible that other models might capture interval timing behavior

and account for MSN neuronal activity, our model does so parsimoniously with relatively few

parameters.

D2-MSNs and D1-MSNs have been proposed to play complementary roles in movement.

For instance, stimulating D1-MSNs facilitates movement, whereas stimulating D2-MSNs

impairs movement[5]. Both populations have been shown to have complementary patterns of

activity during movements[3,4], with MSNs firing at different phases of action initiation and

selection. Further dissection of action selection programs reveals that opposing patterns of

activation among D2-MSNs and D1-MSNs suppress and guide actions, respectively, in the

dorsolateral striatum[2]. Our data extend these findings to interval timing, which is an elementary

cognitive process requiring working memory for temporal rules as well as attention to the

passage of time. A particular advantage of interval timing is that it captures a cognitive behavior

within a single dimension — time. When projected along the time dimension, it is remarkable

that D2-MSNs and D1-MSNs have opposing patterns of activity. Our behavioral results and

computational modeling suggest that disrupting either D2-MSNs or D1-MSNs increased self-

reported estimates of time, indicating that disrupting either D2-MSN or D1-MSN activity

degrades temporal encoding. Our recording experiments do not identify where a possible

response threshold T is instantiated, but downstream basal ganglia structures may have a key role

in setting response thresholds[61].

Since interval timing is reliably disrupted in human diseases of the striatum such as

Huntington's disease, Parkinson's disease, and schizophrenia[16,21,62,63], our results have relevance

to human disease. Furthermore, because many therapeutics targeting dopamine receptors are

used clinically, our results may help illuminate how dopaminergic drugs affect cognitive

function and dysfunction[64,65]. Finally, D2-MSNs and D1-MSNs define indirect and direct

pathways, which are targeted by deep brain stimulation for neurological and psychiatric

disease[66,67].

Our approach has several limitations. First, systemic drug injections will block D2- and

D1-receptors in many different brain regions, including the frontal cortex, which we have shown

to be involved in interval timing[47]. D2 blockade or D1 blockade may have complex effects,

including corticostriatal or network effects that contribute to changes in D2-MSN or D1-MSN

ensemble activity[9]. Second, optogenetic tagging is low-yield, and it is possible that recording

more of these neurons would afford greater opportunity to identify alternative coding schemes,

such as neuronal synchrony[58]. Third, the striatum includes diverse cell types that may play a key

role[68]. Fourth, MSNs can laterally inhibit each other, which may profoundly affect striatal

function[69]. Regardless, we show that cell-type-specific disruption of D2-MSNs and D1-MSNs

both slow timing, implying that these cell types play a key role in interval timing, and our

optogenetic tagging experiments describe their activity at high temporal resolution. Future

experiments may record from non-MSN striatal cell types, including fast-spiking interneurons

that shape basal ganglia output[70]. Fifth, there may be key anatomic differences between dorsal

striatal regions that we did not capture[71]. Finally, we did not deliver stimulation to the striatum,

because our pilot experiments triggered movement artifacts or task-specific dyskinesias[72,73].

Future stimulation approaches carefully titrated to striatal physiology may influence interval

timing without affecting movement[48].

In summary, we examined the role of dorsomedial striatal D2-MSNs and D1-MSNs

during an elementary cognitive behavior, interval timing. We found that disrupting either D2-

MSNs or D1-MSNs impaired estimates of time and degraded temporal encoding by striatal

ensembles. Optogenetic tagging revealed that D2-MSNs and D1-MSNs exhibited opposite,

complementary patterns of neuronal activity. Finally, these patterns could be captured by

computational drift-diffusion models. Collectively, our data provide insight into how the striatum

encodes cognitive information, which could be highly relevant for human diseases that disrupt

the striatum and for next-generation neuromodulation that targets the basal ganglia to avoid

cognitive side effects or to treat cognitive dysfunction.

Methods and Materials

Rodents. All procedures were approved by the Institutional Animal Care and Use Committee (IACUC) at the University of Iowa, and all experimental methods were performed in accordance with applicable guidelines and regulations (Protocol #0062039). We used three mouse lines. For behavioral pharmacology experiments, we used C57BL/6J mice (10 mice (5 female); The Jackson Laboratory, Bar Harbor, ME). For cell-type-specific optogenetics and electrophysiology, we used either *Drd2-cre+* mice derived from Gensat strain ER44 (10 mice (5 female) for optogenetic inhibition; 5 mice (2 female) for optogenetic controls; 4 mice (2 female) for tagging)[74] or *Drd1-cre+* mice derived from Gensat strain EY262 (6 mice (3 female) for optogenetic inhibition; 5 mice (3 female) optogenetic controls; 5 mice (2 female) for tagging). For combined behavioral pharmacology and electrophysiology, we used 11 mice (4 C57BL/6J (2 female), 5 *Drd2-cre+* mice (2 female), and 2 *Drd1-cre+* mice (0 female)).

Interval timing switch task. We used a mouse-optimized interval timing task described in detail previously[33,34,37,40,75,76]. Briefly, mice were trained in sound-attenuating operant chambers, with two nosepoke response ports flanking either side of a food hopper on the front wall, and a third nosepoke response port located at the center of the back wall. The chamber was positioned below an 8-kHz, 72-dB speaker (Fig 1A; MedAssociates, St. Albans, VT). Mice were 85% food restricted and motivated with 20 mg sucrose pellets (Bio-Serv, Flemington, NJ). Mice were initially trained to receive sucrose rewards during fixed ratio nosepoke response trials. After shaping, mice were trained in the "switch" interval timing task[33]. Mice initiated trials at the back response port, after which a tone and nosepoke lights were illuminated simultaneously. Response port entry and exit were captured by infrared beams. On short trials (50% of randomly selected

trials), mice were rewarded after 6 seconds for nosepoke responses at the designated short

response port (left or right). On long trials, mice were rewarded after 18 seconds for responding

first at the short response port and then switching responses to the long response port (left or

right). Critically, the decision to switch from the short to the long response port is a time-based

decision guided by internal estimates of time. All analyses were focused on long switch trials

when animals depart from the short response port. *Switch time* was defined as the moment an

animal exited the short response port prior to entering the long response port. We focused on the

switch time as a representation of temporal control of action. Switch *response duration* was

defined as the time between short response entry and exit for the switch responses only. Switch

traversal time was defined as the duration between short response port exit and long response

port entry.

Behavioral pharmacology procedures. C57BL/6J mice were injected intraperitoneally (IP) 20-

40 minutes before interval timing trials with either SCH23390 ($C_{17}H_{18}ClNO$; D1 antagonist),

sulpiride ($C_{15}H_{23}N_3O_4S$; D2 antagonist), or isotonic saline. The sulpiride dosage was 12.5 mg/kg,

0.01 mL/g, and the SCH23390 was administered at a dosage of 0.05 mg/kg, 0.01 mL/g.

Behavioral performance was compared with interval timing behavior on the prior day when

isotonic saline was injected IP.

Surgical and histological procedures. Surgical procedures were identical to methods described

previously[34]. Briefly, mice were anesthetized using inhaled 4% isoflurane and surgical levels of

anesthesia were maintained at 1-2% for the duration of the surgery. Craniotomies were drilled

above bilateral dorsal striatal anatomical targets, and optogenetic viruses (AAV5-DIO-

eNHPR2.0 (halorhodopsin), AAV5-DIO-ChR2(H134R)-mcherry (ChR), or AAV5-DIO-cherry

(control) from the University of North Carolina Viral Vector Core) were injected (0.5 uL of

virus, +0.9, ML +/-1.3, DV –2.7). Either fiber optics (Doric Lenses, Montreal Quebec; AP +0.9,

ML +/-1.3, DV –2.5) or 4 x 4 electrode or optrode arrays (AP +0.4, ML -1.4, DV -2.7 on the left

side only; Microprobes, Gaithersburg, MD) were positioned in the dorsal striatum. Holes were

drilled to insert skull screws to anchor headcap assemblies and/or ground electrode arrays and

then sealed with cyanoacrylate ("SloZap," Pacer Technologies, Rancho Cucamonga, CA),

accelerated by "ZipKicker" (Pacer Technologies) and methyl methacrylate (AM Systems, Port

Angeles, WA). Following postoperative recovery, mice were trained on the switch task and

acclimated to experimental procedures prior to undergoing experimental sessions.

Immunohistochemistry. Following completion of experiments, mice were transcardially

perfused with ice-cold 1x phosphate-buffered saline (PBS) and 4% paraformaldehyde (PFA)

after anesthesia using ketamine (100 mg/kg IP) and xylazine (10 mg/kg IP). Brains were then

fixed in solutions of 4% PFA and 30% sucrose prior to being cryosectioned on a freezing

microtome. Sections were stained for tyrosine hydroxylase with primary antibodies for >12

hours (rabbit anti-TH; Millipore MAB152; 1:1000) at 4° C. Sections were subsequently

visualized with Alexa Fluor fluorescent secondary antibodies (goat anti-rabbit IgG Alexa 519;

Thermo Fisher Scientific; 1:1,000) matched to host primary by incubating 2 hours at room

temperature. Histological reconstruction was completed using postmortem analysis of electrode

placement by slide-scanning microscopy on an Olympus VS120 microscope (Olympus, Center

Valley, PA; Fig S4).

Electrophysiology. Single-unit recordings were made using a multi-electrode recording system

(Open Ephys, Atlanta, GA), After the experiments, Plexon Offline Sorter (Plexon, Dallas, TX),

was used to remove artifacts. Spike activity was analyzed for all cells that fired at rates above 0.1

Hz. Principal component analysis (PCA) and waveform shape were used for spike sorting. Single

units were defined as those 1) having a consistent waveform shape, 2) being a separable cluster

in PCA space, and 3) having a consistent refractory period of at least 2 milliseconds in interspike

interval histograms. Putative MSNs were further separated from striatal fast-spiking interneurons

(FSIs) based on hierarchical clustering of the waveform peak-to-trough ratio and the half-peak

width (*fitgmdist* and *cluster.m*; Fig S5)[77]. We calculated kernel density estimates of firing rates

across the interval (-4 seconds before trial start to 22 seconds after trial start) binned at 0.1

seconds, with a bandwidth of 1. We used PCA to identify data-driven patterns of z-scored

neuronal activity, as in our past work[28,34,78–80]. Average plots were shown with Gaussian

smoothing for plotting purposes only.

Optogenetics. We leveraged cell-type-specific optogenetics to manipulate D2-MSNs and D1-

MSNs in D2- or D1-cre mice. In animals injected with optogenetic viruses, optical stimulation

was delivered via bilateral patch cables for the entire trial duration of 18 seconds via 589-nm

laser light at 12 mW power on 50% of randomly assigned trials. For optogenetic tagging,

putative D1- and D2-MSNs were optically identified via 473-nm photostimulation. Units with

mean post-stimulation spike latencies of ≤5 milliseconds and a stimulated-to-unstimulated

waveform correlation ratio of >0.9 were classified as putative D2+ or D1+ neurons[81,82].

Modeling. We proposed a four-parameter drift-diffusion model (see Equations 1-3 in Results)

and used it to simulate response times compatible with mice behavioral data. The model was

implemented in MATLAB, starting from initial value b and with discrete computer simulation

steps $x_{new} = x_{old} + (F - x_{old})D\Delta t + \sigma\sqrt{\Delta t}N(0,1)$. Here $N(0,1)$ represents random values

generated from the standardized Gaussian distribution (mean = 0 and standard deviation = 1).

The integration timestep was taken $\Delta t = 0.1$ to be like the 0.1-second bin size used in the

analysis of firing rate data. For each numerical simulation, the "switch time" was defined as the

time t^* when variable x first reached the threshold value $T = F(1 - b/4) + (1 - F)b/4$. For

each condition, we ran 500 simulations of the model (up to 25 seconds per trial) and recorded the

switch times. Examples of firing rate x dynamics are shown in Figure 7a-b. We observed that the

distributions of interval timing response times could be fit by a gamma probability distribution

function $PDF(x) = \frac{\beta^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\beta x}$ with shape α and rate β (see Fig S10). For D2-Cre data, we

found a good model fit: $\alpha = 6.03$, $\beta = 0.67$ (empirical, Laser Off); $\alpha = 6.08$, $\beta = 0.69$ (model,

Laser Off); $\alpha = 7.62$, $\beta = 0.78$ (empirical, Inhibited); and $\alpha = 7.67$, $\beta = 0.79$ (model, Inhibited);

Fig S10, left two columns. The fit for D1-Cre data was found at $\alpha = 5.87$, $\beta = 0.68$ (empirical,

Laser Off); $\alpha = 5.89$, $\beta = 0.69$ (model, Laser Off); $\alpha = 7.13$, $\beta = 0.70$ (empirical, Inhibited); and

$\alpha = 7.31$, $\beta = 0.72$ (model, Inhibited); Fig S10, right two columns.

Statistics. All data and statistical approaches were reviewed by the Biostatistics, Epidemiology,

and Research Design Core (BERD) at the Institute for Clinical and Translational Sciences

(ICTS) at the University of Iowa. All code and data are made available at

http://narayanan.lab.uiowa.edu/article/datasets. We used the median to measure central tendency

and the interquartile range to measure spread. We used Wilcoxon nonparametric signed rank or

rank-sum tests to compare behavior between experimental conditions and Cohen's d to calculate

effect size. When warranted (e.g., for paired comparisons of mean values, as in Fig 2b) we

applied the signed rank test instead. Analyses of putative single-unit activity and basic

physiological properties were carried out using custom routines for MATLAB[28,34,49,78].

Time-dependent ramping slope. To measure time-related ramping, we used generalized linear

models (GLMs) at the individual neuron level in which the response variable was firing rate

binned at 0.1 seconds and the predictor variable was time in the interval[83]. For each neuron, its

time-related "ramping" slope was derived from the GLM fit of firing rate vs time in the interval,

for all trials per neuron (see Fig 6). All models were run at a trial-by-trial level to avoid effects of

trial averaging[51] as in our past work[28,34,78,84].

Bayesian classifier. As in our past work[28,78,79], we used a naïve Bayesian classifier to evaluate

neuronal ensemble decoding from all neurons with at least 20 trials. To preclude edge effects that

might bias classifier performance, we included data from 6 seconds prior to the trial start and 6

seconds after the end of the interval. We used leave-one-out cross-validation to predict an

objective time from the firing rate within a trial. Classifier performance was quantified by

computing the R^2 of objective time vs predicted time, only for bins during the interval (0-6, 6-12,

and 12-18 seconds; see Fig 4). Classifier performance was compared using time-shuffled firing

rates via a Wilcoxon signed rank test.

Data Availability

All raw data are available at http://narayanan.lab.uiowa.edu/article/datasets.

Code Availability

All code is available at http://narayanan.lab.uiowa.edu/article/datasets.

Acknowledgements

This work was funded by NIMH R01MH116043 to NSN

Author Contributions

RAB, MAW, and NN designed the experiments. RAB, CIAB, MAW, RV, and CJ performed all experiments and collected data. The data was analyzed by RAB, MAW, ASB, YK, RC, and NN. RC developed the computational model. AB, MAW, ASB, KS, RC, and NN wrote the manuscript, and all authors reviewed and revised the manuscript.

Extended Data

Figure S1: Fiber optic locations from a) an opsin-expressing mouse with mcherry-tagged

halorhodopsin and bilateral fiber optics, and b) across 10 D2-Cre mice (red) and 6 D1-cre mice

(blue) with fiber optics.

Figure S2: Experiments in D2-Cre mice injected with virus without opsins did not reliably affect

a) cumulative density functions (CDFs) or b) switch times (signed rank $p = 0.44$). Experiments in

D1-Cre mice expressing virus without opsins did not reliably affect c) CDFs or d) switch times

(signed rank $p = 0.81$).

Figure S3: a) Standard deviation of interval timing performance for D2-MSN inhibition sessions

(signed rank test, $p = 0.19$) and b) D1-MSN inhibition sessions ($p = 0.84$), and the number of

total rewards for c) D2-MSN inhibition sessions ($p = 0.07$) and d) D1-MSN inhibition sessions

($p = 0.24$). Data from 10 D2-Cre mice and 6 D1-Cre mice as in Figs 2 and 3.

Figure S4: Recording locations in the dorsomedial striatum. Left panel: electrode

reconstructions for D2-Cre (red), D1-Cre (blue), and wild-type mice (green). Right panel:

reversed image of ChR2 and DAPi stain used for electrode reconstruction (white arrows). Only

the left striatum was implanted with electrodes in all animals.

Figure S5: MSN classification by waveform criteria for sessions with a) pharmacology and

electrophysiology and b) optogenetic tagging.

Figure S6: We sorted and identified the same 99 MSNs across saline sessions and sessions with

D2 blockade or D1 blockade. Each row is a z-scored peri-event time histogram of MSN activity

for all trials during interval timing. Each row is the same neuron identified by waveform and

interspike-intervals across saline, D2 blockade, and D1 blockade sessions.

Figure S7: Early-interval ensemble dynamics respond to D2 and D1 blockade. a) We further

explored early-interval dynamics by PCA in MSN ensembles with matched sorting among

saline, D2 blockade, and D1 blockade (99 neurons; see Fig S6). The matched sorting across

sessions enabled paired signed rank analyses. b) Strikingly, the first component (PC1) explained

75% of variance among MSN ensemble dynamics between 3-6 seconds. c) As with dynamics

between 0-6 seconds, PC1 was shifted relative to saline for D2 blockade (signed rank $p =$

0.000007) and D1 blockade (signed rank $p = 0.002$). We identified two clusters based on PC1,

with cluster 1 (d) loading positively on PC1, and cluster 2 (e) loading negatively on PC1. For

cluster 1 in (d), PCs were distinct from saline for D2 blockade (signed rank $p = 3\text{x}10^{-12}$) and for

D1 blockade (signed rank $p = 10^{-7}$). For cluster 2 in panel (e), PC1 changes were also distinct for

saline for D2 blockade (signed rank $p = 0.00008$) and for D1 blockade (signed rank $p = 0.002$).

Figure S8: a) Electrode tips in the dorsal striatum, where b) we recorded optogenetically

"tagged" neurons that fired action potentials (red) within ≤5 milliseconds of laser onset (red

line).

546
Figure S9: D2- and D1-MSN activity over the whole 18-second interval. a) Peri-event time

histograms from D2-MSNs and b) from D1-MSNs over the whole interval. c) We noticed that on

average, D2-MSNs tended to ramp up (red), whereas D1-MSNs tended to ramp down (blue). d)

Principal component analysis revealed that PC1 exhibited time-dependent ramping over the

whole 18-second interval. This component explained ~45% of variance across tagged MSN

ensembles. e) Over the whole interval, differences between D2-MSNs and D1-MSNs were

captured in PC1, which captures time-dependent ramping (rank sum $p = 0.02$). F) These

differences were also captured via differences in a linear slope of firing rate vs time over the

whole 18-second interval, with D1-MSNs having a more negative slope (rank sum $p = 0.03$).

Data from 32 tagged D2-MSNs in 10 D2-Cre mice and 41 tagged D1-MSNs in 6 D1-Cre mice.

558
559

560
561 **Figure S10:** Histograms of behavioral data from D2-Cre mice with a) Laser Off and b) D2-

562 MSN inhibition (red). Data from D1-Cre mice with c) Laser Off and with d) D1-MSN inhibition

563 (blue). e-h) model predictions. i-j) Comparisons of empirical data vs model. All panels: fits for

564 the gamma distribution with dotted circles. Data from 10 D2-Cre mice and 6 D1-mice from Fig

565 2.

REFERENCES

1. Albin, R. L., Young, A. B. & Penney, J. B. The functional anatomy of basal ganglia disorders. *Trends Neurosci.* **12**, 366–375 (1989).
2. Cruz, B. F. *et al.* Action suppression reveals opponent parallel control via striatal circuits. *Nature* **607**, 521–526 (2022).
3. Cui, G. *et al.* Concurrent activation of striatal direct and indirect pathways during action initiation. *Nature* **494**, 238–242 (2013).
4. Tecuapetla, F., Jin, X., Lima, S. Q. & Costa, R. M. Complementary Contributions of Striatal Projection Pathways to Action Initiation and Execution. *Cell* **166**, 703–715 (2016).
5. Kravitz, A. V. *et al.* Regulation of parkinsonian motor behaviours by optogenetic control of basal ganglia circuitry. *Nature* **466**, 622–626 (2010).
6. Alexander, G. E. & Crutcher, M. D. Functional architecture of basal ganglia circuits: neural substrates of parallel processing. *Trends Neurosci.* **13**, 266–271 (1990).
7. West, A. R. & Grace, A. A. Opposite Influences of Endogenous Dopamine D1 and D2 Receptor Activation on Activity States and Electrophysiological Properties of Striatal Neurons: Studies CombiningIn Vivo Intracellular Recordings and Reverse Microdialysis. *J. Neurosci.* **22**, 294–304 (2002).
8. Jin, X., Tecuapetla, F. & Costa, R. M. Basal ganglia subcircuits distinctively encode the parsing and concatenation of action sequences. *Nat. Neurosci.* **17**, 423–430 (2014).
9. Yun, S. *et al.* Antipsychotic drug efficacy correlates with the modulation of D1 rather than D2 receptor-expressing striatal projection neurons. *Nat. Neurosci.* 1–12 (2023) doi:10.1038/s41593-023-01390-9.
10. Graybiel, A. M. The Basal Ganglia and Cognitive Pattern Generators. *Schizophr. Bull.* **23**, 459–469 (1997).
11. Middleton, F. A. & Strick, P. L. Basal ganglia and cerebellar loops: motor and cognitive circuits. *Brain Res. Rev.* **31**, 236–250 (2000).
12. Wall, N. R., De La Parra, M., Callaway, E. M. & Kreitzer, A. C. Differential Innervation of Direct- and Indirect-Pathway Striatal Projection Neurons. *Neuron* **79**, 347–360 (2013).
13. Averbeck, B. B., Lehman, J., Jacobson, M. & Haber, S. N. Estimates of projection overlap and zones of convergence within frontal-striatal circuits. *J. Neurosci.* **34**, 9497–9505 (2014).
14. Seo, M., Lee, E. & Averbeck, B. B. Action Selection and Action Value in Frontal-Striatal Circuits. *Neuron* **74**, 947–960 (2012).
15. Andreasen, N. C. Understanding the causes of schizophrenia. *N. Engl. J. Med.* **340**, 645–647 (1999).
16. Hinton, S. C. *et al.* Motor timing variability increases in preclinical Huntington's disease patients as estimated onset of motor symptoms approaches. *J. Int. Neuropsychol. Soc.* **13**, 539–543 (2007).
17. Narayanan, N. S. & Albin, R. L. *Cognition in Parkinson's Disease*. (Elsevier, Acad. Press, 2022).
18. Brown, S. W., Collier, S. A. & Night, J. C. Timing and Executive Resources: Dual-Task Interference Patterns Between Temporal Production and Shifting, Updating, and Inhibition Tasks. *J. Exp. Psychol. Hum. Percept. Perform.* (2012) doi:10.1037/a0030484.
19. Parker, K. L., Lamichhane, D., Caetano, M. S. & Narayanan, N. S. Executive dysfunction in Parkinson's disease and timing deficits. *Front. Integr. Neurosci.* **7**, 75 (2013).
20. Merchant, H. & de Lafuente, V. Introduction to the neurobiology of interval timing. *Adv. Exp. Med. Biol.* **829**, 1–13 (2014).
21. Singh, A. *et al.* Timing variability and midfrontal ~4 Hz rhythms correlate with cognition in Parkinson's disease. *NPJ Park. Dis.* **7**, 14 (2021).
22. Merchant, H., Harrington, D. L. & Meck, W. H. Neural Basis of the Perception and Estimation of Time. *Annu. Rev. Neurosci.* **36**, 313–336 (2013).
23. Hamilos, A. E. *et al.* Slowly evolving dopaminergic activity modulates the moment-to-moment probability of reward-related self-timed movements. *eLife* **10**, e62583 (2021).
24. Howard, C. D., Li, H., Geddes, C. E. & Jin, X. Dynamic Nigrostriatal Dopamine Biases Action Selection. *Neuron* **93**, 1436-1450.e8 (2017).

25. Meck, W. H. Neuroanatomical localization of an internal clock: A functional link between
mesolimbic, nigrostriatal, and mesocortical dopaminergic systems. *Brain Res.* **1109**, 93–107 (2006).
26. Soares, S., Atallah, B. V. & Paton, J. J. Midbrain dopamine neurons control judgment of time.
Science **354**, 1273–1277 (2016).
27. Matell, M. S., Meck, W. H. & Nicolelis, M. A. L. Interval timing and the encoding of signal duration
by ensembles of cortical and striatal neurons. *Behav. Neurosci.* **117**, 760–773 (2003).
28. Emmons, E. B. *et al.* Rodent medial frontal control of temporal processing in the dorsomedial
striatum. *J. Neurosci.* **37**, 8718–8733 (2017).
29. Gouvea, T. S. *et al.* Striatal dynamics explain duration judgments. *eLife* **4**, (2015).
30. Wang, J., Narain, D., Hosseini, E. A. & Jazayeri, M. Flexible timing by temporal scaling of cortical
responses. *Nat. Neurosci.* **21**, 102 (2018).
31. Mello, G. B. M., Soares, S. & Paton, J. J. A Scalable Population Code for Time in the Striatum. *Curr.*
Biol. **25**, 1113–1122 (2015).
32. Monteiro, T. *et al.* Using temperature to analyze the neural basis of a time-based decision. *Nat.*
Neurosci. (2023) doi:10.1038/s41593-023-01378-5.
33. Balci, F. *et al.* Interval timing in genetically modified mice: a simple paradigm. *Genes Brain Behav.*
7, 373–384 (2008).
34. Bruce, R. A. *et al.* Experience-related enhancements in striatal temporal encoding. *Eur. J. Neurosci.*
(2021).
35. Buhusi, C. V. & Meck, W. H. What makes us tick? Functional and neural mechanisms of interval
timing. *Nat. Rev. Neurosci.* **6**, 755–765 (2005).
36. Drew, M. R. *et al.* Transient Overexpression of Striatal D2 Receptors Impairs Operant Motivation
and Interval Timing. *J. Neurosci.* **27**, 7731–7739 (2007).
37. Stutt, H. R. *et al.* Sex similarities and dopaminergic differences in interval timing. *BioRxiv Prepr.*
Serv. Biol. 2023.05.05.539584 (2023) doi:10.1101/2023.05.05.539584.
38. De Corte, B. J., Wagner, L. M., Matell, M. S. & Narayanan, N. S. Striatal dopamine and the temporal
control of behavior. *Behav. Brain Res.* **356**, 375–379 (2019).
39. Kamada, T. & Hata, T. Striatal dopamine D1 receptors control motivation to respond, but not interval
timing, during the timing task. *Learn. Mem. Cold Spring Harb. N* **28**, 24–29 (2021).
40. Tosun, T., Gur, E. & Balci, F. Mice plan decision strategies based on previously learned time
intervals, locations, and probabilities. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 787–792 (2016).
41. Foster, N. N. *et al.* The mouse cortico–basal ganglia–thalamic network. *Nature* **598**, 188–194 (2021).
42. Calabresi, P., Picconi, B., Tozzi, A., Ghiglieri, V. & Di Filippo, M. Direct and indirect pathways of
basal ganglia: a critical reappraisal. *Nat. Neurosci.* **17**, 1022–1030 (2014).
43. Bakhurin, K. I. *et al.* Differential Encoding of Time by Prefrontal and Striatal Network Dynamics. *J.*
Neurosci. **37**, 854–870 (2017).
44. Narayanan, N. S. & Laubach, M. Delay activity in rodent frontal cortex during a simple reaction time
task. *J. Neurophysiol.* **101**, 2859–2871 (2009).
45. Kvitsiani, D. *et al.* Distinct behavioural and network correlates of two interneuron types in prefrontal
cortex. *Nature* **498**, 363–366 (2013).
46. Pi, H.-J. *et al.* Cortical interneurons that specialize in disinhibitory control. *Nature* (2013)
doi:10.1038/nature12676.
47. Kim, Y.-C. *et al.* Optogenetic Stimulation of Frontal D1 Neurons Compensates for Impaired
Temporal Control of Action in Dopamine-Depleted Mice. *Curr. Biol. CB* **27**, 39–47 (2017).
48. Kim, Y. *et al.* Phase-adaptive brain stimulation of striatal D1 medium spiny neurons in dopamine-
depleted mice. *Sci. Rep.* **12**, 21780 (2022).
49. Parker, K. L., Chen, K.-H., Kingyon, J. R., Cavanagh, J. F. & Narayanan, N. S. D1-dependent 4 Hz
oscillations and ramping activity in rodent medial frontal cortex during interval timing. *J. Neurosci.*
34, 16774–16783 (2014).
50. Narayanan, N. S. Ramping activity is a cortical mechanism of temporal control of action. *Curr. Opin.*
Behav. Sci. **8**, 226–230 (2016).

51. Latimer, K. W., Yates, J. L., Meister, M. L., Huk, A. C. & Pillow, J. W. Single-trial spike trains in
parietal cortex reveal discrete steps during decision-making. *Science* **349**, 184–187 (2015).
52. Zylberberg, A. & Shadlen, M. N. Cause for pause before leaping to conclusions about stepping.
bioRxiv 085886 (2016) doi:10.1101/085886.
53. Luzardo, A., Ludvig, E. A. & Rivest, F. An adaptive drift-diffusion model of interval timing
dynamics. *Behav. Processes* **95**, 90–99 (2013).
54. Simen, P., Balci, F., de Souza, L., Cohen, J. D. & Holmes, P. A model of interval timing by neural
integration. *J. Neurosci. Off. J. Soc. Neurosci.* **31**, 9238–9253 (2011).
55. Treisman, M. Temporal discrimination and the indifference interval: Implications for a model of the
'internal clock'. *Psychol. Monogr. Gen. Appl.* **77**, 1–31 (1963).
56. Nguyen, Q.-A., Rinzel, J. & Curtu, R. Buildup and bistability in auditory streaming as an evidence
accumulation process with saturation. *PLOS Comput. Biol.* **16**, e1008152 (2020).
57. Meck, W. H., Penney, T. B. & Pouthas, V. Cortico-striatal representation of time in animals and
humans. *Curr Opin Neurobiol* **18**, 145–52 (2008).
58. Matell, M. S. & Meck, W. H. Cortico-striatal circuits and interval timing: coincidence detection of
oscillatory processes. *Cogn. Brain Res.* **21**, 139–170 (2004).
59. Paton, J. J. & Buonomano, D. V. The Neural Basis of Timing: Distributed Mechanisms for Diverse
Functions. *Neuron* **98**, 687–705 (2018).
60. Emmons, E. B. *et al.* Rodent Medial Frontal Control of Temporal Processing in the Dorsomedial
Striatum. *J. Neurosci.* **37**, 8718–8733 (2017).
61. Toda, K. *et al.* Nigrotectal Stimulation Stops Interval Timing in Mice. *Curr. Biol.* **27**, 3763-3770.e3
(2017).
62. Penney, T. B., Meck, W. H., Roberts, S. A., Gibbon, J. & Erlenmeyer-Kimling, L. Interval-timing
deficits in individuals at high risk for schizophrenia. *Brain Cogn.* **58**, 109–118 (2005).
63. Ward, R. D., Kellendonk, C., Kandel, E. R. & Balsam, P. D. Timing as a window on cognition in
schizophrenia. *Neuropharmacology* (2011) doi:10.1016/j.neuropharm.2011.04.014.
64. Buhusi, C. V. & Meck, W. H. Differential effects of methamphetamine and haloperidol on the control
of an internal clock. *Behav. Neurosci.* **116**, 291 (2002).
65. Friedman, J. H. & Lannon, M. C. Clozapine in the treatment of psychosis in Parkinson's disease.
Neurology **39**, 1219–1219 (1989).
66. Bergfeld, I. O. *et al.* Deep Brain Stimulation of the Ventral Anterior Limb of the Internal Capsule for
Treatment-Resistant Depression: A Randomized Clinical Trial. *JAMA Psychiatry* **73**, 456–464
(2016).
67. Okun, M. S. Deep-brain stimulation for Parkinson's disease. *N. Engl. J. Med.* **367**, 1529–1538
(2012).
68. Gokce, O. *et al.* Cellular Taxonomy of the Mouse Striatum as Revealed by Single-Cell RNA-Seq.
Cell Rep. **16**, 1126–1137 (2016).
69. Burke, D. A., Rotstein, H. G. & Alvarez, V. A. Striatal Local Circuitry: A New Framework for
Lateral Inhibition. *Neuron* **96**, 267–284 (2017).
70. Owen, S. F., Berke, J. D. & Kreitzer, A. C. Fast-Spiking Interneurons Supply Feedforward Control of
Bursting, Calcium, and Plasticity for Efficient Learning. *Cell* **172**, 683-695.e15 (2018).
71. Gerfen, C. R. The neostriatal mosaic: compartmentalization of corticostriatal input and striatonigral
output systems. *Nature* **311**, 461–464 (1984).
72. F Hernandez, L., Castela, I., Ruiz-DeDiego, I., Obeso, J. A. & Moratalla, R. Striatal activation by
optogenetics induces dyskinesias in the 6-hydroxydopamine rat model of Parkinson disease. *Mov.*
Disord. Off. J. Mov. Disord. Soc. **32**, 530–537 (2017).
73. Freeze, B. S., Kravitz, A. V., Hammack, N., Berke, J. D. & Kreitzer, A. C. Control of basal ganglia
output by direct and indirect pathway projection neurons. *J. Neurosci. Off. J. Soc. Neurosci.* **33**,
18531–18539 (2013).
74. Gong, S. *et al.* Targeting Cre recombinase to specific neuron populations with bacterial artificial
chromosome constructs. *J. Neurosci. Off. J. Soc. Neurosci.* **27**, 9817–9823 (2007).

75. Larson, T. *et al.* Mice expressing P301S mutant human tau have deficits in interval timing. *Behav.*
Brain Res. **432**, 113967 (2022).
76. Weber, M. A. *et al.* Glycolysis-enhancing α1-adrenergic antagonists modify cognitive symptoms
related to Parkinson's disease. *Npj Park. Dis.* **9**, 1–7 (2023).
77. Berke, J. D. Functional properties of striatal fast-spiking interneurons. *Front. Syst. Neurosci.* **5**, 45
(2011).
78. Emmons, E. *et al.* Temporal Learning Among Prefrontal and Striatal Ensembles. *Cereb. Cortex*
Commun. **1**, (2020).
79. Emmons, E. B., Kennedy, M., Kim, Y. & Narayanan, N. S. Corticostriatal stimulation compensates
for medial frontal inactivation during interval timing. *Sci. Rep.* **9**, 1–9 (2019).
80. Narayanan, N. S. & Laubach, M. Methods for studying functional interactions among neuronal
populations. in *Dynamic Brain Imaging* 135–165 (Springer, 2009).
81. Ryan, M. B., Bair-Marshall, C. & Nelson, A. B. Aberrant striatal activity in parkinsonism and
levodopa-induced dyskinesia. *Cell Rep.* **23**, 3438–3446 (2018).
82. Shin, J. H., Kim, D. & Jung, M. W. Differential coding of reward and movement information in the
dorsomedial striatal direct and indirect pathways. *Nat. Commun.* **9**, 1–14 (2018).
83. Shimazaki, H. & Shinomoto, S. A Method for Selecting the Bin Size of a Time Histogram. *Neural*
Comput. **19**, 1503–1527 (2007).
84. Kim, Y.-C. *et al.* Optogenetic stimulation of frontal D1 neurons compensates for impaired temporal
control of action in dopamine-depleted mice. *Curr. Biol.* **27**, 39–47 (2017).
740