Atypical Biological Motion Perception in Children with Attention Deficit Hyperactivity Disorder: Dissociating the Roles of Local Motion and Global Configuration Processing

Junbin Tian[1], Fang Yang[2,3,4], Ying Wang[2,3,4], Li Wang[2,3,4], Ning Wang[1], Yi Jiang[2,3,4]*, Li Yang[1]*

1 Peking University Sixth Hospital, Peking University Institute of Mental Health, National Clinical Research Center for Mental Disorders (Peking University Sixth Hospital), NHC Key Laboratory of Mental Health (Peking University), Beijing, China

2 State Key Laboratory of Brain and Cognitive Science, CAS Center for Excellence in Brain Science and Intelligence Technology, Institute of Psychology, Chinese Academy of Sciences, Beijing, China

3 Department of Psychology, University of Chinese Academy of Sciences, Beijing, China

4 Chinese Institute for Brain Research, Beijing, China

*For correspondence:

Yi Jiang yijiang@psych.ac.cn

Li Yang yangli_pkuimh@bjmu.edu.cn

Abstract

The ability to perceive biological motion (BM) is crucial for human survival and social interaction. Plentiful studies have found impaired BM perception in autism spectrum disorder characterized by deficits in social interaction. Children with attention deficit hyperactivity disorder (ADHD) often exhibit similar deficits in social interaction, but few studies have investigated BM perception in ADHD. Here, we compared the differences in abilities to process local kinematic and global configurational cues, two fundamental abilities of BM perception, between typically developing (TD) and ADHD children. Then, we further investigated the relationship between BM perception and social interaction skills measured by the Social Responsiveness Scale (SRS) and examined the contributions of potential factors (e.g., gender, age, attention and intelligence) to BM perception. Results revealed that children with ADHD exhibited atypical BM perception with a clear dissociation between local and global BM information processing. Local BM processing ability was largely related to social interaction skills, whereas global BM processing ability would significantly improve with age. Critically, general BM perception (i.e., both local and global BM cues) was affected by sustained attention ability in ADHD. This relationship was mainly mediated by Reasoning Intelligence. These findings elucidate the atypical biological motion perception in ADHD and the potential factors related to BM perception. Moreover, this study provides new evidence for local BM perception as a hallmark of social cognition and advances the comprehensive understanding of the distinct roles of local and global processing in BM perception and social cognitive disorders.

Introduction

Attention deficit/hyperactivity disorder (ADHD) is a common developmental disorder. The prevalence of ADHD in children and adolescents is between 2% and 7%, with an average of around 5%[1]. Besides the well-established core symptoms of ADHD (including a deficit of sustained attention, hyperactivity, and impulsivity), some ASD characteristics, such as dysfunctions in social communication and social interaction, have also been frequently reported in ADHD children[2-4]. However, experimental studies of social cognition in ADHD are still scarce. Some studies described worse

performance in tasks of social cognition. Among them, impaired theory of mind

(ToM) and emotion recognition are most frequently reported[5-7]. It is difficult for

children with ADHD to recognize others' emotions and purposes. Other social

cognition processes in ADHD are unclear. Further exploring diversified social

cognition (e.g., biological motion perception) will provide a new understanding of

impaired social function in ADHD. Moreover, recent studies revealed that the social

cognition in ADHD can potentially vary with different factors at the cognitive,

pathological, or developmental levels, such as general cognitive impairment[5], the

severity of symptoms[8], or age[5]. Yet, how these factors relate to the dysfunctions of

social cognition in ADHD is still in its infancy. Bridging the gap is important because

this knowledge can help depict the developmental trajectory of social cognition and

find effective interventions for impaired social interaction in ADHD.

Biological motion (BM), namely the movement of a living creature, carries much

information beyond bodily movements[9], such as intention[10], emotion[11], gender[12], and

identity[13,14]. Since point-light displays (PLD) technology was used to depict human

motions[15], it is possible to separate biological motion from other characteristics like

shape and color. Considering the seminal impact on cognitive science, developmental

and clinical neuroscience, BM perception has drawn significant attention from

scientists. Some researchers attempted to deconstruct BM processing into more

specific components. Our study concentrated on two fundamental abilities to process

BM cues (Figure 1): the abilities to process local BM cues derived from major joint

motion tracks and global BM cues of human configuration. Separable neural signals

revealed two independent processing stages[16-18]. The ability to process local BM cues

is genetically highly preserved[19]; for example, newly hatched chicks without visual

experience exhibit a spontaneous preference for BM[20]. A similar finding is also

reported in 2-day-old babies[21]. Local BM cues not only carry locomotive direction[22]

but also aid in detecting life in the visual environment[23] without observers' explicit

recognition or attention[24-27]. Moreover, the processing of local BM cues is unaffected

by attention, relatively robust to masking noise and does not show a learning trend[27,28].

In contrast, global BM processing involves top-down processing, and attention plays a

critical role in global BM perception[28,29]. Dispersed attention adversely affects its

performance. Compared with local BM processing, global BM processing is

susceptible to learning and is heavily hindered by increasing mask densities[28]. In

recent years, BM perception has received wide attention in studies of mental

disorders[30], particularly in ASD, characterized by deficits in social communication

and social interaction[31,32]. An important reason is that BM perception is considered a

hallmark of social cognition. Individuals with deficits in BM processing exhibit worse

social perception in daily life[10]. Especially in local BM processing, it was correlated

with autistic traits, especially in the subdimension of social communication[19].

Therefore, it is essential to focus on revealing differences in BM processing between

typical and atypical development, which will enhance our comprehension of social

dysfunction in atypical development.

Few studies focused on BM perception in children with ADHD, compared with

numerous studies of impaired BM perception in ASD. An EEG study found

neuro-electrophysiological changes in processing BM stimuli in ADHD[33], whereas no

behavioral difference existed, which could be improved by more carefully designed

BM stimuli. Another study found that children with ADHD performed worse in BM

detection with moderate ratios of noise[34]. Nevertheless, the accuracy of TD groups

was also very low (i.e., poor performance), which indicated the inferior applicability

of this paradigm. In short, factors such as inapplicable experimental designs may

influence the precision and reliability of the results. Despite some preliminary hints,

systematic evaluation of BM perception in ADHD is warranted.

Therefore, the current study aims to examine BM perception abilities in children with

ADHD. The first question to be addressed by this study is whether the BM perception

abilities differ between TD and ADHD children. As mentioned above, BM perception

is related to social interaction and has two fundamental components. Local BM

processing ability is genetically highly preserved, while global BM processing ability

shows a learning trend and needs the involvement of attention. ADHD is a

neurodevelopmental disorder characterized by attention deficit and mostly

accompanied by impaired social interaction. Therefore, the second question is

whether BM perception is associated with social interaction ability in ADHD, and if

so, which factors (e.g., attention) could affect BM perception in ADHD. To answer

these questions, we utilized a stable and ingenious behavioral paradigm used in a

previous study and made minor modifications to fit it to children[19]. This paradigm

combined an intact and a scrambled point-lighted walker with a noise mask to measure local and global BM information processing abilities. Using this paradigm, in Experiment 1, we compared the performance in processing two different BM cues (i.e., local and global BM cues) as well as their combination (i.e., both local and global cues) between TD and ADHD groups. Then, in Experiment 2, we collected the social interaction trait of participants to examine its correlation with the performance of BM tasks. Finally, we investigated the contributing factors of BM perception in ADHD and explored their potential relations.

Materials and methods

Participants

We recruited a total of one hundred and seventeen children with and without ADHD to enter this study. Eighty-one children met the ADHD diagnostic criteria of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5)[35]. The clinical diagnosis was first made by an experienced child and adolescent psychiatrist in the Child and Adolescent Psychiatric Outpatient Department of the Peking University Sixth Hospital, based on the ADHD Rating Scale. The Chinese version of the kiddie-schedule for affective disorders and schizophrenia-present and lifetime version DSM-5 (K-SADS-PL-C DSM-5)[36,37], a semi-structured interview instrument, was then implemented to confirm the diagnosis. Thirty-six typically developing (TD) children were from ordinary primary schools in Beijing and screened out the presence of ADHD, ASD, affective disorders and behavioral disorders by a trained psychiatrist. All subjects of ADHD groups have full-scale IQ above 75 (5th upper percentile) on Wechsler Intelligence Scale for Children-Fourth Edition, and all TD children had a full-scale IQ that fell above the 5th percentile on the Raven's Standard Progressive Matrices[38] which is used in measuring reasoning ability and regarded as a non-verbal estimate of intelligence. Exclusion criteria for both groups included: (a) neurological diseases; (b) other neurodevelopmental disorders (e.g., ASD, Mental retardation, and tic disorders), affective disorders and schizophrenia; (c) that would impact experiments completion; (d) taking psychotropic drugs or stimulants in 30 days. (e) previous head trauma or neurosurgery.

36 TD children (age = 9.09 ± 2.18, 14 male) and 39 children with ADHD (age = 9.88

± 2.23, 28 male) participated in Experiment 1. Groups did not differ by age (t =

-1.550, p = .126) but differed in sex ($\chi2$ = 8.964, p = 0.004). 42 ADHD children (age

= 9.34 ± 1.89, 27 male) participated in Experiment 2. They were naïve to the task and

did not participate in Experiment 1. Participant demographic characteristics are shown

in Table 1. All individuals of each group had a normal or corrected-to-normal vision

and were naïve to the objectives of experiments. Written informed consent was

obtained from the parents of all children before testing. The Institutional Review

Board of the Peking University Sixth Hospital has approved this study.

Assessment

K-SADS-PL-C DSM-5

K-SADS-PL-C DSM-5 is a semi-structured interview instrument for evaluating

mental disorders in children and adolescents aged 6–18 years[36]. It involved thirty-five

diagnoses based on the diagnostic criteria of DSM-5. A trained psychiatrist concluded

the diagnosis by interviewing the parents and child. Chinese version showed great

psychometric criterion[37].

ADHD Rating Scale

ADHD Rating Scale is adapted from the ADHD diagnostic criteria of DSM[39], which

requires parents or teachers to complete the scale independently. Its Chinese version

has an excellent psychometric criterion and consists of 2 subscales[40]: inattention (IA,

nine items) and hyperactivity-impulsivity (HI, nine items). Each item is rated on a

four-point Likert scale ranging from 1 (the symptom is "never or rarely") to 4 (the

symptom is "very often"). The final results will create three scores: (1) IA dimension

score, (2) HI dimension score, and (3) Total score. Higher scores indicate more severe

ADHD symptoms.

Social Responsiveness Scale

Social Responsiveness Scale (SRS) is a widely used quantitative measure with 65

items to assess the severity of social impairment in many mental disorders[41], and the

psychometric properties of the Chinese version are desirable[42]. It includes five

subdimensions: social awareness, social cognition, social communication, social

motivation, and autistic mannerisms. Each item is rated on a scale from 0 (never true)

to 4 (almost always true), with higher scores indicating worse social ability.

Wechsler Intelligence Scale for Children-Fourth Edition

Wechsler Intelligence Scale for Children-Fourth Edition (WISC-IV) is widely used to

test comprehensive intelligence in individuals aged 6–18. It contains fifteen subtests

which comprise four broad areas of intellectual functioning: Verbal Comprehension,

Perceptual Reasoning, Working Memory, and Processing Speed. Scores of the four

broad areas constitute the full-scale intellectual quotient (FIQ).

QB Test

QB Test is a 15-min Continuous Performance Test (CPT) for assessing inattention

and impulsivity with a high-resolution infrared camera monitoring participant's

activity[43]. Previous psychometric studies have validated its good measurement

properties[44]. After the test was completed, several Q scores were calculated to

summarize participant performance. The Q scores were standardized based on

normative data matching gender and age. The higher Q score implies more abnormal

performance. In this study, we focused on the QbInattention, the Q score responsible

for sustained attention, particularly when children focused on tasks.

Stimuli

Stimuli were presented on a 14-inch monitor using MATLAB together with the

PsychToolbox extensions. All subjects completed experiments in a dim-lit room with

heads on the chinrest to ensure their eyes were 50 cm away from the monitor.

Point-light BM stimuli sequences adopted in this study have been used in previous

studies[45], which were derived from configurations of individual walking motion on a

treadmill and did not contain an overall translation. Each frame of BM sequences

consisted of 13 white dots representing the human head and major joints and was

displayed on a grey background. Each walking cycle lasted 1s with 30 frames. For

each trial, the initial frame of BM sequences was randomized. The whole

point-lighted BM walker presented about 5.7 ° visual angle vertically.

Procedure

In Experiment 1, 36 TD children and 39 children with ADHD were required to

complete three tasks that were similar to but slightly modified from the versions

implemented in our previous study[19] (Figure 2). Each trial began with a fixation cross

(0.6° * 0.6°). After a random 1200-1500 ms interval, the monitor displayed a BM

sequence specific to each task, lasting 2 s (60 frames). Task 1 (BM-Local) assessed

the ability to process local BM cues. 13 dots were randomly relocated within the

range of intact BM walker, which rendered global configuration destroyed and local

kinematics retained instead. After the display, we required children to press a left or

right button to indicate the motion direction of the unidentified creature (i.e.,

scrambled BM walker) as accurately as possible. Children received no feedback on

whether a right or wrong answer was given. There were 30 trials in total and 15 trials

for each condition (left and right). Task 2 (BM-Global) tested the ability to process

the global configuration cues of the BM walker. Children needed to judge whether

there was an intact walker in a mask. Each trial contained a mask consisting of two

scrambled target walkers (26 dots) within the boundary about 1.44 times the intact

walker. And the scrambled or intact version of the target walker was randomly

embedded into the mask. This procedure extracted global configuration cues from the

original BM sequence because two conditions (i.e., scrambled and intact walker)

contained the same information of local kinematics as the mask, and the local motion

cues were hence uninformative. The correct choice asked for using the global

component. To avoid learning the shape of the walker[24], we set target walkers

possibly facing one of five equally spaced directions from left to right. There were 30

trials composed of two conditions (intact or scrambled target). In Task 3

(BM-General), the intact target walker facing left or right was embedded into the

mask. The mask is similar to Task 2. Children were required to judge the facing

direction of the target walker (left or right). Because the mask was set to covey the

local BM cues consistent with the target walker, children could perform the task using

general (i.e., local + global) BM information. Task 3 was composed of 30 trials (15

for each facing directions). Other parameters of Task 2 and 3 were similar to Task 1.

Before each task, children practiced five trials to make sure good understanding. We

performed three tasks in a fixed order so that participants were naïve to the scrambled

walker and only made choices using local BM cues in Task 1.

In Experiment 2, 42 children with ADHD completed the same procedures as in

Experiment 1. Beyond that, their parents should fill in the Social Responsiveness

Scale (SRS) to assess social interaction.

Statistics

We carried out ANOVA to examine the difference in BM perception abilities between

TD and ADHD children and the Pearson correlation analysis to assess the relationship

between the accuracy of each task and the SRS score. General linear models and path

analysis were used to explore the roles of potential factors on BM perception. A p

value < 0.05 was considered statistically significant.

Results

Children with ADHD exhibit atypical BM perception

Figure 3 shows the mean accuracies (ACC) for both groups in the three tasks in

Experiment 1. We tested the difference in ACC between the TD and ADHD groups in

each task using one-way ANOVA (Gender as a covariate). In Task 1, the group

comparison revealed a significant difference ($F_{1,72} = 5.630$, $p = 0.020$, $\eta^2 = 0.073$).

Children are asked to judge the facing direction of the target walker using global BM

cues (Task 2) or general BM cues combining global with local BM cues (Task 3). TD

group have higher accuracies than the ADHD group (Task 2: $F_{1,72} = 12.600$, $p <$

0.001, $\eta^2 = 0.149$; Task 3: $F_{1,72} = 19.926$, $p < 0.001$, $\eta^2 = 0.217$). In all statistical

analyses, the main effects of gender were not significant (all $p > 0.05$).

Atypical perception of local BM information predicted impaired social

interaction in ADHD

Experiment 1 demonstrates that the abilities of BM perception were atypical in

ADHD. Our previous study revealed that local BM processing ability is heritable and

negatively correlated with autistic traits, especially in the subdimension of

communication[19]. In addition, substantial evidence indicates that children with ADHD

often have problems in social interaction. We expected that compromised social

interaction in ADHD might be associated with atypical local BM processing. So, we

recruited 42 naïve children with ADHD to participate in Experiment 2 to explore this

relationship. Their parents or caregivers familiar with children completed SRS (total

scores = 54.64 ± 18.42). Thirty-three TD children had SRS scores (total score = 38.64

± 12.47). The total SRS scores in the TD groups are significantly lower than those in

the ADHD group (t = -4.277, p < 0.001). We found that children with higher total

score of SRS performed worse in Task 1, i.e., the ability of local BM processing was

negatively correlated to SRS total scores (r = -0.264, p = 0.022). Notably, significant

correlations were also identified between SRS total score and global BM processing

and between SRS total score and general BM processing (global BM perception: r =

-0.238, p = 0.039; general BM perception: r = -0.359, p = 0.002).

In the further subgroup analysis, as seen in Figure 4, the correlation between the SRS

total score and the ability to process local cues was only found in the ADHD group

(ADHD: r = -0.461, p = 0.002; TD: r = 0.109, p = 0.547), particularly on subscales of

social awareness, social cognition, social communication, social motivation (see

Table 2 for complete information). However, no correlation was found between SRS

scores and global or general BM processing in either ADHD or TD groups (all p >

0.05).

Reasoning intelligence and attention regulated BM processing

Experiment 2 revealed that local BM processing instead of global BM processing was

related to social interaction. Then, which factors would regulate the difference in

processing global BM cues? Previous meta-analysis studies of BM perception in ASD

found mixed results about the effect of age and IQ on processing BM[31,32]. In addition,

neural responses related to BM processing are strongly regulated by attention[29]. Did

these factors play critical roles in BM perception in ADHD, a neurodevelopmental

disorder characterized by attention deficit? To answer this question, we integrated

data from the ADHD group in Experiment 1 and Experiment 2 (n = 80, one child did

not complete QB test). Two linear models were built to investigate contributing

factors: (a) *BM-global = age + gender + FIQ + QbInattention*; (b) *BM-general = age*

+ gender + FIQ + QbInattention + BM-local + BM-global. We screened factors with

the largest contribution to the models by step-wise regression. For model (a), the

ability to process global BM cues was enhanced with age (standardized $\beta = 0.251$, $p = $

0.025). For model (b), higher FIQ, particularly on subdimension Perceptual

Reasoning (standardized $\beta = 0.271$, $p = 0.005$), and more excellent global

BM processing (standardized $\beta = 0.290$, $p = 0.004$) predicted better performance in

general BM processing. Furthermore, as children grew up, the ability to probe general

BM information increased (standardized $\beta = 0.365$, $p < 0.001$). It is worth noting that

QbInattention had a highly negative correlation with Perceptual Reasoning ($r = -0.355$,

$p = 0.001$) and general BM perception ($r = -0.246$, $p = 0.028$). Given the risk of

collinearity, we further used path analysis to visualize these relationships post hoc

(Figure 5). The result showed sustained attention (i.e., QbInattention) did not predict

performance in BM-General (i.e., Task 3) directly. However, a significant indirect

path via the ability of Perceptual Reasoning was found. In addition, as children with

ADHD grew up, their performance in BM-General (i.e., Task 3) not only improved

directly but enhanced via better processing of global BM cues. Moreover, we built

model (c) to corroborate that the local BM processing was stable with age and

unaffected by attention and IQ: (c) *BM-local = age + gender + FIQ + QbInattention*.

No regressor has remained after step-wise regression.

In addition, we further built three models to explore the effects of Reasoning IQ and

age on BM perception in TD children: (a) *BM-local = age + gender + FIQ*. (b)

BM-global = age + gender + FIQ; (c) *BM-general = age + gender + FIQ +*

BM-local +BM-global. No regressor has remained after step-wise regression in model

(a). For models (b) and (c), there are positive relations between age and performance

(BM-global: standardized $\beta = 0.396$, $p = 0.017$; BM-general: standardized $\beta = 0.330$,

$p = 0.049$).

Discussion

Our study contributes several promising findings on atypical biological motion

perception in ADHD. Specifically, we found that children with ADHD performed

worse in processing local, global and general BM information than TD children.

Notably, there was a clear dissociation between local and global BM information

processing. The ability to process local BM cues, which is highly heritable, could

predict deficits of social interaction in ADHD. By contrast, global BM processing,

which is largely shaped by experience, would significantly improve with age.

Moreover, the results also unraveled the potential factors related to BM perception:

Reasoning Intelligence played a mediating effect on the relationship between

attention and general BM perception.

BM perception is a hot topic in the field of visual cognition. One critical reason is that

BM naturally contains biological and social properties. BM processing has significant

value in successful daily life, particularly in nonverbal communication[11,46] and

adaptive behavior[29,47]. As an early emerging ability, BM can be discriminated in the

newborn baby[21]. In typical development children, there is a linkage between BM

perception and social cognitive abilities[48]. For example, 12-month-old infants showed

social behaviors (i.e., following gaze) elicited by BM displays[49]. It is, therefore, of the

supreme importance of BM to children's development of social cognition. This

"social interpretation" of BM proposed that the ability to process BM may be taken as

a measure of difficulties in social interaction**[19]**. An example is that poor performance

of BM perception is a hallmark of impaired social function in children with ASD[10,50].

That implies that atypical BM perception may contribute to the deficits of social

interaction in ADHD. Albeit we found a correlation between the SRS total score and

the accuracy of each of the tasks in all participants, only a significantly negative

correlation between the SRS total score and the accuracy of processing local cues was

found in ADHD group in the further subgroup analysis, which provides preliminary

evidence for the hypothesis and demonstrates that the local BM processing can

predict impaired social interaction in children with ADHD. This finding is consistent

with previous studies that revealed a common genetic basis for local BM processing

ability and social communication[19]. In the subgroup analysis, however, we did not

find a correlation between social interaction and global BM processing, nor between

social interaction and general BM processing. That is because the ability to process

global BM cues is greatly affected by environment and general cognition functions[19].

BM perception is considered a multi-level phenomenon[51-53]. At least in part,

processing information of local BM and global BM are different[16,19]. Our results also

demonstrate such a dissociation. Local motion processing is stable with age and not

susceptible to attention and IQ, which shows a phylogenetically highly preserved

mechanism[20,21,54,55] and benefits of life detection[22]. On the contrary, a development

tendency seems to exist in global BM processing that tends to be improved with age.

A previous study indicated BM perception was enhanced with age in TD children[56].

This variation is also observed in children with ASD, and some developmental

overlaps were found in the ability to process visual motion information[57].

Performance of BM processing within ASD becomes more aligned with TD children,

and the discrepancy between the two groups alleviates with the increase in age[32].

With regard to ADHD, deficits of social cognition are likely to improve with age in

most individuals [5]. Our study further verified this point in BM perception. It is worth

noting that the ability of global BM processing positively correlated with the

performance of processing general BM cues, i.e., children with ADHD are apt to

catch global BM cues to judge the facing direction of the walker when displayed with

local and global BM cues, which correspond to a hierarchical model[51]. Once a living

creature is detected, an agent (i.e., is it a human?) can be recognized by a coherent,

articulated body structure perceptually organized from its motions (i.e., local BM

cues)[58], which is a top-down processing and probably requires attention[28,59],

especially in the presence of competing information[29]. In addition, cortical processing

of BM is strongly modulated by attention[60]. Therefore, as a disorder characterized by

deficits of sustained attention, we found its severity of inattention was negatively

related to the performance of BM processing.

Interestingly, a compensation strategy was found in children with impaired BM

perception. No difficulty of BM recognition was found in ASD individuals with high

IQ[61], but children with ASD showed weaker adaptation effects for biological motion

than TD children[62]. One possibility is that individuals with high IQ and impaired BM

perception can better create or reason reliable strategies used for BM recognition[63,64].

The current study supported this assumption. Children with higher IQ, particularly

Perceptual Reasoning, have better performance. Because of the effect of attention

deficits on Perceptual Reasoning, the performance within ADHD will not become

aligned with that in TD children.

Overall, our study first revealed two atypical fundamental abilities underlying BM

perception in children with ADHD, which showed a clear dissociation. Significantly,

anomalous local BM processing predicts impaired social interaction in ADHD.

Moreover, these results unraveled the potential contributions of age, IQ and attention

to processing BM information. These findings also shed new light on the direction of future studies, which should explore the performance of more advanced BM processing in children with ADHD, such as emotion and identity recognition in BM tasks. In addition, it is necessary to delineate the profiles for neural processing of biological motion in ADHD. A comparative study is warranted to investigate differences between ADHD and ASD, which is essential for finding the common neuropsychological traits and biomarkers of social cognition impairments.

Acknowledgements

This research was supported by grants from the Beijing Municipal Science and Technology Commission (Z181100001518005), the Ministry of Science and Technology of China (2021ZD0203800), the National Natural Science Foundation of China (31830037), the Interdisciplinary Innovation Team (JCTD-2021-06), and Fundamental Research Funds for the Central Universities.

Author contributions

Junbin Tian: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Visualization, Project administration, Writing – original draft. Fang Yang: Formal analysis, Methodology, Investigation. Ying Wang: Methodology, Software, Writing – review & editing. Li Wang: Methodology, Software, Writing – review & editing. Ning Wang: Investigation. Yi Jiang: Conceptualization, Methodology, Supervision, Writing – review & editing. Li Yang: Conceptualization, Methodology, Supervision, Writing – review & editing.

Declaration of interests

The authors declare no competing interest.

Ethics

All procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. The Institutional Review Board of the Peking University Sixth Hospital has approved this study (reference number for approval: (2020) Ethics Review No.9).

Date availability

The data analyzed during the study is available after publication at

https://osf.io/37p5s/

References

1   Sayal, K., Prasad, V., Daley, D., Ford, T. & Coghill, D. ADHD in children and
         young people: prevalence, care pathways, and service provision. *Lancet
         Psychiatry* **5**, 175-186, doi:10.1016/s2215-0366(17)30167-0 (2018).

2   Grzadzinski, R. *et al.* Examining autistic traits in children with ADHD: does
         the autism spectrum extend to ADHD? *J Autism Dev Disord* **41**, 1178-1191,
         doi:10.1007/s10803-010-1135-3 (2011).

3   Mulligan, A. *et al.* Autism symptoms in Attention-Deficit/Hyperactivity
         Disorder: a familial trait which correlates with conduct, oppositional defiant,
         language and motor disorders. *J Autism Dev Disord* **39**, 197-209,
         doi:10.1007/s10803-008-0621-3 (2009).

4   Reiersen, A. M., Constantino, J. N., Volk, H. E. & Todd, R. D. Autistic traits
         in a population-based ADHD twin sample. *J Child Psychol Psychiatry* **48**,
         464-472, doi:10.1111/j.1469-7610.2006.01720.x (2007).

5   Bora, E. & Pantelis, C. Meta-analysis of social cognition in
         attention-deficit/hyperactivity disorder (ADHD): comparison with healthy
         controls and autistic spectrum disorder. *Psychol Med* **46**, 699-716,
         doi:10.1017/s0033291715002573 (2016).

6   Nejati, V. Reading mind from the eyes in individuals with attention
         deficit-hyperactivity disorder (ADHD): A meta-analysis. *Expert Rev
         Neurother* **22**, 889-896, doi:10.1080/14737175.2022.2151899 (2022).

7   Uekermann, J. *et al.* Social cognition in attention-deficit hyperactivity disorder
         (ADHD). *Neurosci Biobehav Rev* **34**, 734-743,
         doi:10.1016/j.neubiorev.2009.10.009 (2010).

8   McKay, E., Kirk, H., Martin, R. & Cornish, K. Social difficulties in adolescent
         attention deficit hyperactivity disorder: Social motivation, social anxiety and
         symptom severity as contributing factors. *J Clin Psychol* **79**, 1113-1129,
         doi:10.1002/jclp.23462 (2023).

9   Dittrich, W. H. Action categories and the perception of biological motion.
Perception **22**, 15-22, doi:10.1068/p220015 (1993).

10   Pavlova, M. A. Biological Motion Processing as a Hallmark of Social
Cognition. *Cerebral Cortex* **22**, 981-995, doi:10.1093/cercor/bhr156 (2011).

11   Clarke, T. J., Bradshaw, M. F., Field, D. T., Hampson, S. E. & Rose, D. The
perception of emotion from body movement in point-light displays of
interpersonal dialogue. *Perception* **34**, 1171-1180, doi:10.1068/p5203 (2005).

12   Kozlowski, L. T. & Cutting, J. E. Recognizing the sex of a walker from a
dynamic point-light display. *Perception & Psychophysics* **21**, 575-580,
doi:10.3758/BF03198740 (1977).

13   Jokisch, D., Daum, I. & Troje, N. F. Self recognition versus recognition of
others by biological motion: viewpoint-dependent effects. *Perception* **35**,
911-920, doi:10.1068/p5540 (2006).

14   Loula, F., Prasad, S., Harber, K. & Shiffrar, M. Recognizing people from their
movement. *J Exp Psychol Hum Percept Perform* **31**, 210-220,
doi:10.1037/0096-1523.31.1.210 (2005).

15   Johansson, G. Visual perception of biological motion and a model for its
analysis. *Perception & Psychophysics* **14**, 201-211, doi:10.3758/BF03212378
(1973).

16   Duarte, J. V., Abreu, R. & Castelo-Branco, M. A two-stage framework for
neural processing of biological motion. *NeuroImage* **259**, 119403,
doi:https://doi.org/10.1016/j.neuroimage.2022.119403 (2022).

17   Jastorff, J. & Orban, G. A. Human functional magnetic resonance imaging
reveals separation and integration of shape and motion cues in biological
motion processing. *J Neurosci* **29**, 7315-7329,
doi:10.1523/jneurosci.4870-08.2009 (2009).

18   Vangeneugden, J., Peelen, M. V., Tadin, D. & Battelli, L. Distinct neural
mechanisms for body form and body motion discriminations. *J Neurosci* **34**,
574-585, doi:10.1523/jneurosci.4032-13.2014 (2014).

19   Wang, Y. *et al.* Heritable aspects of biological motion perception and its
covariation with autistic traits. *Proceedings of the National Academy of*
Sciences **115**, 1937-1942, doi:10.1073/pnas.1714655115 (2018).

20   Vallortigara, G., Regolin, L. & Marconato, F. Visually inexperienced chicks
481   exhibit spontaneous preference for biological motion patterns. *PLoS Biol* **3**,
482   e208, doi:10.1371/journal.pbio.0030208 (2005).
483   21   Simion, F., Regolin, L. & Bulf, H. A predisposition for biological motion in
484   the newborn baby. *Proc Natl Acad Sci U S A* **105**, 809-813,
485   doi:10.1073/pnas.0707021105 (2008).
486   22   Troje, N. F. & Westhoff, C. The inversion effect in biological motion
487   perception: evidence for a "life detector"? *Curr Biol* **16**, 821-824,
488   doi:10.1016/j.cub.2006.03.022 (2006).
489   23   Chang, D. H. & Troje, N. F. Perception of animacy and direction from local
490   biological motion signals. *J Vis* **8**, 3.1-10, doi:10.1167/8.5.3 (2008).
491   24   Chang, D. H. & Troje, N. F. Characterizing global and local mechanisms in
492   biological motion perception. *J Vis* **9**, 8.1-10, doi:10.1167/9.5.8 (2009).
493   25   Bertenthal, B. I. & Pinto, J. Global Processing of Biological Motions.
494   *Psychological Science* **5**, 221-225, doi:10.1111/j.1467-9280.1994.tb00504.x
495   (1994).
496   26   Wang, L., Zhang, K., He, S. & Jiang, Y. Searching for life motion signals.
497   Visual search asymmetry in local but not global biological-motion processing.
498   *Psychol Sci* **21**, 1083-1089, doi:10.1177/0956797610376072 (2010).
499   27   Chang, D. H. F. & Troje, N. F. Acceleration carries the local inversion effect
500   in biological motion perception. *Journal of Vision* **9**, 19-19,
501   doi:10.1167/9.1.19 (2009).
502   28   Thornton, I. M., Rensink, R. A. & Shiffrar, M. Active versus passive
503   processing of biological motion. *Perception* **31**, 837-853, doi:10.1068/p3072
504   (2002).
505   29   Thompson, J. & Parasuraman, R. Attention, biological motion, and action
506   recognition. *Neuroimage* **59**, 4-13, doi:10.1016/j.neuroimage.2011.05.044
507   (2012).
508   30   Kim, J., Norton, D., McBain, R., Ongur, D. & Chen, Y. Deficient biological
509   motion perception in schizophrenia: results from a motion noise paradigm.
510   *Frontiers in Psychology* **4**, doi:10.3389/fpsyg.2013.00391 (2013).

511   31   Federici, A. *et al.* Anomalous Perception of Biological Motion in Autism: A
512   Conceptual Review and Meta-Analysis. *Scientific Reports* **10**, 4576,
513   doi:10.1038/s41598-020-61252-3 (2020).
514   32   Todorova, G. K., Hatton, R. E. M. & Pollick, F. E. Biological motion
515   perception in autism spectrum disorder: a meta-analysis. *Molecular Autism* **10**,
516   49, doi:10.1186/s13229-019-0299-8 (2019).
517   33   Kröger, A. *et al.* Visual Processing of Biological Motion in Children and
518   Adolescents with Attention-Deficit/Hyperactivity Disorder: An Event Related
519   Potential-Study. *PLOS ONE* **9**, e88585, doi:10.1371/journal.pone.0088585
520   (2014).
521   34   Imanipour, S., Sheikh, M., Shayestefar, M. & Baloochnejad, T. Deficits in
522   Working Memory and Theory of Mind May Underlie Difficulties in Social
523   Perception of Children with ADHD. *Neurology Research International* **2021**,
524   3793750, doi:10.1155/2021/3793750 (2021).
525   35   American Psychiatric Association, D. & Association, A. P. *Diagnostic and*
526   *statistical manual of mental disorders: DSM-5*. Vol. 5 (American psychiatric
527   association Washington, DC, 2013).
528   36   Kaufman, J. *et al.* K-SADS-PL DSM-5 November 2016. *Yale University:*
529   *Advanced Center for Intervention and Services Research (ACISR) for Early*
530   *Onset Mood and Anxiety Disorders-Western Psychiatric Institute and Clinic*
531   *Child and Adolescent Research and Education (CARE) Program* (2016).
532   37   Dun, Y. *et al.* Reliability and validity of the Chinese version of the
533   kiddie-schedule for affective disorders and schizophrenia-present and lifetime
534   version DSM-5 (K-SADS-PL-C DSM-5). *Journal of Affective Disorders* **317**,
535   72-78, doi:https://doi.org/10.1016/j.jad.2022.08.062 (2022).
536   38   Pind, J., Gunnarsdóttir, E. K. & Jóhannesson, H. S. Raven's Standard
537   Progressive Matrices: new school age norms and a study of the test's validity.
538   *Personality and Individual Differences* **34**, 375-386 (2003).
539   39   Reid, R. *et al.* Assessing culturally different students for attention deficit
540   hyperactivity disorder using behavior rating scales. *J Abnorm Child Psychol*
541   **26**, 187-198, doi:10.1023/a:1022620217886 (1998).
542   40   Su, Y. E. *et al.* Parent Ratings of ADHD Symptoms in Chinese Urban
543   Schoolchildren: Assessment With the Chinese ADHD Rating Scale-IV: Home

Version. *J Atten Disord* **19**, 1022-1033, doi:10.1177/1087054712461177
(2015).

41 Constantino, J. & Gruber, C. Social responsive scale (SRS) manual. *Los*
Angeles, CA: Western Psychological Services (2005).

42 Cen, C. Q. *et al.* Investigating the validation of the Chinese Mandarin version
of the Social Responsiveness Scale in a Mainland China child population.
BMC Psychiatry **17**, 51, doi:10.1186/s12888-016-1185-y (2017).

43 QbTech, A. QbTest plus technical manual. *Sweden: Gothenburg* (2010).

44 Hult, N., Kadesjö, J., Kadesjö, B., Gillberg, C. & Billstedt, E. ADHD and the
QbTest: Diagnostic Validity of QbTest. *Journal of Attention Disorders* **22**,
1074-1080, doi:10.1177/1087054715595697 (2015).

45 Vanrie, J. & Verfaillie, K. Perception of biological motion: a stimulus set of
human point-light actions. *Behav Res Methods Instrum Comput* **36**, 625-629,
doi:10.3758/bf03206542 (2004).

46 Dittrich, W. H., Troscianko, T., Lea, S. E. & Morgan, D. Perception of
emotion from dynamic point-light displays represented in dance. *Perception*
25, 727-738, doi:10.1068/p250727 (1996).

47 Pollick, F. E., Kay, J. W., Heim, K. & Stringer, R. Gender recognition from
point-light walkers. *J Exp Psychol Hum Percept Perform* **31**, 1247-1265,
doi:10.1037/0096-1523.31.6.1247 (2005).

48 Kutsuki, A., Kuroki, M., Egami, S., Ogura, T. & Itakura, S. Individual
differences in changes in infants' interest in social signals in relation to
developmental index. *Infant Behav Dev* **32**, 381-391,
doi:10.1016/j.infbeh.2009.06.004 (2009).

49 Yoon, J. M. & Johnson, S. C. Biological motion displays elicit social behavior
in 12-month-olds. *Child Dev* **80**, 1069-1075,
doi:10.1111/j.1467-8624.2009.01317.x (2009).

50 Kaiser, M. D. & Pelphrey, K. A. Disrupted action perception in autism:
behavioral evidence, neuroendophenotypes, and diagnostic utility. *Dev Cogn*
Neurosci **2**, 25-35, doi:10.1016/j.dcn.2011.05.005 (2012).

51 Troje, N. F. & Basbaum, A. Biological motion perception. *The senses: A*
comprehensive reference **2**, 231-238 (2008).

52 Troje, N. F. & Chang, D. H. Shape-independent processing of biological
motion. *People watching: Social, perceptual, and neurophysiological studies*
of body perception, 82-100 (2013).

53 Troje, N. F. What is biological motion? Definition, stimuli and paradigms.
Social perception: Detection and interpretation of animacy, agency, and
intention, 13-36 (2013).

54 Di Giorgio, E. *et al.* Filial responses as predisposed and learned preferences:
Early attachment in chicks and babies. *Behav Brain Res* **325**, 90-104,
doi:10.1016/j.bbr.2016.09.018 (2017).

55 Herman, L. M., Morrel-Samuels, P. & Pack, A. A. Bottlenosed dolphin and
human recognition of veridical and degraded video displays of an artificial
gestural language. *J Exp Psychol Gen* **119**, 215-230,
doi:10.1037//0096-3445.119.2.215 (1990).

56 Ghanouni, P. *et al.* Biological Motion Perception Is Affected by Age and
Cognitive Style in Children Aged 8–15. *Neurology Research International*
2015, 594042, doi:10.1155/2015/594042 (2015).

57 Annaz, D. *et al.* Development of motion processing in children with autism.
Dev Sci **13**, 826-838, doi:10.1111/j.1467-7687.2009.00939.x (2010).

58 Rosch, E. *Principles of categorization.* (Morgan Kaufmann, 1988).

59 Cavanagh, P., Labianca, A. T. & Thornton, I. M. Attention-based visual
routines: sprites. *Cognition* **80**, 47-60, doi:10.1016/s0010-0277(00)00153-0
(2001).

60 Safford, A. S., Hussey, E. A., Parasuraman, R. & Thompson, J. C.
Object-based attentional modulation of biological motion processing:
Spatiotemporal dynamics using functional magnetic resonance imaging and
electroencephalography. *Journal of Neuroscience* **30**, 9064-9073,
doi:10.1523/JNEUROSCI.1779-10.2010 (2010).

61 Rutherford, M. D. & Troje, N. F. IQ Predicts Biological Motion Perception in
Autism Spectrum Disorders. *Journal of Autism and Developmental Disorders*
42, 557-565, doi:10.1007/s10803-011-1267-0 (2012).

62 van Boxtel, J. J., Dapretto, M. & Lu, H. Intact recognition, but attenuated
adaptation, for biological motion in youth with autism spectrum disorder.
Autism Res **9**, 1103-1113, doi:10.1002/aur.1595 (2016).

63   Koldewyn, K., Whitney, D. & Rivera, S. M. The psychophysics of visual

motion and global form processing in autism. *Brain* **133**, 599-610,

doi:10.1093/brain/awp272 (2010).

64   Atkinson, A. P. Impaired recognition of emotions from body movements is

associated with elevated motion coherence thresholds in autism spectrum

disorders. *Neuropsychologia* **47**, 3023-3029,

doi:10.1016/j.neuropsychologia.2009.05.019 (2009).

Figures

Figure 1. Schematic representation of Biological Motion (BM) and scrambled

BM sequence. Intact walker contains information of local kinematics and global

configuration. Local kinematics refers to the motion track of each critical joint

illustrated by the chromatic dot. Global configuration is composed of the relative

locations of each joint. In scrambled BM sequence, global configuration cues have

been deprived, but local kinematics have been retained. (Figure reconstructed from

Wang et al., 2018[19])

 Figure 2. Illustration of the trial sequence. In Task 1 (BM-local), a monitor

 displayed scrambled BM sequences. Participants only judged the facing direction of

 the scrambled walker using local BM cues. In Task 2 (BM-global), each trial only

 showed an intact or scrambled walker (black dots in the figure) embedded in a mask

 containing local BM cues. Because the two conditions contained the same local cues

 that were also present in the mask, a participant must rely on global BM cues to judge

 whether there was an intact walker in the mask. The figure showed one of five

 possible facing directions of an intact walker (i.e., facing participants). In Task 3

 (BM-general), the stimuli sequence consisted of an intact walker (black dots) and a

 mask containing similar target local cues, and children judged the facing direction of

 the walker using general BM cues (local + global). Dots in the figure were rendered in

 black for better illustration and displayed in white in the actual experiments.

 Figure 3. The mean accuracy of the three tasks. Typically developing (TD)

 children had higher accuracies than children with attention deficit hyperactivity

 disorder (ADHD) in three tasks in Experiment 1. Error bars show standard errors. *p

 < 0.05; ***$p < 0.001$

Figure 4. Correlations between the accuracy of Task 1 (BM-Local) and SRS total scores. The correlation between the SRS total score and the ability to process local cues is found in ADHD group. By contrast, this correlation is not found in TD group. The grey shading represents a 95% confidence interval. **$p < 0.01$.

Figure 5. Factors influencing BM perception in ADHD children. Post hoc path analysis confirmed that the effect of sustained attention to performance in BM-General was totally mediated by Perceptual Reasoning, and the ability of global BM processing partly mediated the effect of age on performance in BM-General.

Tables

Table 1. Demographic characteristics of TD and ADHD groups

	Experiment 1		p value	Experiment 2		p value
	TD	ADHD		TD	ADHD	

Sample	36	39		33	42	
Age (years)	9.08±2.18	9.88±2.23	0.126	8.75±1.94	9.34±1.89	0.191
Gender ratio (% male)	38.89%	71.79%	0.004	42.42%	64.28%	0.059

662 **Table 2. The correlation between the ability of local BM processing and**
663 **subdimensions of social ability in ADHD children.**

	Correlation coefficient (r)	*p* value
Social awareness	-0.333	0.031
Social cognition,	-0.416	0.006
Social communication	-0.381	0.013
Social motivation	-0.406	0.008
Autistic mannerisms	-0.245	0.117
Total score	-0.461	0.002

664