Atypical Biological Motion Perception in Children with Attention Deficit Hyperactivity Disorder: Local Motion and Global Configuration Processing

Junbin Tian[1], Fang Yang[2,3,4], Ying Wang[2,3,4], Li Wang[2,3,4], Ning Wang[1], Yi Jiang[2,3,4]*, Li Yang[1]*

1 Peking University Sixth Hospital, Peking University Institute of Mental Health, National Clinical Research Center for Mental Disorders (Peking University Sixth Hospital), NHC Key Laboratory of Mental Health (Peking University), Beijing, China

2 State Key Laboratory of Brain and Cognitive Science, CAS Center for Excellence in Brain Science and Intelligence Technology, Institute of Psychology, Chinese Academy of Sciences, Beijing, China

3 Department of Psychology, University of Chinese Academy of Sciences, Beijing, China

4 Chinese Institute for Brain Research, Beijing, China

*For correspondence:

Yi Jiang yijiang@psych.ac.cn

Li Yang yangli_pkuimh@bjmu.edu.cn

Abstract

Perceiving biological motion (BM) is crucial for human survival and social interaction. Many studies have reported impaired BM perception in autism spectrum disorder, which is characterised by deficits in social interaction. Children with attention deficit hyperactivity disorder (ADHD) often exhibit similar difficulties in social interaction. However, few studies have investigated BM perception in children with ADHD. Here, we compared differences in the ability to process local kinematic and global configurational cues, two fundamental abilities of BM perception, between typically developing and ADHD children. We further investigated the relationship between BM perception and social interaction skills measured using the Social Responsiveness Scale and examined the contributions of latent factors (e.g., sex, age, attention, and intelligence) to BM perception. The results revealed that children with ADHD exhibited atypical BM perception. Local and global BM processing showed distinct features. Local BM processing ability was related to social interaction skills, whereas global BM processing ability significantly improved with age. Critically, general BM perception (i.e., both local and global BM processing) may be affected by sustained attentional ability in children with ADHD. This relationship was primarily mediated by reasoning intelligence. These findings elucidate atypical BM perception in ADHD and the latent factors related to BM perception. Moreover, this study provides new evidence that BM perception is a hallmark of social cognition and advances our understanding of the potential roles of local and global processing in BM perception and social cognitive disorders.

Introduction

Attention deficit hyperactivity disorder (ADHD) is a common developmental disorder with a prevalence ranging from 2% to 7% in children and adolescents, averaging approximately 5%[1]. In addition to the well-established core symptoms of ADHD (including inability to sustain attention, hyperactivity, and impulsivity), some characteristics of autism spectrum disorder (ASD), such as dysfunction in social communication and social interaction, have also been frequently observed in children with ADHD[2-4]. Nevertheless, experimental studies focusing on social cognition in children with ADHD are limited. Some studies have reported poor performance on social cognition tasks. Among these, impaired theory of mind (ToM) and emotion

recognition are the most frequently reported[5-7]. It is difficult for children with ADHD

to recognise the emotions and intentions of others. However, our understanding of

other social cognitive processes in ADHD remains limited. Further exploration of a

diverse range of social cognitions (e.g., biological motion perception) can provide a

fresh perspective on the impaired social function observed in ADHD. Moreover,

recent studies have indicated that social cognition in ADHD may vary depending on

different factors at the cognitive, pathological, or developmental levels, such as

general cognitive impairment[5], symptom severity[8], or age[5]. Nevertheless,

understanding how these factors relate to social cognitive dysfunction in ADHD is

still in its infancy. Bridging this gap is crucial as it can help depict the developmental

trajectory of social cognition and identify effective interventions for impaired social

interaction in individuals with ADHD.

Biological motion (BM), which refers to the movement of a living creature, conveys a

wealth of information beyond bodily movements[9], such as intention[10], emotion[11],

sex[12], and identity[13,14]. The advent of point-light display (PLD) technology, which is

used to depict human motions,[15] allows researchers to separate biological motion

from other characteristics such as shape and colour. Considering its seminal impact on

cognitive, developmental, and clinical neuroscience, BM perception has drawn

significant attention from scientists. Some researchers have attempted to deconstruct

BM processing into more specific components. Our study concentrated on two

fundamental abilities involved in processing BM cues (Figure 1): the ability to

process local BM cues derived from major joint motion tracks, and the ability to

process global BM cues of human configuration. Previous studies revealed

differences between local and global BM perception. Separate neural signals for these

abilities imply two independent BM processing stages[16-18]. Local BM cues not only

help identify locomotive direction[19] but also contribute to the detection of life in a

visual environment[20] without the observers' explicit recognition or attention[21-24]. As a

result, the processing of local BM cues is less affected by attention, relatively robust

to masking noise, and does not show a learning trend[24,25]. In contrast, global BM

processing involves top-down modulation, with attention playing a critical role in its

perception[25,26]. Dispersed attention adversely affects performance. Compared with

local BM processing, global BM processing is susceptible to learning and is heavily

hindered by increased mask densities[25]. These findings suggest that local and global

mechanisms play different roles in BM perception, although the exact mechanism

underlying this distinction remains unclear. Exploring these two components of BM

perception will enhance our understanding of the differences between local and global

BM processing and shed light on the psychological processes involved in atypical BM

perception.

In recent years, BM perception has received significant attention in studies on mental

disorders (e.g., schizophrenia[27]) and developmental disabilities, particularly ASD,

which is characterised by deficits in social communication and social interaction[28,29].

This is because BM perception is considered a hallmark of social cognition.

Individuals with deficits in BM processing exhibit worse social perception in daily

life[10]. Another study found that participants' ability to process BM cues correlated

with their autistic traits, particularly in the subdimension of social communication[30].

Therefore, examining BM perception could enhance our understanding of social

dysfunction in children with ADHD. Compared with the numerous studies examining

impaired BM perception in ASD, few studies have focused on BM perception in

children with ADHD. An EEG study found neuroelectrophysiological changes in the

processing of BM stimuli in children with ADHD[31]. Specifically, compared with the

typically developing (TD) group, children with ADHD showed reduced activity of

motion-sensitive components (N200) while watching biological and scrambled

motions, although no behavioural differences were observed. Another study found

that children with ADHD performed worse in BM detection with moderate noise

ratios than the TD group[32]. This finding may be due to the fact that BM stimuli with

noise dots will increase the difficulty of identification[33], which highlights the

difference in BM processing between TD and ADHD groups[31].

Despite initial findings about atypical BM perception in ADHD, previous studies on

ADHD treated BM perception as a single entity, which may have led to misleading or

inconsistent findings[28]. Hence, it is essential to deconstruct BM processing into

multiple components and motion features. To enhance our understanding of the

ability to process distinct BM cues in ADHD, we employed a carefully designed

behavioural paradigm, as used in our previous study[30], with slight adjustments made

to adapt it for children. This paradigm comprised three tasks: BM-local, BM-global,

and BM-general. BM-local assessed the ability to process local BM cues. Scrambled

BM sequences were displayed and the participants used local BM cues to judge the

direction the scrambled walker was facing. BM-global tested the ability to process the

global configuration cues of the BM walker. Local cues were uninformative, and the

participants used global BM cues to determine the presence of an intact walker. BM-

general tested participants' ability to process general BM cues (local + global cues).

The stimulus sequences consisted of an intact walker and a mask containing similar

target local cues so that the participants could use general BM cues to judge the

direction the walker was facing.

Experiment 1 examined three specific BM perception abilities in children with ADHD.

Children with ADHD show impaired social interaction[2-4], which implies atypical

social cognition. Therefore, we speculated that children with ADHD would perform

worse on the three tasks than TD children. In Experiment 2, we further explored the

relationship between BM perception and social interaction ability in children with

ADHD and identified potential factors (e.g., intelligence quotient [IQ], age, and

attention) that may affect BM perception in this population. We speculated that if the

mechanisms of processing local and global BM cues are indeed distinct, as suggested

by previous studies, then impairment in the ADHD population and the influential

factors behind the impairment may be different.

Materials and methods

Participants

One hundred seventeen children with and without ADHD were recruited for this

study. Eighty-one children met the ADHD diagnostic criteria of the Diagnostic and

Statistical Manual of Mental Disorders (DSM-5)[34]. The clinical diagnosis was first

made by an experienced child and adolescent psychiatrist in the Child and Adolescent

Psychiatric Outpatient Department of Peking University Sixth Hospital, based on the

ADHD Rating Scale. The Chinese version of the Kiddie-Schedule for Affective

Disorders and Schizophrenia-Present and Lifetime Version DSM-5 (K-SADS-PL-C

DSM-5)[35,36], a semi-structured interview instrument, was then implemented to

confirm the diagnosis. Thirty-six TD children from ordinary primary schools in

Beijing were screened for the presence of ADHD, ASD, affective disorders, and

behavioural disorders by a trained psychiatrist. All participants in the ADHD group

had a full-scale IQ >75 (5th upper percentile) on the Wechsler Intelligence Scale for

Children-Fourth Edition, and all TD children had a full-scale IQ above the 5th

percentile on Raven's Standard Progressive Matrices[37], which is used to measure

reasoning ability and is regarded as a non-verbal estimate of intelligence. The

exclusion criteria for both groups were as follows: (a) neurological diseases; (b) other

neurodevelopmental disorders (e.g., ASD, mental retardation, and tic disorders),

affective disorders, and schizophrenia; (c) disorders that would impact the completion

of the experiment; (d) taking psychotropic drugs or stimulants within the past 30 days;

and (e) previous head trauma or neurosurgery.

Thirty-six TD children (age = 9.09 ± 2.18, 14 male) and 39 children with ADHD (age

= 9.88 ± 2.23, 28 male) participated in Experiment 1. The groups did not differ by age

($t = -1.550$, $p = 0.126$) but differed in sex ($\chi2 = 8.964$, $p = 0.004$). Forty-two ADHD

children (age = 9.34 ± 1.89, 27 male) participated in Experiment 2. The participants

did not participate in Experiment 1. The participants' demographic characteristics are

presented in Table 1. Currently, there is no comparable study on ADHD that indicates

effect size as a reference. Studies investigating BM perception in children with ASD

typically have sample sizes ranging from 15 to 35 participants per group[29].

Considering the mild impairment of social function in children with ADHD, we

determined that a sample size of 35 to 40 participants per group was reasonable for

this study. All individuals in each group had normal or corrected-to-normal vision and

were naïve to the experimental objectives. Written informed consent was obtained

from the parents of all the children before testing. This study was approved by the

Institutional Review Boards of Peking University Sixth Hospital and the Institute of

Psychology, Chinese Academy of Sciences.

Assessment

K-SADS-PL-C DSM-5

The K-SADS-PL-C DSM-5 is a semi-structured interview instrument used to evaluate

mental disorders in children and adolescents aged 6–18 years[35]. It involves 35

diagnoses based on the diagnostic criteria of the DSM-5. A trained psychiatrist

confirmed the diagnosis by interviewing the parents and children. The Chinese

version has demonstrated great psychometric properties[36].

ADHD Rating Scale

The ADHD Rating Scale was adapted from the ADHD diagnostic criteria of the DSM[38], which requires parents or teachers to complete the scale independently. Its Chinese version has excellent psychometric properties and consists of 2 subscales[39]: inattention (IA, nine items) and hyperactivity-impulsivity (HI, nine items). Each item is rated on a four-point Likert scale ranging from 1 (the symptom appears "never or rarely") to 4 (the symptom appears "very often"). The final results create three scores: (1) IA dimension score, (2) HI dimension score, and (3) total score. Higher scores indicate more severe ADHD symptoms.

Social Responsiveness Scale

The Social Responsiveness Scale (SRS) is a widely used quantitative measure with 65 items used to assess the severity of social impairment in many mental disorders[40], and the psychometric properties of the Chinese version are reliable[41]. It includes five sub-dimensions: social awareness, social cognition, social communication, social motivation, and autistic mannerisms. Each item is rated on a scale from 0 (never true) to 3 (almost always true), with higher scores indicating worse social ability.

Wechsler Intelligence Scale for Children-Fourth Edition

The Wechsler Intelligence Scale for Children-Fourth Edition (WISC-IV) is widely used to test comprehensive intelligence in individuals aged 6–17 years. It contains 15 subtests comprising four broad areas of intellectual functioning: Verbal Comprehension, Perceptual Reasoning, Working Memory, and Processing Speed. The scores in the four broad areas constitute the full-scale intellectual quotient (FIQ).

QB Test

The QB Test is a 15-minute Continuous Performance Test (CPT) for assessing inattention and impulsivity, with a high-resolution infrared camera monitoring participant's activity[42]. Previous psychometric studies have validated its good measurement properties[43]. After the test is completed, several Q scores are calculated to summarise the participants' performances. The Q scores are standardised based on normative data matched for sex and age. A higher Q score implies more abnormal performance. In this study, we focused on QbInattention, the Q score indicating sustained attention, particularly when children are focused on tasks.

	Stimuli and Procedure

Point-light BM stimuli sequences adopted in this study have been used in previous

studies[44], which were derived from the configurations of individual walking motions

on a treadmill and did not contain an overall translation. Each frame of BM sequences

consisted of 13 white dots representing the human head and major joints and was

displayed on a grey background (see Video 1). Each walking cycle lasted 1 s with 30

frames. For each trial, the initial frame of BM sequences was randomised. The entire

point-lighted BM walker was presented at approximately a 5.7° vertical visual angle.

Stimuli were presented on a 14-inch monitor, and responses were evaluated using

MATLAB together with PsychToolbox extensions. All subjects completed the

experiments in a dimly lit room with their heads on a chinrest to ensure that their eyes

were 50 cm away from the monitor.

In Experiment 1, the children were required to complete three tasks that were similar

to but slightly modified from the versions implemented in our previous study[30]

(Figure 2). Each trial began with a fixation cross (0.6° * 0.6°). Following a random

interval of 1200-1500 ms, the monitor displayed a task-specific BM sequence lasting

for 2 s (60 frames).

BM-Local assessed participants' ability to process local BM cues. During the task, the

monitor displayed only a scrambled walker facing either the left or right (Video 2).

Specifically, the 13 dots constituting the intact walker were randomly relocated within

the original range of the BM walker (randomly presented in 2D). This manipulation

disrupted the global configuration of the intact walker while retaining local

kinematics. After the display, we required the children to press a left or right button to

indicate the direction of motion of the unidentified creature (i.e., the scrambled BM

walker) as accurately as possible. Children did not receive feedback on the accuracy

of each response. Thirty trials were conducted, with 15 trials for each condition (left

and right).

BM-Global tested the ability to process the global configuration cues of the BM

walker. A target walker (scrambled or intact) was displayed within the mask (Video 3)

during this task. The mask consisted of two scrambled target walkers (26 dots) with

the same locomotion direction as the target walker, displayed within a boundary

approximately 1.44 times larger than the intact walker. A scrambled or intact version

of the target walker was randomly embedded in the mask and entirely overlaid. Thus,

the global BM component could be isolated as two conditions (i.e., scrambled and

intact walkers) containing the same local kinematics information, rendering the local

motion cues uninformative. The children were required to judge whether there was an

intact walker on the mask. A correct response relied on the extraction of global cues

from an intact walker. To prevent children from learning the shape of the walker[21], we

set target walkers that possibly faced one of five equally spaced directions from left to

right. Of the five walkers used, two faced straight to the left or right, orthogonal to the

viewing direction. Two walked with their bodies oriented at a 45-degree angle to the

left or right of the observer. The last one walked towards the observer. Video 4 shows

the five facing directions of the walker. Thirty trials were conducted consisting of two

conditions (intact or scrambled target).

BM-General tested participants' ability to process general BM cues (local and global).

In BM-General, the monitor displayed an intact walker (facing either the left or right)

embedded within a mask (see Video 5). The mask used in this task was similar to that

used in BM-Global. The children were required to judge the direction the target

walker was facing (left or right). Because the mask and target walker contained the

same local BM cues and the target walker was presented with additional global

configuration cues, children could rely on general BM information (i.e., a

combination of local and global cues) to perform the task. BM-General consisted of

30 trials, with 15 trials for each facing direction. The other parameters of BM-Global

and BM-General were similar to those of BM-Local. Before each task, the children

practiced for five trials to ensure a good understanding. We performed the three tasks

in a fixed order so that the participants were naïve to the nature of the local BM cues

in BM-Local.

In Experiment 2, 42 children with ADHD completed the same procedure as in

Experiment 1. In addition, the parents completed the SRS to assess social interaction.

Statistics

Two-sample t-tests were used to examine the difference in BM perception abilities

between TD and ADHD children[45,46], and Pearson's correlation analyses were used to

assess the relationship between the accuracy of each task and the SRS score.

Additionally, general linear models and path analyses were used to explore potential

factors influencing BM perception. A p-value < 0.05 was considered statistically

significant. Path analyses were conducted using AMOS, whereas other analyses were

conducted using SPSS.

Results

Children with ADHD exhibit atypical BM perception

Figure 3 displays the mean accuracies (ACC) for both the TD and ADHD groups

across the three tasks in Experiment 1. We examined the difference in the ACC

between the TD and ADHD groups for each task using a two-sample t-test. The

results of BM-Local showed a significant difference (TD: 0.52 ± 0.13, ADHD: $0.44 \pm$

0.09, $t_{73} = 3.059$, $p = 0.003$, Cohen's $d = 0.716$), indicating that children with ADHD

exhibited impaired local BM processing ability. For BM-Global and BM-General,

where children were asked to detect the presence or discriminate the direction the

target walker was facing, the TD group had higher accuracies than the ADHD group

(BM-Global - TD: 0.70 ± 0.12, ADHD: 0.59 ± 0.12, $t_{73} = 3.677$, $p < 0.001$, Cohen's d

$= 0.861$; BM-General - TD: 0.79 ± 0.12, ADHD: 0.63 ± 0.17, $t_{73} = 4.702$, $p < 0.001$,

Cohen's $d = 1.100$). These findings suggest the presence of impaired global and

general BM perception in children with ADHD. To ensure that sex did not influence

the results, we conducted a subsampling analysis with balanced data, and the results

remained consistent (see Supplementary Information).

Atypical perception of local BM information predicts impaired social interaction

in ADHD

Experiment 1 provides evidence of atypical BM perception in children with ADHD.

Previous studies have revealed that BM processing ability is a hallmark of social

cognition[10] and is negatively correlated with social ability[30]. Substantial evidence

indicates that children with ADHD often experience problems with social interaction.

We hypothesised that compromised social interaction in children with ADHD would

be associated with BM processing. To confirm this hypothesis, we recruited 42 naïve

children with ADHD to participate in Experiment 2 and examined the relationship

between their social interaction abilities and BM perception. Parents or caregivers

completed the SRS. The SRS total score of the ADHD group was higher than that of

the TD group (SRS total score - ADHD: 54.64 ± 18.42, TD: 38.64 ± 12.47, $t = -4.277$,

$p < 0.001$). We found that children with higher total SRS scores performed worse on

the three tasks; that is, the abilities of BM processing were negatively correlated with

SRS total score (BM-Local: $r = -0.264$, false discovery rate [FDR]-corrected $p =$

0.033; BM-Global: $r = -0.238$, FDR-corrected $p = 0.039$; BM-General: $r = -0.359$,

FDR-corrected $p = 0.006$).

The correlations encompassing all data from both groups might reflect group

disparities, given the significant distinction in SRS total score between TD and

ADHD children, alongside their marked differences in BM processing abilities.

Therefore, we conducted additional subgroup analysis to further explore the

relationship between social interaction and BM processing ability in children with

ADHD. As depicted in Figure 4, the correlation between the SRS total score and the

ability to process local cues was only found in the ADHD group (ADHD: $r = -0.461$,

FDR-corrected $p = 0.004$; TD: $r = 0.109$, FDR-corrected $p = 0.547$), particularly on

subscales of social awareness, social cognition, social communication, social

motivation (see Table 2 for detailed information). However, we did not find a

statistically significant correlation between the SRS total score and global or general

BM processing in either the ADHD or TD groups (global BM perception, TD: $r = -$

0.020, FDR-corrected $p = 0.910$, ADHD: $r = -0.207$, FDR-corrected $p = 0.374$;

general BM perception, TD: $r = -0.118$, FDR-corrected $p = 0.514$, ADHD: $r = -0.286$,

FDR-corrected $p = 0.134$).

To determine the specificity of the correlation between local BM processing and SRS

total score in the ADHD group, we constructed general linear models to further

compare these correlations (see Supplementary Information). We observed a

significantly stronger correlation between SRS total score and response accuracy in

the ADHD group compare to the TD group ($p = 0.003$) for the BM-Local, but not for

the BM-Global ($p = 0.381$) or BM-General ($p = 0.455$). Additionally, our results

showed trends towards significance, indicating that the correlation between the SRS

total score and response accuracy of the ADHD group in BM-Local was more

negative than that in BM-Global ($p = 0.074$) or in BM-General ($p = 0.073$). These

findings suggest that the atypical local BM processing ability may be specifically

related to the impairment of social interaction in children with ADHD.

Global BM processing develops with age and is regulated by reasoning intelligence and attention function

Many factors can affect social cognition in ADHD, such as general cognitive impairment[5], symptom severity[8], and age[5]. To better understand their role in atypical BM processing, we examined the relationship between BM task performance and factors such as age, IQ, and attention function in the ADHD group. Data from the ADHD group in both Experiments 1 and 2 were integrated, resulting in 80 ADHD participants (one child did not complete the QB test). Three linear models were built to investigate the contributing factors: (a) $ACC_{BM\text{-}Local} = \beta_0 + \beta_1 * age + \beta_2 * gender + \beta_3 * FIQ + \beta_4 * QbInattention$, (b) $ACC_{BM\text{-}Global} = \beta_0 + \beta_1 * age + \beta_2 * gender + \beta_3 * FIQ + \beta_4 * QbInattention$, and (c) $ACC_{BM\text{-}General} = \beta_0 + \beta_1 * age + \beta_2 * gender + \beta_3 * FIQ + \beta_4 * QbInattention + \beta_5 * ACC_{BM\text{-}Local} + \beta_6 * ACC_{BM\text{-}Global}$. $ACC_{BM\text{-}Local}$, $ACC_{BM\text{-}Global}$ and $ACC_{BM\text{-}General}$ refer to the response accuracies of the three tasks in the ADHD group, and *QbInattention* is the standardised score for sustained attention function. We screened the factors with the largest contribution to the models using stepwise regression. In model (a), no variable remained after stepwise regression, suggesting that local BM processing remained stable with age and was not affected by attention or IQ. In model (b), the ability to process global BM cues was enhanced with age (standardised $\beta_1 = 0.251$, $p = 0.025$). In model (c), higher FIQ, particularly on the subdimension of Perceptual Reasoning (standardised $\beta_3 = 0.271$, $p = 0.005$), and better performance in global BM processing (standardised $\beta_6 = 0.290$, $p = 0.004$) predicted better performance in general BM processing. Furthermore, as children aged, the ability to probe general BM information improved (standardised $\beta_1 = 0.365$, $p < 0.001$). It is worth noting that QbInattention showed a strong negative correlation with Perceptual Reasoning ($r = -0.355$, $p = 0.001$) and general BM perception ($r = -0.246$, $p = 0.028$). Owing to the potential collinearity issue, we employed a post hoc path analysis to visualise these relationships (Figure 5). The results indicated that sustained attention (i.e., QbInattention) did not directly predict performance in BM-General but was significantly indirectly predicted by Perceptual Reasoning ability. Furthermore, as children with ADHD aged, their performance in BM-General improved, both directly and through the enhanced processing of global BM cues.

We also built three models to explore further the effects of Reasoning IQ and age on

BM perception in TD children: (d) $ACC_{BM\text{-}Local} = \beta_0 + \beta_1 * age + \beta_2 * gender + \beta_3 *$

FIQ. (e) $ACC_{BM\text{-}Global} = \beta_0 + \beta_1 * age + \beta_2 * gender + \beta_3 * FIQ$; (f) $ACC_{BM\text{-}General} = \beta_0$

$+ \beta_1 * age + \beta_2 * gender + \beta_3 * FIQ + \beta_4 * ACC_{BM\text{-}Local} + \beta_5 * ACC_{BM\text{-}Global}$. In model

(d), no regressor remained significant after stepwise regression. However, in models

(e) and (f), we observed positive relationships between age and performance (model e:

standardized $\beta_1 = 0.396$, $p = 0.017$; model f: standardised $\beta_1 = 0.330$, $p = 0.049$). We

also conducted a path analysis similar to that in the ADHD group and found no

statistically significant mediator effect (Figure S3). The complete information about

models a-f could be found in Table S1.

In summary, our findings suggest that the ability to perceive global and general BM

cues, rather than local BM cues, improves with age in both groups. We speculated

that age-related improvement was different when processing different BM cues and in

different groups. Therefore, we examined the differences between the age-related

improvement in processing local cues and that in processing global and general cues

(see Supplementary Information for detailed methods). In the ADHD group, we

observed that the ability to process general BM cues significantly improved with age

compared to local cues ($p < 0.001$) and a trend that the improvement in processing

global BM cues with age was greater than that in processing local BM cues ($p = 0$.

073). However, these patterns were not observed in the TD group (see Supplementary

Information). In addition, we examined the differences in the improvements in

processing of BM cues with age between the two groups for each task. There was no

difference in the effect of age on the response accuracy between the TD and ADHD

groups for the three tasks (see Supplementary Information).

Discussion

Our study contributes several promising findings concerning atypical BM perception

in children with ADHD. Specifically, we observed atypical local and global BM

perception in children with ADHD. Notably, local and global BM processing

exhibited distinct features. The ability to process local BM cues appears to be

associated with social interaction traits in children with ADHD. In contrast, global

BM processing was associated with age-related development. In addition, general BM

perception may be affected by factors such as attention.

BM perception is a widely studied topic in visual cognition owing to its inherent

biological and social properties. BM processing has significant value in successfully

navigating daily life, particularly in non-verbal communication[11,47] and adaptive

behavior[26,48]. In TD children, there is a clear association between BM perception and

social cognitive abilities[49]. For example, 12-month-old infants exhibit social

behaviours (i.e., following gaze) elicited by BM displays[50]. Therefore, BM perception

plays a crucial role in the development of children's social cognition. This 'social

interpretation' of BM suggests that the difficulties in processing BM may serve as an

indicator of impaired social interaction**[30]**. Our results are consistent with these

findings. We observed atypical BM perception in children with ADHD and a

significant relationship between BM perception performance and the SRS total score.

Further subgroup analysis revealed a significant negative correlation between SRS

total score and the accuracy of local BM processing in the ADHD group. This

correlation was stronger in the ADHD group than in the TD group. The lack of a

significant correlation may be due to the narrow range of SRS scores in the TD group.

Future studies should increase the sample size to explore the correlations among

diverse individuals. These findings suggest that BM processing is a distinct hallmark

of social cognition in ADHD children[10,30].

BM perception is a multi-level phenomenon[51-53]. At least in part, the processing of

local and global BM information appears to involve different neural mechanisms[16].

Sensitivity to local BM cues emerges early in life[54,55] and involves rapid processing in

the subcortical regions[16,56-58]. As a basic pre-attentive feature[23], local BM cues can

guide visual attention spontaneously[59,60]. In contrast, the ability to process global BM

cues is related to slow cortical BM processing and is influenced by many factors such

as attention[25,26] and visual experience[21,51]. As mentioned above, we found a

significant negative correlation between the SRS total score and the accuracy of local

BM processing, specifically in the ADHD group. This could be due to decreased

visual input related to atypical local BM processing, which further impairs global BM

processing. According to the two-process theory of biological motion processing[61],

local BM cues guide visual attention toward BM stimuli[55,62]. Consequently, the visual

input of BM stimuli increases, facilitating the development of the ability to process

global BM cues through learning[21,63]. The latter is a prerequisite for attributing

intentions to others and facilitating social interactions with other individuals[20,64,65].

Thus, atypical local BM processing may contribute to impaired social interactions

through altered visual input. Further empirical studies are required to confirm these

hypotheses.

The ability to process global BM cues develops with age. Previous studies indicated

that global BM perception is enhanced with age in TD children[66,67]. This

developmental trend in global BM processing is also evident in individuals with

impaired BM perception (e.g., children with ASD). BM processing performance in

children with ASD becomes more aligned with that of TD children as they age[29,68-70].

Our study contributes new evidence to the understanding of the development of

global BM processing. We found that the ability to process global and general BM

cues improved significantly with age in both the TD and ADHD groups, implying that

the processing module for global BM cues tends to mature with development. This

finding is akin to the potential age-related improvements observed in certain aspects

of social cognitive deficits in individuals with ADHD[5]. Interestingly, in the ADHD

group, the improvement in processing general and global BM cues was greater than in

processing local BM cues. Few developmental studies have been conducted on local

BM processing. The ability to process local BM cues remained stable and did not

exhibit a learning trend[21,25]. A reasonable interpretation may be that local BM

processing is a low-level mechanism, probably performed by the primary visual

cortex and subcortical regions such as the superior colliculus, pulvinar, and ventral

lateral nucleus[14,56,61], which are not malleable. Although there was no statistical

difference between the improvements in processing local and global BM cues in the

TD group, this may be due to the relatively small sample size or the relatively higher

baseline abilities of BM perception in TD children, resulting in a relatively milder

improvement. It is worth noting that the ability to process global BM cues was

positively correlated with the performance in processing general BM cues in the

ADHD group, whereas no such correlation was found in the TD group. This suggests

that TD children are able to extract and integrate both local and global cues, whereas

children with ADHD may rely more on global BM cues to judge the direction the

walker is facing when presented with both local and global BM cues, which

correspond to a hierarchical model[51]. Once a living creature is detected, an agent (i.e.,

is it a human?) can be recognised by a coherent, articulated body structure that is

perceptually organised based on its motions (i.e., local BM cues)[71]. This involves top-

down processing and probably requires attention[25,72], particularly in the presence of

competing information[26]. Our findings are consistent with those of previous studies

on the cortical processing of BM[73], as we found that the severity of inattention in

children with ADHD was negatively correlated with their performance in global BM

processing, whereas this significant correlation was not found in local BM processing,

which may involve bottom-up processing[61,65] and might not need participants' explicit

attention[21,23,74,75]. However, further studies are needed to verify this hypothesis.

Interestingly, children with impaired BM perception may employ a compensatory

strategy[76]. Previous studies found no impairment in BM recognition in individuals

with autism with high IQ[76], but children with ASD exhibited weaker adaptation

effects for biological motion than TD children[77]. One possibility is that individuals

with a high IQ and impaired BM perception can develop or employ reliable strategies

for BM recognition, compensating for the lack of intuitive social perceptual

processing[78,79]. The current study supports this assumption, as children with a higher

IQ, particularly in Perceptual Reasoning, demonstrated better performance. Owing to

the impact of attention deficits on Perceptual Reasoning, the performance of children

with ADHD did not align with that of TD children.

Overall, our study reveals two distinct and atypical fundamental abilities underlying

BM perception in children with ADHD. Notably, anomalous local BM processing

may predict impaired social interactions in children with ADHD. Moreover, these

results revealed the potential contributions of age, IQ, and attention to BM

information processing. These findings also shed new light for future studies. First,

different developmental trends appear in local and global BM processing. Further

studies are required to explore the relationship between the two fundamental BM

processing abilities, which will contribute to understanding the respective and mutual

neural mechanisms underlying the two types of BM processing. Second, exploring the

performance of more advanced BM processing in children with ADHD, such as

emotion and identity recognition in BM tasks, is necessary to delineate the neural

profiles involved in processing BM in ADHD. Finally, a comparative study between

ADHD and ASD is warranted to identify common neuropsychological traits and
biomarkers of social cognition impairment.

Acknowledgements

We thank the three reviewers for their constructive comments, and Dr. Shuo Gao and
Dr. Li Shen for assistance with manuscript revision. Special thanks to the parents and
the children who took part in the study. This research was supported by grants from
the Beijing Municipal Science and Technology Commission (Z181100001518005),
the Ministry of Science and Technology of China (2021ZD0203800), the National
Natural Science Foundation of China (31830037), the Interdisciplinary Innovation
Team (JCTD-2021-06), and Fundamental Research Funds for the Central Universities.

Author contributions

Junbin Tian: Conceptualization, Data curation, Formal analysis, Investigation,
Methodology, Visualization, Project administration, Writing – original draft. Fang
Yang: Formal analysis, Methodology, Investigation. Ying Wang: Methodology,
Software, Writing – review & editing. Li Wang: Methodology, Software, Writing –
review & editing. Ning Wang: Investigation. Yi Jiang: Conceptualization,
Methodology, Supervision, Writing – review & editing. Li Yang: Conceptualization,
Methodology, Supervision, Writing – review & editing.

Declaration of interests

512 The authors declare no competing interest.

Ethics

All procedures contributing to this work comply with the ethical standards of the
relevant national and institutional committees on human experimentation and with the
Helsinki Declaration of 1975, as revised in 2008. The institutional review boards of
the Peking University Sixth Hospital and the Institute of Psychology, Chinese
Academy of Sciences have approved this study (reference number for approval: (2020)
Ethics Review No.9 and H23030).

Data availability

521 The data analyzed during the study is available at https://osf.io/37p5s/

References

1   Sayal, K., Prasad, V., Daley, D., Ford, T. & Coghill, D. ADHD in children and
young people: prevalence, care pathways, and service provision. *Lancet*
Psychiatry **5**, 175-186, doi:10.1016/s2215-0366(17)30167-0 (2018).

2   Grzadzinski, R. *et al.* Examining autistic traits in children with ADHD: does
the autism spectrum extend to ADHD? *J Autism Dev Disord* **41**, 1178-1191,
doi:10.1007/s10803-010-1135-3 (2011).

3   Mulligan, A. *et al.* Autism symptoms in Attention-Deficit/Hyperactivity
Disorder: a familial trait which correlates with conduct, oppositional defiant,
language and motor disorders. *J Autism Dev Disord* **39**, 197-209,
doi:10.1007/s10803-008-0621-3 (2009).

4   Reiersen, A. M., Constantino, J. N., Volk, H. E. & Todd, R. D. Autistic traits
in a population-based ADHD twin sample. *J Child Psychol Psychiatry* **48**,
464-472, doi:10.1111/j.1469-7610.2006.01720.x (2007).

5   Bora, E. & Pantelis, C. Meta-analysis of social cognition in attention-
deficit/hyperactivity disorder (ADHD): comparison with healthy controls and
autistic spectrum disorder. *Psychol Med* **46**, 699-716,
doi:10.1017/s0033291715002573 (2016).

6   Nejati, V. Reading mind from the eyes in individuals with attention deficit-
hyperactivity disorder (ADHD): A meta-analysis. *Expert Rev Neurother* **22**,
889-896, doi:10.1080/14737175.2022.2151899 (2022).

7   Uekermann, J. *et al.* Social cognition in attention-deficit hyperactivity disorder
(ADHD). *Neurosci Biobehav Rev* **34**, 734-743,
doi:10.1016/j.neubiorev.2009.10.009 (2010).

8   McKay, E., Kirk, H., Martin, R. & Cornish, K. Social difficulties in adolescent
attention deficit hyperactivity disorder: Social motivation, social anxiety and
symptom severity as contributing factors. *J Clin Psychol* **79**, 1113-1129,
doi:10.1002/jclp.23462 (2023).

9   Dittrich, W. H. Action categories and the perception of biological motion.
Perception **22**, 15-22, doi:10.1068/p220015 (1993).

10   Pavlova, M. A. Biological Motion Processing as a Hallmark of Social
Cognition. *Cerebral Cortex* **22**, 981-995, doi:10.1093/cercor/bhr156 (2011).

11   Clarke, T. J., Bradshaw, M. F., Field, D. T., Hampson, S. E. & Rose, D. The
555   perception of emotion from body movement in point-light displays of
556   interpersonal dialogue. *Perception* **34**, 1171-1180, doi:10.1068/p5203 (2005).

557   12   Kozlowski, L. T. & Cutting, J. E. Recognizing the sex of a walker from a
558   dynamic point-light display. *Perception & Psychophysics* **21**, 575-580,
559   doi:10.3758/BF03198740 (1977).

560   13   Jokisch, D., Daum, I. & Troje, N. F. Self recognition versus recognition of
561   others by biological motion: viewpoint-dependent effects. *Perception* **35**, 911-
562   920, doi:10.1068/p5540 (2006).

563   14   Loula, F., Prasad, S., Harber, K. & Shiffrar, M. Recognizing people from their
564   movement. *J Exp Psychol Hum Percept Perform* **31**, 210-220,
565   doi:10.1037/0096-1523.31.1.210 (2005).

566   15   Johansson, G. Visual perception of biological motion and a model for its
567   analysis. *Perception & Psychophysics* **14**, 201-211, doi:10.3758/BF03212378
568   (1973).

569   16   Duarte, J. V., Abreu, R. & Castelo-Branco, M. A two-stage framework for
570   neural processing of biological motion. *NeuroImage* **259**, 119403,
571   doi:https://doi.org/10.1016/j.neuroimage.2022.119403 (2022).

572   17   Jastorff, J. & Orban, G. A. Human functional magnetic resonance imaging
573   reveals separation and integration of shape and motion cues in biological
574   motion processing. *J Neurosci* **29**, 7315-7329, doi:10.1523/jneurosci.4870-
575   08.2009 (2009).

576   18   Vangeneugden, J., Peelen, M. V., Tadin, D. & Battelli, L. Distinct neural
577   mechanisms for body form and body motion discriminations. *J Neurosci* **34**,
578   574-585, doi:10.1523/jneurosci.4032-13.2014 (2014).

579   19   Troje, N. F. & Westhoff, C. The inversion effect in biological motion
580   perception: evidence for a "life detector"? *Curr Biol* **16**, 821-824,
581   doi:10.1016/j.cub.2006.03.022 (2006).

582   20   Chang, D. H. & Troje, N. F. Perception of animacy and direction from local
583   biological motion signals. *J Vis* **8**, 3.1-10, doi:10.1167/8.5.3 (2008).

584   21   Chang, D. H. & Troje, N. F. Characterizing global and local mechanisms in
585   biological motion perception. *J Vis* **9**, 8.1-10, doi:10.1167/9.5.8 (2009).

22   Bertenthal, B. I. & Pinto, J. Global Processing of Biological Motions.
Psychological Science **5**, 221-225, doi:10.1111/j.1467-9280.1994.tb00504.x
(1994).
23   Wang, L., Zhang, K., He, S. & Jiang, Y. Searching for life motion signals.
Visual search asymmetry in local but not global biological-motion processing.
Psychol Sci **21**, 1083-1089, doi:10.1177/0956797610376072 (2010).
24   Chang, D. H. F. & Troje, N. F. Acceleration carries the local inversion effect
in biological motion perception. *Journal of Vision* **9**, 19-19,
doi:10.1167/9.1.19 (2009).
25   Thornton, I. M., Rensink, R. A. & Shiffrar, M. Active versus passive
processing of biological motion. *Perception* **31**, 837-853, doi:10.1068/p3072
(2002).
26   Thompson, J. & Parasuraman, R. Attention, biological motion, and action
recognition. *Neuroimage* **59**, 4-13, doi:10.1016/j.neuroimage.2011.05.044
(2012).
27   Kim, J., Norton, D., McBain, R., Ongur, D. & Chen, Y. Deficient biological
motion perception in schizophrenia: results from a motion noise paradigm.
Frontiers in Psychology **4**, doi:10.3389/fpsyg.2013.00391 (2013).
28   Federici, A. *et al.* Anomalous Perception of Biological Motion in Autism: A
Conceptual Review and Meta-Analysis. *Scientific Reports* **10**, 4576,
doi:10.1038/s41598-020-61252-3 (2020).
29   Todorova, G. K., Hatton, R. E. M. & Pollick, F. E. Biological motion
perception in autism spectrum disorder: a meta-analysis. *Molecular Autism* **10**,
49, doi:10.1186/s13229-019-0299-8 (2019).
30   Wang, Y. *et al.* Heritable aspects of biological motion perception and its
covariation with autistic traits. *Proceedings of the National Academy of*
Sciences **115**, 1937-1942, doi:10.1073/pnas.1714655115 (2018).
31   Kröger, A. *et al.* Visual Processing of Biological Motion in Children and
Adolescents with Attention-Deficit/Hyperactivity Disorder: An Event Related
Potential-Study. *PLOS ONE* **9**, e88585, doi:10.1371/journal.pone.0088585
(2014).
32   Imanipour, S., Sheikh, M., Shayestefar, M. & Baloochnejad, T. Deficits in
Working Memory and Theory of Mind May Underlie Difficulties in Social

Perception of Children with ADHD. *Neurology Research International* **2021**,
3793750, doi:10.1155/2021/3793750 (2021).

33 McKay, L. S. *et al.* Do distinct atypical cortical networks process biological
motion information in adults with Autism Spectrum Disorders? *Neuroimage*
59, 1524-1533, doi:10.1016/j.neuroimage.2011.08.033 (2012).

34 American Psychiatric Association, D. & Association, A. P. *Diagnostic and*
statistical manual of mental disorders: DSM-5. Vol. 5 (American psychiatric
association Washington, DC, 2013).

35 Kaufman, J. *et al.* K-SADS-PL DSM-5 November 2016. *Yale University:*
Advanced Center for Intervention and Services Research (ACISR) for Early
Onset Mood and Anxiety Disorders-Western Psychiatric Institute and Clinic
Child and Adolescent Research and Education (CARE) Program (2016).

36 Dun, Y. *et al.* Reliability and validity of the Chinese version of the kiddie-
schedule for affective disorders and schizophrenia-present and lifetime version
DSM-5 (K-SADS-PL-C DSM-5). *Journal of Affective Disorders* **317**, 72-78,
doi:https://doi.org/10.1016/j.jad.2022.08.062 (2022).

37 Pind, J., Gunnarsdóttir, E. K. & Jóhannesson, H. S. Raven's Standard
Progressive Matrices: new school age norms and a study of the test's validity.
Personality and Individual Differences **34**, 375-386 (2003).

38 Reid, R. *et al.* Assessing culturally different students for attention deficit
hyperactivity disorder using behavior rating scales. *J Abnorm Child Psychol*
26, 187-198, doi:10.1023/a:1022620217886 (1998).

39 Su, Y. E. *et al.* Parent Ratings of ADHD Symptoms in Chinese Urban
Schoolchildren: Assessment With the Chinese ADHD Rating Scale-IV: Home
Version. *J Atten Disord* **19**, 1022-1033, doi:10.1177/1087054712461177
(2015).

40 Constantino, J. & Gruber, C. Social responsive scale (SRS) manual. *Los*
Angeles, CA: Western Psychological Services (2005).

41 Cen, C. Q. *et al.* Investigating the validation of the Chinese Mandarin version
of the Social Responsiveness Scale in a Mainland China child population.
BMC Psychiatry **17**, 51, doi:10.1186/s12888-016-1185-y (2017).

42 QbTech, A. QbTest plus technical manual. *Sweden: Gothenburg* (2010).

43 Hult, N., Kadesjö, J., Kadesjö, B., Gillberg, C. & Billstedt, E. ADHD and the
QbTest: Diagnostic Validity of QbTest. *Journal of Attention Disorders* **22**,
1074-1080, doi:10.1177/1087054715595697 (2015).
44 Vanrie, J. & Verfaillie, K. Perception of biological motion: a stimulus set of
human point-light actions. *Behav Res Methods Instrum Comput* **36**, 625-629,
doi:10.3758/bf03206542 (2004).
45 Fagerland, M. W. t-tests, non-parametric tests, and large studies—a paradox of
statistical practice? *BMC Medical Research Methodology* **12**, 78,
doi:10.1186/1471-2288-12-78 (2012).
46 Rochon, J., Gondan, M. & Kieser, M. To test or not to test: Preliminary
assessment of normality when comparing two independent samples. *BMC
Medical Research Methodology* **12**, 81, doi:10.1186/1471-2288-12-81 (2012).
47 Dittrich, W. H., Troscianko, T., Lea, S. E. & Morgan, D. Perception of
emotion from dynamic point-light displays represented in dance. *Perception*
25, 727-738, doi:10.1068/p250727 (1996).
48 Pollick, F. E., Kay, J. W., Heim, K. & Stringer, R. Gender recognition from
point-light walkers. *J Exp Psychol Hum Percept Perform* **31**, 1247-1265,
doi:10.1037/0096-1523.31.6.1247 (2005).
49 Kutsuki, A., Kuroki, M., Egami, S., Ogura, T. & Itakura, S. Individual
differences in changes in infants' interest in social signals in relation to
developmental index. *Infant Behav Dev* **32**, 381-391,
doi:10.1016/j.infbeh.2009.06.004 (2009).
50 Yoon, J. M. & Johnson, S. C. Biological motion displays elicit social behavior
in 12-month-olds. *Child Dev* **80**, 1069-1075, doi:10.1111/j.1467-
8624.2009.01317.x (2009).
51 Troje, N. F. & Basbaum, A. Biological motion perception. *The senses: A
comprehensive reference* **2**, 231-238 (2008).
52 Troje, N. F. & Chang, D. H. Shape-independent processing of biological
motion. *People watching: Social, perceptual, and neurophysiological studies
of body perception*, 82-100 (2013).
53 Troje, N. F. What is biological motion? Definition, stimuli and paradigms.
*Social perception: Detection and interpretation of animacy, agency, and
intention*, 13-36 (2013).

54   Vallortigara, G., Regolin, L. & Marconato, F. Visually inexperienced chicks
exhibit spontaneous preference for biological motion patterns. *PLoS Biol* **3**,
e208, doi:10.1371/journal.pbio.0030208 (2005).

55   Simion, F., Regolin, L. & Bulf, H. A predisposition for biological motion in
the newborn baby. *Proc Natl Acad Sci U S A* **105**, 809-813,
doi:10.1073/pnas.0707021105 (2008).

56   Chang, D. H. F., Ban, H., Ikegaya, Y., Fujita, I. & Troje, N. F. Cortical and
subcortical responses to biological motion. *NeuroImage* **174**, 87-96,
doi:https://doi.org/10.1016/j.neuroimage.2018.03.013 (2018).

57   Buzzell, G., Chubb, L., Safford, A. S., Thompson, J. C. & McDonald, C. G.
Speed of Human Biological Form and Motion Processing. *PLOS ONE* **8**,
e69396, doi:10.1371/journal.pone.0069396 (2013).

58   Hirai, M., Watanabe, S., Honda, Y. & Kakigi, R. Developmental changes in
point-light walker processing during childhood and adolescence: An event-
related potential study. *Neuroscience* **161**, 311-325,
doi:10.1016/j.neuroscience.2009.03.026 (2009).

59   Thornton, I. M. & Vuong, Q. C. Incidental processing of biological motion.
Current Biology **14**, 1084-1089, doi:10.1016/j.cub.2004.06.025 (2004).

60   Bosbach, S., Prinz, W. & Kerzel, D. A Simon Effect with Stationary Moving
Stimuli. *Journal of Experimental Psychology: Human Perception and*
Performance **30**, 39-55, doi:10.1037/0096-1523.30.1.39 (2004).

61   Hirai, M. & Senju, A. The two-process theory of biological motion processing.
Neurosci Biobehav Rev **111**, 114-124, doi:10.1016/j.neubiorev.2020.01.010
(2020).

62   Bardi, L., Regolin, L. & Simion, F. Biological motion preference in humans at
birth: Role of dynamic and configural properties. *Developmental Science* **14**,
353-359, doi:10.1111/j.1467-7687.2010.00985.x (2011).

63   Grossman, E. D., Blake, R. & Kim, C. Y. Learning to see biological motion:
brain activity parallels behavior. *J Cogn Neurosci* **16**, 1669-1679,
doi:10.1162/0898929042568569 (2004).

64   Frith, C. D. & Frith, U. Interacting minds--a biological basis. *Science* **286**,
1692-1695, doi:10.1126/science.286.5445.1692 (1999).

65   Troje, N. F. & Chang, D. H. F. Life Detection From Biological Motion.
Current Directions in Psychological Science **32**, 26-32,
doi:10.1177/09637214221128252 (2023).
66   Ghanouni, P. *et al.* Biological Motion Perception Is Affected by Age and
Cognitive Style in Children Aged 8–15. *Neurology Research International*
2015, 594042, doi:10.1155/2015/594042 (2015).
67   Annaz, D. *et al.* Development of motion processing in children with autism.
Dev Sci **13**, 826-838, doi:10.1111/j.1467-7687.2009.00939.x (2010).
68   Hubert, B. *et al.* Brief Report: Recognition of Emotional and Non-emotional
Biological Motion in Individuals with Autistic Spectrum Disorders. *Journal of*
Autism and Developmental Disorders **37**, 1386-1392, doi:10.1007/s10803-
006-0275-y (2007).
69   Saygin, A. P., Cook, J. & Blakemore, S.-J. Unaffected Perceptual Thresholds
for Biological and Non-Biological Form-from-Motion Perception in Autism
Spectrum Conditions. *PLOS ONE* **5**, e13491,
doi:10.1371/journal.pone.0013491 (2010).
70   Murphy, P., Brady, N., Fitzgerald, M. & Troje, N. F. No evidence for
impaired perception of biological motion in adults with autistic spectrum
disorders. *Neuropsychologia* **47**, 3225-3235,
doi:https://doi.org/10.1016/j.neuropsychologia.2009.07.026 (2009).
71   Rosch, E. *Principles of categorization*. (Morgan Kaufmann, 1988).
72   Cavanagh, P., Labianca, A. T. & Thornton, I. M. Attention-based visual
routines: sprites. *Cognition* **80**, 47-60, doi:10.1016/s0010-0277(00)00153-0
(2001).
73   Safford, A. S., Hussey, E. A., Parasuraman, R. & Thompson, J. C. Object-
based attentional modulation of biological motion processing: Spatiotemporal
dynamics using functional magnetic resonance imaging and
electroencephalography. *Journal of Neuroscience* **30**, 9064-9073,
doi:10.1523/JNEUROSCI.1779-10.2010 (2010).
74   Thompson, B., Hansen, B. C., Hess, R. F. & Troje, N. F. Peripheral vision:
Good for biological motion, bad for signal noise segregation? *Journal of*
Vision **7**, 12-12, doi:10.1167/7.10.12 (2007).

75   Hirai, M., Saunders, D. R. & Troje, N. F. Allocation of attention to biological
motion: Local motion dominates global shape. *Journal of Vision* **11**, 4-4,
doi:10.1167/11.3.4 (2011).
76   Rutherford, M. D. & Troje, N. F. IQ Predicts Biological Motion Perception in
Autism Spectrum Disorders. *Journal of Autism and Developmental Disorders*
42, 557-565, doi:10.1007/s10803-011-1267-0 (2012).
77   van Boxtel, J. J., Dapretto, M. & Lu, H. Intact recognition, but attenuated
adaptation, for biological motion in youth with autism spectrum disorder.
Autism Res **9**, 1103-1113, doi:10.1002/aur.1595 (2016).
78   Koldewyn, K., Whitney, D. & Rivera, S. M. The psychophysics of visual
motion and global form processing in autism. *Brain* **133**, 599-610,
doi:10.1093/brain/awp272 (2010).
79   Atkinson, A. P. Impaired recognition of emotions from body movements is
associated with elevated motion coherence thresholds in autism spectrum
disorders. *Neuropsychologia* **47**, 3023-3029,
doi:10.1016/j.neuropsychologia.2009.05.019 (2009).
764
765

 Figures

Figure 1. Schematic representation of Biological Motion (BM) and scrambled

BM sequence. Intact walker contains information of local kinematics and global

configuration. Local kinematics refers to the motion tracks of each critical joint

illustrated by the chromatic dot. Global configuration is composed of the relative

locations of each joint. In the scrambled BM sequence, global configuration cues have

been removed, but local kinematics have been retained. (Figure reconstructed from

Wang et al., 2018[30])

Figure 2. Illustration of the trial sequence. In BM-local, a monitor displayed

scrambled BM sequences. Participants only judged the facing direction of the

scrambled walker using local BM cues. In BM-global, each trial only showed an

intact or scrambled walker (black dots in the figure) embedded within a mask

containing local BM cues. Because the two conditions contained the same local cues

that were also present in the mask, the participant must rely on global BM cues to

determine whether an intact walker was present in the mask. The figure shows one of

five possible directions the intact walker could face (i.e., facing participants). In BM-

general, the stimuli sequence consisted of an intact walker (black dots) and a mask

containing similar target local cues, and children judged the direction the walker was

facing using general BM cues (local + global). Dots in the figure are rendered in black

for better illustration but were displayed in white in the actual experiments.

Figure 3. The mean accuracy of the three tasks. Typically developing (TD)

children had higher accuracies than children with attention deficit hyperactivity

disorder (ADHD) in the three tasks in Experiment 1. Error bars show standard

deviations. ***$p < 0.001$

Figure 4. Correlations between response accuracies and SRS total score. The ability to process local cues is significantly correlated with the SRS total score in the ADHD group. The grey shading represents the 95% confidence interval. ** FDR-corrected $p < 0.01$.

Figure 5. Factors influencing BM perception in ADHD children. Post hoc path analysis confirmed that the effect of sustained attention on performance in BM-General was entirely mediated by Perceptual Reasoning, and the ability of global BM processing partly mediated the effect of age on performance in BM-General.

Tables

Table 1. Demographic characteristics of TD and ADHD groups

	Experiment 1		*p* value	Experiment 2		*p* value
	TD	ADHD		TD	ADHD	

Sample	36	39		33	42	
Age (years)	9.09±2.18	9.88±2.23	0.126	8.75±1.94	9.34±1.89	0.191
Sex ratio (% male)	38.89%	71.79%	0.004	42.42%	64.28%	0.059

806 **Table 2. The correlation between the ability of local BM processing and**
807 **subdimensions of SRS in ADHD children.**

	Correlation coefficient (r)	FDR-corrected p value
Social awareness	-0.333	0.039
Social cognition,	-0.416	0.020
Social communication	-0.381	0.022
Social motivation	-0.406	0.020
Autistic mannerisms	-0.245	0.117

Videos

Video 1. An intact walker without a mask. The dots in the video are rendered in
chromatic colors for better illustration and displayed in white in the actual
experiments.

Video 2. An example of BM-Local (a scrambled walker without a mask). The
chromatic dots in this video correspond to the major joints of the intact walker in
Video 1 and are displayed in white in the actual experiments.

Video 3. An example of BM-Global (an intact or scrambled walker with a mask).
The chromatic dots in this video correspond to the major joints of the intact walker in
Video 1 and are displayed in white in the actual experiments.

Video 4. Intact walkers facing five directions in BM-Global. The dots in this video
are rendered in black for better illustration and displayed in white in the actual
experiments.

Video 5. The example of BM-General (an intact walker with a mask). The
chromatic dots in this video correspond to the major joints of the intact walker in
Video 1 and are displayed in white in the actual experiments.