The 14-3-3/spastin pathway coordinates neurite regeneration and recovery of spinal cord injury by targeting microtubule dynamics

Author: Qiuling Liu[1][#], Hua Yang[1][#], Jianxian Luo[1][#], Cheng Peng[1], Ke Wang[1], Guowei Zhang[1], Zhisheng Ji[1]*, Hongsheng Lin[1]*

1 Department of Orthopedics, The First Affiliated Hospital of Jinan University, Guangdong province, Guangzhou 510630, People's Republic of China;

[#]These authors contribute equally to this work.

* Corresponding to:

Zhisheng Ji (jizhisheng0521@163.com)

Hongsheng Lin (tlinhsh@jnu.edu.cn)

Department of Orthopedics, The First Affiliated Hospital of Jinan University, 613 Huangpu Avenue, Guangzhou 510630, People's Republic of China.

Running title: 14-3-3/spastin pathway regulates the recovery of spinal cord injury.

Abstract Axon regeneration is abortive in the central nervous system following injury.

Orchestrating microtubule dynamics has emerged as a promising approach to improve

axonal regeneration. The microtubule severing enzyme spastin is essential for axonal

development and regeneration through remodeling of microtubule arrangement. To

date, however, little is known regarding the mechanisms underlying spastin action in

neural regeneration after spinal cord injury. Here, we use glutathione transferase

pulldown and immunoprecipitation assays to demonstrate that 14-3-3 interacts with

spastin, both *in vivo* and *in vitro,* via spastin Ser233 phosphorylation. Moreover, we

show that 14-3-3 protects spastin from degradation by inhibiting the ubiquitination

pathway and upregulates the spastin-dependent severing ability. Furthermore,

improving the interaction between 14-3-3 and spastin by Fusicoccin (FC-A) promotes

neurite outgrowth and regeneration *in vitro*. Western blot and immunofluorescence

results revealed that 14-3-3 protein is upregulated in the neuronal compartment after

spinal cord injury *in vivo*. In addition, administration of FC-A not only promotes

locomotor recovery, but also nerve regeneration following spinal cord injury in both

contusion and lateral hemisection models. However, application of spastin inhibitor

spastazoline successfully reverses these phenomena. Taken together, these results

indicate that 14-3-3 is a molecular switch that regulates spastin protein levels, and

14-3-3/spastin pathway is an important target for regulation of microtubule dynamics

for nerve regeneration after spinal cord injury.

Keywords: Spastin; Microtubule; Axon regeneration; 14-3-3; Spinal cord injury

Graphical Abstract

Introduction

Regeneration of the adult central nervous system neurons is difficult after injury, a

phenomenon that results in permanent neurological impairments[1]. Researchers have

attributed this not only to the highly inhibitory extrinsic environment at the injury site

but also to limited cell-intrinsic growth capacity. Many studies have demonstrated that

re-programming of growth-associated processes plays a key role in manipulation of

axonal regeneration[2-4]. However, the specific mechanisms underlying axonal

regeneration failure remain largely unknown. Cytoskeleton remodeling, including

actin and microtubule (MT) reorganization, appears to be the key factors directing the

fate of axonal regeneration and sprouting[5]. Notably, the retraction bulb of severed

axons at the lesion site of the spinal cord needs a dynamic cytoskeleton for successful

regeneration. Moreover, formation of axon collaterals from spared axons also needs

dynamic cytoskeleton reorganization to form new neural circuits with the innervated

neurons[6, 7]. Mechanistically, invasion by MTs can not only provide mechanical

force, but also guide intracellular transport to regulate distribution of specific

molecules, including mitochondria, peroxisomes, growth factors and other, and

subsequently mediate axonal extension. Therefore, timely invasion of microtubules

can not only power the formation of a growth cone with competent growth ability from the retraction bulb, but also improve the process of axonal sprouting from the spared axons[8, 9]. Overall, manipulating MT dynamics is critical for boosting axon regeneration.

Spastin is encoded by SPG4 gene and its mutations were identified form the patients of hereditary spastic paraplegia (HSP)[10]. It is a MT severing enzyme (including spastin, katanin and fidgetin) which can cut long MTs into short fragments, a phenomenon that not only endows MTs with high dynamics but also mediates their subsequent turning and regrowth under physiological conditions[11]. Spastin has two major isoforms, namely M1 and M87, coded form different initial sites. Both isoforms are highly expressed in the spinal cord[12]. Moreover, spastin mutation lacking severing activity can lead to the degeneration of the corticospinal tracts in HSP patients. In addition, spastin is highly expressed in the central neuronal system (CNS) and essential for formation of axon branches as well as their extension during neural development[13, 14]. These evidences indicate that spastin plays an important role in axonal development and maintainence in the spinal cord. Intriguingly, several studies have demonstrated that spastin is necessary for successful axon regeneration, but not other severing enzyme like katanin and fidgetin[15]. Moreover, loss spastin function reportedly resulted in an almost complete failure of axon regeneration, suggesting that it plays a vital role in axon regeneration following spinal cord injury. Therefore, manipulating spastin expression may be an effective way to promote axon regeneration. However, efficient regeneration is dependent on the precise spastin dosage, with studies showing that excess spastin is toxic and may destroy the MT network, while low doses have been associated with insufficient MT severing which subsequently limits MT remodeling for regeneration[15, 16]. In this case, direct manipulation of the spastin gene may not be the most appropriate approach for promoting successful axon regeneration after the CNS injury. It is worth noting that posttranslational modifications, protein metabolism and protein-protein interactions are vital for regulating the protein's activity[17]. Therefore, elucidating the specific

molecular mechanism underlying spastin action is imperative to guiding future

development of effective interventions to promote axon regeneration following

damage to the CNS.

14-3-3 protein is a type of adaptor protein that play important roles in many

signaling pathways through interaction with their substrates[18, 19]. For instance, it

inhibits the ROS-induced cell death by interacting with FOXO transcription factors to

prevent upregulation of many pro-apoptotic genes[20]. In mammals, seven isoforms,

namely β、γ、ε、ζ、η、θ and σ, have been identified. They are not only highly preserved

in the mammalian system, but also form a dimeric structure to interact with their

substrates which are mostly phosphoproteins[21]. Previous studies have shown that

spastin is phosphorylated in Ser233 which showed its potential for interaction with

14-3-3[22]. In addition, 14-3-3 protein is also highly expressed in CNS where it plays

a vital role in both axonal growth and guidance. Previous studies have also

demonstrated that 14-3-3 promotes cortical neurite outgrowth and axon

regeneration[23, 24]. To date, however, the molecular mechanisms underlying its

action in promotion of axon regeneration as well as its role in spinal cord injury

remain largely unknown.

In this study, we demonstrate that 14-3-3 interacts with spastin to regulate its

stability, thus controlling both neurite outgrowth and regeneration after injury. Our

results reveal that 14-3-3 interacts with serine 233 in spastin, in a

phosphorylation-dependent manner, thereby preventing its degradation by

ubiquitination and upregulates the severing ability via spastin-mediated activity.

Moreover, we found that enhancing this interaction via Fusicoccin (FC-A) application

resulted in enhanced axon regeneration and locomotor improvement both in contusion

model and in T10 lateral hemi-transection model of spinal cord injury. Collectively,

our findings suggest 14-3-3/spastin pathway is a novel intervention target for axon

regeneration after spinal cord injury.

Materials and methods

Animals

Sprague Dawley rats (P0) were used for primary hippocampal and cortical neuron culture, whereas adult female mice (C57BL/6J, 6-8 weeks) were used to establish the lateral hemi-transection model and the contusion model of spinal cord injury. Animals were housed in a room, maintained at 22°C under a 12:12 h light/dark photoperiod, and given food and water ad libitum. All animal treatments were carried out in strict adherence to the guidelines by the National Institutes of Health's Guide for the Care and Use of Laboratory Animals. The study protocol was approved by Jinan University Institutional Animal Care and Use Committee. The number of animals utilized and suffering were both reduced to the absolute minimum.

Cell culture and transfection. Dissociated hippocampi were digested with 2 mg/mL papain and plated at a density of 1×10^4cells/cm^2 onto poly-D-lysine-coated coverslips for primary hippocampal neuron culture. Cultures were kept alive in an incubator humidified with 5% CO_2 at 37°C using Neurobasal-A media supplemented with 0.5 mM glutamine and 2% B27. Every three days, half of the cultural media was replaced. Various plasmids were transfected with calcium phosphate at 2th day in vitro (DIV). COS7 cells and human embryonic kidney (HEK) 293 cells were grown in a 5% CO_2 incubator at 37°C using Dulbecco's modified Eagle media with 10% fetal bovine serum and 1% penicillin/streptomycin as supplemental ingredients. The constructs were transfected into the COS7 cells and HEK293 cells using Lipofectemine 2000. Cells were cultured for 24-36 hours after transfection before being harvested.

Plasmids and constructs

Several 14-3-3 isoforms were amplified in cDNA from Sprague Dawley rats via Polymerase Chain Reaction (PCR) as previously described[14]. The amplicons were inserted into the pCMV-Tag-2b or pGEX-5X-3 vector (Addgene). Spastin mutants were generated using the Quickchange Kit (Agilent, Santa Clara, CA). R18 was synthetized by IGE Biotechnology LTD （Guangzhou, China） and inserted into pCDNA 3.1 vector. All constructs were verified by sequencing by IGE Biotechnology LTD（Guangzhou, China）.

Expression of recombinant proteins and GST-Pulldown assay

Purification of GST fusion of 14-3-3 proteins and pulldown experiments were
performed as previously described[14, 25]. Briefly, GST-14-3-3s constructs
(Invitrogen) were transformed into *Escherichia coli* strain BL21 (DE3) strains.
Isopropy-D-thiogalactoside was incubated for 6 hours at 30°C to stimulate the
synthesis of fusion proteins. The bacteria were first centrifuged, then resuspended in a
mixture of protease inhibitors (Merck, Whitehouse Station, NJ). The cell suspension
was treated with 0.1% lysozyme, followed by 0.5% deoxycholic acid on ice for 20
min. The contents were, sonicated (10,000 rpm for 30 min), centrifuged and the
supernatant collected, then Glutathione-Sepharose beads (supplemented with 1%
Triton X-100) used to purify GST fusion proteins in the supernatant. Next, 5 μg of
GST-fusion protein was incubated with 400 μg of protein from the spinal cord (1 cm
around T10) of SD rats (P0) under gentle rotation, then centrifugated to collect the
GST-binding proteins. Western blot assay was then performed to quantify
GST-binding proteins.

Immunoprecipitation and western blotting.

Immunoprecipitation (IP) assays were carried out as previously described[26]. Briefly,
total cell extracts or tissues (1 cm around T10) were prepared by soaking them in 400
μL of lysis buffer for 30 minutes at 4 °C. The lysates were centrifuged at 10,000 rpm
for 10 minutes at 4 °C after a brief sonication. The extract was immunoprecipitated
with 2 μg antibodies against spastin, GFP, or Flag, and then incubated for 4 hours at
4 °C with 60 μL of protein G plus protein A/G agarose using continuous gentle
inversion. Three times, the immune complexes were pelleted and washed. Western
blot assay was performed to analyze the precipitated complexes. Western blot analysis
was carried out as previously described[14]. SDS-PAGE was used to separate the
lysates, which were then electrophoretically transferred to a polyvinylidene difluoride
membrane. Membranes were probed with primary antibodies at 4 °C overnight after
being blocked in Tris-buffered saline with 5% milk and 0.05% Tween. The
membranes were cleaned, incubated with goat anti-mouse or anti-rabbit secondary
antibodies that were horseradish peroxidase conjugated, and then visualized with

ECL-prime. Anti-spastin (1:1000, ABN368 from Millipore), anti-Pan 14-3-3 (1:1000, #8312 from Cell Signaling Technology), anti-GFP (1:5000, Ab290 from Abcam), anti-Flag (1:1000, F3165 from Sigma-Aldrich), anti-GAPDH (1:5000, AC002 from ABclonal), and HRP conjugated goat-derived anti-rabbit or anti-mouse were the antibodies used (1:5000, Abclonal).

Scratch assay

Rat cortical neurons were seeded in 96-well plates (30,000 cells per well), then a plastic P10 pipet tip was used to scratch across the center of the well. Half the media was aspirated out and replaced with fresh one every three days until the 7th DIV. The wells were immediately filled with chemicals after scratching. The cultures were fixed and stained with anti-βIII tubulin (1:1000, ab18207) the following day. Images of axons stained with βIII tubulin were collected, imported in ImageJ, and the proportion of neurites stained by βIII tubulin in the central 70% of the scratch were calculated.

Lateral hemi-sectioning and contusion model of SCI

The procedure for T10 lateral hemi-sectioning and contusion model was as previously described[27, 28]. Briefly, mice were first anesthetized with 1.25% 2,2,2-tribromo ethanol (0.2 mL/100 g), via intraperitoneal injection for 3-4 min, then a laminectomy performed near T10 after making a midline incision over the thoracic vertebrae. The T10 level of the spinal cord was recognized by the anatomical landmark which is parallel to the twelfth rib of the mice. For the hemi-sectioning injury model, the right unilateral hemi-section was carefully performed using a scalpel and microscissors, with care taken to avoid damaging the spinal cord dura. For the contusion model, the nitrogen tank controlling the impactor tip was set at 18 psi or 124 kPa. A U-shaped stabilizer with the rat was loaded onto the stage of the Louisville Injury System Apparatus (LISA) and the dura of the spinal cord was adjusted directly under the impactor while monitored by the laser beam. Crash depth was adjusted to 1.0 mm for injury, and with time set to 0.5 s. Finally, the muscles were stitched and the skin was stapled back together.

Pharmacological interference with FC-A or spastin

An equal volume of normal saline was intraperitoneally injected into SCI control

mice. Mice were intraperitoneally injected with FC-A (Enzo life sciences,

ALX-350-115) at a dose of 15 mg/kg starting immediately after SCI and every other

day until end of the experiment. Mice were intraperitoneally injected with

spastazoline (MedChemExpress, HY-111548) at a dose of 20 mg/kg, immediately

after SCI then every other day until end of experiments.

Basso Mouse Scale

Two observers who were blind to the experimental groups conducted the Basso

Mouse Scale (BMS) after the T10 SCI in accordance with a previously described

procedure[29]. Mice were briefly left in an open field for 4 minutes to observe all

visually perceptible signs of locomotor recovery. The scores were based on hind limb

movements performed in an open field and included hind limb joint movement,

weight support, plantar stepping, coordination, paw position, and trunk and tail

control (0, full hind limb paralysis; 9, normal locomotion).

Catwalk analysis

According to previously described methods[30], the CatWalk XT 9.1 automated

quantitative gait analysis system (Noldus Company, Netherlands) was also used to

evaluate the locomotor recovery of mice after injury. Briefly, one week before surgery,

mice were trained to walk in a single direction for the entire length of a darkened

Catwalk chamber in a quiet room five times daily. Seven weeks after surgery, the

mice were assessed five times using the same system. Gait regularity index and

hindlimb contact area were recorded. The regularity index was calculated as the

number of normal strides multiplied by four and then divided by the total number of

strides; this value is ~100% in normal animals.

Footprint analysis

Footprint analysis was conducted before the mice were executed, using a

previously described protocol[31]. Briefly, each animal's hindlimbs were first coated

with nontoxic paint. The mice were then moved from a brightly illuminated starting

box to a darkened one, by allowing them to walk through a narrow-custom-built

plexiglass trough (5 cm wide by 40 cm long), during which they left a trace of their paw prints on the white sheet that overlaid the trough. The footprints were scanned, and digitized images measured using Image J Pro Plus. Stride length was measured as the distance between the adjacent hindlimbs, whereas stride width was taken as the distance between the affected and the contralateral limbs. At least ten footprints per side, from three sessions per animal, were measured for both parameters and their average were calculated.

Foot fault test

The Parallel Rod apparatus was used in this test[32]. Summarily, mice were positioned in the middle of the device, allowed to walk for three minutes during which a video was captured. The test was performed by two observers who were blinded to the experimental groups. One observer recorded the total number of steps taken, while the other counted instances where each right hind limb slipped through the rod. Three runs were recorded for every mouse.

Motor evoked potential (MEP) recordings

To assess SCI recovery, motor evoked potentials (MEPs) were recorded 7 weeks after SCI according to previously described methods[30]. First, mice were anesthetized with 1.25% 2,2,2-tribromo ethanol (0.2 mL/100 g). A craniotomy was then performed to expose the M1 region of the motor cortex. Electrode penetration was guided via a stereotaxic instrument to a depth of 700-1000 mm from the brain surface to target corticospinal neurons in the sensorimotor cortex. The recording electrode was placed on the gastrocnemius muscle and the reference electrode was placed on the paraspinal muscle between the stimulation and recording points. A ground electrode was attached to the tail. A single square-wave stimulus of 0.5 mA, 0.5 ms duration, 2 ms time delay, and 1 Hz was used. Amplitude was measured from the beginning of the first response wave to its maximum point. All potentials were amplified and acquired using a digital oscilloscope (Chengdu Instrument Factory, Chengdu, China).

Immunocytochemistry

Cells were grown on coverslips and processed according to
immunocytochemistry protocols as previously described[14]. Cells were fixed using
freshly prepared 4% paraformaldehyde, followed by permeabilization with 0.1%
Triton X-100 in TBS and blocking in 3% normal donkey serum. The cells on the
coverslips were incubated with anti-GFP (1:1000, Ab290 form Abcam) or
anti-α-tubulin(1:1000, ab7291 from Abcam) and then incubated with secondary
antibody (Alexa Fluor 488 donkey anti-rabbit or Alexa Fluor 647 donkey conjugated
to anti-goat immunoglobulin G (H+L) (Life Technologies). The coverslips were then
mounted with DAPI (diamidino-2-phenylindole)-free Fluoro-Gel on glass slides
(Electron Microscopy Sciences, USA). Cells were scanned using a 63x oil immersion
objective mounted on a confocal microscope (LSM 710 Meta; Carl Zeiss). LSM 710
software was used to process the images after they were captured using sequential
acquisition settings at a resolution of 1024 × 1024 pixels and a 12-bit depth.

Mice were administered consecutive transcardial infusions of 4%
paraformaldehyde in PBS and 0.09% saline after being given a deep anesthetic. After
tissue collection, post-fixation in 4% paraformaldehyde was applied overnight, and
tissues were then immersed in 30% sucrose for two days to dehydrate them. Serial
sagittal cryostat-sections were incubated at 4°C for at least 18 hours before being
stained with primary antibodies in blocking buffer (3% goat serum and 5% donkey
serum in 0.3% PBST). Alexa Fluor-coupled secondary antibodies (Life Technologies)
diluted in blocking buffer were incubated for 2 hours at room temperature to perform
the detection. Anti-GFAP (1:1000, GB11096 from Servicebio), anti-serotonin (1:5000,
AB125 from Sigma-Aldrich), anti-neurofililament (1:200, A19084 from ABclonal)
anti-Myelin Basic Protein (1:200, A11162 from ABclonal) anti-Brdu (1:200, A20790
from ABclonal), anti-Nestin (1:200, A11861 from ABclonal) and anti-NeuN (1:200,
A19086 from ABclonal) were used. Images were scanned using a confocal
microscope.

Statistical analysis

The analyses were carried out by Graphpad 8.4.0 software. Differences across

multiple groups were determined using One-way analysis of variance (ANOVA),

followed by Newman-Keuls post hoc tests for mean separations. $P \leqslant 0.05$ were

considered statistically significant.

Results

14-3-3 protein interacts with spastin

Spastin, a MTs severing enzyme, can cut long MTs into short fragments thereby

contributing to axonal elongation and branching. However, excess spastin levels by

gene manipulation can not only destroy the MT cytoskeleton but are also toxic to cells.

Unraveling the underlying mechanisms of spastin regulation under physiological

conditions may provide novel insights into the mode of axon regeneration following

CNS damage. In the present study, we performed GST-spastin pulldown assays on

lysates from the spinal cord tissue, then analyzed the binding complex via mass

spectrometry to determine the underlying molecular mechanisms of spastin. This

proteomics approach revealed14-3-3 as putative spastin-interacting proteins (Figure

1A & Figure S1A). Notably, 14-3-3 proteins consist of seven isoforms that are highly

conserved and act by targeting phosphoserine and phosphothreonine motifs of

substrate proteins. We have detected different many 14-3-3 peptides in the complex,

including LAEQAER, NLLSVAYK, and AVTEQGAELSNEER (Figure S1A).

Notably, these peptides were highly reserved among different isoforms (Figure S1B).

Next, we employed immunoprecipitation to determine the biochemical interaction

between 14-3-3 protein and spastin, and found that they endogenously interacted in

the spinal cord tissue (Figure 1B). To determine where the 14-3-3/spastin protein

complex functions in neurons, we double stained hippocampal neurons with spastin

and 14-3-3 antibody, and found that 14-3-3 was colocalized with spastin in the entire

cell compartment (Figure 1C), especially in the growth cone (Figure 1C1) and axon

shaft (Figure 1C2). Colocalization profiles are displayed in Figure 1D. To further

identify which isoform of 14-3-3 interacts with spastin, we generated six 14-3-3

isoforms in rats (β、γ、ε、ζ、η、θ), then purified GST fusion 14-3-3 proteins (Figure

1E). Results from a GST pull-down assay revealed that all 14-3-3 isoforms could

interact with spastin (Figure 1F). Subsequent co-immunoprecipitation assay results

confirmed that all 14-3-3 isoforms could form direct complexes with spastin (Figure

1G). Collectively, these results indicated that the 14-3-3 protein interacts with spastin

both *in vitro* and *in vivo*.

14-3-3 binds to phosphorylated Ser 233 in spastin

Next, we characterized the interaction region of spastin/14-3-3 binding. To this

end, we generated several GFP-tagged deletion constructs based on the modular

domain of spastin (Figure 2A), then co-expressed them with Flag-tagged 14-3-3 into

HEK293T cells. Co-immunoprecipitation assay, using GFP-Trap beads, revealed that

spastin fragments containing amino acids 85-316 but not those with 85-214

coprecipitated with 14-3-3 (Figure 2B). Next, we performed immunoprecipitation

assays using MTBD region (215-316 amino acids) of spastin with 14-3-3 and found

that 14-3-3 specifically binds to the MTBD region of spastin (Figure 2C).

Previous studies have shown that phosphorylation of the 14-3-3 substrate

regulates the binding of 14-3-3 proteins, although 14-3-3 can also bind to

non-phosphorylating proteins. To investigate whether spastin's phosphorylation state

affects its binding with 14-3-3, we performed pulldown assays by using GST-14-3-3

and GFP-spastin from transfected HEK293T cell extracts in the presence of broad

protein kinase staurosporine. Results revealed that presence of staurosporine markedly

weakened binding between spastin and 14-3-3, indicating that phosphorylation of

spastin is sufficient for the binding with 14-3-3 (Figure 2D). Next, we predicted five

putative 14-3-3 binding sites in spastin, namely amino acids 213-218, 230-235,

243-249, 425-460, and 559-564, which match the sequence motif of RSX(S/T)XP.

Combination with phosphorylation of spastin reported from previous MS data

(https://www.phosphosite.org/proteinAction.action?id=8325&showAllSites=true),

revealed two putative binding motifs (Ser233 and Ser562) in spastin as putative

targets for 14-3-3 binding (Figure 2E). Then, we mutated single Ser233 and Ser562 or

both to alanine in the spastin tagged with GFP and co-expressed them with

Flag-tagged 14-3-3 in HEK293T cells. Immunoprecipitation of the lysates, using an antibody against GFP, revealed that spastin S562A can bind with 14-3-3, while spastin S233A and spastin Pan A cannot (Figure 2F). Next, we transfected Ser233 mutation of spastin (spastin S233A or spastin S233D) with flag tagged 14-3-3 and generated Pearson's correlation coefficients. Results revealed that spastin S233D was markedly colocalized with 14-3-3, with minimal colocalization with spastin S233A (Figure 2A-B). These results were consistent with those in Figure 3B-C, where Ser233 was localized in the MTBD domain, further indicating that the binding of spastin-14-3-3 requires phosphorylation of spastin Ser233.

To examine whether Ser233 in spastin is involved in neurite outgrowth mediated by spastin, wild type (GFP-spastin) or spastin Ser233 mutant （GFP-spastin S233A or GFP-spastin S233D） was moderately expressed in primary hippocampal neurons at 2th DIV. Consistent with earlier results, spastin promoted neurite outgrowth, as evidenced by both the length and total branches of neurite. A similar effect was observed after transfection with spastin S233A. In contrast, we found a toxic effect following transfection with spastin S233D (Figure 2G-I). Taken together, these data suggests that the binding of 14-3-3 to the Ser233 in spastin has a direct influence on the fate of neurite outgrowth.

14-3-3 protects spastin from degradation via spastin S233 phosphorylation

How does the interaction between 14-3-3 and Ser233 in spastin regulate neurite outgrowth during neural development, and why does spastin S233D expression in neurons cause toxic effect? Our earlier results demonstrated that spastin acts in a dose-dependent manner, with excessive dosages found to destroy the microtubules and generate a toxic effect. Here, we hypothesized that spastin S233D transfection may upregulate expression of the spastin protein relative to WT and spastin S233A. To further test whether 14-3-3 would affect spastin stability, we performed chase-time assays via GFP-spastin overexpression, followed by treatment with Cycloheximide (CHX) to inhibit protein synthesis. Results showed that expression of GFP-spastin protein was markedly downregulated at 9 h after CHX treatment (Figure 3A). Next,

we co-expressed Flag-14-3-3 with GFP-spastin in the same treatment and found that

14-3-3 could significantly protect spastin from degradation after CHX treatment, even

after 24 h (Figure 3B-E). We also investigated whether 14-3-3 inhibited the

degradation in spastin S233 phosphorylation by transfecting with GFP-tagged spastin

S233A or S233D followed by CHX treatment. Results showed that 14-3-3 could

protect spastin (S233D), but not spastin (S233A), from degradation (Figure 3C-F).

A previous study demonstrated that spastin could be polyubiquitinated[33]. To

further determine whether 14-3-3 affected spastin's ubiquitination pathway, we

performed co-immunoprecipitation assays. GFP-spastin and Flag-ubiquitin were

transfected into HEK293T cells, co-expressed with 14-3-3 binding inhibitor R18 or

not . Results indicated that GFP-spastin could be ubiquitinated, while inhibiting the

binding of 14-3-3/spastin promoted spastin ubiquitination (Figure 5G). These results

indicated that 14-3-3 protects spastin from degradation by inhibiting the

ubiquitination pathway.

To evaluate whether spastin mediated microtubule severing activity, we

co-transfected GFP-spastin encoding plasmids into COS7 cells with or without

ubiquitin. COS7 cells are very flat, hence provide more spatial resolution compared to

neurons. Results showed that cells transfected with wild type spastin significantly

induced microtubule severing, as evidenced by almost no MT staining (Figure 3H),

while ubiquitin inhibited severing, leading to increased microtubule staining. Based

on these conditions, further transfection with 14-3-3 plasmid induced more

microtubule staining, indicating that 14-3-3 upregulates microtubule severing and the

process is mediated by spastin. Furthermore, we investigated whether this effect was

dependent on spastin S233 phosphorylation by replacing GFP-spastin with

GFP-spastin (S233A or S233D) during transfection. Results showed that 14-3-3 could

upregulate the severing ability of spastin S233D, but not S233A (Figure 3I), further

indicating that 14-3-3 binds to the phosphorylation of S233 in spastin and promotes

its microtubule severing ability. Collectively, these results suggest that 14-3-3 protein

protects spastin from degradation via spastin S233 phosphorylation.

14-3-3/spastin complex regulates neurite outgrowth and regeneration after injury

Next, we investigated how the interaction between 14-3-3 and spastin regulates neurite outgrowth and regeneration by applying FC-A and R18 to either stabilize or inhibit 14-3-3 binding. Previous studies have shown that FC-A and R18 interact with 14-3-3 protein by either stabilizing or inhibiting the binding of 14-3-3 with their substrates by directly docking to the groove in the 14-3-3 proteins with highly efficiency[34-36]. The structure of FC-A and R18 binding with 14-3-3 are shown in Figure 4A. We also used another drug, spastazoline, to inhibit microtubule severing as previously described[37, 38]. To test our hypothesis, we first explored whether FC-A or R18 could enhance or inhibit the interaction between 14-3-3 and spastin by co-immunoprecipitation with lysates of HEK293T cells, which had been transfected with GFP-spastin and Flag-14-3-3 plasmids whether together with FC-A application or R18 transfection. We found that FC-A could enhance the binding of 14-3-3/spastin, as evidenced by more intense bands compared to no FC-A application, whereas R18 could efficiently inhibit the binding of 14-3-3/spastin with almost no binding was detected (Figure 4B). Subsequently, we used FC-A together with spastazoline to examine the effects of 14-3-3/spastin protein complex on neurite outgrowth and regeneration.

Since 14-3-3 proteins play important roles in inhibition of ROS-induced cell death, we first explored the effect of 14-3-3/spastin binding in neurite outgrowth and repair under glutamate circumstances. Briefly, we established a neural injury model as previously described[39], and found that they had numerous swellings and breakages on the neurite, indicative of successful establishment of an injury model (Figure 4C). After injury, FC-A was administered into primary hippocampal neurons during the developmental stage 2, combination with application of spastin inhibitor spastazoline which had been confirmed by previous study[37]. Results demonstrated that FC-A could promote neurite outgrowth, both with regards to lengths and branches, after glutamate-induced injury, while this promoting effect was abolished following spastazoline administration (Figure 4D-G). A similar effect was observed in neurons

without glutamate induced injury (Figure S4). In contrast, transfection of R18 resulted

in an inhibitory effect on neurite outgrowth, but this was rescued with moderate

spastin expression after injury (Figure 4D-J). A similar trend was observed in neurons

without glutamate-induced injury (Figure S5). Collectively, these results suggest that

14-3-3 protein promotes the repair of injured neuron through the interaction with

spastin.

Next, we investigated the effect of enhanced 14-3-3/spastin binding on neurite

regeneration. Briefly, cortical neurons were grown at the 7th DIV then subjected to

scratch assays by drawing a line over the coverslips using a pipet tip. Neurons were

then treated with either FC-A or spastazoline for 36 hours. Results showed that FC-A

markedly improved neurite regeneration consistent with previous study[34], while

spastazoline application almost abolished the enhanced effects of regeneration (Figure

4K-L). To further quantify the extend of axon regeneration across each group, we

treated cortical neurons with either FC-A or spastazoline for 24 hours, then quantified

the longest regenerative axons. Results revealed similar effects (Figure S6). Taken

together, these results indicate that spastin-dependent severing is a prerequisite for the

enhancement of neurite regeneration mediated by FC-A.

FC-A-mediated nerve regeneration after SCI requires spastin activation by

targeting MTs

Given that 14-3-3 plays vital roles in neurite outgrowth and axonal

regeneration[40], we then hypothesized that 14-3-3 could be involved in the recovery

of spinal cord injury. To test this hypothesis, we established a spinal contusion injury

model and then employed western blot assay to quantify levels of 14-3-3 protein

following spinal cord injury. Results revealed that 14-3-3 was upregulated at 3th and

7th days post injury (DPI), but downregulated at 14 DPI. The 14-3-3 protein level

then reached a peak level at 30 DPI (Figure 5A-B). Moreover, immunostaining results

indicated that 14-3-3 protein upregulation was principally colocalized in the neuronal

compartment (βIII-tubulin labeled) (Figure 5C). Previous studies have demonstrated

that neurons were surrounded by inhibitory microenvironment 2 weeks after spinal

cord injury, our results also showed a slightly decrease at 14 DPI, indicating that 14-3-3 acts as an intrinsic switch for neural repair. Moreover, 14-3-3 was reported to be upregulated and acted as a biomarker for diagnosis of Creutzfeldt-Jakob disease[41]. Our results revealed that levels of 14-3-3 protein remained high even at 30 DPI, indicating that 14-3-3 plays an important role in the recovery of spinal cord injury.

Enhancing the 14-3-3/spastin interaction mediated by FC-A can promote neurite outgrowth and regeneration suggesting that it can be used to promote neurite regrowth after injury *in vivo*. We therefore attempted to investigate whether FC-A could improve nerve regrowth after spinal cord injury in a contusion injury model. FC-A was administrated via intraperitoneal injection following SCI in mice. As shown in figure 5D, H&E and LFB (Luxol fast blue stain) staining showed that the lesion site in the injury group presented the destroyed tissue structure with loss of myelination. Intriguingly, we found the tissue structure appeared to be normal and the area of demyelination in the lesion site strongly decreased in the FC-A group. In addition, in the spastazoline group, the lesion site appeared to be discontiguous, and application of spastazoline reversed the protective effect on spinal cord demyelination. Next, we further investigated nerve regeneration by immunostaining with NF (Neurofilament) and 5-HT, which were co-stained with GFAP (astrocyte marker). As shown in figure 5E, the border of the spinal cord lesion site was marked with astrogliosis (GFAP$^+$), and the neurofilament and 5-HT immunogenicity in the lesion site were almost abolished compared to the uninjured area, indicating that the spinal cord contusion injury model was successfully established. Interestingly, in the FC-A group, we observed that the astrogliosis in the lesion site was attenuated and we found a great deal of neurofilaments and 5-HT signals passing through the lesion site (Figure 5E, boxed in E2), indicating that the application of FC-A enhanced the nerve regeneration after injury. In the spastazoline group, we also observed tissue discontinuity, which is consistent with figure 5D, and almost no NF and 5-HT were found across the lesion site. Moreover, the application of spastazoline abolished the FC-A mediated nerve

regeneration after SCI. These results suggest that FC-A-mediated axonal regeneration

after SCI requires the activation of spastin function.

In order to further confirm whether the intervention of 14-3-3/spastin pathway by

FC-A and spastazoline mediates nerve regeneration after spinal cord injury by

targeting MTs at the site of spinal cord injury, acetylated tubulin and β-tubulin were

stained to label stale MTs (Lacking of dynamics) and total MTs. As shown in figure

S7 A, the intensity of acetylated microtubules was remarkable decreased in the lesion

site. Moreover, there was a clear boundary of acetylated MTs between the lesion site

and the uninjured area. This may be ascribed to the activation of cell proliferation in

the injured area after spinal cord injury which needs dynamic microtubules.

Interestingly, we found the boundary of acetylated microtubules near the damaged

area was unclear after application of FC-A, and the intensity of acetylated

microtubules significantly decreased compared with the injury group, indicating that

microtubules tended to be a dynamic state. However, after the application of

spastazoline, the proportion of acetylated microtubules in the injury site increased

significantly, and the boundary of acetylated microtubules in the injured area was

unclear, indicating that the application of spastazoline led to a significant increase in

the stability of the microtubules. In addition, the application of spastazoline can

significantly reverse the enhancement of microtubule dynamics mediated by FC-A

(Figure 5F). These experiments indicated that the administration of FC-A mediate

nerve regeneration after spinal cord injury through the 14-3-3/spastin pathway by

affecting the MTs dynamics. In addition, these evidences also indicated indirectly that

the FC-A and spastazoline can pass through the blood-brain barrier to reach the site of

spinal cord injury.

FC-A-mediated locomotor function improvement after SCI requires spastin

participation

Since FC-A promotes nerve regeneration after spinal cord injury by affecting the

14-3-3/spastin pathway, we then investigated whether it could mediate the recovery of

locomotor function after SCI. Therefore, we performed several experiments to verify

whether the application of FC-A could affect the locomotor function recovery after

SCI (Fig. 6A). Catwalk analysis showed that the maximum contact area of the mice in

the FC-A group was significantly larger than that in the injury group, and this

phenomenon was reversed after the application of spastazoline (Figure 6B&E). The

regularity index (Calculated as the number of normal strides multiplied by four and

then divided by the total number of strides; this value is ~100% in normal animals)

also had a similar phenomenon. The gait of the hindlimbs of mice in the FC-A group

was more coordinated than that of the mice in the injury group (Fig. 6C), and the

regularity index was statistically enhanced compared with the injury group (Fig. 6G).

This phenomenon was also significantly suppressed after the application of

spastazoline. We further evaluated the motor function after spinal cord injury by BMS

score (Fig. 6D). We found that the BMS scores of the mice in the FC-A group was

significantly improved compared with the mice in the injury group, and peaking after

35 days, while spastazoline reversed the improvement of locomotor function. At the

same time, we used the footprint test to analyze the motor function of the mice (Fig.

6F). The experimental data showed that the paw trailing phenomenon of the

hindlimbs in the FC-A group was significantly improved compared with the injury

group, and the stride length was significantly increased. This improvement was also

significantly suppressed by application of spastazoline (Fig. 6H). Finally, we

attempted to verify whether the spinal neural circuits in mice were remodeled by

examining the motor-evoked potentials (Stimulation at the sensorimotor cortex in the

M1 region and receive signals at the site of gastrocnemius). Our experimental results

showed that the MEP current amplitude decreased significantly after spinal cord

injury, while the current amplitude in the FC-A group significantly increased

compared with the injury group. And the improvement effect was significantly

weakened after the application of spastazoline, indicating that FC-A mediates the

remodeling of neural circuits after spinal cord injury by affecting the 14-3-3/spastin

pathway.

Enhancing the 14-3-3/spastin pathway by FC-A stimulates the repair of SCI in a

T10 lateral hemisection model

We have clarified the important role of 14-3-3/spastin pathway in the repair of spinal cord injury in the contusion model. In order to further observe the axon regeneration after spinal cord injury, we performed a spinal right lateral hemisection at the T10 level to eliminate the right side of axonal projections. Neurotransmission of serotonin (5-HT) in the spinal cord is required for the modulation of sensory, motor, and autonomic functions and can influence the response to spinal cord injury[42, 43]. In addition, studies have reported that the 5-HT fibers have strong regenerative capacity and involved in the regulation of neuronal activity[43]. Thus, we explored the 5-HT immunoactivity in spinal cord tissues isolated on 17 DPI. In the control SCI group, 5-HT-positive axons were significantly upregulated in the injury site, and to some extent in the rostral site. Interestingly, we found much stronger 5-HT staining in the lesion site of FC-A group and rostral site of injury compared with the SCI control group, indicating that the FC-A promoted the regeneration of 5-HT positive axons *in vivo*. However, few 5-HT signals were found in the lesion site, also in the rostral site in the spastazoline group. In addition, the FC-A promoting effects of 5-HT axonal regeneration were significantly weakened by the combined application of spastazoline (Figure 7A&C).

We further investigated the immunoactivity of neurofilament and myelin basic protein (MBP) in the white matter of the lesion site to evaluate axonal regeneration and the extend of remyelination. Neurofilament are cytoskeletal proteins that are expressed abundantly in the cytoplasm of axonal fibers in the CNS[44]. The MBP forms and maintains the structure of the compact myelin sheath which then regulates axonal function[45]. Our results revealed that FC-A could enhance the regeneration of neurofilaments which were largely colocalized with MBP, whereas spastazoline alleviated this effect (Figure 7B&D). Collectively, these results suggest that FC-A promotes axonal regeneration which is dependent on activation of spastin microtubule severing.

Previous studies have reported that the proliferation of neural stem/progenitor

cells is also reactivated and contributes to synapse formation following spinal cord injury[46, 47]. Since spastin is a microtubule severing enzyme involved in cell division. Thus, we explored the possibility that whether 14-3-3/spastin complex participates in proliferation process of neural stem/progenitor cells. Therefore, spinal cord tissues from the lesion site were stained with Nestin, Brdu and NeuN. Intriguingly, we found that FC-A treatment after SCI significantly increased the expression of Nestin (Figure S9A-B) and Brdu (Figure S10), and these effects were abolished by spastazoline treatment. These results demonstrated that the 14-3-3/spastin complex contributed to the proliferation of neural stem/progenitor cells thereby improving recovery after spinal cord injury.

We also further examined whether FC-A administration altered the motor function after SCI in this lateral hemisection injury model. Mice with SCI in all groups exhibited paralysis in the right hindlimb after hemisecion which confirmed the successful establishment of the lateral hemisection SCI model. For mice in SCI control group, the BMS scores gradually improved to ~4 by 16 DPI. In contrast, the scores of mice in the FC-A group indicated an improvement in locomotor function which was impaired in the spastazoline group. Moreover, the improvement in locomotor function following administration of FC-A diminished when spastazoline was administered (Figure 7E). We further analyzed the footprint of mice with SCI and calculated the length or width of strides (Figure S11). We found that stride length improved in mice of the FC-A group but it was reduced in the spastazoline group although not significantly. The improvement induced by FC-A diminished when co-treated with spastazoline (Figure 7G). Notably, there was no significant difference in stride width between the groups (Figure 7H). Similar effects were observed in the foot fault scan experiment. The faults of right hindlimb significantly decreased in the FC-A group and increased in the spastazoline group (Figure 7F). The improvement of hindlimb fault in mice administered with FC-A diminished following the application of spastazoline. The representative movements of mice in each group were recorded in Movie S1 (Control group), Movie S2 (Injury group), Movie S3 (FC-A group),

Movie S4 (Spastazoline group) and Movie S5 (FC-A+Spastazoline group). We also explored whether the stability of MTs was altered by administration of FC-A and spastazoline in the spinal cord injury model. We found that the ratio of acetylated tubulin to total tubulin was significantly increased in the SCI group, decreased in the FC-A group, and upregulated in the spastazoline group, which further confirmed the successful delivery of drugs to the lesion site (Figure S8). These findings suggested that FC-A was effectively delivered to the spinal cord tissue and improved the locomotor function of mice with SCI, through 14-3-3/spastin pathway activation mechanism.

Discussion

The limited cell intrinsic regrowth capability at the injury site of CNS damage or disease results in failure of axonal regeneration. Recent studies have demonstrated that the remodeling of microtubule dynamics was necessary for promoting axon regeneration[48, 49]. Spastin is a microtubule (MT) severing enzyme which cuts long MTs into short fragments for local MTs remodeling and thus could be administrated to promote axon regeneration[50]. In this study, we aimed to reveal the regulatory molecular mechanisms of spastin and its role in the recovery of spinal cord injury. We found that 14-3-3 interacted with spastin by targeting the phosphorylation of Ser233, protecting spastin against ubiquitin-mediated degradation, thus up-regulated the spastin-dependent microtubule severing activity. Moreover, enhancing the 14-3-3/spastin interaction by FC-A promoted the neurite outgrowth and regeneration. In adult mice with of SCI (both in contusion and lateral hemisection spinal cord injury model), we found that enhanced 14-3-3/spastin binding promoted the recovery of locomotor function and axonal regeneration. Therefore, we postulated that the 14-3-3/spastin pathway is an attractive target for improving axonal regeneration and locomotor recovery after SCI.

To date, the severing activity induced by spastin has been well characterized[51, 52] and reported to be essential in the formation of axon branches and axonal

extension[53, 54]. Studies have also shown that spastin regulates axon regeneration in

a dose dependent manner. Of note, during regeneration, the cytoskeleton undergoes

remodeling, suggesting that manipulation of MTs dynamics by spastin may be a

potential approach for controlling neural regeneration. However, efficient

regeneration is dependent on the precise spastin dosage. Therefore, understanding the

underlying regulatory mechanism of spastin is pivotal for identifying therapeutic

targets in CNS. In this study, LC-MS based proteomics analysis was performed and

proteins that bind spastin were screened to investigate the regulatory mechanisms of

spastin during CNS damage (Figure 1A&S1). Among the identified binding proteins,

14-3-3 proteins had high scores and was found to interact with spastin in the spinal

cord (Figure 1B). This revealed the binding of 14-3-3 proteins to spastin in the injured

spinal cord. Of note, MS data from a previous study also showed that the spastin

peptides were present in the binding peptides with 14-3-3 agonist Fusicoccin[50],

which confirmed the important relevance between these two proteins. Considered that

both 14-3-3 and spastin were found to regulate axon outgrowth and regeneration,

colocalization analysis were performed. The results indicated that 14-3-3 interacted

with spastin to affect axon development. In addition, GST pulldown and

co-immunoprecipitation assays further verified the interaction between all 14-3-3

isoforms and spastin (Figure 1F-H). Also, the binding between 14-3-3 and spastin was

further characterized which showed that Ser233 in spastin was responsible for the

binding, and could directly influence the fate of neurite outgrowth (Figure 2).

Previously, it was found that spastin was phosphorylated at Ser233 (human Ser268)

by HIPK2 and this process was vital for successful abscission and the loss of HIPK2

exhibited a protective effect on neurons[55]. Our western blotting results showed that

HIPK2 was expressed at 9th DIV in the brain tissue (not presented), suggesting that

there may be other kinases that regulate Ser233 phosphorylation during neural

development. Therefore, we have well established the model of the 14-3-3/spastin

binding which could be the intervention target for regulating the MT severing

mediated by spastin.

The regulatory mechanisms that control the MT severing activity mediated by spastin are largely unknown. In our previous study, spastin protein expression levels and its severing effect were regulated at post-transcriptional levels, such as micoRNA-30[53] and SUMOylation[14]. In this study, we found that 14-3-3 interacted with spastin from the perspective of protein-protein interactions, and we showed that 14-3-3 prevented the ubiquitin-mediated degradation of spastin by inhibiting the ubuiquitin pathway and phosphorylation of Ser233 was crucial to this process. Our results showed that Ser233 de-phosphorylation lowered the protein stability, consistent with findings from a recent study[33], and our study further illustrated the molecular mechanisms of this phenomenon. In addition, we observed that 14-3-3 up-regulated the microtubule severing activity of spastin through interacting with its phosphorylation at Ser233. Based on this mechanism, we proposed a model in which Ser233 is phosphorylated during the binding of 14-3-3 to spastin, thereby disrupting the spastin/ubiquitin binding, ultimately affecting the stability of spastin and subsequent neurite outgrowth and regeneration. Similar mechanisms were reported for doublecortin[56], cdt2[57], p21[58] and other proteins[59]. For example, 14-3-3ε was found to interact with threonine 42 in doublecortin which protected it from degradation by inhibiting its ubiquitination. In this way, it regulated the doublecortin-regulated microtubule stability and neurite initiation process[57]. In this part, we validated that 14-3-3 stabilized the protein levels of spastin and enhanced the MT severing activity mediated by spastin which was similar to the mechanism of 14-3-3/cdt2 binding.

Since 14-3-3 proteins are adaptor proteins that are highly expressed in neural tissues, they are important regulators of neural development[60, 61]. Using a well-established 14-3-3/spastin interaction model, we found that promoting the MT dynamics by enhancing the interaction of 14-3-3/spastin could enhance neurite outgrowth and regeneration. This accelerated neurite outgrowth and repair following FC-A application by facilitating the binding of 14-3-3/spastin (Figure 4). In accordance with previous findings, 14-3-3 proteins can promote neurite outgrowth

and axonal regeneration[34]. In our study, we further demonstrated that enhanced 14-3-3/spastin interaction by FC-A favored neurite outgrowth and regeneration, but this effect was blunted by spastazoline treatment. Spastazoline was designed to compete with the AAA domain of spastin and this effectively abolished the severing activity mediated by spastin[37]. The results also demonstrated that the normal severing activity mediated by spastin was required for neurite outgrowth and regeneration (Figure 4). These findings open a new window for developing spastin-based agents for accelerating axon regeneration after injury.

So far, few studies have investigated the relationship between spastin and neural regeneration in CNS diseases, such as spinal cord injury or brain injury. However, many studies have elucidated its indispensable role in the process of axon regeneration[50]. Our study first indicate that functional spastin is required for axon regeneration and locomotor recovery after spinal cord injury (Figure 5&6). Moreover, 14-3-3 was upregulated at 3 and 7 DPI, and was even higher at 30 DPI in the neuronal compartment, suggesting that 14-3-3 has an important role in neural repair after spinal cord injury (Figure 5). Using the model of 14-3-3/spastin interaction, we further elucidated the mechanism by which FC-A administration regulates the recovery of spinal cord injury by targeting MTs. Fusicoccin-A (FC-A) is a fusicoccane, a small molecule produced by the *Phomopsis amygdali fungus* that stabilizes the binding of 14-3-3 to its substrates[62]. The crystal structure of FC-A bound to 14-3-3 has already revealed the efficiency of the enhancement of FC-A in the interaction between 14-3-3 and their substrate[63]. In addition, FC-A has been reported to promote axonal regeneration in mammals[34], although the mechanisms are rarely elusive. In this study, we found that FC-A enhanced the 14-3-3/spastin interaction and FC-A administration restored nerve regeneration and locomotor recovery after SCI (in both contusion and lateral hemisection spinal cord injury model), but these improvements were abolished following spastazoline administration (inhibiting the MT severing by direct competing with AAA domain of spastin[37]), suggesting the important role of 14-3-3/pathway in the neural circuts remolding after SCI. This phenomenon was

further confirmed by examined the ratio of acetylated tubulin in the lesion site of SCI,
and our results showed the intervention of 14-3-3/spastin pathway by FC-A and
spastazoline could significantly affect the MTs dynamics in the lesion site which
indirectly demonstrated the efficient delivery of these drugs to the spinal cord across
the blood brain barrier. Since spastin plays an essential role in cell abscission, and the
activation of 14-3-3/spastin pathway by FC-A promoted the proliferation of neural
stem/progenitor cells (Figure S8&S9) thereby support axonal regeneration. These
results also support that promoting 14-3-3/spastin interaction by FC-A enhanced
axonal regeneration after spinal cord injury, and the microtubule severing independent
of spastin was important in this process. Therefore, our findings present
14-3-3/spastin pathway as a novel intervention for accelerating nerve regeneration
after spinal cord injury.

In summary, we proposed the 14-3-3/spastin binding model for manipulating
microtubule dynamics and can be an effective intervention for promoting recovery of
spinal cord injury. In addition, our results indicate that 14-3-3 and spastin are
important regulators for nerve regeneration and modulating the interaction between
14-3-3 and spastin may be an effective strategy for the treatment of spinal cord injury.
750
751

Abbreviations
5-HT, 5-hydroxytryptamine; Cdt2, cell division cycle protein; CNS, central neuronal
system; DPI, days post injury; ECL, enhanced chemiluminescence; FOXO, forkhead
box transcription factors; GFAP, glial fibrillary acidic protein; GFP, green fluorescent
protein; HIPK2, homeodomain interacting protein kinase 2; HSP, hereditary spastic
paraplegia; IP, Immunoprecipitation; MBP, Myelin basic protein; MS, mass
spectrometry; MT, microtubule; ROS, reactive oxygen species;
759

760 **Acknowledgement**
761   This work was supported by the Natural Science Foundation of China (grant nos.

82102314, 31900691, and 32170977), the Natural Science Foundation of Guangdong Province (grant nos. 2022A1515010438 and 2022A1515012306), Science and Technology Projects in Guangzhou (grant nos. 202201020018) and Project funded by China Postdoctoral Science Foundation (2023M731320).

Competing interests

The authors declare no conflict of interest.

Data availability

The datasets obtained and/or analysed in the current study are available from the corresponding author upon reasonable request.

Reference

1. Curcio M, Bradke F. Axon Regeneration in the Central Nervous System: Facing the Challenges from the Inside. Annual review of cell and developmental biology. 2018; 34: 495-521.

2. Hilton BJ, Husch A, Schaffran B, Lin TC, Burnside ER, Dupraz S, et al. An active vesicle priming machinery suppresses axon regeneration upon adult CNS injury. Neuron. 2022; 110: 51-69.e7.

3. Mahar M, Cavalli V. Intrinsic mechanisms of neuronal axon regeneration. Nature reviews Neuroscience. 2018; 19: 323-37.

4. Clark CEJ, Liu Y, Cooper HM. The Yin and Yang of Wnt/Ryk axon guidance in development and regeneration. Science China Life Sciences. 2014; 57: 366-71.

5. Jin Y. Unraveling the mechanisms of synapse formation and axon regeneration: the awesome power of C. elegans genetics. Science China Life Sciences. 2015; 58: 1084-8.

6. Pinto-Costa R, Sousa SC, Leite SC, Nogueira-Rodrigues J, Ferreira da Silva T, Machado D, et al. Profilin 1 delivery tunes cytoskeletal dynamics toward CNS axon regeneration. The Journal of clinical investigation. 2020; 130: 2024-40.

7. Crunkhorn S. CNS injury: Microtubule stabilizer repairs spinal cord injury. Nature reviews Drug discovery. 2015; 14: 310.

8. Chen L, Rolls MM. Microtubule deacetylation sets the stage for successful axon regeneration. The

EMBO journal. 2012; 31: 3033-5.

9.   Vargas EJM, Matamoros AJ, Qiu J, Jan CH, Wang Q, Gorczyca D, et al. The microtubule

regulator ringer functions downstream from the RNA repair/splicing pathway to promote axon

regeneration. Genes & development. 2020; 34: 194-208.

10.   Shribman S, Reid E, Crosby AH, Houlden H, Warner TT. Hereditary spastic paraplegia: from

diagnosis to emerging therapeutic approaches. The Lancet Neurology. 2019; 18: 1136-46.

11.   Liu Q, Zhang G, Ji Z, Lin H. Molecular and cellular mechanisms of spastin in neural development

and disease (Review). International journal of molecular medicine. 2021; 48.

12.   Salinas S, Carazo-Salas RE, Proukakis C, Schiavo G, Warner TT. Spastin and microtubules:

Functions in health and disease. J Neurosci Res. 2007; 85: 2778-82.

13.   Lopes AT, Hausrat TJ, Heisler FF, Gromova KV, Lombino FL, Fischer T, et al. Spastin depletion

increases tubulin polyglutamylation and impairs kinesin-mediated neuronal transport, leading to

working and associative memory deficits. PLoS biology. 2020; 18: e3000820.

14.   Ji ZS, Liu QL, Zhang JF, Yang YH, Li J, Zhang GW, et al. SUMOylation of spastin promotes the

internalization of GluA1 and regulates dendritic spine morphology by targeting microtubule dynamics.

Neurobiology of disease. 2020; 146: 105133.

15.   Stone MC, Rao K, Gheres KW, Kim S, Tao J, La Rochelle C, et al. Normal spastin gene dosage is

specifically required for axon regeneration. Cell reports. 2012; 2: 1340-50.

16.   Rao K, Stone MC, Weiner AT, Gheres KW, Zhou C, Deitcher DL, et al. Spastin, atlastin, and ER

relocalization are involved in axon but not dendrite regeneration. Molecular biology of the cell. 2016;

27: 3245-56.

17.   Martins-Marques T, Ribeiro-Rodrigues T, Batista-Almeida D, Aasen T, Kwak BR, Girao H.

Biological Functions of Connexin43 Beyond Intercellular Communication. Trends in cell biology. 2019;

29: 835-47.

18.   Fu W, Hu W, Yi YS, Hettinghouse A, Sun G, Bi Y, et al. TNFR2/14-3-3ε signaling complex

instructs macrophage plasticity in inflammation and autoimmunity. The Journal of clinical investigation.

2021; 131.

19.   Fu W, Hettinghouse A, Chen Y, Hu W, Ding X, Chen M, et al. 14-3-3 epsilon is an intracellular

component of TNFR2 receptor complex and its activation protects against osteoarthritis. Annals of the

rheumatic diseases. 2021; 80: 1615-27.

20.   Nielsen MD, Luo X, Biteau B, Syverson K, Jasper H. 14-3-3 Epsilon antagonizes FoxO to control

growth, apoptosis and longevity in Drosophila. Aging cell. 2008; 7: 688-99.

21.   Park E, Rawson S, Li K, Kim BW, Ficarro SB, Pino GG, et al. Architecture of autoinhibited and

active BRAF-MEK1-14-3-3 complexes. Nature. 2019; 575: 545-50.

22.   Pisciottani A, Biancolillo L, Ferrara M, Valente D, Sardina F, Monteonofrio L, et al. HIPK2

Phosphorylates the Microtubule-Severing Enzyme Spastin at S268 for Abscission. Cells. 2019; 8.

23.   Kaplan A, Morquette B, Kroner A, Leong S, Madwar C, Sanz R, et al. Small-Molecule

Stabilization of 14-3-3 Protein-Protein Interactions Stimulates Axon Regeneration. Neuron. 2017; 93:

1082-93 e5.

24.   Yam PT, Kent CB, Morin S, Farmer WT, Alchini R, Lepelletier L, et al. 14-3-3 proteins regulate a

cell-intrinsic switch from sonic hedgehog-mediated commissural axon attraction to repulsion after

midline crossing. Neuron. 2012; 76: 735-49.

25.   Ji Z, Zhang G, Chen L, Li J, Yang Y, Cha C, et al. Spastin Interacts with CRMP5 to Promote

Neurite Outgrowth by Controlling the Microtubule Dynamics. Developmental neurobiology. 2018; 78:

1191-205.

26.   Ji ZS, Li JP, Fu CH, Luo JX, Yang H, Zhang GW, et al. Spastin interacts with collapsin response

mediator protein 3 to regulate neurite growth and branching. Neural regeneration research. 2021; 16:

2549-56.

27.   Patel M, Li Y, Anderson J, Castro-Pedrido S, Skinner R, Lei S, et al. Gsx1 promotes locomotor

functional recovery after spinal cord injury. Molecular therapy : the journal of the American Society of

Gene Therapy. 2021; 29: 2469-82.

28.   Liu Y, Wang X, Li W, Zhang Q, Li Y, Zhang Z, et al. A Sensitized IGF1 Treatment Restores

Corticospinal Axon-Dependent Functions. Neuron. 2017; 95: 817-33.e4.

29.   Brown AR, Martinez M. Ipsilesional Motor Cortex Plasticity Participates in Spontaneous

Hindlimb Recovery after Lateral Hemisection of the Thoracic Spinal Cord in the Rat. The Journal of

neuroscience : the official journal of the Society for Neuroscience. 2018; 38: 9977-88.

30.   Sun G, Yang S, Cao G, Wang Q, Hao J, Wen Q, et al. γδ T cells provide the early source of IFN-γ

to aggravate lesions in spinal cord injury. The Journal of experimental medicine. 2018; 215: 521-35.

31. Zhou K, Zheng Z, Li Y, Han W, Zhang J, Mao Y, et al. TFE3, a potential therapeutic target for

Spinal Cord Injury via augmenting autophagy flux and alleviating ER stress. Theranostics. 2020; 10:

9280-302.

32. Kamens HM, Crabbe JC. The parallel rod floor test: a measure of ataxia in mice. Nature protocols.

2007; 2: 277-81.

33. Sardina F, Pisciottani A, Ferrara M, Valente D, Casella M, Crescenzi M, et al. Spastin recovery in

hereditary spastic paraplegia by preventing neddylation-dependent degradation. Life science alliance.

2020; 3.

34. Kaplan A, Morquette B, Kroner A, Leong S, Madwar C, Sanz R, et al. Small-Molecule

Stabilization of 14-3-3 Protein-Protein Interactions Stimulates Axon Regeneration. Neuron. 2017; 93:

1082-93.e5.

35. Saponaro A, Porro A, Chaves-Sanjuan A, Nardini M, Rauh O, Thiel G, et al. Fusicoccin Activates

KAT1 Channels by Stabilizing Their Interaction with 14-3-3 Proteins. The Plant cell. 2017; 29:

2570-80.

36. Petosa C, Masters SC, Bankston LA, Pohl J, Wang B, Fu H, et al. 14-3-3zeta binds a

phosphorylated Raf peptide and an unphosphorylated peptide via its conserved amphipathic groove.

The Journal of biological chemistry. 1998; 273: 16305-10.

37. Pisa R, Cupido T, Kapoor TM. Designing Allele-Specific Inhibitors of Spastin, a

Microtubule-Severing AAA Protein. Journal of the American Chemical Society. 2019; 141: 5602-6.

38. Pisa R, Cupido T, Steinman JB, Jones NH, Kapoor TM. Analyzing Resistance to Design Selective

Chemical Inhibitors for AAA Proteins. Cell chemical biology. 2019; 26: 1263-73.e5.

39. Savolainen KM, Loikkanen J, Eerikäinen S, Naarala J. Glutamate-stimulated ROS production in

neuronal cultures: interactions with lead and the cholinergic system. Neurotoxicology. 1998; 19:

669-74.

40. Kajiwara Y, Buxbaum JD, Grice DE. SLITRK1 binds 14-3-3 and regulates neurite outgrowth in a

phosphorylation-dependent manner. Biological psychiatry. 2009; 66: 918-25.

41. Zerr I. Laboratory Diagnosis of Creutzfeldt-Jakob Disease. The New England journal of medicine.

2022; 386: 1345-50.

42. van den Brand R, Heutschi J, Barraud Q, DiGiovanna J, Bartholdi K, Huerlimann M, et al.

Restoring voluntary control of locomotion after paralyzing spinal cord injury. Science (New York, NY).

2012; 336: 1182-5.

43.   Perrin FE, Noristani HN. Serotonergic mechanisms in spinal cord injury. Experimental neurology.

2019; 318: 174-91.

44.   Yuan A, Rao MV, Veeranna, Nixon RA. Neurofilaments and Neurofilament Proteins in Health and

Disease. Cold Spring Harbor perspectives in biology. 2017; 9.

45.   Liu B, Xin W, Tan JR, Zhu RP, Li T, Wang D, et al. Myelin sheath structure and regeneration in

peripheral nerve injury repair. Proceedings of the National Academy of Sciences of the United States of

America. 2019; 116: 22347-52.

46.   Ceto S, Sekiguchi KJ, Takashima Y, Nimmerjahn A, Tuszynski MH. Neural Stem Cell Grafts

Form Extensive Synaptic Networks that Integrate with Host Circuits after Spinal Cord Injury. Cell stem

cell. 2020; 27: 430-40.e5.

47.   Venkatesh K, Ghosh SK, Mullick M, Manivasagam G, Sen D. Spinal cord injury:

pathophysiology, treatment strategies, associated challenges, and future implications. Cell and tissue

research. 2019; 377: 125-51.

48.   Ko SH, Apple EC, Liu Z, Chen L. Age-dependent autophagy induction after injury promotes axon

regeneration by limiting NOTCH. Autophagy. 2020; 16: 2052-68.

49.   Griffin JM, Bradke F. Therapeutic repair for spinal cord injury: combinatory approaches to

address a multifaceted problem. EMBO molecular medicine. 2020; 12: e11505.

50.   Lin YF, Xie Z, Zhou J, Chen HH, Shao WW, Lin HD. Effect of exogenous spastin combined with

polyethylene glycol on sciatic nerve injury. Neural regeneration research. 2019; 14: 1271-9.

51.   Sarbanes SL, Zehr EA, Roll-Mecak A. Microtubule-severing enzymes. Current biology : CB.

2022; 32: R992-r7.

52.   Vemu A, Szczesna E, Zehr EA, Spector JO, Grigorieff N, Deaconescu AM, et al. Severing

enzymes amplify microtubule arrays through lattice GTP-tubulin incorporation. Science (New York,

NY). 2018; 361.

53.   Jiang T, Cai Z, Ji Z, Zou J, Liang Z, Zhang G, et al. The lncRNA MALAT1/miR-30/Spastin Axis

Regulates Hippocampal Neurite Outgrowth. Frontiers in cellular neuroscience. 2020; 14: 555747.

54.   Costa AC, Sousa MM. The Role of Spastin in Axon Biology. Frontiers in cell and developmental

biology. 2022; 10: 934522.

55.   Zhang J, Pho V, Bonasera SJ, Holtzman J, Tang AT, Hellmuth J, et al. Essential function of HIPK2

in TGFbeta-dependent survival of midbrain dopamine neurons. Nature neuroscience. 2007; 10: 77-86.

56.   Cornell B, Wachi T, Zhukarev V, Toyo-Oka K. Regulation of neuronal morphogenesis by

14-3-3epsilon (Ywhae) via the microtubule binding protein, doublecortin. Human molecular genetics.

2016; 25: 4405-18.

57.   Dar A, Wu D, Lee N, Shibata E, Dutta A. 14-3-3 proteins play a role in the cell cycle by shielding

cdt2 from ubiquitin-mediated degradation. Molecular and cellular biology. 2014; 34: 4049-61.

58.   Lee JH, Lu H. 14-3-3Gamma inhibition of MDMX-mediated p21 turnover independent of p53.

The Journal of biological chemistry. 2011; 286: 5136-42.

59.   Chan TY, Egbert CM, Maxson JE, Siddiqui A, Larsen LJ, Kohler K, et al. TNK1 is a

ubiquitin-binding and 14-3-3-regulated kinase that can be targeted to block tumor growth. Nature

communications. 2021; 12: 5337.

60.   Kast DJ, Dominguez R. Mechanism of IRSp53 inhibition by 14-3-3. Nature communications.

2019; 10: 483.

61.   Cornell B, Toyo-Oka K. 14-3-3 Proteins in Brain Development: Neurogenesis, Neuronal

Migration and Neuromorphogenesis. Frontiers in molecular neuroscience. 2017; 10: 318.

62.   Marra M, Camoni L, Visconti S, Fiorillo A, Evidente A. The Surprising Story of Fusicoccin: A

Wilt-Inducing Phytotoxin, a Tool in Plant Physiology and a 14-3-3-Targeted Drug. Biomolecules. 2021;

11.

63.   Liu D, Bienkowska J, Petosa C, Collier RJ, Fu H, Liddington R. Crystal structure of the zeta

isoform of the 14-3-3 protein. Nature. 1995; 376: 191-4.

Figures and legends:

Figure 1. 14-3-3 physically interacts with spastin in vivo and vitro. (A) Mass spectrum of peptides which was pull-downed by GST-spastin from proteins of the spinal cord tissue (near 1 cm around the T10 level), the representative peptides of 14-3-3 were shown. (B) Immunoprecipitation assay using spastin antibody was performed and Pan 14-3-3 antibody was used for western blotting. (C) Hippocampal

neurons were stained with 14-3-3s (green) and spastin (red) antibodies, the growth

cone (C1) and axon shaft (C2) profile were shown. (D) Localization of spastin (red

line) and 14-3-3 (green line) protein in the hippocampal neurons were analyzed and

the intensity of the white dotted line across the neurite compartment were measured.

(E) Purified GST and GST-14-3-3s fusion proteins were subjected to sodium

dodecyl-sulfate polyacrylamide gel electrophoresis and coomassie blue staining was

confirmed the successful purification of relative proteins. (F) GST fusion proteins

were used to pull down the lysates of the spinal cord and spastin antibody was used

for western blotting. (G) HEK 293T cells were transfected with Flag tagged 14-3-3s

and GFP tagged spastin, the cell lysates were subjected to GFP-Trap assay, Flag

antibody was used to detect the presence of 14-3-3s.

Figure 2. Ser233 in spastin is essential for the interaction with 14-3-3 and important for neurite outgrowth. (A) Schematic illustration of the gene truncation strategy for the modular domain of spastin. (B) Truncation mutants of spastin together with Flag tagged 14-3-3 were transfected into HEK 293T cells, and cell lysates were immunoprecipitated by GFP-Trap assay. Flag antibody were used for detecting the

flag tagged 14-3-3 fusion protein. (C) The microtubule binding domain (MTBD) truncated GFP tagged spastin were transfected together with Flag tagged 14-3-3 into HEK 293T cells, and subjected to GFP trap assay. (D) HEK 293T cells were transfected with GFP-spastin and then treated with or without staurosporine for 1 hour. Cell lysates were pull down by GST 14-3-3 fusion protein and GFP antibody were used for western blotting. (E) The binding sites between 14-3-3 and spastin were predicted using the scansite 2.0 software and whether these sites could be phosphorylated were confirmed by phosphosite plus website. (F) Serine 233 and Serine 562 of spastin were mutated to alanine, these GFP fusion mutations were transfected together with Flag tagged 14-3-3, then cell lysates were subjected to GFP-trap assay. Flag antibody was used to detect the binding ability between spastin mutations and 14-3-3. (G) Hippocampal neurons were grown on DIV 2 and transfected with spastin Ser233 mutations (Spastin S233A, Spastin S233D), then fixed on DIV 3. Representative images of transfected neurons as indicated were shown. Quantitative analysis of the length of neurites (H) and the number of neurites (I) were shown (n≥25 per group). All error bars are SEM. *P <0.05. Scale bar: 100 μm.

Figure 3. 14-3-3 protein protects spastin by Ser233 phosphorylation from degradation through ubiquitin pathway. (A) GFP tagged spastin together with Flag tagged 14-3-3 (B) or without was transfected into HEK 293T cells. Following the

addition of protein synthesis inhibitor (cycloheximide, CHX), cell lysates were subjected to western blot to detect the GFP tagged spastin protein levels. (C, D) Western blot analysis of HEK 293T cell lysates transfected with spastin mutations (S233A or S233D) after treating with cycloheximide at different indicated times. (E) Quantitative analysis of spastin protein levels after treating by cycloheximide with or without 14-3-3 protein. (F) Quantitative analysis of GFP tagged spastin protein levels upon transfected with different spastin mutations. (G) HEK 293T cells were transfected with GFP tagged spastin and Flag tagged ubiquitin, together with R18 or not, cells were harvested and subjected to immunoprecipitation using GFP antibody. Flag antibody was used to detect the ubiquitin band. (H) COS7 cells were transfected with indicated GFP tagged spastin with or without Flag tagged ubiquitin for 24 hours. Then, cells were fixed and stained with tubulin antibody to visualize microtubule arrangements. (I) The normalized quantitative analysis of microtubule fluorescence intensity among indicated groups (n≥20 pictures per group). $*P < 0.05$, $**P < 0.01$, $***P < 0.001$. Scale bar: 50 μm, 10 μm.

Figure 4. 14-3-3 agonist Fusicoccin (FC-A) promotes the repair of neurite outgrowth and regeneration after injury via spastin. (A) The surface structure and (B) secondary structure of molecular docking model between 14-3-3 protein with FC-A (PDB:2O98) or R18 (PDB:1A38). (C) HEK 293T cells were transfected with GFP tagged spastin and Flag tagged 14-3-3, cell lysates were incubated with FC-A or R18 and then subjected to GFP-trap assay. Flag antibody were used to detect the 14-3-3 binding. (C) Hippocampal neurons (DIV2) were transfected with GFP to visualize its morphology. Neurons were then incubated with glutamate (120 μM) to induce injury, the neurite breakages and swellings were found in C1 and C2. (D) The injured neurons were transfected with spastin and incubated with FC-A, R18 or spastazoline for 24 hours, then cells were fixed and images were taken by confocal microscopy. Neurons were traced using Image J Pro Plus and quantitative analysis of the length of primary neurites (E), secondary neurites (F) and total neurites (G) were performed. Quantitative analysis of the number of primary neurites (H), secondary neurites (I) and total neurites (J). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. Scale bar: 50 μm. (K) Primary cortical neurons were grown and scratch assays were performed at DIV7. FC-A or spastazoline were applied and incubated for 36 hours. Then, cells were stained with βIII-tubulin to visualize the morphology of neurites. (L) Quantitative analysis of the normalized axon regeneration rate. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. Scale bar: 100 μm.

1013

1014

Figure 5. 14-3-3/spastin pathway is involved in nerve regeneration after spinal cord injury by targeting MTs. (A) After spinal cord contusion, the spinal cord tissues of the lesion site (near 1 cm) at indicated times (1, 3, 7 14, and 30 DPI) were ground and lysed, then subjected to western blot. (B) Quantitative analysis of the 14-3-3 protein expression (n=3 per group). (C) The lesion site of spinal tissue in the sham group and the SCI group were stained with Pan 14-3-3 (red) and β III tubulin (green). The arrows mean the elevated 14-3-3 protein in the neuronal compartment. (D) 14-3-3 agnoist FC-A and spastin inhibitor spastazoline were admisnistrated after spinal cord contusion. At 45 DPI (Days post-injury), the tissues were fixed and embedded, then cut horizontally and longitudinally. H&E and LFB (Luxol fast blue stain) were stained. The spinal cord demyelination was shown inside the area of the dotted line. Scale bar，200 μm. (E) The slices were then subjected to immunofluorescence staining, and GFAP (green) was stained to label astrocytes, NF (red) was stained to label neurons and 5-HT (magenta) was stained to label monoaminergic axons. The lesion site in different groups was boxed and enlarged in (E1)(E2)(E3)(E4). Scale bar, 400 μm, and 100 μm. (F) The spinal cord slices were stained with acetylated tubulin (Stable MTs which lack of dynamics) and β -tubulin (total tubulin). Scale bar，50 μm. (G) The normalized intensity of NF was quantified (n=5 animals per group). Mean ± SEM. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. (H) The normalized ratio of acetylation tubulin to total tubulin was calculated by the intensity value of acetylated tubulin divided by total tubulin (n=5 pictures per group). Mean ± SEM. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Figure 6. 14-3-3/spastin pathway coordinates locomotor recovery after spinal cord contusion. (A) Schematic illustration of relative treatments and examinations at different indicated times. (B) The max contact areas of the right hindlimb were recorded and analyzed by the Catwalk software. (C) The footfall patterns were visualized by the Catwalk software. The blue dot represents the start of a pattern. The

red dot represents the gait was not part of a pattern. The yellow dot represents that the

gait was not taken into account. The green one means the gait belongs to a normal

part of a pattern. (D) BMS score at different indicated days after injury and data were

represented as mean±SEM. (n=5 group, data were analyzed by one way ANOVA and

Bonferroni's multiple-comparison test were used for post hoc comparisons. (E) The

quantitative analysis of the hindlimb (right) max contact area. (n=5 animals per

group). (F) Footprint analysis of the mice at 6 weeks after spinal cord contusion.

The forelimbs were marked with blue and the hindlimbs were marked with red. Scale

bar: 2.5 cm. (G) The normalized regularity index of the footfall pattern calculated by

Catwalk software (~100 % in the sham group, n=5 animals per group). (H)

Quantatitive analysis of the stride length in the footprint assay. The mean stride length

of each walk was counted, n=5 animals per group). (I) Motor evoked potentials

(MEPs) were recorded at 7 weeks after spinal cord contusion. The craniotomy was

performed and the M1 region of the sensorimotor cortex were stimulated by single

square-wave stimulus of 0.5 mA, 0.5 ms duration, 2 ms time delay, and 1 Hz. The

MEP were recorded with the signals detected by the electrode on the gastrocnemius

muscle. The representative histograms of the amplitude-times are shown. The arrows

indicated the stimulus. (J) The current amplitudes of the MEP were quantified, n=5

animals per group. Mean $\pm$ SEM. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

1062

1063

Figure 7. FC-A promotes axon regeneration in a T10 lateral hemisection spinal cord injury model. (A) The spinal cords of adult mice were subjected to lateral hemisection and administrated by intraperitoneal injection with FC-A or spastazoline. Sagittal view of 5-HT-immunoreactive raphespinal fibers in the spinal cord on 17 DPI. Scale bar, 500 μm and 100 μm. (B) Sagittal view of neurofilaments which was MBP positive in the white manner of the lesion site of the spinal cord. Scale bar, 50 μm. (C) The normalized quantification of 5-HT immunoreactive fluorescence intensity (0.5 mm caudal to the lesion site, n=5 animals per group). (D) The normalized quantification of NF (Neurofilament) immunoreactive fluorescence intensity in the lesion site (n=5 animals per group). (E) The locomotor function of mice after SCI were scored at the indicated time points according to the standard of Basso Mouse Scale (BMS). (F) Foot fault test was performed and the total steps and steps drop out of the right hindlimb were recorded. The rate of the foot errors was analyzed, n=5 animals per group. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$. (G) (H)The footprint assay was performed and stride length (G) and stride width (H) quantitative analyzed. Data were presented as mean $\pm$ SEM, n=5 animals per group, the mean value of each animal's walk were calculated.