The 14-3-3/spastin pathway coordinates neurite regeneration and

recovery of spinal cord injury by targeting microtubule dynamics

Author: Qiuling Liu[1#], Hua Yang[1#], Jianxian Luo[1#], Cheng Peng[1], Ke Wang[1], Guowei

Zhang[1], Zhisheng Ji[1]*, Hongsheng Lin[1]*

1 Department of Orthopedics, The First Affiliated Hospital of Jinan University,

Guangdong province, Guangzhou 510630, People's Republic of China;

[#]These authors contribute equally to this work.

* Corresponding to:

Zhisheng Ji (jizhisheng0521@163.com)

Hongsheng Lin (tlinhsh@jnu.edu.cn)

Department of Orthopedics, The First Affiliated Hospital of Jinan University, 613

Huangpu Avenue, Guangzhou 510630, People's Republic of China.

Running title: 14-3-3/spastin pathway regulates the recovery of spinal cord injury.

Abstract Axon regeneration is abortive in the central nervous system following injury. Orchestrating microtubule dynamics has emerged as a promising approach to improve axonal regeneration. The microtubule severing enzyme spastin is essential for axonal development and regeneration through remodeling of microtubule arrangement. To date, however, little is known regarding the mechanisms underlying spastin action in neural regeneration after spinal cord injury. Here, we use glutathione transferase pulldown and immunoprecipitation assays to demonstrate that 14-3-3 interacts with spastin, both *in vivo* and *in vitro,* via spastin Ser233 phosphorylation. Moreover, we show that 14-3-3 protects spastin from degradation by inhibiting the ubiquitination pathway and upregulates the spastin-dependent severing ability. Furthermore, improving the interaction between 14-3-3 and spastin by Fusicoccin (FC-A) promotes neurite outgrowth and regeneration *in vitro*. Western blot and immunofluorescence results revealed that 14-3-3 protein is upregulated in the neuronal compartment after spinal cord injury *in vivo*. In addition, administration of FC-A not only promotes locomotor recovery, but also nerve regeneration following spinal cord injury in both contusion and lateral hemisection models. However, application of spastin inhibitor spastazoline successfully reverses these phenomena. Taken together, these results indicate that 14-3-3 is a molecular switch that regulates spastin protein levels, and 14-3-3/spastin pathway is an important target for regulation of microtubule dynamics for nerve regeneration after spinal cord injury.

Keywords: Spastin; Microtubule; Axon regeneration; 14-3-3; Spinal cord injury

Graphical Abstract

Introduction

Regeneration of the adult central nervous system neurons is difficult after injury, a

phenomenon that results in permanent neurological impairments[1]. Researchers have

attributed this not only to the highly inhibitory extrinsic environment at the injury site

but also to limited cell-intrinsic growth capacity. Many studies have demonstrated that

re-programming of growth-associated processes plays a key role in manipulation of

axonal regeneration[2-4]. However, the specific mechanisms underlying axonal

regeneration failure remain largely unknown. Cytoskeleton remodeling, including

actin and microtubule (MT) reorganization, appears to be the key factors directing the

fate of axonal regeneration and sprouting[5]. Notably, the retraction bulb of severed

axons at the lesion site of the spinal cord needs a dynamic cytoskeleton for successful

regeneration. Moreover, formation of axon collaterals from spared axons also needs

dynamic cytoskeleton reorganization to form new neural circuits with the innervated

neurons[6, 7]. Mechanistically, invasion by MTs can not only provide mechanical

force, but also guide intracellular transport to regulate distribution of specific

molecules, including mitochondria, peroxisomes, growth factors and other, and

subsequently mediate axonal extension. Therefore, timely invasion of microtubules

can not only power the formation of a growth cone with competent growth ability

from the retraction bulb, but also improve the process of axonal sprouting from the

spared axons[8, 9]. Overall, manipulating MT dynamics is critical for boosting axon

regeneration.

Spastin is encoded by SPG4 gene and its mutations were identified from the

patients of hereditary spastic paraplegia (HSP)[10]. It is a MT severing enzyme

(including spastin, katanin and fidgetin) which can cut long MTs into short fragments,

a phenomenon that not only endows MTs with high dynamics but also mediates their

subsequent turning and regrowth under physiological conditions[11]. Spastin has two

major isoforms, namely M1 and M87, coded form different translation initiation

codon (1st and 87th methionine). Both isoforms are highly expressed in the spinal

cord[12]. Moreover, spastin mutation lacking severing activity can lead to the

degeneration of the corticospinal tracts in HSP patients. In addition, spastin is highly

expressed in the central neuronal system (CNS) and essential for formation of axon

branches as well as their extension during neural development[13, 14]. These

evidences indicate that spastin plays an important role in axonal development and

maintainence in the spinal cord. Intriguingly, a study conducted in *Drosophila* has

reported that excess or deficient dosage of spastin is detrimental to axon

regeneration[15]. Furthermore, another study in *Drosophila* has also indicated the

significance of spastin in coordinating the concentration of the endoplasmic reticulum

(ER) and microtubules at the growing axon tips during axon regeneration[16]. In a

separate study, researchers administered spastin conjugated with polyethylene glycol

into the epineurium and observed its potential in mediating the recovery of motor

function following sciatic nerve injury. However, this study lacks the requisite

histological and molecular biological evidence[17]. Currently, there is insufficient

research to definitively establish the role of spastin in axon regeneration after central

nervous system injury in adult mammals. Manipulating spastin expression seems to be

an effective way to promote axon regeneration. However, efficient regeneration is

dependent on the precise spastin dosage, with studies showing that excess spastin is

toxic and may destroy the MT network, while low doses have been associated with

insufficient MT severing which subsequently limits MT remodeling for

regeneration[15, 16]. Therefore, direct manipulation of the spastin gene may not be

the most appropriate approach for promoting successful axon regeneration after the

CNS injury. It is worth noting that post-translational modifications, protein

metabolism and protein-protein interactions are vital for regulating the protein's

activity[18]. Thus, elucidating the specific molecular mechanism underlying spastin

action is imperative to guiding future development of effective interventions to

promote axon regeneration following damage to the CNS.

14-3-3 protein is a type of adaptor protein that play important roles in many

signaling pathways through interaction with their substrates[19, 20]. For instance, it

inhibits the ROS-induced cell death by interacting with FOXO transcription factors to

prevent upregulation of many pro-apoptotic genes[21]. In mammals, seven isoforms,

namely β、γ、ε、ζ、η、θ and σ, have been identified. They are not only highly conserved

in the mammalian system, but also form a dimeric structure to interact with their

substrates which are mostly phosphoproteins[22]. Previous studies have shown that

spastin is phosphorylated in Ser233 which showed its potential for interaction with

14-3-3[23]. In addition, 14-3-3 protein is also highly expressed in CNS where it plays

a vital role in both axonal growth and guidance. Previous studies have also

demonstrated that 14-3-3 promotes cortical neurite outgrowth and axon

regeneration[24, 25]. To date, however, the molecular mechanisms underlying its

action in promotion of axon regeneration as well as its role in spinal cord injury

remain largely unknown.

In this study, we demonstrate that 14-3-3 interacts with spastin to regulate its

stability, thus controlling both neurite outgrowth and regeneration after injury. Our

results reveal that 14-3-3 interacts with serine 233 in spastin, in a

phosphorylation-dependent manner, thereby preventing its degradation by

ubiquitination and upregulates the severing ability via spastin-mediated activity.

Moreover, we found that enhancing this interaction via Fusicoccin (FC-A) application

resulted in enhanced axon regeneration and locomotor improvement both in contusion

model and in T10 lateral hemi-transection model of spinal cord injury. Collectively,

our findings suggest 14-3-3/spastin pathway is a novel intervention target for axon

regeneration after spinal cord injury.

Materials and methods

Animals

Sprague Dawley rats (P0) were used for primary hippocampal and cortical neuron

culture, whereas adult female mice (C57BL/6J, 6-8 weeks) were used to establish the

lateral hemi-transection model and the contusion model of spinal cord injury. Animals

were housed in a room, maintained at 22°C under a 12:12 h light/dark photoperiod,

and given food and water ad libitum. All animal treatments were carried out in strict

adherence to the guidelines by the National Institutes of Health's Guide for the Care

and Use of Laboratory Animals. The study protocol was approved by Jinan University

Institutional Animal Care and Use Committee. The number of animals utilized and

suffering were both reduced to the absolute minimum.

Cell culture and transfection. Dissociated hippocampi were digested with 2 mg/mL

papain and plated at a density of 1×10^4cells/cm^2 onto poly-D-lysine-coated coverslips

for primary hippocampal neuron culture. Cultures were kept alive in an incubator

humidified with 5% CO_2 at 37°C using Neurobasal-A media supplemented with 0.5

mM glutamine and 2% B27. Every three days, half of the cultural media was replaced.

Various plasmids were transfected with calcium phosphate at 2th day in vitro (DIV).

COS7 cells and human embryonic kidney (HEK) 293 cells were grown in a 5% CO_2

incubator at 37°C using Dulbecco's modified Eagle media with 10% fetal bovine

serum and 1% penicillin/streptomycin as supplemental ingredients. The constructs

were transfected into the COS7 cells and HEK293 cells using Lipofectemine 2000.

Cells were cultured for 24-36 hours after transfection before being harvested.

Plasmids and constructs

Several 14-3-3 isoforms were amplified in cDNA from Sprague Dawley rats via

Polymerase Chain Reaction (PCR) as previously described[14]. The amplicons were

inserted into the pCMV-Tag-2b or pGEX-5X-3 vector (Addgene). Spastin mutants

were generated using the Quickchange Kit (Agilent, Santa Clara, CA). R18 was

synthetized by IGE Biotechnology LTD（Guangzhou, China）and inserted into

pCDNA 3.1 vector. All constructs were verified by sequencing by IGE Biotechnology

LTD（Guangzhou, China）.

Expression of recombinant proteins and GST-Pulldown assay

Purification of GST fusion of 14-3-3 proteins and pulldown experiments were

performed as previously described[14, 26]. Briefly, GST-14-3-3s constructs

(Invitrogen) were transformed into *Escherichia coli* strain BL21 (DE3) strains.

Isopropy-D-thiogalactoside was incubated for 6 hours at 30°C to stimulate the

synthesis of fusion proteins. The bacteria were first centrifuged, then resuspended in a

mixture of protease inhibitors (Merck, Whitehouse Station, NJ). The cell suspension

was treated with 0.1% lysozyme, followed by 0.5% deoxycholic acid on ice for 20

min. The contents were, sonicated (10,000 rpm for 30 min), centrifuged and the

supernatant collected, then Glutathione-Sepharose beads (supplemented with 1%

Triton X-100) used to purify GST fusion proteins in the supernatant. Next, 5 μg of

GST-fusion protein was incubated with 400 μg of protein from the spinal cord (1 cm

around T10) of SD rats (P0) under gentle rotation, then centrifugated to collect the

GST-binding proteins. Western blot assay was then performed to quantify

GST-binding proteins.

Immunoprecipitation and western blotting.

Immunoprecipitation (IP) assays were carried out as previously described[27]. Briefly,

total cell extracts or tissues (1 cm around T10) were prepared by soaking them in 400

μL of lysis buffer for 30 minutes at 4 °C. The lysates were centrifuged at 10,000 rpm

for 10 minutes at 4 °C after a brief sonication. The extract was immunoprecipitated

with 2 μg antibodies against spastin, GFP, or Flag, and then incubated for 4 hours at

4 °C with 60 μL of protein G plus protein A/G agarose using continuous gentle

inversion. Three times, the immune complexes were pelleted and washed. Western

blot assay was performed to analyze the precipitated complexes. Western blot analysis

was carried out as previously described[14]. SDS-PAGE was used to separate the

lysates, which were then electrophoretically transferred to a polyvinylidene difluoride

membrane. Membranes were probed with primary antibodies at 4 °C overnight after

being blocked in Tris-buffered saline with 5% milk and 0.05% Tween. The

membranes were cleaned, incubated with goat anti-mouse or anti-rabbit secondary

antibodies that were horseradish peroxidase conjugated, and then visualized with

ECL-prime. Anti-spastin (1:1000, ABN368 from Millipore), anti-Pan 14-3-3 (1:1000,

#8312 from Cell Signaling Technology), anti-Pan phosphoserine/threonine

(Cat#612548 form BD life science, anti-GFP (1:5000, Ab290 from Abcam), anti-GFP

(1:10000, 50430-2-AP from Proteintech), anti-Flag (1:1000, F3165 from

Sigma-Aldrich), anti-GAPDH (1:5000, AC002 from ABclonal), and HRP conjugated

goat-derived anti-rabbit or anti-mouse were the antibodies used (1:5000, Abclonal).

Scratch assay

Rat cortical neurons were seeded in 96-well plates (30,000 cells per well), then a

plastic P10 pipet tip was used to scratch across the center of the well. Half the media

was aspirated out and replaced with fresh one every three days until the 7th DIV. The

wells were immediately filled with chemicals after scratching. The cultures were fixed

and stained with anti-βIII tubulin (1:1000, ab18207) the following day. Images of

axons stained with βIII tubulin were collected, imported in ImageJ, and the proportion

of neurites stained by βIII tubulin in the central 70% of the scratch were calculated.

Lateral hemi-sectioning and contusion model of SCI

The procedure for T10 lateral hemi-sectioning and contusion model was as

previously described[28, 29]. Briefly, mice were first anesthetized with 1.25%

2,2,2-tribromo ethanol (0.2 mL/100 g), via intraperitoneal injection for 3-4 min, then

a laminectomy performed near T10 after making a midline incision over the thoracic

vertebrae. The T10 level of the spinal cord was recognized by the anatomical

landmark which is parallel to the twelfth rib of the mice. For the hemi-sectioning

injury model, the right unilateral hemi-section was carefully performed using a scalpel

and microscissors, with care taken to avoid damaging the spinal cord dura. For the contusion model, the nitrogen tank controlling the impactor tip was set at 18 psi or 124 kPa. A U-shaped stabilizer with the rat was loaded onto the stage of the Louisville Injury System Apparatus (LISA) and the dura of the spinal cord was adjusted directly under the impactor while monitored by the laser beam. Crash depth was adjusted to 1.0 mm for injury, and with time set to 0.5 s. Finally, the muscles were stitched and the skin was stapled back together.

Pharmacological interference with FC-A or spastin

An equal volume of normal saline was intraperitoneally injected into SCI control mice. Mice were intraperitoneally injected with FC-A (Enzo life sciences, ALX-350-115) at a dose of 15 mg/kg starting immediately after SCI and every other day until end of the experiment. Mice were intraperitoneally injected with spastazoline (MedChemExpress, HY-111548) at a dose of 20 mg/kg, immediately after SCI then every other day until end of experiments.

Basso Mouse Scale

Two observers who were blind to the experimental groups conducted the Basso Mouse Scale (BMS) after the T10 SCI in accordance with a previously described procedure[30]. Mice were briefly left in an open field for 4 minutes to observe all visually perceptible signs of locomotor recovery. The scores were based on hind limb movements performed in an open field and included hind limb joint movement, weight support, plantar stepping, coordination, paw position, and trunk and tail control (0, full hind limb paralysis; 9, normal locomotion).

Catwalk analysis

According to previously described methods[31], the CatWalk XT 9.1 automated quantitative gait analysis system (Noldus Company, Netherlands) was also used to evaluate the locomotor recovery of mice after injury. Briefly, one week before surgery, mice were trained to walk in a single direction for the entire length of a darkened Catwalk chamber in a quiet room five times daily. Seven weeks after surgery, the mice were assessed five times using the same system. Gait regularity index and

hindlimb contact area were recorded. The regularity index was calculated as the number of normal strides multiplied by four and then divided by the total number of strides; this value is ~100% in normal animals.

Footprint analysis

Footprint analysis was conducted before the mice were executed, using a previously described protocol[32]. Briefly, each animal's hindlimbs were first coated with nontoxic paint. The mice were then moved from a brightly illuminated starting box to a darkened one, by allowing them to walk through a narrow-custom-built plexiglass trough (5 cm wide by 40 cm long), during which they left a trace of their paw prints on the white sheet that overlaid the trough. The footprints were scanned, and digitized images measured using Image J Pro Plus. Stride length was measured as the distance between the adjacent hindlimbs, whereas stride width was taken as the distance between the affected and the contralateral limbs. At least ten footprints per side, from three sessions per animal, were measured for both parameters and their average were calculated.

Foot fault test

The Parallel Rod apparatus was used in this test[33]. Summarily, mice were positioned in the middle of the device, allowed to walk for three minutes during which a video was captured. The test was performed by two observers who were blinded to the experimental groups. One observer recorded the total number of steps taken, while the other counted instances where each right hind limb slipped through the rod. Three runs were recorded for every mouse.

Motor evoked potential (MEP) recordings

To assess SCI recovery, motor evoked potentials (MEPs) were recorded 7 weeks after SCI according to previously described methods[31]. First, mice were anesthetized with 1.25% 2,2,2-tribromo ethanol (0.2 mL/100 g). A craniotomy was then performed to expose the M1 region of the motor cortex. Electrode penetration was guided via a stereotaxic instrument to a depth of 700-1000 mm from the brain surface to target corticospinal neurons in the sensorimotor cortex. The recording

electrode was placed on the gastrocnemius muscle and the reference electrode was

placed on the paraspinal muscle between the stimulation and recording points. A

ground electrode was attached to the tail. A single square-wave stimulus of 0.5 mA,

0.5 ms duration, 2 ms time delay, and 1 Hz was used. Amplitude was measured from

the beginning of the first response wave to its maximum point. All potentials were

amplified and acquired using a digital oscilloscope (Chengdu Instrument Factory,

Chengdu, China).

Immunocytochemistry

Cells were grown on coverslips and processed according to

immunocytochemistry protocols as previously described[14]. Cells were fixed using

freshly prepared 4% paraformaldehyde, followed by permeabilization with 0.1%

Triton X-100 in TBS and blocking in 3% normal donkey serum. The cells on the

coverslips were incubated with anti-GFP (1:1000, Ab290 form Abcam) or

anti-α-tubulin(1:1000, ab7291 from Abcam) and then incubated with secondary

antibody (Alexa Fluor 488 donkey anti-rabbit or Alexa Fluor 647 donkey conjugated

to anti-goat immunoglobulin G (H+L) (Life Technologies). The coverslips were then

mounted with DAPI (diamidino-2-phenylindole)-free Fluoro-Gel on glass slides

(Electron Microscopy Sciences, USA). Cells were scanned using a 63x oil immersion

objective mounted on a confocal microscope (LSM 710 Meta; Carl Zeiss). LSM 710

software was used to process the images after they were captured using sequential

acquisition settings at a resolution of 1024 × 1024 pixels and a 12-bit depth.

Mice were administered consecutive transcardial infusions of 4%

paraformaldehyde in PBS and 0.09% saline after being given a deep anesthetic. After

tissue collection, post-fixation in 4% paraformaldehyde was applied overnight, and

tissues were then immersed in 30% sucrose for two days to dehydrate them. Serial

sagittal cryostat-sections were incubated at 4°C for at least 18 hours before being

stained with primary antibodies in blocking buffer (3% goat serum and 5% donkey

serum in 0.3% PBST). Alexa Fluor-coupled secondary antibodies (Life Technologies)

diluted in blocking buffer were incubated for 2 hours at room temperature to perform

the detection. Anti-GFAP (1:1000, GB11096 from Servicebio), anti-serotonin (1:5000,

AB125 from Sigma-Aldrich), anti-neurofililament (1:200, A19084 from ABclonal),

anti-Myelin Basic Protein (1:200, A11162 from ABclonal), anti-Brdu (1:200, A20790

from ABclonal), anti-Nestin (1:200, A11861 from ABclonal) and anti-NeuN (1:200,

A19086 from ABclonal) were used. Images were scanned using a confocal

microscope.

Statistical analysis

The analyses were carried out by Graphpad 8.4.0 software. Differences across

multiple groups were determined using One-way analysis of variance (ANOVA),

followed by Newman-Keuls post hoc tests for mean separations. $P \leq 0.05$ were

considered statistically significant.

Results

14-3-3 protein interacts with spastin

Spastin, a MTs severing enzyme, can cut long MTs into short fragments thereby

contributing to axonal elongation and branching. However, excess spastin levels by

gene manipulation can not only destroy the MT cytoskeleton but are also toxic to

neurons. Unraveling the underlying mechanisms of spastin regulation under

physiological conditions may provide novel insights into the mode of axon

regeneration following CNS damage. In our previous study[27], GST-spastin

pulldown assays with proteins lysed from the spinal cord tissue were performed, and

the obtained proteins were analyzed via mass spectrometry to determine the

underlying molecular mechanisms of spastin. The peptides that interacted with GST

were subtracted from those interacting with GST-spastin, and the peptides of proteins

that interacts with spastin were obtained. We revealed 14-3-3 as putative

spastin-interacting proteins (Figure 1A & Figure S1A). Notably, 14-3-3 proteins

consist of seven isoforms that are highly conserved and act by targeting

phosphoserine and phosphothreonine motifs of substrate proteins. We have detected

many different 14-3-3 peptides in the complex, including LAEQAER, NLLSVAYK,

and AVTEQGAELSNEER (Figure S1A). Notably, these peptides were highly

conserved among different isoforms (Figure S1B). Next, we employed

immunoprecipitation to determine the biochemical interaction between 14-3-3 protein

and spastin, and found that they endogenously interacted in the spinal cord tissue and

the cortical neurons (Figure 1B&1C). To elucidate the distribution of 14-3-3 and

spastin within neurons, hippocampal neurons were double stained with spastin and

14-3-3 antibodies. The results revealed that 14-3-3 exhibited a ubiquitous cellular

distribution, while spastin showed colocalization with 14-3-3 in both axon shafts and

growth cones (Figure 1D). Colocalization profiles are displayed in Figure 1E. 14-3-3

proteins consist of seven isoforms, and their protein spatial structure are highly

conserved (Figure 1F). To further identify which isoform of 14-3-3 interacts with

spastin, we generated six 14-3-3 isoforms in rats (β、γ、ε、ζ、η、θ), then purified GST

fusion 14-3-3 proteins (Figure 1G). Results from a GST pull-down assay revealed that

all 14-3-3 isoforms could interact with spastin (Figure 1H). Subsequent

co-immunoprecipitation assay results also confirmed that all 14-3-3 isoforms could

form complexes with spastin (Figure 1I). Collectively, these results indicated that the

14-3-3 protein interacts with spastin both *in vitro* and *in vivo*.

14-3-3 binds to phosphorylated Ser 233 in spastin

Next, we characterized the interaction region of spastin/14-3-3 binding. To this

end, we generated several GFP-tagged deletion constructs based on the modular

domain of spastin (Figure 2A), then co-expressed them with Flag-tagged 14-3-3 into

HEK293T cells. Co-immunoprecipitation assay, using GFP-Trap beads, revealed that

spastin fragments containing amino acids 85-316 but not those with 85-214

coprecipitated with 14-3-3 (Figure 2B). Next, we performed immunoprecipitation

assays using MTBD region (215-316 amino acids) of spastin with 14-3-3 and found

that 14-3-3 specifically binds to the MTBD region of spastin (Figure 2C).

Previous studies have shown that phosphorylation of the 14-3-3 substrate

regulates the binding of 14-3-3 proteins, although 14-3-3 can also bind to

non-phosphorylating proteins. To investigate whether spastin's phosphorylation state

affects its binding with 14-3-3, we performed pulldown assays by using GST-14-3-3 and GFP-spastin from transfected HEK293T cell extracts in the presence of broad protein kinase staurosporine. To validate whether staurosporine could decrease the phosphorylation levels of spastin, we enriched spastin through immunoprecipitation and used Pan phosphoserine/threonine antibody to observe spastin's phosphorylation. The results showed that staurosporine significantly reduces the phosphorylation levels of spastin (Figure. 2D). Furthermore, our results revealed that the presence of staurosporine markedly weakened the binding between spastin and 14-3-3, indicating that phosphorylation of spastin is sufficient for the binding with 14-3-3 (Figure 2E).

Next, we predicted five putative 14-3-3 binding sites in spastin, namely amino acids 213-218, 230-235, 243-249, 425-460, and 559-564, which match the motif of RSX(S/T)XP. Combination with phosphorylation of spastin reported from MS data (https://www.phosphosite.org/proteinAction.action?id=8325&showAllSites=true), we revealed two putative binding motifs (Ser233 and Ser562) in spastin as putative targets for 14-3-3 binding (Figure 2F). Widely employed mutation techniques have showed that mutating a protein's phosphorylation site to alanine prevents phosphorylation, while mutating it to aspartic acid can mimic phosphorylation. Then, we mutated single Ser233 and Ser562 or both to alanine in the spastin tagged with GFP and co-expressed them with Flag tagged 14-3-3 in HEK293T cells. Immunoprecipitation of the lysates, using an antibody against GFP, revealed that spastin S562A can bind with 14-3-3, while spastin S233A and spastin Pan A cannot (Figure 2G). To further illustrate the significance of phosphorylation of spastin at S233 in its interaction with 14-3-3, the mutation of Ser 233 in spastin to aspartic acid was also transfected for immunoprecipitation, the results revealed that spastin S233D exhibited enhanced binding to 14-3-3 compared to both wild type and spastin S233A, and this interaction was even insensitive to the staurosporine treatment (Figure 2H). This evidence further suggests that phosphorylation of spastin at the S233 site is a critical event in its interaction with 14-3-3. Also, these results were consistent with those in Figure 3B-C, where Ser233 was localized in the region of 215-316 amino

acids in spastin, further indicating that the binding of spastin/14-3-3 requires

phosphorylation of Ser233 in spastin.

To examine whether Ser233 in spastin is involved in neurite outgrowth, wild

type (GFP-Spastin) or spastin Ser233 mutant (GFP-Spastin S233A or GFP-Spastin

S233D) was moderately expressed in primary hippocampal neurons at 2th DIV. It is

important to note that overexpression of spastin can lead the severing of the overall

microtubule cytoskeleton in neurons, leading to neuronal toxicity. To achieve a

moderate level of spastin expression, the transfection dosage and duration were

meticulously controlled. Consistent with earlier results[34, 35], spastin promoted

neurite outgrowth, as evidenced by both the length and branches of neurite. A similar

effect was observed after transfection with spastin S233A. In contrast, we found a

toxic effect following transfection with spastin S233D (Figure 2G-I). Following

conventional logic, it is anticipated that spastin S233A, which cannot bind to 14-3-3,

would inhibit neuronal growth, while spastin S233D, which binds to 14-3-3, would

promote neuronal growth; however, we found that both wild-type spastin and S233A

could promote neurite outgrowth primarily, and this phenomenon may be ascribed to

their substantial expression using an overexpressing method (regardless of whether it

is WT or spastin S233A), and the spastin concentration was far beyond the

physiological concentration. Consequently, we did not uncover the expected role of

spastin S233 in neurite outgrowth. However, spastin S233D exhibits an opposing

effect on neurons, suggesting that phosphorylation of S233 in spastin influences

spastin's function and the development of neurite outgrowth.

14-3-3 protects spastin from degradation via spastin S233 phosphorylation

How does the interaction between 14-3-3 and Ser233 in spastin regulate neurite

outgrowth during neural development, and why does spastin S233D expression in

neurons cause a toxic effect? Hence, we first investigated whether the

phosphorylation of spastin at S233 affects its microtubule-severing capability. As

illustrated in figure S2, both spastin S233A and S233D could effectively severed

microtubules. Furthermore, the S233 site is situated outside of spastin's MTBD

domain (amino acids 270-328) and AAA (amino acids 342-599) domains. Consequently, we speculated that phosphorylation of the spastin S233 site does not influence spastin's microtubule-severing ability. Notably, we observed a significantly higher green fluorescence intensity of GFP-spastin (S233D) compared to spastin S233A. Combined with the fact that spastin acts in a dose-dependent manner and spastin S233D expression in neurons caused a toxic effect, we therefore hypothesized that phosphorylation of Ser233 in spastin may affect the protein levels of spastin. To further test whether 14-3-3 would affect spastin stability, we performed chase-time assays via GFP-spastin overexpression, followed by treatment with cycloheximide (CHX) to inhibit protein synthesis. Results showed that expression of GFP-spastin protein was markedly downregulated at 9 h after CHX treatment (Figure 3A). Next, we co-expressed Flag-14-3-3 with GFP-spastin in the same treatment and found that 14-3-3 could significantly protect spastin from degradation after CHX treatment, even after 12 h (Figure 3B). We also investigated whether spastin S233 phosphorylation by transfecting with GFP tagged spastin S233A or S233D followed by CHX treatment. Due to the presence of endogenous 14-3-3, we did not overexpress 14-3-3 with S233 mutations that mimicked different phosphorylation state. The results demonstrated that phosphorylation of spastin at S233 (S233D) protected spastin from degradation (Figure 3C&D&I). Since 14-3-3 can inhibit spastin protein turnover, does 14-3-3 also increase the protein level of spastin? Therefore, we conducted chase-time assays by overexpressing GFP-spastin without addition of CHX. The results showed that 14-3-3 could upregulate the protein levels of spastin (Figure 3E&F&J). We also observed that spastin S233D significantly increased the protein level compared to the wild type spastin (Figure 3G&K). These findings suggest that 14-3-3 can inhibit protein turnover by binding to phosphorylated spastin and upregulate the protein levels of spastin.

A previous study demonstrated that spastin could be polyubiquitinated[36]. Therefore, we employed MG132, a proteasome inhibitor, to suppress the ubiquitin-proteasome degradation and assess whether it impacts the degradation

pathway of spastin in neuronal compartments. The results revealed that in the

presence of MG132, the protein levels of spastin significantly increased. However, in

the presence of CHX, the protein levels of spastin decreased, while the inhibition of

ubiquitin-proteasome degradation by MG132 did not show a significant decrease

(Figure 3L&M). These findings indicate the presence of a proteasomal degradation

pathway for spastin.

Previous studies have shown that FC-A and R18 interact with 14-3-3 protein by

either stabilizing or inhibiting the binding of 14-3-3 with their substrates by directly

docking to the groove in the 14-3-3 proteins with highly efficiency[37-39]. The

structure of FC-A and R18 binding with 14-3-3 are shown in Figure 4A. To test our

hypothesis, we first explored whether FC-A or R18 could enhance or inhibit the

interaction between 14-3-3 and spastin by co-immunoprecipitation with lysates of

HEK293T cells, which had been transfected with GFP-spastin and Flag-14-3-3

plasmids whether together with FC-A application or R18 transfection. We found that

FC-A could enhance the binding of 14-3-3/spastin, as evidenced by more intense

bands compared to no FC-A application, whereas R18 could efficiently inhibit the

binding of 14-3-3/spastin with almost no binding was detected (Figure 4B). To further

determine whether 14-3-3 affected spastin's ubiquitination pathway, we performed

co-immunoprecipitation assays. GFP tagged spastin ,GFP tagged spastin S233A ,GFP

tagged spastin S233D and Flag tagged ubiquitin were transfected into HEK293T cells,

together with 14-3-3 binding inhibitor R18 or not. Results indicated that GFP-spastin

could be ubiquitinated, while the ubiquitination of non-phosphorylated spastin

(S233A) was significantly enhanced, and the addition of the 14-3-3 inhibitor R18

elevated the ubiquitination levels of spastin (Figure 3N&O). These results indicated

that 14-3-3 protects spastin from degradation by inhibiting the ubiquitination pathway.

To assess the impact of 14-3-3 on the microtubule-severing ability of spastin, we

co-transfected GFP tagged spastin into COS7 cells with or without ubiquitin. COS7

cells are very flat, hence provide more spatial resolution compared to neurons. Results

showed that cells transfected with wild type spastin significantly induced microtubule

severing, as evidenced by almost no MT staining (Figure 3P), while ubiquitin inhibited severing, leading to increased microtubule staining. Based on these conditions, further transfection with 14-3-3 plasmid induced more microtubule staining, indicating that 14-3-3 upregulates microtubule severing and the process is mediated by spastin. Furthermore, we investigated whether this effect was dependent on spastin S233 phosphorylation by replacing GFP-spastin with GFP-spastin (S233A or S233D) during transfection. Results showed that 14-3-3 could upregulate the severing ability of spastin S233D, but not S233A (Figure 3Q), further indicating that 14-3-3 binds to the phosphorylation of S233 in spastin and promotes its microtubule severing ability. Collectively, these results suggest that 14-3-3 protein protects spastin from degradation via spastin S233 phosphorylation.

14-3-3/spastin complex regulates neurite outgrowth and regeneration after injury

Next, we investigated how the interaction between 14-3-3 and spastin regulates neurite outgrowth and regeneration by applying FC-A and R18 to either stabilize or inhibit 14-3-3 binding. Another drug, spastazoline, introduces an electropositive group to the N386 in human spastin side chain, leading to specific and efficient inhibition of the ATPase activity of AAA domain to abolish spastin's microtubule-severing function.[40, 41]. Subsequently, we used FC-A or R18 together with spastazoline to examine the effects of 14-3-3/spastin protein complex on neurite outgrowth and regeneration.

Since 14-3-3 proteins play important roles in inhibition of ROS-induced cell death, we first explored the effect of 14-3-3/spastin binding in neurite outgrowth and repair under glutamate circumstances. Following neural injury, neurons undergo lysis, resulting in the release of a substantial quantity of excitatory glutamate[42]. Glutamate, upon binding to its receptors, initiates calcium overload, subsequently causing mitochondrial membrane potential collapse, nuclear swelling, and ultimately inducing neuronal necrosis. This process is characterized by neurite swelling and breakage [43, 44]. Our results showed that the injured neurons had numerous swellings and breakages on the neurite, indicative of successful establishment of an

injury model (Figure 4C). After injury, FC-A was administered into primary hippocampal neurons during the developmental stage 2, combination with application of spastin inhibitor spastazoline. Results demonstrated that FC-A could promote neurite outgrowth, both with regards to lengths and branches, after glutamate-induced injury, while this promoting effect was abolished following spastazoline administration (Figure 4D-G). A similar effect was observed in neurons without glutamate induced injury (Figure S4). In contrast, transfection of R18 resulted in an inhibitory effect on neurite outgrowth, but this was rescued with moderate spastin expression after injury (Figure 4D-J). A similar trend was observed in neurons without glutamate-induced injury (Figure S5). Collectively, these results suggest that 14-3-3 protein promotes the repair of injured neuron through the interaction with spastin.

Next, we investigated the effect of enhanced 14-3-3/spastin binding on neurite regeneration. Briefly, cortical neurons were grown at the 7th DIV then subjected to scratch assays by drawing a line over the coverslips using a pipet tip. Neurons were then treated with either FC-A or spastazoline for 36 hours. Results showed that FC-A markedly improved neurite regeneration consistent with previous study[37], while spastazoline application almost abolished the enhanced effects of regeneration (Figure 4K-L). To further quantify the extend of axon regeneration, we treated cortical neurons with either FC-A or spastazoline for 36 hours, then quantified the longest regenerative axons. Results revealed similar effects (Figure S6). To further elucidate whether FC-A affects neuronal process by impacting on spastin, cortical neurons cultured *in vitro* were administrated with FC-A and the protein expression levels of spastin were detected. The results indicated that FC-A could enhance the expression levels of spastin within cortical neurons (Figure. 4M). Taken together, these results indicate that spastin-dependent function is a prerequisite for the enhancement of neurite regeneration mediated by FC-A.

FC-A-mediated nerve regeneration after SCI requires spastin activation by targeting MTs

Given that 14-3-3 plays vital roles in neurite outgrowth and axonal

regeneration[45], we then hypothesized that 14-3-3 could be involved in the recovery

of spinal cord injury. To test this hypothesis, we established a spinal contusion injury

model and then employed immunostaining assay to quantify expression levels of

14-3-3 protein following spinal cord injury. Given the scarcity of neural compartment

at the injury center, we select tissue slices as close as possible to lesion core to

illustrate the relationship between 14-3-3 and the injured neurons. The results

indicated that 14-3-3 protein was upregulated and the upregulated compartments were

principally colocalized in the neuronal compartment (βIII-tubulin labeled) (Figure

5A). Western blot experiments also reported that 14-3-3 was upregulated at 3th and

7th days post injury (DPI), but downregulated at 14 DPI. The 14-3-3 protein level

then reached a peak level at 30 DPI (Figure 5B&E). Moreover, previous studies have

demonstrated that neurons were surrounded by inhibitory microenvironment 2 weeks

after spinal cord injury, our results also showed a slightly decrease at 14 DPI,

indicating that 14-3-3 acts as an intrinsic switch for neural repair. Moreover, 14-3-3

was reported to be upregulated and acted as a biomarker for diagnosis of

Creutzfeldt-Jakob disease[46]. Our results revealed that levels of 14-3-3 protein

remained high even at 30 DPI, indicating that 14-3-3 may plays an important role in

the process of recovery after spinal cord injury.

So, how does the expression of spastin change following spinal cord injury? As

previously mentioned, the drug FC-A is known to enhance the binding affinity

between 14-3-3 and spastin, thereby increasing spastin's protein stability. Therefore,

can the administration of FC-A after spinal cord injury upregulate the protein levels of

spastin? To address these questions, tissues were collected and western blot

experiments were conducted to assess the expression levels of spastin after spinal cord

injury. Our findings revealed that the expression of spastin exhibited a sharp decrease

immediately after spinal cord injury (1 day), followed by a gradual increase during the

acute phase (1-7 days), eventually returning to levels similar to the sham group at 30

days (Figure 5C). Notably, the expression trend of spastin is similar to that of 14-3-3

protein after spinal cord injury (Figure 5H), suggesting potential involvement of

14-3-3 and spastin in the same pathway. Moreover, we observed that the application

of FC-A after spinal cord injury significantly augmented spastin's expression (Figure

5D&G&I). These results indicated that the administration of FC-A may have the

potential to enhance spastin's protein stability after spinal cord injury.

Enhancing the 14-3-3/spastin interaction mediated by FC-A can promote neurite

outgrowth and regeneration suggesting that it can be used to promote neurite regrowth

after injury *in vivo*. We therefore attempted to investigate whether FC-A could

improve nerve regrowth after spinal cord injury in a contusion injury model. FC-A

was administrated via intraperitoneal injection following SCI in mice. As shown in

figure 5J, H&E and LFB (Luxol fast blue stain) staining showed that the lesion site in

the injury group presented the destroyed tissue structure with loss of myelination.

Intriguingly, we found the tissue structure appeared to be normal and the area of

demyelination in the lesion site strongly decreased in the FC-A group. In addition, in

the spastazoline group, the lesion site appeared to be discontiguous, and application of

spastazoline reversed the protective effect of FC-A on spinal cord demyelination

(Figure 5K). Since spastin plays a crucial role not only in neuronal development but

also in mitosis, spastazoline may inhibit the proliferation of stormal cells in the lesion

site, thereby impeding the wound healing process following spinal cord injury. Next,

we further investigated nerve regeneration by immunostaining with NF

(Neurofilament) and 5-HT, which were co-stained with GFAP (astrocyte marker). As

shown in figure 5E, the border of the spinal cord lesion site was marked with

astrogliosis (GFAP$^+$), and the neurofilament and 5-HT immunogenicity in the lesion

site were almost abolished compared to the uninjured area, indicating that the spinal

cord contusion injury model was successfully established. Interestingly, in the FC-A

group, we observed that the astrogliosis in the lesion site was attenuated and we found

a great deal of neurofilaments and 5-HT signals in the lesion site (Figure 5L, boxed in

L2), indicating that the application of FC-A enhanced the nerve regeneration after

injury. In the spastazoline group, we also observed tissue discontinuity, which is

consistent with figure 5J, and almost no NF and 5-HT signals were found in the lesion

site. Moreover, the application of spastazoline abolished the FC-A mediated nerve

regeneration after SCI. These results suggest that FC-A-mediated axonal regeneration

after SCI requires the activation of spastin function.

In order to further confirm whether the intervention of 14-3-3/spastin pathway by

FC-A and spastazoline mediates nerve regeneration after spinal cord injury by

targeting MTs at the site of spinal cord injury, acetylated tubulin and β-tubulin were

stained to label stale MTs (Lacking of dynamics) and total MTs. As shown in figure

S7 A, the intensity of acetylated microtubules was remarkable decreased in the lesion

site. Moreover, there was a clear boundary of acetylated MTs between the lesion site

and the uninjured area. This may be ascribed to the activation of cell proliferation in

the injured area after spinal cord injury which needs dynamic microtubules.

Interestingly, we found the boundary of acetylated microtubules near the damaged

area was unclear after application of FC-A, and the intensity of acetylated

microtubules significantly decreased compared with the injury group, indicating that

microtubules tended to be a dynamic state. However, after the application of

spastazoline, the proportion of acetylated microtubules in the injury site increased

significantly, and the boundary of acetylated microtubules in the injured area was

unclear, indicating that the application of spastazoline led to a significant increase in

the stability of the microtubules. In addition, the application of spastazoline can

significantly reverse the enhancement of microtubule dynamics mediated by FC-A

(Figure 5M&O). These experiments indicated that the administration of FC-A mediate

nerve regeneration after spinal cord injury through the 14-3-3/spastin pathway by

affecting the MTs dynamics. In addition, these evidences also indicated that the FC-A

and spastazoline can pass through the blood-brain barrier to reach the site of spinal

cord injury.

FC-A-mediated locomotor function improvement after SCI requires spastin

participation

Since FC-A promotes nerve regeneration after spinal cord injury by affecting the

14-3-3/spastin pathway, we then investigated whether it could mediate the recovery of

locomotor function after SCI. Therefore, we performed several experiments to verify whether the application of FC-A could affect the locomotor function recovery after SCI (Fig. 6A). Catwalk analysis showed that the maximum contact area of the mice in the FC-A group was significantly larger than that in the injury group, and this phenomenon was reversed after the application of spastazoline (Figure 6B&E). The regularity index (Calculated as the number of normal strides multiplied by four and then divided by the total number of strides; this value is ~100% in normal animals) also had a similar phenomenon. The gait of the hindlimbs of mice in the FC-A group was more coordinated than that of the mice in the injury group (Fig. 6C), and the regularity index was statistically enhanced compared with the injury group (Fig. 6G). This phenomenon was also significantly suppressed after the application of spastazoline. We further evaluated the motor function after spinal cord injury by BMS (Basso mouse scale) score (Fig. 6D). BMS is a 9-point scale that provides a gross indication of locomotor ability and determines the phases of locomotor recovery. We found that the BMS scores of the mice in the FC-A group was significantly improved compared with the mice in the injury group, and peaking after 35 days, while spastazoline reversed the improvement of locomotor function. At the same time, we used the footprint test to analyze the motor function of the mice (Fig. 6F). The experimental data showed that the paw trailing phenomenon of the hindlimbs in the FC-A group was significantly improved compared with the injury group, and the stride length was significantly increased. This improvement was also significantly suppressed by application of spastazoline (Fig. 6H). Finally, we attempted to verify whether the spinal neural circuits in mice were remodeled by examining the motor-evoked potentials (Stimulation at the sensorimotor cortex in the M1 region and receive signals at the site of gastrocnemius). Our experimental results showed that the MEP current amplitude decreased significantly after spinal cord injury, while the current amplitude in the FC-A group significantly increased compared with the injury group. And the improvement effect was significantly weakened after the application of spastazoline, indicating that FC-A mediates the remodeling of neural circuits after

645   spinal cord injury by affecting the 14-3-3/spastin pathway.

Enhancing the 14-3-3/spastin pathway by FC-A stimulates the repair of SCI in a

T10 lateral hemisection model

We have clarified the important role of 14-3-3/spastin pathway in the repair of

spinal cord injury in the contusion model. In order to further observe the axon

regeneration after spinal cord injury, we performed a spinal right lateral hemisection

at the T10 level to eliminate the right side of axonal projections. Neurotransmission of

serotonin (5-HT) in the spinal cord is required for the modulation of sensory, motor,

and autonomic functions and can influence the response to spinal cord injury[47, 48].

In addition, studies have reported that the 5-HT fibers have strong regenerative

capacity and involved in the regulation of neuronal activity[48]. Thus, we explored

the 5-HT immunoactivity in spinal cord tissues isolated on 17 DPI. In the control SCI

group, 5-HT-positive axons were significantly upregulated in the injury site, and to

some extent in the rostral site. Interestingly, we found much stronger 5-HT staining in

the lesion site of FC-A group and rostral site of injury compared with the SCI control

group, indicating that the FC-A promoted the regeneration of 5-HT positive axons *in*

vivo. However, few 5-HT signals were found in the lesion site, also in the rostral site

in the spastazoline group. In addition, the FC-A promoting effects of 5-HT axonal

regeneration were significantly weakened by the combined application of spastazoline

(Figure 7A&C).

We further investigated the immunoactivity of neurofilament and myelin basic

protein (MBP) in the white matter of the lesion site to evaluate axonal regeneration

and the extend of remyelination. Neurofilament are cytoskeletal proteins that are

expressed abundantly in the cytoplasm of axonal fibers in the CNS[49]. The MBP

forms and maintains the structure of the compact myelin sheath which then regulates

axonal function[50]. Our results revealed that FC-A could enhance the regeneration of

neurofilaments which were largely colocalized with MBP, whereas spastazoline

alleviated this effect (Figure 7B&D). Collectively, these results suggest that FC-A

promotes axonal regeneration which is dependent on activation of spastin microtubule

severing.

Previous studies have reported that the proliferation of neural stem/progenitor

cells is also reactivated and contributes to synapse formation following spinal cord

injury[51, 52]. Since spastin is a microtubule severing enzyme involved in cell

division. Thus, we explored the possibility that whether 14-3-3/spastin complex

participates in proliferation process of neural stem/progenitor cells. Therefore, spinal

cord tissues from the lesion site were stained with Nestin, Brdu and NeuN.

Intriguingly, we found that FC-A treatment after SCI significantly increased the

expression of Nestin (Figure S9A-B) and Brdu (Figure S10), and these effects were

abolished by spastazoline treatment. These results demonstrated that the

14-3-3/spastin complex contributed to the proliferation of neural stem/progenitor cells

thereby improving recovery after spinal cord injury.

We also further examined whether FC-A administration altered the motor

function after SCI in this lateral hemisection injury model. Mice with SCI in all

groups exhibited paralysis in the right hindlimb after hemisecion which confirmed the

successful establishment of the lateral hemisection SCI model. For mice in SCI

control group, the BMS scores gradually improved to ~4 by 16 DPI. In contrast, the

scores of mice in the FC-A group indicated an improvement in locomotor function

which was impaired in the spastazoline group. Moreover, the improvement in

locomotor function following administration of FC-A diminished when spastazoline

was administered (Figure 7E). We further analyzed the footprint of mice with SCI and

calculated the length or width of strides (Figure S11). We found that stride length

improved in mice of the FC-A group but it was reduced in the spastazoline group

although not significantly. The improvement induced by FC-A diminished when

co-treated with spastazoline (Figure 7G). Notably, there was no significant difference

in stride width between the groups (Figure 7H). Similar effects were observed in the

foot fault scan experiment. The faults of right hindlimb significantly decreased in the

FC-A group and increased in the spastazoline group (Figure 7F). The improvement of

hindlimb fault in mice administered with FC-A diminished following the application

of spastazoline. We also explored whether the stability of MTs was altered by administration of FC-A and spastazoline in the spinal cord injury model. We found that the ratio of acetylated tubulin to total tubulin was significantly increased in the SCI group, decreased in the FC-A group, and upregulated in the spastazoline group, which further confirmed the successful delivery of drugs to the lesion site (Figure S8). These findings suggested that FC-A was effectively delivered to the spinal cord tissue and improved the locomotor function of mice with SCI, through 14-3-3/spastin pathway activation mechanism.

Discussion

The limited cell intrinsic regrowth capability at the injury site of CNS damage or disease results in failure of axonal regeneration. Recent studies have demonstrated that the remodeling of microtubule dynamics was necessary for promoting axon regeneration[53, 54]. Spastin is a microtubule (MT) severing enzyme which cuts long MTs into short fragments for local MTs remodeling and thus could be administrated to promote axon regeneration[17]. In this study, we aimed to reveal the regulatory molecular mechanisms of spastin and its role in the recovery of spinal cord injury. We found that 14-3-3 interacted with spastin by targeting the phosphorylation of Ser233, protecting spastin against ubiquitin-mediated degradation, thus up-regulated the spastin-dependent microtubule severing activity. Moreover, enhancing the 14-3-3/spastin interaction by FC-A promoted the neurite outgrowth and regeneration. In adult mice with of SCI (both in contusion and lateral hemisection spinal cord injury model), we found that enhanced 14-3-3/spastin binding promoted the recovery of locomotor function and axonal regeneration. Therefore, we postulated that the 14-3-3/spastin pathway is an attractive target for improving axonal regeneration and locomotor recovery after SCI.

To date, the severing activity induced by spastin has been well characterized[55, 56] and reported to be essential in the formation of axon branches and axonal extension[34, 57]. Studies have also shown that spastin regulates axon regeneration in

a dose dependent manner. Of note, during regeneration, the cytoskeleton undergoes remodeling, suggesting that manipulation of MTs dynamics by spastin may be a potential approach for controlling neural regeneration. However, efficient regeneration is dependent on the precise spastin dosage. Therefore, understanding the underlying regulatory mechanism of spastin is pivotal for identifying therapeutic targets in CNS. In this study, LC-MS based proteomics analysis was performed and proteins that bind spastin were screened to investigate the regulatory mechanisms of spastin during CNS damage (Figure 1A&S1). Among the identified binding proteins, 14-3-3 proteins had high scores and was found to interact with spastin endogenously in the spinal cord (Figure 1B&1C). Of note, MS data from a previous study also showed that the spastin peptides were present in the binding peptides with 14-3-3 agonist Fusicoccin[17], which confirmed the important relevance between these two proteins. Considered that both 14-3-3 and spastin were found to regulate axon outgrowth and regeneration, the localizations of both proteins were characterized within hippocampal neurons. In addition, GST pulldown and co-immunoprecipitation assays further verified the interaction between all 14-3-3 isoforms and spastin (Figure 1F-H). Also, the binding between 14-3-3 and spastin was further characterized which showed that Ser233 in spastin was responsible for the binding, and could directly influence the fate of neurite outgrowth (Figure 2). Previously, it was found that spastin was phosphorylated at Ser233 (human Ser268) by HIPK2 and this process was vital for successful abscission and the loss of HIPK2 exhibited a protective effect on neurons[58]. Our western blotting results showed that HIPK2 was expressed at 9th DIV in the brain tissue (not presented), suggesting that there may be other kinases that regulate Ser233 phosphorylation during neural development. Therefore, we have well established the model of the 14-3-3/spastin binding which could be the intervention target for regulating the MT severing mediated by spastin.

The regulatory mechanisms that control the MT severing activity mediated by spastin are largely unknown. In our previous study, spastin protein expression levels and its severing effect were regulated at post-transcriptional levels, such as

micoRNA-30[34] and SUMOylation[14]. In this study, we found that 14-3-3

interacted with spastin from the perspective of protein-protein interactions, and we

showed that 14-3-3 prevented the ubiquitin-mediated degradation of spastin and

upregulated the spastin protein levels by inhibiting the ubuiquitin pathway and

phosphorylation of Ser233 was crucial in this process. Our results showed that Ser233

de-phosphorylation lowered the protein stability, consistent with findings from a

recent study[36], and our study further illustrated the molecular mechanisms of this

phenomenon. In addition, we observed that 14-3-3 up-regulated the microtubule

severing activity of spastin through interacting with its phosphorylation at Ser233.

Based on this mechanism, we proposed a model in which Ser233 is phosphorylated

during the binding of 14-3-3 to spastin, thereby disrupting the spastin/ubiquitin

binding, ultimately affecting the stability of spastin and subsequent neurite outgrowth

and regeneration. Similar mechanisms were reported for doublecortin[59], cdt2[60],

p21[61] and other proteins[62]. For example, 14-3-3ε was found to interact with

threonine 42 in doublecortin which protected it from degradation by inhibiting its

ubiquitination. In this way, it regulated the doublecortin-regulated microtubule

stability and neurite initiation process[60]. In this part, we validated that 14-3-3

stabilized and upregulated the protein levels of spastin and enhanced the MT severing

activity mediated by spastin which was similar to the mechanism of 14-3-3/cdt2

binding.

Since 14-3-3 proteins are adaptor proteins that are highly expressed in neural

tissues, they are important regulators in neural development[63, 64]. Using a

well-established 14-3-3/spastin interaction model, we found that promoting the MT

dynamics by enhancing the interaction of 14-3-3/spastin could enhance neurite

outgrowth and regeneration. This accelerated neurite outgrowth and repair following

FC-A application by facilitating the binding of 14-3-3/spastin (Figure 4). In

accordance with previous findings, 14-3-3 proteins can promote neurite outgrowth

and axonal regeneration[37]. In our study, we further demonstrated that enhanced

14-3-3/spastin interaction by FC-A favored neurite outgrowth and regeneration, but

this effect was blunted by spastazoline treatment. Spastazoline was designed to

compete with the AAA domain of spastin and this effectively abolished the severing

activity mediated by spastin[40]. The results also demonstrated that the normal

severing activity mediated by spastin was required for neurite outgrowth and

regeneration (Figure 4). These findings open a new window for developing

spastin-based agents for accelerating axon regeneration after injury.

So far, few studies have investigated the relationship between spastin and neural

regeneration in CNS diseases, such as spinal cord injury or brain injury. However,

many studies have elucidated its indispensable role in the process of axon

regeneration[17]. In this study, we first observed a downregulation of spastin

expression at 1st day post-spinal cord injury, which gradually increased in the

recovery process. We also indicate that functional spastin is required for axon

regeneration and locomotor recovery after spinal cord injury (Figure 5&6). Moreover,

14-3-3 was upregulated at 3 and 7 DPI, and was even higher at 30 DPI in the neuronal

compartment, indicating that 14-3-3 may plays an important role in the process of

recovery after spinal cord injury (Figure 5). Using the model of 14-3-3/spastin

interaction, we further elucidated the mechanism by which FC-A administration

regulates the recovery of spinal cord injury by targeting MTs. Fusicoccin-A (FC-A) is

a fusicoccane, a small molecule produced by the *Phomopsis amygdali fungus* that

stabilizes the binding of 14-3-3 to its substrates[65]. The crystal structure of FC-A

bound to 14-3-3 has already revealed the efficiency of the enhancement of FC-A in

the interaction between 14-3-3 and their substrate[66]. In addition, FC-A has been

reported to promote axonal regeneration in mammals[37], although the mechanisms

are rarely elusive. In this study, We found that FC-A can enhance the expression of

spastin after spinal cord injury, and FC-A administration restored nerve regeneration

and locomotor recovery after SCI (in both contusion and lateral hemisection spinal

cord injury model), but these improvements were abolished following spastazoline

administration (inhibiting the MT severing by direct competing with AAA domain of

spastin[40]), suggesting the important role of 14-3-3/pathway in the neural circuts

remolding after SCI. This phenomenon was further confirmed by examined the ratio

of acetylated tubulin in the lesion site of SCI, and our results showed the intervention

of 14-3-3/spastin pathway by FC-A and spastazoline could significantly affect the

MTs dynamics in the lesion site which indirectly demonstrated the efficient delivery

of these drugs to the spinal cord across the blood brain barrier. Since spastin plays an

essential role in cell abscission, and the activation of 14-3-3/spastin pathway by FC-A

promoted the proliferation of neural stem/progenitor cells (Figure S9) thereby support

axonal regeneration. These results also support that promoting 14-3-3/spastin

interaction by FC-A enhanced axonal regeneration after spinal cord injury, and the

microtubule severing independent of spastin was important in this process. Therefore,

our findings present 14-3-3/spastin pathway as a novel intervention for accelerating

nerve regeneration after spinal cord injury.

In summary, we proposed the 14-3-3/spastin binding model for manipulating

microtubule dynamics and can be an effective intervention for promoting recovery of

spinal cord injury. In addition, our results indicate that 14-3-3 and spastin are

important regulators for nerve regeneration and modulating the interaction between

14-3-3 and spastin may be an effective strategy for the treatment of spinal cord injury.

Abbreviations

5-HT, 5-hydroxytryptamine; Cdt2, cell division cycle protein; CNS, central neuronal

system; DPI, days post injury; ECL, enhanced chemiluminescence; FOXO, forkhead

box transcription factors; GFAP, glial fibrillary acidic protein; GFP, green fluorescent

protein; HIPK2, homeodomain interacting protein kinase 2; HSP, hereditary spastic

paraplegia; IP, Immunoprecipitation; MBP, Myelin basic protein; MS, mass

spectrometry; MT, microtubule; ROS, reactive oxygen species;

Acknowledgement

This work was supported by the Natural Science Foundation of China (grant nos.

82102314, 82372504 and 32170977), the Guangdong Basic and Applied Basic

Research Foundation (grant nos. 2022A1515010438 and 2022A1515012306), Science

and Technology Projects in Guangzhou (grant nos. 2023A04J1284, 2023A03J1024,

202201020018), the Clinical Frontier Technology Program of the First Affiliated

Hospital of Jinan University, China, No. JNU1AF- CFTP- 2022- a01206 and Project

funded by China Postdoctoral Science Foundation (2023M731320).

Competing interests

The authors declare no conflict of interest.

Data availability

The datasets obtained and/or analyzed in the current study are available from the

corresponding author upon reasonable request.

Reference

1.   Curcio M, Bradke F. Axon Regeneration in the Central Nervous System: Facing

the Challenges from the Inside. Annual review of cell and developmental biology.

2018; 34: 495-521.

2.   Hilton BJ, Husch A, Schaffran B, Lin TC, Burnside ER, Dupraz S, et al. An

active vesicle priming machinery suppresses axon regeneration upon adult CNS injury.

Neuron. 2022; 110: 51-69.e7.

3.   Mahar M, Cavalli V. Intrinsic mechanisms of neuronal axon regeneration. Nature

reviews Neuroscience. 2018; 19: 323-37.

4.   Clark CEJ, Liu Y, Cooper HM. The Yin and Yang of Wnt/Ryk axon guidance in

development and regeneration. Science China Life Sciences. 2014; 57: 366-71.

5.   Jin Y. Unraveling the mechanisms of synapse formation and axon regeneration:

the awesome power of C. elegans genetics. Science China Life Sciences. 2015; 58:

1084-8.

6.   Pinto-Costa R, Sousa SC, Leite SC, Nogueira-Rodrigues J, Ferreira da Silva T,

Machado D, et al. Profilin 1 delivery tunes cytoskeletal dynamics toward CNS axon

regeneration. The Journal of clinical investigation. 2020; 130: 2024-40.

7.   Crunkhorn S. CNS injury: Microtubule stabilizer repairs spinal cord injury.

Nature reviews Drug discovery. 2015; 14: 310.

8.   Chen L, Rolls MM. Microtubule deacetylation sets the stage for successful axon

regeneration. The EMBO journal. 2012; 31: 3033-5.

9.   Vargas EJM, Matamoros AJ, Qiu J, Jan CH, Wang Q, Gorczyca D, et al. The

microtubule regulator ringer functions downstream from the RNA repair/splicing

pathway to promote axon regeneration. Genes & development. 2020; 34: 194-208.

10. Shribman S, Reid E, Crosby AH, Houlden H, Warner TT. Hereditary spastic

paraplegia: from diagnosis to emerging therapeutic approaches. The Lancet Neurology.

2019; 18: 1136-46.

11. Liu Q, Zhang G, Ji Z, Lin H. Molecular and cellular mechanisms of spastin

in neural development and disease (Review). International journal of molecular

medicine. 2021; 48.

12. Salinas S, Carazo-Salas RE, Proukakis C, Schiavo G, Warner TT. Spastin and

microtubules: Functions in health and disease. J Neurosci Res. 2007; 85: 2778-82.

13. Lopes AT, Hausrat TJ, Heisler FF, Gromova KV, Lombino FL, Fischer T, et al.

Spastin depletion increases tubulin polyglutamylation and impairs kinesin-mediated

neuronal transport, leading to working and associative memory deficits. PLoS biology.

2020; 18: e3000820.

14. Ji ZS, Liu QL, Zhang JF, Yang YH, Li J, Zhang GW, et al. SUMOylation of

spastin promotes the internalization of GluA1 and regulates dendritic spine

morphology by targeting microtubule dynamics. Neurobiology of disease. 2020; 146:

105133.

15. Stone MC, Rao K, Gheres KW, Kim S, Tao J, La Rochelle C, et al. Normal

spastin gene dosage is specifically required for axon regeneration. Cell reports. 2012;

2: 1340-50.

16. Rao K, Stone MC, Weiner AT, Gheres KW, Zhou C, Deitcher DL, et al. Spastin,

atlastin, and ER relocalization are involved in axon but not dendrite regeneration.

Molecular biology of the cell. 2016; 27: 3245-56.

17. Lin YF, Xie Z, Zhou J, Chen HH, Shao WW, Lin HD. Effect of exogenous spastin

combined with polyethylene glycol on sciatic nerve injury. Neural regeneration

research. 2019; 14: 1271-9.

18. Martins-Marques T, Ribeiro-Rodrigues T, Batista-Almeida D, Aasen T, Kwak BR,

Girao H. Biological Functions of Connexin43 Beyond Intercellular Communication.

Trends in cell biology. 2019; 29: 835-47.

19. Fu W, Hu W, Yi YS, Hettinghouse A, Sun G, Bi Y, et al. TNFR2/14-3-3ε

signaling complex instructs macrophage plasticity in inflammation and autoimmunity.

The Journal of clinical investigation. 2021; 131.

20. Fu W, Hettinghouse A, Chen Y, Hu W, Ding X, Chen M, et al. 14-3-3 epsilon is

an intracellular component of TNFR2 receptor complex and its activation protects

against osteoarthritis. Annals of the rheumatic diseases. 2021; 80: 1615-27.

21. Nielsen MD, Luo X, Biteau B, Syverson K, Jasper H. 14-3-3 Epsilon antagonizes

FoxO to control growth, apoptosis and longevity in Drosophila. Aging cell. 2008; 7:

688-99.

22. Park E, Rawson S, Li K, Kim BW, Ficarro SB, Pino GG, et al. Architecture of

autoinhibited and active BRAF-MEK1-14-3-3 complexes. Nature. 2019; 575: 545-50.

23. Pisciottani A, Biancolillo L, Ferrara M, Valente D, Sardina F, Monteonofrio L, et

al. HIPK2 Phosphorylates the Microtubule-Severing Enzyme Spastin at S268 for

Abscission. Cells. 2019; 8.

24. Kaplan A, Morquette B, Kroner A, Leong S, Madwar C, Sanz R, et al.

Small-Molecule Stabilization of 14-3-3 Protein-Protein Interactions Stimulates Axon

Regeneration. Neuron. 2017; 93: 1082-93 e5.

25. Yam PT, Kent CB, Morin S, Farmer WT, Alchini R, Lepelletier L, et al. 14-3-3

proteins regulate a cell-intrinsic switch from sonic hedgehog-mediated commissural

axon attraction to repulsion after midline crossing. Neuron. 2012; 76: 735-49.

26. Ji Z, Zhang G, Chen L, Li J, Yang Y, Cha C, et al. Spastin Interacts with CRMP5

to Promote Neurite Outgrowth by Controlling the Microtubule Dynamics.

Developmental neurobiology. 2018; 78: 1191-205.

27. Ji ZS, Li JP, Fu CH, Luo JX, Yang H, Zhang GW, et al. Spastin interacts with
collapsin response mediator protein 3 to regulate neurite growth and branching.
Neural regeneration research. 2021; 16: 2549-56.

28. Patel M, Li Y, Anderson J, Castro-Pedrido S, Skinner R, Lei S, et al. Gsx1
promotes locomotor functional recovery after spinal cord injury. Molecular therapy :
the journal of the American Society of Gene Therapy. 2021; 29: 2469-82.

29. Liu Y, Wang X, Li W, Zhang Q, Li Y, Zhang Z, et al. A Sensitized IGF1
Treatment Restores Corticospinal Axon-Dependent Functions. Neuron. 2017; 95:
817-33.e4.

30. Brown AR, Martinez M. Ipsilesional Motor Cortex Plasticity Participates in
Spontaneous Hindlimb Recovery after Lateral Hemisection of the Thoracic Spinal
Cord in the Rat. The Journal of neuroscience : the official journal of the Society for
Neuroscience. 2018; 38: 9977-88.

31. Sun G, Yang S, Cao G, Wang Q, Hao J, Wen Q, et al. $\gamma\delta$ T cells provide the early
source of IFN-γ to aggravate lesions in spinal cord injury. The Journal of experimental
medicine. 2018; 215: 521-35.

32. Zhou K, Zheng Z, Li Y, Han W, Zhang J, Mao Y, et al. TFE3, a potential
therapeutic target for Spinal Cord Injury via augmenting autophagy flux and
alleviating ER stress. Theranostics. 2020; 10: 9280-302.

33. Kamens HM, Crabbe JC. The parallel rod floor test: a measure of ataxia in mice.
Nature protocols. 2007; 2: 277-81.

34. Jiang T, Cai Z, Ji Z, Zou J, Liang Z, Zhang G, et al. The lncRNA
MALAT1/miR-30/Spastin Axis Regulates Hippocampal Neurite Outgrowth. Frontiers
in cellular neuroscience. 2020; 14: 555747.

35. Riano E, Martignoni M, Mancuso G, Cartelli D, Crippa F, Toldo I, et al.
Pleiotropic effects of spastin on neurite growth depending on expression levels.
Journal of neurochemistry. 2009; 108: 1277-88.

36. Sardina F, Pisciottani A, Ferrara M, Valente D, Casella M, Crescenzi M, et al.

Spastin recovery in hereditary spastic paraplegia by preventing neddylation-dependent

degradation. Life science alliance. 2020; 3.

37. Kaplan A, Morquette B, Kroner A, Leong S, Madwar C, Sanz R, et al.

Small-Molecule Stabilization of 14-3-3 Protein-Protein Interactions Stimulates Axon

Regeneration. Neuron. 2017; 93: 1082-93.e5.

38. Saponaro A, Porro A, Chaves-Sanjuan A, Nardini M, Rauh O, Thiel G, et al.

Fusicoccin Activates KAT1 Channels by Stabilizing Their Interaction with 14-3-3

Proteins. The Plant cell. 2017; 29: 2570-80.

39. Petosa C, Masters SC, Bankston LA, Pohl J, Wang B, Fu H, et al. 14-3-3zeta

binds a phosphorylated Raf peptide and an unphosphorylated peptide via its

conserved amphipathic groove. The Journal of biological chemistry. 1998; 273:

16305-10.

40. Pisa R, Cupido T, Kapoor TM. Designing Allele-Specific Inhibitors of Spastin, a

Microtubule-Severing AAA Protein. Journal of the American Chemical Society. 2019;

141: 5602-6.

41. Pisa R, Cupido T, Steinman JB, Jones NH, Kapoor TM. Analyzing Resistance to

Design Selective Chemical Inhibitors for AAA Proteins. Cell chemical biology. 2019;

26: 1263-73.e5.

42. Verma M, Lizama BN, Chu CT. Excitotoxicity, calcium and mitochondria: a triad

in synaptic neurodegeneration. Translational neurodegeneration. 2022; 11: 3.

43. Savolainen KM, Loikkanen J, Eerikäinen S, Naarala J. Glutamate-stimulated

ROS production in neuronal cultures: interactions with lead and the cholinergic

system. Neurotoxicology. 1998; 19: 669-74.

44. Ankarcrona M, Dypbukt JM, Bonfoco E, Zhivotovsky B, Orrenius S, Lipton SA,

et al. Glutamate-induced neuronal death: a succession of necrosis or apoptosis

depending on mitochondrial function. Neuron. 1995; 15: 961-73.

45. Kajiwara Y, Buxbaum JD, Grice DE. SLITRK1 binds 14-3-3 and regulates

neurite outgrowth in a phosphorylation-dependent manner. Biological psychiatry.

2009; 66: 918-25.

46. Zerr I. Laboratory Diagnosis of Creutzfeldt-Jakob Disease. The New England

journal of medicine. 2022; 386: 1345-50.

47. van den Brand R, Heutschi J, Barraud Q, DiGiovanna J, Bartholdi K, Huerlimann

M, et al. Restoring voluntary control of locomotion after paralyzing spinal cord injury.

Science (New York, NY). 2012; 336: 1182-5.

48. Perrin FE, Noristani HN. Serotonergic mechanisms in spinal cord injury.

Experimental neurology. 2019; 318: 174-91.

49. Yuan A, Rao MV, Veeranna, Nixon RA. Neurofilaments and Neurofilament

Proteins in Health and Disease. Cold Spring Harbor perspectives in biology. 2017; 9.

50. Liu B, Xin W, Tan JR, Zhu RP, Li T, Wang D, et al. Myelin sheath structure and

regeneration in peripheral nerve injury repair. Proceedings of the National Academy

of Sciences of the United States of America. 2019; 116: 22347-52.

51. Ceto S, Sekiguchi KJ, Takashima Y, Nimmerjahn A, Tuszynski MH. Neural Stem

Cell Grafts Form Extensive Synaptic Networks that Integrate with Host Circuits after

Spinal Cord Injury. Cell stem cell. 2020; 27: 430-40.e5.

52. Venkatesh K, Ghosh SK, Mullick M, Manivasagam G, Sen D. Spinal cord injury:

pathophysiology, treatment strategies, associated challenges, and future implications.

Cell and tissue research. 2019; 377: 125-51.

53. Ko SH, Apple EC, Liu Z, Chen L. Age-dependent autophagy induction after

injury promotes axon regeneration by limiting NOTCH. Autophagy. 2020; 16:

2052-68.

54. Griffin JM, Bradke F. Therapeutic repair for spinal cord injury: combinatory

approaches to address a multifaceted problem. EMBO molecular medicine. 2020; 12:

e11505.

55. Sarbanes SL, Zehr EA, Roll-Mecak A. Microtubule-severing enzymes. Current

biology : CB. 2022; 32: R992-r7.

56. Vemu A, Szczesna E, Zehr EA, Spector JO, Grigorieff N, Deaconescu AM, et al.

Severing enzymes amplify microtubule arrays through lattice GTP-tubulin

incorporation. Science (New York, NY). 2018; 361.

57. Costa AC, Sousa MM. The Role of Spastin in Axon Biology. Frontiers in cell and

developmental biology. 2022; 10: 934522.

58. Zhang J, Pho V, Bonasera SJ, Holtzman J, Tang AT, Hellmuth J, et al. Essential

function of HIPK2 in TGFbeta-dependent survival of midbrain dopamine neurons.

Nature neuroscience. 2007; 10: 77-86.

59. Cornell B, Wachi T, Zhukarev V, Toyo-Oka K. Regulation of neuronal

morphogenesis by 14-3-3epsilon (Ywhae) via the microtubule binding protein,

doublecortin. Human molecular genetics. 2016; 25: 4405-18.

60. Dar A, Wu D, Lee N, Shibata E, Dutta A. 14-3-3 proteins play a role in the cell

cycle by shielding cdt2 from ubiquitin-mediated degradation. Molecular and cellular

biology. 2014; 34: 4049-61.

61. Lee JH, Lu H. 14-3-3Gamma inhibition of MDMX-mediated p21 turnover

independent of p53. The Journal of biological chemistry. 2011; 286: 5136-42.

62. Chan TY, Egbert CM, Maxson JE, Siddiqui A, Larsen LJ, Kohler K, et al. TNK1

is a ubiquitin-binding and 14-3-3-regulated kinase that can be targeted to block tumor

growth. Nature communications. 2021; 12: 5337.

63. Kast DJ, Dominguez R. Mechanism of IRSp53 inhibition by 14-3-3. Nature

communications. 2019; 10: 483.

64. Cornell B, Toyo-Oka K. 14-3-3 Proteins in Brain Development: Neurogenesis,

Neuronal Migration and Neuromorphogenesis. Frontiers in molecular neuroscience.

2017; 10: 318.

65. Marra M, Camoni L, Visconti S, Fiorillo A, Evidente A. The Surprising Story of

Fusicoccin: A Wilt-Inducing Phytotoxin, a Tool in Plant Physiology and a

14-3-3-Targeted Drug. Biomolecules. 2021; 11.

66. Liu D, Bienkowska J, Petosa C, Collier RJ, Fu H, Liddington R. Crystal structure

of the zeta isoform of the 14-3-3 protein. Nature. 1995; 376: 191-4.

1048

1049

 Figure legends:

 Figure 1. 14-3-3 physically interacts with spastin *in vivo* and *in vitro*. (A) Proteins

 from the spinal cord tissue (approximately 1 cm around the T10 level) were

 pull-downed by purified GST and GST-Spastin proteins and the proteins obtained

 were subjected to mass spectrometry analysis. The 306 peptides were identified in the

 GST group and 699 peptides in the GST-Spastin group. The peptides that were

 common to both the GST and GST-Spastin groups were excluded, and 531 peptides

 were the peptides of proteins that interact with spastin. Three 14-3-3 protein peptides

 were shown. (B) Immunoprecipitation assay using spastin antibody was performed

 from spinal cord lysates and Pan 14-3-3 antibody was used for western blotting. (C)

 Immunoprecipitation assay using spastin antibody was performed from cortical neural

 lysates and Pan 14-3-3 antibody was used for western blot. (D) Hippocampal neurons

 were stained with 14-3-3 (green) and spastin (red) antibodies, the growth cone (D1)

 and axon shaft (D2) profile were shown. (E) Localization of spastin (red line) and

 14-3-3 (green line) protein in the hippocampal neurons were analyzed and the

 intensity of the white dotted line across the neurite compartment were measured. (F)

 The highly conserved spatial conformation of different 14-3-3 protein isoforms were

 shown. (G) Purified GST and GST-14-3-3s fusion proteins were subjected to sodium

 dodecyl-sulfate polyacrylamide gel electrophoresis and coomassie blue staining was

 confirmed the successful purification of relative proteins. (H) GST fusion proteins

 were used to pull down the lysates of the spinal cord and spastin antibody was used

 for western blotting. (I) HEK 293T cells were transfected with Flag tagged 14-3-3s

 and GFP tagged spastin, the cell lysates were subjected to GFP-Trap assay, Flag

 antibody was used to detect the presence of 14-3-3s.

 Figure 2. Ser233 in spastin is essential for the interaction with 14-3-3 and

 important for neurite outgrowth. (A) Schematic illustration of the gene truncation

 strategy for the modular domain of spastin. (B) Truncation mutants of spastin together

 with Flag tagged 14-3-3 were transfected into HEK 293T cells, and cell lysates were

immunoprecipitated by GFP-Trap assay. Flag antibody were used for detecting the flag tagged 14-3-3 fusion protein. (C) The microtubule binding domain (MTBD) truncated GFP tagged spastin were transfected together with Flag tagged 14-3-3 into HEK 293T cells, and subjected to GFP trap assay. (D) GFP tagged spastin was transfected into HEK293T cells by treatment with staurosporine (a broad spectrum protein kinase inhibitor) or not. GFP-Trap beads were used to enrich GFP-Spastin, and the phosphorylation levels of spastin were detected using a Pan phosphoserine/threonine antibody. (E) HEK 293T cells were transfected with GFP-spastin and then treated with or without staurosporine for 1 hour. Cell lysates were pull downed by GST 14-3-3 fusion protein and GFP antibody were used for western blotting. (F) The binding sites between 14-3-3 and spastin were predicted using the scansite 2.0 software and whether these sites could be phosphorylated were confirmed by phosphosite plus website. (G) Serine 233 and Serine 562 of spastin were mutated to alanine, these GFP fusion mutations were transfected together with Flag tagged 14-3-3, then cell lysates were subjected to GFP-trap assay. Flag antibody was used to detect the binding ability between spastin mutations and 14-3-3. (H) GFP tagged wild-type spastin, spastin S233A (mimicking dephosphorylation), or spastin S233D (mimicking phosphorylation) were co-transfected with Flag tagged 14-3-3 into HEK293T cells. The group transfected with GFP-Spastin S233D was additionally treated with staurosporine. Cells were lysed and immunoprecipitation were conducted using GFP-Trap, and western blot analysis was performed using Flag antibodies to investigate the impact of Ser233 phosphorylation in spastin on its interaction with 14-3-3. (I) Hippocampal neurons were grown on DIV 2 and transfected with spastin Ser233 mutations (Spastin S233A, Spastin S233D), then fixed at DIV 3. Representative images of transfected neurons as indicated were shown. Quantitative analysis of the length of neurites (J) and the number of neurites (K) were shown. Data were represented as mean value of three independent experiments. All error bars are SEM. *P <0.05. Scale bar: 100 μm.

Figure 3. 14-3-3 protein protects spastin by Ser233 phosphorylation from degradation through ubiquitin pathway. (A,B) GFP tagged spastin together with Flag or with Flag tagged 14-3-3 was transfected into HEK 293T cells. Following the addition of protein synthesis inhibitor (cycloheximide, CHX), cell lysates were subjected to western blot to detect the GFP tagged spastin protein levels. (C, D) Western blot analysis of HEK 293T cell lysates transfected with spastin mutations (S233A or S233D) after treating with cycloheximide at different indicated times. (E,F) GFP tagged spastin with Flag or Flag tagged 14-3-3 were co-transfected into HEK293T cells. Cell protein lysates from different time points were collected for western blot. (G) GFP tagged spastin S233D was transfected into HEK293T cells, and western blot experiments were conducted to observe the protein expression levels of GFP tagged spastin at different time points. (H) Quantitative analysis of spastin protein levels after addition of cycloheximide with or without 14-3-3 protein. (I) Quantitative analysis of GFP tagged spastin protein levels upon transfected with different spastin mutations. (J) Quantitative analysis of spastin protein levels with or without 14-3-3 protein. (K) Quantitative analysis of GFP-Spastin (S233D) protein levels at different time points. (L) Cultured cortical neurons were treated with proteasome inhibitor MG132 or/and CHX for 36 h, cell lysates were subjected to western blot analysis. (M) Quantification analysis of spastin protein levels. (N) HEK 293T cells were transfected with GFP tagged spastin or its S233 mutants and Flag tagged ubiquitin, together with R18 or not, cells were harvested and subjected to immunoprecipitation using GFP antibody. Flag antibody was used to detect the ubiquitin band. (O) Quantitative analysis of the ubiquitination levels of GFP-Spastin. (P) COS7 cells were transfected with indicated GFP tagged spastin with or without Flag tagged ubiquitin for 24 hours. Then, cells were fixed and stained with tubulin antibody to visualize microtubule arrangements. (Q) The normalized quantitative analysis of microtubule fluorescence intensity among indicated groups. Data were represented as mean value of three independent experiments. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. Scale bar: 50 µm, 10 µm.

1137

Figure 4. 14-3-3 agonist Fusicoccin (FC-A) promotes the repair of neurite outgrowth and regeneration after injury via spastin. (A) The surface structure and secondary structure of molecular docking model between 14-3-3 protein with FC-A (PDB:2O98) or R18 (PDB:1A38). (B) HEK 293T cells were transfected with GFP tagged spastin and Flag tagged 14-3-3, cell lysates were incubated with FC-A or R18 and then subjected to GFP-trap assay. Flag antibody were used to detect the 14-3-3 binding. (C) Hippocampal neurons (DIV2) were transfected with GFP to visualize its morphology. Neurons were then incubated with glutamate (120 μM) to induce injury, the neurite breakages and swellings were found in C1 and C2. (D) The injured neurons were transfected with spastin and incubated with FC-A, R18 or spastazoline for 24 hours, then cells were fixed and images were taken by confocal microscopy. Neurons were traced using Image J Pro Plus and quantitative analysis of the length of primary neurites (E), secondary neurites (F) and total neurites (G) were performed. Quantitative analysis of the number of primary neurites (H), secondary neurites (I) and total neurites (J). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. Scale bar: 50 μm. (K) Primary cortical neurons were grown and scratch assays were performed at DIV7. FC-A or spastazoline were applied and incubated for 36 hours. Then, cells were stained with βIII-tubulin to visualize the morphology of neurites. (L) Cortical neurons cultured in vitro were treated with FC-A for 36 hours, and western blot experiments were conducted to observe the protein levels of spastin. (M) Quantitative analysis of the normalized axon regeneration rate. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. Scale bar: 100 μm.

1160

Figure 5. 14-3-3/spastin pathway is involved in nerve regeneration after spinal cord injury by targeting MTs. (A) After spinal cord contusion, the lesion site of spinal tissue in the sham group and the SCI group were stained with Pan 14-3-3 (red) and βIII tubulin (green). The arrows mean the elevated 14-3-3 protein in the neuronal compartment. (B)(C) After spinal cord contusion, the spinal cord tissues of the lesion

site (near 1 cm) at indicated times (1, 3, 7 14, and 30 DPI) were ground and lysed, then subjected to western blot with 14-3-3 and spastin antibody. (D) After spinal cord injury, intraperitoneal injections of FC-A were administered. Tissue from the lesion site was collected at indicated time points and subjected to western blot to observe the spastin protein levels. (E,F) Quantitative analysis of the 14-3-3 protein expression and spastin protein expression (n=3 per group). (G) Quantitative analysis of the spastin protein expression administrated with FC-A after spinal cord injury (n=3 per group). (H) Analysis of the expression trends of 14-3-3 and spastin proteins following spinal cord injury. (I) Trend analysis of spastin protein expression levels in lesion site of mice with spinal cord injury, comparing the injury control group with the FC-A treatment group. (J) 14-3-3 agonist FC-A and spastin inhibitor spastazoline were admisnistrated after spinal cord contusion. At 45 DPI (Days post-injury), the tissues were fixed and embedded, then cut horizontally and longitudinally. H&E and LFB (Luxol fast blue stain) were stained. The spinal cord demyelination was shown inside the area of the dotted line. Scale bar , 200 μm. (K) Quantitative analysis of demyelinated area in the lesion site among different groups. (L) The slices were then subjected to immunofluorescence staining, and GFAP (green) was stained to label astrocytes, NF (red) was stained to label neurons and 5-HT (magenta) was stained to label monoaminergic axons. The lesion site in different groups was boxed and enlarged in (L1)(L2)(L3)(L4). Scale bar, 400 μm, and 100 μm. (M) The spinal cord slices were stained with acetylated tubulin (Stable MTs which lack of dynamics) and β-tubulin (total tubulin). Scale bar , 50 μm. (N) The normalized intensity of NF was quantified (n=5 animals per group). Mean ± SEM. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. (O) The normalized ratio of acetylation tubulin to total tubulin was calculated by the intensity value of acetylated tubulin divided by total tubulin (n=5 per group). Mean ± SEM. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Figure 6. 14-3-3/spastin pathway coordinates locomotor recovery after spinal cord contusion. (A) Schematic illustration of relative treatments and examinations at

different indicated times. (B) The max contact areas of the right hindlimb were recorded and analyzed by the Catwalk software. (C) The footfall patterns were visualized by the Catwalk software. The blue dot represents the start of a pattern. The red dot represents the gait was not part of a pattern. The yellow dot represents that the gait was not taken into account. The green one means the gait belongs to a normal part of a pattern. (D) BMS score at different indicated days after injury and data were represented as mean±SEM. (n=5 group, data were analyzed by one way ANOVA and Bonferroni's multiple-comparison test were used for post hoc comparisons. (E) The quantitative analysis of the hindlimb (right) max contact area. (n=5 animals per group). (F) Footprint analysis of the mice at 6 weeks after spinal cord contusion. The forelimbs were marked with blue and the hindlimbs were marked with red. Scale bar: 2.5 cm. (G) The normalized regularity index of the footfall pattern calculated by Catwalk software (~100 % in the sham group, n=5 animals per group). (H) Quantatitive analysis of the stride length in the footprint assay. The mean stride length of each walk was counted, n=5 animals per group). (I) Motor evoked potentials (MEPs) were recorded at 7 weeks after spinal cord contusion. The craniotomy was performed and the M1 region of the sensorimotor cortex were stimulated by single square-wave stimulus of 0.5 mA, 0.5 ms duration, 2 ms time delay, and 1 Hz. The MEP were recorded with the signals detected by the electrode on the gastrocnemius muscle. The representative histograms of the amplitude-times are shown. The arrows indicated the stimulus. (J) The current amplitudes of the MEP were quantified, n=5 animals per group. Mean ± SEM. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$.

Figure 7. FC-A promotes axon regeneration in a T10 lateral hemisection spinal cord injury model. (A) The spinal cords of adult mice were subjected to lateral hemisection and administrated by intraperitoneal injection with FC-A or spastazoline. Sagittal view of 5-HT-immunoreactive raphespinal fibers in the spinal cord on 17 DPI. Scale bar, 500 μm and 100 μm. (B) Sagittal view of neurofilaments which was MBP positive in the white manner of the lesion site of the spinal cord. Scale bar, 50 μm. (C)

The normalized quantification of 5-HT immunoreactive fluorescence intensity (0.5

mm caudal to the lesion site, n=5 animals per group). (D) The normalized

quantification of NF (Neurofilament) immunoreactive fluorescence intensity in the

lesion site (n=5 animals per group). (E) The locomotor function of mice after SCI

were scored at the indicated time points according to the standard of Basso Mouse

Scale (BMS). (F) Foot fault test was performed and the total steps and steps drop out

of the right hindlimb were recorded. The rate of the foot errors was analyzed, n=5

animals per group. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. (G) (H)The footprint assay

was performed and stride length (G) and stride width (H) quantitative analyzed. Data

were presented as mean±SEM, n=5 animals per group, the mean value of each animal'

s walk were calculated.

1235

1236

1237

A LC-MS from proteins of GST pull-down
Number of peptides identified

Species: Rattus

14-3-3 peptide sequence

1.LAEQAER

2.NLLSVAYK

3.AVTEQGAELSNEER

GST 138 (16.5%) 168 (20.1%) GST-Spastin 531 (63.4%)

B Spinal cord lysates

IP: IgG Spastin 4% Input

IB: Pan14-3-3 — 33

IB: Spastin — 70

C Cortical neuronal lysates

IP: IgG Spastin 4% Input

— 33

— 70

D

D1 D2

Pan 14-3-3 Spastin DAPI Merged 50 μm

D1 D2 5 μm

E

Intensity(a.u.) vs Distance

1 2 3 4 5 6

Spastin (red), 14-3-3 (green)

pixels(a.u.)

F 14-3-3 protein isoforms 3D structure (PDB number, isoform)

2C23,β 6BYJ,γ 3UAL,ε

2BTP,θ 2C63,η 6ZFG,ζ

G

GST GST-14-3-3

MW: β γ ε ζ η θ

H GST-14-3-3 Pull-down

GST GST-14-3-3

MW: β γ ε ζ η θ 4% lysate

IB: Spastin

I GFP-Spastin co-overexpressed with

Flag Flag-14-3-3

IP:Flag β γ ε ζ η θ

IB:GFP 95 70

IB:Flag 43 35

Input 95 70