Disease exacerbation by fibroblast inclusion in Duchenne Muscular Dystrophy MYOrganoids reveals limitations of microdystrophin therapeutic efficacy

Laura Palmieri [1,2], Louna Pili[1,2], Abbass Jaber, [1,2] Ai Vu Hong [1,2], Matteo Marcello[1,2], Riyad El-Khoury [1,2,3], Guy Brochiet [3,4], Anne Bigot [5], David Israeli [1,2], Isabelle Richard [1,2], Sonia Albini [1,2*]

[1]Genethon, 91100 Evry, France; [2]Université Paris-Saclay, Univ Evry, Inserm, Généthon, Integrare research unit UMR_S951, 91000, Evry, France. [3]Institut de Myologie, Neuromuscular Morphology Unit, Groupe Hospitalier Pitié-Salpêtrière, Paris, France.[4]AP-HP, Centre de Référence de Pathologie Neuromusculaire Nord/Est/Ile de France, Groupe Hospitalier Pitié-Salpêtrière, Paris, France. [5]Sorbonne Université, Inserm, Institut de Myologie, Centre de Recherche en Myologie, F-75013 Paris, France.

[*] **Corresponding author**. Email: salbini@genethon.fr, ORCID: 0000-0001-9502-1004

SUMMARY

The yet incurable Duchenne muscular dystrophy (DMD) is caused by the absence of dystrophin, a protein essential to preserve muscle integrity continuously challenged by contractions. Adeno-associated virus (AAV) delivery of truncated forms of dystrophin is currently the most promising therapeutic approach; however, patient outcomes differed from animal studies, emphasizing the necessity for models predictive of human response. Here, we describe the generation of MYOrganoids, a 3D muscle platform derived from human induced pluripotent stem cells (iPSC), whose structural and functional maturation is enhanced by fibroblasts incorporation. Importantly, a pro-fibrotic signature, elicited by fibroblast contact with dystrophic muscle cells, was pivotal to exacerbate muscle force loss and fatiguability of DMD MYOrganoids, enabling their use as therapeutic readouts. Remarkably, gene transfer of the gold standard microdystrophin (μDys) in DMD MYOrganoids significantly improved muscle resistance, partially rescued membrane stability while failing to reduce fibrotic traits, in line with the marginal correction of the DMD transcriptional signature achieved. In particular, the persistent profibrotic activity of fibroblasts within μDys-treated DMD MYOrganoids, indicates lack of effect from μDys receiving cells. This study highlights human MYOrganoids' potential to unravel limitations of current treatments and investigate alterations in cell communications under fibrotic conditions, thereby accelerating the discovery of more effective strategies.

Keywords: iPSC, DMD, fibroblasts, MYOrganoids, AAV, microdystrophin

Introduction

Duchenne muscular dystrophy (DMD; ONIM: #310200) is an X-linked disorder that affects one in every 5000 male births [1] with no resolutive cure up to date. It is characterized by progressive muscle wasting affecting skeletal muscles primarily and cardiac and respiratory muscles later, thereby causing premature death [2]. DMD is caused by genetic mutations in the *DMD* gene, leading to the absence of Dystrophin, an essential protein that provides physical support to myofibers by linking them to the extracellular matrix through the Dystrophin Glycoprotein Complex (DGC) [3–6]. The lack of Dystrophin results in a series of muscle membrane breakdowns and repairs, leading to subsequent secondary issues like chronic inflammation and fibrosis [7–10]. Fibrosis, an excessive deposition of extracellular matrix components, is a critical driver of DMD progression. Indeed, the secretion and the production of TGF-β (Transforming Growth Factor beta) stimulates the production of matrix components, like fibronectin and collagen, from fibroblasts and myofibers [11–13].

Gene therapy using adeno-associated virus (AAV) is currently the most promising treatment for Duchenne muscular dystrophy. Ongoing clinical trials use AAV to deliver short forms of Dystrophin, known as microdystrophin (μDys), which encodes a truncated but functional protein [14–20]. However, while the therapeutic effects were unequivocally achieved in DMD animal models [21], the results from clinical trials revealed only partial therapeutic efficacy in terms of gain of muscle function, besides the appearance of unexpected safety issues [22,23]. These observations confirm the limited translatability of results obtained in animal models to human patients. It appears therefore crucial to develop high throughput models mimicking human DMD pathology, suitable to quickly assess the efficacy of therapeutics, more reliably and with reduced animal waste and costs. In this context, *in vitro* modeling based on human cells is a valuable option. In particular, the induced pluripotent stem cells (iPSC) technology offers the opportunity to derive an unlimited number of specialized cells from patients for disease modeling and drug screening [24,25]. Among the *in vitro* cellular models, organoid-like structures are becoming invaluable for disease modeling as the use of 3D cultures and biomaterials allows the

reconstitution of tissue architecture and microenvironment that are instrumental for pathophysiological

evaluations [26,27].

Although various *in vitro* 3D systems are accessible for modeling Duchenne muscular dystrophy in

skeletal muscle engineering [28–31] their throughput use is limited by the long duration and variability

related to the complexity of cellular composition achieved. Additionally, the absence of disease-specific

readouts for muscle function analysis readouts is still a challenge for assessing the effectiveness of

therapeutics. It is also worth noting that tissue engineering applications for AAV gene therapy have been

exploited mostly in the context of retinopathies [25,32–34] while only limitedly explored for muscular

disorders, for which a high-throughput human muscle system would be an invaluable resource for gene

therapy screening.

Here, we described the rapid generation of iPSC-derived muscle organoid structure, named hereafter

MYOrganoids. We employed and adapted an engineered muscle platform to generate MYOrganoids

using a previously reported method for direct iPSC conversion into 2D skeletal muscle cells [35–37]. We

further included fibroblasts to increase the structural and functional maturation required for

pathophysiological studies. As such, fibroblasts and ECM compose the connective tissue which is a key

regulator of differentiation and muscle structure as well as of the muscle niche that undergoes

pathological remodeling during disease [38,39]. In this context, fibroblasts play a major role in muscle

assembly and act as a source of microenvironment cues exerted by their secretory activity [38,40–

42]Additionally, we demonstrated that fibroblast inclusion and direct interaction with muscle cells, during

iPSC-derived MYOrganoid generation, was required to reveal key DMD hallmarks related to loss of

muscle resistance and fibrotic signature.

Our study also evaluates for the first time the therapeutic efficacy of AAV-mediated µDys gene transfer,

in engineering muscle tissues, as proof of concept of their suitability for studying disease mechanisms

and evaluating potential therapeutics. As such, while AAV9 infectivity in 3D engineered muscles has

been already evaluated [43], in vitro µDys AAV gene transfer, to the best of our knowledge, has not yet

been reported. By using different doses of µDys in DMD MYOrganoids, we observed a dose-dependent

response in restoring muscle function while only a partial effect at the level of membrane stability and

fibrotic signature in DMD muscles. Our findings indicate that patient-derived MYOrganoids, whose

pathogenic traits are exacerbated, are suitable for studying the fibrotic process orchestrated by either

muscle or fibroblast population and its interplay with gene transfer approaches. Our system has therefore

the potential to identify molecular mechanisms driving the dystrophic process and accelerate the

identification of effective therapeutics for DMD.

Results

Generation of structurally organized 3D human MYOrganoids by direct conversion of iPSC and inclusion of fibroblasts

MYOrganoids were generated from human iPSC committed to differentiating into the myogenic lineage by inducible expression of MyoD and BAF60C [35], from now on referred to as iPSC$^{BM.}$. MYOrganoids were prepared starting from iPSC^{BM} after one day from the induction of myogenic genes. Casting procedure was performed through adaptation of an engineered muscle system [31,44] which results in the growth of the tissue in a ring format supported by two flexible silicon stretchers. The 3D cultures were kept for 2 days in growth medium, afterwards medium was replaced for differentiation for another 12 days (**Figure 1A**). The differentiation protocol was optimized from the conditions previously reported [11,36,37] using myogenic commercial media that ensured the highest expression of myogenic markers and myogenic differentiation in monolayer conditions (**Figure S1A-B**). Since cellular heterogenicity, especially of mesenchymal origin, is important for muscle formation [31,38,39,45], we included human fibroblasts during the casting procedure, to assess whether this would affect muscle organization. For this aim, casting was performed using iPSC^{BM} cells in the presence or absence of human fibroblasts. Achieving alignment and differentiation simultaneously necessitates a delicate balance between fibroblasts and muscle cells (as noted by N. Rao et al., 2013). To replicate physiological conditions accurately, we incorporated a fibroblast concentration that mirrors the stromal population detectable through single cell and single nuclei-RNA seq analysis of muscles [46–49]. We found that including 10% fibroblasts, accelerated the condensation and growth over time of the muscle rings into a compact structure 0.8 mm long and 1 mm thick at day 14 (**Figure 1B).** By performing immunofluorescence in whole-mount tissues for sarcomeric α-actinin (SAA), we could detect an enrichment of SAA-positive myotubes throughout the ring-shaped micro-tissue (**Figure 1C**). We then aimed to assess the impact of fibroblast inclusion on muscle structure. Since the organization of muscle cells within the ECM plays a key role in fusion and maturation [50], both alignment of myotube and circularity, a consequence of their

parallelism, were evaluated. Staining for myosin heavy chain (MyHC, myotube marker) and vimentin

(fibroblasts marker) on longitudinal sections showed fibroblast recruitment near muscle fibers (**Figure**

1D). Myotubes alignment was determined by measuring the angles in between myotubes [51] and showed

a significant decrease towards 0 degrees upon fibroblast inclusion, which indicates parallelism while, in

the MYOrganoids without fibroblasts, we detected a disordered pattern of muscular cells (**Figure 1E**).

Additionally, the circularity of myofibers was measured from transversal sections stained for the

membrane marker wheat germ agglutinin (WGA), using the ratio between X and Y Feret diameters

(**Figure 1D**). MYOrganoids including fibroblasts had an improved circularity (ratio closer to 1) when

compared to control (**Figure 1E**). Improved myotube circularity and alignment as shown, indicate that

fibroblast incorporation during the casting procedure guides skeletal cell orientation providing structural

support for MYOrganoids, a prerequisite for maturation.

Figure 1. Generation of iPSC-derived MYOrganoids and impact of fibroblast inclusion on muscle organization. (A) Scheme of the protocol used to generate muscle artificial tissues (MYOrganoids) from iPSC committed towards the myogenic lineage by 24h treatment with doxycycline for inducible expression of Myod and BAF60C transgenes (iPSC^BM). The casting procedure included: committed iPSC, fibroblasts when indicated (+/-fibro) and a collagen-based scaffold, within a 48-well plate equipped with silicon pillars. After 2 days in growth medium, the 3D structures were shifted to differentiation medium until day 14 for histological analysis. (B) Condensation kinetics of MYOrganoids +/-fibro. (C) Whole-mount staining of MYOrganoids with SARCOMERIC α-ACTININ (SAA) and 3D reconstruction of the ring-shaped constructs using confocal imaging. Scale bar: 1 mm. (D) Representative longitudinal and cross sections of MYOrganoids +/- fibroblasts, immunostained for Myosin Heavy Chain (MYHC) and VIMENTIN or for wheat germ agglutinin (WGA). Nuclei were visualized with DAPI. Scale bars 200 µm (left panel) and 10 µm (middle and right panel). Arrows indicate fibroblasts recruited adjacently to the muscle fibers; α is the angle formed between myotubes; lines indicate aligned myotubes, crosses represent X/Y myotubes diameters. (E) The alignment was calculated based on the angle (α) formed between myotubes (α close to 0 corresponds to aligned myotubes, while far from 0 corresponds to not aligned myotubes), while circularity from X/Y myotubes diameters ratio (ratio 1 circular, far from 1 not circular). Ratio X/Y diameter is represented by a QQ normality plot. Data were collected from 3 independent experiments with at least 3 replicates. N=35. Unpaired two-tailed t-test was used (***p ≤ 0.001).

Increased structural and functional maturation of fibroblast-including MYOrganoids

Since muscle maturation is strictly dependent on the internal myofiber organization, we evaluated the differentiation of our 3D MYOrganoids by looking at the sarcomere structure. Transversal and longitudinal sections were used to monitor dystrophin (DYS) expression at the sarcolemma (**Figure 2A**) and sarcomeric α-actinin (SAA) localization for assessment of the striation pattern typical of mature myotubes (**Figure 2B**). Dystrophin was properly localized to the muscle membrane of myotubes from MYOrganoids including fibroblasts and was significantly more expressed than MYOrganoids without fibroblasts (**Figure 2D**). Remarkably, the maturation index, reported as a percentage of the number of nuclei included in striated myotubes, was significantly superior in MYOrganoids including fibroblasts as compared to MYOrganoids without fibroblasts which appear very disorganized with a rare appearance of striations (**Figure 2E**). Increased maturation of myotubes within MYOrganoids including fibroblast was also supported by analysis of the fusion index, indicating a significantly higher percentage of multinucleated myotubes (>3 nuclei) and a lower percentage of mononucleated ones (**Figure 2F**). This evidence highlights the positive role of fibroblasts in the maturation process through fusion and multinucleation. The proper sarcomeric organization was also confirmed by electron microscopy where we could detect longer, properly formed Z patterning and the presence of I and A bands along with an overall increase of sarcomeric density and alignment (**Figure 2C**). Consistently, MYOrganoids containing fibroblasts showed wider Z-line (**Figure 2H**), index of higher maturation of sarcomeres [52]. We further performed gene expression analysis for terminal differentiation markers such as muscle creatine kinase (*MCK*), myosin heavy chain (*MYH*) isoforms, such as *MYH2*, representative of fast adult fiber type, and *MYH7*, as a slow fibers marker (**Figure 2G**). Higher expression of all genes in MYOrganoids containing fibroblasts confirms the acquisition of a more mature state, compared to MYOrganoids without fibroblasts.

We then assessed whether our MYOrganoids were functional by evaluating their physiological response to contraction stimulations, using a muscle organ bath system based on electrical pacing [53]. To evaluate

muscle force, MYOrganoids were transferred to the muscle strip chamber and stretched until the optimal

length (Lo) for functional analysis (**Figure 2I**). Isometric force analysis revealed significantly higher

tetanic force in MYOrganoids containing fibroblasts compared to the control (**Figure 2J**). Values were

then normalized for the cross-sectional area (CSA) using the weight and optimal length of contraction

established for each MYOrganoid [41,54] and expressed as specific tetanic force (mN/mm^2) (**Figure 2K**).

In particular, the highest force with fibroblasts had peak values ranging from 0.3 to 0.5 mN versus 0.1

to 0.2 mN in MYOrganoids without fibroblasts after normalization (**Figure 2K**). These data

demonstrated that MYOrganoids plus fibroblasts have an improved structural organization and

functional maturation that enables force contraction studies by electrical pacing.

*Figure 2. Structural and functional maturation in fibroblast-including MYOrganoids. (**A**) Representative transversal sections stained for Dystrophin (DYS). Scale bar: 40 µm. Nuclei were visualized with DAPI. (**B**) Representative longitudinal sections of MYOrganoids +/- fibroblasts (fibro), immunostained for Sarcomeric α-Actinin (SAA) and nuclei visualized with DAPI. Scale bars: 40 µm, enlargement 10 µm. (**C**) Transmission electron microscopy images showing sarcomeric structures. Orange arrows: Z-lines. Scale bar: 500 nm. (**D**) DYSTROPHIN staining quantification represented as mean intensity fluorescence and expressed as fold change to the negative control (sections stained without first antibody). N=2 (**E**) Quantification of maturation index was performed by calculating the nuclei inside striated myofibers visualized by SAA staining as a percentage of the total number of myofibers. N=3-5 (**F**) Quantification of fusion index calculated as % Myosin Heavy Chain positive (MYHC+) myotubes containing different numbers of nuclei (n) as depicted. N=3 (**G**) Gene expression analysis of MCK, MYH2 and MYH7, reported as gene expression relative to MYOD expressing population. N=2-5 (**H**) Width of Z-line visualized by Electron microscopy, in MYOrganoids with and without inclusion of fibroblasts. N=16 (**I**) Contractile muscle force analysis of MYOrganoids using a muscle-strip-based organ bath system. Lo, the optimal length used for normalization of force data (see methods). (**J**) Representative tetanic traces of MYOrganoids with and without the inclusion of fibroblasts. N=3 (**K**) Normalized tetanic force peak in MYOrganoids +/- fibroblasts. N=8-15). Data are presented as means +/- SEM. Unpaired t-test was applied for statistical analysis (*p ≤0.05, **p ≤ 0.01, ***p ≤ 0.001, ****p ≤ 0.0001).*

Inclusion of fibroblasts in DMD iPSC-derived MYOrganoids leads to increased muscle fatigue and pro-fibrotic signature

The improved muscle organization and functional maturation shown by MYOrganoids including fibroblasts, prompted us to exploit fibroblast features in disease modeling for DMD, where their role in disease progression is well known [12,13]. We incorporated DMD fibroblasts to recapitulate the pathogenic microenvironment arising from their profibrotic activity exerted by tissue remodeling and matrix deposition [13,55]. For that purpose, we used three DMD iPSC with different *DMD* mutations, a deletion of exon 45 (DMDdEx45), a deletion of exons 8-43 (DMDdEx8-43) and a deletion of exons 8-9 (DMDdEx8-9) with their isogenic control, the DMD dEx6-9 iPSC corrected to restore dystrophin expression [56] (hereafter called IsoCTR). We additionally used two control iPSC lines derived from healthy patients (Ctr1, Ctr2). Control and DMD MYOrganoids were generated from control and DMD iPSC with healthy or DMD human immortalized fibroblasts for functional and transcriptomic analysis (**Figure 3A**). The myogenic differentiation ability of both Ctr and DMD iPSC lines was first evaluated in 2D cultures. All cell lines showed comparable myogenic potential, as expected using a direct myogenic conversion protocol bypassing any defective developmental steps (**Figure S2A**). Histological characterization in MYOrganoids showed the absence of dystrophin protein in DMD MYOrganoids and efficient myogenic differentiation and maturation in Ctr and DMD MYOrganoids, as shown by the striated pattern visualized by SAA-stained sections, confirming the positive role of fibroblasts in enhancing maturation also in a DMD context (**Figure 3B**).

To assess whether DMD MYOrganoids display hallmarks of DMD pathophysiology, we evaluated muscle function, which represents one of the most difficult challenges in establishing therapeutic readouts with *in vitro* systems. To identify reliable force parameters reflecting the defective DMD muscle performance, MYOrganoids were subjected to isometric contractions to measure tetanic force, and to eccentric contractions, which plays a critical role in the disease progression of DMD[57] by triggering the membrane degeneration/regeneration cycles[58] to assess muscle strength and fatiguability.

No significant changes were observed in the tetanic isometric force between Ctr and DMD

MYOrganoids (**Figure S2B**), as expected for muscles never been challenged by contractions and with

the same myogenic potential, while DMD MYOrganoids showed higher drop force in the eccentric

contraction repetitions exclusively when including DMD fibroblasts starting from the 4th eccentric

repetition (**Figure 3C**). On the other side, Ctr and DMD MYOrganoids not including fibroblasts, show

significant differences in drop force just at the tenth repetition (**Figure 3C**). To accurately quantify

muscle fatigue, we calculated the fatigue index as the drop of force between the isometric contractions

performed before and after the 10 repetitions of eccentric exercise. The analysis showed a significantly

higher fatigue index in DMD MYOrganoids as compared to Ctr MYOrganoids and remarkably, this

phenomenon was highly accentuated by the presence of fibroblasts. (**Figure 3D**). To check whether the

impact on disease exacerbation and increase of difference in fatiguability between Ctr and DMD

MYOtissue was affected by the different genetic backgrounds of the fibroblast source, we employed

three additional fibroblast cell lines from either healthy and DMD individuals to generate Ctr and DMD

MYOrganoids. Muscle force analysis confirmed that the fatigue index in DMD MYOrganoids was

significantly higher than Ctr MYOrganoids, regardless of the genetic source of either control or DMD

fibroblasts. This analysis also confirmed the display of higher and significant muscle fatiguability in

fibroblasts-including MYOrganoids (**Figure S2C**). Collectively, these data indicate that DMD

MYOrganoids including DMD fibroblasts, through their pro-fibrotic activity, display exacerbated loss

of muscle resistance and increase in fatiguability (**Figure 3A-D**). The finding also demonstrates that

eccentric-based drop force evaluation is a meaningful and reliable therapeutic readout, as significant

differences were detected between Ctr and DMD MYOrganoids including fibroblasts.

We then performed transcriptomic analysis in the isogenic iPSC lines (IsoCTR and DMDdEx8-9) to

better characterize the pathogenic hallmarks introduced by fibroblast incorporation within the organoids.

Analysis of significant (p.adj < 0.05) and relevantly different transcripts (abs log2FoldChange > 1)

revealed 4610 differentially expressed genes (DEGs) between DMDdEx8-9 and IsoCTR MYOrganoids,

of which 2181 upregulated and 2429 downregulated (**Figure S3A-B**). Of those, we identified genes

involved in decreased muscular contraction (e.g. muscle system process, muscle contraction,

sarcoplasmic reticulum calcium ion transport and actin-myosin filament sliding) as well as reduced

energetic molecular process (e.g. oxidative phosphorylation, mitochondrial respiratory chain complex

assembly, ATP synthesis coupled electron transport), as depicted by gene ontology biological process

(GOBP) enrichment analysis (**Figure 3E, Table S1**). Remarkably, between the upregulated pathways in

DMD MYOrganoids, we identified higher collagen fibril organization with an enrichment score of +0.6

(**Figure 3E**). This GO analysis was also confirmed by Kyoto encyclopedia of genes and genome

(KEGG) enrichment analysis, which showed a downregulation of oxidative phosphorylation and an

upregulation of ECM receptor interaction, such as Integrin alpha and Beta (**Figure S3C**). Interestingly,

the analysis of genes involved in the ECM remodeling revealed an upregulation of genes coding for

collagens, laminins, metalloproteases and TGF-β related genes such as TGF-B receptor (TGFBR1) and

the ncRNA inhibitor of TGF-β (TGFB2-AS1) in DMD MYOrganoids compared to isogenic control

(**Figure 3F**). Additionally, we found increased levels of latent TGF-β binding proteins (LTBP 2 and 4)

in dystrophic MYOrganoids, indicating that TGF-β is activated in the DMD context and contributes to

ECM remodeling (**Table S1**).

The increased profibrotic signature in DMD iPSC-derived MYOrganoids was then confirmed in all CTR

and DMD iPSC-derived MYOrganoids, by evaluation of the expression of two key fibrotic markers,

Fibronectin-1 (*FN1*) and Collagen-1 (*COL1A1*). Importantly, the differences in the expression of fibrotic

markers between Ctr and DMD MYOrganoids were significant only when including fibroblasts (**Figure**

3G). We further examined histologically the presence of activated TGF-β signaling in DMD organoids

by looking at phosphorylated SMAD3 (pSMAD3), the transcriptional effector of the canonical TGF-β

pathway. Notably, DMDdEx8-9-derived MYOrganoids showed increased pSMAD3 positive nuclei both

in myotubes (MyHC positive staining) and fibroblasts (MyHC negative cells) (**Figure 3H**). This

observation indicates a crosstalk between fibroblasts and myofibers and supports the potential of using

fibroblasts in tissue engineering as a source of ECM and pro-fibrotic cues under pathological conditions.

Figure 3. Fibroblast inclusion in DMD iPSC-MYOrganoids induces exacerbation of profibrotic signature and fatiguability. (A) *Overview of MYOrganoids generation from control iPSC (Ctr1 and Ctr2), DMD iPSC (DMDdEx45, DMDdEx8-43) and isogenic iPSC (IsoCTR and DMDdEx8-9) including control or dystrophic fibroblasts respectively for histological characterization, force and transcriptomic analysis.* ***(B)*** *Immunostaining of MYOrganoids cross sections for Dystrophin (Dys), Sarcomeric a-actinin (SAA), and DAPI for nuclei. Scale bar: 30 μm* ***(C)*** *Drop force over 10 repetitions of eccentric (ECC) contractions in all Ctr and all DMD MYOrganoids without (-) or with (+) fibroblasts*

N=4-6. (D) Fatigue index calculated as % of drop force between two isometric contractions (ISO) performed before and after 10 repetitions of ECC in all Ctr and all DMD MYOrganoids +/- respective fibroblasts. N=5-10. E) Gene Ontology enrichment analysis (GO TERM) from RNA-seq data in isogenic iPSC-derived CTR (IsoCTR) and DMD (DMDdEx8-9). The scale of color is representative of the p adjusted (p.adj) F) Clustered heatmap of fibrotic Differentially Expressed Genes (DEG) relative to DMD vs IsoCTR (left). Color scale represents the Z-score (red is higher, blue lower) G) Relative gene expression of Fibronectin (FN1) and Collagen (COL1A1) in all iPSC CTR and DMD-derived MYOrganoids with or without fibroblasts. H) Representative images showing activation of TGF-β signaling by immunostaining for pSMAD3+ cells in MyHC+ myotubes (arrows), and in fibroblasts (asterisk), considered as MyHC- cells, and relative quantification. Scale bar: 10 μm, n=4 G). Data are presented as mean ± SEM. Unpaired t-test was performed for statistical purposes (*p ≤0.05, **p ≤ 0.01, ***p ≤ 0.001, ****p ≤ 0.0001, ns= not significant).

Juxtracrine role of fibroblast in the modeling of DMD phenotype

We then sought to gain insights into the contribution of fibroblasts in the recapitulation of key

pathogenic hallmarks of DMD. Given that fibroblasts are known to primarily act through their secreted

molecules [59–61] we analyzed the secretome of both control (Ctr) and DMD MYOrganoids (MyoT +

Fibro) and compared it to that of single-cell 3D cultures (Fibro-only and MyoT-only) (**Figure 4A**).

Analysis of culture media at day 14 showed elevated levels of secreted TGF-β and collagen 4 (COL4)

in DMD MYOrganoids cocultured with fibroblasts compared to control conditions. No significant

differences were observed between control and DMD in Fibro-only and MyoT-only 3D cultures

(**Figure 4B**). These data indicate that DMD MYOrganoids in coculture with fibroblast exhibit an

increased release of fibrotic factors.

To determine whether fibroblasts amplify the fibrotic phenotype through their secretory activity, we collected conditioned media (CM) from control (Ctr) and DMD Fibro-only 3D cultures on day 7 of *in vitro* growth. We then applied this CM to treat Ctr and DMD MyoT-only tissues for additional 7 days, assessing the secretion of fibrotic factors, gene expression, and muscle force (**Figure 4C**). ELISA assay on secreted TGF-β and fibronectin 1 (FN1) revealed no difference in MyoT-only 3D cultures treated with the CM from either Ctr or DMD Fibro-only tissues (**Figure 4D**). Congruently, gene expression levels of TGF-β and Col1A1 were similar in Ctr and DMD MyoT-only cultures, regardless of the treatment with the CM (**Figure 4E**), while those were higher just in DMD cocultures (MyoT + Fibro) compared to control conditions (**Figure S4A**). The lack of effect of the CM indicates that the profibrotic

increase seen in DMD MYOrganoids in coculture with fibroblast occurs by a juxtracrine effect, as it comes from contacts between the two cell types.

We further evaluated the requirement of fibroblasts-myotubes coculture in disease exacerbation by assessing muscle force analysis following CM treatment, which confirmed no difference in fatiguability in myotube-only 3D cultures treated with control or DMD Fibro-derived CM, unlike higher fatigue index in DMD organoids containing both fibroblasts and myotubes in coculture (**Figure 4F**). Interestingly, MYOrganoids including fibroblasts displayed a higher tetanic force compared to myotubes-only cultures treated or not with fibroblasts CM, also supporting their cell contact-dependent role for enhanced functional maturation (**Figure S4B**). Collectively, analysis of secreted protein and force under CM experiments showed that the dystrophic fatiguability and fibrosis traits emerged only when fibroblasts and muscle cells were cocultured, indicating that contact-dependent cellular communication between muscle and fibroblasts is required to reveal the DMD traits of profibrotic secretion and fatiguability.

Figure 4. Fibroblast role in the modeling of DMD phenotype requires direct contact with DMD iPSC-derived muscle cells. *A) Scheme of engineered 3D cultures created using either iPS-derived myotubes (MyoT), fibroblasts (Fibro), or a combination of both (MyoT + Fibro=MYOrganoids) from CTR or DMD cells.* *B) TGF-β and COL1A1 ELISA performed in Fibro only, MyoT only or combination (Fibro+MyoT).* *C) Scheme of conditioned medium (CM) exchange.* *From day 7, (CM) was collected daily from 3D cultures with fibroblasts (Fibro) alone and used daily in MyoT 3D cultures compared to untreated MyoT only or MyoT+Fibro co-cultured MYOrganoids. D) TGF-β and Fibronectin 1 (FN1) levels detected by ELISA in CTR or DMD only MyoT only organoids after 7 days of treatment with CM from either wild-type fibroblasts (CTR-CM) or DMD fibroblasts (DMD-CM) and analyzed at day 14. E) TGF-β and COL1A1 gene expression in in CTR or DMD only MyoT only organoids after 7 days of treatment with CM from either wild-type fibroblasts (CTR-CM) or DMD fibroblasts (DMD-CM) and analyzed at day 14. F) Fatigue index in MyoT + Fibro organoids, and in MyoT only tissues treated or not with CTR-CM and DMD-CM. Ctr conditions are indicated in blue and DMD organoids in orange. For all panels, data are represented as average ± SEM. *p < 0.05; ***p<0.0001; ns = not significant (unpaired Student's t test).*

AAV-microdystrophin gene transfer rescues muscle resistance while partially restoring Dystrophin Glycoprotein Complex components in DMD MYOrganoids

As proof of concept that MYOrganoids were suitable as a screening platform for gene therapy, we used AAV-mediated delivery of microdystrophin (µDys) and assessed its therapeutic efficacy in the DMD context. We used AAV9 capsid for this aim, as this viral serotype was used in recent clinical trials and showed a high transduction rate in patients' myofibers [62,63]. We infected the organoids with a codon-optimized µDys gene (dR4-23, same protein product used in clinical trials [64]) under the control of the muscle-specific spc512 promoter [17] for gene transfer using AAV9 (AAV9-µDys) in all DMD MYOrganoids iPSC (dEx45, dEx8-43 and dEx8-9 with isogenic control, IsoCTR) and assessed gene transfer efficiency, muscle force analysis and membrane stability assessment (**Figure 5A**). We first optimized the infection conditions using the reporter AAV9-CMV-GFP in Ctr MYOrganoids including fibroblasts. Infection was performed on day 7 of the differentiation protocol, diluting AAV particles directly in the medium, and maintained for an additional 7 days (**Figures S5A-C**). Optimal low and high AAV9-µDys doses were established based on previous dosing studies to have intermediate transduction levels at low doses (1E+09 vg/MYOrganoid), and high transduction levels at high doses (5E+10 vg/MYOrganoid) (**Figure S5D**). Gene transfer efficiency was evaluated by quantification of viral copy number (VCN) on genomic DNA and by the expression level of the transgene and the encoded protein, showing a clear dose-dependent entry and expression of µDys in DMD MYOrganoids (**Figure 5B, Figures S6A-B**). As expected by using a muscle-specific promoter, only the muscle cells (MyoT) but not the fibroblasts expressed the transgene after infection of AAV9-uDys in the single populations (**Figure S6C**). We then tested whether µDys affected the contractility and fatigue resistance of the infected organoids. Isometric tetanic force analysis did not reveal any significant changes in DMD MYOrganoids following µDys delivery compared to not-infected conditions (**Figure S6D**), as expected by first-time contracting muscle tissues. We then challenged the muscles with repeated eccentric repetitions, and we observed a dose-dependent attenuation of the drop force in DMD MYOrganoids after

μDys infection (**Figure 5C**). Consistently, the fatigue index was greatly reduced in a dose-dependent

manner in all DMD MYOrganoids treated with a high dose of μDys, although reaching significance only

for DMDdEx45 and DMDdEx8-9 MYOrganoids (**Figure 5D**). To confirm that the observed effects were

specifically due to μDys expression rather than AAV infection alone, we infected DMD MYOrganoids

with a control vector containing a dystrophin fragment lacking a promoter (hereafter called empty

vector), which infection did not result in protein expression (**Figure S6E-G**). Particularly, the

MYOrganoids infected with the empty vector did not show any differences in the fatigue index analysis,

at both low and high dose, while we observed a decreased fatigue index in DMD MYOrganoids infected

with an AAV containing μDys (**Figure S6H**). Overall, the muscle force data confirmed that ectopic

μDys expression rescues muscle endurance and fatiguability.

Because dystrophin exerts its biomechanical support by holding the Dystrophin Glycoprotein Complex

(DGC) at the membrane, we monitored key components of this complex, such as the transmembrane β-

Dystroglycan (β-DG), directly binding Dystrophin, and the extra-cellular α-Dystroglycan (α-DG), whose

proper expression and localization are impaired in the absence of dystrophin [65]. To this aim, DMD

MYOrganoids infected with low and high dose of μDys, were subjected to histological analysis.

Quantification of μDys showed around 25-30% of dystrophin-positive myotubes at low dose compared

to 85% in the high dose condition in DMD dEx8-9 (**Figure 5E-F**) and dEx45 and 65% in the DMD

dEx8-43 MYOrganoids (**Figure S7A-B**). Immunostaining on transversal MYOrganoids showed a dose-

dependent yet not complete restoration of β-DG in all DMD iPSC-derived MYOrganoids. Interestingly,

even high doses ensuring nearly total Dystrophin transduction in all the DMD MYOrganoids did not

significantly restore α-DG (around 10% in DMDdEx8-9 and DMDdEx8-43 and around 30% in

DMDdEx45) suggesting a potential therapeutic limitation of μDys (**Figure 5E-F**). Importantly, these

data demonstrate the limited restoration of Dystrophin-associated components to the sarcolemma after

successful μDys gene transfer.

Figure 5. μDys gene transfer-mediated improvement of muscle resistance and partial rescue of DCG complex. (A) Scheme of AAV9-μDys infection and doses used in DMD iPSC-derived MYOrganoids. (B) Evaluation of gene transfer efficiency by viral copy number (VCN) analysis (N=3) and protein expression analysis by capillary western blot in isogenic iPSC DMD dEx8-9-derived MYOrganoids infected with low and high dose of μDys versus non-infected (-) IsoCTR (N=2-3). (C) Drop force over 10 repetitions of eccentric (ECC) contractions in isogenic cell lines and Ctr1, Ctr2 and DMDdEx45, DMDdEx8-43-derived MYOrganoids receiving low and high doses of μDys. N=6-7 (D) Fatigue index in all Ctr and DMD MYOrganoids treated or not with m-Dystrophin at low and high dose. N=5-6. (E) Histological evaluation of Dystrophin Glycoprotein Complex (DGC) upon infection, by immunostaining for Dystrophin (DYS), α-Dystroglycan (α-DG) and β- Dystroglycan (β-DG) in cross sections of Isogenic Ctr and DMD-iPSC-derived MYOrganoids. Scale bar: 40 µm and relative quantification in all control and iPSC-derived MYOrganoids (F). Data

are presented as means +/- SEM. Statistical analysis was performed with an ordinary one-way ANOVA test (*p ≤0.05, **p ≤ 0.01, ***p ≤0.001, ****p ≤ 0.0001, ns = not significant) with multiple comparisons corrected with Tukey's test.

Optimal microdystrophin gene transfer partially corrects the transcriptomic profile of DMD MYOrganoids and does not rescue the profibrotic signature

To examine the effect of microdystrophin (µDys) on the signaling pathways and molecular processes

characteristics of dystrophic pathology, we performed bulk RNA-seq analysis comparing the corrected

DMD dEx8-9 (IsoCTR) and DMDdEx8-9 MYOrganoids untreated and treated with a high dose of

AAV9-µDys (n=4).

Principal component analysis (PCA) of the top 10000 genes with the highest variance showed a distinct

transcriptomic profile of DMD organoids treated with µDys (dEx8-9 + µDys high) and untreated (dEx8-

9) compared to isogenic controls (IsoCTR) (**Figure S8A**). To evaluate the effect of µDys in restoring

key pathways involved in DMD, we firstly compared the transcriptomic profile of µDys-treated DMD

organoids to untreated DMD organoids (dEx8-9 µDys vs dEx8-9), and then to the isogenic control

(dEx8-9 µDys vs IsoCTR). Particularly, µDys-treated DMD organoids showed a total of 327 DEGs, of

which 154 upregulated and 173 downregulated, compared to untreated organoids (**Figure 6A, Table**

S2). Of those 327 DEGs, 52% (170/327) are in common with the DEGs that were found dysregulated in

DMD organoids when compared to the isogenic control (**Figure S8B**). Therefore, we performed the

gene ontology enrichment analysis of the common category to see the effect of µDys on the DMD-

dysregulated pathways. This analysis revealed increased expression of genes involved in the microtubule

organization (GOBP: Microtubule organizing center organization NES = +2.31), cytoskeleton

organization (GOBP: Protein localization to cytoskeleton NES=+1.99), and a decreased expression of

genes involved in the inflammation (GOBP: Regulation of inflammatory response, NES = 1.45) (**Figure**

6B).

We then analyzed the DEGs between the treated organoids and the isogenic control to check the

transcriptomic signature not restored by µDys (dEx8-9-µDys vs IsoCTR). Particularly, µDys-treated

organoids present 7818 DEGs (4004 upregulated and 3814 downregulated) when compared to the

corrected isogenic organoids (**Figure 6C**). Gene ontology analysis of the DEGs between µDys-treated

DMD organoids and IsoCTR organoids revealed the pathways that are not restored by the gene transfer.

Particularly, treated organoids showed persistent downregulation of genes involved in calcium transport

(GOBP: Sarcoplasmic reticulum calcium ion transport, NES = -2.15), oxidative phosphorylation

(GOBP: Oxidative phosphorylation, NES = -2.29) and upregulation of TGFβ signaling (Hallmark TGFβ

signaling, NES = +1.57), when compared to isogenic controls (**Figure 6D, Table S3**). Consistently, a

clustered heatmap of all 1716 most significant differentially expressed genes (DEGs) further highlighted

the transcriptomic differences between control and DMD organoids (**Figure 6E**). Particularly, analysis

of Z-score of fibrotic genes among the three conditions (IsoCTR, DMDdEx8-9 and DMDdEx8-9 treated

with µDys) revealed a persistent but variegated profibrotic signature. In particular, we revealed a slight

decrease in the expression of ADAMs (i.e. ADAM17, ADAMTS7, ADAMTS16) of collagen genes (e.g.

COL11A1, COL4A6) and of the TGF-β receptor 1 gene (TGFBR1) in µDys-treated DMD organoids,

when compared to the untreated condition, although still significantly different from isogenic control

(**Figure 6E, right panel).** Additionally, other fibrotic genes, like *FN1* and *COL1A*1, Integrins or

cytokines (e.g.IL1A), showed no expression changes compared to untreated control (**Figure 6E, right**

panel). Further validation of the expression of *COL1A1*, *FN1,* not decreased by the µDys treatment, and

of *TGF-β*, partially downregulated, confirmed the persistent upregulation of those genes in DMD-treated

MYOrganoids (**Figure 6F**).

These data demonstrate that gene transfer of µDys in DMD MYOrganoids normalizes only a small

portion of the dysregulated transcriptome and does not reduce the transcription of key fibrotic ECM and

secreted proteins.

Figure 6: Transcriptomic analysis in μDys-treated isogenic DMD MYOrganoids. (A,C) Volcano plot showing the differential gene expression in treated (dEx8-9-μDys) versus untreated (dEx8-9) DMD MYOrganoids(A) or DMD dEx8-9-μDys vs isogenic control (IsoCTR (C). Upregulated genes are shown in red, while downregulated genes are shown in blue. Numbers indicate the amount of downregulated and upregulated genes. *(B,D)* Gene ontology enrichment analysis of treated (dEx8-9-μDys) versus untreated (dEx8-9) DMD MYOrganoids (B) or dEx8-9-μDys vs IsoCtr (D). Normalized enrichment score (NES) and adjusted p-value (p.adj) are indicated for each subcategory. *(E)* Clustered Heat map and enlargement depicting relative transcript levels of differentially expressed genes in corrected DMD (IsoCTR) and DMD MYOrganoids transcriptomics treated or not with μDystrophin (dEx8-9 and dEx8-9 μDys high). Lower and higher expressions are depicted in blue and red, respectively. The data points above the significance threshold (p adjusted < 0.05, fold2change > 2) are marked in blue (downregulated) and red (upregulated), and others are marked in gray (not significant). *(F)* Relative gene expression of TGF-β, ColA1 and FN1 in Isogenic control, DMD dEx8-9 untreated or tretated with μDys. Data are presented as means +/- SEM. Statistical analysis was performed with an ordinary one-way ANOVA test (*p ≤0.05, **p ≤ 0.01, ***p ≤ 0.001, ****p ≤ 0.0001, ns = not significant).

Microdystrophin gene therapy does not reduce fibroblast profibrotic activity.

The persistence of profibrotic signature in µDys-treated MYOrganoids, identified by bulk RNA-seq, led us to gain further insights into the therapeutic effect exerted by µDys in the specific subpopulation of the MYOrganoid. Since µDys is not expressed in fibroblasts (**Figure S6C**), we wondered if the ameliorated myotubes can have a paracrine effect on fibroblasts secretory activity.

Thus, we evaluated the effect of the secretome of the DMD MYOrganoids treated with µDys on the fibroblasts-only 3D cultures or fibroblasts-including MYOrganoids (MyoT+Fibro) (**Figure 7A**). For that, we treated both DMD MyoT+ Fibro and only Fibro 3D cultures with the conditioned media (CM) of DMD MYOrganoids (MyoT + Fibro) treated with high doses of AAV9-µDys (**Figure 7A**). ELISA on the cell culture media of DMD MYOrganoids treated with µDys showed a decreased concentration of TGF-β comparable to the isogenic control (**Figure 7B**). Moreover, we observed a decreased secretion of TGF-β protein in DMD MYOrganoids treated with the CM of µDys-infected organoids, which remained yet significantly different from the fibroblast-including MYOrganoids (**Figure 7B**). Interestingly, DMD fibroblasts only 3D culture treated with the CM of µDys infected MYOrganoids did not show any differences in the secretion of TGF-β, showing any effect from the µDys modulated secretome on the fibroblast population (**Figure 7B**). Next, we assessed the expression of key fibrotic genes, such as *COL1A1* and *FN1* (downstream genes activated in fibrotic pathways that remain unaltered in µDys-infected organoids). In DMD MYOrganoids, *COL1A1* and *FN1* expression remain upregulated following µDys treatment, in accordance with RNA-seq data, and the same effect was also seen with CM from µDys-treated organoids (**Figure 7C**). Together these data suggested a lack of paracrine effect of the milieu of µDys treated- muscle cells in the secretion of ECM proteins and a partial effect for the upstream signaling protein TGF-β exclusively within fibro-including MYOrganoids.

To understand the mechanisms behind these results, we evaluated the reduction in TGF-β pathway activation within both cell subpopulations in DMD MYOrganoids following µDys treatment. Histological staining revealed a significant decrease in pSMAD3-positive nuclei within the skeletal

muscle subpopulation of µDys-treated organoids, whereas no difference was observed in fibroblasts after

µDys treatment (**Figure 7D**). This result suggests a cell-autonomous correction of this signaling within

the muscle population receiving µDys, but no paracrine effect on fibroblast activity. The persistence of

activated TGF-β signaling in the fibroblast subpopulation after µDys gene transfer (**Figures 3H and 7D**)

can therefore explain the uncorrected profibrotic signature and poses the basis for future investigation

aimed at improving or driving beneficial cell-cell communication.

Figure 7. Persistent profibrotic activity in fibroblasts contained in µDys-MYOrganoids A) Diagram of
conditioned media (CM) experiments following µDys gene transfer. Engineered tissues were created using either
iPS-derived myotubes in coculture with fibroblasts (MyoT + Fibro) or with fibroblasts alone (Fibro). MyoT + Fibro
MYOrganoids have been infected with AAV9-µDys at high dose (5E+10 vg/org) at day 7-8 and their CM have been
used as treatment for not-infected MyoT+Fibro and Fibro engineered tissues. B) TGF-β ELISA on Ctr MyoT +
Fibro and DMD + Fibro, treated or not with µDys or µDys-CM organoids media. In green, DMD Fibro only
engineered tissue TGF-β released in the media treated or not with the µDys-treated organoids CM. C) Collagen
1A1 (COL1A1) and Fibronectin 1 (FN1) gene expression in Ctr and DMD MYOrganoids, treated or not with either
AAV9-µDys and µDys-treated organoids CM. In green, DMD fibro-only tissues treated or not with µDys-CM. D)
Histological staining of phospho-SMAD3 (pSMAD3) staining in green, myosin heavy chain (MyHC) in red and
nuclei with DAPI in µDys-treated DMD MYOrganoids, and relative quantification of pSMAD3 positive nuclei in
*percentage. Scale bar = 10 µm. For all panels, data are represented as average ± SEM. *P < 0.05; ** p < 0,005;*
*** p < 0,001; ns = not significant (unpaired Student's t test).

Discussion

Here, we report the generation of 3D-engineered muscles called MYOrganoids, composed of a homogeneous population of skeletal muscle cells [11,36] in coculture with fibroblasts, a major source of connective tissue as well as key players for muscle tissue assembly [38–40]. We proved that fibroblast inclusion in the MYOrganoids [38–40] results in improved maturation and organization of the muscle tissue and high functional performance under muscle force evaluation. Importantly, fibroblast incorporation was pivotal in exacerbating muscle force loss and fatiguability of DMD MYOrganoids, enabling their use as therapeutic readouts. This is important considering that the capacity to evaluate muscle function in a high throughput manner remains limited, further hindering progress in testing or developing effective treatments. As such, while several remarkable studies reported on muscle force in 3D models [30,31,56,66,67], evaluating muscle function still presents challenges due to the complexity of identifying disease-specific force parameters and the variability of *in vitro* models. Moreover, the use of our direct differentiation method which leads to comparable myogenic potential of CTR and DMD MYOrganoids, enabled us to perform muscle force analysis without bias introduced by either intrinsic myogenic defects, as we start from equally functional myofibers, or by heterogenous cell populations that may affect the interpretation of results [31,52,68].

This study also sheds new light on the mechanism underlying the contribution of fibroblasts to the modeling of DMD. We showed that direct interaction of DMD fibroblast with muscle cells is essential to reproducing key pathogenic hallmarks, such as a profibrotic signature and increased muscle fatigue in DMD MYOrganoids. Interestingly, fibroblast role in the recapitulation of DMD hallmarks was not exerted by their profibrotic secretory activity per se and coculture with iPSC-derived muscle cells was required, as observed with the lack of effect of CM. In particular, we observed that, in a DMD environment, fibroblasts show higher secretion of fibrotic factors like TGF-β, fibronectin, and collagen through juxtracrine signaling. This result aligns with other studies demonstrating a higher profibrotic secretory activity from both DMD muscle and fibrotic cells [59,69,70]. It also supports the notion that cell

communications between the two cell populations, fibroblast and muscle fiber, is crucial to reveal

pathogenic traits, underscoring the relevance of fibroblast inclusion to help mimicking the fibrotic

microenvironment and aggravate muscle endurance. One possible hypothesis is that fibroblasts

secretory activity is enhanced by alterations in cell communications with dystrophin-deficient muscle

cells whose membrane stability is impaired, thereby causing increased matrix deposition and

susceptibility to contraction-induced damage and fatigue.

Overall, the exacerbated severity of DMD MYOrganoids enabled the evaluation of the therapeutic

potential of uDys, currently employed in clinical trials. Unexpectedly, μDys^{6970}[16,17] delivery in DMD

MYOrganoids did not fully rescue DMD phenotype as it led to incomplete restoration of the

components of the dystroglycan complex and no reduction in fibrotic genes transcriptome. Our findings

reveal a discrepancy to what was observed in animal models, where full rescue of muscle transcriptome

and membrane stability is achieved with uDys transfer [71,72]. This evidence supports the relevance of a

human platform to unravel the potential limitations of the current treatments and consider genetic

variations. Notably, we observed a reduction in the TGF-β pathway activation exclusively on muscle

cells, while no effect was observed in fibroblasts. This can explain the persistence of ECM protein

production and secretion in μDys treated DMD MYOrganoids, while we observed an effect in the

restoration of muscle contractility. Up to now, no other work revealed the impact of μDys on the

fibroblast population of the muscle, revealing the potential of our system in highlighting specific cell-

cell communications in response to treatment. Importantly, the persistence of dysregulated pathways

not corrected by gene transfer, could be related either to the lack of key functional domains or the

necessity to explore complementary strategies correcting additional parameters or to empower μDys

efficacy to achieve full functional recovery.

While our work reveals the importance of analyzing the cell-cell communication between fibroblasts

and muscle cells, further studies need to be performed to fully understand the role of fibroblasts in the

dystrophic process. For example, how the absence of dystrophin in the myotubes is affecting the

440 fibroblasts and how the reintroduction of µDys is affecting the composition of the ECM, still need to

441 be elucidated. Finally, pro-fibrotic DMD MYOrganoids might provide a valuable tool either for the

442 investigation of mechanisms driving dystrophic process and as a human *in vitro* counterpart to animal

443 *in vivo* preclinical studies, with the potential to accelerate the identifications of new treatments as well

444 as to unravel therapeutic limitations of current DMD treatments.

Acknowledgments

We express our gratitude to Rene Hummel (Danish MyoTechnology), Guillaume Tanniou, and Nicolas Guerchet for their invaluable technical support in muscle force evaluation. We thank Guillaume Corre for the aid with RNA-seq analysis. We thank the service of bioproduction of Genethon, for preparing the AAV particles used in this study. We also extend our appreciation to Jérémie Cosette for the assistance in imaging and to the histology team for their technical expertise. Finally, we would like to thank Frederique Magdinier for generously providing the DMD iPSC dEx45 and Ctr1 iPSC.

Fundings

This study was financially supported by the Institut National de la Sante et de la Recherche Medicale (INSERM) and by the "Association Française contre les Myopathies" (AFM).

Author contributions

SA and LPa designed the experiments. LPa performed most of the experiments, including iPSC culture, organoid generation, imaging, and muscle force analysis. LPi performed force analysis, gene expression analysis and western blot experiments. AJ analyzed conditioned media biomarkers and performed secretome analysis. LPa, MM contributed to muscle force analysis. AVH analyzed RNA-seq data. RE and GB performed electron microscopy analysis. AB generated and provided the immortalized human fibroblasts. SA and LPa performed data analysis and figure preparation. SA conceived, supervised the project, and wrote the manuscript. All authors discussed the results. SA, LPa, DI and IR reviewed and edited the manuscript.

Declaration of interests

All authors declare they have no competing interests.

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Sonia Albini (salbini@genethon.fr).

Material availability

Ctr2, IsoCTR, DMDdEx8-9 and DMDdEx8-43 hIPS cell lines used in this study are restricted due to MTAs (Ref: MTA 2022 GENETHON - AIM _DMD, MMTA202109-0107, 2021-0296_MTA_CECS). This study did not generate new unique reagents.

Data and code availability

The data supporting the findings of this study are available within the article and its supplemental information. The RNA sequencing datasets generated in this study can be found in the NCBI Bioproject database (https://www.ncbi.nlm.nih.gov/bioproject) using the access number PRJNA1044721.

Microscopy and functional data reported in this paper will be shared by the lead contact upon request.

Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODELS

Induced pluripotent stem cells (iPSCs)

Human iPSCs used in the study were as follows: Ctr1 and Ctr2 from healthy individuals, DMDdEx45, DMDdEx8-43 from DMD patients and the isogenic iPSC DMDdEx8-9 and corrected control (CRISPR-mediated skipping of mutated exons dEx6-9, isogenic control) (**Table 1**). The Ctr1 and DMDdEx45 hiPSC lines were generated from skin fibroblasts from Coriell (**Table 1**) and were obtained from Marseille Stem Cells platform in Marseille Medical Genetics laboratory [73].

iPSCs were maintained using mTeSRplus medium (Stem Cell Technologies) and passaged using ReleSR (STEMCELL Technologies) on Matrigel-coated wells (Corning). iPSC engineering and muscle differentiation were performed adapting a transgene-based method previously described[69,74]. Quality control was performed regularly with Oct-4 and Nanog expression profile and mycoplasma testing, and cells were used within 30 passages. Additionally, antibiotic and antimycotic solution (Merk-Millipore) was regularly added to each cell culture media. All iPSC lines are engineered for transgene-based myotubes differentiation. Briefly, cells were nucleofected (4D Nucleofector, Lonza) with two enhanced versions of piggyBAC (ePB) containing respectively MyoD and BAF60c genes under tetracycline responsive promoter [69,74]. Additionally, iPSCs were regularly selected for the presence of the two ePBs with puromycin (10 µg/ml) and blasticidin (20 µg/ml) at the same time (Sigma-Aldrich).

Human immortalized fibroblasts

Human immortalized fibroblasts from control (AB1191C16PV, AIM) and DMD patient (AB1024DMD11Q, AIM) were generated and obtained from Myobank-AFM of Myology Institute from muscle biopsies (**Table 1**). Control and DMD human immortalized fibroblasts, used in co-culture with hiPSC in 3D MYOrganoids, were maintained in culture in DMEM High Glucose Lglut supplemented with 20% fetal bovine serum (FBS) and passaged using TrypleE (Gibco).

Human primary fibroblasts

Human primary fibroblasts from control (3 lines: 5-8898, 11-1610, 11-1973) and from DMD patients (3 lines, 5-11618, 5-12608, 5-11613) were obtained from DNA bank of Genethon (**Table 1**). Fibroblasts were maintained in DMEM high glucose (Gibco) supplemented with 20 ng/ml recombinant human fibroblast growth factor (h-FGF) (Peprotech) and 20% fetal bovine serum (Gibco). Primary fibroblasts were passaged using TrypleE (Gibco)

Table 1: Cell line list

	Current name	**Reference**	**Healthy individuals and patients**
Human induced pluripotent stem cells (hiPSCs)	Ctr1	AG08C5 Coriell	Skin fibroblast, 1-year-old healthy male
	Ctr2	M180 CECS	Skeletal myoblasts, healthy male
	IsoCTR	*Kyrychenko et al 2017*	Peripheral blood mononuclear cells healthy male
	DMDdEx45	GM25313 Coriell	Skin fibroblast, 13-year-old DMD male patient, deletion ex45
	DMDdEx8-9	*Kyrychenko et al 2017*	Peripheral blood mononuclear cells healthy male
	DMDdEx8-43	M202	Skeletal myoblasts, DMD male patient, out-of-frame deletion exon 8-43
Immortalized skin fibroblasts (hTERT/CDK4)	Ctr fibroblasts	AB1191C16PV	Skin, 16-year-old healthy male
	DMD fibroblasts	AB1024DMD11Q	, Skin 11-year-old DMD male, stop-codon exon 59
Primary skin fibroblasts	Fibroblasts control 1	5-8898 DNA Bank Genethon	Skin fibroblasts healthy patient

	Fibroblasts control 2	11-1610 DNA Bank Genethon	Skin fibroblasts healthy patient
	Fibroblasts control 3	11-1973 DNA Bank Genethon	Skin fibroblasts healthy patient
	Fibroblasts DMD 1	5-11618 DNA Bank Genethon	Skin fibroblasts DMD patient
	Fibroblasts DMD 2	5-12608 DNA Bank Genethon	Skin fibroblasts DMD patient
	Fibroblasts DMD 3	5-11613 DNA Bank Genethon	Skin fibroblasts DMD patient

METHOD DETAILS

Generation of MYOrganoids

MYOrganoids from iPSCs were generated adapting the protocol described for engineered heart tissue[75] .First, iPSC were induced with doxycycline 200 ng/µl for 24 hours to induce the expression of MyoD and Baf60c. The day after, 1.25×10^6 iPSC-committed were resuspended in 77 µl of growth media (SKM02, AMSbiokit) supplemented with hES cell Recovery (Stemgent) and molded in hydrogel composed by 40 µl of Bovine Collagen solution 6mg/ml (Sigma-Aldrich), 17.8 µl of Matrigel Growth Factor reduced (GFR, Corning) 10% v/v 3), 40µl 2X DMEM (Gibco) 4) and 5.2 µl of NaOH 0.05 N). For the generation of MYOrganoids including fibroblasts, 1.25×10^5 (ratio 1:10) fibroblasts were included in the iPSC-committed mix during hydrogel preparation. 180µl of hydrogel was casted into 48-well plate TM5 MyrPlate (Myriamed), containing in each well a pair of flexible poles (static stretchers) that support the growth of the engineered tissue in a ring shape. After 1 hour of polymerization at 37°C, growth media (SKM02, AMSbiokit) was added for 24 hours. On day 2 of the 3D, growth media (SKM02, AMSbiokit) was replaced by differentiation media (SKM03plus, AMSbiokit) and changed every day until day 14.

For Condition Media (CM) experiments, single cell-population organoid-like structures were made using the same protocol described above. CM was collected from Ctr or DMD fibroblasts-only 3D structure and diluted 1:10 in fresh differentiation media (SKM03plus, AMSbiokit). The diluted media was then used to culture Ctr or DMD myotubes-only organoids. CM was collected and used to treat 3D cultures from day 7 until day 13.

Muscle force analysis

Functional analyses were carried out on day 14 after 3D casting. Contraction experiments were performed using the MyoDynamics Muscle Strip System 840 MD (Danish Myo Technology A/S) and CS4 stimulator (Danish Myo Technology A/S). All functional analysis were performed at 37°C, 5% CO_2 95% O_2, in Tyrode's solution supplemented with 25 mM $NaHCO_3$. Optimal muscle length was determined by gradually stretching the muscle until 1.0 mN of passive tension registered. Functional tests were performed under isometric and eccentric conditions. MYOrganoids were electrically stimulated with 250 pulses of 30V, 4 ms width at the 125 Hz of frequency for both isometric and eccentric contractions. For eccentric analysis, MYOrganoids were 1 mm stretched at the 6.5 mm/s speed during the muscular contraction. Data collection and analysis was done by PowerLab device and LabChart software (ADInstruments, New Zealand) respectively. Each artificial tissue was subjected to 1 isometric contraction, 10 eccentric contractions and 1 isometric contraction. Fatigue is represented as percentage drop force between the first and the last isometric contraction. Where indicated, force is indirectly normalized for the CSA (Cross Section Area) calculated as muscle force (mN) x Lo (mm) x density (mg/mm3)/weight (mg) and expressed as mN/mm2. Engineered muscle density is experimentally determined as 2.089 mg/mm3.

Immunofluorescence

MYOrganoids that did not undergo functional analysis have been analyzed for immunohistochemistry. Briefly, the artificial tissues are fixed in 4% methanol-free

paraformaldehyde (PFA) overnight on day 14. For whole mount staining, fixed MYOrganoids were permeabilized, stained and cleared with the MACS clearing kit (Miltenyi) according to manufacturer's instructions. Whole mount-stained organoids are then imaged with confocal microscope (LEICA STED SP8) at 10X magnification. Antibodies used for staining on whole mount MYOrganoid are listed in **Table 2**.

For staining on transversal or longitudinal sections, fixed MYOrganoids are dehydrated with a gradient of sucrose (7.5%-30%) over-day and embedded in OCT matrix in plastic mold. After 24 hours, embedded MYOrganoids are processed with the cryostat (LEICA) with 14 μm thick sections. Slices were then dried and fixed again with 4% methanol-free PFA (Invitrogen). Fixed sections are then blocked with serum cocktail (5% Goat serum and 5% Fetal bovine serum), before being stained overnight at +4°C with primary antibody. After that, slices are washed three times in PBS and hybridized with AlexaFluor secondary antibody accordingly to the host species of the first antibody. Stained slides were then covered with Fluoromont + Dapi (SouthernBiotech) and glass slide 1.5H. For imaging, sections are scanned with AxioScan microscope and confocal Leica SP8. Antibodies used for staining on MYOrganoid sections are listed in **Table 2**.

For 2D staining, cells are grown on μ-Dish 35 mm (Ibidi) and then fixed in 4% methanol-free PFA for 7 minutes. For membrane and cytoplasmatic stainings, cells are permeabilized with 0.15% Triton X-100 for 10 minutes and then washed with PBS for 5 minutes. For nuclear staining, cells are permeabilized with 0.25% Triton X-100 for 15 minutes. Permeabilized cells are then blocked with serum cocktail (5% Goat serum and 5% Fetal bovine serum), before being stained overnight at +4°C with primary antibody. Cells are then washed three times with PBS for 5 minutes and then hybridized with AlexaFluor secondary antibody according to the host species of the first antibody. After three washing of 5 minutes in PBS, nuclei are stained for 10 minutes with Hoechst 33342 (Invitrogen) at the final dilution of 2 μg/ml. For imaging, sections are scanned confocal Leica SP8. Antibodies used for 2D staining are listed in **Table 2**.

Table 2: Antibodies used for immunofluorescence and capillary western blot

Antibodies	Supplier	Reference
Mouse monoclonal Myosin Heavy Chain (used at 1:10)	DSHB	Cat#MF20, RRID:AB_2147781
Rabbit polyclonal Vimentin (used at 1:100)	Proteintech	Cat#10366-1-AP, RRID:AB_2273020
Mouse monoclonal Dystrophin (used at 1:20)	Leica Biosystem	Cat#NCL-DYSB, RRID:AB_563691
Mouse monoclonal Fibronectin (used at 1:200)	Sigma-Aldrich	Cat# F7387, RRID:AB_476988
Rabbit monoclonal pSMAD-3 (used at 1:100)	Abcam	Cat# ab52903, RRID:AB_882596

Mouse monoclonal α-Dystroglycan (used at 1:100)	Millipore	Cat# 05-298, RRID:AB_309674
Mouse monoclonal β-Dystroglycan (used at 1:100)	Leica Biosystems	Cat#NCL-b-DG, RRID:AB_442043
Rabbit monoclonal alpha-actinin 2 (used at 1:100)	Thermo Fisher Scientific	Cat# 701914, RRID:AB_2688290
Mouse monoclonal Oct3/4 (used at 1:200)	Santa Cruz Biotechnology	Cat# sc-5279, RRID: AB_628051
Mouse monoclonal Pax7 (used at 1:100)	DSHB	Cat# PAX7, RRID:AB_2299243
Goat anti-Mouse IgG1 Cross-Adsorbed Secondary Antibody, Alexa Fluor 594	Thermo Fisher Scientific	Cat#A-21125, RRID:AB_2535767
Goat anti-Rabbit IgG (Heavy chain), Superclonal™ Recombinant Secondary Antibody, Alexa Fluor 488	Thermo Fisher Scientific	Cat# A27034, RRID:AB_2536097
Goat anti-Mouse IgG2a Cross-Adsorbed Secondary Antibody, Alexa Fluor 594	Thermo Fisher Scientific	Cat#A-21135, RRID:AB_2535774
Goat anti-Mouse IgG2b Cross-Adsorbed Secondary Antibody, Alexa Fluor 647	Thermo Fisher Scientific	Cat#A-21242, RRID:AB_2535811

Electron Microscopy studies

Electron microscopy analysis was prospectively performed on MYOtissue specimens that were fixed with glutaraldehyde (2.5%, pH 7.4), post fixed with osmium tetroxide (2%), dehydrated in a graded series of ethanol ranging from 30% to absolute solution and embedded in resin (EMBed-812, Electron Microscopy Sciences, USA). 80 nm thick sections from at least four blocks from Ctr iPSC-derived MYOrganoids in presence or absence of Ctr fibroblasts were stained with uranyl acetate and lead citrate. The grids were observed using a "JEOL 1400 Flash" electron microscope (120 kV) and were photo documented using a Xarosa camera (Soft Imaging System, France). Images covering whole longitudinal sections were assessed and representative images were used.

Images analysis

Myotubes alignment

FIJI and CellPose were used for image analysis. Myotube alignment was determined by angle measurement and by myotube circularity from cross-section cuts of MYOrganoids. Angles between two myotubes were measured with the "angle tool" function in FIJI, by drawing two lines perpendicular to two adjacent myotubes membrane. Myotubes circularity was determined by custom FIJI script. Briefly, myotubes cross section area was first segmented by pre-trained Cellpose2 cyto2 model [76] and then converted into Regions of Interest (ROIs) by Labels_To_Rois.py plugin (Waisman et al. 2021) for subsequent quantification on FIJI (Ferret diameters X and Y). After that, the ratio between Feret's diameter on axis X and axis Y was assessed. At least 4 frames for 3 biological replicates have been analyzed.

Maturation and fusion index

Maturation index was assessed counting nuclei contained in striated myotubes identified by sarcomeric alpha actinin SAA staining in FIJI software and normalized for the total number of nuclei contained in myotubes. Fusion index has been calculated counting myotubes myosin heavy chain (MyHC) positive containing 1, 2, 3 or more than 3 nuclei as a percentage of the total number of nuclei. At least 4 frames for 3 biological replicates have been analyzed.

Z-disk length analysis

Z-disk length has been assessed by drawing straight lines in FIJI and by measuring the length in at least 10 mature myotubes per condition.

Phospho-SMAD3 quantification

Phospho-SMAD3 (pSMAD3) has been measured as a percentage of positive nuclei of myosin heavy chain positive cells (MyHC+, i.e. myotubes) or myosin heavy chain negative cells (MyHC-, i.e. fibroblasts). At least 4 frames for 3 biological replicates have been analyzed.

Dystrophin, α-dystroglycan and β-dystroglycan quantification

MYOtissue cross sections were stained sarcomeric alpha-actinin (SAA) for myotubes cytosol labeling. The Cellpose2 cyto2 model 10 was fine-tuned on manually myofiber-labeled images based on SAA staining (hyperparameters: n_epochs=200, learning_rate=0.05, weight_decay=0.0001). The labeled dataset used in fine-tuning was prepared in such a way that the model can simultaneously segment myotubes and ignore low-quality staining areas. Fine-tuned models were then used to extract myotubes masks. Reconstruction of myotubes masks was done using the cellpose package (Stringer et al. 2021). Reconstructed masks were then converted into Regions of Interest (ROIs) for subsequent quantification (each ROI corresponds to an individual myofiber) using the Labels_To_Rois.py FIJI plugin [77]. The generated ROIs were used for subsequent quantification using FIJI macro. Positive ROIs for dystrophin / alpha-dystroglycan / beta-dystroglycan signal were counted and represented as a percentage of the total ROIs of one image.

Gene expression analysis

For gene expression analysis, bulk RNA was isolated from MYOrganoids by RNeasy micro kit (QIAGEN) according to manufacturer's instructions, controlled and quantified by Nanodrop. Around 0.5-1µg of RNA was retro-transcribed to cDNA thanks to the RevertAid H Minus First Strand cDNA Synthesis Kit (Invitrogen). Droplet digital PCR was performed to assess the expression of myogenic factors (MYOGENIN, MYH2, MYH7, MCK), of fibrotic markers (COL1A1, FN1, TGF-β1) and µDystrophin, thanks to the QX200™ ddPCR™ EvaGreen Supermix (Biorad). Gene expression results in copy/µl (cp/µl) are represented as fold change to the expression of GAPDH housekeeping gene. Primers used are listed in **Table 3**.

Table 3: List of primers used for viral copy number analysis and gene expression

Primers for gene expression analysis		
Gene	**Forward**	**Reverse**
GAPDH	ACAACTTTGGTATCGTGGAAGG	GCCATCACGCCACAGTTTC
MYOGENIN	AATGCAGCTCTCACAGCGCCTC	TCAGCCGTGAGCAGATGATCC
MCK	TGGAGAAGCTCTCTGTGGAAGCT	TCCGTCATGCTCTTCAGAGGGTA
MYH2	AGAAACTTCGCATGGACCTAGA	CCAAGTGCCTGTTCATCTTCA
MUH7	ACTGCCGAGACCGAGTATG	GCGATCCTTGAGGTTGTAGAGC
µDYS	GCACCACCAGATGCACTAT	GTGTAGGCGTAGCTCTTGAAT
COL1A1	AAGAGGAAGGCCAAGTCGAG	GTTTCCACACGTCTCGGTCA
FN1	AGCCGAGGTTTTAACTGCGA	CCCACTCGGTAAGTGTTCCC
TGF-β1	GCCTGAGGCCGACTACTA	CTGTGTGTACTCTGCTTGAACT
Primers for VCN analysis		
Gene	**Forward**	**Reverse**
P0	CTCCAAGCAGATGCAGCAGA	ATAGCCTTGCGCATCATGGT
µDYS	GCACCACCAGATGCACTAT	GTGTAGGCGTAGCTCTTGAAT

ELISA assay

Media was collected before muscle functional analysis after being in culture for 24 hours. Secreted human Fibronectin 1 (FN1), TGF-β and Collagen IV were assessed by ELISA kit (**Table 4**) from MYOrganoids supernatant.

Table 4: ELISA kit used for secreted protein analysis

Target protein	Supplier	Reference
Fibronectin 1	Abcam	ab219046
TGF-β1	Abcam	ab100647
Collagen IV	Bio-techne	NBP2-75864

AAV production and MYOrganoids infection

Recombinant AAVs were produced as previously described (Bourg et al. 2022) using AAV9 serotype. Purification was performed using affinity chromatography and titration was done by ddPCR using transgene-specific primers. For optimization of infection, an AAV9-CMV-GFP construct was used. The micro-dystrophin transgene used in the study, under the control of spc512 promoter, was an optimized version of the construct used for GENETHON's preclinical investigation and clinical trial [71], with deletion from spectrin-like repeats 4 to 23 and full C-terminal truncation, here referred as μDys. Infection in MYOrganoids was performed by delivering the AAV9 particles diluted into the differentiation media at day 7, at two different doses: 1E+9 vg/MYOtissue (low dose) and 5E+10 vg/MYOtissue (high dose). Media was replaced after 24 hours from the infection and changed daily until day 14. For conditioned media (CM) experiment, media was collected from μDys-infected DMD MYOrganoids from day 7 until day 13 and used to culture DMD MYOrganoids (myotubes + fibro) or DMD fibro-only 3D cultures.

Viral copy number analysis

Viral DNA was extracted from mature MYOrganoids by NucleoMag Pathogen kit (Macherey Nagel) using Kingfisher instrument (Thermofisher). DNA yield and purity was assessed by Nanodrop; viral copy number (VCN) was identified by droplet digital PCR using supermix for probe (Biorad). Results are shown as copy number variation using P0 as reference DNA. Primers used are listed in **Table 3**.

Capillary western blot analysis

MYOrganoids proteins were extracted in RIPA buffer supplemented with Protease Inhibitor Cocktail EDTA-free (Roche) and Benzonase by homogenization. Total proteins were then quantified by BCA method, thanks to the Pierce 660 protein assay kit (Invitrogen) according to manufacturer's instructions. Protein detection has been performed by capillary western blot, thanks to the JESS protein simple (Bio-techne), according to manufacturer's directions. Dystrophin

685 detection (both full-length and μDystrophin) has been performed by the antibody DysB (**Table 2**)
686 and its expression has been quantified by total protein normalization.

RNA sequencing and transcriptomic analysis

The RNA quality of samples was verified using the Bioanalyzer 2100 (Agilent) and Qubit
fluorometric quantification (ThermoFisher Scientific). The samples that had an RNA integrity
number higher than 9 were used for RNA sequencing (Genewiz). The Stranded Total RNA Library
Prep Kit (Illumina) was used to create sequencing libraries, which were sequenced following the
Illumina protocol on the NovaSeq instrument (Illumina), resulting in approximately 20 million
paired-end reads per library. The paired-end reads were filtered and subjected to quality control
using fastp (Chen et al. 2018). They were then mapped to the GRCh38/hg38 genome using
HISAT2 [78] count tables were generated using htseq-count [79]. Differentially expressed genes (DEGs)
were identified using the DESeq2 R package with p value adjusted by Benjamin-Hochberg
procedure less than 0.05. Pathway analysis was carried out in R-Studio (version 4.0.3) using
functional class scoring with Gene Set Enrichment Analysis [80,81].

Statistical analysis

All data were analyzed by GraphPad Prism 9.5.1 software. Parametric tests such as t-tests and
ANOVA were used for statistical comparison. To compare two groups, unpaired one tail t-test was
used; to compare multiple groups, we used one-way ANOVA with Tukey's correction for multiple
comparison tests. Results were considered significantly different at $p < 0.05$. Graphs were
generated using Graphpad Prism v9 or R version 3.6.2, The figures display the mean ± standard
error of the mean (SEM) as described in the figure legend.

References

There are no sources in the current document.

1. Mendell, J. R. *et al.* Evidence-based path to newborn screening for Duchenne muscular dystrophy. *Ann Neurol* **71**, 304–313 (2012).
2. Hoffman, E. P., Brown, R. H. & Kunkel, L. M. Dystrophin: the protein product of the Duchenne muscular dystrophy locus. *Cell* **51**, 919–928 (1987).
3. Ervasti, J. M. & Campbell, K. P. A role for the dystrophin-glycoprotein complex as a transmembrane linker between laminin and actin. *J Cell Biol* **122**, 809–823 (1993).
4. Petrof, B. J., Shrager, J. B., Stedman, H. H., Kelly, A. M. & Sweeney, H. L. Dystrophin protects the sarcolemma from stresses developed during muscle contraction. *Proc Natl Acad Sci U S A* **90**, 3710–3714 (1993).
5. Hoffman, E. P., Brown, R. H. & Kunkel, L. M. Dystrophin: the protein product of the Duchenne muscular dystrophy locus. *Cell* **51**, 919–928 (1987).
6. Hoffman, E. P. The discovery of dystrophin, the protein product of the Duchenne muscular dystrophy gene. *FEBS Journal* **287**, 3879–3887 (2020).
7. Davies, K. E. & Nowak, K. J. Molecular mechanisms of muscular dystrophies: old and new players. *Nat Rev Mol Cell Biol* **7**, 762–773 (2006).
8. Ennen, J. P., Verma, M. & Asakura, A. Vascular-targeted therapies for Duchenne muscular dystrophy. *Skelet Muscle* **3**, (2013).
9. Porter, J. D. *et al.* A chronic inflammatory response dominates the skeletal muscle molecular signature in dystrophin-deficient mdx mice. *Hum Mol Genet* **11**, 263–272 (2002).
10. Amor, F. *et al.* Cholesterol metabolism is a potential therapeutic target in Duchenne muscular dystrophy. *J Cachexia Sarcopenia Muscle* **12**, 677–693 (2021).
11. Caputo, L. *et al.* Acute conversion of patient-derived Duchenne muscular dystrophy iPSC into myotubes reveals constitutive and inducible over-activation of TGFβ-dependent pro-fibrotic signaling. *Skelet Muscle* **10**, (2020).
12. Wynn, T. A. Cellular and molecular mechanisms of fibrosis. *Journal of Pathology* **214**, 199–210 (2008).
13. Kharraz, Y., Guerra, J., Pessina, P., Serrano, A. L. & Muñoz-Cánoves, P. Understanding the process of fibrosis in Duchenne muscular dystrophy. *Biomed Res Int* **2014**, 965631 (2014).
14. Mendell, J. R. *et al.* duan. *Mol Ther* **29**, 464–488 (2021).
15. Aguti, S., Malerba, A. & Zhou, H. The progress of AAV-mediated gene therapy in neuromuscular disorders. *Expert Opin Biol Ther* **18**, 681–693 (2018).
16. Duan, D. Systemic AAV Micro-dystrophin Gene Therapy for Duchenne Muscular Dystrophy. *Molecular Therapy* **26**, 2337–2356 (2018).
17. Le Guiner, C. *et al.* Long-term microdystrophin gene therapy is effective in a canine model of Duchenne muscular dystrophy. *Nat Commun* **8**, (2017).
18. Elangkovan, N. & Dickson, G. Gene Therapy for Duchenne Muscular Dystrophy. *J Neuromuscul Dis* **8**, S303–S316 (2021).
19. Harper, S. Q. *et al.* Modular flexibility of dystrophin: implications for gene therapy of Duchenne muscular dystrophy. *Nat Med* **8**, 253–261 (2002).
20. Wang, B., Li, J. & Xiao, X. Adeno-associated virus vector carrying human minidystrophin genes effectively ameliorates muscular dystrophy in mdx mouse model. *Proc Natl Acad Sci U S A* **97**, 13714–13719 (2000).
21. Fabb, S. A., Wells, D. J., Serpente, P. & Dickson, G. Adeno-associated virus vector gene transfer and sarcolemmal expression of a 144 kDa micro-dystrophin effectively restores the dystrophin-associated protein complex and inhibits myofibre degeneration in nude/mdx mice. *Hum Mol Genet* **11**, 733–741 (2002).
22. Duan, D. Duchenne Muscular Dystrophy Gene Therapy in 2023: Status, Perspective, and Beyond. *Hum Gene Ther* **34**, 345–349 (2023).
23. Elangkovan, N. & Dickson, G. Gene Therapy for Duchenne Muscular Dystrophy. *J Neuromuscul Dis* **8**, S303–S316 (2021).
24. Takahashi, K. *et al.* Induction of Pluripotent Stem Cells from Adult Human Fibroblasts by Defined Factors. *Cell* **131**, 861–872 (2007).
25. Tornabene, P. *et al.* Intein-mediated protein trans-splicing expands adeno-associated virus transfer capacity in the retina. *Sci Transl Med* **11**, (2019).
26. Clevers, H. Modeling Development and Disease with Organoids. *Cell* **165**, 1586–1597 (2016).
27. Hofer, M. & Lutolf, M. P. Engineering organoids. *Nat Rev Mater* **6**, 402–420 (2021).
28. Shin, M. K. *et al.* Generation of Skeletal Muscle Organoids from Human Pluripotent Stem Cells to Model Myogenesis and Muscle Regeneration. *Int J Mol Sci* **23**, (2022).

29.   Pinton, L. *et al.* 3D human induced pluripotent stem cell-derived bioengineered skeletal muscles for tissue,
disease and therapy modeling. *Nat Protoc* **18**, (2023).
30.   Rao, L., Qian, Y., Khodabukus, A., Ribar, T. & Bursac, N. Engineering human pluripotent stem cells into
a functional skeletal muscle tissue. *Nat Commun* **9**, (2018).
31.   Shahriyari, M. *et al.* Engineered skeletal muscle recapitulates human muscle development, regeneration
and dystrophy. *J Cachexia Sarcopenia Muscle* **13**, 3106–3121 (2022).
32.   Garita-Hernandez, M. *et al.* AAV-Mediated Gene Delivery to 3D Retinal Organoids Derived from Human
Induced Pluripotent Stem Cells. *Int J Mol Sci* **21**, (2020).
33.   Kruczek, K. *et al.* Gene Therapy of Dominant CRX-Leber Congenital Amaurosis using Patient Stem Cell-
Derived Retinal Organoids. *Stem Cell Reports* **16**, 252–263 (2021).
34.   Khabou, H. *et al.* Noninvasive gene delivery to foveal cones for vision restoration. *JCI Insight* **3**, (2018).
35.   Caputo, L. *et al.* Acute conversion of patient-derived Duchenne muscular dystrophy iPSC into myotubes
reveals constitutive and inducible over-activation of TGFβ-dependent pro-fibrotic signaling. *Skelet Muscle*
10, (2020).
36.   Albini, S. *et al.* Epigenetic Reprogramming of Human Embryonic Stem Cells into Skeletal Muscle Cells
and Generation of Contractile Myospheres. *Cell Rep* **3**, 661–670 (2013).
37.   Albini, S. & Puri, P. L. Generation of Myospheres From hESCs by Epigenetic Reprogramming. *Journal of
Visualized Experiments* (2014) doi:10.3791/51243.
38.   Mathew, S. J. *et al.* Connective tissue fibroblasts and Tcf4 regulate myogenesis. *Development* **138**, 371–
384 (2011).
39.   Bersini, S. *et al.* Engineering an Environment for the Study of Fibrosis: A 3D Human Muscle Model with
Endothelium Specificity and Endomysium. *Cell Rep* **25**, 3858-3868.e4 (2018).
40.   Li, M., Dickinson, C. E., Finkelstein, E. B., Neville, C. M. & Sundback, C. A. The role of fibroblasts in
self-assembled skeletal muscle. *Tissue Eng Part A* **17**, 2641–2650 (2011).
41.   Santos, G. L. *et al.* Fibroblast Derived Human Engineered Connective Tissue for Screening Applications.
J Vis Exp **2021**, (2021).
42.   Kühl, U. *et al.* Role of muscle fibroblasts in the deposition of type-IV collagen in the basal lamina of
myotubes. *Differentiation* **28**, 164–172 (1984).
43.   Pinton, L. *et al.* 3D human induced pluripotent stem cell-derived bioengineered skeletal muscles for tissue,
disease and therapy modeling. *Nat Protoc* (2023) doi:10.1038/S41596-022-00790-8.
44.   Tiburcy, M., Meyer, T., Soong, P. L. & Zimmermann, W. H. Collagen-based engineered heart muscle.
Methods Mol Biol **1181**, 167–176 (2014).
45.   Rao, N. *et al.* Fibroblasts influence muscle progenitor differentiation and alignment in contact independent
and dependent manners in organized co-culture devices. *Biomed Microdevices* **15**, 161 (2013).
46.   Saleh, K. K. *et al.* Single cell sequencing maps skeletal muscle cellular diversity as disease severity
increases in dystrophic mouse models. *iScience* **25**, (2022).
47.   Giordani, L. *et al.* High-Dimensional Single-Cell Cartography Reveals Novel Skeletal Muscle-Resident
Cell Populations. *Mol Cell* **74**, 609-621.e6 (2019).
48.   Petrany, M. J. *et al.* Single-nucleus RNA-seq identifies transcriptional heterogeneity in multinucleated
skeletal myofibers. *Nat Commun* **11**, (2020).
49.   Karlsen, A. *et al.* Distinct myofibre domains of the human myotendinous junction revealed by single-
nucleus RNA sequencing. *J Cell Sci* **136**, (2023).
50.   Thorsteinsdottir, S., Deries, M., Cachaço, A. S. & Bajanca, F. The extracellular matrix dimension of
skeletal muscle development. *Dev Biol* **354**, 191–207 (2011).
51.   Costantini, M. *et al.* Biofabricating murine and human myo-substitutes for rapid volumetric muscle loss
restoration. *EMBO Mol Med* **13**, (2021).
52.   Al Tanoury, Z. *et al.* Prednisolone rescues Duchenne muscular dystrophy phenotypes in human pluripotent
stem cell-derived skeletal muscle in vitro. *Proc Natl Acad Sci U S A* **118**, (2021).
53.   Minato, K. *et al.* Measurement of Lateral Transmission of Force in the Extensor Digitorum Longus
Muscle of Young and Old Mice. *Int J Mol Sci* **22**, (2021).
54.   Erdogan, B. R. *et al.* Normalization of organ bath contraction data for tissue specimen size: does one
approach fit all? *Naunyn Schmiedebergs Arch Pharmacol* **393**, 243–251 (2020).
55.   Mann, C. J. *et al.* Aberrant repair and fibrosis development in skeletal muscle. *Skelet Muscle* **1**, 21 (2011).
56.   Kyrychenko, V. *et al.* Functional correction of dystrophin actin binding domain mutations by genome
editing. *JCI Insight* **2**, (2017).
57.   Childers, M. K. *et al.* Eccentric contraction injury in dystrophic canine muscle. *Arch Phys Med Rehabil*
83, 1572–1578 (2002).
58.   Elangkovan, N. & Dickson, G. Gene Therapy for Duchenne Muscular Dystrophy. *J Neuromuscul Dis* **8**,
S303 (2021).

59. Zanotti, S., Gibertini, S. & Mora, M. Altered production of extra-cellular matrix components by muscle-
derived Duchenne muscular dystrophy fibroblasts before and after TGF-beta1 treatment. *Cell Tissue Res*
339, 397–410 (2010).
60. Weber, K. T., Sun, Y., Bhattacharya, S. K., Ahokas, R. A. & Gerling, I. C. Myofibroblast-mediated
mechanisms of pathological remodelling of the heart. *Nat Rev Cardiol* **10**, 15–26 (2013).
61. Giuliani, G., Rosina, M. & Reggio, A. Signaling pathways regulating the fate of fibro/adipogenic
progenitors (FAPs) in skeletal muscle regeneration and disease. *FEBS J* **289**, 6484–6517 (2022).
62. Manini, A., Abati, E., Nuredini, A., Corti, S. & Comi, G. Pietro. Adeno-Associated Virus (AAV)-
Mediated Gene Therapy for Duchenne Muscular Dystrophy: The Issue of Transgene Persistence. *Front*
Neurol **12**, 814174 (2021).
63. Aguti, S., Malerba, A. & Zhou, H. The progress of AAV-mediated gene therapy in neuromuscular
disorders. *Expert Opin Biol Ther* **18**, 681–693 (2018).
64. Elangkovan, N. & Dickson, G. Gene Therapy for Duchenne Muscular Dystrophy. *J Neuromuscul Dis* **8**,
S303 (2021).
65. Lapidos, K. A., Kakkar, R. & McNally, E. M. The dystrophin glycoprotein complex: signaling strength
and integrity for the sarcolemma. *Circ Res* **94**, 1023–1031 (2004).
66. Madden, L., Juhas, M., Kraus, W. E., Truskey, G. A. & Bursac, N. Bioengineered human myobundles
mimic clinical responses of skeletal muscle to drugs. *Elife* **4**, (2015).
67. Khodabukus, A. *et al.* Electrical stimulation increases hypertrophy and metabolic flux in tissue-engineered
human skeletal muscle. *Biomaterials* (2018) doi:10.1016/j.biomaterials.2018.08.058.
68. Chal, J. *et al.* Differentiation of pluripotent stem cells to muscle fiber to model Duchenne muscular
dystrophy. *Nat Biotechnol* **33**, 962–9 (2015).
69. Caputo, L. *et al.* Acute conversion of patient-derived Duchenne muscular dystrophy iPSC into myotubes
reveals constitutive and inducible over-activation of TGFβ-dependent pro-fibrotic signaling. *Skelet Muscle*
10, (2020).
70. Granados, A. *et al.* SETDB1 modulates the TGFβ response in Duchenne muscular dystrophy myotubes.
Sci Adv **10**, (2024).
71. Le Guiner, C. *et al.* Long-term microdystrophin gene therapy is effective in a canine model of Duchenne
muscular dystrophy. *Nat Commun* **8**, (2017).
72. Duan, D. Systemic AAV Micro-dystrophin Gene Therapy for Duchenne Muscular Dystrophy. *Mol Ther*
26, 2337–2356 (2018).
73. Badja, C. *et al.* Efficient and Cost-Effective Generation of Mature Neurons From Human Induced
Pluripotent Stem Cells. *Stem Cells Transl Med* **3**, 1467–1472 (2014).
74. Albini, S. *et al.* Epigenetic Reprogramming of Human Embryonic Stem Cells into Skeletal Muscle Cells
and Generation of Contractile Myospheres. *Cell Rep* **3**, 661–670 (2013).
75. Tiburcy, M., Meyer, T., Liaw, N. Y. & Zimmermann, W. H. Generation of Engineered Human
Myocardium in a Multi-well Format. *STAR Protoc* **1**, (2020).
76. Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist algorithm for cellular
segmentation. *Nat Methods* **18**, 100–106 (2021).
77. Waisman, A., Norris, A. M., Elías Costa, M. & Kopinke, D. Automatic and unbiased segmentation and
quantification of myofibers in skeletal muscle. *Sci Rep* **11**, (2021).
78. Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory requirements.
Nat Methods **12**, 357–360 (2015).
79. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome Biol* **15**, (2014).
80. Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for interpreting
genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545–15550 (2005).
81. Mootha, V. K. *et al.* PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately
downregulated in human diabetes. *Nat Genet* **34**, 267–273 (2003).
872

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

A

B

Secreted TGF-β

C

COL1A1

FN1

D

DMDdEx8-9 + µDys high

Myotubes (MyHC+)   *Fibroblasts (MyHC -)*

10 µm

Fibroblasts (MyhC-)

Myotubes (MyhC+)