Facial expression is a distinctive behavioural marker of pain processing in the brain

Marie-Eve Picard[1,2], Miriam Kunz[3], Jen-I Chen[1,2], Michel-Pierre Coll[4], Étienne Vachon-Presseau[5,6,7], Tor D. Wager[8], Pierre Rainville[2,9]

[1] Department of psychology, Université de Montréal, Montreal, Quebec, Canada

[2] Centre de recherche de l'institut universitaire de gériatrie de Montréal, Montreal, Quebec, Canada

[3] Department of medical psychology and sociology, Medical faculty, University of Augsburg, Augsburg, Germany

[4] School of psychology, Université Laval, Quebec, Quebec, Canada

[5] Faculty of Dentistry, McGill University, Montreal, Quebec, Canada

[6] Department of Anesthesia, McGill University, Montreal, Quebec, Canada

[7] Alan Edwards Centre for Research on Pain, McGill University, Montreal, Quebec, Canada

[8] Department of psychological and brain sciences, Dartmouth College, Hanover, New Hampshire, United States

[9] Stomatology department, Faculté de médecine dentaire, Université de Montréal, Montreal, Quebec, Canada

*Corresponding author: Marie-Eve Picard

E-mail: marie-eve.picard.2@umontreal.ca

Abstract

Pain is a private experience observable through various verbal and non-verbal behavioural manifestations. Despite the importance of understanding the cerebral mechanisms underlying those manifestations, there is currently limited knowledge on the neural correlates of facial expression of pain. Here, we applied a brain decoding approach to functional magnetic resonance imaging (fMRI) data to predict the facial expression of pain during noxious heat stimulation in healthy volunteers. Results revealed the inability of previously developed pain neurosignatures to predict the facial expression of pain. We thus propose a Facial Expression of Pain Signature (FEPS) conveying distinctive information about the brain response to nociceptive stimulations with minimal overlap with other pain-relevant brain signatures. The FEPS provides a better characterization of the distributed cerebral representations of non-verbal pain communication. This underscores the complexity of pain phenomenology by reinforcing the view that neurosignatures conceived as biomarkers must be interpreted in relation to the specific pain manifestation predicted.

Introduction

Theories of pain communication highlight the diversity in pain manifestations, which occur through multiple channels: verbal reports, vocal complaints, change of postures, and facial expressions. From an evolutionary perspective, several manifestations of pain appear to be preserved across vertebrate phyla and reflect various functional roles to preserve the integrity of the organism (Sneddon 2019). Withdrawal behaviour allows the individual to move away from the noxious source, while facial expressions in social species convey information about the presence of a potential threat and an appeal for assistance (Hadjistavropoulos et al. 2011). Functionally distinct manifestations imply at least partly segregated neurophysiological processing (Sliwa et al. 2022). Previous functional magnetic resonance imaging (fMRI) studies investigating the neural correlates of acute pain have suggested that spontaneous or induced fluctuations in pain facial expression partly reflect changes in activity within the cortical targets of the spino-thalamo-cortical pathways. These fluctuations are, however, independent from changes in stimulus intensity and are inversely related to activity in prefrontal regions (Kunz et al. 2011; Vachon-Presseau et al. 2016; Kunz, Chen, and Rainville 2020). This suggests that pain facial expression may reflect the integration of activity across distributed brain networks processing ascending nociceptive signals, determining action policy, and gating efferent facial motor outputs.

The interest in developing pain neuromarkers has led researchers to use multivariate pattern analysis to investigate the distributed brain mechanisms underlying the experience of pain evoked by acute nociceptive stimuli. However, functional magnetic resonance imaging (fMRI) studies have revealed not one but several brain signatures of acute experimental pain that may reflect the diversity and complexity of pain-related function. The Neurological Pain Signature (NPS; Wager et al. 2013) was developed to predict changes in pain induced by variations in stimulus intensity and captured by subjective reports, reflecting primarily the

cerebral contributions to acute nociceptive pain (Krishnan et al. 2016; Van Oudenhove et al. 2020; Wager et al. 2013). To account for spontaneous fluctuations in the perception of pain intensity, the Stimulus-Independent Intensity of Pain Signature (SIIPS-1) was trained on noxious thermal trials after removing the effects of the stimulus intensity and the NPS response (Woo et al. 2017). More recently, the affective dimension of pain has received more attention, resulting in a multivariate pattern predictive of negative affect ratings to thermal pain, referred here as the Thermal Pain Aversive Signature (TPAS; Čeko et al. 2022). Finally, a signature was elaborated to characterise the neuronal representations associated with the valuation of pain (PVP) in the context of a decision task involving a cost-benefit analysis of future pain against a monetary reward (Coll et al. 2022). Taken together, those signatures have contributed to improve our understanding of the neurobiological mechanisms of pain, as reflected in self-report or explicit decision making.

Facial expression has been used as a reliable behavioural measure of pain across different mammal species (Costa et al. 2014; Craig 1992; Evangelista et al. 2019; Langford et al. 2010; Sotocina et al. 2011), but few studies have investigated the brain mechanisms associated with the spontaneous non-verbal communication of pain in human (Kunz et al. 2011; Kunz, Chen, and Rainville 2020). As an automatic behavioral manifestation, pain facial expression might be an indicator of activity in nociceptive systems, perceptual and evaluative processes, or general negative affect. We assessed the association between pain facial expression and the available pain-relevant brain signatures and we applied multivariate analysis with machine learning techniques to develop a predictive brain activation model of the facial responses to pain.

Results and Discussion

The Facial Action Coding System (FACS; Ekman and Friesen 1978) was used to quantify the facial expression of pain in healthy participants while brain responses evoked by brief moderately painful heat stimulation were recorded using fMRI. For each trial, the intensity and the frequency of pain-related action units were scored and combined into a FACS composite score (Methods). The association with the NPS, the SIIPS-1, the PVP, and the TPAS was assessed across the whole brain using the correlation between the FACS scores and the dot product computed between each signature and the activation maps for each individual trial.

Pain facial expression was not significantly associated with NPS expression (*pearson-r* = .01; *p* = .86), TPAS expression (*pearson-r* = -.03; *p* = .55), or PVP expression (*pearson-r* = .08; *p* = .07). Low but significant positive correlation was found with the SIIPS-1 expression (*pearson-r* = .10; *p* = .03). These low values indicate that the available pain-relevant brain signatures show poor sensitivity to the facial expression of pain. This motivated the development of a new multivariate brain pattern to predict pain expression.

We used a multivariate approach at the voxel level across the whole brain to develop the FEPS (Methods). A LASSO principal component regression was applied to predict the FACS composite scores from the trial-by-trial fMRI contrast images. The FEPS was able to predict the FACS composite scores with a performance significantly above chance level (averaged cross-validated prediction across 10 folds: *pearson-r* = .49 ± .11; R^2 = .17 ± .08; RMSE = 6.54 ± 0.99; *p* < .001 compared to a permuted distribution; Figure 1A-B). These results indicate that we were able to develop a multivariate brain pattern accounting for the facial responses related to pain.

Figure 1: FEPS: a brain signature of the facial expression of pain. A) Relationship between the actual and the predicted FACS composite scores for each cross-validation fold. B) Distribution of the Pearson's r scores across the cross-validation folds. C) Predictive weight map of pain expression thresholded at *p-uncorrected* < .05 using bootstrap tests performed with 5000 samples (with replacement). The thresholded map is shown for visualization and interpretation purposes only, although the prediction was made using voxel weights across the whole brain. MNI coordinates of the clusters with the related z-score can be found in Table S1. The colour bar represents the z-scored regression weights reflecting the positive and negative association with the magnitude of the FACS composite score of pain expression.

The distributed pattern of activity predicting pain expression was projected back on the brain to examine the spatial distribution of higher weights contributing to the prediction (Figure 1C). Positive weight clusters were found in the primary motor cortex (M1; bilateral), the frontal pole, the right posterior parietal cortex, the aMCC and the dorsal part of the parietal operculum, adjacent to the secondary somatosensory cortex (S2). These regions are

respectively associated with motor control, reward and affective value, attentional processes, and nociceptive pain processing (Table S1)(Price 2000; Rushworth et al. 2011; Shackman et al. 2011). Notably, separate analysis indicated that M1 activity was not sufficient to achieve significant prediction of facial responses (Figure 1S). Regions showing negative weights included the dorso-lateral PFC (dlPFC), the ventro-lateral PFC (vlPFC), the mid cingulate cortex (MCC), the subgenual ACC, the ventral part of the parietal operculum, the precuneus, and the vmPFC (Table S1). Negative weights imply that increased activity in those regions is associated with decreased facial response, consistent with a role in the inhibition of pain expression (Kunz et al. 2011). These results are consistent with the distributed nature of brain activity associated with pain facial expression reflecting in part the ascending nociceptive response and the ensuing affective processes, as well as top-down socio-affective regulation underlying the implementation of learned display rules (Karmann et al. 2016).

Several regions identified in the FEPS have also been reported in other pain-related brain signatures. Regions predictive of pain facial expression and pain intensity (NPS and SIIPS-1) include S2, the vmPFC, and the precuneus. The vlPFC is a region that does not receive direct spino-thalamo-cortical nociceptive afferents, and was reported both in the FEPS and in the SIIPS-1. Overlap between the FEPS and the PVP (Pain Value Pattern) include regions associated with reward and affect (i.e. OFC). Finally, the primary motor cortex, and S2 were also reported as contributing regions in the TPAS. The weak positive association between both the SIIPS-1 and the facial expression in addition with the spatial comparison showing some common regions across these pain-relevant signatures suggest possible shared features with the FEPS.

Figure 2: Spatial similarity between the FEPS and other pain-related signatures. A) Pattern similarity between the FEPS and other pain-related brain signatures using the weights of the full brain patterns. Pattern similarities were computed at the voxel level using the cosine similarity; a value of 1 reflects proportional patterns; a value of 0 reflects orthogonal patterns; a value of -1 reflects patterns of opposite direction. The left panel shows the similarity matrix, and the right panel shows only the significant similarities between the pain-related signatures (* $p < .05$; ** $p < .01$; *** $p < .001$). B) Deconstructing the pattern similarity with regards to seven cortical networks as defined in the Yeo atlas[24]: Visual Network (VN); Somatomotor Network (SMN); Dorsal Attention Network (DAN); Ventral Attention Network (VAN); Limbic Network (LN); Frontoparietal Network (FPN); Default Mode Network (DMN). Null distributions computed using permutation tests are shown, and the actual similarity values are represented by the vertical bar. Significant similarity values were found in the FPN (similarity = .24; $p = .001$), and the DMN (similarity = .18; $p = .005$) for the SIIPS-1, in the LN (similarity = .23; $p = .003$), and DMN (similarity = .11; $p = .05$) for the PVP, and in the LN (similarity = .11; $p = .04$) for the TPAS.

We computed the cosine similarity between the FEPS and other pain-related brain signatures to further examine the shared and specific representations between those predictive patterns (Methods). Cosine similarity ranging from -.03 to .09 was found between the FEPS and the other pain-related brain signatures reflecting the overall low similarity between the

signatures at the whole-brain level (Figure 2A). Highest similarity values with the FEPS were found for the SIIPS-1 and the PVP, consistent with the notion that the facial expression of pain may reflect at least partly changes in brain responses associated with spontaneous fluctuations in pain experience captured by pain ratings and with the assessment of pain value. Similarity with the FEPS was further assessed across different cortical networks. The significant positive similarity with the SIIPS-1 at the frontoparietal level may suggest common mechanisms in prediction and emotional regulation of the pain experience that would be reflected in both facial expression and subjective reports (Pan et al. 2018). Recruitment of the frontoparietal network may also be involved in the conscious representation of the pain context, making nociceptive information available for integration into decision-making processes (Coll et al. 2022; Bastuji et al. 2016; Cul, Baillet, and Dehaene 2007; Zheng et al. 2020). Finally, the convergent similarities in the limbic network with the PVP, and the TPAS, is consistent with a key role of affective pain processing influencing facial expression, perceived pain value, and reported unpleasantness or aversiveness of pain (Čeko et al. 2022; Garcia-Larrea and Peyron 2013; Roy et al. 2009).

In research and clinical practice, verbal reports of perceived pain intensity are considered to be the gold standard for measuring pain. Other measures that are often weakly correlated with those subjective reports, like facial expressions of pain, are often considered a less valid metric of the experience of pain even though they provide important complementary information on pain-related processes (Hadjistavropoulos et al. 2011). The Facial Expression of Pain Signature (FEPS) was able to predict the magnitude of the facial expression of pain. Regions that contribute to the prediction include motor and pain-related areas associated with both sensory and affective processing of pain. Although it shares, to some extent, similar representations with other pain-relevant signatures within various cerebral networks, the FEPS is distinctive from these other signatures. Results of this study

provide unique evidence of the complementary information provided by facial expression on pain-related brain processes. Future studies must provide a more comprehensive account of diverse pain manifestations and their related function to better capture the pain phenomenon in its entirety.

Methods

Participants

Secondary analyses of brain imaging data acquired in 34 participants were performed in this study (18 women, 16 men; age mean±SD = 23.4±2.5 years)(Kunz et al. 2011). No participants reported having a history of neurological or psychiatric disease, nor chronic pain. Participants reported not using alcohol nor analgesic for at least 24h before the experimental session. All participants provided written informed consent and received monetary compensation for their participation. The information about the video recording of the face was indicated in the consent form but this was not emphasized and it was not mentioned at the time of data acquisition. All procedures were approved by the ethics committee of the Centre de recherche de l'institut universitaire de gériatrie de Montréal.

Study design

Pre-experimental session

Participants were submitted to a pre-experimental session to assess the range of thermal pain sensitivity using a magnitude estimation procedure. All participants included in this study had a normal range of thermal pain sensitivity. The degree of facial expressiveness was also evaluated.

Thermal stimuli

Thermal stimulations were induced using a Medoc TSA-2001 thermosensory stimulator with a MRI-compatible 3 x 3 cm^2 contact probe (Medoc), positioned at the level of the lower left leg. Thermal stimuli lasted 9 seconds (2s ramp up, 5s plateau at targeted temperature, 2s *ramp down*) and were followed by an interstimulus interval of 18-25 seconds. The experiment was programmed using E-prime software (Psychology Software Tools, Pittsburgh, PA). A baseline temperature of 38°C was applied between stimuli for all participants. The target temperatures were determined individually before the MRI scans to

induce a warm non-painful sensation in the control condition, and a moderate to strong self-reported pain intensity (targeting 75-80/100 on the pain scale; temperature (mean±SD) = 47.8±0.90 °C). Participants were not aware that the warm and painful temperatures remained constant across trials. The order of the control condition and experimental condition was pseudorandomized. There were 8 trials for each experimental condition per run, for a total of 16 trials per condition and a total of 544 pain trials across all participants (34 participants x 16 pain trials). After each stimulus, participants rated the warm or pain sensation by moving a cursor with an MRI-compatible response key on a computerized visual analog scale. Because we were interested in the facial expression of pain, only the painful trials were considered in the analyses.

Facial expression

The facial expression of the participants was recorded using an MRI-compatible camera (MRC Systems) mounted onto the head coil of the MRI scanner. To be able to quantify facial expressions that occurred during the stimulation, a signal was sent from the stimulator to the sound card to automatically mark the onset of each stimulus on the video recording. Two certified FACS-coders evaluated the video recordings to rate the frequency, and intensity (on a 5-point scale) of pain-related action units (AUs; AU4, AU6-7, AU9-10, and AU43) for each trial (see Kunz et al., 2011 for details about the AUs selection). From these scores, a composite score (FACS composite score) was computed by taking the product between the mean AU frequency and mean AU intensity values, reflecting pain expression for each painful trial. The FACS composite scores were then used as the predictive variable. To examine whether the facial expression could be a confounding with the pain reports, we predicted the trial-by-trial FACS composited scores from the pain ratings using a mixed effect model with the participants as a random effect. Pain ratings were only weakly associated to facial responses (R^2_{GLMM} = .01; β = 0.80±0.25; $t(518.07)$ = 3.21; $p < .001$).

Anatomical and functional acquisition

MRI images were acquired using a 12-channel head coil 3T Siemens TRIO scanner. T1-weighted structural data were collected using a MP-RAGE sequence (TR = 2300ms, TE = 2.91ms, flip angle = 9°, FOV = 256mm, matrix size = 240×256, voxel size = 1×1×1.2mm, 170 whole-brain volumes). Functional data were acquired using an EPI T2*-weighted sequence (TR = 3000ms, TE = 30ms, flip angle = 90°, FOV = 220×220mm^2, voxel size = 3.44×3.44×3.40mm, 40 interleaved axial slices).

Preprocessing of fMRI data

The fMRI data were preprocessed using SPM8 (Statistical Parametric Mapping, Version 8; Wellcome Department of Imaging Neuroscience, London, United Kingdom) on MATLAB 7.4 (The MathWorks Inc, Natick, Massachusetts, United States). Preprocessing steps included a slice-timing correction, a correction for head movements, and co-registration between functional and anatomical images for each participant. Functional images were normalized into the MNI space. A spatial smoothing procedure (6mm FWHM gaussian kernel) and a high pass filter (128s) were also applied.

BOLD signal was modelled using a canonical hemodynamic response function. First level analyses were computed using the GLM to obtain a pain activation map for each trial using SPM8. Additionally, the six movements parameters and averaged signals from the white matter and the cerebrospinal fluid were included as nuisance regressors. Eleven trials were discarded due to excessive movements and a total of 533 activation maps of individual painful trials were used to develop the Facial Expression of Pain Signature (FEPS).

Analyses

Association between the facial expression of pain and pain-related brain signatures

The dot product between the Neurologic Pain Signature (NPS; Wager et al. 2013), the Stimulus Intensity Independent Pain Signature-1 (SIIPS-1; Woo et al. 2017), the Predictive

Value of Pain (PVP; Coll et al. 2022), the Thermal Pain Aversive Signature (TPAS; Čeko et al. 2022), and the trial-by-trial activation maps was computed to derive a measure of similarity (pattern expression) between the signatures and the maps. These scalar values were then correlated with the FACS composite scores to assess the association between the facial expression of pain and the NPS, the SIIPS-1, the PVP, and the TPAS. Pearson-r correlation coefficients, and p-values are reported.

Multivariate pattern analysis

We applied a Least Absolute Shrinkage and Selection Operator Principal Component Regression (LASSO-PCR) with a 10-fold cross-validation procedure for brain decoding (Wager et al. 2011) using scikit-learn implementation (Pedregosa et al. 2011). The analyses were performed using the trial-by-trial contrast images as input to the model and the participants as a grouping factor (i.e. data from a given participant could only be either in the training set or the testing set for a given cross-validation fold). This procedure was used to predict FACS composite scores from contrast images. The performance of each regression model was evaluated using Pearson's correlation coefficient (*pearson-r*), coefficient of determination (R^2), and root mean square error (RMSE). The scikit-learn's implementation of the coefficient of determination was used, allowing negative values if the model showed worse performance compared to an unfitted model (i.e. a horizontal line). The averaged performance metrics across folds are reported for each analysis. To test if the models performed significantly above chance, permutation tests were computed using 5000 iterations, leading to a p-value corresponding to the probability that the particular metric would be obtained by chance. A bootstrap resampling procedure was also performed to evaluate the stability of the voxel contribution to the model performance. This procedure consists of randomly sampling 5000 times the dataset with replacement. The resulting samples contain the same number of observations as the dataset. The LASSO-PCR procedure

as described above is then applied on each sample. Z-scores and p-values are calculated on the overall regression coefficients.

This analysis procedure was first applied at the whole brain level. It was repeated using a spatial mask including only the precentral region bilaterally. This mask was derived from the Oxford-Harvard Cortical Atlas (Caviness et al. 1996). This secondary analysis was conducted to verify if the pattern of activity within the primary motor cortex (M1) might be sufficient to predict facial expression. This model was not able to predict the FACS composite scores of pain expression and is not discussed further (Figure S1A-B). To ensure that the whole brain model prediction of the facial responses was not confounded with the pain ratings, we predicted the facial composite scores from the FEPS pattern expression scores, and the trial-by-trial pain ratings using a mixed effect model including the participants as a random effect. Variance in the facial composite scores was significantly explained by the FEPS pattern expression scores (β = 5.18±0.25; t(237.2) = 20.62; p < .001), but not by the pain ratings (β =0.24±0.20; t(484.9) = 1.21; p = .23).

Spatial similarity across the FEPS and pain-related brain signatures

Similarity between the FEPS and other pain-related brain signatures was assessed using the cosine similarity computed across all voxels ($Similarity(X, Y) = \frac{X \cdot Y}{|X||Y|}$). This metric was computed on the NPS, the SIIPS-1, the PVP, and the TPAS. To further explore the cortical similarities between the FEPS and other pain-related brain signatures, we also computed the cosine similarity across different cortical networks (i.e. visual, somatomotor, dorsal attention, ventral attention, limbic, frontoparietal, and default mode)(Yeo et al. 2011). Permutation tests (n = 1000) using generative null models preserving the spatial autocorrelation of a target brain map were used to assess the significance of the similarity between the brain signatures (Burt et al. 2020). P-values were calculated from the generated

null distributions as the fraction of permuted samples where the absolute cosine similarity was equal or larger than the cosine similarity obtained on the original signatures.

Figure S1: Predictive performance of the M1-based model. A) Relationship between the actual and the predicted FACS composite scores for each cross-validation fold (k=10) using only the activity from the primary motor cortex as defined by the Oxford-Harvard Cortical Atlas (Caviness et al. 1996). B) Distribution of the Pearson-r scores across the 10 cross-validation folds (*Pearson-r* = .11 ± .08; R^2 = -.29 ± .33; RMSE = 8.02 ± 0.96; *p* = .58).

Table S1. Peak regions contributing to the prediction of the facial expression scores.

Regions	x	y	z	Clusters size (mm³)	Peaks z-score
Regions with positive weights					
Frontal pole (bilateral)	12	70	-2	728	3.94
aMCC	6	18	40	200	2.31
Precentral gyrus (M1; right)	44	-14	38	1552	4.78
Precentral gyrus (M1; left)	-42	-18	40	520	3.70
Parietal operculum (right)	48	-26	34	32	3.23
Posterior cingulate gyrus (right)	20	-38	46	128	3.48
Inferior parietal lobule (right)	56	-28	42	56	3.20
Superior parietal lobule (right)	26	-56	68	8	3.12
Occipitotemporal/ Inferior temporal gyrus (left)	-48	-80	8	80	3.49
Regions with negative weights					
vlPFC (right)	38	54	-2	152	-3.40
vlPFC (left)	-40	48	-4	96	-3.48
dlPFC (right)	40	46	20	280	-3.62
vmPFC/subgenual ACC (bilateral)	-4	34	-10	1096	-3.60
Middle frontal gyrus (premotor area; left)	-34	0	56	8	-3.07
MCC (right)	4	4	42	120	-3.52
Post central (S1)/inferior parietal lobule (right)	48	-38	50	32	-3.13
Precuneus (left)	-14	-68	42	112	-3.44
Precuneus (right)	12	-66	54	16	-2.96
Fusiform gyrus (right)	26	-60	-14	56	-3.39

The clusters are shown in Figure 1.

References

Bastuji, Hélène, Maud Frot, Caroline Perchet, Michel Magnin, and Luis Garcia-Larrea. 2016. "Pain Networks from the inside: Spatiotemporal Analysis of Brain Responses Leading from Nociception to Conscious Perception." *Human Brain Mapping* 37 (12): 4301–15. https://doi.org/10.1002/hbm.23310.

Burt, Joshua B., Markus Helmer, Maxwell Shinn, Alan Anticevic, and John D. Murray. 2020. "Generative Modeling of Brain Maps with Spatial Autocorrelation." *NeuroImage* 220 (October): 117038. https://doi.org/10.1016/j.neuroimage.2020.117038.

Caviness, V. S., J. Meyer, N. Makris, and D. N. Kennedy. 1996. "MRI-Based Topographic Parcellation of Human Neocortex: An Anatomically Specified Method with Estimate of Reliability." *Journal of Cognitive Neuroscience* 8 (6): 566–87. https://doi.org/10.1162/jocn.1996.8.6.566.

Čeko, Marta, Philip A. Kragel, Choong-Wan Woo, Marina López-Solà, and Tor D. Wager. 2022. "Common and Stimulus-Type-Specific Brain Representations of Negative Affect." *Nature Neuroscience* 25 (6): 760–70. https://doi.org/10.1038/s41593-022-01082-w.

Coll, Michel-Pierre, Hocine Slimani, Choong-Wan Woo, Tor D. Wager, Pierre Rainville, Étienne Vachon-Presseau, and Mathieu Roy. 2022. "The Neural Signature of the Decision Value of Future Pain." *Proceedings of the National Academy of Sciences* 119 (23): e2119931119. https://doi.org/10.1073/pnas.2119931119.

Costa, Emanuela Dalla, Michela Minero, Dirk Lebelt, Diana Stucke, Elisabetta Canali, and Matthew C. Leach. 2014. "Development of the Horse Grimace Scale (HGS) as a Pain Assessment Tool in Horses Undergoing Routine Castration." *PLOS ONE* 9 (3): e92281. https://doi.org/10.1371/journal.pone.0092281.

Craig, Kenneth D. 1992. "The Facial Expression of Pain Better than a Thousand Words?" *APS Journal* 1 (3): 153–62. https://doi.org/10.1016/1058-9139(92)90001-S.

Cul, Antoine Del, Sylvain Baillet, and Stanislas Dehaene. 2007. "Brain Dynamics Underlying the Nonlinear Threshold for Access to Consciousness." *PLOS Biology* 5 (10): e260. https://doi.org/10.1371/journal.pbio.0050260.

Ekman, Paul, and Wallace V. Friesen. 1978. *Facial Action Coding System*. Consulting Psychologists Press.

Evangelista, Marina C., Ryota Watanabe, Vivian S. Y. Leung, Beatriz P. Monteiro, Elizabeth O'Toole, Daniel S. J. Pang, and Paulo V. Steagall. 2019. "Facial Expressions of Pain in Cats: The Development and Validation of a Feline Grimace Scale." *Scientific*

Reports 9 (1): 19128. https://doi.org/10.1038/s41598-019-55693-8.

Garcia-Larrea, Luis, and Roland Peyron. 2013. "Pain Matrices and Neuropathic Pain Matrices: A Review." *PAIN*, Biennial Review of Pain, 154 (December): S29–43. https://doi.org/10.1016/j.pain.2013.09.001.

Hadjistavropoulos, Thomas, Kenneth D. Craig, Steve Duck, Annmarie Cano, Liesbet Goubert, Philip L. Jackson, Jeffrey S. Mogil, et al. 2011. "A Biopsychosocial Formulation of Pain Communication." *Psychological Bulletin* 137 (6): 910–39. https://doi.org/10.1037/a0023876.

Karmann, Anna Julia, Christian Maihöfner, Stefan Lautenbacher, Wolfgang Sperling, Johannes Kornhuber, and Miriam Kunz. 2016. "The Role of Prefrontal Inhibition in Regulating Facial Expressions of Pain: A Repetitive Transcranial Magnetic Stimulation Study." *The Journal of Pain* 17 (3): 383–91. https://doi.org/10.1016/j.jpain.2015.12.002.

Krishnan, Anjali, Choong-Wan Woo, Luke J Chang, Luka Ruzic, Xiaosi Gu, Marina López-Solà, Philip L Jackson, Jesús Pujol, Jin Fan, and Tor D Wager. 2016. "Somatic and Vicarious Pain Are Represented by Dissociable Multivariate Brain Patterns." Edited by Peggy Mason. *ELife* 5 (June): e15166. https://doi.org/10.7554/eLife.15166.

Kunz, Miriam, Jen-I. Chen, Stefan Lautenbacher, Etienne Vachon-Presseau, and Pierre Rainville. 2011. "Cerebral Regulation of Facial Expressions of Pain." *Journal of Neuroscience* 31 (24): 8730–38. https://doi.org/10.1523/JNEUROSCI.0217-11.2011.

Kunz, Miriam, Jen-I. Chen, and Pierre Rainville. 2020. "Keeping an Eye on Pain Expression in Primary Somatosensory Cortex." *NeuroImage* 217 (August): 116885. https://doi.org/10.1016/j.neuroimage.2020.116885.

Langford, Dale J., Andrea L. Bailey, Mona Lisa Chanda, Sarah E. Clarke, Tanya E. Drummond, Stephanie Echols, Sarah Glick, et al. 2010. "Coding of Facial Expressions of Pain in the Laboratory Mouse." *Nature Methods* 7 (6): 447–49. https://doi.org/10.1038/nmeth.1455.

Pan, Junhao, Liying Zhan, ChuanLin Hu, Junkai Yang, Cong Wang, Li Gu, Shengqi Zhong, et al. 2018. "Emotion Regulation and Complex Brain Networks: Association Between Expressive Suppression and Efficiency in the Fronto-Parietal Network and Default-Mode Network." *Frontiers in Human Neuroscience* 12. https://www.frontiersin.org/articles/10.3389/fnhum.2018.00070.

Pedregosa, Fabian, Gaël Varoquaux, Alexandre Gramfort, Vincent Michel, Bertrand Thirion, Olivier Grisel, Mathieu Blondel, et al. 2011. "Scikit-Learn: Machine Learning in

Python." *Journal of Machine Learning Research* 12 (85): 2825–30.

Price, Donald D. 2000. "Psychological and Neural Mechanisms of the Affective Dimension of Pain." *Science* 288 (5472): 1769–72. https://doi.org/10.1126/science.288.5472.1769.

Roy, Mathieu, Mathieu Piché, Jen-I. Chen, Isabelle Peretz, and Pierre Rainville. 2009. "Cerebral and Spinal Modulation of Pain by Emotions." *Proceedings of the National Academy of Sciences* 106 (49): 20900–905. https://doi.org/10.1073/pnas.0904706106.

Rushworth, Matthew F. S., MaryAnn P. Noonan, Erie D. Boorman, Mark E. Walton, and Timothy E. Behrens. 2011. "Frontal Cortex and Reward-Guided Learning and Decision-Making." *Neuron* 70 (6): 1054–69. https://doi.org/10.1016/j.neuron.2011.05.014.

Shackman, Alexander J., Tim V. Salomons, Heleen A. Slagter, Andrew S. Fox, Jameel J. Winter, and Richard J. Davidson. 2011. "The Integration of Negative Affect, Pain and Cognitive Control in the Cingulate Cortex." *Nature Reviews Neuroscience* 12 (3): 154–67. https://doi.org/10.1038/nrn2994.

Sliwa, Julia, Marion Mallet, Maëlle Christiaens, and Daniel Y. Takahashi. 2022. "Neural Basis of Multi-Sensory Communication in Primates." *Ethology Ecology & Evolution* 34 (3): 322–43. https://doi.org/10.1080/03949370.2021.2024266.

Sneddon, Lynne U. 2019. "Evolution of Nociception and Pain: Evidence from Fish Models." *Philosophical Transactions of the Royal Society B: Biological Sciences* 374 (1785): 20190290. https://doi.org/10.1098/rstb.2019.0290.

Sotocina, Susana G, Robert E Sorge, Austin Zaloum, Alexander H Tuttle, Loren J Martin, Jeffrey S Wieskopf, Josiane CS Mapplebeck, et al. 2011. "The Rat Grimace Scale: A Partially Automated Method for Quantifying Pain in the Laboratory Rat via Facial Expressions." *Molecular Pain* 7 (January): 1744-8069-7–55. https://doi.org/10.1186/1744-8069-7-55.

Vachon-Presseau, Etienne, Mathieu Roy, Choong-Wan Woo, Miriam Kunz, Marc-Olivier Martel, Michael J. Sullivan, Philip L. Jackson, Tor D. Wager, and Pierre Rainville. 2016. "Multiple Faces of Pain: Effects of Chronic Pain on the Brain Regulation of Facial Expression." *Pain* 157 (8): 1819–30. https://doi.org/10.1097/j.pain.0000000000000587.

Van Oudenhove, Lukas, Philip A. Kragel, Patrick Dupont, Huynh Giao Ly, Els Pazmany, Paul Enzlin, Amandine Rubio, et al. 2020. "Common and Distinct Neural Representations of Aversive Somatic and Visceral Stimulation in Healthy Individuals." *Nature*

Communications 11 (1): 5939. https://doi.org/10.1038/s41467-020-19688-8.

Wager, Tor D., Lauren Y. Atlas, Lauren A. Leotti, and James K. Rilling. 2011. "Predicting Individual Differences in Placebo Analgesia: Contributions of Brain Activity during Anticipation and Pain Experience." *Journal of Neuroscience* 31 (2): 439–52. https://doi.org/10.1523/JNEUROSCI.3420-10.2011.

Wager, Tor D., Lauren Y. Atlas, Martin A. Lindquist, Mathieu Roy, Choong-Wan Woo, and Ethan Kross. 2013. "An FMRI-Based Neurologic Signature of Physical Pain." *New England Journal of Medicine* 368 (15): 1388–97. https://doi.org/10.1056/NEJMoa1204471.

Woo, Choong-Wan, Liane Schmidt, Anjali Krishnan, Marieke Jepma, Mathieu Roy, Martin A. Lindquist, Lauren Y. Atlas, and Tor D. Wager. 2017. "Quantifying Cerebral Contributions to Pain beyond Nociception." *Nature Communications* 8 (1): 14211. https://doi.org/10.1038/ncomms14211.

Yeo, Thomas B. T., Fenna M. Krienen, Jorge Sepulcre, Mert R. Sabuncu, Danial Lashkari, Marisa Hollinshead, Joshua L. Roffman, et al. 2011. "The Organization of the Human Cerebral Cortex Estimated by Intrinsic Functional Connectivity." *Journal of Neurophysiology* 106 (3): 1125–65. https://doi.org/10.1152/jn.00338.2011.

Zheng, Weihao, Choong-Wan Woo, Zhijun Yao, Pavel Goldstein, Lauren Y Atlas, Mathieu Roy, Liane Schmidt, et al. 2020. "Pain-Evoked Reorganization in Functional Brain Networks." *Cerebral Cortex* 30 (5): 2804–22. https://doi.org/10.1093/cercor/bhz276.

Acknowledgements

We wish to thank André Cyr and Carollyn Hurst from the Unité de Neuroimagerie Fonctionnelle for their help with the data acquisition, and Andréanne Proulx, François Paugman and Pierre Bellec for their insights on the analyses as part of BrainHack School. We also want to acknowledge Compute Canada for the computational resources allocated to conduct the analyses. M.-E.P. was supported by scholarships from the Fonds de Recherche du Québec-Nature et Technologies and from the Institut de Valorisation des Données. This study was funded by a discovery grant from the Natural Sciences and Engineering Research (RGPIN-2018-06799) to P.R. and the NIH (R01 DA035484) to T.D.W.

Credit statement

Marie-Eve Picard: Conceptualization, Software, Formal analysis, Writing-Original draft, Visualization. **Miriam Kunz**: Conceptualization, Methodology and data acquisition, Data curation. **Jen-I Chen**: Methodology and data acquisition, Data curation, Formal analysis. **Michel-Pierre Coll**: Software, Validation, Visualization. **Étienne Vachon-Presseau**: Conceptualization, Methodology, Validation. **Tor D. Wager**: Conceptualization, Validation, Funding acquisition. **Pierre Rainville**: Conceptualization, Methodology and data acquisition, Writing-Original draft, Funding acquisition. All authors reviewed and approved the final version.

Data Availability

Participants did not provide informed consent for the sharing of individual data but unthresholded group maps are available upon request. Unthresholded patterns are available on Neurovault (https://neurovault.org/collections/13924/).

Code Availability

All custom Python scripts used to produce the analyses are available at https://github.com/me-pic/picard_feps_2022.