Regulation of late-acting operons by three transcription factors and a CRISPR-Cas component

during *Myxococcus xanthus* development

Shreya Saha and Lee Kroos*

Department of Biochemistry and Molecular Biology, Michigan State University, East Lansing, MI

48824, USA.

Running title: Complex regulation of sporulation genes during *M. xanthus* development

*For correspondence. E-mail kroos@msu.edu; Tel. (+1) 517 355 9726; Fax (+1) 517 353 9334.

Summary

Upon starvation rod-shaped *Myxococcus xanthus* bacteria form mounds and then differentiate

into round stress-resistant spores. Little is known about the regulation of late-acting operons

important for spore formation. C-signaling has been proposed to activate FruA, which binds

DNA cooperatively with MrpC to increase transcription of many genes. We report that this

model can explain regulation of the *fadIJ* operon involved in spore metabolism, but not that of

the spore coat biogenesis operons *exoA-I*, *exoL-P*, and *nfsA-H*. Rather, a mutation in *fruA*

increased the transcript levels from these operons early in development, suggesting negative

regulation by FruA initially, and a mutation in *mrpC* affected transcript levels from each operon

differently. FruA bound to all four promoter regions *in vitro*, but strikingly each promoter

region was unique in terms of whether or not MrpC and the DNA-binding domain of Nla6

bound, and in terms of cooperative binding. Furthermore, the DevI component of a CRISPR-Cas

system is a negative regulator of all four operons, based on transcript measurements. Our

results demonstrate complex regulation of sporulation genes by three transcription factors and

a CRISPR-Cas component, which we propose thwarts viral intrusion while making spores suited

to withstand starvation and environmental insults.

Introduction

The Gram-negative soil bacterium *Myxococcus xanthus* provides an attractive model system to study signal-induced gene regulation and bacterial community behavior (1). Upon nutrient depletion, cells move on solid surfaces and form mounds, within which some of the rod-shaped cells differentiate into round, stress-resistant spores. During this multicellular developmental process of fruiting body formation, a majority of the population undergoes lysis, while some cells remain outside of fruiting bodies as peripheral rods (2, 3). Transcriptomic analyses indicate that approximately 1500 (4, 5) to 4000 (6) genes (i.e., 20-50% of the genes) are up- or down-regulated during *M. xanthus* development.

A signal-responsive gene regulatory network (GRN) governs the developmental process of *M. xanthus* (7). Starvation triggers production of the intracellular secondary messenger molecules (p)ppGpp (8, 9) and c-di-GMP (10), which lead to production of the extracellular A- and C-signals (9, 11-14) and exopolysaccharide (10), respectively. Other signals are implicated in the developmental process, but are less well-characterized in terms of their identity and/or their impact on the GRN (15).

Among the better-characterized developmental signals of *M. xanthus*, early studies showed that C-signal acts later than (p)ppGpp and A-signal (9, 11, 16, 17). The exchange of C-signal between cells required their movement into alignment, possibly into contact (18-20). An increasing level of C-signaling was proposed to ensure that cells first build mounds and then form spores, resulting in multicellular fruiting bodies (21-23). The C-signal appears to be a proteolytic fragment of the CsgA protein (24-26) and/or diacylglycerols produced by cardiolipin phospholipase activity of full-length CsgA (27), but the mechanism(s) of signal transduction

remain to be elucidated. C-signaling appears to posttranslationally activate the transcription

factor FruA (28, 29) (Fig. 1), but the mechanism is unknown. FruA is similar to response

regulators of two-component signal-transduction systems (30). Response regulators are

typically activated by phosphorylation by a histidine kinase (28). However, for several reasons,

phosphorylation is unlikely to be the mechanism by which FruA is activated in response to C-

signaling (29, 31).

Transcription of the *fruA* gene is regulated by a cascade of starvation-responsive

transcription factors that includes MrpC (7, 32) (Fig. 1), which binds upstream of the *fruA*

promoter and stimulates transcription (33, 34). In addition to transcriptional control of the

mrpC gene by starvation (35), the level of MrpC protein is regulated by proteolysis (Fig. 1),

slowing the pace of development under the normal conditions of starvation (36, 37) and halting

development if sufficient nutrients are added back before rods become committed to spore

formation (38, 39). An N-terminally truncated form of MrpC called "MrpC2" was proposed to

be important for development (34), but recent work indicated that MrpC2 is likely an artifact of

the method of cell lysate preparation and that full-length MrpC is the active form *in vivo* (35), at

least in part due to a threonine/serine motif near its N-terminus (40). Both MrpC and MrpC2

bind to DNA *in vitro* (34, 35, 41).

In vitro, MrpC2 and FruA bind cooperatively to the promoter regions of many

developmentally-regulated genes (41). Mutational analyses of five such promoter regions (*dev*,

fmgA, *fmgBC*, *fmgD*, and *fmgE*) suggest that cooperative binding of MrpC and FruA to a site just

upstream of the promoter stimulates transcription *in vivo*, forming a feed-forward loop (Fig. 1,

only *dev* is shown), whereas binding to nearby sites can increase or decrease transcription (31,

42-45). The different arrangements of binding sites with different affinities for MrpC and FruA

may explain the observed differences between genes in terms of their spatiotemporal patterns

of expression and their extent of dependence on C-signaling for expression (7, 17, 46). The

levels of MrpC, FruA, and CsgA normally rise in starving rods engaged in mound building (3), but

the addition of sufficient nutrients triggers rapid proteolysis of MrpC and prevents the normal

increase in transcripts for proteins involved in sporulation (38, 39). Combinatorial control of

gene expression by MrpC and FruA appears to ensure that only starving rods (in which MrpC

and FruA are synthesized and stably accumulate) aligned in mounds (which enhances short-

range C-signaling between rods, presumably activating FruA within them) commit to spore

formation (7, 19, 31, 38, 46).

Among the genes combinatorially controlled by MrpC and FruA, early studies showed that

mutations in the *dev* operon can cause severe defects in sporulation (47-50). The *dev* operon

includes eight genes comprising a CRISPR-Cas system that has been proposed to protect

developing *M. xanthus* against bacteriophage lysogenization (51). The first gene in the *dev*

operon, *devI*, codes for a 40-residue protein that strongly inhibits sporulation when

overproduced (52), delays sporulation by about 6 h when produced at the normal level (29, 53),

and exerts weak positive autoregulation on *dev* transcript accumulation (29). In contrast, in-

frame deletions in three genes of the *dev* operon (*devTRS*) increase accumulation of the *dev*

transcript 10-fold during development (29, 52, 53), indicating that DevTRS proteins exert strong

negative autoregulation (Fig. 1). The negative autoregulation by DevTRS proteins is crucial in

order to prevent DevI overproduction from inhibiting sporulation (52, 53) (Fig. 1). The DevTRS

proteins are similar to Cas8a1, Cas7, and Cas5, respectively, of the CRISPR-associated complex

for antiviral defense (Cascade) (51-54).

The products of several other operons act late during the developmental process, impacting

the metabolism of nascent spores (*fadIJ*) and the biogenesis of their coat polysaccharide (*exoA-*

I, *exoL-P*, and *nfsA-H*), but the regulation of these operons is not fully understood (Fig. 1). The

putative *fadIJ* operon is normally induced twofold in sporulating cells, but not in *csgA*, *fruA*, or

mrpC mutants (55). Recently published data support that *fadIJ* are co-transcribed as an operon

(56). FadIJ are enzymes in a fatty acid β-oxidation pathway that can impact spore structure and

resistance properties (55). The *nfsA-H* operon is up-regulated in spores formed during the

starvation-induced developmental process and in spores induced artificially by glycerol addition

(57). The starvation-induced up-regulation fails in a *csgA* mutant, is diminished in a *devRS*

mutant, and, unusually, is greater in a *fruA* mutant, indicative of negative regulation by FruA

(57). The NfsA-H proteins form a complex involved in biogenesis of the coat polysaccharide to

generate compact and rigid stress-resistant spores (58-60). The *exoA-I* operon (58) is normally

induced late in development, but not in *csgA* or *devRS* mutants (61). Transcription depends on

binding of FruA (62) and Nla6 (63) to the *exoA-I* promoter region. Interestingly, Nla6 is part of a

cascade of enhancer-binding proteins (EBPs) that responds to starvation early in development

(32) and leads to production of MrpC and FruA (7) (Fig. 1). The Nla6 DNA-binding domain (DBD)

also binds to the promoter region of the predicted *MXAN_3259–MXAN_3263* operon and

appears to directly regulate transcription (63). Recently published data support that the five

genes are co-transcribed as an operon (56), which was renamed the *exoL-P* operon based on

functional studies (64). The ExoA-I and ExoL-P proteins are involved in the synthesis, export,

and modification of the spore coat polysaccharide (58, 59, 64).

Here, we report systematic investigation of the late-acting *fadIJ*, *nfsA-H*, *exoA-I*, and *exoL-P*

operons with respect to the phenotypes of mutants and the regulation of gene expression. We

found that the *nfs* and *exo* mutants exhibit defects in sporulation earlier than established

previously, whereas a *fad* mutant initially made more sonication-resistant spores than a "wild-

type" (WT) strain, but later made a similar number of mature spores. Cooperative binding of

MrpC and C-signal-activated FruA appears to explain up-regulation of *fadIJ* late in development,

but not the regulation of *nfsA-H*, *exoA-I*, and *exoL-P.* FruA negatively regulates all four operons

early in development and MrpC affects each operon differently. Nla6 up-regulates all four

operons and we discovered that its DBD binds cooperatively with FruA to the *nfsA-H*, *exoL-P*,

and *exoA-I* promoter regions, and also binds cooperatively with MrpC to the *nfsA-H* promoter

region. We also found that all four operons are subject to down-regulation by a CRISPR-Cas

system, especially when the DevI component is overexpressed, as in a *devS* mutant. We

conclude that three transcription factors and a CRISPR-Cas component regulate the late-acting

operons and we propose that complex, differential regulation of genes involved in spore

metabolism and coat biogenesis produces spores suited to withstand environmental insults.

Results

**nfs *and* exo *mutants exhibit defects early in the sporulation process*

In previous work, we established methods to systematically analyze *M. xanthus* development

under submerged culture conditions (29). Our method of submerged culture involves growing

cells in nutrient medium followed by sedimentation, resuspension in starvation buffer, and

incubation in a plastic container. Cells attach to the container bottom and develop under a

layer of buffer.

To determine the effects of mutations in the late-acting operons under our conditions of

submerged culture, we compared the development of mutants with that of WT laboratory

strain DK1622 (65) from which the mutant strains were derived. Specifically, we examined the

effects of an insertion in *fadI* (55), a deletion of the entire *nfsA-H* operon (57), an insertion in

exoC (58, 61), and an insertion in *exoL* (63, 64). As expected (29), the WT strain formed mounds

by 18 h poststarvation (PS) and the mounds began to darken by 30 h (Fig. 2). Darkening

typically correlates with spore formation. The *fadI* mutant was indistinguishable from the WT

strain. The *nfsA-H* mutant was delayed by about 3 h in mound formation and appeared to be

delayed and reduced in mound darkening. The *exoC* and *exoL* mutants formed normal-looking

mounds by 18 h, but subsequent mound darkening was reduced compared to the WT strain.

To quantify changes at the cellular level, we harvested samples from submerged culture and

either treated the samples with glutaraldehyde to fix cells or left the samples untreated (29).

Using the untreated samples, we quantified "sonication-resistant spores" microscopically and

"mature spores" that were heat- and sonication-resistant and capable of germination and

colony formation on nutrient agar. We used the fixed samples to quantify "sonication-sensitive

cells" microscopically (i.e., the total number of cells observed in the fixed sample minus the

number of sonication-resistant spores observed in the corresponding untreated sample). The

vast majority of sonication-sensitive cells were rod-shaped, but we also observed a small

percentage of "transitioning cells" (TCs) intermediate in morphology between rods and spores

(46) in some of the fixed samples.

As expected (29), we first observed sonication-resistant spores at 27 h PS in samples of the

WT strain, and the spores rose to ~2% at 48 h (Fig. 3). The *fadI* mutant formed more spores

than the WT strain. Consistent with the observed mound darkening defects (Fig. 2), the *nfsA-H*

mutant formed less spores than the WT strain, and the *exoC* and *exoL* mutants did not form

sonication-resistant spores (at a detection limit of 0.05%) (Fig. 3 and Table S1). Neither did the

exoC and *exoL* mutants form mature spores at 72 h (at a detection limit of 0.00001%), while the

nfsA-H mutant made ~15% as many mature spores as the WT strain, and the *fadI* mutant made

a similar number of mature spores as the WT strain (Table S1).

The WT strain exhibited a decline of sonication-sensitive cells during development similar to

that reported previously (29), with only ~30% of the cells remaining at 18 h PS and only ~6%

remaining at 48 h (Fig. S1A), presumably reflecting lysis of the majority of cells during

submerged culture development (3). The mutants showed similar decreases in cell number as

the WT strain (Fig. S1A). As noted above, we observed a small percentage of TCs in some of the

fixed samples. For the WT strain, ~3% of the cells were TCs at 24 h (Fig. S1B), when ~20% of the

cells remained (Fig. S1A). Thus, TCs comprised 15% of the total population at 24 h, before

sonication-resistant spores were observed (Fig. 3), in agreement with *in situ* observations using

confocal microscopy (46). The *fadI* mutant showed about half as many TCs as the WT strain at

most time points (Fig. S1B), perhaps due to the *fadI* mutant forming more sonication-resistant

spores (Fig. 3). The *exo* and *nfs* mutants exhibited less TCs at 24 h (~0.4-0.7%), but in each case

the percentage rose to at least 1% later (Fig. S1B), suggesting that these mutants begin to

change shape, but are impaired early in the process of making spores.

We conclude that the *exoC* and *exoL* mutants showed stronger defects in sporulation than

the *nfsA-H* mutant, whereas the *fadI* mutant initially made more sonication-resistant spores

than the WT strain, but later made a similar number of mature spores. All four mutants

exhibited less TCs than the WT strain early in the sporulation process.

Transcript levels from the late-acting operons are very low in the absence of C-signaling

To determine the effects of loss of C-signaling on expression of the late-acting operons, we

used methods established previously (29) to measure transcript levels in the WT strain and a

csgA insertion mutant at 18-30 h PS, the period leading up to and including the time that many

cells commit to spore formation and sonication-resistant spores begin to be observed (38) (Fig.

3). As expected for the WT strain (39), transcript levels rose during development (Fig. 4). As

noted previously (53), the *exoA* transcript level varied greatly between biological replicates (Fig.

4A), which we do not understand. The *exoL* transcript level likewise varied greatly (Fig. 4B).

The *nfsA* (Fig. 4C) and *fadI* (Fig. 4D) transcript levels varied less and on average increased only

~fourfold, much less than the *exoA* (~24-fold) and *exoL* (~70-fold) transcript levels.

In the *csgA* mutant, the transcript levels did not increase during development (Fig. 4,

asterisks indicate $p < 0.05$ in Student's two-tailed *t*-tests comparing the mutant to the WT strain

at each time point). Asterisks are absent above the *exoA* and *exoL* transcript levels of the *csgA*

mutant because the statistical test yielded $p > 0.05$ due to the large variation between

biological replicates of the WT strain. Nevertheless, the *exoA* and *exoL* transcript levels of the

csgA mutant were low for all replicates at each time point. The low transcript levels could be

due to decreased synthesis and/or increased degradation in the *csgA* mutant compared with

the WT strain. To measure the transcript degradation rates, we added rifampicin to inhibit

transcription at 30 h PS and determined the transcript levels at intervals thereafter. The

transcript degradation rates did not differ significantly between the *csgA* mutant and the WT

strain (Fig. 5), suggesting that decreased synthesis primarily accounts for the low transcript

levels in the absence of C-signaling. In agreement, reporter activity from fusions to *exoC* (61),

nfsA (57), and *fadI* (55) was very low in *csgA* mutants relative to WT strains during

development.

Considerable evidence supports a model in which C-signaling activates FruA

posttranslationally in order to increase transcription of genes during *M. xanthus* development

(28, 29). We showed previously that the FruA level is about twofold lower in the *csgA* mutant

than in the WT strain at 18-30 h PS (29). Boosting the FruA level in the *csgA* mutant to the WT

level using a vanillate-inducible promoter (P*van*) fused to *fruA* did not increase the transcript

levels of five genes or operons (*dev*, *fmgA*, *fmgBC*, *fmgD*, and *fmgE*) known to be under

combinatorial control of FruA and MrpC. Neither did boosting the level of FruA D59E (with a

phosphomimetic substitution in its receiver domain) using a P*van*-*fruA D59E* fusion.

We tested whether boosting the level of FruA or its D59E variant in the *csgA* mutant

increases the transcript levels of the late-acting operons. The transcript levels remained low in

all cases (Fig. S2), suggesting that neither a low level of FruA nor a lack of D59 phosphorylation

causes the transcript levels to remain low in the *csgA* mutant. Rather, these results are

consistent with evidence that C-signaling regulates FruA activity by a mechanism other than

phosphorylation of its receiver domain (29, 31).

Transcript levels from the late-acting operons are unexpectedly high in a* fruA *mutant

If C-signaling activates FruA and the activated FruA increases transcription of the late-acting

operons, then transcript levels are expected to be low in a *fruA* insertion mutant, as observed

for the *dev* operon (29). Instead, the *exoA* and *exoL* transcript levels were greater in the *fruA*

mutant than in the WT strain at 18 to 24 h PS (Fig. 4A and 4B), and the *nfsA* transcript level in

the *fruA* mutant exceeded that in the WT strain at 18 h (Fig. 4C), suggesting that FruA is a

negative regulator of some late-acting operons early in development. In contrast, the *fadI*

transcript level in the *fruA* mutant was similar to that in the WT strain at 18 h and less than that

in the WT strain at later times (Fig. 4D), consistent with a model in which C-signaling activates

FruA and the activated FruA increases *fadI* transcription. Activated FruA may also increase *nfsA*

transcription at 27 and 30 h, since the *nfsA* transcript level in the *fruA* mutant was less than that

in the WT strain at those times (Fig. 4C).

Interestingly, the average transcript levels from the late-acting operons were greater in the

fruA mutant than in the *csgA* mutant, except for the very low *fadI* transcript level in both

mutants at 27 and 30 h PS (Fig. 4). As noted above, FruA is present, but its level is about

twofold lower in the *csgA* mutant than in the WT strain at 18-30 h (29). In the *csgA* mutant,

FruA is expected to remain unactivated due to the absence of C-signaling (28, 29). Therefore, it

is possible that unactivated FruA negatively regulates the late-acting operons in the *csgA*

mutant. Negative regulation by unactivated FruA may also account for the lower levels of *exoA*,

exoL, and *nfsA* transcripts we observed early in development of the WT strain as compared with

the *fruA* mutant (Fig. 4) (see Discussion).

The exoA*, *exoL*, and* nfsA *transcript levels of* mrpC *and* fruA *mutants differ

MrpC appears to directly stimulate transcription from the *fruA* promoter (33). In agreement,

FruA was not detected in an *mrpC* in-frame deletion mutant at 18-30 h PS (29). Hence, the

mrpC mutant lacks both MrpC and a detectable level of FruA. To compare the *mrpC* mutant

with the *fruA* mutant (lacking FruA but with MrpC present at its normal level) (29), we

measured the transcript levels from the late-acting operons in the mutants and the WT strain in

parallel. Strikingly, the *exoA*, *exoL*, and *nfsA* transcript levels in the *mrpC* and *fruA* mutants

differed, and in each case the pattern of effects was unique. The average *exoA* transcript level

was greater in the *mrpC* mutant than in the *fruA* mutant or in the WT strain at all times (Fig. 6A,

asterisks above brackets indicate *p* < 0.05 in Student's two-tailed *t*-tests comparing the mutants

at the same time point and asterisks above error bars compare a mutant to the WT strain at the

same time point). The comparison of mutants suggests that MrpC negatively regulates the

exoA transcript level independently of FruA, since the loss of both transcription factors in the

mrpC mutant increased the *exoA* transcript level more than loss of only FruA in the *fruA*

mutant. The *exoL* transcript level in the *mrpC* mutant exceeded that in the WT strain at 18 and

24 h, and on average exceeded that in the *fruA* mutant at 18 h, but on average was less than in

the *fruA* mutant at 24 h, and at 30 h the difference met a test of statistical significance (Fig. 6B).

In this case, comparison of the mutants suggests that, independently of FruA, MrpC negatively

regulates the *exoL* transcript level early in development and positively regulates it later. The

average *nfsA* transcript level was lower in the *mrpC* mutant than in the *fruA* mutant or in the

WT strain at all times (Fig. 6C), suggesting that MrpC positively regulates the *nfsA* transcript

level independently of FruA (i.e., opposite the regulation of *exoA*). The *fadI* transcript level was

lower in both mutants than in the WT strain at all times (Fig. 6D), consistent with positive

combinatorial control by MrpC and FruA, as observed for *dev* (42), *fmg* (31, 43-45), and likely

many other developmentally-regulated genes or operons (41).

To determine whether the lack of both MrpC and a detectable level of FruA in the *mrpC*

mutant affects the transcript degradation rates, we added rifampicin to inhibit transcription at

18 h PS and determined the transcript levels at intervals thereafter. We chose 18 h for this

analysis since the *exoA* (Fig. 6A) and *exoL* (Fig. 6B) transcript levels were elevated in the *mrpC*

mutant relative to the WT strain at that time. The transcript degradation rates did not differ

significantly between the *mrpC* mutant and the WT strain (Fig. S3), suggesting that increased

synthesis (rather than decreased degradation) primarily accounts for the elevated *exoA* and

exoL transcript levels in the *mrpC* mutant.

FruA can positively regulate* exoA*, * exoL*, and* nfsA *in the absence of MrpC

To examine the effects of FruA in the absence of MrpC, we used the P$_{van}$-*fruA* fusion mentioned

previously (29) (Fig. S2) to produce FruA in the *mrpC* mutant. The inducer (vanillate) was added

during growth and at 0 h PS. At 6 h, the FruA level in the *mrpC* P$_{van}$-*fruA* mutant was about

threefold greater than in the WT strain, but their FruA level was similar at 12 and 18 h (Fig.

S4A). Interestingly, the *mrpC* P$_{van}$-*fruA* mutant formed immature mounds by 12 h (Fig. S5),

although subsequent mound darkening was reduced compared to the WT strain (Fig. S6) and

spores were not detected (Table S1). Nevertheless, mound formation suggests that C-signaling

is activating FruA to some extent, since a complete block, as in *mrpC*, *fruA*, and *csgA* mutants,

results in no mound formation (29) (Fig. S5).

To determine the effects of FruA on regulation of the late-acting operons in the absence of

MrpC, we measured the transcript levels in the *mrpC* P$_{van}$-*fruA* mutant in parallel with those in

the *mrpC* and *fruA* mutants and the WT strain at 6-18 h PS. We chose to collect samples during

that period of development since FruA was present (Fig. S4A) and appeared to be active (Fig.

S5) in the *mrpC* P$_{van}$-*fruA* mutant and the WT strain, as described above. The *exoA* (Fig. 7A) and

exoL (Fig. 7B) transcript levels in the *mrpC* P$_{van}$-*fruA* mutant exceeded those in the WT strain at

all times, and on average exceeded those in the *mrpC* and *fruA* mutants, except the *exoL*

transcript level in the *mrpC* mutant was similar to that in the *mrpC* P$_{van}$-*fruA* mutant at 18 h.

We conclude that FruA can positively regulate the *exoA* and *exoL* transcript levels in the

absence of MrpC. This regulation may involve C-signal-activated FruA since the *mrpC* P$_{van}$-*fruA*

mutant formed immature mounds (Fig. S5 and S6), whereas *mrpC*, *fruA*, and *csgA* mutants fail

to form mounds (29) (Fig. S5). The *nfsA* transcript level in the *mrpC* P$_{van}$-*fruA* strain exceeded

that in the WT strain at 6 h and that in the *mrpC* mutant at 6 and 12 h (Fig. 7C), also indicative

of positive regulation by FruA (perhaps activated FruA) in the absence of MrpC. On the other

hand, the *fadI* transcript level in the *mrpC* P$_{van}$-*fruA* strain was less than that in the WT strain

(Fig. 7D), as observed for the *mrpC*, *fruA*, and *csgA* mutants (Fig. 4D, 6D, and 7D), supporting

that positive regulation requires both MrpC and activated FruA.

Nla6 positively regulates the transcript levels of the late-acting operons

The Nla6 transcription factor appears to be a direct regulator of the *exoA-I* and *exoL-P* operons

since a protein with the *E. coli* maltose-binding protein (MBP) fused to the Nla6 DBD (MBP-Nla6

DBD) bound to promoter region DNA fragments (63). A comparison of the *exoA* and *exoL*

transcript levels in an *nla6* insertion mutant and a WT strain suggested that Nla6 positively

regulates both during the first 8 h PS, and negatively regulates both at 24 h (63). To facilitate

planned strain construction, we constructed a new *nla6* insertion mutant and compared it with

the one characterized previously (63, 66). The new mutant is tetracycline-resistant (Tc^r) and

the one described previously is kanamycin-resistant (Km^r). Both mutants formed immature

mounds by 12 h, but failed to progress to more mature mounds with distinct, round edges by

18 h (Fig. S5). Later during development, the Km^r *nla6* mutant mounds matured somewhat at

24-30 h, but failed to darken by 36-48 h (Fig. S6). The Tc^r *nla6* mutant mounds did not mature

until 36 h and also failed to darken by 48 h. Spores were not detected for either mutant (Table

S1).

The two *nla6* mutants were indistinguishable in terms of the molecular markers we tested.

Since our results showed that FruA and MrpC impact transcript levels of the late-acting operons

at 18 h PS (Fig. 4 and 6) and in some cases earlier during development (Fig. 7), we measured the

levels of FruA, MrpC, and transcripts from the late-acting operons in the *nla6* mutants and the

WT strain at 6-18 h. The FruA and MrpC protein levels in the *nla6* mutants were similar to

those in the WT strain at 6-18 h (Fig. S4). The *exoA*, *nfsA*, and *fadI* transcript levels in the WT

strain increased at 18 h, but those in the *nla6* mutants failed to increase (Fig. 8), suggesting that

Nla6 positively regulates those transcript levels. The *exoL* transcript level in the WT strain did

not increase at 18 h, but on average that in the *nla6* mutants was less (Fig. 8B), suggesting that

Nla6 also positively regulates the *exoL* transcript level. Because the *exoA*, *exoL*, and *nfsA*

transcript levels of the *mrpC* and/or *fruA* mutants exceeded those of the WT strain early in

development (Fig. 4, 6, and 7), we tried to construct *mrpC nla6* and *fruA nla6* double mutants,

but our efforts were unsuccessful, so we were unable to determine whether positive regulation

by Nla6 could account for the elevated transcript levels in the *mrpC* and *fruA* mutants relative

to the WT strain.

Binding of MBP-Nla6 DBD, MrpC, and FruA differs at the promoter region of each of the late-

acting operons

As a possible indication of direct regulation by Nla6, MrpC, and FruA, we tested for binding to a

DNA fragment upstream of the predicted start codon of the first gene in each of the late-acting

operons. Figure S7 shows features of each promoter region fragment based on previous work.

FruA and MrpC were shown to bind cooperatively to the *dev* promoter region fragment (42)

(Fig. S7A), which was used here as a control. The FruA DBD was shown to bind to at least two

sites in the *exoA-I* promoter region fragment (62) and MBP-Nla6 DBD was also shown to bind,

presumably *via* interaction with DNA sequences similar to a half-site consensus sequence (63)

(Fig. S7B). MBP-Nla6 DBD was also shown to bind to the *exoL-P* promoter region fragment (63)

and three matches to the half-site consensus sequence were identified (Fig. S7C). ChIP-seq

analysis suggested that MrpC binds to a site upstream of *nfsA-H* at 18 h PS (41) (Fig. S7D). Here,

we used electrophoretic mobility shift assays (EMSAs) to test for binding of purified MBP-Nla6

DBD, MrpC, and FruA separately, and in combination because MrpC and FruA were shown

previously to bind cooperatively to the promoter regions of *dev* (42), *fmg* (31, 43-45), and other

developmentally-regulated genes or operons (41).

As expected (42), MrpC and FruA bound individually to the *dev* promoter region fragment

(Fig. 9A, lanes 4, 5, 7, 8), and the combination of MrpC and FruA exhibited greater than additive

binding (lanes 11, 14), indicative of cooperative binding. MrpC and FruA DBD also appeared to

bind cooperatively (Fig. S8A, lane 4), indicating the N-terminal domain of FruA was not

required. Binding of MBP-Nla6 DBD was not detected at 4 μM (Fig. 9A, lane 2), consistent with

previous reports (32, 63), but binding was detected at 16 μM and increased at greater

concentrations (Fig. S8A, lanes 8-10).

The *fadIJ* promoter region fragment yielded results that were in some ways similar to those

for the *dev* fragment. MrpC and FruA bound individually to the *fadIJ* fragment (Fig. 9B, lanes 3,

4), and the combination of MrpC and FruA appeared to bind cooperatively (lane 7), but with

much less cooperative binding than to the *dev* fragment. The order of MrpC and FruA addition

did not affect cooperative binding to either fragment (Fig. 9A, lanes 11, 14 and Fig. S8B, lanes 5,

6). However, MrpC and FruA DBD did not bind cooperatively to the *fadIJ* fragment (Fig. S8B,

lane 10), indicating a requirement for the FruA N-terminal domain, in contrast to the *dev*

fragment. On the other hand, for both fragments, MBP-Nla6 DBD binding was not detected at

4 μM and at that concentration did not affect binding of MrpC or FruA (Fig. 9A, lanes 2, 9, 10,

12, 13 and Fig. 9B, lanes 2, 5, 6). Also like the *dev* fragment, MBP-Nla6 DBD binding to the *fadIJ*

fragment was detected at 16 μM and increased at higher concentrations (Fig. S8B, lanes 14-16).

The apparent cooperative binding of MrpC and FruA to the *fadIJ* fragment, together with our

measurements of the *fadI* transcript level, support a similar mechanism of developmental

regulation of the *fadIJ* operon as proposed for the *dev* operon (29, 42), the *fmg* genes (31, 43-

45), and likely many other genes (41), in which C-signaling activates FruA to bind DNA

cooperatively with MrpC and increase transcription.

MBP-Nla6 DBD at 2 μM bound to the *exoA-I* promoter region fragment (Fig. 9C, lane 2), as

expected (63). FruA also bound (Fig. 9C, lanes 7, 8), as expected based on FruA DBD binding

observed previously (62). Strikingly, MBP-Nla6 DBD and FruA appeared to bind cooperatively

and the order of addition made no difference (Fig. 9C, lanes 10, 13 and Fig. S8C, lanes 4-6).

FruA DBD exhibited very little cooperative binding with MBP-Nla6 DBD (Fig. S8C, lane 10),

suggesting the N-terminal domain of FruA plays an important role. Although MrpC appeared to

negatively regulate the *exoA* transcript level independently of FruA (Fig. 6A), binding of MrpC

was not detected at 4 μM (Fig. 9C, lane 5) or at concentrations ranging up to 20 μM (Fig. S8C,

lanes 12-16), and MrpC did not affect MBP-Nla6 DBD or FruA binding (Fig. 9C, lanes 9, 11, 12,

14). Taken together, our results suggest that MrpC regulates *exoA* indirectly both

independently of FruA (Fig. 6A) and because MrpC is required for synthesis of FruA (29, 62),

which appears to bind cooperatively with Nla6 to the promoter region and increase *exoA-I*

transcription.

We discovered that FruA binds to the *exoL-P* promoter region fragment (Fig. 9D, lane 4).

Binding of MBP-Nla6 DBD was not detected at 4 μM in the experiment shown in Figure 9D (lane

2), but was detected at 4 μM in two other experiments (Fig. S8D, lanes 2, 8). Interestingly,

MBP-Nla6 DBD and FruA appeared to bind cooperatively and the order of addition made no

difference (Fig. 9D, lane 6 and Fig. S8D, lanes 4-6), similar to our results for the *exoA-I* fragment.

Both fragments also yielded similar results in other experiments; FruA DBD exhibited very little

cooperative binding with MBP-Nla6 DBD to the *exoL-P* fragment (Fig. S8D, lane 10) and MrpC

did not affect MBP-Nla6 DBD or FruA binding (Fig. 9D, lanes 5, 7). However, the two fragments

differed in terms of MrpC binding, which was not detected for the *exoA-I* fragment at MrpC

concentrations ranging up to 20 μM (Fig. S8C, lanes 12-16). In contrast, for the *exoL-P*

fragment, although binding of MrpC was not detected at 4 μM in the experiment shown in

Figure 9D (lane 3), binding was detected at 4 μM and increased at higher concentrations in

another experiment (Fig. S8D, lanes 12-16). Therefore, MrpC (as well as FruA and Nla6, which

bind cooperatively) appears to directly regulate *exoL-P*, differentiating its regulation from that

of *exoA-I* (see Discussion).

We found that MBP-Nla6 DBD, MrpC, and FruA could each bind individually to the *nfsA-H*

promoter region fragment (Fig. 9E, lanes 2-7). In combination with MBP-Nla6 DBD at 4 μM,

individual binding of MrpC was not detected and the intensity of the shifted complex increased

at the lagging edge (Fig. 9E, lane 9), suggestive of cooperative binding. In agreement, MBP-Nla6

DBD at 2 μM appeared to bind cooperatively with MrpC, and the order of addition made little

or no difference (Fig. S8E, lanes 4-6). We note that the large sizes of MBP-Nla6 DBD and the

DNA fragment likely explain the small difference in migration of the putative MBP-Nla6

DBD/MrpC/DNA complex compared to the MBP-Nla6 DBD/DNA complex (Fig. S8E, lanes 2, 4-6).

Like MrpC, FruA appeared to bind cooperatively with MBP-Nla6 DBD (Fig. 9E, lane 10 and Fig.

S8E, lanes 10-12). In the mixtures with MBP-Nla6 DBD, individual binding of FruA was detected,

suggestive of less cooperativity than in the case of MrpC. FruA DBD did not appear to bind

cooperatively MBP-Nla6 DBD (Fig. S8E, lane 16), and FruA did not appear to bind cooperatively

with MrpC (Fig. S8E, lanes 20-22). We conclude that MBP-Nla6 DBD, MrpC, and FruA appear to

directly regulate *nfsA-H* and we note that the apparent cooperative binding of MBP-Nla6 DBD

and MrpC to the *nfsA-H* fragment distinguishes it from the *exoA-I* and *exoL-P* fragments.

Altogether, the results of our EMSAs and transcript measurements suggest that *fadIJ*

regulation is similar to that of *dev* (42), *fmg* (31, 43-45), and likely many other genes or operons

(41), involving cooperative binding of MrpC and C-signal-activated FruA. In contrast, regulation

of the other late-acting operons appears to involve cooperative binding of Nla6 and FruA, and

MrpC acting indirectly (*exoA-I*), directly but not binding cooperatively with Nla6 or FruA (*exoL-*

P), or binding cooperatively with Nla6 (*nfsA-H*).

DevI negatively regulates the transcript levels of the late-acting operons

Reporter activity from fusions to *exoC* (61) and *nfsA* (57) was reduced in a *devRS* insertion

mutant compared with a WT strain during development. The *exoA* transcript level was also

reduced in a *devS* in-frame deletion mutant compared with a WT strain during development

(53). To further investigate the effects of mutations in the *dev* operon, which comprises a

CRISPR-Cas system (51), on regulation of the late-acting operons, we measured transcript levels

in *devI* and *devS* in-frame deletion mutants. Sporulation occurs about 6 h earlier than normal in

the *devI* mutant (29, 53), suggesting that DevI delays sporulation of the WT strain. Sporulation

is severely impaired in the *devS* mutant (29, 51) due to loss of negative autoregulation of *dev*

transcription and resulting overproduction of DevI, which strongly inhibits sporulation (52).

We found that the *exoA*, *exoL*, *nfsA*, and *fadI* transcript levels remain low in the *devS*

mutant at 18-30 h PS (Fig. 10). We infer that DevI overproduction in the *devS* mutant inhibits

accumulation of transcripts from the late-acting operons. Although Student's two-tailed *t*-tests

rarely yielded $p < 0.05$ comparing transcript levels in the *devS* mutant with the WT strain at the

corresponding time PS, this was due to the large variation between biological replicates of the

WT strain. We emphasize that transcript levels were low in all biological replicates of the *devS*

mutant at all times PS.

The average transcript levels were greater in the *devI* mutant than the WT strain, except the

nfsA transcript level was similar in both at 30 h PS (Fig. 10). These results are consistent with

the notion that DevI inhibits accumulation of transcripts from the late-acting operons. Here

again, Student's two-tailed *t*-tests rarely yielded $p < 0.05$, but in this case there was large

variation between biological replicates of both the *devI* mutant and the WT strain. Even so,

taken together, the measurements of transcript levels in the *devI* and *devS* mutants support

that DevI is a negative regulator of the late-acting operons.

Discussion

We investigated the function and regulation of genes whose products act near the end of the

M. xanthus developmental process to change metabolism of the spore and protect its contents

by depositing a polysaccharide coat on its surface. We found that the *exoA-I*, *exoL-P*, and *nfsA-*

H operons involved in spore coat biogenesis impact the formation of resistant spores earlier

than shown previously, but do not prevent the initial cellular shape change associated with

sporulation. We discovered that the regulation of the late-acting operons is unusually complex.

It involves three transcription factors that bind to the promoter regions in different

combinations, allowing each promoter to be regulated uniquely. Combinatorial regulation of

this complexity is common in higher eukaryotes (67-69), but not in bacteria. Our results also

show that the late-acting operons are controlled by a CRISPR-Cas system. These systems

typically defend bacteria against viral infection, but are often repurposed for a variety of

functions, including regulation of endogenous gene expression (70, 71). Collectively, our results

suggest that *M. xanthus* evolved complex eukaryotic-like combinatorial control of transcription

and a link to a CRISPR-Cas system to thwart viral intrusion while making spores suited to

withstand starvation and environmental insults.

New insights into the functions of the late-acting operons

Our results show that mutations in *exoC*, *nfsA-H*, and *exoL* reduce the formation of sonication-

resistant spores beginning at 27 h PS (Fig. 3), but do not prevent the initial cellular shape

change associated with starvation-induced sporulation (Fig. S1B). Previously, these mutants

were examined for sporulation at 120 h (58, 63, 64). Our findings indicate a much earlier role of

the ExoA-I, NfsA-H, and ExoL-P proteins during development than established previously. The

nfsA-H mutant made about twofold less sonication-resistant spores than the WT strain at 27-48

h, whereas the *exoC* and *exoL* mutants made less than the detection limit (Fig. 3). The milder

sporulation defect of the *nfsA-H* mutant compared with the *exoC* and *exoL* mutants (Fig. 3 and

Table S1) is consistent with previous reports (58, 63, 64). The *exoC* mutant fails to export spore

coat polysaccharide, whereas the *nfsA-H* mutant appears to export it, but fails to assemble it

properly, producing an amorphous, unstructured coat that nevertheless appears to provide

some heat- and sonication-resistance (58). The *fadI* mutant made about twofold more

sonication-resistant spores than the WT strain (Fig. 3). The *fadI* insertion mutant presumably

has a reduced rate of fatty acid β-oxidation (55), so perhaps altered metabolism enhanced

formation of sonication-resistant spores (Fig. 3), albeit not mature spores at 72 h (Table S1).

Building on the recent observation of TCs in nascent fruiting bodies (46), we devised a

method that allowed us to visualize and enumerate TCs in samples of the starved mutants (Fig.

S1B). These TCs likely resemble cells of *exo* and *nfs* mutants that fail to complete

morphogenesis upon chemical induction of sporulation in liquid culture (58, 59, 64). The lower

percentage of TCs we observed for *exoC*, *nfsA-H*, and *exoL* mutants as compared with the WT

strain at 24 and 27 h (Fig. S1B) may reflect reduced ability to initiate and/or maintain the

cellular shape change associated with sporulation.

Combinatorial regulation by three signal-responsive transcription factors links expression of

late-acting operon to environmental conditions

Eukaryotic-like signaling and gene regulation have been known in *M. xanthus* for decades (72,

73). More recently, combinatorial control, a common theme in eukaryotic GRNs (67-69),

emerged from studies of *M. xanthus* gene regulation (7, 35). Combinatorial control allows

signal integration by utilizing two or more signal-responsive transcription factors to directly

regulate the same gene. Our transcript measurements demonstrate unique regulation of each

late-acting operon and our DNA-binding assays implicate combinatorial control of each

promoter region by two or three transcription factors (Fig. 11). The late-acting operons are

shown twice to indicate differential regulation early (~6-18 h PS) and late (~21-30 h) in

development (gray boxes at right). The early period approximates when mound formation

occurs in the WT strain (Fig. 2, S5, S6, and the cartoon at left in Fig. 11). The late period

includes times leading up to and including the beginning of sporulation (Fig. 3 and 11). Our

transcript measurements focused on the early (Fig. 7 and 8) or late (Fig. 4, 6, 10, and S2) period

and both included 18 h. The results show that transcript levels began to increase markedly by

18 h (*exoA*, *nfsA*, and *fadIJ*) or 24 h (*exoL*) and continued to increase until 30 h. For simplicity,

our model ignores this difference in timing of transcript increase and we focus primarily on

explaining negative regulation during the early period and positive regulation during the late

period. In the model, green arrows and red lines with a barred end indicate positive and

negative regulation, respectively, based on our transcript measurements. Solid versus dashed

arrows/lines represent our interpretation of our EMSA results as indicative of direct versus

indirect regulation. We interpret binding of transcription factors at relatively low

concentrations (≤ 4 μM) to a promoter region fragment as indicative of direct regulation, and

either no detectable binding or binding at relatively high concentrations (≥ 16 μM) as indicative

of indirect regulation. Cartoons below the operons in the model also take into account our

DNA sequence analysis of potential binding sites for the transcription factors (Fig. S9), in an

effort to illustrate how independent, cooperative, and competitive binding could explain the

complex regulation observed. Importantly, much more work will be required to test our model.

For example, binding sites will need to be identified and mutated, and the effects on

transcription measured.

To explain negative regulation of the late-acting operons early in development, our model

proposes that unactivated FruA directly opposes stimulation of transcription by Nla6-P at the

exoA-I, *exoL-P*, and *nfsA-H* promoter regions (Fig. 11). This model could explain a large increase

in the *exoA-I* and *exoL-P* transcript levels between 0 and 1 h PS followed by a comparable

decrease by 12 h in a WT strain (63). FruA is present at 6 h and its level increases threefold by

12 h (Fig. S4A). FruA* appears to be present by 18 h (29), but the kinetics of FruA activation are

unknown. Presumably, mound formation brings cells into proximity, enabling C-signal-

dependent activation of FruA (19, 46), so FruA* increases and unactivated FruA decreases

during mound formation, which occurs primarily between 12 and 18 h under our conditions

(Fig. S5). Therefore, we expect unactivated FruA to be abundant by 6 h (possibly earlier),

perhaps opposing Nla6-P stimulation after the initial increase of *exoA-I* and *exoL-P* transcript

levels (63).

Our model proposing negative regulation by unactivated FruA early in development (Fig. 11)

is consistent with two prior reports of negative regulation by FruA. Expression of a reporter

fused to the *nfsA-H* promoter region was elevated in a *fruA* mutant compared with a WT strain

at 12-18 h PS (57). While our work was in progress, transcriptomic analysis identified > 1500

genes with an elevated transcript level in the *fruA* mutant compared to the WT strain at 12 h

(6). Based on the data of McLoon *et al.* (6), the transcript levels of all genes in the late-acting

operons were elevated in the *fruA* mutant relative to the WT strain at 12 h (Table S2). In our

study, the *exoA-I*, *exoL-P*, and *nfsA-H* transcript levels were elevated in the *fruA* mutant relative

to the WT strain at 18 h (Fig. 4, 6, and 7), but not at 12 h (Fig. 7), suggesting a temporal

difference in development between the studies. The *fadI* transcript level was lower in the *fruA*

mutant than in the WT strain at all times tested in our experiments (Fig. 4D, 6D, and 7D), but its

level in the *fruA* mutant exceeded that in the *csgA* mutant by a small, reproducible amount at

18 h (Fig. 4D, p = 0.002 in a Student's two-tailed *t*-test), supporting weak negative regulation by

unactivated FruA (Fig. 11). Our model proposes that unactivated FruA directly opposes

stimulation of *fadIJ* transcription by cooperative binding of MrpC and FruA* early in

development (Fig. 11 cartoon), rather than opposing stimulation by Nla6-P, for two reasons.

First, a high concentration of MBP-Nla6 DBD was required to detect binding to the *fadIJ*

promoter region fragment (Fig. S8B), suggesting indirect regulation by Nla6-P (Fig. 11). Second,

our analysis revealed potential binding sites for MrpC and FruA or FruA*, but not for Nla6,

upstream of the *fadIJ* promoter (Fig. S9A).

Our model also proposes that MrpC affects transcription of the *exoA-I*, *exoL-P*, and *nfsA-H*

operons early in development (Fig. 11). The *exoA* transcript level was on average at least

twofold greater in the *mrpC* mutant than the *fruA* mutant at 18 h PS (Fig. 6A and 7A),

suggesting negative regulation by MrpC independently of FruA, but we did not detect MrpC

binding to the *exoA-I* promoter region fragment (Fig. S8C) or potential binding sites (Fig. S9D).

Therefore, we propose that MrpC indirectly decreases transcription of the *exoA-I* operon early

in development of the WT strain (Fig. 11). In contrast, our results imply that MrpC directly

decreases *exoL-P* transcription and directly increases that of *nfsA-H* early in development (Fig.

6, 7, 9, S8, and 11).

The cartoons below the *exoA-I*, *exoL-P*, and *nfsA-H* operons in Figure 11 propose that

unactivated FruA and MrpC engage in cooperative or competitive binding with Nla6-P or each

other primarily to inhibit transcription early in development of the WT strain. Our EMSAs (Fig.

9 and S8) and our analysis revealing adjacent potential binding sites (Fig. S9) support the

cooperative binding depicted. We infer competitive binding at overlapping potential binding

sites (Fig. S9). The arrangement of transcription factors in the cartoons (Fig. 11) mimics the

arrangement of potential binding sites (Fig. S9) proposed to play important roles (Supporting

Information, additional discussion). Not all the potential binding sites shown in Figure S9 are

depicted in the Figure 11 cartoons.

To explain positive regulation of the late-acting operons late in development, our model

proposes that C-signaling switches FruA from a negative regulator to a positive regulator, FruA*

(Fig. 11). The switch is most evident comparing the *nfsA* transcript levels in the *fruA* mutant

with those in the WT strain as development proceeds (i.e., the transcript level increases in the

WT strain and decreases in the *fruA* mutant) (Fig. 4C). The switch is less evident from a

comparison of *exoA* transcript levels due to large variation between biological replicates of the

WT strain late in development, but on average the transcript level increases in the WT strain

and decreases in the *fruA* mutant (Fig. 4A). Our findings that ectopic FruA expression in the

mrpC mutant allowed formation of immature mounds (Fig. S5) and greatly increased the *exoA*

transcript level (Fig. 7A) also support a switch to stimulation of transcription by FruA* (Fig. 11).

Similar findings suggest that FruA* increases the *exoL* transcript level (Fig. 7B and 11).

However, the *exoL* transcript level does not decrease in the *fruA* mutant late in development

(Fig. 4B). Therefore, we propose that MrpC can act as a direct positive regulator independently

of FruA* (Fig. 11). In contrast, our results suggest that cooperative binding of MrpC and FruA*

stimulates *fadIJ* transcription late in development (Fig. 11).

The cartoons in Figure 11 shown below the operons late in development reflect our EMSA

results (Fig. 9 and S8) and our analysis of potential binding sites (Fig. S9), using similar reasoning

as explained above, except we propose that FruA* may differ from unactivated FruA in terms of

affinity for potential binding sites and the effect of cooperative binding with Nla6-P or

dephosphorylated Nla6 (Supporting Information, additional discussion). In short, we propose

590 that MrpC and FruA* bind cooperatively near the *fadIJ* promoter to stimulate transcription late

591 in development, whereas binding of MrpC and unactivated FruA farther upstream weakly

592 inhibits transcription stimulation earlier (Fig. 11 cartoons). The cartoons below the *exoA-I*,

593 *exoL-P*, and *nfsA-H* operons propose that FruA* and MrpC bind independently and/or engage in

594 cooperative or competitive binding with Nla6-P (or dephosphorylated Nla6, which is not shown

595 for simplicity) primarily to stimulate transcription late in development of the WT strain.

596  We conclude that each late-acting operon is uniquely regulated by two or three

597 transcription factors acting combinatorially. The positions and affinities of binding sites for

598 Nla6-P or dephosphorylated Nla6, MrpC, and FruA or activated FruA* in each promoter region

599 determine the transcriptional output from independent, cooperative, and competitive binding.

600 This regulatory complexity was masked in transcriptomic studies. The four late-acting operons

601 we investigated were clustered in one (5) or two (4, 6) groups with hundreds of other genes,

602 but our transcript measurements in several mutants revealed unique regulation of each operon

603 (Fig. 4, 6, and 7), our DNA-binding assays showed different binding of transcription factors to

604 each promoter region (Fig. 9 and S8), and our analysis of potential binding sites uncovered

605 diverse, complex architectures (Fig. S9). Collectively, our results and the transcriptomic studies

606 predict that tremendous regulatory complexity will be uncovered by genome-wide

607 measurements of transcript levels in mutants, transcription factor binding studies, and

608 bioinformatic analyses.

609  Since combinatorial control allows signal integration, what are the signals to which the late-

610 acting operons respond? Starvation triggers a cascade of EBPs that includes Nla6-P and Nla28-P

611 (32), and leads to the production of MrpB-P (74), another EBP that in turn leads to production

of MrpC (75) (Fig. 11). However, more specific molecular signals controlling these steps are

unknown. In particular, it is unknown whether Nla6 responds to different signals early and late

in development, or how it switches from positive to negative regulation of the *exo* operons (63)

(Fig. 11). Nutrients and an unknown signal sensed by the Esp system stimulate proteolysis of

MrpC (Fig. 11) to delay or halt development (36-39). The presumed advantage of such a

developmental checkpoint is to ensure that cellular physiology and interactions with other cells,

as well as environmental conditions, warrant commitment to spore formation (38, 76, 77). C-

signaling appears to require the movement of cells into alignment or possibly into contact, and

lead to the formation of active FruA* (18-20, 28, 29, 46). Early in development, unactivated

FruA appears to function as a global negative regulator (6) (Fig. 11), preventing premature

production of some proteins that would be detrimental during mound formation (e.g., Nfs and

Exo proteins involved in polysaccharide export), in addition to conserving cellular resources.

Altogether, our knowledge about the specific signals to which the late-acting operons respond

is quite limited, but in general, nutrient limitation and movement of cells into proximity appear

to play important roles, and are likely influenced by many other factors in the environment of

the developing population.

We propose that combinatorial control of the late-acting operons and hundreds of other

genes by signal-responsive transcription factors allows proper temporal and spatial expression

of genes involved in spore metabolism, coat biogenesis, and other functions in order to

produce spores that are densely packed in fruiting bodies and suited to withstand

environmental insults. Numerous studies show that environmental conditions affect the

resistance and surface characteristics of endospores from *Bacillus* species (78-80). In *Bacillus*

subtilis, several transcription factors positively and negatively fine-tune the expression of

hundreds of genes during sporulation (81-83), ensuring that the resulting spores are endowed

with resistance and surface properties tailored for their environment (84-86). Understanding

how ecology and evolution impact the developmental GRN of *M. xanthus* and related species is

an ongoing and future challenge (6, 87, 88).

The DevI component of a CRISPR-Cas system is a negative regulator of the late-acting operons

Our results confirm and extend the role of DevI as a negative regulator of late-acting operons

during development. We confirmed that the *exoA* transcript level is very low in the *devS* null

mutant (53) and we observed similar results for the *exoL*, *nfsA*, and *fadI* transcript levels (Fig.

10). The cumulative effect of failure to express the late-acting operons may explain the inability

of the *devS* mutant to undergo normal cellular shape change (53). Alternatively, one or more

other genes subject to negative regulation by DevI overproduction may be responsible. We

also found that transcript levels from the late-acting operons were slightly greater on average

in a *devI* mutant than in the WT strain, supporting our conclusion that DevI is a negative

regulator of the late-acting operons (Fig. 11); however, the mechanism of negative regulation

by DevI is unknown.

The *dev* CRISPR-Cas system appears to have evolved recently (52), perhaps in niches where

protection from bacteriophage infection and/or delayed sporulation were advantageous (51,

53). Most natural isolates of *M. xanthus* lack *devI* and their *dev* promoter region is predicted to

be nonfunctional (52). The *dev* operon is dispensable for development of the WT laboratory

strain used in this work (53). However, the *dev* CRISPR-Cas system may protect against

intrusion by bacteriophage Mx8 during development, since a 37-bp sequence in the CRISPR

matches a sequence in the Mx8 integrase gene (51). This hypothesis remains to be tested, as

do the potential benefits of such protection.

In summary, our investigation uncovered complex regulation of genes involved in spore

metabolism and spore coat biogenesis by three transcription factors and a CRISPR-Cas

component during *M. xanthus* development. Further elucidating the signals to which Nla6,

MrpC, and FruA respond, understanding how these transcription factors interact at promoter

regions, and elucidating how DevI inhibits the cellular shape change associated with sporulation

are important goals for the future.

Experimental procedures

Bacterial strains, plasmids and primers

The strains, plasmids and primers used in this study are listed in Table S3.

M. xanthus strain MSS2 with P$_{van}$-*fruA* integrated ectopically was constructed by

electroporating (89) pSS10 into strain DK1622, selecting transformants on CTT agar (1%

Casitone, 10 mM Tris-HCl [pH 8.0], 1 mM KH$_2$PO$_4$-K$_2$HPO$_4$, 8 mM MgSO$_4$ [final pH 7.6] solidified

with 1.5% agar) containing 15 μg/mL tetracycline (90), and verification by colony PCR using

primers pMR3691 MCS G-F and pMR3691 MCS G-R.

M. xanthus strain MSS10 with a plasmid insertion mutation in *nla6* was constructed by

electroporating pSS11 into strain DK1622, selecting transformants on CTT agar containing

tetracycline (15 μg/mL), and verification by colony PCR using primers PMR3487 Rev, Nla6 Fwd4,

and Nla6 Fwd5. To construct pSS11, primers Nla6 Fwd and Nla6 Rev were used to generate a

PCR product using chromosomal DNA from *M. xanthus* strain DK1622 as a template. The

product was combined with DNA amplified from pMR3487 using primers PMR3487G Fwd and

PMR3487G Rev primers, and a Gibson assembly reaction was used to enzymatically join the

overlapping DNA fragments (91). The reaction mixture was transformed into *E. coli* strain

DH5α, transformants were selected at 37°C on Luria-Bertani (LB) (92) 1.5% agar containing 15

μg/mL tetracycline, and the cloned DNA sequence was verified using primers 3487 seq Fwd1,

3487 seq Fwd2, 3487 seq Fwd3, 3487 seq Fwd4, and 3487 seq Fwd5.

To provide a DNA template for amplification of the *nfsA-H* promoter region, pSS14 was

constructed as follows. Primers Nfs -290G and Nfs +83G were used to generate a PCR product

using chromosomal DNA from *M. xanthus* strain DK1622 as a template. The product was

combined with DNA amplified from pMR3487, the overlapping DNA fragments were joined

using a Gibson assembly reaction, the reaction mixture was transformed into *E. coli* strain

DH5α, transformants were selected, and the cloned DNA sequence was verified as described

above for pSS11.

To express MBP-Nla6 DBD for purification, primers Nla DBD For and Nla DBD Rev were used

to generate a PCR product using *nla6* DNA as a template. The PCR product was digest with *Sbf*I

and ligated to *Xmn*I-*Sbf*I-digested pMAL-c5x (NEB), resulting in pMAL-c5x/MBP-ΔCNla6 DBD,

which lacked a CG bp at the beginning of the Nla6 DBD-encoding segment. The missing CG bp

was added by performing site-directed mutagenesis using the QuikChange strategy (Stratagene)

with primers MBP-Nla6 add C fwd and MBP-Nla6 add C rev, resulting in pMAL-c5x/MBP-Nla6

DBD, and the DNA sequence was verified using primer mbp fwd colpcr.

Growth and development of M. xanthus

Strains of *M. xanthus* were grown at 32°C in CTTYE liquid medium (CTT medium with 0.2% yeast

extract) with shaking at 350 rpm. CTT agar was used for growth on solid medium and was

supplemented with 40 µg/mL of kanamycin sulfate or 15 µg/mL of tetracycline as required.

Fruiting body development under submerged culture conditions was performed using MC7 (10

mM morpholinepropanesulfonic acid [MOPS; pH 7.0], 1 mM $CaCl_2$) as the starvation buffer as

described previously (38). Briefly, cells in the mid-exponential phase of growth were collected

by centrifugation, the supernatant was removed, and the cells were resuspended in MC7 buffer

at a density of approximately 1,000 Klett units. A 96-µL sample (designated T_0) was removed

and 4 µL of glutaraldehyde was added from a 50% stock solution to achieve a 2% final

concentration. The sample was stored at 4°C for at least 24 h in order to fix the cells for later

quantification of total cells as described below. For each developmental sample, 1.5 mL of the

1,000-Klett-unit cell suspension plus 10.5 mL of MC7 buffer was added to an 8.5-cm-diameter

plastic petri plate. Upon incubation at 32°C, cells adhere to the bottom of the plate and

undergo development. At the indicated times, developing populations were photographed

using a Leica Wild M8 microscope equipped with an Olympus E-620 digital camera.

Sample collection

At the indicated times PS, the MC7 buffer overlay was replaced with 5 mL of fresh MC7 buffer

or 5 mL of MC7 buffer containing 50 µg/mL of rifampicin to inhibit transcription for

measurement of transcript degradation rates. Developing cells were scraped from the bottom

of the plates, the entire contents were collected in a 15-mL centrifuge tube, and samples were

mixed thoroughly as described previously (29). A 96-μL sample was removed, glutaraldehyde

was added, and the sample was stored as described above for the T_0 sample, fixing the cells for

quantification of total cells as described below. A 400 μL sample was removed and stored at -

20°C for measurement of sonication-resistant spores as described below. For the experiments

shown in Figures 8 and S4, a 100 μl sample was removed and added to an equal volume of 2×

sample buffer (0.125 M Tris-HCl [pH 6.8], 20% glycerol, 4% sodium dodecyl sulfate [SDS], 0.2%

bromophenol blue, 0.2 M dithiothreitol), boiled for 5 min, and stored at -20°C for immunoblot

analysis. Immediately after collecting the samples just described, the rest of the 1,000-Klett-

unit cell suspension was mixed with 0.5 mL of RNase stop solution (5% phenol [pH < 7] in

ethanol), followed by rapid cooling in liquid nitrogen until almost frozen, centrifugation at 8,700

× g for 10 min at 4°C, removal of the supernatant, freezing of the cell pellet in liquid nitrogen,

and storage at -80°C until RNA extraction.

Quantification of total cells, transitioning cells, and sonication-resistant spores

The total number of cells, including rods, elliptical TCs, and round spores, was determined using

the glutaraldehyde-fixed samples collected as described above. Each sample was thawed and

mixed by vortexing and pipetting, diluted with MC7 buffer, sonicated for 10 s, and then all cells

were counted microscopically as described previously (29), except noting the number of cells

that were elliptical or round. The number of sonication-resistant spores in the 400-μL samples

collected as described above was quantified as described previously (38). These samples were

not glutaraldehyde-fixed and were sonicated for 10 s intervals three times with cooling on ice in

between, leaving only round spores (i.e., neither rods nor elliptical TCs were observed). The

total cell number minus the number of sonication-resistant spores was designated the number

of sonication-sensitive cells (consisting primarily of rods and a small percentage of TCs) and was

expressed as a percentage of the total cell number in the corresponding T_0 sample (consisting

only of rods). The number of cells that were elliptical or round in the glutaraldehyde-fixed

sample minus the number of sonication-resistant spores in the corresponding sample that was

not glutaraldehyde-fixed, was designated the number of TCs and was also expressed as a

percentage of the total cell number in the corresponding T_0 sample.

RNA extraction and analysis

RNA was extracted using the hot-phenol method, followed by digestion with DNase I (Roche) as

described previously (36). The RNA (1 μg) was then subjected to cDNA synthesis using

Superscript III reverse transcriptase (InVitrogen) and random primers (Promega), as instructed

by the manufacturers. In parallel, RNA (1 μg) was subjected to cDNA synthesis reaction

conditions without Superscript III reverse transcriptase, as a control. One μl of cDNA at the

appropriate dilution (as determined empirically) and 20 pmol of each primer were subjected to

qPCR in a 25 μl reaction using 2× reaction buffer as described previously (29). qPCR was done

in quadruplicate for each cDNA using a LightCycler® 480 System (Roche). In parallel, a standard

curve was generated for each pair of qPCR primers using genomic DNA (gDNA) from *M. xanthus*

WT strain DK1622 and gene expression was quantified using the relative standard curve

method (user bulletin 2; Applied Biosystems). 16S rRNA was used as the internal standard for

each sample. The primers used for RT-qPCR analysis of 16S rRNA (93) and the late-acting

operons (39) have been described previously.

Relative transcript levels for mutants are the average of three biological replicates after

each replicate was normalized to the transcript level observed for one replicate of the WT

strain at the earliest time point (i.e., either 6 or 18 h PS) in the same experiment (29).

Transcript levels for the WT strain at other times PS were likewise normalized to that observed

for the WT strain at the earliest time point in the same experiment. For the WT strain at the

earliest time point, the transcript levels of at least three biological replicates from different

experiments were normalized to their average, which was set as 1.

To measure transcript degradation rates, samples were collected immediately after addition

of rifampicin (designated t_0) and 8 and 16 min later for RNA extraction and analysis as described

above. For each biological replicate the transcript levels after 8 and 16 min were normalized to

the transcript level at t_0, which was set as 1, and the natural log of the resulting values was

plotted versus minutes after rifampicin treatment (29). The slope of a linear fit of the data was

used to compute the mRNA half-life.

Expression and purification of proteins

To express His-tagged proteins, *E. coli* strain BL21(DE3) was transformed with pET11a/FruA-

DBD-H_8, pET11km/FruA-H_6, or pPH158 (for H_6-MrpC) and transformants were selected on LB

1.5% agar containing 50 μg/mL ampicillin or kanamycin sulfate. In each case, a single colony

was used to inoculate 10 mL of LB supplemented with antibiotic, followed by overnight

incubation at 37°C with shaking. Five mL of overnight culture was used to inoculate 500 mL of

the same medium, followed by continued shaking at 37°C until the culture reached 60-80 Klett

units. IPTG (1 mM final concentration) was added to induce synthesis of the recombinant

protein. After 2 h, cells were harvested as reported previously (94) and stored at -80°C.

To purify His-tagged proteins, a modification of a protocol described previously (95) was

used. Each cell pellet was resuspended in 35 mL of lysis buffer (50 mM Na-phosphate buffer pH

8.0, 500 mM NaCl, 14 mM β-mercaptoethanol, protease inhibitors [1 Roche Mini EDTA-free

tablet/10 mL]). Cells were disrupted by sonication (4 times for 1 min with intermittent cooling

on ice). After centrifugation at 18,000 × g for 10 min at 4°C, the supernatant was mixed with 5

mL of lysis buffer containing 10% w/v Triton X-100, 1 mL of 1 M imidazole (pH 8.0), enough lysis

buffer for 50 mL total, and finally 0.5 mL of Ni-NTA agarose (Qiagen) that had been washed 3

times with lysis buffer containing 20 mM imidazole. The mixture was rotated for 1 h at 4°C.

The Ni-NTA agarose was collected by centrifugation at 700 × g for 3 min at 4°C and washed 4

times with 50 mL of lysis buffer containing 20 mM imidazole, 0.5% w/v Triton X-100, and 25%

w/v glycerol. The Ni-NTA agarose was rotated for 30 min at 4°C with 10 mL of elution buffer

(50 mM Na-phosphate buffer pH 8.0, 500 mM NaCl, 1.4 mM β-mercaptoethanol, protease

inhibitors [1 Roche Mini EDTA-free tablet], 250 mM imidazole, 0.5% w/v Triton X-100, 25% w/v

glycerol). Eluates were dialyzed twice at 4°C against 1 L of storage buffer (10 mM Tris-HCL pH

8.0, 100 mM NaCl, 1 mM β-mercaptoethanol, 10% w/v glycerol) and aliquots were stored at -

80°C. The protein concentration was determined using a Bradford (96) assay kit (Bio-Rad).

To express MBP-Nla6 DBD, *E. coli* strain BL21(DE3) was transformed with pMAL-c5x/MBP-

Nla6 DBD and transformants were selected on LB 1.5% agar containing 100 μg/mL ampicillin.

Five colonies were used to inoculate 80 mL of LB supplemented with antibiotic and 0.2%

dextrose, followed by incubation at 37°C with shaking at 350 rpm until the culture reached 60

Klett units. IPTG (0.3 mM final concentration) was added to induce synthesis of the

recombinant protein. After 2 h, cells were harvested, the cell pellet was resuspended in 10 mL

of column buffer (20 mM Tris-HCl pH 7.5, 200 mM NaCl, 1 mM EDTA, protease inhibitors [1

Roche Mini EDTA-free tablet], 10 mM β-mercaptoethanol), and the cell suspension was stored

at -20°C.

MBP-Nla6 DBD was purified according to instructions in the pMAL™ Protein Fusion &

Purification System manual (NEB). Briefly, the cell pellet was thawed and cells were disrupted

by sonication (8 times for 15 s with intermittent cooling on ice). After centrifugation at 18,000

$\times$ g for 20 min at 4°C, the supernatant was loaded onto a 1 mL amylose gravity-flow column,

which had been prewashed with 5 mL of column buffer. The loaded column was washed with

12 mL of column buffer, then eluted with 2.5 mL of column buffer supplemented with 10 mM

maltose, collecting fractions of 4 drops each (approximately 200 μL). Fractions were subjected

to SDS-PAGE with Coomassie Blue staining. Fractions 4-9, with maximal purified MBP-Nla6

DBD, were pooled, dialyzed as described above, and aliquots were stored at -80°C. The protein

concentration was determined by measuring the absorbance at 280 nm and using the

calculated extinction coefficient of MBP-Nla6 DBD (1.32 [mg/mL]$^{-1}$cm^{-1}).

Electrophoretic mobility shift assays

^{32}P-labelled DNA fragments were generated by PCR using primers labelled with [γ-^{32}P]ATP using

T4 polynucleotide kinase (NEB) according to the manufacturer's instructions. The templates for

PCR were either plasmid DNA or gDNA from *M. xanthus* WT strain DK1622. The primers and

templates were as follows: primers LK1298 and LK1331 with template pPV391 for *dev*, primers

exoA-I -171 fwd and *exoA-I* -1 rev with template gDNA for *exoA-I*, primers *fadIJ* -214 fwd and

fadIJ -5 rev with template gDNA for *fadIJ*, primers *exoL-P* -217 fwd and *exoL-P* +56 rev with

template gDNA for *exoL-P*, and primers *nfsA-H* -166 fwd and *nfsA-H* +35 with template pSS14

for *nfsA-H.* The labeled DNA fragments were purified by electrophoresis on 15%

polyacrylamide gels, followed by visualization using autoradiography, excision, and overnight

elution by soaking in TE buffer (10 mM Tris-HCl pH 8.0, 1 mM EDTA) as described previously

(94).

Binding reactions (10 μL) were performed as reported previously (94), except the reaction

mixtures were incubated at 37°C for 10 min prior to loading on 8% polyacrylamide gels. To

determine whether the order of protein addition was important, one protein was added to the

reaction mixture during an initial 10 min incubation at 37°C, followed by addition of a second

protein and another 10 min incubation at 37°C, prior to loading on 8% polyacrylamide gels.

Gels were dried and exposed to X-ray film for autoradiography as described (94).

Acknowledgements

We thank Monique Floer for advice about high-throughput qPCR and for use of the LightCycler®

480 System. We thank Anthony Garza, Lawrence Shimkets, Ramya Rajagopalan, Wenyuan Shi,

Sumiko Inouye, Montserrat Elias-Arnanz, and Penelope Higgs for sharing strains. We thank

Muqing Ma and Anthony Garza for constructing pMAL-c5x/MBP-ΔCNla6 DBD. This work was

supported by the National Science Foundation (awards MCB-1411272 and IOS-1951025) and by

salary support for L.K. from Michigan State University AgBioResearch.

Author contributions

Conception or design of the study: LK, SS

Acquisition of the data: SS, LK

Analysis or interpretation of the data: SS, LK

Writing of the manuscript: LK, SS

References

1. Yang Z, Higgs P, editors. Myxobacteria: genomics, cellular and molecular biology. Norfolk, UK: Caister Academic Press; 2014.
2. O'Connor KA, Zusman DR. Behavior of peripheral rods and their role in the life cycle of *Myxococcus xanthus*. J Bacteriol. 1991;173(11):3342-55.
3. Lee B, Holkenbrink C, Treuner-Lange A, Higgs PI. *Myxococcus xanthus* developmental cell fate production: heterogeneous accumulation of developmental regulatory proteins and reexamination of the role of MazF in developmental lysis. J Bacteriol. 2012;194(12):3058-68.
4. Munoz-Dorado J, Moraleda-Munoz A, Marcos-Torres FJ, Contreras-Moreno FJ, Martin-Cuadrado AB, Schrader JM, et al. Transcriptome dynamics of the *Myxococcus xanthus* multicellular developmental program. eLife. 2019;8:e50374.
5. Sharma G, Yao AI, Smaldone GT, Liang J, Long M, Facciotti MT, et al. Global gene expression analysis of the *Myxococcus xanthus* developmental time course. Genomics. 2021;113(1 Pt 1):120-34.
6. McLoon AL, Boeck ME, Bruckskotten M, Keyel AC, Sogaard-Andersen L. Transcriptomic analysis of the *Myxococcus xanthus* FruA regulon, and comparative developmental transcriptomic analysis of two fruiting body forming species, *Myxococcus xanthus* and *Myxococcus stipitatus*. BMC Genomics. 2021;22(1):784.
7. Kroos L. Highly signal-responsive gene regulatory network governing *Myxococcus* development. Trends Genet. 2017;33:3-15.
8. Manoil C, Kaiser D. Accumulation of guanosine tetraphosphate and guanosine pentaphosphate in *Myxococcus xanthus* during starvation and myxospore formation. J Bacteriol. 1980;141:297-304.
9. Harris BZ, Kaiser D, Singer M. The guanosine nucleotide (p)ppGpp initiates development and A-factor production in *Myxococcus xanthus*. Genes Dev. 1998;12(7):1022-35.
10. Skotnicka D, Smaldone GT, Petters T, Trampari E, Liang J, Kaever V, et al. A minimal threshold of c-di-GMP is essential for fruiting body formation and sporulation in *Myxococcus xanthus*. PLoS Genet. 2016;12(5):e1006080.
11. Singer M, Kaiser D. Ectopic production of guanosine penta-and tetraphosphate can initiate early developmental gene expression in *Myxococcus xanthus*. Genes and Dev. 1995;9:1633-44.
12. Crawford EW, Jr., Shimkets LJ. The *Myxococcus xanthus socE* and *csgA* genes are regulated by the stringent response. Mol Microbiol. 2000;37(4):788-99.

13. Konovalova A, Lobach S, Sogaard-Andersen L. A RelA-dependent two-tiered regulated proteolysis
cascade controls synthesis of a contact-dependent intercellular signal in *Myxococcus xanthus*. Mol
Microbiol. 2012;84(2):260-75.
14. Konovalova A, Wegener-Feldbrugge S, Sogaard-Andersen L. Two intercellular signals required for
fruiting body formation in *Myxococcus xanthus* act sequentially but non-hierarchically. Mol
Microbiol. 2012;86(1):65-81.
15. Bretl DJ, Kirby JR. Molecular mechanisms of signaling in *Myxococcus xanthus* development. J Mol
Biol. 2016;428:3805-30.
16. Kuspa A, Kroos L, Kaiser D. Intercellular signaling is required for developmental gene expression in
Myxococcus xanthus. Dev Biol. 1986;117:267-76.
17. Kroos L, Kaiser D. Expression of many developmentally regulated genes in *Myxococcus* depends on
a sequence of cell interactions. Genes Dev. 1987;1:840-54.
18. Kim SK, Kaiser D. Cell motility is required for the transmission of C-factor, an intercellular signal that
coordinates fruiting body morphogenesis of *Myxococcus xanthus*. Genes Dev. 1990;4:896-905.
19. Kim SK, Kaiser D. Cell alignment required in differentiation of *Myxococcus xanthus*. Science.
1990;249:926-8.
20. Kroos L, Hartzell P, Stephens K, Kaiser D. A link between cell movement and gene expression argues
that motility is required for cell-cell signaling during fruiting body development. Genes Dev.
1988;2:1677-85.
21. Kim SK, Kaiser D. C-factor has distinct aggregation and sporulation thresholds during *Myxococcus*
development. J Bacteriol. 1991;173:1722-8.
22. Kruse T, Lobedanz S, Berthelsen NM, Sogaard-Andersen L. C-signal: a cell surface-associated
morphogen that induces and co-ordinates multicellular fruiting body morphogenesis and
sporulation in *Myxococcus xanthus*. Mol Microbiol. 2001;40(1):156-68.
23. Li S-F, Lee B, Shimkets LJ. *csgA* expression entrains *Myxococcus xanthus* development. Genes Dev.
1992;6:401-10.
24. Kim SK, Kaiser D. C-factor: a cell-cell signaling protein required for fruiting body morphogenesis of
M. xanthus. Cell. 1990;61:19-26.
25. Lobedanz S, Sogaard-Andersen L. Identification of the C-signal, a contact-dependent morphogen
coordinating multiple developmental responses in *Myxococcus xanthus*. Genes Dev.
2003;17(17):2151-61.
26. Rolbetzki A, Ammon M, Jakovljevic V, Konovalova A, Sogaard-Andersen L. Regulated secretion of a
protease activates intercellular signaling during fruiting body formation in *M. xanthus*. Dev Cell.
2008;15(4):627-34.
27. Boynton TO, Shimkets LJ. *Myxococcus* CsgA, *Drosophila* Sniffer and human HSD17B10 are
cardiolipin phospholipases. Genes Dev. 2015;29:1903-14.
28. Ellehauge E, Norregaard-Madsen M, Sogaard-Andersen L. The FruA signal transduction protein
provides a checkpoint for the temporal co-ordination of intercellular signals in *Myxococcus xanthus*
development. Mol Microbiol. 1998;30:807-17.
29. Saha S, Patra P, Igoshin O, Kroos L. Systematic analysis of the *Myxococcus xanthus* developmental
gene regulatory network supports posttranslational regulation of FruA by C-signaling. Mol
Microbiol. 2019;111:1732-52.
30. Ogawa M, Fujitani S, Mao X, Inouye S, Komano T. FruA, a putative transcription factor essential for
the development of *Myxococcus xanthus*. Mol Microbiol. 1996;22(4):757-67.
31. Mittal S, Kroos L. A combination of unusual transcription factors binds cooperatively to control
Myxococcus xanthus developmental gene expression. Proc Natl Acad Sci USA. 2009;106:1965-70.

32.   Giglio KM, Caberoy N, Suen G, Kaiser D, Garza AG. A cascade of coregulating enhancer binding
proteins initiates and propagates a multicellular developmental program. Proc Natl Acad Sci USA.
2011;108:E431-E9.
33.   Ueki T, Inouye S. Identification of an activator protein required for the induction of *fruA*, a gene
essential for fruiting body development in *Myxococcus xanthus*. Proc Natl Acad Sci USA.
2003;100(15):8782-7.
34.   Nariya H, Inouye S. A protein Ser/Thr kinase cascade negatively regulates the DNA-binding activity
of MrpC, a smaller form of which may be necessary for the *Myxococcus xanthus* development. Mol
Microbiol. 2006;60(5):1205-17.
35.   McLaughlin PT, Bhardwaj V, Feeley BE, Higgs PI. MrpC, a CRP/Fnr homolog, functions as a negative
autoregulator during the *Myxococcus xanthus* multicellular developmental program. Mol Microbiol.
2018;109:245-61.
36.   Higgs PI, Jagadeesan S, Mann P, Zusman DR. EspA, an orphan hybrid histidine protein kinase,
regulates the timing of expression of key developmental proteins of *Myxococcus xanthus*. J
Bacteriol. 2008;190:4416-26.
37.   Schramm A, Lee B, Higgs PI. Intra- and inter-protein phosphorylation between two hybrid histidine
kinases controls *Myxococcus xanthus* developmental progression. J Biol Chem. 2012;287:25060–72.
38.   Rajagopalan R, Kroos L. Nutrient-regulated proteolysis of MrpC halts expression of genes important
for commitment to sporulation during *Myxococcus xanthus* development. J Bacteriol.
2014;196:2736-47.
39.   Hoang Y, Kroos L. Ultrasensitive response of developing *Myxococcus xanthus* to the addition of
nutrient medium correlates with the level of MrpC. J Bacteriol. 2018;200:e00456-18.
40.   Feeley BE, Bhardwaj V, McLaughlin PT, Diggs S, Blaha GM, Higgs PI. An amino-terminal
threonine/serine motif is necessary for activity of the Crp/Fnr homolog, MrpC and for *Myxococcus*
xanthus developmental robustness. Mol Microbiol. 2019;112(5):1531-51.
41.   Robinson M, Son B, Kroos L. Transcription factor MrpC binds to promoter regions of many
developmentally-regulated genes in *Myxococcus xanthus*. BMC Genomics. 2014;15:1123.
42.   Campbell A, Viswanathan P, Barrett T, Son B, Saha S, Kroos L. Combinatorial regulation of the *dev*
operon by MrpC2 and FruA during *Myxococcus xanthus* development. J Bacteriol. 2015;197:240-51.
43.   Lee J, Son B, Viswanathan P, Luethy P, Kroos L. Combinatorial regulation of *fmgD* by MrpC2 and
FruA during *Myxococcus xanthus* development. J Bacteriol. 2011;193:1681-9.
44.   Mittal S, Kroos L. Combinatorial regulation by a novel arrangement of FruA and MrpC2
transcription factors during *Myxococcus xanthus* development. J Bacteriol. 2009;191:2753-63.
45.   Son B, Liu Y, Kroos L. Combinatorial regulation by MrpC2 and FruA involves three sites in the *fmgE*
promoter region during *Myxococcus xanthus* development. J Bacteriol. 2011;193:2756-66.
46.   Hoang Y, Franklin JL, Dufour YS, Kroos L. Cell density, alignment, and orientation correlate with C-
signal-dependent gene expression during *Myxococcus xanthus* development. Proc Natl Acad Sci
USA. 2021;118(45):e2111706118.
47.   Kroos L, Kuspa A, Kaiser D. A global analysis of developmentally regulated genes in *Myxococcus*
xanthus. Dev Biol. 1986;117:252-66.
48.   Kroos L, Kuspa A, Kaiser D. Defects in fruiting body development caused by Tn*5 lac* insertions in
Myxococcus xanthus. J Bacteriol. 1990;172:484-7.
49.   Boysen A, Ellehauge E, Julien B, Sogaard-Andersen L. The DevT protein stimulates synthesis of FruA,
a signal transduction protein required for fruiting body morphogenesis in *Myxococcus xanthus*. J
Bacteriol. 2002;184(6):1540-6.
50.   Thony-Meyer L, Kaiser D. *devRS*, an autoregulated and essential genetic locus for fruiting body
development in *Myxococcus xanthus*. J Bacteriol. 1993;175:7450-62.

51. Viswanathan P, Murphy K, Julien B, Garza AG, Kroos L. Regulation of *dev*, an operon that includes
genes essential for *Myxococcus xanthus* development and CRISPR-associated genes and repeats. J
Bacteriol. 2007;189:3738-50.
52. Rajagopalan R, Wielgoss S, Lippert G, Velicer GJ, Kroos L. *devI* is an evolutionarily young negative
regulator of *Myxococcus xanthus* development. J Bacteriol. 2015;197(7):1249-62.
53. Rajagopalan R, Kroos L. The *dev* operon regulates the timing of sporulation during *Myxococcus*
xanthus development. J Bacteriol. 2017;199(10):e00788-16.
54. Makarova KS, Aravind L, Wolf YI, Koonin EV. Unification of Cas protein families and a simple
scenario for the origin and evolution of CRISPR-Cas systems. Biol Direct. 2011;6:38.
55. Bullock HA, Shen H, Boynton TO, Shimkets LJ. Fatty acid oxidation is required for *Myxococcus*
xanthus development. J Bacteriol. 2018;200(10):e00572-17.
56. Kuzmich S, Blumenkamp P, Meier D, Szadkowski D, Goesmann A, Becker A, et al. CRP-Like
transcriptional regulator MrpC curbs c-di-GMP and 3',3'-cGAMP nucleotide levels during
development in *Myxococcus xanthus*. mBio. 2022;13:e00044-22.
57. Muller FD, Treuner-Lange A, Heider J, Huntley SM, Higgs PI. Global transcriptome analysis of spore
formation in *Myxococcus xanthus* reveals a locus necessary for cell differentiation. BMC Genomics.
2010;11:264.
58. Muller FD, Schink CW, Hoiczyk E, Cserti E, Higgs PI. Spore formation in *Myxococcus xanthus* is tied
to cytoskeleton functions and polysaccharide spore coat deposition. Mol Microbiol.
2012;83(3):486-505.
59. Holkenbrink C, Hoiczyk E, Kahnt J, Higgs PI. Synthesis and assembly of a novel glycan layer in
Myxococcus xanthus spores. J Biol Chem. 2014;289(46):32364-78.
60. Wartel M, Ducret A, Thutupalli S, Czerwinski F, Le Gall AV, Mauriello EM, et al. A versatile class of
cell surface directional motors gives rise to gliding motility and sporulation in *Myxococcus xanthus*.
PLoS Biol. 2013;11(12):e1001728.
61. Licking E, Gorski L, Kaiser D. A common step for changing cell shape in fruiting body and starvation-
independent sporulation of *Myxococcus xanthus*. J Bacteriol. 2000;182(12):3553-8.
62. Ueki T, Inouye S. Identification of a gene involved in polysaccharide export as a transcription target
of FruA, an essential factor for *Myxococcus xanthus* development. J Biol Chem.
2005;280(37):32279-84.
63. Giglio KM, Zhu C, Klunder C, Kummer S, Garza AG. The enhancer binding protein Nla6 regulates
developmental genes that are important for *Myxococcus xanthus* sporulation. J Bacteriol.
2015;197(7):1276-87.
64. Perez-Burgos M, Garcia-Romero I, Valvano MA, Sogaard Andersen L. Identification of the Wzx
flippase, Wzy polymerase and sugar-modifying enzymes for spore coat polysaccharide biosynthesis
in *Myxococcus xanthus*. Mol Microbiol. 2020;113(6):1189-208.
65. Kaiser D. Social gliding is correlated with the presence of pili in *Myxococcus xanthus*. Proc Natl Acad
Sci USA. 1979;76:5952-6.
66. Caberoy NB, Welch RD, Jakobsen JS, Slater SC, Garza AG. Global mutational analysis of NtrC-like
activators in *Myxococcus xanthus*: identifying activator mutants defective for motility and fruiting
body development. J Bacteriol. 2003;185(20):6083-94.
67. Davidson EH. Genomic Regulatory Systems. San Diego: Academic Press; 2001.
68. Rickels R, Shilatifard A. Enhancer logic and mechanics in development and disease. Trends Cell Biol.
2018;28(8):608-30.
69. Panigrahi A, O'Malley BW. Mechanisms of enhancer action: the known and the unknown. Genome
Biol. 2021;22(1):108.
70. Mohanraju P, Saha C, van Baarlen P, Louwen R, Staals RHJ, van der Oost J. Alternative functions of
CRISPR-Cas systems in the evolutionary arms race. Nat Rev Microbiol. 2022;20(6):351-64.

71. Koonin EV, Makarova KS. Evolutionary plasticity and functional versatility of CRISPR systems. PLoS
Biol. 2022;20(1):e3001481.
72. Munoz-Dorado J, Inouye S, Inouye M. A gene encoding a protein serine/threonine kinase is
required for normal development of *M. xanthus*, a Gram-negative bacterium. Cell. 1991;67:995-
1006.
73. Kroos L. Eukaryotic-like signaling and gene regulation in a prokaryote that undergoes multicellular
development. Proc Natl Acad Sci USA. 2005;102(8):2681-2.
74. Rajagopalan R, Sarwar Z, Garza AG, Kroos L. Developmental gene regulation. In: Yang Z, Higgs P,
editors. Myxobacteria: genomics, cellular and molecular biology. Norfolk, UK: Caister Academic
Press; 2014. p. 105-26.
75. Sun H, Shi W. Genetic studies of *mrp*, a locus essential for cellular aggregation and sporulation of
Myxococcus xanthus. J Bacteriol. 2001;183(16):4786-95.
76. Wireman JW. Developmental induction of *Myxococcus xanthus* myxospores. J Bacteriol.
1979;140:147-63.
77. Kroos L. Bacterial development in the fast lane. J Bacteriol. 2008;190:4373-6.
78. Bressuire-Isoard C, Broussolle V, Carlin F. Sporulation environment influences spore properties in
Bacillus: evidence and insights on underlying molecular and physiological mechanisms. FEMS
Microbiol Rev. 2018;42(5):614-26.
79. Isticato R, Lanzilli M, Petrillo C, Donadio G, Baccigalupi L, Ricca E. *Bacillus subtilis* builds structurally
and functionally different spores in response to the temperature of growth. Environ Microbiol.
2020;22(1):170-82.
80. Hamiot A, Lemy C, Krzewinski F, Faille C, Dubois T. Sporulation conditions influence the surface and
adhesion properties of *Bacillus subtilis* spores. Front Microbiol. 2023;14:1219581.
81. Eichenberger P, Fujita M, Jensen ST, Conlon EM, Rudner DZ, Wang ST, et al. The program of gene
transcription for a single differentiating cell type during sporulation in *Bacillus subtilis*. PLoS Biol.
2004;2(10):1664-83.
82. Kroos L. The *Bacillus* and *Myxococcus* developmental networks and their transcriptional regulators.
Ann Rev Genet. 2007;41:13-39.
83. Arrieta-Ortiz ML, Hafemeister C, Bate AR, Chu T, Greenfield A, Shuster B, et al. An experimentally
supported model of the *Bacillus subtilis* global transcriptional regulatory network. Mol Syst Biol.
2015;11(11):839.
84. McKenney PT, Driks A, Eichenberger P. The *Bacillus subtilis* endospore: assembly and functions of
the multilayered coat. Nat Rev Microbiol. 2013;11(1):33-44.
85. Shuster B, Khemmani M, Abe K, Huang X, Nakaya Y, Maryn N, et al. Contributions of crust proteins
to spore surface properties in *Bacillus subtilis*. Mol Microbiol. 2019;111(3):825-43.
86. Shuster B, Khemmani M, Nakaya Y, Holland G, Iwamoto K, Abe K, et al. Expansion of the spore
surface polysaccharide layer in *Bacillus subtilis* by deletion of genes encoding glycosyltransferases
and glucose modification enzymes. J Bacteriol. 2019;201(19):e00321-19.
87. La Fortezza M, Rendueles O, Keller H, Velicer GJ. Hidden paths to endless forms most wonderful:
ecology latently shapes evolution of multicellular development in predatory bacteria. Commun
Biol. 2022;5(1):977.
88. Kalathera J, Patel V, Pande S. Differential evolution of cooperative traits in aggregative multicellular
bacterium *Myxococcus xanthus* driven by varied population bottleneck sizes. bioRxiv. 2023;doi:
https://doi.org/10.1101/2023.09.20.558552.
89. Kashefi K, Hartzell P. Genetic supression and phenotypic masking of a *Myxococcus xanthux frzF*-
defect. Molec Microbiol. 1995;15:483-94.

90. Iniesta AA, Garcia-Heras F, Abellon-Ruiz J, Gallego-Garcia A, Elias-Arnanz M. Two systems for
conditional gene expression in *Myxococcus xanthus* inducible by isopropyl-β-D-
thiogalactopyranoside or vanillate. J Bacteriol. 2012;194(21):5875-85.

91. Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, 3rd, Smith HO. Enzymatic assembly of
DNA molecules up to several hundred kilobases. Nat Methods. 2009;6(5):343-5.

92. Sambrook J, Fritsch E, Maniatis T. Molecular Cloning: a laboratory manual, 2nd ed. Cold Spring
Harbor: Cold Spring Harbor Laboratory Press; 1989.

93. Ossa F, Diodati ME, Caberoy NB, Giglio KM, Edmonds M, Singer M, et al. The *Myxococcus xanthus*
Nla4 protein is important for expression of stringent response-associated genes, ppGpp
accumulation, and fruiting body development. J Bacteriol. 2007;189(23):8474-83.

94. Yoder-Himes D, Kroos L. Regulation of the *Myxococcus xanthus* C-signal-dependent Ω4400
promoter by the essential developmental protein FruA. J Bacteriol. 2006;188:5167-76.

95. Nariya H, Inouye S. Identification of a protein Ser/Thr kinase cascade that regulates essential
transcriptional activators in *Myxococcus xanthus* development. Mol Microbiol. 2005;58(2):367-79.

96. Bradford M. A rapid and sensitive method for the quantitation of microgram quantities of protein
utilizing the principle of protein-dye binding. Anal Biochem. 1976;72:248-54.

97. Sun H, Shi W. Analyses of *mrp* genes during *Myxococcus xanthus* development. J Bacteriol.
2001;183(23):6733-9.

98. Viswanathan P, Ueki T, Inouye S, Kroos L. Combinatorial regulation of genes essential for
Myxococcus xanthus development involves a response regulator and a LysR-type regulator. Proc
Natl Acad Sci USA. 2007;104:7969-74.

1098

1099
1100
Figure 1. Model of the gene regulatory network governing formation of *M. xanthus* fruiting
bodies. Starvation increases the level of phosphorylated Nla6 (Nla6-P), Nla28-P, MrpB-P, and
MrpC in a cascade of transcription factors (7, 32, 35). Nutrients (38, 39) and Esp signaling (36,
37) can negatively regulate MrpC *via* proteolysis, and MrpC negatively autoregulates by
competing with MrpB-P for binding to the *mrpC* promoter region (35). MrpC positively
regulates C-signaling (97) and *fruA* transcription (33), leading to production of unactivated FruA.
C-signaling activates FruA to FruA* by an unknown mechanism (28, 29). FruA* positively
regulates mound formation (30), which enhances C-signaling (18-20), forming a positive
feedback loop. FruA* also positively regulates the *dev* operon (98) together with MrpC (42),
thus forming a feed-forward loop that has been speculated to also increase transcription of
genes involved in cellular shape change (29). DevTRS proteins negatively autoregulate,
decreasing DevI, which negatively regulates cellular shape change (52, 53) and may inhibit
negative autoregulation by DevTRS (29). Here, we investigate possible regulation by Nla6-P,
MrpC, FruA or FruA*, and DevI (lines to gray box with **?**) of late-acting operons involved in spore
metabolism and spore coat biogenesis.

Figure 2. Development of *M. xanthus* strains. The WT strain and its indicated mutant derivatives were subjected to starvation under submerged culture condition and microscopic images were obtained at the indicated times PS. The WT strain and all the mutants except *nfsA-H* formed compact mounds by 18 h (an arrow points to one in each panel). The *nfsA-H* mutant formed compact mounds by 21 h. Mounds of the WT strain began to darken by 30 h. Mounds of the *nfsA-H*, *exoC*, and *exoL* mutants darkened by 30 h but remained slightly less dark than mounds of the WT strain and the *fadI* mutant by 48 h. Bar, 100 μm. Similar results were observed in at least three biological replicates.

1127
1128
Figure 3. Sonication-resistant spores during *M. xanthus* development. The WT strain and its
indicated mutant derivatives were subjected to starvation under submerged culture conditions.
Samples were collected at the indicated times PS for quantification of sonication-resistant
spores. Values are expressed as a percentage of the number of rod-shaped cells present at the
time starvation initiated development (T_0) (Table S1). Bars show the average of three biological
replicates and error bars indicate one standard deviation.
1135

1136
1137
Figure 4. Transcript levels in the WT strain and the *csgA* and *fruA* mutants late in *M. xanthus*
development. The WT strain and its mutant derivatives were subjected to starvation under
submerged culture conditions. Samples were collected at the indicated times PS for
measurement of the *exoA* (A), *exoL* (B), *nfsA* (C), and *fadI* (D) transcript levels by RT-qPCR.
Graphs show the data points and the average of at least three biological replicates, relative to
the WT strain at 18 h PS, and error bars indicate one standard deviation. Asterisks indicate a
difference ($p < 0.05$ in Student's two-tailed *t*-tests) from the WT strain at the corresponding
time PS.
1146
1147

1148
1149

1150
1151

E

Transcript	Half-life in the WT strain (min)	Half-life in the *csgA* mutant (min)	*p* value
exoA	12 ± 2	8 ± 3	*p* = 0.13
exoL	8 ± 3	10 ± 2	*p* = 0.20
nfsA	5 ± 2	7 ± 1	*p* = 0.26
fadI	7 ± 1	9 ± 2	*p* = 0.26

1152
1153

Figure 5. Transcript stability in the WT strain and the *csgA* mutant late in *M. xanthus*
development. The WT strain and the *csgA* mutant were subjected to starvation under
submerged culture conditions for 30 h. The overlay was replaced with fresh starvation buffer
containing rifampicin (50 μg/mL) and samples were collected immediately (t_0) and at the
indicated times (t_x) for measurement of the *exoA* (A), *exoL* (B), *nfsA* (C), and *fadI* (D) transcript
levels by RT-qPCR. Transcript levels at t_x were normalized to that at t_0 for each of three
biological replicates and used to determine the transcript half-life for each replicate. The graph
shows the average ln(t_x/t_0) and one standard deviation for the three biological replicates of the
WT strain (black dashed line) and the *csgA* mutant (gray solid line). The average half-life and
one standard deviation, as well as the *p* value from a Student's two-tailed *t*-test, are reported in
(E).
1165

1166

Figure 6. Transcript levels in the WT strain and the *mrpC* and *fruA* mutants late in *M. xanthus*

development. The WT strain and its mutant derivatives were subjected to starvation under

submerged culture conditions. Samples were collected at the indicated times PS for

measurement of *exoA* (A), *exoL* (B), *nfsA* (C), and *fadI* (D) transcript levels by RT-qPCR. Graphs

show the data points and the average of at least three biological replicates, relative to the WT

strain at 18 h PS, and error bars indicate one standard deviation. Asterisks above error bars

indicate a difference ($p < 0.05$ in Student's two-tailed *t*-tests) from the WT strain at the

corresponding time PS and asterisks above brackets indicate a difference between the mutants.

1175

1176
1177
Figure 7. Transcript levels in the WT strain, the *mrpC* mutant with inducible *fruA*, and the *mrpC*
and *fruA* mutants early in *M. xanthus* development. The WT strain and its mutant derivatives
were subjected to starvation under submerged culture conditions. Samples were collected at
the indicated times PS for measurement of *exoA* (A), *exoL* (B), *nfsA* (C), and *fadI* (D) transcript
levels by RT-qPCR. Expression of *fruA* fused to P_{van} was induced with vanillate (0.5 mM) during
growth and at 0 h PS. Graphs show the data points and the average of at least three biological
replicates, relative to the WT strain at 6 h PS, and error bars indicate one standard deviation.
Asterisks above error bars indicate a difference (*p* < 0.05 in Student's two-tailed *t*-tests) from
the WT strain at the corresponding time PS and asterisks above brackets indicate a difference
between the mutants.
1188

1189

1190

Figure 8. Transcript levels in *nla6* mutants early in *M. xanthus* development. The WT strain and

its mutant derivatives were subjected to starvation under submerged culture conditions.

Samples were collected at the indicated times PS for measurement of *exoA* (A), *exoL* (B), *nfsA*

(C), and *fadI* (D) transcript levels by RT-qPCR. Graphs show the data points and the average of

at least three biological replicates, relative to the WT strain at 6 h PS, and error bars indicate

one standard deviation. Asterisks indicate a difference ($p < 0.05$ in Student's two-tailed *t*-tests)

from the WT strain at the corresponding time PS and the asterisk above the bracket indicates a

difference between the mutants.

1199

1200
1201
Figure 9. Binding of the Nla6 DBD, MrpC, and FruA to the promoter regions of the late-acting operons. EMSAs were performed with ^{32}P-labeled DNA fragments (2 nM) of the *dev* (-114 to -19) (A), *fadIJ* (-214 to -5) (B), *exoA-I* (-171 to -1) (C), *exoL-P* (-217 to +56) (D), and *nfsA-H* (-166 to +35) (E) promoter regions, and MBP-Nla6 DBD, His$_6$-MrpC, and FruA-His$_6$ at the μM concentrations indicated by numbers above each autoradiographic image. A white number on a black background indicates a protein that was added to the DNA-binding reaction 10 min later than the other protein. The positions of migration of shifted complexes are indicated along the side of each image by arrowheads for binding of individual proteins and by arrows for binding of two proteins (see boxed key). Lanes are numbered below images.

1211

1212
1213
Figure 10. Transcript levels in *devI* and *devS* mutants during *M. xanthus* development. The WT
strain and its mutant derivatives were subjected to starvation under submerged culture
conditions. Samples were collected at the indicated times PS for measurement of *exoA* (A),
exoL (B), *nfsA* (C), and *fadI* (D) transcript levels by RT-qPCR. Graphs show the data points and
the average of at least three biological replicates, relative to the WT strain at 18 h PS, and error
bars indicate one standard deviation. Asterisks indicate a significant difference ($p < 0.05$ in
Student's two-tailed *t*-tests) from the WT strain at the corresponding time PS.
1221

1222
1223
Figure 11. Model for regulation of late-acting operons during *M. xanthus* development. See
the Figure 1 legend for explanation of the parts of the GRN that were known prior to this study.
The new findings of our study are shown in color and are explained in the text. Briefly, the
cartoon at left depicts a timeline of the developmental process with mounds forming by 18 h PS
and spore-filled fruiting bodies forming by 36 h. Green arrows indicate positive regulation and
red lines with a barred end indicate negative regulation. Solid arrows/lines indicate direct
regulation and dashed arrows/lines indicate indirect regulation. Cartoons below the operons
illustrate how independent, cooperative, and competitive binding of transcription factors to the
promoter regions (see boxed key) could explain negative regulation early in development and
positive regulation late in development. These cartoons are further explained in the Supporting
Information.