Grow now, pay later: when should a bacterium go into debt?

Jaime G. Lopez[1,2] **and Amir Erez** [2]✉

[1]Department of Bioengineering, Stanford University, Stanford 94305, United States;
[2]Racah Institute of Physics, The Hebrew University, Jerusalem 9190401, Israel

✉ **For correspondence:**
amir.erez1@mail.huji.ac.il

Data availability: All code and data are available at
github.com/AmirErez/MicrobialDebt

Competing interests: The authors declare no competing interests.

Abstract

Microbes grow in a wide variety of environments and must balance growth and stress resistance. Despite the prevalence of such trade-offs, understanding of their role in non-steady environments is limited. In this study, we introduce a mathematical model of "growth debt", where microbes grow rapidly initially, paying later with slower growth or heightened mortality. We first compare our model to a classical chemostat experiment, validating our proposed dynamics and quantifying *Escherichia coli*'s stress resistance dynamics. Extending the chemostat theory to include serial-dilution cultures, we derive phase diagrams for the persistence of "debtor" microbes. We find that debtors cannot coexist with non-debtors if "payment" is increased mortality but can coexist if it lowers enzyme affinity. Surprisingly, weak noise considerably extends the persistence of resistance elements, pertinent for antibiotic management. Our microbial debt theory, broadly applicable across many environments, bridges the gap between chemostat and serial dilution systems.

Introduction

Microorganisms have shaped the world we live in and have adapted to thrive almost anywhere, including the human body, desert soils, forest soils, and coral reefs (*1–4*). Competition for limited resources is a central characteristic of microbial life, and this fierce competition, ongoing since the first cell emerged billions of years ago, has led microbes to explore astonishing ways to gain a growth advantage. However, our understanding of the core principles behind microbial competition remains mostly limited to steady environments.

In the microbial world, organisms commonly make a trade-off between hedging against future adverse events and maximizing immediate growth (*5*). This trade-off manifests in a variety of ecological contexts, many of them relevant to human health. For example, carrying the genes necessary to resist antibiotics slows a microorganism's growth in the absence of antibiotic, but the organism's growth will be maintained if antibiotics enter the environment (*6–9*). Similarly, the malaria parasite *Plasmodium falciparum* incurs a substantial growth defect by carrying a mutation that protects from anti-malarial drugs (*10*). This trade-off is not limited to drug resistance: mutations in regulatory elements, such as *Escherichia coli*'s *rpoS*, allow organisms to trade-off nutritional competence and general stress resistance (*11*, *12*). To understand the core principle behind this "grow now, pay later" phenomenon, we introduce the metaphor of microbial debt—in which lacking the response capacity to future adverse events, reaps an immediate speedup in growth that is then "paid back" in the future. So, debtors are species that shift cost to the future (similar to interest paid on a loan) to gain a benefit in the present. Yet, despite the ubiquity of debt-like mechanisms in nature, theoretical understanding of debt in non-steady environments is missing.

Debt causes a shift of cost from the present to the future, and its effect is thereby strongly tied

to temporal variation. Temporal variation is a known driver of eco-evolution, impacting systems ranging from the microbiota of the Hadza people of East Africa ([13]), the mucosa of mice ([14]), and antibiotic resistance evolution ([15]). Evolutionarily, temporal variation can reduce the efficiency of natural selection and increase the fixation probability of mutations ([16]). Perhaps the simplest, yet most experimentally prevalent, example of temporal variation is serial-dilution batch culture - in which microbes are subject to periodic dilution and nutrient replenishment. This is substantially different from the chemostats — another common experimental system in which nutrient supply and dilution occur continuously ([17]). Studying and comparing the behaviors of chemostat and serial dilution systems, which represent two extremes of environmental variation, can provide insights into the environmental fluctuations present in real ecosystems. Although serial-dilution culturing is ubiquitous in experimental work ([18], [19]), modeling efforts in this direction are still incomplete ([20]–[23]). In the context of serial dilution cultures, we have recently shown that the effect of temporal variation can be understood through the "early-bird" effect ([20]). The early-bird effect arises from the temporal gap between the introduction of nutrients and the inoculating mixture of microbes (at $t = 0$, simultaneously); and the transfer of a fraction of the culture to inoculate the next batch (a dilution event, at $t = t_f$). During this time, the early-bird species exhibit rapid growth at the onset of the batch cycle. Although their per-capita growth diminishes later in the batch cycle, the significant size of the 'early-bird' species can still consume nutrients, consequently inhibiting the growth of 'late-bird' species. This effect can cause shifts in community structure, shifts that do not occur in an equivalent chemostat. The early-bird effect has recently been shown to control the assembly of gut microbiome-derived *ex vivo* communities ([24]). It suggests a possible advantage, unique to serial-dilution cultures: to pursue a 'debt' or 'grow now, pay later' strategy which accelerates early growth by shifting costs to the future.

Much theory has focused on the dynamics of debt trade-offs in fixed, chemostat-like conditions, showing that that debt trade-offs can enable coexistence of strains ([25], [26]). In some cases, predictions from these chemostat theories have been qualitatively verified experimentally ([27]). While these trade-offs have been explored extensively in steady environments, less is understood about their dynamics in fluctuating environments. Contemporary understanding of the ecology of antibiotic resistance suggests that variation can substantially alter the value of debt ([28]), highlighting the need for further work. Biophysical theories have been developed that predict optimal growth behaviors of microbes under environmental fluctuations, frameworks which can be applied to stress resistance phenotypes ([29], [30]). However, despite recent progress ([31]), there remains a pressing need for a comprehensive theoretical framework for trade-offs in serial dilution systems, bridging across commonly used experimental setups.

In this manuscript we ask: what are the consequences of accruing debt for a microbe within the context of a consumer-resource model? How does debt impact the growth dynamics of microbial communities under serial-dilution conditions compared to those in chemostat environments? We develop a modeling framework to explore microbial debt and demonstrate its use to analyze previously published experimental data. Our theory makes no assumptions about the shape of the debt trade-off surface ([9], [27], [32]). Though we consider only fixed and unchanging trade-offs, it is possible to extend the model to account for non-genetic adaptation ([33]). Our proposed model mimics a typical serial-dilution experimental procedure, while also being directly applicable to chemostats. We study the competition between two species, a debtor and a non-debtor, and find that when debt is compensated for by increased mortality, coexistence cannot occur between debtor and non-debtor, regardless of whether the mortality occurs in a punctuated or continuous manner. However, when debt is paid through decreased metabolic enzyme affinity, a stable coexistence state emerges. Intriguingly, when the cost of debt is subject to noise, the losing species can survive for a much longer time when compared to an equivalent deterministic system. Therefore, noise can enhance the persistence of antibiotic resistance elements.

Results

Chemostat dynamics reveal growth-debt regulation in *E. coli*

Before introducing our serial dilution model, we first present the equivalent chemostat model as a baseline for the serial dilution results and as a bridge to the existing, chemostat-dominated literature. A single nutrient is provided at flux S, and its concentration within the chemostat is denoted $c(t)$. The chemostat is well-mixed and constantly diluted with dilution rate δ. Let Species N be a non-debtor (a microbe with resistance to the stressor) with biomass density $\rho_N(t)$. Nutrient uptake is controlled by $g[c(t)]$, and growth is proportional to nutrient uptake. The per-capita growth rate of Species N is $E\,g[c(t)]$ where E defines the maximal per-capita growth rate. In contrast, Species D, the debtor species, with biomass density $\rho_D(t)$, has higher maximal growth rate, $E + \Delta E$ due to sacrificing its stress resistance capability and diverting these resources to immediate growth. The lack of stress resistance leads to constant stress-induced death at rate ω. The equations governing these chemostat dynamics are therefore: $\frac{1}{\rho_N}\frac{d\rho_N}{dt} = Eg[c] - \delta$; $\frac{1}{\rho_D}\frac{d\rho_D}{dt} = (E + \Delta_E)g[c] - \delta - \omega$; and $\frac{dc}{dt} = S - \rho_D(E + \Delta_E)g[c] - \rho_N Eg[c]$. Note that we neglect δc in the nutrient dynamics as the dominant nutrient depletion term comes from the bacterial consumption (*25*, *34*).

To validate our model assumptions and quantitatively show the existence of debt trade-offs, we apply this chemostat model to the classical experiments of Notley-McRobb et al. (*11*). These experiments studied competition between wild-type (WT) *Escherichia coli* and mutants with defective *rpoS* genes, the transcription factor governing stress response in *E. coli*. Here, the WT plays the role of non-debtor and the mutants are debtors. In *Figure 1*A, we show the relative abundance of the non-debtor WT strain in chemostats operating at different dilution rates. In these low-stress chemostats, the community is rapidly overtaken by the debtor *rpoS* mutants, well-captured by our model fit (see SI Appendix *section 4* for details). However, these debtors have traded growth rate against stress resistance, and are now far more vulnerable to stress. In *Figure 1*B we show the temperature stress response of the community from one chemostat as the debtor population increases. Initially, the population is composed of non-debtors and many bacteria survive the stressor. However, once debtors sweep the population almost no bacteria survive the stressor. These stress survival dynamics are well-described by our model fit. Thus, debt trade-offs occur within *E. coli* and are quantitatively consistent with our model assumptions

In this chemostat formalism, when is growth debt worthwhile? For this calculation, we define $g[c]$ as the classical Monod function $\frac{c}{c+K}$, where K is the half-saturation coefficient. By comparing the competitive ability of the non-debtor and debtor, we determine that a stressor death penalty ω is worthwhile if the corresponding benefit Δ_E satisfies:

$$\omega/\delta < \Delta_E/E, \tag{1}$$

(for details see SI Appendix *section 1*). Thus, at a higher dilution rate, δ, the relative cost of debt decreases. *Equation 1* predicts a linear relationship between inverse dilution rate $1/\delta$ and the growth advantage Δ_E/E, consistent with the debtor mutants observed by Notley-McRobb (*11*). In *Figure 1*C, we show the growth advantage of debtor mutants as a function of inverse dilution rate. The debtor mutant Δ_E/E is a reflection of *E. coli*'s investment in stress resistance. A large debtor advantage indicates that the non-debtor invests more resources in stress response, while a small advantage indicates little investment. The linear relationship shown in *Figure 1*C indicates *E. coli* invests less in stress resistance at higher growth rates in a manner consistent with our theory.

Similar chemostat theories to those presented here have been further developed. For example, it has been shown that with additional information about the form of the *rpoS* trade-off surface, one can qualitatively predict the outcome of *E. coli* evolution within a chemostat (*27*). However, what happens away from chemostat conditions in a serial-dilution, fluctuating environment?

Figure 1. Application of chemostat theory to experiments of Notley-McRobb *et al.* validates model assumptions and shows existence of trade-offs between maximal growth rate and stress resistance in *E. coli* (*11*). (*A*) Fit of chemostat theory to *rpoS* evolution trajectories in pH 7 glucose-limited chemostats at three different dilution rates δ. Fit parameters: $\Delta_E/E = [1.52 \pm 0.52, 0.37 \pm 0.1, 0.13 \pm 0.04]$ at $\delta = [0.1, 0.3, 0.6]$ h^{-1}, respectively and $x_D(t=0) = 0.025$. (*B*) Fit of theory to community temperature stress assay at 60°C. Samples taken from glucose-limited $\delta = 0.3$ h^{-1} chemostats at different days. Fit parameters: $\omega_N = 0.084 \pm 0.014$ min^{-1} and $\omega_D = 0.99 \pm 1.92$ min^{-1}. (*C*) Plot of debtor mutant advantage Δ_E/E as a function of inverse dilution rate $1/\delta$. Values from fit in *A*.

Serial dilution model formulation

We now establish our serial dilution model where two species compete for a single, limited nutrient in an environment influenced by a stressor. In a serial dilution culture, microbes are grown in a well-mixed environment for a given period of time, t_f, before being diluted into fresh media and allowed to grow again. We refer to the new nutrient added after dilution as the 'nutrient bolus'. A single type of nutrient is provided, and its concentration within a batch is denoted $c(t)$ with $c(t = 0) = c_0$. As in the chemostat, let Species N be a non-debtor (a microbe with resistance to the stressor) with biomass density $\rho_N(t)$, with per-capita growth rate $E\,g[c]$. In contrast, Species D, the debtor species, with biomass density $\rho_D(t)$, has higher maximal growth rate, $E + \Delta E$ due to sacrificing its stress resistance capability. The lack of stress resistance leads to two possible forms of stress-induced death: a continuous death rate per unit time (ω), and/or, an increased dilution during dilution events (Ω). The within-batch dynamics are therefore:

$$
\begin{aligned}
\frac{dc}{dt} &= -g[c]\left(E\,\rho_N + (E + \Delta_E)\,\rho_D\right), \\
\frac{1}{\rho_N}\frac{d\rho_N}{dt} &= E\,g[c] - \omega_N, \\
\frac{1}{\rho_D}\frac{d\rho_D}{dt} &= (E + \Delta_E)\,g[c] - \omega_D.
\end{aligned}
\tag{2}
$$

Here, ω_N and ω_D are the non-debtor and debtor within-batch death rates, respectively. These death rates could be the result of an antibiotic in the media or another stressor, such as temperature, salinity, or pH. Note that the terms on the right hand side of the equations for $\rho_{N/D}$ are the per-capita growth rates. It is possible to render these equations dimensionless by measuring time in units of E, and concentration in units of K (cf. SI Appendix *section 3*). However, since our model is easily applicable to experimental data, we keep the dimensionful version of these dynamics for more direct data comparison.

The second source of mortality in the model is not continuous, but rather, punctuated, happening only at the transition between batches. We refer to the transition from batch to batch as 'dilution', reflecting what occurs in laboratory serial dilution, but it represents any number of potential punctuated stressors, such as sudden heat or osmotic shock. When dilution happens, each species is diluted by its species-specific factor Ω_N or Ω_D. Thereby, dilution in our model generalizes the typical scenario where a small volume of a well-mixed culture is used to inoculate the next batch, diluting each species equally. In this generalized version, the total inoculum size, $\rho_N(0) + \rho_D(0)$, can vary from batch to batch until steady state is reached. A schematic of these mechanisms is shown in *Figure 2*A. At the end time (t_f) of batch b, each species gets diluted differently, and so, at the beginning of batch $b + 1$,

$$
\begin{aligned}
\rho_N^{(b+1)}(t = 0) &= e^{-\Omega_N t_f}\left(\frac{\rho_0}{\rho_0 + c_0}\right)\rho_N^{(b)}(t_f), \\
\rho_D^{(b+1)}(t = 0) &= e^{-\Omega_D t_f}\left(\frac{\rho_0}{\rho_0 + c_0}\right)\rho_D^{(b)}(t_f).
\end{aligned}
\tag{3}
$$

In the rest of this manuscript, we set $\Omega_N = \omega_N = 0$ and drop the subscript from Ω_D and ω_D.

Nutrient utilization, $g[c(t)]$. We considered two general forms of the utilization function $g[c]$: utilization that is proportional to the nutrient concentration, $c(t)$, and utilization that obeys Hill-like non-linearity. Specifically,

$$
g[c] =
\begin{cases}
\frac{c^h}{c^h + K^h} & \text{Hill, } h \\
\frac{c}{K} & \text{Linear}
\end{cases}
\tag{4}
$$

The $h = 1$ case is known as the Monod function (*17*), and its $c \ll K$ limit corresponds to linear consumption. At the opposite limit, at high nutrient concentration, linear consumption is nonphysical

Figure 2. Overview of serial dilution ecosystem with debt. (*A*) Schematic of the serial dilution system with debt. In the continuous death case, the debtor is able to initially grow more quickly, but loses biomass continuously. In the punctuated death case, the debtor grows more quickly but is diluted more severely at the end of the batch. (*B*) Example of growth functions. Linear is $g[c] = Ec$, Monod is $g[c] = E\left(\frac{c}{c+K}\right)$, and Hill-2 is $g[c] = E\left(\frac{c^2}{c^2+K^2}\right)$. (*C*) Time necessary for a single non-debtor species to deplete 99% of the supplied nutrient t^* for different growth functions. In order to choose a single t_f that is sufficiently long, we select the t^* for $c_0/K = 10^4$ with a Monod growth function.

as it implies that a huge quantity of nutrient would be consumed almost instantly. Additionally, if $c \ll K$ and $h > 1$, at low nutrient levels, nutrient consumption effectively grinds to a halt. The Monod case, $h = 1$, extends between these two behaviors and is well-established to model microbial nutrient utilization (*Figure 2*B). For the rest of this manuscript, we thus focus on the Monod form.

Batch time, t_f. When modelling serial dilution, it is possible to choose the batch time t_f to be as long as necessary to consume all resources (*20*, *21*). However, here we also considered within-batch mortality, and therefore arbitrarily long batches would lead to the extinction of the susceptible species. A more realistic view, in line with experimental practices, sets a fixed batch time, t_f. The time needed for consumption of most of the nutrient depends on how much nutrient is provided. In this study, we varied the supplied nutrient concentration, c_0, over six orders of magnitude. To ensure that most of the nutrient is consumed in all relevant scenarios, we define t_f based on the nutrient utilization at high c_0. Assuming Monod utilization with only one species present, t_f is defined as the time it takes to consume 99% of c_0 with $c_0/K = 10^4$. Thereby, we ensure that at least 99% of the nutrient will be consumed for all relevant scenarios explored in this work. The dependence of t_f on c_0, while maintaining $\frac{c_0 - c(t_f)}{c_0} = 0.99$, is shown in *Figure 2*C.

Figure 3. Serial dilution results in (*A*) Phase diagram of competition between debtor and non-debtor as a function of bolus size c_0/K and normalized death penalty. The dashed line represents the phase boundary for the equivalent chemostat system, diverging substantially from the serial dilution phase boundary at high c_0/K. This phase diagram captures both continuous and punctuated death, with the y-axis being equivalently Ω/Δ_E or ω/Δ_E (*B*) Within-batch dynamics of a single species along the Δ_E, ω trade-off. Note that while the debtor species grows rapidly initially, its final biomass may be much lower due to death. *Inset*: (same axes). In competition with a non-debtor, the increased early growth of the debtor consumes the nutrient and denies nutrient from the non-debtor.

Equivalent value of debt represented as continuous and punctuated mortality

In our model, when is microbial debt worthwhile? Does it matter if the debt incurred is paid for in a continuous or punctuated fashion? We quantify debt by considering the dynamics within a single batch, i.e., a single growth and dilution cycle. At a given batch b, it is possible to integrate Eq. 2, given the initial inocula concentrations $\rho_{1,2}^{(b)}(0)$. At the end of this batch, $t = t_f$, we have the following populations:

$$
\begin{aligned}
\rho_N^{(b)}(t_f) &= \rho_N^{(b)}(0)e^{E \int_0^{t_f} g[c(t')]\,dt'}, \\
\rho_D^{(b)}(t_f) &= \rho_D^{(b)}(0)e^{E(1+\frac{\Delta_E}{E}) \int_0^{t_f} g[c(t')]\,dt' - \omega t_f}.
\end{aligned}
\tag{5}
$$

When going to the next batch, the debtor species gets diluted by a multiplicative factor $e^{-\Omega t_f}$ beyond the $\frac{\rho_0}{\rho_0+c_0}$ factor both species get diluted by. Therefore, at the beginning of batch $b+1$,

$$
\begin{aligned}
\rho_N^{(b+1)}(0) &= \frac{\rho_0}{\rho_0 + c_0}\,\rho_N^{(b)}(t_f), \\
\rho_D^{(b+1)}(0) &= e^{-\Omega t_f}\frac{\rho_0}{\rho_0 + c_0}\,\rho_D^{(b)}(t_f).
\end{aligned}
\tag{6}
$$

At steady state, equal growth implies that both species are increased and decreased by the same fold-change. Therefore, coexistence implies that,

$$
\underbrace{\left(\frac{1}{t_f}\int_0^{t_f} g[c(t')]\,dt'\right)}_{\text{Time-averaged nutrient utilization}} \underbrace{\Delta_E}_{\text{Benefit per utilization}} = \underbrace{\omega + \Omega}_{\text{Cost}}.
\tag{7}
$$

Equation 7 states that coexistence balances growth cost with benefit: the growth benefit per nutrient utilized times the time-averaged nutrient utilization equals the cost. In essence, the benefit from accruing the debt in the form of increased growth must be equal to the death cost incurred. This expression defines the border between regimes in which debt is favorable and those in which it is unfavorable (*Figure 3*A). In this choice of parameterization, the two forms of debt, continuous (ω) and punctuated (Ω), are equivalent and their rates are additive.

Importantly, in the single species case, even away from steady state, the nutrient utilization integral is independent of $g[c]$ if the vast majority of the nutrients are consumed within the batch, with $\int_0^{t_f} g[c(t)]\,dt \approx \frac{1}{E}\log\left(1 + \frac{c_0}{\rho_0}\right)$ by mass balance. Note that $1 + \frac{c_0}{\rho_0}$ is the dilution factor of the serial dilution system, a property determined by the experimentalist. This is a tremendous simplification, since knowledge of $g[c]$ is unnecessary. It is especially useful to parameterize experimental mono-culture dynamics, without having to measure $g[c]$. For example, application to an antibiotic resistance measurement requires only the element's carriage cost and the death rate of cells without the resistance elements.

Serial dilution cultures show diminishing return on debt at high nutrient concentrations when compared to a chemostat

We have determined the phase diagram for debt favorability in a serial dilution system, but how does this compare to a system with continuously supplied nutrient, i.e., a chemostat? If we add a very small bolus, $c_0 \ll K$, then dilute the culture by a fraction close to 1, and repeat, it stands to reason that the limiting behavior becomes continuous in time. As $c_0/K \to 0$, the serial dilutions become effectively a chemostat, and so, we expect that the continuous and punctuated dilution cases have the same chemostat limit. To consider $c_0 \ll K$, the chemostat limit, we follow ([20]), and linearize the dynamics around the small parameter c_0/K (for details see the *SI Appendix section 4*). The chemostat cost-benefit tradeoff then becomes $\frac{\Omega + \omega}{\Delta_E} \approx \frac{c_0}{\rho_0 E t_f}$, shown as the dashed line in *Figure 3*A. It is satisfying to note that this is equivalent to the linear chemostat debt relation in Equation 1 and thus also consistent with the experimental analysis shown in *Figure 1*C.

Why does the boundary between debtor and non-debtor dominance in the serial-dilution culture deviate from its equivalent chemostat at large c_0? There is a diminishing return on debt as more nutrient is supplied. The diminishing return and its relation to the early-bird effect can be understood from *Figure 3*B, showing time-series of a debtor in a single batch, grown in mono-culture, with several values of (ω, Δ_E) pairs. The pairs lie along the coexistence boundary with the non-debtor (*Figure 3*A). As ω, Δ_E are increased, the species grow more rapidly initially, (higher slope in the semi-log plot), then followed by an exponential loss because of ω (downward slope). The initial debt, which gives the rapid rise, is offset by the incurred death later in the batch. Indeed, when the non-debtor species is introduced in a co-culture, both species reach the same level at the end of the batch (inset of *Figure 3*B). This demonstrates the early-bird effect since the debtor deprives the non-debtor from nutrient in the early growth phase, though finally the debtor settles at a much lower population size because of the debt payoff.

Debts involving a growth affinity trade-off allow coexistence

Thus far, we have examined trade-offs where the maximum growth rate under normal conditions is counterbalanced by mortality in the presence of a microbicidal stressor. However, there are other forms of growth stress and payment, e.g., there exist microbiostatic stressors that do not kill but rather inhibit the growth of microbes. The inhibition can be considered as a penalty on the maximum growth rate. Similarly, resistance mechanisms can affect growth properties other than the maximum growth rate. For example, resistance mutations may lead to a growth defect that manifests most strongly at low nutrient concentrations. Interestingly, the resistant phenotype (with higher maximal growth rate and lower enzyme affinity) is now playing the role of the debtor in a bacteriostatic environment, because it "grows now, pays later". The debtor in the bacteriostatic environment mutates its enzymes to avoid the bacteriostatic effect, thereby sacrificing its future growth at low nutrient concentration because the mutated enzymes have lower affinity.

In our framework, growth penalties that are most severe in low nutrient conditions would manifest in the Monod affinity, K, increasing to $K + \Delta_K$, with a possible biophysical mechanism being mutations in proteins that reduce antibiotic binding but also reduce binding affinity for the natural substrate. ([28], [35]). Therefore, we shift our perspective and study the debt trade-off in which the

debtor species has mutated so that it pays a growth cost in low-nutrient environment, but gains a maximal growth rate ΔE. How does this trade-off impact the value of debt?

We deduce the affinity-growth trade-off using an invasibility calculation: when would a minuscule amount of an 'invader' species succeed in expanding under nutrient dynamics dictated by a single dominant non-debtor? Namely, a species with lower enzyme affinity, $K + \Delta_K$, but also higher maximal growth rate, $E + \Delta_E$, is introduced to a system dominated by the single-species dynamics of an unperturbed (K, E) species. How large does Δ_E need to be to offset the cost of the decreased affinity $K + \Delta_K$? Equivalent growth implies that $E \int_0^{t_f} \frac{c}{c+K} dt' = E \ln\left(1 + \frac{c_0}{\rho_0}\right) = \left(E + \Delta_E\right) \int_0^{t_f} \frac{c}{c+K+\Delta_K} dt'$. Expanding the perturbed Monod function at small Δ_K / K, with $f(K + \Delta_K) = f(K) + \frac{df}{dK}\big|_K \Delta_K + O(\Delta_K^2)$, we write to leading order,

$$\underbrace{\int_0^{t_f} \frac{c}{c + K + \Delta_K} dt'}_{I(t_f)} = \underbrace{\int_0^{t_f} \frac{c}{c + K} dt'}_{I(t_f)} - \underbrace{\Delta_K \int_0^{t_f} \frac{c}{(c + K)^2} dt'}_{\Delta_I(t_f)}, \tag{8}$$

$$\frac{\Delta_E}{E} = \frac{\Delta_I}{I} = \Delta_K \frac{\int_0^{t_f} \frac{c}{(c+K)^2} dt'}{\int_0^{t_f} \frac{c}{c+K} dt'}.$$

At equilibrium, the relative gain in maximal growth rate, Δ_E / E, is offset by the relative loss in utilization, Δ_I / I. It is possible to approximate this relation at the low and high c_0 limits, (see *SI Appendix section 5*). The integral Δ_I and its approximate forms are shown in **Figure 4**A. To leading order, we have,

$$\frac{\Delta_E}{E} \approx \begin{cases} \frac{\Delta_K}{K} & c_0 \ll K \\ \frac{\Delta_K}{c_0}\left(1 + \frac{\ln(c_0/K)}{\ln(c_0/\rho_0)}\right) & c_0 \gg K \end{cases} \tag{9}$$

The relative cost at equilibrium behaves differently in the low c_0 (chemostat) and high c_0 limits. At low c_0, as in a chemostat, the cost scales with Δ_K. Therefore, for a given value of relative gain, Δ_E / E there is a fixed cost Δ_K / K to maintain equilibrium, no matter how much nutrient is provided. There is little enough nutrient provided such that the growth function is linear in the nutrient amount, leaving the difference in affinity, Δ_K, as the dominant term. A different picture emerges in the high c_0 limit, where nutrient is initially saturating. Here, for a fixed gain, $\Delta_E / E =$ const, the cost $\Delta_K \sim c_0$, meaning that to maintain equilibrium, if more nutrient is provided, a larger enzyme affinity penalty can be supported. Said differently, the debtor takes over for a fixed Δ_K if c_0 increases. The debtor advantage over the chemostat limit is shown in the horizontal tendency of the Δ_K / c_0 boundary in **Figure 4**B. This is because 'payment' happens mostly when $c \approx K + \Delta_K$ and when $c_0 \gg K$ and Δ_K, a larger c_0 gives the debtor an increasing span of growth with effectively no payment.

To complete the invasibility test, one must consider the opposite situation, where the debtor dominates the culture, dictating the nutrient dynamics with $(K + \Delta_K, E + \Delta_E)$, and an invasion attempt by the non-debtor. When can the non-debtor invade? A similar derivation gives $\frac{\Delta_E}{E + \Delta_E} = \frac{\Delta_{I[K + \Delta_K]}}{I_{[K + \Delta_K]}}$, with the I and Δ_I integrals evolved under the debtor monoculture. As with the previous boundary, we may approximate, (*SI Appendix section 5*). Interestingly, a coexistence region emerges at high c_0 / K (**Figure 4**B). Coexistence is due to the different effect of the dominant species: Δ_E is measured against E when the non-debtor is dominant but against $E + \Delta_E$ when the debtor is dominant. This mode of coexistence is similar to that seen in another recent serial dilution theory (*36*).

Stochastic stressor dynamics significantly lengthen lifetime of ecologically unstable species

The models explored in the prior sections exhibit temporal variability, but in a deterministic manner: dilution factors and stressor levels staying the same in all batch cycles. However, in real ecosystems the magnitude of the stressor effect is variable. For example, antibiotic concentration fluctu-

Figure 4. The tradeoff between maximal growth rate, $E + \Delta_E$ and reduced affinity due to $K + \Delta_K$. (A) Δ_I/Δ_K depicts the loss of growth per biomass, and its low and high c_0/K approximations. (B) Phase diagram of the coexistence and dominance regimes, and its leading order analytic approximation (Eq. 9, white dashes). Black dashes: the chemostat limit of the phase boundary. $E = \Delta_E = 1$ and $K = 1$. A coexistence region where both species support a nonzero population emerges high c_0.

Figure 5. The effect of noise in Ω. (A) Full curves—mean ρ_N fraction. Shaded area: standard error of the mean, averaged over 1000 stochastic trajectories. Green curve—deterministic convergence to non-debtor take-over. (B) Time-series of the distribution of $\log_{10} \rho_N/\rho_D$, colors—distribution timeseries calculated over 50,000 batches, from the first batch until 4×10^5 batches. The widening corresponds to a diffusion process and the drift the extinction of ρ_D. Inset: the constant drift in the median value of ρ_N/ρ_D, identical for all three noise levels, equivalent to the mode of the histograms.

ates in the environment (*15*), leaving a debtor species disproportionately susceptible to extreme dilution. What is the effect of such stochastic noise on the debt cost-benefit trade-off? Let Ω fluctuate, drawing Ω in each batch from a uniform distribution, $\Omega \in [\Omega - \Delta_\Omega, \Omega + \Delta_\Omega]$. This corresponds to 'pink' or $1/\Omega$ noise where extreme events occur at very low probability.

We consider a system with $\Omega > \Omega^*$ such that the debtor is outcompeted in the deterministic version of the model. We show this deterministic trajectory, batch to batch, plotting the relative abundance of the non-debtor (*Figure 5*A). To explore how this extinction process is affected by noise, we plot the stochastic trajectory averages of the equivalent system with two different levels of Ω noise. Intriguingly, the introduction of even a small level of noise, constituting 10% of Ω, significantly decelerates the extinction dynamics. With 10% noise, $\Omega \pm 10\%$, the time to extinction is approximately tripled, while with 50% noise, the time to extinction is lengthened by orders of magnitude (*Figure 5*A). This behavior is symmetric to a reversal in roles: the same behavior occurs with non-debtor extinction if the environment favors debtors. Thus, noise in stressor severity can drastically extend the lifetime of ecologically unstable microbes.

How does this seemingly minor noise in Ω produce such drastic departures from the deterministic behavior? The key observation is that Ωt_f (and ωt_f) act on the population abundances

in logarithmic space (*Equation 3* and *Equation 5*). Thus, rather than inducing a random walk of ρ_N in linear space (which would preserve the deterministic dynamics of the mean), this process is instead a random walk in $\log(\rho_N/\rho_D)$. In *Figure 5*B we plot the evolution of the distribution of the *balance* variable, $\log(\rho_N/\rho_D)$ (*37*). As can be seen, the balance undergoes a canonical biased random walk, with the distribution widening and simultaneously its mean drifting. The drift is due to the induced extinction because unfavorable debt conditions, $\Omega > \Omega^*$, were chosen. As expected, the mean $\log(\rho_N/\rho_D)$ of the different noise levels, including the deterministic dynamics, are all equal (*Figure 5*B inset). As is known, the balance or log-ratio is the appropriate degree of freedom to consider when studying the dynamics of a composition (*38*, *39*). Indeed, balances would be the appropriate variables to consider in multi-species generalization of our framework. This random walk in logarithmic space has long tails for the relative abundances, $\frac{\rho_N}{\rho_N+\rho_D}$, a quantity typically relevant ecologically and measured in experiments. Thus, large, rare events significantly influence an observed mean fraction even when noise is relatively small. This suggests a broad distribution of stress resistance traits even within a single species, consistent with what has been observed in *E. coli* (*12*).

Conclusions and Discussion

Real-world microbes can incur debt by increasing immediate growth while neglecting protection from future events. Here, we proposed a theory for the impact of such debt trade-offs on microbial competition. We began by formulating chemostat debt dynamics and validating them with data from a classical chemostat experiment, thereby exposing the dynamic debt regulation behavior of *E. coli*. Notably, we validated the predicted linear relationship between the inverse dilution rate and the debtor advantage proposed by our chemostat theory. Having established the basic debt dynamics, we extended our theory to serial-dilution cultures. We found that growth debt where the 'payment' occurs continuously and where it occurs at the end of the batch are equivalent and can be unified in a single framework.

We constructed a phase diagram that delineates the boundary where debt becomes competitively favorable, as a function of nutrient amount in the system. This phase diagram encompasses both serial dilution and chemostat dynamics, unifying the two experimental systems. In the chemostat, the trade-off line is proportional to the total nutrient supply. Conversely, in the serial dilution setup, the addition of nutrients in a saturating concentration results in a diminishing return on debt. The effect of diminished return is considerable. For instance, a non-debtor that would typically go extinct in a high-nutrient chemostat may dominate an equivalent serial dilution culture, completely reversing the competition outcome.

We found that not all forms of debt are the same: debt associated with enzyme affinity allows the coexistence of two species on a single resource, violating the competitive exclusion principle by forming temporal niches (*36*). Mutating metabolic enzymes to withstand environmental stressors allows the debtor to boost its maximum growth rate, an advantageous strategy when nutrients are abundant, at the cost of growth when nutrient levels fall below $K + \Delta K$. Here, the role of the debtor is reversed compared to the case studied first: the debtor is now resistant to a bacteriostatic agent, while the non-debtor species is susceptible.

Stochastic stressor dynamics, a realistic extension of the model, revealed unexpected behavior. Even small fluctuations about the mean stress can significantly delay the extinction of a species, due to the compositional nature of the serial-dilution process. To decouple the compositional effects a log-ratio is used (*37*–*39*), here, $\log \frac{\rho_N}{\rho_D}$. As the system relaxes to steady state, the log-ratio, batch to batch, is the same in the deterministic and stochastic systems. Conversely, the relative fraction, $\frac{\rho_N}{\rho_N+\rho_D}$, survival time is greatly extended with the level of noise.

The outcomes of our stochastic dynamics study hold considerable implications for the strategic management and control of antibiotic resistance elements within environmental contexts. Rotating antibiotics to reduce the prevalence of resistance elements has been proposed (*40*). Our find-

ings indicate that the timescale for the depletion of resistance elements in the environment may be significantly longer than deterministic models (*41*) would estimate— when based solely on the fitness cost of the resistance element due to sub-inhibitory antibiotic concentration fluctuations. The variability induces large and rare events that significantly increase the relative fraction of the resistant population. Our findings extend beyond antibiotic resistance and imply that stress resistance genes could persist within communities for extended periods despite a net loss of fitness, highlighting their potential longevity. Noisy debt dynamics may broadly influence the prevalence of accessory genes in bacteria, which may be compounded by the noise in the expression of stress resistance elements (*42*).

Chemical stressors may change during the batch, e.g., antibiotics can be produced by other microbes, resulting in nontrivial stressor dynamics (for a chemostat analysis of these dynamics, see (*25*)). This may lead to large departures from our predicted behaviors: for example, such antisymmetric microbial interactions have been found in theory to lead to rock-paper-scissor dynamics (*43*). Furthermore, we have not considered the case where the dead bacteria contribute nutrient to the environment. Allowing such recycling may weaken the debtor advantage since instead of depriving the competitor of nutrient the debtor would then simply store and then release the nutrient. In summary, while this manuscript covers the impact of exogenous stressors, to elucidate the impact of endogenous stressors on debt will require further work.

Our framework predicts that the competitive value of debt in serial dilution ecosystems will decrease as nutrient concentrations increase. Such experiments involve measuring the cost of the resistance element relative to baseline growth, and the death rate by the stressor. In the chemostat analysis (*Figure 1*), we demonstrated how to deduce both values using existing experimental techniques, even from data published decades ago. We assumed first-order stressor kinetics, but our predictions are generalizable. A potential experiment would monitor the persistence of an antibiotic resistance plasmid in a community across a range of antibiotic and nutrient concentrations to construct an empirical version of the phase diagram in *Figure 3*A. The debtor could be a wild-type *E. coli* with the non-debtor being that same strain with an antibiotic resistance plasmid. Ideally this plasmid would confer resistance that does not inactivate the antibiotic (e.g. an efflux transporter), to prevent benefit to non-resistant organisms (*44*). This experiment could be performed at high-throughput in multiwell plates, with perpendicular gradients of antibiotic concentrations and nutrient levels, measuring the plasmid abundance using qPCR. One could measure death kinetics as a function of antibiotic concentration, and in the absence of antibiotic use a competition experiment to infer the cost of the plasmid. The resulting phase diagram should feature a boundary defined by *Equation 7*, increasing linearly with c_0 at low values of c_0, then plateauing at high c_0. Evolution may frustrate such an experiment—discussed below.

The influence of stressor noise on the lifespan of resistance elements can be experimentally tested. Here, the model predicts that across many independent populations of cells, populations subject to a more noisy stressor are expected to have longer mean persistence of resistance elements (*Figure 5*). One approach is to first identify a constant stressor level at which the resistance element goes extinct over multiple serial passages. For an antibiotic resistance element, this would be a sub-MIC level of antibiotic that impedes growth, but less than the cost of a resistance element. With baseline growth in a constant stressor environment established, run high-throughput batch cultures with each well receiving a noisy amount of antibiotic each passage, and control wells receiving the same mean amount. Resistance is measured as a population mean, as in Notley-McRobb *et al.* (*11*). The model predicts that the mean level of resistance in the noisy communities will exceed that in the constant-stressor controls. Even in the presence of additional mechanisms evolving over time, the impact of noise on overall resistance levels should still be observable through community-wide assays. The distribution of stress resistance levels observed in this experiment could also be compared to those found in nature (*12*).

Focusing on the fundamental mechanisms underlying microbial debt, we considered only two species and a single nutrient. Yet, natural microbial communities typically feature many coexisting

species. The mortality (ω, Ω) debt trade-off is not capable of supporting two species, so it would not support multiple species coexistence. Conversely, the affinity debt trade-off may support more than two species coexisting but may require fine-tuning (36). Investigating debt dynamics within complex, randomly assembled ecosystems (45) may shed light on the functioning of multi-species communities, such as the gut microbiome. Natural microbial communities also tend to feature more than the single resource assumed in our model. Previously, we demonstrated the 'early-bird' effect, where the presence of multiple resources can amplify the benefit of an early growth advantage within a batch (20). This 'early-bird' microbe can leverage its larger population size later in the batch to deplete resources it may not specialize in, depriving competitors. Thus, we expect debt dynamics to influence ecosystems with multiple species and resources in nontrivial ways.

Evolution is an important factor to consider, as the emergence of new variants of debtor and non-debtor occurs by mutation and subsequent competition with their ancestors. A promising direction to analyze the evolution of debt strategies is by using adaptive dynamics (46). Evolution may lead to the dominance of a single strain in the chemostat context, as it tends to minimize the steady-state nutrient level— the phenomenon of 'pessimization' (46). Pessimization is consistent with the results of Notley-McRobb *et al.* (11), where the debtor dominates at pH 7. At other pH values, a third strain emerged, with partial *rpoS* mutants that in our framework can be less extreme debtors. These chemostat experiments did not reach steady state, so we cannot conclude if coexistence occurred between the three strains. Later work from the same authors demonstrated that it is possible to qualitatively predict the outcomes of such evolution experiments using a chemostat framework (27). In a serial dilution context, it unclear whether the chemostat 'pessimization' intuition holds. A serial dilution experiment by Tenaillon *et al.* (47) featured a large number of replicate *E. coli* communities evolving in parallel. While Notley-McRobb *et al.* found that mutations in *rpoS* were ubiquitous, Tenaillon *et al.* found that *rpoS* mutations were rare, constituting only three of 1131 identified mutations (all stress resistance genes accounted for only 27 mutations in total). This result is consistent with our theory, as we predict that debtors (i.e., *rpoS* mutants) should be less competitive in serial dilution conditions. However, it is difficult to compare these two experiments, as the switch from serial dilution to chemostat also likely introduces additional stressors: *E. coli* tends to alter the pH of its environment, exposing the community to stress at the end of the batch. Such pH stress effectively increases the cost of debt beyond what is currently in our model.

Through the ages, debt has been an ever-present part of human life (48). Yet the omnipresence of debt in human life mirrors a much older and more basic notion. When competition for resources is fierce and resource levels change with time, a "grow now, pay later" strategy can be exploited to gain an advantage over one's competitors, a manifestation of the "early-bird" effect (20). Such an early-bird strategy is built on depriving competitors of resources by growing quickly, even if it means paying a significant future cost. Our work has shed light on how such fundamental trade-offs influence the world.

Acknowledgments

This work was partially funded by AE's startup grant at the Hebrew University. Both JGL and AE conceived the idea, carrier out calculations, and wrote the manuscript. We thank Nathalie Balaban, Po-Yi Ho, and Anthony Lyndon Shiver for critically reading this manuscript.
The authors declare that they have no competing interests.
All code and data are available at github.com/AmirErez/MicrobialDebt.

References

1. J. Lloyd-Price, G. Abu-Ali, C. Huttenhower, *Genome Medicine* **8**, 51, ISSN: 1756-994X, (https://doi.org/10.1186/s13073-016-0307-y) (2016).

2. A. Crits-Christoph *et al.*, *Microbiome* **1**, 28, ISSN: 2049-2618 (2013).

3. S. Lladó, R. López-Mondéjar, P. Baldrian, *Microbiology and Molecular Biology Reviews* **81**, e00063–16 (2017).

4. L. W. Kelly, A. F. Haas, C. E. Nelson, *mSystems* **3**, e00162–17, eprint: https://journals.asm.org/doi/pdf/10.1128/mSystems.00162-17, (https://journals.asm.org/doi/abs/10.1128/mSystems.00162-17) (2018).

5. F. Abram, T. Arcari, D. Guerreiro, C. P. O'Byrne, in ed. by R. K. Poole, D. J. Kelly (Academic Press, 2021), vol. 79, pp. 133–162, (https://www.sciencedirect.com/science/article/pii/S006529112100014X).

6. M. Linkevicius, J. M. Anderssen, L. Sandegren, D. I. Andersson, *Journal of Antimicrobial Chemotherapy* **71**, 1307–1313 (2016).

7. V. Patel, N. Matange, *eLife* **10**, e70931 (2021).

8. A. H. Melnyk, A. Wong, R. Kassen, *Evolutionary applications* **8**, 273–283 (2015).

9. K. Phan, T. Ferenci, *The ISME journal* **11**, 1472–1482 (2017).

10. S. Nair *et al.*, *Antimicrobial Agents and Chemotherapy* **66**, e01529–21 (2022).

11. L. Notley-McRobb, T. King, T. Ferenci, *Journal of bacteriology* **184**, 806–811 (2002).

12. E. Y. Valencia, J. P. Barros, T. Ferenci, B. Spira, *Microbial ecology* **83**, 68–82 (2022).

13. S. A. Smits *et al.*, *Science* **357**, 802–806, ISSN: 0036-8075, (http://science.sciencemag.org/content/357/6353/802) (2017).

14. J. F. Brooks *et al.*, *Cell* **184**, 4154–4167.e12, ISSN: 0092-8674 (2021).

15. V. Diwan, C. S. Lundborg, A. J. Tamhankar, *PLOS ONE* **8**, e68715, ISSN: 1932-6203 (2013).

16. I. Cvijović, B. H. Good, E. R. Jerison, M. M. Desai, *Proceedings of the National Academy of Sciences* **112**, E5021–E5028 (2015).

17. J. Monod, *Recherches sur la croissance des cultures bacteériennes* (Hermann and Cie, 1942).

18. J. E. Goldford *et al.*, *Science* **361**, 469, (http://science.sciencemag.org/content/361/6401/469.abstract) (2018).

19. N. Meroz, N. Tovi, Y. Sorokin, J. Friedman, *Nature Communications* **12**, 2891, ISSN: 2041-1723 (2021).

20. A. Erez, J. G. Lopez, B. G. Weiner, Y. Meir, N. S. Wingreen, *eLife* **9**, ed. by D. Weigel, A. Sanchez, A. Succurro, L. Dai, e57790, ISSN: 2050-084X, (https://doi.org/10.7554/eLife.57790) (2020).

21. A. Erez, J. G. Lopez, Y. Meir, N. S. Wingreen, *Physical Review E* **104**, 044412 (2021).

22. P.-Y. Ho, B. H. Good, K. C. Huang, *eLife* **11**, ed. by N. Segata, W. S. Garrett, S. Gibbons, e75168, ISSN: 2050-084X (2022).

23. L. Pacciani-Mori, A. Giometto, S. Suweis, A. Maritan, *PLOS Computational Biology* **16**, e1007896, ISSN: 1553-7358 (2020).

24. A. Aranda-Diaz *et al.*, *bioRxiv*, 2023–01 (2023).

25. S.-B. Hsu, P. Waltman, *Mathematical biosciences* **187**, 53–91 (2004).

26. R. E. Lenski, S. E. Hattingh, *Journal of Theoretical Biology* **122**, 83–93 (1986).

27. R. Maharjan *et al.*, *Ecology letters* **16**, 1267–1276 (2013).

28. A. D. Letten, A. R. Hall, J. M. Levine, *Nature ecology & evolution* **5**, 431–441 (2021).

29. M. Basan *et al.*, *Nature* **584**, 470–474 (2020).

30. E. Kussell, S. Leibler, *Science* **309**, 2075–2078 (2005).

31. J. Lin, M. Manhart, A. Amir, *Genetics* **215**, 767–777 (2020).

32. K. Phan, T. Ferenci, *The ISME Journal* **7**, 2034–2043 (2013).

33. Y. Kaplan *et al.*, *Nature* **600**, 290–294, ISSN: 1476-4687 (2021).

34. A. Posfai, T. Taillefumier, N. S. Wingreen, *Phys. Rev. Lett.* **118**, 028103, (https://link.aps.org/doi/10.1103/PhysRevLett.118.028103) (2 2017).

35. T. Song *et al.*, *Molecular microbiology* **91**, 1106–1119 (2014).

36. Y. Fridman, Z. Wang, S. Maslov, A. Goyal, *PLOS Computational Biology* **18**, e1010244, ISSN: 1553-7358 (2022).

37. J. Rivera-Pinto *et al.*, *mSystems* **3**, e00053–18 (2018).

38. J. Aitchison, *Journal of the Royal Statistical Society: Series B (Methodological)* **44**, 139–160, ISSN: 2517-6161 (1982).

39. J. Friedman, E. J. Alm, *PLOS Computational Biology* **8**, e1002687, ISSN: 1553-7358 (2012).

40. E. M. Brown, D. Nathwani, *Journal of antimicrobial chemotherapy* **55**, 6–9 (2005).

41. G. F. Webb, E. M. D'Agata, P. Magal, S. Ruan, *Proceedings of the National Academy of Sciences* **102**, 13343–13348 (2005).

42. O. Patange *et al.*, *Nature communications* **9**, 5333 (2018).

43. E. D. Kelsic, J. Zhao, K. Vetsigian, R. Kishony, *Nature* **521**, 516–519 (2015).

44. R. A. Sorg *et al.*, *PLoS biology* **14**, e2000631 (2016).

45. R. Marsland III *et al.*, *PLoS computational biology* **15**, e1006793 (2019).

46. O. Diekmann, *BANACH CENTER PUBLICATIONS* **63**, 47–86 (2002).

47. O. Tenaillon *et al.*, *Science* **335**, 457–461 (2012).

48. D. Graeber, *Debt: The First 5,000 Years* (Melville House, BROOKLYN, NY, Reprint edition, 2012), ISBN: 978-1-61219-129-4.

Appendix 1

Chemostat debt trade-offs

We consider a single debtor species growing within a chemostat and compute the values of Δ_E and ω at which this species has equivalent competitive ability as a non-debtor species.

$$\frac{d\rho}{dt} = \rho(E + \Delta_E)\frac{c}{c + K} - (\delta + \omega)\rho \tag{10}$$

$$\frac{dc}{dt} = S - \rho(E + \Delta_E)\frac{c}{c + K} \tag{11}$$

The outcome of competition within a chemostat is determined by the steady-state nutrient concentration, c^*. Species that can survive at the lower c^* will invade and exclude species with higher c^*. From Eq. 10, the c^* for a debtor species will be

$$c^* = \frac{K(\delta + \omega)}{(E + \Delta_E) - (\delta + \omega)}. \tag{12}$$

We now determine the values for Δ_E and ω at which the debtor has the same c^* as a non-debtor:

$$\frac{K(\delta + \omega)}{(E + \Delta_E) - (\delta + \omega)} = \frac{K\delta}{E - \delta}. \tag{13}$$

Solving the above expression yields:

$$\omega = \delta\frac{\Delta_E}{E}. \tag{14}$$

Thus, in a chemostat, a growth benefit of Δ_E is worthwhile if it comes with $\omega < \delta\frac{\Delta_E}{E}$.

Note that the requirement $\delta < E$ must be satisfied otherwise the chemostat washes out. Therefore, if c_0 increases, the steady state ρ must also increase, since increasing the nutrient concentration S while keeping the dilution rate constant gives $S/\delta = \rho$.

Chemostat model fitting to experimental data

To demonstrate applying our theory to empirical population dynamics, we fit data from Fig. 1A of Notley-McRobb et al. (*11*). We focus on these data as the chemostats at other pH values did not reach steady state. We first digitally extracted data from the figures. We utilize the chemostat equations in Eq. 11 to model these data. As these populations are not exposed to external stress, we set $\omega = 0$. Chemostats near steady state operate in a low-nutrient regime, so we approximate the growth function to be linear $E\left(\frac{c}{c+K}\right) \approx Ec$, absorbing the factor $1/K$ into E. We do not explicitly model mutation, but rather assume mutants arise very early in the system and fit an initial relative abundance of the non-debtor species $x_D(t = 0)$. Dilution rate δ is set based on the specified chemostat dilution rate. To convert from "generations" to physical time, we multiply by $\ln(2)/\delta$. Due to the fact that the input data are relative abundances, we can only estimate $\Delta E/E$ and also set $S = 1$. $\Delta E/E$ was estimated separately for each dilution. The model fitting and confidence/prediction interval estimation was performed using the same methods as in (*20*).

To apply our data to stressor deal kinetics, we fit data from Fig. 3A of Notley-McRobb et al., digitally extracting the data as above. We model the surviving fraction of the total microbial population as $f = (1 - x_N(t = 0))\exp(-\omega_D t) + x_N(t = 0)\exp(-\omega_N t)$, where $x_N(t = 0)$ is set using measurements from Notley-McRobb et al. All data is fit using a single value of ω_N and ω_D. Fitting and confidence/prediction interval estimation is performed as above. Note, however, that our current theory is not able to model all stressors: Notley-McRobb *et al.* also subjected their cultures to hydrogen peroxide stress and the resulting cell death kinetics deviate substantially from the first-order kinetics we assume.

> ## Serial dilution model dynamics in dimensionless variables
>
> It is sometimes useful to cast the dynamics in dimensionless variables. Our starting point is:
>
> $$\begin{aligned}
> \frac{dc}{dt} &= -g[c]\left(E\,\rho_N + (E + \Delta_E)\,\rho_D\right), \\
> \frac{1}{\rho_N}\frac{d\rho_N}{dt} &= E\,g[c] - \omega_N, \\
> \frac{1}{\rho_D}\frac{d\rho_D}{dt} &= (E + \Delta_E)\,g[c] - \omega_D.
> \end{aligned} \tag{15}$$
>
> By defining time in units of the growth rate of the normal non-debtor, $\bar{t} = Et$, similarly, nutrient in units of $\bar{c} = c/K$. We annotate $\bar{\Delta}_E = \Delta_E/E$ and $\bar{\omega}_N = \omega_N/E$. Explicitly using the Monod form for $g[c]$ we have,
>
> $$\begin{aligned}
> \frac{d\bar{c}}{d\bar{t}} &= -\frac{\bar{c}}{\bar{c} + 1}\left(\rho_N + (1 + \bar{\Delta}_E)\,\rho_D\right), \\
> \frac{1}{\rho_N}\frac{d\rho_N}{d\bar{t}} &= \frac{\bar{c}}{\bar{c} + 1} - \bar{\omega}_N, \\
> \frac{1}{\rho_D}\frac{d\rho_D}{d\bar{t}} &= (1 + \bar{\Delta}_E)\frac{\bar{c}}{\bar{c} + 1} - \bar{\omega}_D.
> \end{aligned} \tag{16}$$

Derivation of the chemostat limit of the serial dilutions, $c_0 \ll K$

Let species N be the normal, non-debtor species, growing at maximal rate E, and diluted by $\frac{\rho_0}{\rho_0 + c_0}$ to the next batch. Let the debtor species D grow at maximal rate $E + \Delta_E$ and get diluted by $\frac{\rho_0}{\rho_0 + c_0} e^{-\Omega t_f}$. We expand the low-nutrient limit,

$$
\begin{aligned}
\rho_\sigma(t) &= \rho_\sigma(0) + \phi \rho_\sigma^{(1)}(t) + O(\phi^2), \quad \left[\rho_\sigma^{(1\dots)}(0) = 0\right] \\
c(t) &= \phi c^{(1)}(t) + O(\phi^2).
\end{aligned}
\tag{17}
$$

We note that in the $c_0/K \ll 1$ limit (i.e., $\phi \ll 1$), we have,

$$
\frac{c}{c + K} = \phi \frac{c^{(1)}(t)}{K} + O(\phi^2)
\tag{18}
$$

The nutrient dynamics to order ϕ are,

$$
\begin{aligned}
\phi \frac{dc^{(1)}}{dt} + O(\phi^2) &= -\phi \frac{c^{(1)}}{K} \left(\rho_N(0) + \left(1 + \frac{\Delta_E}{E}\right) \rho_D(0) \right) \\
c^{(1)}(t) &= c^{(1)}(0) e^{-\gamma t}, \\
\gamma &= E \frac{\rho_N(0) + \left(1 + \frac{\Delta_E}{E}\right) \rho_D(0)}{K}.
\end{aligned}
\tag{19}
$$

We may now solve for the time-course integral,

$$
\begin{aligned}
\int_0^{t_f} \frac{c}{c + K} \, dt' &= \phi \int_0^{t_f} \frac{c^{(1)}(t)}{K} + O(\phi^2) \\
&= \phi \frac{c^{(1)}(0)}{K\gamma} \left(1 - e^{-\gamma t_f}\right) + O(\phi^2) \\
&\approx \frac{c_0}{K\gamma} + O(\phi^2) \quad \text{, at } t_f \gg 1/\gamma
\end{aligned}
\tag{20}
$$

The species abundance dynamics follow,

$$
\frac{d\rho_N}{dt} = \rho_N E \frac{c}{c + K} = \phi \rho_N(0) E \frac{c^{(1)}(t)}{K} + O(\phi^2)
\tag{21}
$$

Matching order ϕ and integrating both sides until time t_f, and using $\rho_N^{(1)}(0) = 0$, gives us

$$
\begin{aligned}
\rho_N^{(1)}(t_f) &= \rho_N(0) E \int_0^{t_f} \frac{c^{(1)}(t')}{K} \, dt' \\
&= \rho_N(0) E \frac{c^{(1)}(0)}{K\gamma} \left(1 - e^{-\gamma t_f}\right) \\
\rho_N(t_f) &= \rho_N(0) \left(1 + \frac{c_0 E}{K\gamma} \left(1 - e^{-\gamma t_f}\right)\right) + O(\phi^2) \\
\rho_D(t_f) &= \rho_D(0) \left(1 + \frac{c_0 E}{K\gamma} \left(1 + \frac{\Delta_E}{E}\right) \left(1 - e^{-\gamma t_f}\right)\right) + O(\phi^2)
\end{aligned}
\tag{22}
$$

At time (t_f) in batch b, each species gets diluted by a different fraction. Therefore, at the beginning of batch $b + 1$,

$$
\begin{aligned}
\rho_N^{\text{batch } (b+1)}(0) &= \frac{\rho_0}{\rho_0 + c_0} \rho_N^{\text{batch } (b)}(t_f), \\
\rho_D^{\text{batch } (b+1)}(0) &= e^{-\Omega t_f} \frac{\rho_0}{\rho_0 + c_0} \rho_D^{\text{batch } (b)}(t_f).
\end{aligned}
\tag{23}
$$

We require steady state, meaning that $\rho_\sigma^{\text{batch } (b+1)} = \rho_\sigma^{\text{batch } (b)}$. This gives,

$$\gamma = E \frac{\rho_0}{K}$$

$$\Omega t_f = \ln\left(\frac{\rho_0}{\rho_0 + c_0}\right) + \ln\left(1 + \frac{c_0}{\rho_0}\left(1 + \frac{\Delta_E}{E}\right)\right)$$

$$\frac{\Omega}{\Delta_E} \approx \frac{c_0}{\rho_0 E t_f} \tag{24}$$

Therefore, $\Omega/\Delta_E \sim c_0$ in the chemostat limit, $c_0 \ll K, \rho_0$ as shown in **Figure 3**.

In the language of the chemostat, the steady state requires $\omega = \delta\frac{\Delta_E}{E}$ (**Equation 14**). In the serial dilution framework, $\delta = \frac{c_0}{\rho_0 t_f}$ and $S = \frac{c_0}{t_f} = \rho_0\delta$ (20), therefore,

$$\frac{\omega}{\Delta_E} = \frac{c_0}{\rho_0 E t_f} . \tag{25}$$

This is precisely the relation derived in the leading-order c_0/K expansion, Eq. 24.

Debt trade-offs with differing dilution in the limit of high c_0

Here, we compute the growth-debt trade-offs in the limit of high c_0. In this scenario, the $c(t) \gg K$ during the vast majority of growth. Thus, we approximate $\frac{c}{c+K} \approx 1$ in this computation, a unit step function that suddenly terminates and gives zero. This approximation yields the following set of ODEs governing the batch dynamics:

$$\frac{d\rho_N}{dt} = \rho_N E, \tag{26}$$

$$\frac{d\rho_D}{dt} = \rho_D(E + \Delta_E), \tag{27}$$

$$\frac{dc}{dt} = -\rho_N E - \rho_D(E + \Delta_E). \tag{28}$$

Both populations are therefore described by simple exponential functions, $\rho_N(t) = \rho_N(0)e^{Et}$ and $\rho_D(t) = \rho_N(0)e^{(E+\Delta_E)t}$. Substituting these functions into the nutrient equation and solving yields $c(t) = c_0 + \rho_N(0)(1 - e^{Et}) + \rho_D(0)(1 - e^{(E+\Delta_E)t})$.

Response to Δ_K

We may evolve $c(t)$ according to the single species dynamics and calculate the definite integral. If the single species is a non-debtor, at high c_0/K, we have $\rho(t) \approx \rho_0 e^{Et}$ and therefore, $c(t) = c_0 + \rho_0(1 - e^{Et})$. Accordingly, $t^* \leq \frac{1}{E}\ln\left(1 + \frac{c_0}{\rho_0}\right)$.

$$E \int_0^{t_f} \frac{c}{(c+K)^2}\, dt' = \begin{cases} \frac{1}{K}\ln\left(1 + \frac{c_0}{\rho_0}\right) & c_0 \ll K \\ \frac{1}{c_0}\ln\frac{c_0^2}{K\rho_0} & c_0 \gg K \end{cases} \tag{29}$$

Therefore, at low c_0/K we have $\ln(1 + c_0/\rho_0) \approx c_0/\rho_0$ and,

$$\frac{\Delta_E}{E} \approx \begin{cases} \frac{\Delta_K}{K} & c_0 \ll K \\ \frac{\Delta_K}{c_0}\left(1 + \frac{\ln(c_0/K)}{\ln(c_0/\rho_0)}\right) & c_0 \gg K \end{cases} \tag{30}$$

Similarly, we may consider the opposite situation, where we evolve under the debtor species, $(K + \Delta_K, E + \Delta_E)$, and find the condition that the normal species can invade. A similar derivation gives $\frac{\Delta_E}{E+\Delta_E} = \frac{\Delta_{I[K+\Delta_K]}}{I_{[K+\Delta_K]}}$, with the I and Δ_I integrals evolved under the debtor species. Again, we may approximate, $\frac{\Delta_E}{E+\Delta_E}\ln\left(1 + \frac{c_0}{\rho_0}\right) = \Delta_K(E + \Delta E)\int \frac{c}{(c+K+\Delta_K)^2}\, dt'$, giving,

$$\frac{\Delta_E}{E + \Delta_E} \approx \begin{cases} \frac{\Delta_K}{K} & c_0 \ll K \\ \frac{\Delta_K}{c_0}\left(1 + \frac{\ln(c_0/(K+\Delta_K))}{\ln(c_0/\rho_0)}\right) & c_0 \gg K \end{cases} \tag{31}$$

The chemostat equivalent for a K debtor

We can consider the chemostat equivalent directly. We consider a single debtor species growing within a chemostat and compute the values of Δ_E and Δ_K at which this species has equivalent competitive ability as a non-debtor species.

$$\frac{d\rho}{dt} = \rho(E + \Delta_E)\frac{c}{c + K + \Delta_K} - \delta\rho, \tag{32}$$

$$\frac{dc}{dt} = S - \rho(E + \Delta_E)\frac{c}{c + K + \Delta_K} \tag{33}$$

The equilibrium for a debtor with the c^* of a non-debtor is,

$$\frac{(K + \Delta_K)\delta}{E + \Delta_E - \delta} = \frac{K\delta}{E - \delta}. \tag{34}$$

Solving the above expression yields $\frac{\Delta_E}{E} = \frac{\Delta_K}{K}\left(1 - \frac{\delta}{E}\right) = \frac{\Delta_K}{K}\left(1 - \frac{1}{Et_f}\ln(1 + c_0/\rho_0)\right)$. In the language of the serial-dilutions framework, $\delta = \frac{1}{t_f}\ln(1 + c_0/\rho_0) \approx \frac{c_0}{\rho t_f}$ and $S = \frac{c_0}{t_f} \approx \rho\delta$ (20), therefore,

$$\frac{\Delta_E}{E} \approx \frac{\Delta_K}{K}. \tag{35}$$

This is the relation derived in the leading-order c_0/K expansion.

Appendix 6—figure 1. Comparing the effect of noise in t_f and Ω. (*a*) Full curves - mean ρ_N fraction, averaged over 1000 stochastic trajectories. Green curve - deterministic convergence to non-debtor take-over. The three stochastic regimes show a similar, and much slower trend to mean ρ_N taking over the population than the deterministic case. (*b*) The median relative species log-fold-difference ρ_N/ρ_D. Viewed this way, the drift towards ρ_N takeover is the same for both the deterministic and the three stochastic regimes.

Appendix 6

Comparing noise in t_f with noise in Ω

If we keep all conditions the same, fluctuations in t_f are different from fluctuations in Ω because t_f affects both species through the utilization integral. We sought out to understand if fluctuations in t_f therefore lead to different behavior. We simulated a system that drifts to the debtor going extinct, by imposing a small positive Ω on top of the equilibrium Ω value that fits the debtor's 5% gain in maximal growth rate. *Figure 1* compares the effect of noise in t_f with noise in Ω. As is apparent, the noise in t_f acts the same way.