1   **Megakaryocyte maturation involves activation of the adaptive unfolded protein response**

2   [1]Faiz, M., [2]Kalev-Zylinska, M.L, [3]Singleton, D.C, [3]Hay, M.P. and [1*]Ledgerwood, E.C.

3   [1]Department of Biochemistry, School of Biomedical Sciences, University of Otago, Dunedin,
4   New Zealand

5   [2]Blood and Cancer Biology Laboratory, Department of Molecular Medicine & Pathology,
6   Faculty of Medical and Health Sciences, The University of Auckland, Auckland, New Zealand.

7   [3]Auckland Cancer Society Research Centre, Faculty of Medical and Health Sciences, The
8   University of Auckland, Auckland, New Zealand

9   *Corresponding author: liz.ledgerwood@otago.ac.nz

10 ORCID IDs

11 MF: 0000-0001-8681-6286
12 MLK-Z: 0000-0001-8378-8048
13 DCS: 0000-0002-2044-8078
14 MPH: 0000-0002-7087-9087
15 ECL: 0000-0002-4956-6043

16

17 **Short title: Megakaryopoiesis and the UPR**

18 **Keywords**: megakaryopoiesis, endoplasmic reticulum (ER) stress, unfolded protein response
19 (UPR), calcium, IRE1α, XBP1

20

Abstract

Endoplasmic reticulum stress triggers the unfolded protein response (UPR) to promote cell survival or apoptosis. Transient endoplasmic reticulum stress activation has been reported to trigger megakaryocyte production, and UPR activation has been reported as a feature of megakaryocytic cancers. However, the role of UPR signaling in megakaryocyte biology is not fully understood. We studied the involvement of UPR in human megakaryopoiesis using PMA (phorbol 12-myristate 13-acetate)-induced maturation of megakaryoblastic cell lines and thrombopoietin-induced differentiation of human peripheral blood-derived progenitors. Our results demonstrate that the adaptive UPR is a feature of megakaryocytic maturation, independent of ER stress-induced apoptosis, with the IRE1α UPR pathway being particularly important. Differentiation did not alter the ability of megakaryocytic cells to respond to the endoplasmic reticulum stressors DTT or thapsigargin. However, thapsigargin, but not DTT, inhibited differentiation, consistent with the involvement of Ca^{2+} signaling in megakaryopoiesis.

Introduction

Megakaryocytes are bone marrow cells that produce platelets. The differentiation of hematopoietic stem cells (HSC) to megakaryocytes (megakaryopoiesis) and subsequent platelet release (thrombopoiesis) is controlled by a number of transcription factors and cytokines. HSC undergo multiple rounds of endomitosis to produce large polyploid mature megakaryocytes with an extensive internal membrane reservoir for platelet formation (Machlus et al., 2014; Tilburg et al., 2022). Recent reports suggest that activation of the endoplasmic reticulum (ER) stress response is a feature of both normal megakaryopoiesis and thrombopoiesis (Kovuru et al., 2020; Lopez et al., 2013; Morishima & Nakanishi, 2016), and abnormal megakaryopoiesis in *CALR*-mutated myeloproliferative neoplasms (El-Khoury et al., 2020; Fosselteder et al., 2023; Jutzi et al., 2023; Nam et al., 2019).

The ER is involved in folding and post-translational modification of secreted and transmembrane proteins, storage of calcium, and lipid membrane biogenesis, with disruptions of these processes leading to ER stress (Braakman & Hebert, 2013; Clapham, 2007; Fagone & Jackowski, 2009). In response to ER stress, eukaryotic cells trigger the unfolded protein response (UPR) that initially helps cells to overcome the stress by upregulating the expression of chaperones to promote folding, repressing mRNA translation to temporarily inhibit global protein synthesis, and initiating ER-associated degradation (ERAD) pathways (adaptive UPR) (Feral et al., 2021; Hetz et al., 2020). If the adaptive UPR is unable to restore protein homeostasis, UPR signaling events may cause apoptosis (terminal UPR). There are three key UPR signal activators in mammals: IRE1α (inositol-requiring enzyme one alpha), PERK (protein kinase R-like ER kinase), and ATF6 (activating transcription factor 6). Activation of the endonuclease activity of IRE1α catalyzes removal of an intron from *XBP1* mRNA, resulting in production of Xbp1s, a transcription factor that drives the expression of genes involved in restoring protein homeostasis, localization and lipid synthesis (Adams et al., 2019). IRE1α also triggers regulated IRE1-dependent decay (RIDD), leading to RNA degradation in a cell and stimulus specific manner to regulate functions including apoptosis, metabolism and inflammation (Abdullah & Ravanan, 2018). In the PERK pathway, PERK phosphorylates eIF2α to inhibit global translation and at the same time promote translation of the ATF4 transcription factor. In the ATF6 pathway, ATF6 translocates to the Golgi network where is it cleaved by site 1 protease (S1P) to release the

ATF6p50 transcription factor. ATF4 and ATF6p50 control the expression of an overlapping set

of genes, including ones encoding proteins that contribute to protein folding in the ER (Hetz et

al., 2020).

UPR signaling events, and in particular the Ire1α pathway, are involved in the differentiation of

many cell types, including myoblasts (He et al., 2021), fibroblasts (Baek et al., 2012), osteoblasts

(Saito et al., 2011), neutrophils (Tanimura et al., 2018), plasma cells (Reimold et al., 2001),

mammary epithelial cells (Tsuchiya et al., 2017), eosinophils (Bettigole et al., 2015), dendritic

cells and macrophages (Bettigole & Glimcher, 2016). The UPR also has a vital role in sustaining

the activity of specialized secretory cells such as plasma cells and pancreatic β cells (Hetz &

Papa, 2018).

The role of UPR signaling in megakaryopoiesis and thrombopoiesis is unclear. It has been

proposed that ER stress activation triggers apoptotic-like caspase activation during

megakaryocytic differentiation of human megakaryoblastic cell lines, and may drive proplatelet

formation (Kovuru et al., 2020; Lopez et al., 2013; Morishima & Nakanishi, 2016). However,

while early reports suggested that localized caspase activation is required for megakaryopoiesis

(Clarke et al., 2003; De Botton et al., 2002), the role of caspase activation in megakaryopoiesis is

uncertain. Indeed genetic studies in mice convincingly showing that intrinsic apoptosis must be

suppressed for successful differentiation (Josefsson et al., 2014; McArthur et al., 2018). The

studies suggesting a role for ER stress-induced apoptosis in megakaryopoiesis have mostly used

PMA-induced maturation of various cell lines. However, the PMA concentrations used to induce

maturation and draw conclusions about the role of UPR signaling also resulted in extensive cell

death (Kovuru et al., 2020; Lopez et al., 2013; Morishima & Nakanishi, 2016). Therefore, it is

unclear whether UPR activation in these studies was a specific feature of megakaryocytic

maturation or a generalized response to cell stress. Here we investigate the involvement of UPR

signaling pathways during in vitro maturation of megakaryocytic cell lines and human peripheral

blood (PB) derived progenitors, under conditions that induce maturation in the absence of cell

death.

Materials and Methods

Cell culture and differentiation

Human myeloid leukemic cells, MEG-01 (ATCC CRL-2021) (Ogura et al., 1985) and K-562

(ATCC CRL-243) (Tabilio et al., 1983), were maintained in RPMI-1640 medium supplemented

with 10% fetal bovine serum (FBS), sodium bicarbonate (2 g/L), penicillin (100 U/mL), and

streptomycin (100 µg/mL) at 37°C in a 5% CO_2 humidified atmosphere. The identity of the cell

lines was confirmed by STR analysis (DNA Diagnostics, NZ) and the cells were confirmed to be

mycoplasma free. In vitro megakaryocytic maturation was induced by treating K-562 or MEG-

01 cells with 0.5 nM phorbol-12-myristate-13-acetate (PMA).

$CD45^+$ cells were isolated and purified from peripheral blood mononuclear cells from individual

donors as previously described (Balduini et al., 2009; Ong et al., 2017). To induce progenitor

differentiation into megakaryocytes, isolated $CD45^+$ cells were cultured in StemSpan^{TM} medium

(StemCell Technologies) supplemented with 10 ng/ml thrombopoietin (TPO), interleukin (IL) 6

and IL 11 (StemCell Technologies). Cultures were maintained at 37°C in a 5% CO_2 humidified

atmosphere for 14 days with media changes on days 5, 8, and 11.

ER stress modulators

The conventional ER stressors thapsigargin (Tg) (T9033; Sigma-Aldrich) or dithiothreitol (DTT)

(D0632; Sigma-Aldrich) were used at the indicated concentrations and time points. The

following compounds were used as UPR inhibitors: PF429242-dihydrochloride (SML0667;

Sigma-Aldrich), an ATF6 inhibitor (Hay et al., 2007); AMG-PERK-44 (HY-12661A,

MedChemExpress), a PERK inhibitor (Smith et al., 2015); and SN34221 (Supplementary data),

an IRE-1α endonuclease inhibitor (Patterson & Lonergan, 2008). The chemicals were dissolved

in DMSO or distilled water, and DMSO used as a control when appropriate.

Analysis of megakaryocytic maturation by flow cytometry

Cells were harvested, washed with Dulbecco's PBS, resuspended in 100 µL of Dulbecco's PBS,

and stained for 30 min in the dark at room temperature. Antibodies used were anti-CD61

phycoerythrin (PE) (1:20; 1075384; BD Biosciences), mouse IgG1 PE (isotype control;

7223589; BD Biosciences), anti-CD41a V450 (1:20; 1011309; BD Biosciences), V450 IgG1

(isotype control; 7128864; BD Biosciences). Live cells were determined with Zombie NIR

(1:100; B312060; BioLegend) or Zombie Green (1:100; B334686; BioLegend). After incubation,

the cells were washed with Dulbecco's PBS or wash buffer (1X Dulbecco's PBS, 1 mM EDTA,

3% FBS), resuspended in FACS buffer (0.01% sodium azide, 0.1% BSA in Dulbecco's PBS)

analyzed using the Guava® easyCyte 5 HPL benchtop flow cytometer (MerckMillipore) or BD

LSRFortessa[TM] cell analyzer (BD Biosciences). Data analysis was performed using FlowJo

version 10 (FlowJo, Ashland, USA). Data reported are geometric mean fluorescence intensity

(MFI) of the live cell (Zombie negative) population.

RNA isolation and RT-qPCR

Total RNA was isolated from K-562 and MEG-01 cells using TRIzol (Invitrogen). The

concentration of the RNA was measured using Qubit[TM] RNA HS Assay Kit (ThermoFisher

Scientific). First-strand cDNA synthesis was performed with 1 µg of total RNA in 20 µL

reaction volume using SuperScript[TM] IV VILO[TM] (SSIV VILO) master mix. Real-time

quantitative RT-PCR reactions were run on LightCycler[TM] 480 Instrument II System (Roche Life

Science) using PowerUp[TM] SYBER[TM] Green Master Mix (Thermo Fisher Scientific). All the

procedures were followed as recommended by the manufacturer. The qPCR primer sequences

are shown in Table S1. RT-qPCR was performed in triplicate, and relative expression was

quantified using the Pfaffl method (Pfaffl, 2001). Data were normalized to hypoxanthine

phosphoribosyl transferase 1 (*HPRT*) and glyceraldehyde 3-phoshate dehydrogenase (*GAPDH*)

using basket normalization (Vandesompele et al., 2002).

Semi-quantitative PCR analysis for XBP1 splicing detection

To detect *XBP1* splicing, *XBP1* mRNA was amplified (forward primer: 5′

CCTTGTAGTTGAGAACCAGG 3′; reverse primer 5′ AGGGGCTTGGTATATATGTGG 3′),

and the reactions were carried out in the C1000 Touch[TM] thermal cycler (Bio-Rad). Briefly, first-

strand cDNA was synthesized with 2 µg of total RNA in 20 µL reaction volume. To amplify

XBP1, 5 µL of undiluted cDNA was used as a template in a 50 µL reaction mixture with an

initial denaturation step for 3 min at 94 °C, followed by 30 cycles of denaturation for 30 s at 94

°C, annealing for 30 s at 54 °C, and extension for 60 s at 72 °C, and a final extension step for 10

min at 72 °C. PCR products were resolved on a 2% (w/v) agarose gel, stained with 0.5 µg/mL

ethidium bromide, and visualized using the Gel Doc[TM] XR+ Imaging System (Bio-Rad).

Statistical analysis

Statistical analyses were performed using GraphPad Prism software version 8.0 (GraphPad

Software, San Diego, CA, USA). The comparison of means between different groups was

performed using one sample t-test, one-way or two-way ANOVA, or two-tailed Student's t-test.

$P < 0.05$ was considered significant.

Results

The unfolded protein response is activated during megakaryocytic maturation of K-562

and MEG-01 cells

The K-562 and MEG-01 cell lines undergo megakaryocytic maturation upon PMA treatment

(Herrera et al., 1998; Kamal et al., 2018). Previous studies of the ER stress response in various

megakaryocytic cell lines have used concentrations of PMA (10-40 nM) that cause high levels

(>30%) of cell death (Kovuru et al., 2020; Lopez et al., 2013). We confirmed that treatment with

0.5 nM PMA stimulated maturation as assessed by an increase in the expression of the

transcription factor *GATA1* (Figure S1A&B) and the cell surface protein CD61 (Figure S1C).

Importantly there was no change in cell viability during maturation (Figure S1D), providing an

experimental system in which we could study the role of the UPR in megakaryocytic maturation

in the absence of confounding cell death.

To determine whether the UPR is activated during PMA-induced maturation of K-562 and

MEG-01 cells we examined the expression of UPR target genes (Figure 1). Expression of the

IRE1α target *ERdj4* (*DNAJB9*) and the ATF6 targets *HERPUD1* and *XBP1* increased in response

to PMA. The expression of *CHOP* (*DDIT3*), which is induced in response to all three UPR

pathways also increased, whereas the expression of the PERK target *GADD34* increased in K-

562 but not Meg-01 cells. These results suggest that the activation of UPR signaling pathways

occurs during megakaryocytic maturation. Since these changes in expression occurred without a

loss of cell viability, this most likely represents activation of the adaptive rather than terminal

UPR.

Figure 1. UPR target gene expression is upregulated during PMA-induced maturation of K-562 and MEG-01 cells into megakaryocytes. K-562 (A) or MEG-01 (B) cells were treated with 0.5 nM PMA and expression of *ERdj4, HERPUD1*, *XBP1*, *CHOP* and *GADD34* mRNA at days 0, 1, 2 and 3 was determined by RT-qPCR. Relative expression was calculated using the value at day 0 as 1. N=3 ± SD. *$P < 0.05$, **$P < 0.01$ compared to day 0 (one sample t-test).

Inhibition of IRE1α endonuclease activity, but not PERK activity or ATF6 processing, inhibits megakaryocytic maturation

We next used inhibitors of the three UPR signaling pathways to determine whether activation of one or more contributes to megakaryocytic maturation. SN34221 inhibits the endonuclease activity of IRE1α (Patterson & Lonergan, 2008), AMG PERK 44 inhibits PERK (Smith et al., 2015), and PF-429242 inhibits the site-1 protease that cleaves ATF6 to release ATF6p50 (Hay et al., 2007). The inhibitors showed the expected specificity with SN34221 decreasing expression of *ERdj4*, AMG PERK 44 decreasing expression of *CHOP*, *HERPUD1* and possibly *GADD34*, and PF-429242 decreasing expression of *CHOP* and *HERPUD1* (Figure S2). Inhibition of IRE1α had a negative impact on maturation, with 25% lower CD61 expression and 40% lower *GATA1* expression at day 3 at the highest concentration of the inhibitor (Figure 2A&D). Treatment with SN34221 was associated with a small decrease in cell viability (Figure S2D). In contrast CD61 but not *GATA1* expression was increased when PERK was inhibited (Figure 2B&D), although AMG PERK 44 also led to up to 50% loss in cell viability (Figure S2E) which may confound the result. Finally, inhibition of the ATF6 pathway had no impact on maturation (Figure 2C&D) or cell viability (Figure S2F).

Figure 2. The effect of pharmacological inhibition of UPR on K-562 maturation and peripheral
blood CD45+ cell differentiation. A - C. K-562 cells were treated with 0.5 nM PMA in the presence of
SN34221 (A), AMG PERK 44 (B) of PF429242 (C) and expression of CD61determined (fold change
MFI compared to day 0). N=3 ± SD *$P < 0.05$, ***$P < 0.001$, ****$P < 0.0001$ (two-way ANOVA). D. K-
562 cells were treated with 0.5 nM PMA in the presence (white bars) or absence (grey bars) of 50 μM
SN34221, 2 μM AMG PERK 44 or 1 μM PF429242 and relative *GATA1* expression measured at day 3.
N=3 ± SD. *$P < 0.05$, ** $P<0.01$, ***$P<0.001$ (unpaired t-test). E. % CD61+/CD41a+ cells at days 11
and 14 of TPO-induced megakaryocyte differentiation of CD45+ peripheral blood cells in the presence of
DMSO or 50 μM SN34221. N=3 ± SD *$P < 0.05$ compared to TPO + DMSO (two-way ANOVA).

Having observed the potential involvement of the IRE1α UPR pathway in megakaryocytic

maturation of K-562 cells, we next determined the impact of IRE1α inhibition on TPO-induced

megakaryocyte differentiation of CD45+ cells isolated from human peripheral blood. The IRE1α

endonuclease inhibitor SN34221 significantly decreased the percentage of CD61+/CD41a+

megakaryocytes at days 11 and 14 (Figure 2E). Consistent with this result, when cultures were

examined by bright field microscopy the cells treated with SN34221 were generally smaller and

there was no evidence of proplatelet formation or membrane blebbing (Figure S3C). These

results suggest that the IRE1α signaling branch of UPR may be involved in PMA-induced

megakaryocytic maturation of K-562 cells and TPO-induced megakaryocyte differentiation of

primary cells.

Activation of ER stress by thapsigargin, but not DTT, inhibits PMA-induced maturation of K-562 cells

Our data suggests that activation of the UPR, in particular the IRE1α pathway, may be involved

in megakaryopoiesis. We next determined whether activation of UPR signalling by an exogenous

ER stress had any impact on megakaryocyte maturation. We confirmed that the ER stressors

DTT, a small-molecule redox reagent for reducing protein disulfide bonds, and thapsigargin, a

sarco/endoplasmic reticulum ATPase (SERCA) inhibitor, induced UPR as shown by the splicing

of the IRE1α target *XBP1* (Figure S4). Treatment with DTT had no impact on megakaryocytic

maturation (Figure 3A) or viability (Figure 3B) of K-562 cells. In contrast, treatment with

thapsigargin caused a dose-dependent reduction in PMA-induced maturation of K-562 cells

(Figure 3C) with no change in cell viability (Figure 3D). Thapsigargin also inhibited TPO-

induced differentiation of CD45+ peripheral blood cells (Figure 3E) and subsequent budding and

proplatelet production (Figure S3D). The difference in effect of the two ER stressors suggests

that while exogenous UPR activation does not enhance megakaryocyte maturation, ER Ca^{2+}

homeostasis plays a central role.

Figure 3. The effect of UPR activation on K-562 maturation and peripheral blood CD45+ cell. A - D.
K-562 cells were treated with 0.5 nM PMA for 2 days in the presence of DTT (A and B) or Tg (C and D)
and analyzed for expression of CD61 (fold change MFI compared to day 0) (A and C) or cell viability (B
and D). N=3 ± SD. *$P < 0.05$, **$P < 0.01$ (one way ANOVA). **E.** % CD61+/CD41a+ cells at days 11 and
14 of TPO-induced megakaryocyte differentiation of CD45+ peripheral blood cells in the presence of
DMSO or 1 μM Tg. N=3 ± SD *$P < 0.05$ compared to TPO + DMSO (two-way ANOVA).

Megakaryocytic maturation of K-562 or MEG-01 cells does not alter the response to

exogenous ER stressors

Finally, we examined whether maturation alters the ability of the K-562 or MEG-01 cells to

respond to exogenous ER stressors. K-562 cells or MEG-01 cells were treated with DTT or

thapsigargin before or 3 days after PMA treatment, and the expression of UPR target genes was

quantified by RT-qPCR. DTT treatment activated the UPR as indicated by a strong induction of

XBP1 splicing, and *CHOP* and *BIP* expression, in immature (untreated) and mature (PMA

treated) K-562 cells (Figure 4A). Similarly, thapsigargin treatment induced *XBP1* splicing and

increased *CHOP, BIP* and *GADD34* expression in immature and mature K-562 cells (Figure 4B).

Similar results were obtained with MEG-01 cells where both ER stressors activated UPR

signaling in immature and mature cells (Figure S5). These results indicate that even though UPR

pathways are activated during PMA-induced maturation, this does not alter the ability of the cells

to further activate UPR signaling in response to exogenous ER stressors.

Figure 4. PMA treated and untreated K-562 cells are similarly responsive to ER stress induced by DTT or thapsigargin (Tg). Immature (no PMA) or mature (3 days after PMA treatment) K-562 cells were treated with 10 mM DTT for 0-2 h (A) or 0.1 µM Tg for 0-12 h (B). Expression of *XBP1*s and *XBP1*t, *BIP*, *CHOP* and *GADD34* mRNA was quantified by RT-qPCR. Expression is relative to 0 h. N=3 ± SD. *$P < 0.05$, **$P < 0.01$ (one sample t-test).

Discussion

In the current study we used a low concentration of PMA that induced megakaryocytic

maturation of K-562 and MEG-01 cell lines in the absence of cell death. Under these conditions

maturation was accompanied by increased transcription of UPR target genes *ERDj4, HERPUD1*,

XBP1, *CHOP*, and *GADD34,* indicating activation of all three UPR signaling pathways. These

results are consistent with previous studies reporting an increase in expression of UPR target

genes in various models of megakaryocytic maturation (Kovuru et al., 2020; Lopez et al., 2013;

Morishima & Nakanishi, 2016). However, in contrast to these earlier studies, increased

expression of UPR target genes occurred in the absence of cell death, demonstrating that UPR-

induced apoptosis is not required for megakaryocytic maturation. The terminal stages of

megakaryopoiesis involve endomitosis and polyploidization with an accompanying high demand

for synthesis of proteins and membrane expansion and remodeling prior to proplatelet formation

and platelet release (de Jonckheere et al., 2023; Mazzi et al., 2018). Therefore, it is unsurprising

that the adaptive UPR is activated in maturing megakaryocytes to assist terminal

megakaryopoiesis by increasing the biosynthesis of chaperone proteins, increasing membrane

synthesis, and boosting the protein folding capacity of the ER.

To determine whether activation of a specific arm of the adaptive UPR was particularly

important, we inhibited IRE1α endonuclease activity (SN34221), PERK autophosphorylation

(AMG PERK 44) and ATF6 cleavage (PF-429242). The IRE1α inhibitor SN34221 decreased

PMA-induced maturation of K-562 cells and TPO-induced differentiation of peripheral blood

CD45+ cells, implicating this UPR pathway in supporting megakaryocytic maturation. Whether

XBP1 splicing and/or RIDD activation is relevant for human megakaryopoiesis remains an open

question. XBP1s is involved in maturation of other hematopoietic cell types (Bettigole &

Glimcher, 2016; Reimold et al., 2001), and is an important regulator of specialist secretory cells

which require a high level of protein synthesis (Lee et al., 2005; Reimold et al., 2001). XBP1s

regulates gene expression by forming heterodimers with other transcription factors including

MIST1, an important regulator of differentiation (Acosta-Alvear et al., 2007; Hetz & Papa,

2018). A recent characterization of mice with megakaryocyte/platelet-specific deletion of *XBP1*

found normal platelet counts but a higher percentage of circulating reticulated (newly released)

platelets (Jain et al., 2022) suggesting possible alteration in megakaryopoiesis and/or

thrombopoiesis. RIDD has also been implicated in controlling differentiation-associated

processes (He et al., 2021; Levi-Ferber et al., 2021) but has not been directly studied in the

context of megakaryocyte biology. Our observations that both immature and mature K-562 and

MEG-01cells show similar responses to the exogenous ER stressors DTT and thapsigargin, and

that DTT has no impact on megakaryocytic maturation, suggest that the potential role of IRE1α

response is distinct from a canonical ER stress response to accumulation of unfolded proteins.

Further studies are needed to tease out the roles of XBP1s and RIDD in human

megakaryopoiesis and thrombopoiesis.

Inhibition of the PERK pathway enhanced maturation, suggesting that this pathway may act to

restrain megakaryocyte maturation. This result is consistent with inhibition of PMA-induced

megakaryocytic maturation by salubrinal, a drug which potentiates eIF2α phosphorylation

(Boyce et al., 2005; Lopez et al., 2013), but in contrast to the decreased platelet count seen in

mice with megakaryocyte/platelet-specific deletion of *PERK* (Jain et al., 2022). The high level of

cell death we saw with PERK inhibition may explain the decreased platelet count observed by

Jain et al (2022).

Treatment with thapsigargin inhibited both PMA-induced maturation of K-562 cells and TPO-

induced differentiation of peripheral blood-derived CD45+ cells. Thapsigargin blocks SERCA,

inducing protein misfolding and the UPR, and altering Ca^{2+} homeostasis (Jaskulska et al., 2020;

Sehgal et al., 2017). Since DTT had no impact on megakaryocytic maturation, it is likely that

dysregulation of Ca^{2+} homeostasis underlies inhibition of maturation by thapsigargin. This is

consistent with the known roles of intracellular Ca^{2+} in megakaryopoiesis. Activation of

phospholipase C generates inositol trisphosphate (IP3) triggering release of Ca^{2+} from the ER via

the IP3 receptor. Subsequent store-operated calcium entry (SOCE) induces expression of nuclear

factor of activated T cells (NFAT), a transcriptional regulator implicated in megakaryopoiesis

(Hogan et al., 2003; Zaslavsky et al., 2013; Zhu et al., 2018). Additionally, Ca^{2+}entry through a

membranous Ca^{2+} ion channel N-methyl-D-aspartate receptor supports megakaryocyte

differentiation (Hearn et al., 2020; Hearn et al., 2022). Ca^{2+}-dependent activation of

Ca^{2+}/calmodulin-dependent protein kinase (CaMK) induces transcription of CREB (c-AMP

response element-binding protein) to activate the genes with CRE elements, including

megakaryocyte-lineage specific genes such as *FLI1*, *HOXC6*, *MXD3*, *PRDM16*, *FOXB1*, *TBX6*,

HDAC11 and *NPAS1* (Hogan et al., 2003; Zhang et al., 2005; Zhu et al., 2018).

Megakaryopoiesis and thrombopoiesis are complex process that require the activation of many

synergistic pathways. Overall our findings firstly indicate that the adaptive UPR, potentially

involving IRE1α activation, rather than terminal UPR is a feature of megakaryocyte

differentiation, and secondly provide further evidence for the importance of calcium signaling in

megakaryopoiesis.

References

Abdullah, A., & Ravanan, P. (2018). The unknown face of IRE1α - Beyond ER stress. *Eur J Cell*
Biol, 97(5), 359-368. https://doi.org/10.1016/j.ejcb.2018.05.002

Acosta-Alvear, D., Zhou, Y., Blais, A., Tsikitis, M., Lents, N. H., Arias, C., Lennon, C. J.,
Kluger, Y., & Dynlacht, B. D. (2007). XBP1 controls diverse cell type- and condition-
specific transcriptional regulatory networks. *Mol Cell, 27*(1), 53-66.
https://doi.org/10.1016/j.molcel.2007.06.011

Adams, C. J., Kopp, M. C., Larburu, N., Nowak, P. R., & Ali, M. M. U. (2019). Structure and
Molecular Mechanism of ER Stress Signaling by the Unfolded Protein Response Signal
Activator IRE1. *Front Mol Biosci, 6*, 11. https://doi.org/10.3389/fmolb.2019.00011

Baek, H. A., Kim, D. S., Park, H. S., Jang, K. Y., Kang, M. J., Lee, D. G., Moon, W. S., Chae, H.
J., & Chung, M. J. (2012). Involvement of endoplasmic reticulum stress in myofibroblastic
differentiation of lung fibroblasts. *Am J Respir Cell Mol Biol, 46*(6), 731-739.
https://doi.org/10.1165/rcmb.2011-0121OC

Balduini, A., Malara, A., Pecci, A., Badalucco, S., Bozzi, V., Pallotta, I., Noris, P., Torti, M., &
Balduini, C. L. (2009). Proplatelet formation in heterozygous Bernard-Soulier syndrome
type Bolzano. *J Thromb Haemost, 7*(3), 478-484. https://doi.org/10.1111/j.1538-
7836.2008.03255.x

Bettigole, S. E., & Glimcher, L. H. (2016). Novel roles for XBP1 in hematopoietic development.
Cell Cycle, 15(13), 1653-1654. https://doi.org/10.1080/15384101.2016.1172470

Bettigole, S. E., Lis, R., Adoro, S., Lee, A. H., Spencer, L. A., Weller, P. F., & Glimcher, L. H.
(2015). The transcription factor XBP1 is selectively required for eosinophil differentiation.
Nat Immunol, 16(8), 829-837. https://doi.org/10.1038/ni.3225

Boyce, M., Bryant, K. F., Jousse, C., Long, K., Harding, H. P., Scheuner, D., Kaufman, R. J.,
Ma, D., Coen, D. M., Ron, D., & Yuan, J. (2005). A selective inhibitor of eIF2α

dephosphorylation protects cells from ER stress. *Science, 307*(5711), 935-939. https://doi.org/10.1126/science.1101902

Braakman, I., & Hebert, D. N. (2013). Protein folding in the endoplasmic reticulum. *Cold Spring Harb Perspect Biol, 5*(5), a013201. https://doi.org/10.1101/cshperspect.a013201

Clapham, D. E. (2007). Calcium signaling. *Cell, 131*(6), 1047-1058. https://doi.org/10.1016/j.cell.2007.11.028

Clarke, M. C., Savill, J., Jones, D. B., Noble, B. S., & Brown, S. B. (2003). Compartmentalized megakaryocyte death generates functional platelets committed to caspase-independent death. *J Cell Biol, 160*(4), 577-587. https://doi.org/10.1083/jcb.200210111

De Botton, S., Sabri, S., Daugas, E., Zermati, Y., Guidotti, J. E., Hermine, O., Kroemer, G., Vainchenker, W., & Debili, N. (2002). Platelet formation is the consequence of caspase activation within megakaryocytes. *Blood, 100*(4), 1310-1317. https://doi.org/10.1182/blood-2002-03-0686

de Jonckheere, B., Kollotzek, F., Munzer, P., Gob, V., Fischer, M., Mott, K., Coman, C., Troppmair, N. N., Manke, M. C., Zdanyte, M., Harm, T., Sigle, M., Kopczynski, D., Bileck, A., Gerner, C., Hoffmann, N., Heinzmann, D., Assinger, A., Gawaz, M., Stegner, D., Schulze, H., Borst, O., & Ahrends, R. (2023). Critical shifts in lipid metabolism promote megakaryocyte differentiation and proplatelet formation. *Nat Cardiovasc Res, 2*, 835-852. https://doi.org/10.1038/s44161-023-00325-8

El-Khoury, M., Cabagnols, X., Mosca, M., Vertenoeil, G., Marzac, C., Favale, F., Bluteau, O., Lorre, F., Tisserand, A., Rabadan Moraes, G., Ugo, V., Ianotto, J. C., Rey, J., Solary, E., Roy, L., Rameau, P., Debili, N., Pasquier, F., Casadevall, N., Marty, C., Constantinescu, S. N., Raslova, H., Vainchenker, W., & Plo, I. (2020). Different impact of calreticulin mutations on human hematopoiesis in myeloproliferative neoplasms. *Oncogene, 39, 5323-5337*. https://doi.org/10.1038/s41388-020-1368-3

Fagone, P., & Jackowski, S. (2009). Membrane phospholipid synthesis and endoplasmic reticulum function. *J Lipid Res, 50 Suppl*(Suppl), S311-316. https://doi.org/10.1194/jlr.R800049-JLR200

Feral, K., Jaud, M., Philippe, C., Di Bella, D., Pyronnet, S., Rouault-Pierre, K., Mazzolini, L., & Touriol, C. (2021). ER Stress and Unfolded Protein Response in Leukemia: Friend, Foe, or Both? *Biomolecules, 11*(2), 199. https://doi.org/10.3390/biom11020199

Fosselteder, J., Pabst, G., Sconocchia, T., Schlacher, A., Auinger, L., Kashofer, K., Beham-Schmid, C., Trajanoski, S., Waskow, C., Scholl, W., Sill, H., Zebisch, A., Wolfler, A., Thomas, D., & Reinisch, A. (2023). Human gene-engineered calreticulin mutant stem cells recapitulate MPN hallmarks and identify targetable vulnerabilities. *Leukemia, 37*(4), 843-853. https://doi.org/10.1038/s41375-023-01848-6

Hay, B. A., Abrams, B., Zumbrunn, A. Y., Valentine, J. J., Warren, L. C., Petras, S. F., Shelly, L. D., Xia, A., Varghese, A. H., Hawkins, J. L., Van Camp, J. A., Robbins, M. D., Landschulz, K., & Harwood, H. J., Jr. (2007). Aminopyrrolidineamide inhibitors of site-1 protease. *Bioorg Med Chem Lett, 17*(16), 4411-4414. https://doi.org/10.1016/j.bmcl.2007.06.031

He, S., Fu, T., Yu, Y., Liang, Q., Li, L., Liu, J., Zhang, X., Zhou, Q., Guo, Q., Xu, D., Chen, Y., Wang, X., Chen, Y., Liu, J., Gan, Z., & Liu, Y. (2021). IRE1α regulates skeletal muscle regeneration through Myostatin mRNA decay. *J Clin Invest, 131*(17). https://doi.org/10.1172/JCI143737

Hearn, J. I., Green, T. N., Chopra, M., Nursalim, Y. N. S., Ladvanszky, L., Knowlton, N., Blenkiron, C., Poulsen, R. C., Singleton, D. C., Bohlander, S. K., & Kalev-Zylinska, M. L. (2020). N-Methyl-D-Aspartate Receptor Hypofunction in Meg-01 Cells Reveals a Role for Intracellular Calcium Homeostasis in Balancing Megakaryocytic-Erythroid Differentiation. *Thromb Haemost, 120*(4), 671-686. https://doi.org/10.1055/s-0040-1708483

Hearn, J. I., Green, T. N., Hisey, C. L., Bender, M., Josefsson, E. C., Knowlton, N., Baumann, J., Poulsen, R. C., Bohlander, S. K., & Kalev-Zylinska, M. L. (2022). Deletion of Grin1 in mouse megakaryocytes reveals NMDA receptor role in platelet function and proplatelet formation. *Blood, 139*(17), 2673-2690. https://doi.org/10.1182/blood.2021014000

Herrera, R., Hubbell, S., Decker, S., & Petruzzelli, L. (1998). A Role of the MEK/MAPK Pathway in PMA-induced Cell Cycle Arrest: Modulation of Megakaryocytic Differentiation of K-562 Cells. *Exp Cell Res, 238*, 407-414.

Hetz, C., & Papa, F. R. (2018). The Unfolded Protein Response and Cell Fate Control. *Mol Cell, 69*(2), 169-181. https://doi.org/10.1016/j.molcel.2017.06.017

Hetz, C., Zhang, K., & Kaufman, R. J. (2020). Mechanisms, regulation and functions of the unfolded protein response. *Nat Rev Mol Cell Biol, 21*(8), 421-438. https://doi.org/10.1038/s41580-020-0250-z

Hogan, P. G., Chen, L., Nardone, J., & Rao, A. (2003). Transcriptional regulation by calcium, calcineurin, and NFAT. *Genes Dev, 17*(18), 2205-2232. https://doi.org/10.1101/gad.1102703

Jain, K., Tyagi, T., Du, J., Hu, X., Patell, K., Martin, K. A., & Hwa, J. (2022). Unfolded Protein Response Differentially Modulates the Platelet Phenotype. *Circ Res, 131*(4), 290-307. https://doi.org/10.1161/CIRCRESAHA.121.320530

Jaskulska, A., Janecka, A. E., & Gach-Janczak, K. (2020). Thapsigargin-From Traditional Medicine to Anticancer Drug. *Int J Mol Sci, 22*(1). https://doi.org/10.3390/ijms22010004

Josefsson, E. C., Burnett, D. L., Lebois, M., Debrincat, M. A., White, M. J., Henley, K. J., Lane, R. M., Moujalled, D., Preston, S. P., O'Reilly, L. A., Pellegrini, M., Metcalf, D., Strasser, A., & Kile, B. T. (2014). Platelet production proceeds independently of the intrinsic and extrinsic apoptosis pathways. *Nat Commun, 5*, 3455. https://doi.org/10.1038/ncomms4455

Jutzi, J. S., Marneth, A. E., Jimenez-Santos, M. J., Hem, J., Guerra-Moreno, A., Rolles, B., Bhatt, S., Myers, S. A., Carr, S. A., Hong, Y., Pozdnyakova, O., van Galen, P., Al-Shahrour, F., Nam, A. S., & Mullally, A. (2023). CALR-mutated cells are vulnerable to combined inhibition of the proteasome and the endoplasmic reticulum stress response. *Leukemia, 37*(2), 359-369. https://doi.org/10.1038/s41375-022-01781-0

Kamal, T., Green, T. N., Hearn, J. I., Josefsson, E. C., Morel-Kopp, M. C., Ward, C. M., During, M. J., & Kalev-Zylinska, M. L. (2018). N-methyl-d-aspartate receptor mediated calcium influx supports in vitro differentiation of normal mouse megakaryocytes but proliferation of leukemic cell lines. *Res Pract Thromb Haemost, 2*(1), 125-138. https://doi.org/10.1002/rth2.12068

Kovuru, N., Raghuwanshi, S., Sharma, D. S., Dahariya, S., Pallepati, A., & Gutti, R. K. (2020). Endoplasmic reticulum stress induced apoptosis and caspase activation is mediated through mitochondria during megakaryocyte differentiation. *Mitochondrion, 50*, 115-120. https://doi.org/10.1016/j.mito.2019.10.009

Lee, A. H., Chu, G. C., Iwakoshi, N. N., & Glimcher, L. H. (2005). XBP-1 is required for biogenesis of cellular secretory machinery of exocrine glands. *EMBO J, 24*(24), 4368-4380. https://doi.org/10.1038/sj.emboj.7600903

Levi-Ferber, M., Shalash, R., Le-Thomas, A., Salzberg, Y., Shurgi, M., Benichou, J. I., Ashkenazi, A., & Henis-Korenblit, S. (2021). Neuronal regulated *ire-1*-dependent mRNA decay controls germline differentiation in *Caenorhabditis elegans*. *Elife, 10*. https://doi.org/10.7554/eLife.65644

Lopez, J. J., Palazzo, A., Chaabane, C., Albarran, L., Polidano, E., Lebozec, K., Dally, S., Nurden, P., Enouf, J., Debili, N., & Bobe, R. (2013). Crucial role for endoplasmic

reticulum stress during megakaryocyte maturation. *Arteriosc, Thromb Vasc Biol, 33*(12), 2750-2758. https://doi.org/10.1161/ATVBAHA.113.302184

Machlus, K. R., Thon, J. N., & Italiano, J. E., Jr. (2014). Interpreting the developmental dance of the megakaryocyte: a review of the cellular and molecular processes mediating platelet formation. *Br J Haematol, 165*(2), 227-236. https://doi.org/10.1111/bjh.12758

Mazzi, S., Lordier, L., Debili, N., Raslova, H., & Vainchenker, W. (2018). Megakaryocyte and polyploidization. *Exp Hematol, 57*, 1-13. https://doi.org/10.1016/j.exphem.2017.10.001

McArthur, K., Chappaz, S., & Kile, B. T. (2018). Apoptosis in megakaryocytes and platelets: the life and death of a lineage. *Blood, 131*(6), 605-610. https://doi.org/10.1182/blood-2017-11-742684

Morishima, N., & Nakanishi, K. (2016). Proplatelet formation in megakaryocytes is associated with endoplasmic reticulum stress. *Genes Cells, 21*(7), 798-806. https://doi.org/10.1111/gtc.12384

Nam, A. S., Kim, K. T., Chaligne, R., Izzo, F., Ang, C., Taylor, J., Myers, R. M., Abu-Zeinah, G., Brand, R., Omans, N. D., Alonso, A., Sheridan, C., Mariani, M., Dai, X., Harrington, E., Pastore, A., Cubillos-Ruiz, J. R., Tam, W., Hoffman, R., Rabadan, R., Scandura, J. M., Abdel-Wahab, O., Smibert, P., & Landau, D. A. (2019). Somatic mutations and cell identity linked by Genotyping of Transcriptomes. *Nature, 571*(7765), 355-360. https://doi.org/10.1038/s41586-019-1367-0

Ogura, M., Morishima, Y., Ohno, R., Kato, Y., Hirabayashi, N., Nagura, H., & Saito, H. (1985). Establishment of a novel human megakaryoblastic leukemia cell line, MEG-01, with positive Philadelphia chromosome. *Blood, 66*(6), 1384-1392. https://doi.org/10.1182/blood.V66.6.1384.1384

Ong, L., Morison, I. M., & Ledgerwood, E. C. (2017). Megakaryocytes from CYCS mutation-associated thrombocytopenia release platelets by both proplatelet-dependent and -independent processes. *Br J Haematol, 176*(2), 268-279. https://doi.org/10.1111/bjh.14421

Patterson, J. B., & Lonergan, D. G. (2008). *IRE-1alpha inhibitors* PCT WO 2008 154484 A1.

Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res, 29*(9), e45. https://doi.org/10.1093/nar/29.9.e45

Reimold, A. M., Iwakoshi, N. N., Manis, J., Vallabhajosyula, P., Szomolanyi-Tsuda, E., Gravallese, E. M., Friend, D., Grusby, M. J., Alt, F., & Glimcher, L. H. (2001). Plasma cell

differentiation requires the transcription factor XBP-1. *Nature, 412*(6844), 300-307. https://doi.org/10.1038/35085509

Saito, A., Ochiai, K., Kondo, S., Tsumagari, K., Murakami, T., Cavener, D. R., & Imaizumi, K. (2011). Endoplasmic reticulum stress response mediated by the PERK-eIF2(alpha)-ATF4 pathway is involved in osteoblast differentiation induced by BMP2. *J Biol Chem, 286*(6), 4809-4818. https://doi.org/10.1074/jbc.M110.152900

Sehgal, P., Szalai, P., Olesen, C., Praetorius, H. A., Nissen, P., Christensen, S. B., Engedal, N., & Moller, J. V. (2017). Inhibition of the sarco/endoplasmic reticulum (ER) Ca(2+)-ATPase by thapsigargin analogs induces cell death via ER Ca(2+) depletion and the unfolded protein response. *J Biol Chem, 292*(48), 19656-19673. https://doi.org/10.1074/jbc.M117.796920

Smith, A. L., Andrews, K. L., Beckmann, H., Bellon, S. F., Beltran, P. J., Booker, S., Chen, H., Chung, Y. A., D'Angelo, N. D., Dao, J., Dellamaggiore, K. R., Jaeckel, P., Kendall, R., Labitzke, K., Long, A. M., Materna-Reichelt, S., Mitchell, P., Norman, M. H., Powers, D., Rose, M., Shaffer, P. L., Wu, M. M., & Lipford, J. R. (2015). Discovery of 1H-pyrazol-3(2H)-ones as potent and selective inhibitors of protein kinase R-like endoplasmic reticulum kinase (PERK). *J Med Chem, 58*(3), 1426-1441. https://doi.org/10.1021/jm5017494

Tabilio, A., Pelicci, P. G., Vinci, G., Mannoni, P., Civin, C. I., Vainchenker, W., Testa, U., Lipinski, M., Rochant, H., & Breton-Gorius, J. (1983). Myeloid and megakaryocytic properties of K-562 cell lines. *Cancer Res, 43*(10), 4569-4574. https://www.ncbi.nlm.nih.gov/pubmed/6349795

Tanimura, A., Miyoshi, K., Horiguchi, T., Hagita, H., Fujisawa, K., & Noma, T. (2018). Mitochondrial Activity and Unfolded Protein Response are Required for Neutrophil Differentiation. *Cell Physiol Biochem, 47*(5), 1936-1950. https://doi.org/10.1159/000491464

Tilburg, J., Becker, I. C., & Italiano, J. E. (2022). Don't you forget about me(gakaryocytes). *Blood, 139*(22), 3245-3254. https://doi.org/10.1182/blood.2020009302

Tsuchiya, M., Koizumi, Y., Hayashi, S., Hanaoka, M., Tokutake, Y., & Yonekura, S. (2017). The role of unfolded protein response in differentiation of mammary epithelial cells. *Biochem Biophys Res Commun, 484*(4), 903-908. https://doi.org/10.1016/j.bbrc.2017.02.042

Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A., & Speleman, F. (2002). Accurate normalization of real-time quantitative RT-PCR data by geometric

averaging of multiple internal control genes. *Genome Biol, 3*(7), research0034.1-0034.11
https://doi.org/10.1186/gb-2002-3-7-research0034

Zaslavsky, A., Chou, S. T., Schadler, K., Lieberman, A., Pimkin, M., Kim, Y. J., Baek, K. H.,
Aird, W. C., Weiss, M. J., & Ryeom, S. (2013. The calcineurin-NFAT pathway negatively
regulates megakaryopoiesis. *Blood, 121*(16), 3205-3215. https://doi.org/10.1182/blood-
2012-04-421172

Zhang, X., Odom, D. T., Koo, S. H., Conkright, M. D., Canettieri, G., Best, J., Chen, H., Jenner,
R., Herbolsheimer, E., Jacobsen, E., Kadam, S., Ecker, J. R., Emerson, B., Hogenesch, J.
B., Unterman, T., Young, R. A., & Montminy, M. (2005). Genome-wide analysis of
cAMP-response element binding protein occupancy, phosphorylation, and target gene
activation in human tissues. *Proc Natl Acad Sci U S A, 102*(12), 4459-4464.
https://doi.org/10.1073/pnas.0501076102

Zhu, F., Feng, M., Sinha, R., Seita, J., Mori, Y., & Weissman, I. L. (2018). Screening for genes
that regulate the differentiation of human megakaryocytic lineage cells. *Proc Natl Acad Sci
U S A, 115*(40), E9308-E9316. https://doi.org/10.1073/pnas.1805434115

Supplementary Materials

The supplementary data file contains the following:

Synthesis of SN34221

Figure S1. PMA induces megakaryocyte maturation of K-562 and MEG-01 cells.

Figure S2. Impact of UPR inhibitors on cell viability and maturation of K-562 cells.

Figure S3. Impact of SN34221 or thapsigargin on proplatelet production.

Figure S4. Thapsigargin and DTT induce *XBP1* splicing in K-562 cells.

Figure S5. UPR activation in immature and mature MEG-01 cells in response to DTT or thapsigargin.

Figure S6. SN34221 inhibits *XBP1* splicing.

Table S1. RT-qPCR primers used in the study.

Statements and Declarations

Funding: This work was supported by Leukaemia & Blood Cancer New Zealand (to ECL), University of Otago School of Biomedical Sciences Dean's Bequest Fund (to ECL), and Auckland Medical Research Foundation (to MK-Z). Author MF was supported by a University of Otago Doctoral Scholarship. Authors MPH and DCS were supported by Cancer Society Auckland/Northland.

Competing interests: The authors have no competing interests to declare.

Author contributions: ECL and MK-Z conceived the study. ECL, MK-Z and MF designed the study. Experimental work was performed by MF. All authors contributed to data analysis. The first draft of the manuscript was written by MF and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Data availability: Data and materials included within the article or Supplementary Materials are available from the authors on request.

Preprint: This article has been deposited on BioRxiv: https://www.biorxiv.org/content/10.1101/2023.07.26.550730v1.full

Ethics approval: The study was approved by the University of Otago Human Ethics Committee (NZ) (H20/132).

Consent to participate: Written informed consent was obtained from all individual participants included in the study.