Artificial selection optimizes pollutant-degrading bacterial communities

Flor I. Arias-Sánchez[1,2], Björn Vessman[2], Alice Haym[2], Géraldine Alberti[2], and Sara Mitri[2, 3]

[1]BIH Center for Regenerative Therapies (BCRT), Charité - Universitätsmedizin Berlin, Berlin, Germany
[2]Département de Microbiologie Fondamentale, Université de Lausanne, 1015 Lausanne, Switzerland
[3]Swiss Institute of Bioinformatics

Abstract. Artificial selection is a promising way to improve microbial community functions, but previous experiments have only shown moderate success. Here, we experimentally evaluate a new method that was inspired by genetic algorithms to artificially select small bacterial communities of known species composition based on their degradation of an industrial pollutant. Starting from 29 randomly generated four-species communities, we repeatedly grew communities for four days, selected the 10 best-degrading communities, and rearranged them into 29 new communities with species compositions that resembled those of the most successful ones. The best community after 18 such rounds of selection degraded the pollutant better than the best community in the first round. It featured member species that degrade well, species that degrade badly alone but improve community degradation, and free-rider species that did not contribute to community degradation. Most species in the evolved communities did not differ significantly from their ancestors, suggesting that genetic evolution plays a small role at this time scale. These experiments show that artificial selection on microbial communities can work in principle, and inform on how to improve future experiments.

Introduction

Microbial communities naturally provide us with many ecosystem functions like digesting inaccessible nutrients or cleaning wastewater. Being able to design such multi-species communities from scratch to optimize ecosystem functions would be a major biotechnological breakthrough, but knowing which species to combine and how such a choice will affect ecological and evolutionary dynamics and thereby functional dynamics is a very challenging problem.

A first intuitive approach is to collect candidate species, study their capacities through genomic and phenotypic analyses and then combine them in clever ways that are likely to result in high function (1–4). An alternative is to automate the optimization process while remaining blind to the properties of each species. This blind approach can be taken using artificial selection (5, 6).

Artificial selection – also known as "directed evolution" or simply "breeding" – is a powerful approach that takes inspiration from natural selection. Not only has it revolutionized agriculture (7), but artificial selection has also been successfully applied in chemistry to optimize industrial enzymes (8, 9), or in pharmacy to reduce HIV drug production costs (10). These success stories have sparked the idea of artificially selecting microbial communities, promising to enhance human and ecosystem health, as well as many industrial applications.

In the year 2000, Swenson et al. (11, 12) published two studies selecting natural microbial communities to increase plant biomass, to degrade an environmental pollutant or to alter the pH of an aquatic ecosystem. Although selected communities occasionally improved over time, they also observed improvements in some control lines and overall, community performance didn't differ significantly from the start of the experiments. Many studies have since followed, selecting for various host effects (13–18), production or consumption of chemicals (19–21) or simply for population size (22, 23). The success of these experiments has also been limited (6, 24), often showing inconsistent results between repeats or only a moderate increase in function.

One fundamental difficulty with artificially selecting communities is that selection is applied at the group level – rather than the individual level as with conventional breeding – with individual organisms going through several generations within each selection round, resulting in little control over ecological and evolutionary dynamics occurring within each community and within each species (5, 25, 26). This means that over time, (i) competition between species may lead to the extinction of slower-growing species that may contribute to community function (6, 27), and (ii) assuming a trade-off between function and growth, competition within species selects for cheater mutants that do not contribute to the function and sweep to fixation (27, 28).

A second problem with the existing approaches lies in how "offspring" communities are generated from their "parents"

at every round: parent communities are either simply diluted to make offspring (low abundant species may go extinct) or pooled together and then distributed over the offspring communities. Both approaches result in offspring communities that are very similar to one another, and do not deviate much from the communities at the start of the experiment [20, 29]. The resulting lack of variability between communities gives little material for artificial selection to work on. The challenge then is to develop a selection method that favors cooperation within and between species, while maintaining between-community variability and selecting for increased function at the community level.

Here we address these fundamental problems by experimentally testing a novel selection approach called "disassembly selection" that was inspired by optimization algorithms from the computational sciences called genetic algorithms [30, 31] and that we have evaluated theoretically [29]. We use a computational algorithm to guide lab experiments in real-time to automatically explore the species composition search space: we randomly generate communities of known species composition, and then repeatedly select the best-scoring communities, disassemble their member species and re-assemble new communities that differ slightly in their composition for the next round. This approach improves performance while maintaining between-community variability [29]. To limit competition within communities and avoid aggressive species that exclude all others, we penalize communities where species extinctions occur.

We use our approach to find a community that can optimally degrade industrial pollutants called Metal Working Fluids (MWFs), a challenge we have previously studied using a single four-species bacterial community [32]. As this original community could only degrade 44.4% of the MWF on average, we hypothesized that there would be room for improvement.

After 18 rounds of selection, we found a four-species community that degraded 75.1% of the MWF on average, significantly better than our original community [32], the best community in the first round, and a random control. Despite this successful outcome, we separately found a species pair that performed at least as well as the top community, suggesting that our approach can still be simplified and improved.

Results

Degradation efficiency increased over 18 rounds.
Briefly, we designed a community selection method where 11 species were first randomly combined into 29 communities of four species each. We let these 29 communities grow for four days, scored them according to their degradation ability (penalizing for species extinctions), "disassembled" the top ten by selective plating, sampled viable cells of each species and used them to rebuild a new round of 29 communities that resembled the best-scoring ones. Resemblance was achieved by either rebuilding the exact same communities as in the previous round – even with the same starting population sizes for each species – or by randomly exchanging one member species in a winning community to introduce some variability and to ensure that all 11 species remained in the meta-community. We carried out this procedure 18 times, with one round per week (Methods, Fig. 1A).

To test whether our selection approach could find communities that degraded better than the random species combinations at the start and that this was due to community-level selection, we included a control treatment where communities propagated to the next round were selected randomly rather than based on their degradation score. We compared the five best-degrading communities in the two treatments ("random" and "selection") at each round. Only considering the top five allows us to exclude the noise introduced by composing new, possibly poorly-performing, communities (Fig. S1). The top five communities in the last round of the selection treatment scored higher than the top five initial communities from both treatments (round 0: 62.28%±4.92 vs. round 18 selection: 73.42%±7.38, Wilcoxon rank-sum test with continuity correction, $df = 15$, $p = 0.012$), and than those in the last round of the random treatment (round 18 random: 63.47%±5.49, random vs. selection $df = 9$, $p = 0.033$). In contrast, the top 5 from the last round of the random treatment did not degrade significantly better than the initial communities (random vs. initial $df = 15$, $p = 0.85$, Fig. 1B).

Throughout our experiment, we tested 167 different combinations of four species (141 in the selection and 156 in the random treatment, with some overlap) out of 174 possible permutations of 11 species (some species combinations were avoided as they were indistinguishable using selective plates, see Methods). The selection treatment tested high-performing communities more often than the random treatment, and this occurred preferentially in the later rounds of the experiment (Fig. 1C, S2), showing selection for improved degradation and the maintenance of high-performing communities. Our approach also continued to explore the search space by testing many new communities at each round: in round 18 there were still communities with low degradation scores (Fig. 1C).

We next asked whether the communities with the highest degradation scores resembled each other in terms of species

Fig. 1. Selection method and its performance. A) Illustration of the selection method (see Methods for details). Each tube represents a community of four species (two colors drawn for illustrative purposes): 1) Define 29 communities of randomly drawn species and inoculate each community in MWF+AA. 2) Following growth, measure degradation score as the difference in pollution load to an abiotic control, illustrated by the gray field at the top of each tube. 3) Select the communities with top 10 degradation scores for disassembly (illustrated by tubes 2 and 5 here) and plate these on selective media to separate their member species. Plating allows to document extinctions and calculate final community scores (combining degradation scores and extinction data). 4) Sample viable cells of each species, only from the corresponding single community with the highest final score. Adjust population size and freeze down. 5a) Generate new communities in proportion to their final community scores. 5b) Randomly choose 21/29 of the new communities (illustrated with 4) for species exchange. Remove one resident species at random and introduce a new species in its place. Assemble the new communities in the lab using the frozen species and repeat from step 2). B) Degradation scores of the 5 best communities in each round for the selection (green triangles) and random (orange circles) treatments, with lines through the average of the 5. C) Community composition (x-axis) vs. degradation score (hue, color bar) for each community over the 18 rounds of selection (y-axis) in the selection (top panel) and random (bottom panel) treatments. The x-axis is ordered by increasing degradation scores (averaged over all instances of the same species composition). Note that these are degradation, not final community scores (extinctions not considered). D) Community composition corresponding to panel C, showing the presence (dark blue) or absence (white/grey) of each species, and illustrating the difference in composition by the Hamming distance (i.e. the number of substitutions needed to transform a given community to another) to the community with the highest degradation score at the bottom.

composition. We calculated the Hamming distance to the best community (number of species that one must exchange in a given community to get the same species composition as the best community), which can also be seen as a measure of the ruggedness of the "fitness landscape" (4). At first glance, there was no obvious pattern between the similarity in community composition to the top community and degradation score. However, the best 5 communities had a distribution of Hamming distances that was significantly different from the distribution of distances between all pairs of communities in our study (Student's t-test, $p = 8.2 \times 10^{-5}$, Fig. S3). It also appears that some species, such as *C. testosteroni* (Ct) and *A. faecalis* (Af) tended to be found in the winning communities, and *P. fulva* (Pf) was rarely in badly-performing communities (Fig. 1D). We explore this more quantitatively below.

Selection reduced extinctions, but did not increase evenness or total biomass. We first explore whether selection has favored certain community properties: population growth, community evenness or species survival. Since the selection method penalizes extinctions (we scaled the degradation scores by the fraction of surviving species), we quantified extinctions in each round by plating 10 communities per treatment (see Methods) on selective media on day 4 and comparing the presence of each species to how we composed the community on day 0. The distribution of extinctions per round was significantly lower in the selection compared to the random control treatment (Kolmogorov–Smirnov test, $p = 0.013$, Fig. 2A, S4, Table S1). As a control, we also counted the number of contamination events (any species that was present at day 4, despite not being inoculated at day 0), which we did not expect to vary significantly between the treatments. Indeed, we found no significant difference in contamination events per round between the two treatments (Kolmogorov–Smirnov, $p = 0.8$). Despite this difference, another explanation could be that fewer extinctions occurred in the selection than the random treatment because selected

Fig. 2. Number of extinctions, evenness and total population size over time. A) Number of extinctions per round (solid lines) and cumulative (dashed lines) in the 10 plated communities of the selection and random treatments. B-C) Mean (lines) $\pm$ SD (shaded areas) values of the 10 plated communities at each round where B) shows evenness (the effective species number divided by its theoretical maximum value) and C) total population size in CFU/ml. D-E) Degradation percent plotted against D) evenness and E) total population size with the selection treatment in triangles and the random treatment in circles, and color representing selection rounds. Population size, growth or evenness could only be calculated for the 10 communities per treatment that we plated (see Methods).

communities more often contained strong growers that promote the survival of others and increase degradation score (we highlight communities lacking strong growers in Table S1).

Next, we ask if communities in the selection treatment were more even than in the random control. We might expect selection to favor evenness, since species in diverse communities may complement one another while communities dominated by a single species risk excluding others that could contribute to degradation. Calculating evenness as the effective species number relative to its maximum value (Methods, Eq. (1),(33)), the evenness of the 10 communities whose populations we quantified increased with time in both treatments (Fig. 2B). The correlation was stronger in the selection treatment compared to the random control (Spearman's $\rho = 0.45$ and 0.24, respectively), but the correlations were not very predictive (ordinary least squares regression between evenness and round, $R^2 = 0.19$, 0.06 at $p = 1.9 \times 10^{-10}$, 7.5×10^{-4} for selection and random, respectively).

Finally, we might expect the total biomass in communities to influence degradation for two reasons: (i) degradation could

be the aggregated effect of individual cells assuming that all species contribute to degradation, and (ii) as species adapt to the medium, they might increase their growth rates, which should increase degradation. We calculated the total population size on day 4 per community at each round of selection, but found no significant effect of total biomass or selection treatment on degradation: Total biomass did not correlate strongly with time (Spearman's $\rho = 0.04, 0.07$, for selection and random, respectively, Fig. 2C) and was not significantly different between the two treatments. Indeed, degradation score did not even correlate with total biomass (Fig. 2E, Spearman's $\rho = -0.0062$, $p = 0.904$).

In sum, selection seems to have favored communities whose members are less likely to drive each other extinct, but no other community features could explain the increase in degradation scores or the difference between treatments.

Successful communities were composed of good degraders, their facilitators and freeriders. Noticing that certain species were often found in the winning communities, we next explored which species features were selected

Fig. 3. A) Species representation and corresponding percentage of degradation in the last 5 rounds of the evolution experiment. As both measures can be quantified in percent, we display them on the same y-axis. The dashed line represents the average frequency at which we expect to see a given species in the last 5 rounds by chance, and the shaded area one standard deviation away from that average. Points that are outside the shaded area are more or less represented than expected by chance. The violin plots show the degradation scores of communities containing that species. B) Degradation percent on day 3 in monocultures, co-cultures, top communities and 11 species together, using species taken from ancestral strains or strains isolated at the end of the random or selection treatment. Data-points are ordered according to the average degradation % and interesting cases are highlighted with a colored background and arrows corresponding to data shown in panels D and E. C) Experiment to determine whether Ac might be a "free-rider". Boxplots show the distribution of degradation scores, while dots show population sizes ($\log_{10}$CFU/ml, right-hand y-axis) of different species at day 3 of co-cultures as indicated on the x-axis. Ac reduces the degradation score of the communities it is in, or increases their variance. D) Matrix of degradation percentage in mono- (diagonal elements) and co-cultures of ancestral strains only (average of dots in panel B). E) Matrix of population sizes ($\log_{10}$CFU/ml) in mono- (diagonal elements highlighted with white squares) and co-cultures of ancestral strains only. In panels B, D and E, we highlight interesting cases in blue and light green that are further discussed in the text.

and whether community degradation scores depended on the presence of specific species or species combinations.

First, we analyzed which species were over- or under-represented in the meta-community compared to what one would expect by chance. For each treatment, we quantified how often each species appeared among plated communities in the last 5 rounds of the experiment ($n = 10$ at each round). If a species' frequency was more than one standard deviation above or below the frequency one would expect by chance (18.18), we designate it as over- or under-represented, respectively (mean$\pm$SD$= 18.18 \pm 11.8$; Fig. 3A dashed line, shaded area). Over-represented species were: Ct, Af and Ac in the selection treatment, and Pf and Pr in the random treatment, while Ml and At2 were under-represented in the selection treatment and Ac in the random treatment. The communities that contained the over-represented species tended to be associated with high degradation scores (Fig. 3A).

The best-scoring community in the selection treatment (At1+Ct+Af+Ac) contained all 3 over-represented species, which partially explains their over-representation. However, it does not answer how its member species were contributing to the score. High degradation in these communities could either be due to single species degrading well, or to synergistic effects between the species. To find the answer, we grew all 11 species alone and in most pair-wise co-cultures and ranked them from best to worst degradation. We included four of the best 4-species communities and all eleven species grown together, as a reference (Fig. 3B). We observed a wide variation in degradation abilities, and to our surprise, all 11 species together ranked 35th (dashed line in Fig. 3B), which is well below what even single species could achieve.

The best individual degraders were Pf, Ct and At1 (mean degradation: 66%, 58% and 50% respectively), while Af, Ea and At2 were the worst (mean degradation: 2%, 2% and 4% respectively, Fig. 3B, D). Interestingly, Af which is one of the worst degraders, was present in many winning communities.

This may be because when combined with At1, it achieves one of the highest degradation scores (Fig. 3B, D blue highlight). Compared to their growth in monoculture, At1 promoted the growth of Af by more than 3 logs, although Af reduced the growth of At1 (Fig. 3E blue highlight).

Surprisingly, not all good degraders were over-represented in the selection treatment. Pf and At1 each featured in only 2 of the 10 best communities (Fig. 1D), despite Pf being the best-performing species alone and featuring in 8 of the 10 best pairs (Fig. 3B, D). In contrast, Ct was present in 7 out of the 10 winning communities (Fig. 1D).

We also analyzed which species were most present when extinctions occurred, as we selected against extinctions. In the 20 communities of the selection treatment where extinctions occurred, the species most often found were At1, Pr and Pf (13, 12 and 12, respectively, Fig. S4). In 9 of the 20 communities, At1 and Pr were both present, which may explain why they do not feature together in the best 10 communities (Fig. 1D), despite being one of the best degrading pairs (Fig. 3B, D blue highlight). In contrast, only 2 extinction events occurred in the selection treatment when Ct was present. Even if At1 was often associated with extinctions, it greatly increased the growth of Af and Pr, resulting in the best-degrading pairs (Fig. 3B, D blue highlights), of which one (At1+Af) was present in the winning community.

The third over-represented species is Ac, which on its own was one of the worst degraders (mean degradation: 11%). And although its degradation improved greatly when together with Pf and Ct (mean degradation: 50% and 46%, respectively), these degradation scores were lower than what Pf (mean degradation: 66%) and Ct (58%) could achieve alone. Interestingly, Ac's growth was also significantly promoted by the three degrader species (Fig. 3E, Ac row, light green highlight) and while it did not reduce the growth of the degraders much, it greatly reduced their capacity to degrade, particularly for At1 (mean degradation: 13% as opposed to 50% when grown alone). These results suggest that Ac may have acted as a "free-rider" species that got carried along with the best communities. We tested this idea by removing Ac from the winning community and observed a reduced variability in its performance (Fig. 3C). Removing it from the fourth-best community also significantly increased the community degradation score (degradation of Ct+Af+Ac+Ea: 30.2±4.5%, vs. Ct+Af+Ea: 47.3±3.5%, t-test, $p = 0.0012$), whereas removing Ct from this community drastically reduced its degradation score (Af+Ac+Ea: 4.7 ± 1.2%, $p < 0.001$, Fig. 3C).

In sum, selection appears to have favored communities with at least one good degrader species, especially if its score could be enhanced by "weaker" species, as long as they did not cause extinctions. This approach does not seem to eliminate free-riders that appear in the final communities despite their deleterious effects on degradation scores.

Did the best communities improve compared to their ancestors? Up to this point, we have viewed our disassembly selection approach as a way to recombine different species in our original set in a way that increases the degradation scores. As bacteria undergo many cell divisions over the course of the experiment, however, we may also expect within-species genetic evolution to have modified the participating species themselves compared to the ancestral strains we started with. Within-species selection may act to increase growth rates, for example, or to reduce antagonistic genotypes that would cause the extinction of other species in the community, bringing down community score.

To determine whether the species at the end of the experiment grew or degraded differently from their ancestors, we compared the 11 species isolated from the different treatments to their ancestors, measuring their degradation and population sizes (Fig. 4A, B). We found that Ct isolated from the selection treatment grew significantly better than its ancestor ($3.8 \times 10^8 \pm 2.2 \times 10^8$ vs. $3.3 \times 10^7 \pm 4.1 \times 10^7$, Wilcoxon rank sum, $df = 79$, $p = 4.72 \times 10^{-16}$) and than its counterpart isolated from the random control treatment ($9.92 \times 10^7 \pm 8.68 \times 10^7$, $df = 71$, $p = 1.1 \times 10^{-12}$). The strain of Ct from the selection treatment also degraded significantly worse than its ancestor (51.7±5.2 vs. 56.4±5.5%, $df = 79, p = 0.00054$), suggesting that it may have evolved to invest more into biomass and less into degradation, but we do not explore this idea further. Other than that, no significant effects were observed.

We were also curious to see whether inter-species interactions had changed throughout the selection treatment. Indeed, we had chosen to conduct this experiment in growth medium containing casamino acids (see Methods), as we knew from previous work that competition was stronger in this environment compared to MWF without casamino acids (32) and we wondered whether selection could reduce competition. We first used the population size data to estimate these interactions in the ancestral species (Fig. 4C), where we confirmed that competition was common. We also confirmed previous findings that species that could not grow well alone tended to be affected positively by others, particularly by strong growers (Fig. 4C, 3E) (32, 34).

We then selected a few pairs from the random and selec-

Fig. 4. Effect of within-species evolution. A) degradation percent or B) population size ($\log_{10}$CFU/ml) of each species at day 3 in three conditions: the ancestral strains before the experiment, strains harvested after 18 rounds of selection treatment (S) and strains after 18 rounds of the random control treatment (R). Data from mono- (alone) and pairwise co-cultures are shown (with partner). Significant differences are calculated using a generalized linear model with biological replicate as random variable and number of species in culture as an explanatory variable, significant p-values with a Bonferroni correction for multiple comparisons are shown. C)-E) Interactions between ancestral species (C), the species evolved in the random treatment (D) and the selection treatment (E) defined as the $\log_2$ fold-change of the focal species in CFU/ml of day 3 in co-culture with the companion species vs mono-culture. Interactions that were not significant (no significant difference between growing alone or with companion species) are shaded. Positive (facilitative) interactions are in blue, while negative interactions are shown in red. White squares are ones that we did not measure. Overall, we saw very few changes between ancestral and evolved species.

tion treatment isolates for which we conducted mono- and
co-cultures to estimate the interactions between the evolved
species. While some interactions differed after evolution
(Fig. 4D, E), we found little evidence that competition had
been weakened, and more generally, no overall pattern. One
exception was Ct isolated from the selection treatment, which
was no longer inhibited by any species. The reduction in
competition can be explained by our finding that it could
grow better alone (Fig. 4B). We therefore conclude that evo-
lution at this timescale has not had profound effects on the
species' phenotypes.

Discussion

Previous community selection experiments have struggled to
show consistent improvements in community functions com-
pared to controls (6, 24). We have now devised and tested

a selection method to improve the degradation of MWF pol-
lutants in small synthetic bacterial communities. The disas-
sembly method automatically searches for species combina-
tions with high degradation scores, while selecting against
species that cause the extinction of other community mem-
bers. The best community found using this approach per-
formed significantly better than the best initial communities
(Wilcoxon rank sum test, $p < 0.001$) and 69% better than the
community studied in our previous work (Fig. S5, Wilcoxon
rank sum test, $p = 0.007$) (32).

Further investigating some of the top communities and many
species pairs revealed that successful degradation could be
achieved by combining strong degraders with other species
that might not be able to survive alone, but enhanced the
degradation score when paired with the strong degrader. This
simple heuristic revealed that the best overall performance

was achieved by two species in co-culture: At1 and Af. Adding two more species to this pair (Ct and particularly the free-rider Ac) increased the variance in community performance, and combining all 11 species performed particularly poorly (Fig. 3). It appears then that our optimal community of four species is in fact too rich.

Another key observation is that despite our efforts to favor within-species evolution – we sampled many colonies when disassembling communities through plating to include sufficient within-species diversity, and used a competition-promoting medium to give room for interactions to evolve to become less negative or more positive – it did not have a large effect on final population sizes, degradation abilities or inter-species interactions. One explanation may be that species are changing their biotic environment too often for selection to favor any particular interactions. In agreement with this, the only change we observed is that Ct evolved in the selection treatment grew better than its ancestor alone, and was less negatively affected by others than its ancestor (Fig. 4). It could be that Ct evolved to invest less into degradation and more into growth, although we currently have no evidence to back this up. The up-side of finding only minor changes is that one may not need to be too concerned that species will evolve to become more competitive or invest less into community function, at least on this time-scale.

Given what we have learned, would we now perform artificial selection differently? After all, the approach was quite cumbersome and would not be easy to set up for a new problem.

A first question is whether the artificial selection approach is useful at all, or whether we could have predicted the composition of the best community using fewer culture experiments. To explore this question, we performed an additional analysis using a simple linear model that predicts the degradation score based on species presence/absence (4). Including the data from all our experiments, the linear model had a reasonable fit ($R^2 = 0.75$) and would have chosen a community that performed relatively well (degradation score: 69.2%). However, if we only used mono- and co-culture data to fit the model, performance dropped ($R^2 = 0.26$) and the best predicted communities ranged in performance from 40.3% to 83.1% (Fig. 5A, B). This analysis suggests that the "fitness landscape" of MWF-degrading communities is quite rugged, in line with the non-uniformity of the Hamming distance analysis in Fig. 1D. It would be interesting to determine the minimal amount of data needed to achieve a good prediction and explore whether other prediction methods would perform better (e.g. (35)).

A second lesson could be to focus on the strength of our method as a search algorithm to efficiently explore the space of possible species combinations, analogous to a genetic algorithm (30, 31). As allowing the species to evolve within these communities did not seem to change much, it would be simpler to start communities at every round from frozen stocks and avoid the challenging experimental step of disassembling communities (as illustrated in Fig. 5C). We would then no longer need selective media for all species in the pool and it would suffice to know whether species went extinct, which could be achieved through amplicon sequencing. Another important modification would be to allow community size to change, as opposed to restricting it to four species as we have done here. This would involve removing or adding species independently, allowing communities to grow or shrink in size. This decoupling would increase the search space of possible combinations, but might find better solutions, for example by avoiding free-rider species like Ac to establish in so many communities.

Allowing community size to change automatically would also answer an important question for community function: how many species are actually needed to solve the problem of interest? In our previous work (32), a mathematical model predicted that in harsher environments, more species are needed to achieve maximal community function compared to permissive environments. Experimentally, degradation saturated at two species in the more permissive MWF with casamino acids, compared to three species in MWF alone (32), which is consistent with our best solution here having only two species. In hindsight, a more challenging environment might have shown a stronger improvement over the experiment and required a larger optimal community.

A final important limitation of our approach is that we fixed the initial population size of all species at each round, in order to select against cheater strains that grow quickly without contributing, and to improve heritability of the community function (27). It would be interesting to see how our best-performing communities would equilibrate over a few rounds of growth and dilution, when we do not adjust the initial population sizes. It is conceivable that the performance of the community at equilibrium would be different. Ideally, community stability should be part of the community score, although this would require substantial revision to the selection algorithm.

In summary, we have tested an approach to artificially select amongst communities composed of different combinations of culturable species. Our approach found a four-species community that is efficient at degrading MWF pollutants and is

Fig. 5. A-B) Linear model analysis. We use a linear model (code taken from (4)) that uses species presence/absence to predict degradation percent based on A) all the data we generated, or B) only the mono- and co-culture data. Community richness is shown in color. Each dot is one degradation score measurement, such that biological replicates and technical replicates, if available, are all represented. C) Proposing a new artificial selection method. Rather than disassembling communities, we propose to use the winning communities as templates to generate the offspring communities in the next round. These communities would then be seeded by taking the clonal ancestral species from the freezer, such that there would be no within-species evolution over rounds. Step 3) would be to select the top 10 communities, 4a) to generate communities in proportion to their community scores and 4b) to randomly choose 21/29 of the new communities (illustrated with 4) for either species removal or introduction (see white asterisks in step 4, 4a and 4b are shown in one step). Freezer icon created by SAM Designs from Noun Project.

superior to the performance of all species in our pool grown together. However, the selection experiment was relatively complex and a smaller community was also found by testing species pairs and comparing them to the winning community. Going forward, we propose a simpler, more effective approach (Fig. 5C). Even though the challenges of ensuring ecological and evolutionary stability remain open, we argue that this first proof-of-concept supports the blind approach to automate the breeding of bacterial communities with optimal functions.

Methods and Materials

Bacterial species and culture conditions. We used 11 bacterial species listed in Table 1. At1, Ct and Ml were previously isolated from MWF as previously described (32, 36, 37). Note that Ml (*Microbacterium liquefaciens*) was previously referred to as *Microbacterium saperdae* but a more recent classification has led us to refer to it differently. At2 was kindly donated by Justine Collier (plant associated) and the remaining species were isolated from MWF and kindly donated to us by Peter Küenzi from Blaser Swisslube AG. The species were identified at Blaser Swisslube AG by MALDI-TOF, and confirmed by PCR amplification and 16S gene se-

quencing. All experiments were performed in 6ml batch cultures containing 0.5% (v/v) Castrol Hysol™ XF MWF (acquired in 2016) diluted in water with added salts, metal traces (Tables 2, 3), and supplemented with 1% Casamino Acids (Difco, UK). Cultures were incubated at 28°C, shaken at 200 rpm.

Species	Our acronym
Staphylococcus warneri	Sw
Agrobacterium tumefaciens MWF001	At1
Comamonas testosteroni MWF001	Ct
Microbacterium liquefaciens MWF001	Ml
Alcaligenes faecalis	Af
Aeromonas caviae	Ac
Enterococcus avium	Ea
Klebsiella pneumoniae	Kp
Pseudomonas fulva	Pf
Providencia rettgeri	Pr
Agrobacterium tumefaciens C58	At2

Table 1. Bacterial species used in the experiment and the acronyms we use to we refer to them throughout the manuscript.

Compound	Amount
H_2O	1000 ml
K_2HPO_4	6 g
KH_2PO_4	6 g

Table 2. Phosphate solution (1%) for the MWF+AA medium as described in Table 3.

Compound	Amount
H_2O	405 ml
Phosphate solution	60 ml
NaCl 1% solution	60 ml
Casamino 1% acids solution	60 ml
Hutner's vitamin-free mineral base	12 ml
Castrol Hysol 100%	3 ml

Table 3. For 600 ml of MWF+AA medium, mix in the above order, top to bottom. The phosphate solution is found in Table 2. The MWF needs to be added carefully, one drop at a time to allow mixing.

Selective media. We designed 10 selective media that allow the growth of only one or two of the 11 species at a time. Some species combinations (Ct & At2, Af & Ct, Ml & Sw, Ml & Ea, Ac & Pf, Ct & Kp) cannot be easily distinguished on these media, and we avoided combining these species in the communities (Fig. S6A). This means that instead of the 330 combinations of 4 species out of 11, we have 174 possible communities. Our selective media are generally composed of a rich base and at least one antibiotic (details in Tables S2 and S3). The disassembly plates consist of two 24-well plates where we poured 1.5ml of each selective media into 4 wells (as shown in the 24-well templates in Fig. S6B). Because temperature was helpful to distinguish some species, we incubated some media at 28°C and others at 37°C. Disassembly was achieved by plating droplets of each diluted community on all the selective media (more details below). For each round, we prepared the disassembly plates one week in advance and stored them at 4°C in the dark until they were used. Every week, the selectivity of the media was verified by inoculating 10μl droplets from a dilution series of 2-day old cultures of all 11 ancestral species in square plates of all selective media.

Artificial Selection. Each round of the selection experiment lasted one week and consisted of five steps (Fig. 1): (1) assembling communities and letting them grow, (2) measuring pollution load, (3) selecting top communities and disassembling them on agar plates, (4) freezing down species samples, and (5) generating species compositions for the next round.

Community assembly. In each round, we used 60 10ml glass tubes, 29 were assigned to communities of the selection treatment, 29 to the random treatment and two tubes were abiotic controls. The first round started with the same 29 randomly generated communities of 4 species each in the two treatments. These were drawn such that all 11 species were present in at least one community and such that species that we cannot separate with selective plates never appear in the same community.

Communities for the first round were assembled as follows: Single colonies of each of the 11 species were picked and grown overnight in 5mL of TSB at 28°C, shaken at 200rpm. The next day, cultures were adjusted to an OD_{600} of 0.05 in 10ml of PBS in a 15ml falcon tube. For subsequent rounds, similar 15ml tubes containing each of the 11 species for each treatment at OD_{600}=0.05 were taken from the freezer (see below) and thawed. The cells were then washed by centrifuging at 4000rpm for 15 minutes and resuspended in 10ml of MWF+AA medium (see above). For each community culture in the experiment (29 for each treatment) and the abiotic controls, 6ml of MWF+AA were prepared in the 10ml glass tubes and 100μl of each species were added, yielding a total of 400μl of four species of similar relative abundances. All 60 tubes were then incubated at 28°C and shaken at 200 rpm for four days.

Measuring degradation scores. On day 4, as a proxy for pollution load, we measured the chemical oxygen demand (COD) using NANOCOLOR COD tube tests (detection range 1-15 g/l by Macherey-Nagel (ref: 985 038), see (32) for more details. We used these measurements to calculate degradation scores as $(1 - COD_4(sample)/COD_4(control))\times100$, i.e. the COD of the community after 4 days relative to the COD of the abiotic control after 4 days, in percent. Data shown in Fig. 3C was generated using expired COD tubes, which explains why their values are different from those of the other experiments. However, given that the important comparison is between treatments within that experiment, we decided not to repeat it.

Selecting and disassembling top communities. We selected the 10 out of 29 communities with the highest degradation scores from the selection treatment and 10 out of 29 communities at random from the random treatment. To disassemble the communities and determine species' population sizes, we plated dilutions ($10^{-1}, 10^{-2}, 10^{-4}$ and 10^{-6}) of each community onto all selective media (see above, Fig. S6), incubated the selective plates for two days (either at 28°C or 37°C), and counted colony-forming units (CFUs) for each species. This allowed us to disassemble all community mem-

bers, estimate population sizes and identify extinction and contamination events (species that were inoculated on day 0 but did not appear on their selective media, and species that were not inoculated in a given community but grew on selective media, respectively). We penalized extinction by scaling the degradation score of each community by the fraction of surviving species (contaminants are not counted) $0 < f < 1$ (e.g. $f = 0.5$ if only two of the four inoculated species are detected). The final community score was then calculated as $(1 - \text{COD}_4(\text{sample})/\text{COD}_4(\text{control})) \times 100 \times f$.

Freezing down species. At every round of selection, we froze down a representative of each species by isolating it from the highest-scoring community where that species was present. We sampled several CFUs from the highest dilution in the relevant selective plate by adding PBS to the selected well and re-suspending by pipetting. We then adjusted the OD_{600} of the samples to 0.05 in a total volume of 10 ml of PBS with 25% glycerol, then aliquoted $2 \times 1\text{ml}$ for long-term storage in cryo tubes, 3ml for use in the following round and 5ml as backup in 15ml falcon tubes and froze all samples at $-80°\text{C}$. If a species went extinct in a round of selection, we recovered it from its frozen stock collected in a previous round.

Generating new species compositions. For the following round of selection, we used a script that calculates a probability distribution from the community scores of the 10 disassembled communities and generates offspring communities by randomly sampling 29 times with replacement in proportion to this distribution. Communities with higher scores are more likely to be selected. In the random control, we sampled 29 times with uniform probability from the 10 disassembled communities.

To introduce variability into these newly generated communities, out of the 29 generated communities in each treatment, we randomly chose 21 to receive an invader species that replaced one of the four members. Both the invader and the species to be removed were chosen by uniform probability, with a few exceptions: We first chose as invaders species that were not yet represented in any offspring communities, adding them to random receiving communities; once all 11 species were represented at least once in the new communities, we chose the remaining invaders at random but avoided invading species that were already present in the receiving community, and species that are indistinguishable from resident species on our selective media. Selection and invasion thereby result in 2×29 lists of four species each, sampled in proportion to degradation scores (or not for the random treatment) and with 21/29 of them having exchanged an

old community member for a new one. We then assembled the communities in the lab from the frozen species record as described above. The script used to automatically generate offspring communities is written in python 3 (38) and can be found at https://github.com/Mitri-lab/disassembly_selection_experiment.git.

Comparing ancestral and evolved strains. Following the artificial selection experiment, we conducted follow-up experiments to better understand why the selection algorithm favored certain species combinations. For each species, the frozen stocks from round 18 of the selection and random treatments were plated and incubated. Single colonies were picked and grown overnight in 5mL of TSB at 28°C, shaken at 200rpm. The next day, cultures were adjusted to an OD_{600} of 0.05 in 10ml of TSB and grown for a further 3h. The cells were then washed at 4000rpm for 15 minutes and re-suspended in 10ml of MWF+AA medium. For each culture, 6ml of MWF+AA were prepared in 10ml glass tubes and $100\mu l$ of each species were added. These cultures were incubated at 28°C, shaken at 200rpm for 3 days. CFUs were measured through serial dilution and plating on days 0, 1, 2 and 3 using the appropriate selective media (Fig. S6). We measured CODs of an abiotic control culture at day 0 and 3, and the culture tubes at day 3. The degradation scores were calculated as before. To estimate interactions between species, we grew each strain alone or with a given partner strain and compared the population size of each focal strain as the $\log_2$ fold-change in CFU/ml on day 3 in the presence or absence of the partner species. CFU/ml were quantified on selective media (Fig. S6), but on round agar plates considering all dilutions, giving higher resolution compared to the selection experiment. We used LB agar for Af and Pf instead of their selective media, as requirements were less stringent (we only need to count them, not disassemble them) and there appeared to be differences in growth between the ancestral and evolved strains on the selective media for those species. These experiments were performed by two different authors (GA and BV), which is accounted for in the statistical analysis (see below).

Data analysis. We used the Hamming distance between two communities to quantify the difference in species composition between them (Fig. 1). The community is in this case represented by the presence and absence of each of the 11 species, and the Hamming distance is the fraction of species mismatches. We used the implementation from the SciPy library in python (39).

We calculated evenness as the effective species number, or Hill number of order 1 ([40]):

$$^1D = \exp\left(-\sum_k p_k \log(p_k)\right), \tag{1}$$

divided by its maximum value (similar to Pielou's evenness ([33])), where p_k is the relative abundance of species k in the community. Ordinary least squares regression between evenness and round was calculated using the python package *statsmodels* ([41]).

We used parametric and non-parametric tests for significant differences between groups, preferring the Student's t-test for the former and the Wilcoxon rank sum test for the latter, and compared distributions using the Kolmogorov–Smirnov test. We measured correlations using Spearman's ρ and quantified regressions using the ordinary least-squares implementation in the python library *statsmodels*, ([41]). When relevant, we corrected for multiple comparisons using the Bonferroni method.

To compare the growth and interactions of evolved and ancestral strains, we took into account that experiments were performed by two different people. To calculate statistical significant differences in growth or degradation, *experimentalist* was taken to be a random factor in a generalized linear model. To calculate interactions (one species growing in mono- versus co-culture), we only used data collected by the same experimentalist. If it was not available (only one person had measured the mono-cultures), we used this instead (see dataset 1).

Contributions and acknowledgements

FAS, SM and BV conceived the project. FAS developed the experimental methods. BV implemented the selection method. AH performed the selection experiment and FAS, BV and GA follow-up experiments. FAS, BV and SM analysed the data. FAS, BV and SM wrote the paper. Thanks to members of the Mitri lab for discussions, especially to Margaret Vogel and Afra Salazar for detailed comments, to Peter Küenzi (Blaser Swisslube) and Justine Collier (UNIL) for bacterial strains. We sincerely thank Samuele E. A. Testa, who trained BV in the lab and helped with experiments. Thanks to Marc Garcia-Garcerà and Bastien Vallat for helping identify the strains fom Blaser.

1. Christopher E. Lawson, William R. Harcombe, Roland Hatzenpichler, Stephen R. Lindemann, Frank E. Löffler, Michelle A. O'Malley, Héctor García Martín, Brian F. Pfleger, Lutgarde Raskin, Ophelia S. Venturelli, David G. Weissbrodt, Daniel R. Noguera, and Katherine D. McMahon. Common principles and best practices for engineering microbiomes. *Nature Reviews Microbiology*, 17(12):725–741, 2019.

2. Bryce M. Connors, Sarah Ertmer, Ryan L. Clark, Jaron Thompson, Brian F. Pfleger, and Ophelia S. Venturelli. Model-guided design of the diversity of a synthetic human gut community. *bioRxiv*, page 2022.03.14.484355, 2022.

3. Alvaro Sanchez, Djordje Bajic, Juan Diaz-Colunga, Abigail Skwara, Jean C.C. Vila, and Seppe Kuehn. The community-function landscape of microbial consortia. *Cell Systems*, 14 (2):122–134, 2023.

4. Abigail Skwara, Karna Gowda, Mahmoud Yousef, Juan Diaz-Colunga, Arjun S. Raman, Alvaro Sanchez, Mikhail Tikhonov, and Seppe Kuehn. Learning the functional landscape of microbial communities. *bioRxiv*, 534159:2023.03.24.534159, 2023.

5. Flor I. Arias-Sánchez, Björn Vessman, and Sara Mitri. Artificially selecting microbial communities: If we can breed dogs, why not microbiomes? *PLOS Biology*, 17(8):e3000356, 2019.

6. Álvaro Sánchez, Jean C.C. Vila, Chang-Yu Chang, Juan Diaz-Colunga, Sylvie Estrela, and María Rebolleda-Gomez. Directed evolution of microbial communities. *Annual Review of Biophysics*, 50(1):null, 2021. PMID: 33646814.

7. Jikun Huang, Carl Pray, and Scott Rozelle. Enhancing the crops to feed the poor. *Nature 2002 418:6898*, 418(6898):678–684, 2002.

8. Keqin Chen and Frances H. Arnold. Enzyme engineering for nonaqueous solvents: Random mutagenesis to enhance activity of subtilisin E in polar organic media. *Nature Biotechnology*, 9:1073–1077, 1991.

9. Sheenam Verma and Dinakar M. Salunke. Directed evolution – bringing the power of evolution to the laboratory : 2018 Nobel Prize in Chemistry. *Current Science*, 115(9):1627–1630, 2018.

10. Janet Karpinski, Ilona Hauber, Jan Chemnitz, Carola Schäfer, Maciej Paszkowski-Rogacz, Deboyoti Chakraborty, Niklas Beschorner, Helga Hofmann-Sieber, Ulrike C. Lange, Adam Grundhoff, Karl Hackmann, Evelin Schrock, Josephine Abi-Ghanem, M. Teresa Pisabarro, Vineeth Surendranath, Axel Schambach, Christoph Lindner, Jan Van Lunzen, Joachim Hauber, and Frank Buchholz. Directed evolution of a recombinase that excises the provirus of most HIV-1 primary isolates with high specificity. *Nature Biotechnology 2016 34:4*, 34(4): 401–409, 2016.

11. William Swenson, Jeff Arendt, and David Sloan Wilson. Artificial selection of microbial ecosystems for 3-chloroaniline biodegradation. *Environmental Microbiology*, 2(5):564–571, 2000.

12. W. Swenson, D. S. Wilson, and R. Elias. Artificial ecosystem selection. *Proceedings of the National Academy of Sciences*, 97(16):9110–9114, 2000.

13. Jigyasa Arora, Margaret Mars Brisbin, and Alexander S. Mikheyev. The microbiome wants what it wants: microbial evolution overtakes experimental host-mediated indirect selection. *bioRxiv*, page 706960, 2019.

14. Samuel Jacquiod, Aymé Spor, Shaodong Wei, Victoria Munkager, David Bru, Søren J. Sørensen, Christophe Salon, Laurent Philippot, and Manuel Blouin. Artificial selection of stable rhizosphere microbiota leads to heritable plant phenotype changes. *bioRxiv*, 2021.

15. Michael D. Jochum, Kelsey L. McWilliams, Elizabeth A. Pierson, and Young Ki Jo. Host-mediated microbiome engineering (HMME) of drought tolerance in the wheat rhizosphere. *PLOS ONE*, 14(12):e0225933, 2019.

16. Ulrich G. Mueller, Thomas E. Juenger, Melissa R. Kardish, Alexis L. Carlson, Kathleen M. Burns, Joseph A. Edwards, Chad C. Smith, Chi-Chun Fang, and David L. Des Marais. Artificial Selection on Microbiomes To Breed Microbiomes That Confer Salt Tolerance to Plants. *mSystems*, 2021.

17. Kevin Panke-Buisse, Angela C Poole, Julia K Goodrich, Ruth E Ley, and Jenny Kao-Kniffin. Selection on soil microbiomes reveals reproducible impacts on plant function. *The ISME Journal*, 9(4):980–989, 2015.

18. Kevin Panke-Buisse, Stacey Lee, and Jenny Kao-Kniffin. Cultivated Sub-Populations of Soil Microbiomes Retain Early Flowering Plant Trait. *Microbial Ecology*, 73(2):394–403, 2017.

19. Manuel Blouin, Battle Karimi, Jérôme Mathieu, and Thomas Z. Lerch. Levels and limits in artificial selection of communities. *Ecology Letters*, 18(10):1040–1048, 2015.

20. Chang-Yu Chang, Melisa L. Osborne, Djordje Bajic, and Alvaro Sanchez. Artificially selecting bacterial communities using propagule strategies†. *Evolution*, 74(10):2392–2403, 2020.

21. Robyn J Wright, Matthew I Gibson, and Joseph A Christie-Oleza. Understanding microbial community dynamics to improve optimal microbiome selection. *Microbiome*, 7(1):85, 2019.

22. Tiffany Raynaud, Marion Devers, Aymé Spor, and Manuel Blouin. Effect of the Reproduction Method in an Artificial Selection Experiment at the Community Level. *Frontiers in Ecology and Evolution*, 7:416, 2019.

23. Tiffany Raynaud, Marion Devers-Lamrani, Aymé Spor, and Manuel Blouin. Community diversity determines the evolution of synthetic bacterial communities under artificial selection. *Evolution*, 76(8):1883–1895, 2022.

24. U.G. Mueller and J.L. Sachs. Engineering Microbiomes to Improve Plant and Animal Health. *Trends in Microbiology*, 23(10):606–617, 2015.

25. C J Goodnight and L Stevens. Experimental studies of group selection: what do they tell us about group selection in nature? *The American naturalist*, 150 Suppl(S1):S59–79, 1997.

26. C J Goodnight. Heritability at the ecosystem level. *Proceedings of the National Academy of

Sciences of the United States of America, 97(17):9365–6, 2000.

27. Li Xie, Alex E. Yuan, and Wenying Shou. Simulations reveal challenges to artificial community selection and possible strategies for success. *PLOS Biology*, 17(6):e3000295, 2019.

28. Guilhem Doulcier, Amaury Lambert, Silvia De Monte, and Paul B Rainey. Eco-evolutionary dynamics of nested darwinian populations and the emergence of community-level heredity. *eLife*, 9:e53433, 2020.

29. Björn Vessman, Pablo Guridi-Fernández, Flor Inés Arias-Sánchez, and Sara Mitri. Novel artificial selection method improves function of simulated microbial communities. *bioRxiv*, page 2023.01.08.523165, 2023.

30. John H. Holland. *Adaptation in natural and artificial systems: An introductory analysis with applications to biology, control, and artificial intelligence.* 1976. .

31. Alexander Lalejini, Emily Dolson, Anya E Vostinar, and Luis Zaman. Artificial selection methods from evolutionary computing show promise for directed evolution of microbes. *eLife*, 11, 2022.

32. Philippe Piccardi, Björn Vessman, and Sara Mitri. Toxicity drives facilitation between 4 bacterial species. *Proceedings of the National Academy of Sciences*, 116(32):15979–15984, 2019.

33. E. C. Pielou. *Ecological diversity.* .

34. Jared Kehe, Anthony Ortiz, Anthony Kulesa, Jeff Gore, Paul C. Blainey, and Jonathan Friedman. Positive interactions are common among culturable bacteria. *Science Advances*, 7 (45):7159, 2021.

35. Juan Diaz-Colunga, Abigail Skwara, Jean C C Vila, Djordje Bajic, and ´Alvaro S ´Anchez. Global epistasis and the emergence of ecological function. *bioRxiv*, page 2022.06.21.496987, 2023.

36. Christopher J van der Gast and Ian P Thompson. Effects of pH amendment on metal working fluid wastewater biological treatment using a defined bacterial consortium. *Biotechnology and Bioengineering*, 89(3):357–66, 2005.

37. Christopher J. van der Gast and Ian P. Thompson. US 8,703,475 B2.

38. Guido Van Rossum and Fred L. Drake. *Python 3 Reference Manual.* CreateSpace, Scotts Valley, CA, 2009. .

39. Pauli Virtanen, Ralf Gommers, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni Burovski, Pearu Peterson, Warren Weckesser, Jonathan Bright, Stéfan J. van der Walt, Matthew Brett, Joshua Wilson, K. Jarrod Millman, Nikolay Mayorov, Andrew R. J. Nelson, Eric Jones, Robert Kern, Eric Larson, C J Carey, İlhan Polat, Yu Feng, Eric W. Moore, Jake VanderPlas, Denis Laxalde, Josef Perktold, Robert Cimrman, Ian Henriksen, E. A. Quintero, Charles R. Harris, Anne M. Archibald, Antônio H. Ribeiro, Fabian Pedregosa, Paul van Mulbregt, and SciPy 1.0 Contributors. SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. *Nature Methods*, 17:261–272, 2020.

40. Anne Chao, Chun-Huo Chiu, Lou Jost, A Chao, C.-H Chiu, and L Jost. Phylogenetic Diversity Measures and Their Decomposition: A Framework Based on Hill Numbers. *Topics in Biodiversity and Conservation*, 14:141–172, 2016.

41. Skipper Seabold and Josef Perktold. statsmodels: Econometric and statistical modeling with python. In *9th Python in Science Conference*, .