Systematic exploration of prokaryotic form I rubisco maximal carboxylation rates

Benoit de Pins[1], Lior Greenspoon[1], Yinon M. Bar-On[1,†], Melina Shamshoum[1], Roee Ben-Nissan[1], Eliya Milshtein[1], Dan Davidi[2,‡], Itai Sharon[3], Oliver Mueller-Cajar[4], Elad Noor[1], Ron Milo[1,*].

[1] Department of Plant and Environmental Sciences, Weizmann Institute of Science, Rehovot 76100, Israel
[2] Department of Genetics, Harvard Medical School, Boston, MA 02115, USA
[3] Migal Galilee Research Institute, Kiryat Shmona 11016, Israel
[4] School of Biological Sciences, Nanyang Technological University, Singapore 637551, Singapore

† Currently: Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA 91125, USA
‡ Currently: Aleph, Tel Aviv-Yafo 6688210, Israel
*Corresponding author: ron.milo@weizmann.ac.il

Abstract

Autotrophy is the basis for complex life on Earth. Its physiological limits are largely constrained by rubisco - the enzyme that catalyzes almost all carbon fixation on the planet. Yet, with only a small fraction of rubisco diversity kinetically characterized so far, the underlying biological factors driving the evolution of fast rubiscos in nature remain unclear. We conducted a high-throughput kinetic characterization of over 100 prokaryotic form I rubiscos, the most ubiquitous group of rubisco sequences in nature, to uncover the determinants of rubisco's carboxylation velocity. We show that phototrophy and the presence of a carboxysome CO_2 concentrating mechanism correlate with faster rubiscos (median rate difference of 3-5 fold). In contrast to previous assumptions, rubiscos originating from α-cyanobacteria exhibit the highest carboxylation rates among form I enzymes (≈ 10 s^{-1} median versus <7 s^{-1} in other groups). Our study ties biological and environmental insights associated with kinetic variation across rubiscos from nature.

Introduction

Biological carbon fixation is the keystone of food production and energy storage in the living world. Over 99% of global carbon fixation is catalyzed by rubisco[1], the most abundant enzyme in the biosphere[2]. Rubisco is mainly divided into four distinct forms (I, II, II/III, and III) and can be found in all domains of life, from plants to algae through autotrophic bacteria and archaea[3]. Within this diversity, form I is by far the most abundant of the four forms: it is used by all plants and cyanobacteria and is responsible for almost all CO_2 fixation in nature[4].

Paradoxically, while being the most abundant, rubisco is probably the slowest (i.e. low maximum carboxylation rate - $k_{cat,C}$) central metabolic enzyme[5,6]. In addition to its slow carboxylation rate, its active site can react with oxygen through a competitive oxygenase activity that ultimately leads to the release of 20-50% of the incorporated CO_2[7]. These properties are puzzling because of

44 rubisco's central role, which would lead us to expect a fast and efficient enzyme. Why has rubisco
45 not evolved to be a more efficient enzyme? An extensive sampling of rubisco's genetic diversity
46 is crucial to grasp the boundaries of its natural kinetic space.
47
Our group has recently developed a novel approach to systematically mine the kinetic space of
natural rubiscos. We use computational methods to select representative rubiscos from the
tremendous sequence diversity space. Gene synthesis is then used to generate expression
constructs encoding many rubiscos for purification and kinetic characterization. In previous work
we showed the feasibility of this approach by exploring form II and II/III rubisco variants[6]. Indeed,
we found an uncharacterized rubisco that has a $k_{cat,C}$ two-fold higher than all previously-known
rubiscos - demonstrating the potential of this approach to stretch the kinetic boundaries of this
pivotal enzyme.
56
In this work, we expand our search to form I rubiscos, which represent 97% of known sequences[6].
Moreover, their immense ecological and sequence diversity[8–10], limited kinetic data, and higher
carboxylation rates in comparison to plant rubiscos[5], make prokaryotic form I variants particularly
interesting for exploring the evolution of fast carboxylating rubiscos. By conducting a first-of-its-
kind large-scale study of uncharacterized prokaryotic from I rubiscos, and leveraging available
meta-data on their sequences, we uncover the main biological and environmental determinants
behind the evolution of fast-carboxylating rubiscos.
64

Results
Large scale survey of prokaryotic form I rubisco
67 To map the natural diversity of rubisco sequences, we performed an exhaustive search for rubisco
68 homologs across the major genomic and metagenomic public databases. A total of 4300 unique
69 sequences were identified as prokaryotic form I rubiscos (see Materials and Methods for more
70 details).
71
By clustering at different sequence identity thresholds and across different rubisco subgroups (as
detailed in the Materials and Methods and Supplementary Fig. 1), we selected 144 rubisco
variants that represent the full spectrum of prokaryotic form I rubisco sequence diversity. Notably,
prior work has so far characterized only 11 prokaryotic form I rubiscos using active site
quantification, a method that allows for precise measurements of turnover rates while considering
the enzyme's activation state (Fig. 1A).
78
In contrast to form II, II/III and III rubiscos which are composed of a ≈50 kDa large (L) subunit
organized as a homodimer, form I rubiscos comprise an additional ≈15 kDa small subunit (S) in
an L_8S_8 stoichiometry. Due to this complex oligomeric structure, recombinant expression of form
I rubiscos is challenging[11]. To improve correct folding in *Escherichia coli*, we coexpressed
rubiscos with the chaperone GroEL-GroES, which is known to help reconstitution of prokaryotic
form I rubiscos[12–14]. In the case of variants originating from β-cyanobacteria, we coexpressed *rbcL*
and *rbcS* together with their cognate chaperone *rbcX* whenever that gene was present in the
operon. In addition, because these variants initially showed critical issues of solubility (with only
a third of them soluble), we screened different homologs of rubisco accumulation factor 1 (Raf1),

a chaperone that mediates the assembly of β-cyanobacterial rubiscos[15–18]. We identified one
homolog, from the bacteria *Euhalothece natronophila*, whose co-expression nearly doubled the
number of solubly-expressed β-cyanobacterial rubiscos (Supplementary Fig. 2).
91
The carboxylation rates of each expressed rubisco were determined using a modified version of
the spectroscopic coupled assay reported at Davidi *et al.*[6]. Here we directly assayed the crude
cell lysates without purifying the enzyme in order to best preserve the quaternary structure of form
I rubiscos. The method allows determining the specific carboxylation rate even without purification
thanks to the use of CABP (see Methods and Supplementary Note 1). Since the assay is likely
saturating the enzyme for CO_2, measured rates should approximate $k_{cat,C}$ values as suggested by
the good match between the rates of five rubisco variants measured in our lab and their previously
published values (Supplementary Fig. 3).
100
101 Out of 144 rubisco variants tested, 112 were successfully expressed and soluble. Of which, 98
102 exhibited significant catalytic activity (which we define as >0.5 reactions per active site per
103 second). The median rate among active form I rubiscos was 5.4 s^{-1}, similar to the median $k_{cat,C}$ of
104 plant rubisco literature values (4.7 s^{-1} when corrected to 30°C by assuming a Q10 value of 2.2[19]),
105 with the fastest rate being 14 s^{-1} (Fig. 1B).
106
107 Altogether, our measurements complete a nearly tenfold increase in the number of kinetically
108 characterized prokaryotic form I rubiscos. As described below, our results allow for the discovery
109 of features associated with high-performing form I prokaryotic rubiscos.
110
111

114 **Fig. 1. Systematic exploration of the diversity of prokaryotic form I rubisco.** *A) Number of prokaryotic*
115 *form I rubisco variants with a carboxylation rate reported across the literature and in this study. B) Histogram*
116 *of the carboxylation rates measured in this study and across the literature.*
117
118

Form I rubiscos from phototrophic bacteria are faster carboxylases than those from chemoautotrophs

Form I rubisco-expressing bacteria are facultative or obligate autotrophs: they can convert oxidized inorganic carbon (CO_2) into the reduced organic compounds forming their biomass. To fuel the energy-intensive reactions involved in carbon fixation, autotrophs can draw upon two different energy sources: light (photo-autotrophy) or chemical reactions (chemo-autotrophy). We collected information, when available, on the bacteria expressing the studied rubiscos and classified them into phototrophs and chemotrophs, to investigate whether one of these trophic modes was linked to higher carboxylation rates. To refrain from selection biases, we aimed to uniformly sample rubiscos from both classes (see Materials and Methods). We find that rubiscos originating from phototrophs have a carboxylation rate of 6.5 s^{-1} [4.4-7.9 s^{-1}] (median and interquartile range), three times faster than rubiscos derived from chemoautotrophs (2.0 s^{-1} [1.3-3.9 s^{-1}]) (Fig. 2A, Mann-Whitney U test, $p < 0.01$).

Potential confounding variables (e.g., phylogeny), limit our ability to indicate causality. Yet this result could serve as a basis to investigate if phototrophic bacteria may harbor faster carbon fixation as part of coping with the sun's high rate of energy delivery. In particular, phototrophic bacteria typically inhabit environments with fluctuating light intensity, which may require faster metabolic responses to adapt to changes in light availability. It is possible that rubisco adapted to this need.

139
140
141 ***Fig. 2. Large-scale analysis of the biological parameters associated with fast carboxylating rubiscos***
142 *Box and cumulative distribution plots of rubisco carboxylation rates from different clusters: chemo- and*
143 *phototrophic bacterial rubiscos (A), carboxysome-associated rubiscos and their counterparts (B), α- and β-*
144 *cyanobacterial, and carboxysome-associated proteobacterial rubiscos (C). To ensure unbiased study of*
145 *every group, we selected 20 and 9 class-representative chemo- and phototroph-associated rubiscos, 9 and*
146 *9 class-representative carboxysome-associated and non-associated rubiscos, and 15, 19 and 20 class-*

152
153

Carboxysome-associated form I rubiscos are significantly faster

Another biological parameter that could likely have influenced rubisco evolution is the presence
of a carboxysome-based CO_2 concentrating mechanism (CCM). This cellular mechanism
combines the active transport of inorganic carbon into the cell and the colocalization of carbonic
anhydrase and rubisco inside subcellular proteinaceous microcompartments called
carboxysomes, locally increasing CO_2 concentration around rubisco[20]. High CO_2 levels inhibit
oxygenation by competitive inhibition, which might permit the use of faster, but less CO_2-specific,
rubisco variants.

To compare carboxysomal and non-carboxysomal form I rubiscos, we uniformly sampled rubiscos
from each class and compared their measured carboxylation kinetics (see Materials and
Methods). We found that with a median catalytic rate of 7.4 s^{-1} [5.2-8.3 s^{-1}], carboxysome
associated rubiscos are more than 5 times faster than their non-carboxysomal counterparts (1.3
s^{-1} [1.1-2.1 s^{-1}]) (Fig. 2B, Mann–Whitney U test, $p < 0.01$).

Carboxysomes likely evolved during the Proterozoic eon - in the context of the rise of oxygen and
the decrease of carbon dioxide in our atmosphere[20] - to maintain carboxylase activity in this
changing atmosphere. One evolutionary strategy is the emergence of carbon-concentrating
mechanisms (CCMs), including carboxysomes, that locally concentrate CO_2 around rubisco and
therefore maintain local gas concentrations favorable to carboxylation. Another strategy consists
of the evolution of rubisco towards stronger affinity for CO_2. It has long been postulated that
rubisco is a constrained enzyme, limited by a catalytic tradeoff between its carboxylation rate and
affinity for CO_2 (illustrated by the positive correlation between $k_{cat,C}$ and $K_{M,CO2}$)[21,22]. Following this
hypothesis[23], rubiscos with stronger affinity for CO_2 tend to exhibit lower carboxylation rates. In
contrast, rubiscos which evolved in the context of a CCM with less pressure towards stronger CO_2
affinity, should present higher carboxylation rates. Our findings support this conjecture using the
most comprehensive kinetic sampling to date.

Alpha-cyanobacteria express the fastest form I rubiscos across the tree of life

Carboxysomes are expressed by some proteobacteria (like the thiosulfate oxidizer
Halothiobacillus neapolitanus or the purple sulfur bacteria *Thiorhodococcus drewsii*) and by all
cyanobacteria. Cyanobacteria can be divided into two sub-clades, α- and β-cyanobacteria, based
on the monophyletic grouping of their rubisco and carboxysome genes[24,25]. It has been posited
that α- and β-cyanobacteria rubiscos had identical catalytic rates[26] or even, that β-cyanobacteria
rubiscos are faster[22], as they include the fastest form I rubisco characterized up to now (from
Synechococcus elongatus PCC 6301)[23]. However, such statements were made from scarce
measurements, especially among α-cyanobacterial enzymes as only two $k_{cat,C}$ values are
currently available[27,28]. We now reevaluate this hypothesis using a wider kinetic sampling of these

different rubisco subforms. We have class representative subsets of 15, 19 and 23 variants
uniformly covering α-, β-cyanobacteria, and proteobacteria carboxysome-associated rubiscos
diversity from our dataset (see Materials and Methods).
195
196   We find that, in contrast to previous statements, α-cyanobacterial rubiscos show the highest
197   carboxylation rates (9.8 s^{-1} [8.6-10.5 s^{-1}]) among all prokaryotic form I rubiscos, ≈50% higher than
198   β-cyanobacterial rubiscos and their proteobacterial counterparts (6.3 s^{-1} [5.6-7.5 s^{-1}] and 6.6 s^{-1}
199   [4.7-7.5 s^{-1}] respectively, Fig. 2C, Kruskal-Wallis test followed by Dunn multiple comparison test,
200     $p < 0.001$).
201
When oxygenic photosynthesis emerged in an ancestor of cyanobacteria ≈3 billion years ago[29,30],
it alleviated the dependence of these autotrophs on relatively scarce inorganic reductants and
probably contributed to the emergence of faster rubiscos. However, we do not find β-
cyanobacterial rubiscos to be faster than carboxysome-associated proteobacterial counterparts,
in contrast to α-cyanobacterial rubiscos. One possible difference between α- and β-cyanobacteria
could be their cell size. Indeed, α-cyanobacteria are known to encompass many members of the
so-called picocyanobacteria, the smallest cyanobacteria on Earth[31,26]. We therefore collected from
the literature the size dimensions of cyanobacteria, when available. With a median cell volume of
0.5 μm^3 (0.3-1.0 μm^3), α-cyanobacteria are more than 25 times smaller in volume than their β
counterparts (13.6 μm^3 [3.9-25 μm^3]) (Supplementary Fig. 4, Mann–Whitney U test, $p < 0.0001$).
Smaller cells offer higher surface-to-volume ratio and increased exchange with the medium[32],
which could support a higher supply of nutrients, and contribute to the evolution of faster rubiscos
in these specific cyanobacteria (see Supplementary Note 2 for more details).
215

Discussion

217   Rubisco is among the enzymes that have probably shaped Earth's biosphere and geosphere the
218   most. Its kinetic parameters result from billions of years of evolution, following (and causing)
219   changes in the atmosphere and climate. We present here a systematic large-scale survey of ≈140
220   rubisco variants covering the genetic diversity of prokaryotic form I rubiscos.
221
222   We note that the present study is limited by the risk of underestimating some rates if, for instance,
223   enzymes are partially denatured in the expression conditions used. Yet, a denatured enzyme
224   probably could not interact with RuBP or CABP, and more generally, we have no reason to
225   presume *a priori* a systematic bias affecting some of the studied groups over the others.
226
We find that on average, form I rubiscos from phototrophic bacteria are three times faster than
those from chemotrophs. A possible explanation could be that phototrophs tend to benefit from a
higher energy flux than chemotrophs, thus more easily driving the ATP-dependent reactions of
the Calvin-Benson-Bassham cycle. However, such a case mostly represents oxygenic
photosynthesis (cyanobacteria) which is further shown to be associated with the fastest rubiscos
in our work. The small number of rubiscos associated with anoxygenic photosynthesis in our
dataset prevents us from better studying this specific trophic mode. The latter is associated with
different situations and constraints, and most likely provides lower energy levels to cells. Our
previous survey[6] spanning the diversity of form II and form II/III rubiscos mostly identified one

group of fast carboxylating rubiscos, emerging from microaerophilic chemotrophs. The very small number of phototrophs in this previous study - absence of oxygenic phototrophs (exclusively composed of cyanobacteria) and low number of anoxygenic phototrophs - made this kind of comparison impossible with that data source.

In the context of significant shifts in oxygen and carbon dioxide levels since the Great Oxygenation Event, biophysical CCMs like carboxysomes evolved within some autotrophs, allowing for maintenance of high CO_2 levels around rubisco. We find that carboxysome-associated rubiscos are, on average, more than 5 times faster than others. Among the main parameters examined in this study, the presence of a carboxysome exhibits the strongest association with the occurrence of fast carboxylating rubiscos (Supplementary Fig. 5). This finding underscores the constraints imposed by CO_2 levels on rubisco carboxylation rate, as lowering this pressure appears to reliably facilitate the development of higher carboxylation rates. A compelling example within our dataset is the gammaproteobacteria *Hydrogenovibrio kuenenii*, which expresses two form I rubiscos. The carboxysomal rubisco has a rate of 8.3 s^{-1}, which is twice as fast as its non-carboxysomal counterpart (4.2 s^{-1}, Supplementary Data 1), exemplifying the major impact of carboxysome presence on rubisco carboxylation. Although we have not collected data regarding rubisco's Michaelis constant for CO_2 ($K_{M,CO2}$) in this study, the high CO_2 levels inside carboxysomes likely allow for fast but weaker-affinity rubiscos to evolve, in line with the hypothesis that a tradeoff exists between these two kinetic parameters[20].

We further analyzed the correlation between rubisco carboxylation rate and various biological and environmental parameters such as bacterial environmental source, rubisco subtype, bacterial halotolerance, pH, oxygen-sensitivity, or optimal growth temperature. As presented in Supplementary Fig. 6-11, these showed weaker or no correlations.

Overall, rubisco carboxylation rates seem to correlate with a select few environmental parameters, providing insights into autotrophic metabolism constraints. The fitness of autotrophic organisms largely depends on carbon fixation rates. This process relies on three necessary inputs: energy, water, and CO_2. When water and energy are highly abundant (aquatic phototrophs), the main limiting factor is the flow of carbon into the metabolic network. CO_2 needs to first reach the cell membrane, diffuse to the active site of rubisco, and successfully be involved in a catalytic event. Small cells (increased exchange) with a carboxysome (higher probability of catalysis) are therefore the ones most limited by the carboxylation rate of rubisco, as all other bottlenecks are relatively smaller in that case. The fact that the fastest rubiscos found in our study were from α-cyanobacteria, the smallest carboxysome-expressing oxygenic phototrophs in the microbial world, is consistent with this rationale and could be related to the finding that they account for the majority of cyanobacteria globally[26].

This study provides a systematic exploration of prokaryotic form I rubisco maximal rates and its relationship with various contextual factors that could have shaped the evolution of this most abundant enzyme on Earth. Gaining understanding of this dynamic process is crucial for predicting the evolution of autotrophs in a changing climate and the ability to engineer it for human needs.

Acknowledgments

We thank Yoav Peleg, Ron Sender, Noam Prywes, Ralf Steuer, Avi Flamholz and David Savage for important conversations and productive feedback on this manuscript. We thank Michelle Gehring for additional information about unpublished data from their laboratory. This research was supported by the Mary and Tom Beck Canadian Center for Alternative Energy Research, Miel de Botton, the Schwartz Reisman Collaborative Science Program, and the Charles and Louise Gartner Professorial Chair.

Materials and Methods

Rubisco sequence collection

Rubisco large subunit sequences were collected from (i) the NCBI's nr database[33] downloaded in December 2020 and searched following the method described in Davidi *et al.*[6]; (ii) in-house assemblies of the 244 samples from the Tara Oceans expedition[34]; (iii) assemblies and rubisco sequences published by [35–39]. Sequences outside the length range of 300-700 amino acids were removed. The remaining sequences were then clustered at an 80% sequence identity threshold using USEARCH algorithm[40]. Cluster representatives were aligned using MAFFT (v7.475, default parameters)[41], and columns with more than 95% gaps were removed using trimAl (v1.4.rev15, -gt 0.05)[42]. A phylogenetic tree was constructed using FastTree (v2.1.10, default parameters)[43]. To identify the different rubisco forms, we relied on annotated sequences from NCBI, Tabita *et al.*[44], and Banda *et al.*[39]. This process resulted in a total of 72,395 sequences, including 56,161 form I rubiscos, and most notably for this study, 4,302 non-eukaryotic form I rubiscos. The latter were further re-clustered at 90% identity using USEARCH algorithm (custom python script; see below), and a phylogenetic tree was constructed using RAxML[45].

Rubisco variants selection for characterization

Form I rubiscos are divided into 5 separate groups[9,38,46–48] (see Supplementary Fig. 7A): forms IA and IB (forming the "green" type, found in cyanobacteria and some proteobacteria); forms ICD and IE (the "red" type, found in proteobacteria); and the recently discovered form I "Anaero" (found in bacteria related to anaerobic, thermophilic Chloroflexaeota and Firmicutes)[49]. To comprehensively sample the sequence space of form I rubisco diversity, an iterative clustering approach of the large subunit gene was employed using USEARCH algorithm. The resulting representative sequences from each cluster were selected for characterization. Initially, 32 rubiscos were chosen to cover the entire diversity of form I rubisco at a threshold of 75% identity. Subsequently, further clustering was performed on smaller groups of rubiscos of particular interest, with increasing threshold percentages. Throughout the study, this successively included 38 rubiscos representing the diversity of form IA and B rubisco at 85% identity, 13 rubiscos representing the diversity of cyanobacterial rubisco at 88% identity, 31 rubiscos representing the diversity of IB rubisco at 91% identity, 20 rubiscos representing the diversity of cyanobacterial IA rubisco at 97.5% identity, and 23 rubiscos representing the diversity of proteobacteria carboxysome-associated rubisco at 90% identity (See Supplementary Fig. 1). These representative sequences could sometimes overlap, resulting in a total of 129 different rubisco sequences tested in this study. Additionally, 15 rubiscos were arbitrarily selected for setting-up the experimental pipeline. In total, 144 different rubisco variants were selected for characterization in this study.

Gene synthesis

For each chosen rubisco variant, the complete rubisco operon, encompassing rubisco large and small subunit genes, as well as the chaperone rbcX gene for IB rubiscos, was retrieved. The operons were then codon-optimized for expression in *E. coli* (Twist Codon Optimization tool) and synthesized by Twist Bioscience. Following synthesis, these operons were cloned into a pET-29b(+) overexpression vector (NdeI_XhoI insertion sites). Validation of gene synthesis and cloning was conducted through next-generation sequencing as part of the Twist bioscience service.

High-throughput rubisco expression

Chemocompetent BL21(DE3) cells, previously transformed with a pESL plasmid coding for the chaperone GroEL-GroES[13], were transformed with the rubisco library and incubated at 37°C, 250 rpm in 8 ml of LB media supplemented with 30 μg/ml chloramphenicol and 50 μg/ml kanamycin. Growth was performed in 24-deep-well plates. When cells reached an OD600 of 0.6, GroEL-GroES expression was induced by adding arabinose (0.2% final) and incubating at 23°C for 45 min. Rubisco expression was then induced by adding 0.2 mM IPTG (isopropyl β-d-thiogalactoside) and incubating at 23°C for 21 h. For protein extraction, cells were harvested by centrifugation (15 min; 4,000 g; 4°C) and pellets were lysed with BugBuster® ready mix (Millipore) for 25 min at room temperature; 70 μl BugBuster master mix (EMD Millipore) was added to each sample. Crude extracts were then centrifuged for 30 min at 4,000 g, 4°C to remove the insoluble fraction. For quality control, all samples were run on an SDS–PAGE gel (Supplementary Fig. 12).

Raf1-IB rubisco co-expression

For IB rubiscos (originating from β-cyanobacteria), a pilot study was performed to find an homologue of the chaperone Rubisco accumulation factor 1 (Raf1), that could help solubly-express these rubisco variants. 3 different Raf1 were first tested with their cognate rubisco, from 3 β-cyanobacteria (namely, *Trichormus variabilis*, *Pseudanabaena sp. PCC 6802*, and *Euhalothece natronophila*). Codon-optimized *raf1* genes were synthesized and cloned by Twist Bioscience in polycistron with their cognate rubisco genes into pET-29b(+) vectors. Proteins were expressed and tested *in vitro*, as described above, and Raf1 from *Euhalothece natronophila* was chosen, based on ability to solubilize its cognate rubisco. The gene was therefore cloned into a pESL plasmid, in the same operon as GroEL-ES genes, and transformed into chemocompetent BL21(DE3) cells. IB rubiscos were then transformed and expressed in these cells as described previously.

High-throughput determination of rubisco carboxylation rates

To determine rubisco carboxylation rates, we performed kinetic assays directly from the soluble fraction of prepared lysates since purifying both large and small rubisco subunits together was not feasible in a high-throughput manner. Soluble fractions were incubated with 4% CO_2 and 0.4% O_2 for rubisco activation (15 min; 4°C; plate shaker at 250 rpm). Following activation, rubisco carboxylase activities were tested as described in Davidi *et al.*[6]. Briefly, 10 μl of the activated rubisco sample was added to six aliquots of 80 μl of assay mix (a detailed list of all assay components and their sources is provided in Supplementary Table 1) containing different concentrations of CABP (0, 0, 10, 20, 30, and 90 nM). The mix was incubated for 15 minutes at

30°C in a plate reader (Infinite® 200 PRO; TECAN) connected to a gas control module (TECAN
pro200) ensuring atmospheric conditions of 4% CO_2 and 0.4% O_2. Rubisco carboxylation activity
was initiated by adding 10 µl RuBP to each sample (final concentration of 1 mM and a total volume
of 100 µl). The carboxylation rate was determined through a coupled reaction[50,51]. In brief, 3-
phosphoglycerate, the product of ribulose 1,5-bisphosphate carboxylation, was phosphorylated
and subsequently reduced into glyceraldehyde 3-phosphate, involving NADH oxidation. The latter
could be monitored through 340 nm absorbance for 15 minutes at 2-second intervals. To convert
from NADH to rubisco reactions per second, we assumed a stoichiometric ratio of 2:1 between
NADH and carboxylation reactions. The active-site concentration was determined by fitting a
linear regression model (custom python script; see below) to the measured reaction rates as a
function of the CABP inhibitor concentrations (See Supplementary Note 1 and Supplementary
Fig. 13 for more details). Due to the use of soluble fractions of total cell lysates, the initial
concentration of rubisco could not be determined beforehand, which often led to saturation of the
first assay with rubisco. To overcome this issue, we adjusted the concentration of lysates by
dilution, to obtain a rubisco concentration that enabled measurable inhibition by CABP, allowing
for an accurate quantification of the active-site concentration. We finally obtained the rate per
active site by dividing the activity with no CABP by the concentration of active sites. As was done
in Davidi *et al.*[6], and since not all variants were tested on the same day, the form-II rubisco from
R. rubrum, commonly used as a standard in rubisco kinetic assays, was consistently included as
an internal reference in every measurement.
388
Rubisco variants selection for comparative analysis
For comparing rubisco carboxylation rates between groups without sampling bias, we aimed to
select representative sets of variants uniformly covering the different groups studied. For the
comparison of rubiscos originating from phototrophic versus chemotrophic organisms, we
selected a set of 32 rubiscos, covering the entire diversity of form I rubiscos at a 75% similarity
threshold, enriched with a set of 36 rubiscos, covering the entire diversity of form IA and B
rubiscos at an 85% similarity threshold. Among these 68 variants, phototrophic rubiscos were
distinguished from chemotrophic ones based on their co-occurrence with photosystem genes in
the respective genome. This resulted in 53 chemotropic and 15 phototrophic rubiscos, of which
20 and 9 respectively were soluble and active *in vitro*.
For the comparison of carboxysome and non-carboxysome-associated rubiscos, we selected a
set of 14 rubiscos covering the entire diversity of carboxysome-associated rubiscos at an 85%
similarity threshold, and of 30 rubiscos covering the entire diversity of non-carboxysome-
associated rubiscos at a 75% similarity threshold. 9 and 9 of them were soluble and active *in vitro*.
For the comparison of α- and β- cyanobacterial, and carboxysome-associated proteobacterial
rubiscos, we selected 19 rubiscos covering the entire diversity of α-cyanobacteria rubiscos at a
97.5% similarity threshold, 29 rubiscos covering the entire diversity of β-cyanobacteria rubiscos
at a 91% similarity threshold, and 23 rubiscos covering the entire diversity of carboxysome-
associated proteobacterial rubiscos at a 90% similarity threshold. 15, 19 and 20 of them
respectively were soluble and active *in vitro*.
409

Data and Code Availability

All the data supporting the findings of this study as well as the codes used for generating our list of rubiscos and for analyzing the results is open source and can be found in the following link: https://gitlab.com/milo-lab-public/rubisco-F1.

References

1. Raven, J. A. Contributions of anoxygenic and oxygenic phototrophy and chemolithotrophy to carbon and oxygen fluxes in aquatic environments. *Aquat. Microb. Ecol.* **56**, 177–192 (2009).

2. Bar-On, Y. M. & Milo, R. The global mass and average rate of rubisco. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 4738–4743 (2019).

3. Prywes, N., Phillips, N. R., Tuck, O. T., Valentin-Alvarado, L. E. & Savage, D. F. Rubisco Function, Evolution, and Engineering. *Annu. Rev. Biochem.* (2023) doi:10.1146/annurev-biochem-040320-101244.

4. Wildman, S. G. Along the trail from Fraction I protein to Rubisco (ribulose bisphosphate carboxylase-oxygenase). *Photosynth. Res.* **73**, 243–250 (2002).

5. Flamholz, A. I. *et al.* Revisiting Trade-offs between Rubisco Kinetic Parameters. *Biochemistry* vol. 58 3365–3376 Preprint at https://doi.org/10.1021/acs.biochem.9b00237 (2019).

6. Davidi, D. *et al.* Highly active rubiscos discovered by systematic interrogation of natural sequence diversity. *EMBO J.* **39**, e104081 (2020).

7. Mann, C. C. Genetic Engineers Aim to Soup Up Crop Photosynthesis. *Science* **283**, 314–316 (1999).

8. Tabita, F. R. Microbial ribulose 1, 5-bisphosphate carboxylase/oxygenase: a different perspective. *Photosynth. Res.* **60**, 1–28 (1999).

9. Badger, M. R. & Bek, E. J. Multiple Rubisco forms in proteobacteria: their functional significance in relation to CO2 acquisition by the CBB cycle. *Journal of Experimental Botany* vol. 59 1525–1541 Preprint at https://doi.org/10.1093/jxb/erm297 (2008).

10. Witte, B. *et al.* Functional prokaryotic RubisCO from an oceanic metagenomic library. *Appl. Environ. Microbiol.* **76**, 2997–3003 (2010).

11. Aigner, H. *et al.* Plant RuBisCo assembly in E. coli with five chloroplast chaperones including BSD2. *Science* **358**, 1272–1278 (2017).

12. Liu, C. *et al.* Coupled chaperone action in folding and assembly of hexadecameric Rubisco. *Nature* **463**, 197–202 (2010).

13. Goloubinoff, P., Christeller, J. T., Gatenby, A. A. & Lorimer, G. H. Reconstitution of active
dimeric ribulose bisphosphate carboxylase from an unfolded state depends on two
chaperonin proteins and Mg-ATP. *Nature* **342**, 884–889 (1989).

14. Lin, Z. & Rye, H. S. Expansion and compression of a protein folding intermediate by GroEL.
Mol. Cell **16**, 23–34 (2004).

15. Huang, F. *et al.* Rubisco accumulation factor 1 (Raf1) plays essential roles in mediating
Rubisco assembly and carboxysome biogenesis. *Proceedings of the National Academy of*
Sciences **117**, 17418–17428 (2020).

16. Xia, L.-Y. *et al.* Molecular basis for the assembly of RuBisCO assisted by the chaperone
Raf1. *Nature Plants* **6**, 708–717 (2020).

17. Hauser, T. *et al.* Structure and mechanism of the Rubisco-assembly chaperone Raf1. *Nat.*
Struct. Mol. Biol. **22**, 720–728 (2015).

18. Kolesinski, P., Belusiak, I., Czarnocki-Cieciura, M. & Szczepaniak, A. Rubisco
Accumulation Factor 1 from Thermosynechococcus elongatus participates in the final
stages of ribulose-1,5-bisphosphate carboxylase/oxygenase assembly in Escherichia coli
cells and in vitro. *FEBS J.* **281**, 3920–3932 (2014).

19. Cen, Y.-P. & Sage, R. F. The regulation of Rubisco activity in response to variation in
temperature and atmospheric CO2 partial pressure in sweet potato. *Plant Physiol.* **139**,
979–990 (2005).

20. Flamholz, A. & Shih, P. M. Cell biology of photosynthesis over geologic time. *Curr. Biol.* **30**,
R490–R494 (2020).

21. Jordan, D. B. & Ogren, W. L. Species variation in the specificity of ribulose biphosphate
carboxylase/oxygenase. *Nature* **291**, 513–515 (1981).

22. Iñiguez, C. *et al.* Evolutionary trends in RuBisCO kinetics and their co-evolution with CO2
concentrating mechanisms. *Plant J.* **101**, 897–918 (2020).

23. Savir, Y., Noor, E., Milo, R. & Tlusty, T. Cross-species analysis traces adaptation of
Rubisco toward optimality in a low-dimensional landscape. *Proc. Natl. Acad. Sci. U. S. A.*
107, 3475–3480 (2010).

24. Badger, M. R. & Price, G. D. CO2 concentrating mechanisms in cyanobacteria: molecular
components, their diversity and evolution. *J. Exp. Bot.* **54**, 609–622 (2003).

25. Whitehead, L., Long, B. M., Price, G. D. & Badger, M. R. Comparing the in vivo function of
α-carboxysomes and β-carboxysomes in two model cyanobacteria. *Plant Physiol.* **165**,
398–411 (2014).

26. Cabello-Yeves, P. J. *et al.* α-cyanobacteria possessing form IA RuBisCO globally dominate

aquatic habitats. *ISME J.* (2022) doi:10.1038/s41396-022-01282-z.

27. Shih, P. M. *et al.* Biochemical characterization of predicted Precambrian RuBisCO. *Nat.*
Commun. **7**, 10382 (2016).

28. Long, B. M. *et al.* Carboxysome encapsulation of the CO2-fixing enzyme Rubisco in
tobacco chloroplasts. *Nat. Commun.* **9**, 3570 (2018).

29. Fournier, G. P. *et al.* The Archean origin of oxygenic photosynthesis and extant
cyanobacterial lineages. *Proceedings of the Royal Society B: Biological Sciences* **288**,
20210675 (2021).

30. Sánchez-Baracaldo, P. & Cardona, T. On the origin of oxygenic photosynthesis and
Cyanobacteria. *New Phytol.* **225**, 1440–1446 (2020).

31. Jasser, I. & Callieri, C. Picocyanobacteria. in *Handbook of Cyanobacterial Monitoring and*
Cyanotoxin Analysis 19–27 (John Wiley & Sons, Ltd, 2017).

32. Young Kevin D. The Selective Value of Bacterial Shape. *Microbiol. Mol. Biol. Rev.* **70**, 660–
703 (2006).

33. Benson, D. A. *et al.* GenBank. *Nucleic Acids Res.* **41**, D36–42 (2013).

34. Sunagawa, S. *et al.* Structure and function of the global ocean microbiome. *Science* **348**,
1261359 (2015).

35. Wrighton, K. C. *et al.* Fermentation, hydrogen, and sulfur metabolism in multiple
uncultivated bacterial phyla. *Science* **337**, 1661–1665 (2012).

36. Brown, C. T. *et al.* Unusual biology across a group comprising more than 15% of domain
Bacteria. *Nature* **523**, 208–211 (2015).

37. Anantharaman, K. *et al.* Thousands of microbial genomes shed light on interconnected
biogeochemical processes in an aquifer system. *Nat. Commun.* **7**, 1–11 (2016).

38. Tabita, F. R., Satagopan, S., Hanson, T. E., Kreel, N. E. & Scott, S. S. Distinct form I, II, III,
and IV Rubisco proteins from the three kingdoms of life provide clues about Rubisco
evolution and structure/function relationships. *J. Exp. Bot.* **59**, 1515–1524 (2008).

39. Banda, D. M. *et al.* Novel bacterial clade reveals origin of form I Rubisco. *Nat Plants* **6**,
1158–1166 (2020).

40. Edgar, R. C. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*
26, 2460–2461 (2010).

41. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

42. Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973

(2009).

43. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2--approximately maximum-likelihood

trees for large alignments. *PLoS One* **5**, e9490 (2010).

44. Tabita, F. R., Hanson, T. E., Satagopan, S., Witte, B. H. & Kreel, N. E. Phylogenetic and

evolutionary relationships of RubisCO and the RubisCO-like proteins and the functional

lessons provided by diverse molecular forms. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **363**,

2629–2640 (2008).

45. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

46. Spreitzer, R. J. & Salvucci, M. E. Rubisco: structure, regulatory interactions, and

possibilities for a better enzyme. *Annu. Rev. Plant Biol.* **53**, 449–475 (2002).

47. Park, S. W. *et al.* Presence of duplicate genes encoding a phylogenetically new subgroup

of form I ribulose 1,5-bisphosphate carboxylase/oxygenase in Mycobacterium sp. strain

JC1 DSM 3803. *Res. Microbiol.* **160**, 159–165 (2009).

48. Grostern, A. & Alvarez-Cohen, L. RubisCO-based CO2 fixation and C1 metabolism in the

actinobacterium Pseudonocardia dioxanivorans CB1190. *Environ. Microbiol.* **15**, 3040–

3053 (2013).

49. Schulz, L. *et al.* Evolution of increased complexity and specificity at the dawn of form I

Rubiscos. *Science* **378**, 155–160 (2022).

50. Lilley, R. M. & Walker, D. A. The reduction of 3-phosphoglycerate by reconstituted

chloroplasts and by chloroplast extracts. *Biochimica et Biophysica Acta (BBA) -

Bioenergetics* **368**, 269–278 (1974).

51. Kubien, D. S., Brown, C. M. & Kane, H. J. Quantifying the amount and activity of Rubisco in

leaves. *Methods Mol. Biol.* **684**, 349–362 (2011).