Use of linear features by red-legged partridges in an intensive agricultural landscape: implications for landscape management in farmland

Charlotte Perrot[1*], Léo Séranne[1,], Antoine Berceaux[2], Mathias Noel[2], Beatriz Arroyo[3], Léo Bacon[1]

[1] DRAS OFB, French Biodiversity Agency, 147 av de Lodève Juvignac, France
[2] Regional Federation of Hunters of Occitanie, 23 Chem. de Laveran, 31390 Carbonne, France
[3] Instituto de Investigación en Recursos Cinegéticos (IREC, CSIC-UCLM-JCCM), Ronda de Toledo 12, 13005 Ciudad Real, Spain.

Corresponding author
Correspondence: C. Perrot – mail : charlotte.perrot@ofb.gouv.fr

ABSTRACT

Current agricultural practices and change are the major cause of biodiversity loss. An important change associated with the intensification of agriculture in the last 50 years is the spatial homogenization of the landscape with substantial loss of such biodiversity-rich elements as seminatural linear feature (hedgerows, field margins, grass strips, etc.). In Europe, management prescriptions serve to increase heterogeneity by the creation of these seminatural and linear features which are not being used primarily for agricultural production. However, these elements are not equal in their support for biodiversity according to their structure and composition. The aim of this study is to determine the importance of landscape heterogeneity and specifically linear features on the spatial distribution of red-legged partridges, a small game species in decline in Europe. Through GPS-monitoring of adult birds, we first assess home range size throughout the year and during the breeding season, in relation to breeding status and to linear features density (semi-natural linear vegetation and tracks-roads for human traffic). Then, we focus on habitat selection in relation to linear landscape features during the breeding period. We found that linear elements shape the use of space by red-legged partridges according to their reproductive status. Traffic routes and semi-natural features structured by both herbaceous and woody cover, negatively influenced home range size. Further, breeding birds select linear elements with herbaceous cover while non-breeders select linear elements with woody cover, underlining the different needs of birds according to their breeding status. All birds selected areas near tracks, but non-breeders seemed to avoid roads. This study shows the importance for this species of the linear components that structure the agricultural landscape. We propose recommendations to promote the presence of the red-legged partridge in this agricultural environment but also of the biodiversity in general.

Keywords: Farmland bird, intensive agriculture, selection habitat, semi-natural linear element

Introduction

Agriculture is the most important land use activity, covering over a third of the world's land surface
(World Bank 2017). Current agricultural practices and changes are the biggest cause of biodiversity loss
(Dudley and Alexander 2017) by having broad scale negative effects with the conversion of natural
ecosystems into farms (Foley et al. 2005) and the release of pollutants (Brühl and Zaller 2019). An important
change associated with the intensification of agriculture in the last 50 years is the spatial homogenization of
the landscape with only a few sown crop types distributed in large uniform fields (Šálek et al. 2018) and with
substantial loss of such biodiversity-rich elements as hedgerows, field margins and other non-cropped areas
(Benton et al. 2003). For example, since 1950, 70% of the hedgerows have disappeared from the French
bocage and continue to decrease. A fundamental concept in landscape ecology is that spatial homogeneity
affects negatively ecological systems (Wiens 2002) which is also true for agricultural ecosystems.

In Europe, one response over biodiversity loss concerns in farmland has been the introduction of agri-
environment schemes (AES), in which farmers are funded to modify their farming practices to provide
environmental benefits (Batáry et al. 2015). Management prescriptions generally serve to increase
heterogeneity (Benton et al. 2003) by the maintenance, creation or management of seminatural features
which are not being used primarily for agricultural production. This includes for instance many linear
elements like field margins, road verges, ditch banks, hedgerows, and wooded banks (Grashof-Bokdam and
van Langevelde 2005). In intensively managed lowland landscapes, these linear features contain the greatest
botanical diversity and provide different functions for wildlife. They act as corridors, facilitating species
movement, providing shelters, food and nesting habitat for invertebrates (Lagerlöf et al. 1992; Delattre et al.
2013; Coulthard et al. 2016), reptiles (Knierim et al. 2018; Balouch et al. 2022), amphibians (Maes et al.
2008), mammals (Dondina et al. 2016; Pelletier-Guittier et al. 2020), and birds (Heath et al. 2017). The
structure (height and width) and plant diversity of semi-natural linear features have also been shown to
influence the biodiversity they support (Graham et al. 2018). Seminatural linear features are not equal in
their support for biodiversity and this highlights the importance of guidelines for local stakeholders to
manage them in line with their goals (Vickery et al. 2009; Montgomery et al. 2020).

The red-legged partridge (*Alectoris rufa*) is a typical species of agricultural land in southwestern Europe,
where it has considerable socioeconomic value in rural environments as one of the principal small game
species, including in France with ca 1.3 M birds hunted during the season 2013/14 (Aubry et al. 2016).
However, despite its importance as a quarry species, populations have suffered marked declines during the
last century throughout its range, including France (Souchay et al. 2022). Several reasons may explain these
declines, but the most important appears to be habitat alteration, particularly changes occurring during
recent decades in agrarian management systems (Delibes-Mateos et al. 2012). Previous studies on habitat
preferences of partridges in agricultural environments have highlighted the importance of seminatural
features for this species. For example, breeding density of red-legged partridge was found to correlate with
the length of permanent field boundaries including hedgerows (Rands 1986; Meriggi et al. 1991) and with
the amount of natural vegetation (Cabodevilla et al. 2021). Using radiotracking of breeding birds, Casas and
Viñuela (2010) showed that herbaceous strips among fields were positively selected as nesting sites, with
high nesting success. Today, to improve habitat selection studies, we can use new technological advances for
higher spatial and temporal accuracy. Indeed, recent integration of high-resolution Global Positioning System
(GPS) into miniaturized tracking tags has dramatically improved our ability to study habitat selection by
providing us with accurate locations at high temporal frequency (Kays et al. 2015).

The aim of this study was to determine the importance of landscape heterogeneity and specifically linear
features on the spatial distribution of red-legged partridges using GPS monitoring. Specifically, we first
assessed home range size throughout the year and during the breeding season, in relation to breeding status
and to linear features density. Then, we focused also on habitat selection in relation to linear landscape
features during the breeding period, and in relation to their structure: linear features considered included
seminatural linear elements with either herbaceous or woody structure, and traffic routes since they are
likely to influence wildlife habitat selection (Fahrig and Rytwinski 2009; Ascensão et al. 2012). We expected
breeding birds to be more selective than non-breeders as they are more constrained in their movements
during the incubation and rearing period. Finally, we looked at how the nests of the tracked birds were
located in relation to these linear features. Using the red-legged partridge as a flagship species, we discuss

our results in terms of recommendations for landscape management in farmland areas to maximize benefits
for breeding farmland birds.

Methods

Study area description and population.

The study was conducted at two sites 56 km apart in Southwest France. The first is located in the village
of Garganvillar in the Tarn-et-Garonne department (43°58'N, 1°30'E, 28.82 km2, hereafter TG fig.1). The
second is located within the villages of Azas, Montpitol and Roquesérière in the Haute-Garonne department
(43°43'N, 1°39'E, 43.76 km2, hereafter HG, Fig. 1). Both sites are characterized by an intensive agricultural
landscape dominated by cereal crops (wheat, barley, oats), corn crops, leguminous crops (lentil, chickpeas)
and oilseed crops (sunflower, rape). Their topography is relatively flat. These two sites benefit from a warm-
temperate, oceanic climate with Mediterranean tendencies, characterized by a hot and fairly dry summer, a
sunny autumn, a mild winter and a spring marked by recurring rainfall. Density of red-legged partridge is 2
pairs per 100 ha in TG and 1.5 pair per 100 ha in HG which is low. On both sites, captive-reared red-legged
partridges are released for population reinforcement every year in July. Survival of released red-legged
partridges to the following spring is lower than 0.05 (Souchay et al. 2018). Spring is also the period of capture
for this study. Thus, we considered captured birds in spring as wild birds regardless of their origin. In the
same way, because TG and HG have similar characteristics (topography, land use, climate and game
management), and are relatively close, we did not differentiate the birds according to where they reside.

The red-legged partridge exhibits a peculiar breeding system of double-nesting behaviour (Green 1984a;
Casas et al. 2009). Male incubation behaviour appears common in this species. Females lay eggs in two
different nests, and when both clutches are laid, females usually start incubation of the second clutch while
males incubate the first one and then raise their chicks (Green 1984a; Casas et al. 2009). In Southwest
France, predators of red-legged partridge include red foxes (*Vulpes vulpes*), feral cats (*Felis catus*), stone
martens (*Martes foina*), Free-roaming dogs (*Canis lupus familiaris*), common buzzards (*Buteo buteo*).

Figure 1 - Map of the two study sites illustrating the intensive agricultural landscape

Linear structure classification

Linear landscape structures include green linear features such as hedge and grass strips but also human traffic routes. Green linear features were surveyed in the field from June to July 2018. They were classified into 4 categories according to the percentage of cover of the herbaceous stratum and the presence of a woody stratum. *H Group* was characterized by an herbaceous cover above 25% and by the absence of woody plants. *W Group* was characterized by the presence of woody plants and by an herbaceous cover below 25%. *HW Group* was characterized by an herbaceous cover above 25% and by the presence of woody plants. Finally, the last group was characterized by an herbaceous cover lower than 25% and by the absence of woody stratum. However, this group represented less than 1% of the vegetated linear feature, so was not considered in this analysis. Next, all green linear features were georeferenced using QGIS software.

Human traffic routes include roads (paved traffic routes) and tracks (unpaved traffic routes). To map roads and tracks in our two sites, we used vectors from IGN cartographic database (ROUTE 500® edition 2020) and from OpenStreetMap database (OSM). Then, for both sites and for all linear structures, distance-to-linear-feature rasters were created, allowing to consider the different biological functions that these structures can have for red-legged partridges (i.e., proximity to a refuge, feeding, moving). Working with distances also allows to dilute the noise due to the imprecision of the GPS locations of the tracked birds, which is about 8 meters.

Animal captures and GPS data

Wild red-legged partridges were caught using decoy traps (Smith et al. 1981). Traps are observed from about 100 meters so that the trapped partridge remained in the trap as less as possible. We equipped 16 birds in March and April 2018, and 17 birds in February, March and April 2019 with solar GPS-tags coupled to a UHF radio transmitters (e-obs digital telemetry, 10g). The device was attached to the bird with a field-adjustable leg-loop harness using Teflon ribbon. Birds were morphologically sexed. The origin of birds (wild vs. farmed) was also checked by looking at the condition of their feathers. However, this is only possible the year after their release because once the breeding moult is done it is no longer possible to differentiate them. After release, birds were radio-tracked by triangulation using a portable Yagi antenna to download GPS data via the UHF system and monitor reproduction. Duration of monitoring was very variable between individuals (mean ± SD = 111 ± 159 days), due to the natural mortality of birds, UHF signal loss and GPS device dysfunctioning. All the animals were treated according to the ethical conditions detailed in the specific accreditations delivered to the Office Français de la Biodiversité by the Préfecture de Paris (prefectorial decree n°2009- 014) in agreement with the French environmental code (Art. R421-15 to 421-31 and R422-92 to 422-94-1).

From March to October, GPS devices recorded locations every hour from 3:00 to 21:00 UTC, that is 19 locations per day. From November to February, as days are shorter, we lightened the programming to avoid draining the battery. During this time, GPS devices recorded locations every hour from 6h00 to 18h00 UTC, that is 13 locations per day. For all GPS data, we screened them for erroneous locations from the individuals' trajectories (Bjørneraas et al. 2010). To avoid noise from the capture and death of the bird, we removed the first and last day of monitoring.

Reproductive status of partridges was estimated by the detection of incubation. An incubating bird was considered as breeding and conversely a non-incubating bird was considered as non-breeding. Detection of incubation was done by two methods. First, in 2018, birds were radio-tracked and thus observed twice a week to detect if they are incubating or not. When a bird was located 3 times successively in the same place, it was considered in incubation and the nest location was recorded. In 2019, for lack of human resources, field observations could not be done. We therefore identified nesting attempts and nest location from bird trajectories using the *find_nests* function of the *nestR* package (Picardi et al. 2020).

Analysis

Annual space use

For monthly home range size estimation, we selected individuals tracked at least one full month, that is 17 individuals (8 females with 4 breeders and 4 non-breeders, and 9 males with 3 breeders and 6 non-breeders). Mean monitoring length of these birds was 4.9 (± 4.2 SD) months. We derived home range size

for each individual for each month from an utilization distribution (UD) computed using the Brownian bridge
movement model (Horne et al. 2007). BBMM is a continuous time stochastic model of movement that
incorporates an animal's movement path and time between locations to calculate UD. We calculated
monthly UDs at 95% using the *R adehabitat* package (Calenge 2011). For each home range, the density of
each linear structure (roads, tracks and green linear features) was calculated in meters of linear structure per
hectare using QGIS software (version 3.10.4-A Coruña).
Next, we investigated if the month of the year, the individual characteristics and the density of linear
structures had an influence on the variability of monthly home range size using linear models. Since we had a
relatively small sample size and since males are likely to incubate and participate in the rearing of young, we
chose to consider only the reproductive status of the individual and not their sex. In other words, if a male
or female individual was seen incubating, it was considered a breeder, if not as a non-breeder. In the same
way, as we did not know the origin (wild or farmed bird) for all individuals, we could not test its influence on
the monthly home ranges size. Therefore, the complete model contained the month of the year, the
reproductive status of individual and density of the five linear structures as explanatory variables. Since there
may be multiple monthly home ranges per individual, bird identity was included as a random effect. Next, we
tested the significance of this random effect by the function *ranova* of the R package *lmerTest* (Kuznetsova et
al. 2017) to decide on the relevance of maintaining this effect in the model. To satisfy the assumptions of
the model (i.e. normality and homoscedasticity of residuals), home range size was log-transformed.
Following recent recommendations to produce model estimates comparable between and within studies, we
standardized all explanatory variables by centring and dividing by two standard deviations. To prevent
overparameterization, we respected the sample size rule-of thumb of 10:1 subjects to predictors in multiple
regression (Harrell 2015). Model selection was based on Akaike Information Criterion corrected for small
sample size (AICc). When several models were within a Δ AICc of 2 from the best model, we employed a
model averaging approach, using the natural average method implemented in the *MuMIn* package (Grueber
et al. 2011) of R on models within two points of AICc from the best one. This allowed us to account for model
selection uncertainty in order to obtain robust parameter estimates.

Space use and habitat selection during the breeding period
To study space use during the breeding period we focused on home range size, nest location and habitat
selection during breeding. We chose the period when most individuals were monitored simultaneously and
over the same duration in 2018 and 2019. This period ranges from March 26 to June 5 in 2018 and 2019,
therefore 71 days of monitoring for 10 tracked individuals (4 females all breeding and 6 males with 2
breeders et 4 non-breeders). It includes the end of the pair formation, nest construction and incubation.

Home range
We calculated home range size for every individual during this period using the same method as for
monthly home range size. As before, we investigated if the month of the year, individual characteristics and
the density of linear structures had an influence on the variability of this home range size. First, we tested
our hypothesis that non-breeding birds should express a larger home range than breeding birds. Hence, we
compared these two groups using the non-parametric Mann-Withney-Wilcoxon test using the alternative
"greater." Next, we tested the influence of month and green linear features on home range size using linear
models. Since the sample size is small (N=10), we tested each variable one by one, which makes six different
models. To satisfy the assumptions of the model (i.e. normality and homoscedasticity of residuals), home
range size was log-transformed.

Nest habitat
Distance from each nest to each linear structure was calculated. When distance between the nest and
the green linear feature was less than 8 meters (GPS accuracy), we considered that the nest was within this
linear feature. Then we checked if the nest was close to a road or a track.

Habitat selection
Since red-legged partridges are mainly active during the day, and since we had few night locations, we
used only the locations recorded during daytime. To evaluate how linear structures influence habitat
selection, observed 60-min movement steps (i.e. straight lines connecting successive GPS locations) were

compared to random steps; the former are typically used to estimate habitat use, and the latter as an
estimate of habitat availability (Thurfjell et al. 2014). Habitat selection and movement are interlinked, with
movement rates influencing selection and vice versa (Forester et al. 2009). Therefore, integrated step-
selection analysis was used to compare observed and random steps to evaluate habitat selection while
simultaneously accounting for the differing movement behaviours (Avgar et al. 2016).
Twenty-five random steps for each observed step were generated using analytical distributions that were
parameterized based on the observed animal movement patterns. Observed movement patterns were
described using gamma distribution of step length and a Von Mises distribution of turning angles from each
individual.
Habitat selection was evaluated as a function of distances to linear structures: green linear features (H,
W, HW), roads and tracks. Habitat attributes at the end of each observed step were compared to attributes
at the end of random steps. Distances were transformed using the natural logarithm because we expected
animals to respond more strongly to features closer to them, with the response declining at an unknown rate
(Dickie et al. 2020). Each individual was modelled separately using conditional logistic regression (*amt*
package in R; Signer et al. 2019). The relative probability of selection was modelled as a function of distance
linear structures. The step length, natural logarithm of step length, cosine of turning angle were included as
modifiers of the observed movement parameter.
Individual selection and movement responses were summarized to evaluate consistency in responses
based on model coefficients and their 95% confidence intervals (CIs). Selection is defined as occurring if use
was higher than availability, resulting in positive selection coefficients, and avoidance if use was less than
availability, resulting in negative selection coefficients. CIs overlapping with zero were interpreted as
indifference, and non-overlapping CIs as significant selection/avoidance depending on the variable of interest
(Dickie et al. 2020; Fieberg et al. 2021).
Then, to test the influence of breeding status on habitat selection, we used inverse variance-weighted
linear modelling to obtain population-level averages for both groups (Murtaugh, 2007). Finally, to
understand how strongly linear features influenced selection, the relative selection strength was calculated
(Avgar et al. 2017). For all linear structures, we calculated the relative probability of selecting an area
according to its distance with these structures.
Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Maecenas porttitor congue massa. Fusce
posuere, magna sed pulvinar ultricies, purus lectus malesuada libero, sit amet commodo magna eros quis
urna. Nunc viverra imperdiet enim. Fusce est. Vivamus a tellus. Pellentesque habitant morbi tristique
senectus et netus et malesuada fames ac turpis egestas. Proin pharetra nonummy pede. Mauris et orci.
Aenean nec lorem.

Results

Linear structure length

259   On both sites, 147.25 km of roads and 63.91 km of tracks were recorded. Concerning green linear
260 features, 52.39 km, 11,18 km and 65.68 km were respectively recorded for *H*, *W* and *HW group*.

Annual space use

262   The mean size of monthly home range was 25.14 ha ± 1.28 SE. Four models explained home range size
263 almost equally well (Table 1). Model averaging indicated a negative effect of the density of the two traffic
264 ways, although this effect is less marked for roads than for paths (Table 2, Fig 2a, 2b). In the same way,
265 density of HW linear negatively influenced home range size (Table 2, Fig 2c). We did not detect any effect of
266 reproductive status, month and linear density of H and W group on monthly home range size.

Figure 2 - Effect of a) track density, b) road density, c) density of linear of HW group on monthly home range size in red-legged partridge according to the model averaging of the top model set (estimate ± 95%IC). In grey, points represent raw data.

268 **Table 1** - Model selection of the factors influencing monthly home range size in red legged partridge. Models with a Δ AICc ≤ 2
269 from the best model are represented.
270

Response variable	Models	df	logLik	AICc	delta	weight
Home range size (log)						
	track density + road density + HW density	5	-41.14	92.95	0	0.37
	track density + HW density + H density	5	-41.35	93.38	0.43	0.30
	track density + road density + HW density + H density	6	-40.73	94.41	1.46	0.18
	track density + road density + HW density + W density	6	-40.82	94.60	1.65	0.16

271
272
273

274 **Table 2** - Model-averaged estimates ± SE and 95%CI of parameters explaining variations in monthly home range size in red-
275 legged partridge. The relative importance of each factor is calculated by summing the AIC weights across the top models (Table
276 1) where the given factor appears (last column).

Response variable	Parameters	Estimate	SE	Confidence interval	Sum of weight
Home range size (log)					
	Track density	**-0.50**	**0.08**	**(-0.66 ; -0.33)**	**1**
	Road density	**-0.24**	**0.11**	**(-0.47 ; -0.02)**	**0.70**
	HW density	**-0.37**	**0.09**	**(-0.55 ; -0.20)**	**1**
	H density	-0.21	0.13	(-0.46 ; 0.04)	0.47
	W density	-0.06	0.08	(-0.22 ; 0.10)	0.16

278
279
280

Reproductive space use

Home range

While not statistically significant at the 0.05 level, the Mann-Withney-Wilcoxon test indicated that the
home range size of non-breeders tends to be larger than that of breeders (W = 20, p = 0.057). During the
reproductive period ranging from March 26 to June 5, non-breeders had a mean home range size of 53.19 ha
± 12.60 SE while breeders had a mean home range size of 22.34 ha ± 3.27 SE (Fig. 3). Concerning linear
structures, track density and HW group density negatively influenced home range size (tracks: F_{18} = 7.33, p =
0.03; HW group: F_{18} = 27.51, p = 0.0008, Fig. 4a ,4b). We did not detect any effect of road and linear density
of H and W groups on reproductive home range size.
290
291

Figure 3 - Mean home range size ± SE of red-legged partridge for non-breeders (N=4) and breeders (N=6).

299
300 **Figure 4** - Effect of a) track density, b) density of linear of HW group on home range size during breeding
301 in red-legged partridge according to linear regression models (estimate ± 95%IC). In grey, points
302 represent raw data.

 Nest habitat

9 nest locations of 7 different breeders were identified either by field observations or by studying the
trajectory of the birds using the *nestR* package. 4 of these nests were located in green linear features of *HW*
group, of which 2 were close to a road (distance = 6.26 m and 7.14 m) and 2 close to a track (distance = 1.18
m and 6.02 m). 4 other nests were located in green linear features of *H group*, all near a road (distance = 3.72
m, 3.20 m, 4.47 m and 7.71 m). The last nest was located in a garden containing grass and bushes, but no
linear structures.

Linear features selection
Breeding red-legged partridges selected to be closer to green area features of *H* and *HW groups* and to
tracks. However, they did not show any selection or avoidance behaviours concerning *W group* features and
roads (Table 3). Concerning distance to *H group* feature, it is 1.91 times more likely that an individual will use
areas 5 m away from this linear feature than an area 50 m away, which itself is 1.48 times more attractive
than an area 200 m away. For distance to *HW group* feature, it is 1.71 times more likely that an individual
uses an area 5 m away from this linear than an area 50 m away, which itself is 1.38 times more attractive
than an area 200 m away (Fig. 5a). In the same way, the probability of using an area 5 meters from a track is
1.22 times more likely than an area 50 meters from the track, which itself is 1.12 times more attractive than
an area 200 meters away (Fig. 5a).
Non-breeder birds selected to be closer to green area features of *W* and *HW groups* and to tracks but
they were indifferent to *H group* features. (Table 3). They also tended to avoid the areas closest to roads
(Fig. 5b). An area 5 meters away from a feature of *W group* is 2.38 times more attractive than an area
located 50 meters away, which is itself 1.68 times more attractive than an area located 200 meters away (Fig.
5b). For *HW group* feature, an area 5 meters away from is 2.03 times more attractive than an area located 50
meters away, which is itself 1.53 times more attractive than an area located 200 meters away (Fig. 5b). In the
same way, the probability of using an area 5 meters from a track is 1.37 times more likely than using an area
50 meters from the track, which itself is 1.20 times more attractive than an area 200 meters away (Fig. 5b).
328

329 **Table 3** - Model-averaged estimates ± SE and 95%CI of parameters explaining variations in monthly home range size in red-
330 legged partridge. The relative importance of each factor is calculated by summing the AIC weights across the top models (Table
331 1) where the given factor appears (last column).

Status	Component	Variable	Coefficient	SE	CI	n
Breeder	Movement	Step length	0.0005	0.0003	[0.0002 ; 0.0009]	6
		ln(Step length)	0.0626	0.0298	[0.0191 ; 0.1061]	6
		Cos(Turn angle)	-0.0091	0.0190	[-0.0368 ; 0.0186]	6
	Selection	ln(distance to W)	-0.0293	0.0975	[-0.1716 ; 0.1130]	6
		ln(distance to HW)	-0.2327	0.0388	[-0.2893 ; -0.1761]	6
		ln(distance to H)	-0.2812	0.0496	[-0.3536 ; -0.2088]	6
		ln(distance to track)	-0.0853	0.0548	[-0.1653 ; -0.0053]	6
		ln(distance to road)	0.0802	0.1066	[-0.0753 ; 0.2358]	6
Non-breeder	Movement	Step length	0.0005	0.0001	[0.0003 ; 0.0007]	4
		ln(Step length)	0.0764	0.0576	[-0.0078 ; 0.1605]	4
		Cos(Turn angle)	-0.0622	0.0514	[-0.1373 ; 0.0128]	4
	Selection	ln(distance to W)	-0.3771	0.1175	[-0.5487 ; -0.2056]	4
		ln(distance to HW)	-0.3065	0.0386	[-0.3629 ; -0.2501]	4
		ln(distance to H)	-0.0888	0.0845	[-0.2121 ; 0.0345]	4
		ln(distance to track)	-0.1364	0.0721	[-0.2418 ; -0.0311]	4
		ln(distance to road)	0.0919	0.0655	[-0.0037 ; 0.1875]	4

333
334 **Figure 5** - Relative selection strength of green linear feature of H (dotted line), HW (dotted-dashed line),
335 W (dashed line) groups and of tracks (grey long dashed line) and roads (grey solid line) for a) breeder b) non-
336 breeders red-legged partridge.
337

Discussion

The reproductive status of birds, as well as the structure of the habitats in which they live, significantly influence their use of space (Rolando 2002). Our study is consistent with this, as tracked red-legged partridges occupied space differently depending on their reproductive status. First, during the breeding season, breeding birds tended to have a smaller home range than non-breeders, probably due to the constraints of breeding. Indeed, during the incubation period, the movements of the nesting males and females are limited to the nest and its close environment. Then, during the rearing period, parents must adapt to the low movement abilities of their non-flying chicks (Green 1984b).

Second, reproductive status also shapes habitat selection. In spring, semi-natural linear features with herbaceous cover were highly selected by breeders. Moreover, all nests monitored but one were located in this type of linear feature, i.e. those with an herbaceous stratum. These results are consistent with previous studies on partridges that have shown the importance of grass cover for nesting (Rands 1986; Casas and Viñuela 2010). Indeed, field margins with herbaceous plants such as hedgerows or grass strips, provide a cover to hide the nest from predators, food for adults (grasses and seeds) and for chicks (arthropods) (Vickery et al. 2009). However, field margins are known to attract predators because of their foraging behaviours toward linear features (Morris and Gilroy 2008). For example, during foraging, field margins and hedgerows facilitate movement of foxes, a partridge predator, through the landscape (Feber et al. 2019). Hence, nesting in these linear landscape elements could be an ecological trap like for the grey partridge (Bro et al. 2004; Rantanen et al. 2010). Conversely, non-breeding birds chose linear elements with a woody structure (bushes and/or trees) with or without grass cover. Non-breeders chose these hedgerows probably because the woody structures offer a good safety against raptors and they can exploit nearby resources or access to locations which might otherwise be too risky to use at all (Hinsley and Bellamy 2000). Hedgerows with a woody and herbaceous stratum were therefore selected by red-legged partridges regardless of their reproductive status. Furthermore, we found in our tracked partridges that the higher the density of these hedges, the more the monthly and breeding home range were reduced. These two results together underline the importance of these hedges with woody and herbaceous strata for the red-legged partridge, probably for its multiple functions (shelter, food resource, movement facilitator).

In the same way, both breeders and non-breeders selected tracks or areas very close to them and home range size was negatively correlated with track density. There are several non-exclusive hypotheses for the attraction of unpaved roads. Firstly, tracks may be unobstructed ways that facilitate the movement of ground-dwelling birds such as partridges. Secondly, tracks may also be associated with grassy ditches and banks, or hedgerows. Thus, partridges potentially find shelter and food closer to tracks. Finally, tracks are a source of grit used by partridges mainly for grinding hard food items but also for its nutritional role as a source of minerals (Gionfriddo and Best 1999).

Compared to tracks, roads have more intense traffic which is a source of disturbance due to the noise, but also a source of mortality (Forman and Alexander 1998). Hence, in our study, non-breeding red-legged partridges tended to avoid these noisy and hazardous habitats probably to ensure their survival. This result is consistent with the study of Cooke et al. (2020) that found a negative relationship between red-legged partridge abundance and road exposure. Moreover, the regular discovery of red-legged partridges killed on the roads by collision with vehicles in our study site, gives weight to this hypothesis (unpublished data). However, breeding red-legged partridges did not avoid the roads contrary to non-breeding red-legged partridges. In addition, four of the eight nests found were located within 8 meters of a road. Thus, the road environment appears to attract breeding individuals during reproduction. Similar to tracks, roads are frequently surrounded by ditches and grassy banks full of seeds and invertebrates, providing nesting habitat as well as food for adults and chicks. However, many studies demonstrate that road-side productivity and plant diversity attract wildlife while at the same time inflicting mortality so that roads can act as ecological traps (Coffin 2007). Surprisingly, the partridges with a small home range were the ones with the highest road density in their home range. Thus, we could think that the road environment provides the necessary resources for red-legged partridges during the year. But it is more likely that roads act as dangerous barriers that impede movement and thus limit the

extent of bird territories. Red-legged partridges seem affected by the habitat fragmentation through the
development of the road network like in other terrestrial bird species (Laurance et al. 2004).
Red-legged partridge seems also impacted by the homogenization of agricultural landscapes due to
the development of intensive agriculture. This homogenization results into larger fields, and thus rarer
semi-natural lines (hedgerows, field sides and unpaved roadsides). Red-legged partridges must move
further to cover their basic needs (food, nest, resting place, etc.) leading to an increase in their home
range size as in other farmland species (Michel et al. 2017; Mayer et al. 2022). This could explain why
home ranges estimated in our study are up to 2.7 times larger than those previous studies during the
breeding season (Ricci 1985; Mauvy et al. 1991). But monitoring methods may also be the cause of this
size discrepancy. GPS tracking allows the recording of a greater number of locations than the radio
tracking used in previous studies, which may explain the larger home range in our study.
Finally, we found no seasonal effect on monthly home range size, which is surprising because (1) birds
generally have different movement constraints in relation to their phenology (breeding, dispersal, fall
company) (2) the seasonal rotation of agricultural crops changes the distribution of food resources for
farmland species during the year. Due to the small number of birds monitored, the lack of power in our
analyses could explain the lack of detection of this seasonal variability.

405
Management recommendations
Orłowski (2008) proposed to move bushy vegetation away from the roads to reduce the likelihood
that birds would cross the road and die. In the same way, we recommend the creation of a vegetation-
free buffer zone along roadsides to prevent ground-nesting birds from colliding with vehicles but in
roadkill hotspots only. Vegetation removal in roadkill hotspots should always be combined with small-
mesh fencing to prevent passage of not only red-legged partridges, but also their predators (foxes) and
small species mammals that inhabit the roadside (Galantinho et al. 2022). However, on roads with less
risk of collision, this roadside management does not seem appropriate because it would be to the
detriment of many species, since vegetated roadsides are favourable to the establishment of small
mammals (Ascensão et al. 2012), insects (New et al. 2021) and contribute to the plant diversity (Lázaro-
Lobo and Ervin 2019). Hence, collision risk mapping in this intensive agricultural environment would
greatly facilitate roadside management.
For the red-legged partridge we believe that the major problem is not road layout but the lack of
semi-natural elements in these intensive agricultural landscapes, so that birds are forced to use non-
optimal habitats such as roadsides. These results underline the importance of developing a network of
semi-natural elements in agricultural areas, such as hedges, grassy strips or embankments for the
preservation of the red-legged partridge and of biodiversity in general. We have seen that, depending on
their reproductive status, red-legged partridges did not select the same kind of field margins because
they do not provide the same resources (nesting site, food, shelter, corridor, etc.). More broadly,
quantity of seminatural elements and their heterogeneity in shape, size, structural complexity, and also
in their plant composition are essential to meet the needs of the different species living in this
agricultural environment. For example, hedgerows favoured the presence of forest edge birds (Mallet et
al. 2022) but may also be detrimental for some grassland bird species, such as skylark (*Alauda arvensis*)
for which grass strips are beneficial (Josefsson et al. 2013). Equally, some bats are more active in
hedgerows composed mainly of trees (Lacoeuilhe et al. 2018), while the hazel dormouse (*Muscardinus*
avellanarius) and the European badger (*Meles meles*) prefer hedgerows with a high shrub cover (Dondina
et al. 2016).
Agricultural policies should therefore support the increase of linear semi-natural areas in agricultural
landscapes without forgetting to consider their complexity and heterogeneity so has to better meet
management and conservation objectives. The maintenance over time of these seminatural linear
elements should be part of this policy, so that they continue to fulfil their function for biodiversity. For
this to work, it is essential to get farmers on board with this biodiversity conservation effort. So, it is
necessary to measure the effectiveness of this habitat management, by monitoring the fauna and flora
and to transfer these results from scientists to local stakeholders.

Acknowledgements

We would like to thank all the people involved in the fieldwork among sites and over years and in particular Frédéric Le Capitaine, Benoît Laboup, Arnaud Gaujard, Julie Grezeleau, Romain Mongeot, Estella Vignon, Anne-Charlotte Pommier-Petit, Soumaya Belghali and Lisa Gili. We would also like to thank our partners for their support in this study: the Federation of Hunters of Haute-Garonne (FDC 31) and the Federation of Hunters of Tarn-et-Garonne (FDC82). We thank Aude Géraud for her logistic support during the project. We also would like to thank Françoise Ponce, Jean-Bernard Puchala and Luc Fruitet for their help in setting up this project. We are grateful to Michel Ruas for the donation of 2 farmed red-legged partridges which allowed us to improve our skills in equipping birds with GPS tags.

Data, scripts, code, and supplementary information availability

Data are available online: https://doi.org/10.5281/zenodo.8144149
Scripts and code are available online: https://doi.org/10.5281/zenodo.8144131

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Funding

This project was funded by the OFB, Agrifaune and the Regional Federation of Hunters of Occitanie.

References

Ascensão F, Clevenger AP, Grilo C, Filipe J, Santos-Reis M. 2012. Highway verges as habitat providers for small mammals in agrosilvopastoral environments. Biodivers Conserv. 21(14):3681–3697. doi:10.1007/s10531-012-0390-3.

Aubry P, Anstett L, Ferrand Y, Reitz F, Klein F, Ruette S, Sarasa M, Arnauduc J-P, Migot P. 2016. Enquête nationale sur les tableaux de chasse à tir. :8.

Avgar T, Lele SR, Keim JL, Boyce MS. 2017. Relative Selection Strength: Quantifying effect size in habitat- and step-selection inference. Ecology and Evolution. 7(14):5322–5330. doi:10.1002/ece3.3122.

Avgar T, Potts JR, Lewis MA, Boyce MS. 2016. Integrated step selection analysis: bridging the gap between resource selection and animal movement. Methods in Ecology and Evolution. 7(5):619–630. doi:https://doi.org/10.1111/2041-210X.12528.

Balouch S, Driscoll DA, Naseer A, Rais M, Doherty TS. 2022. Impacts of land cover on reptile movement and habitat use in farming landscapes. Animal Conservation. 25(6):837–848. doi:10.1111/acv.12789.

Batáry P, Dicks LV, Kleijn D, Sutherland WJ. 2015. The role of agri-environment schemes in conservation and environmental management. Conservation Biology. 29(4):1006–1016. doi:10.1111/cobi.12536.

Benton TG, Vickery JA, Wilson JD. 2003. Farmland biodiversity: is habitat heterogeneity the key? Trends in Ecology & Evolution. 18(4):182–188. doi:10.1016/S0169-5347(03)00011-9.

Bjørneraas K, Van Moorter B, Rolandsen CM, Herfindal I. 2010. Screening Global Positioning System Location Data for Errors Using Animal Movement Characteristics. The Journal of Wildlife Management. 74(6):1361–1366. doi:10.1111/j.1937-2817.2010.tb01258.x.

Bro E, Mayot P, Corda E, Reitz F. 2004. Impact of habitat management on grey partridge populations: assessing wildlife cover using a multisite BACI experiment. Journal of Applied Ecology. 41(5):846–857. doi:10.1111/j.0021-8901.2004.00939.x.

Bro E, Reitz F, Clobert J. Nest-site selection of grey partridge (Perdix perdix) on agricultural lands in north-central France. :10.

Brühl CA, Zaller JG. 2019. Biodiversity Decline as a Consequence of an Inappropriate Environmental Risk Assessment of Pesticides. Frontiers in Environmental Science. 7. doi:10.3389/fenvs.2019.00177. https://www.frontiersin.org/articles/10.3389/fenvs.2019.00177.

Cabodevilla X, Estrada A, Mougeot F, Jimenez J, Arroyo B. 2021. Farmland composition and farming practices explain spatio-temporal variations in red-legged partridge density in central Spain. Science of The Total Environment. 799:149406. doi:10.1016/j.scitotenv.2021.149406.

Calenge C. 2011. Home Range Estimation in R: the adehabitatHR Package. Office national de la chasse et de la faune sauvage.:61.

Casas F, Mougeot F, Viñuela J. 2009. Double-nesting behaviour and sexual differences in breeding success in wild Red-legged Partridges Alectoris rufa. Ibis. 151(4):743–751. doi:10.1111/j.1474-919X.2009.00958.x.

Casas F, Viñuela J. 2010. Agricultural practices or game management: which is the key to improve red-legged partridge nesting success in agricultural landscapes? Environmental Conservation. 37(2):177–186. doi:10.1017/S0376892910000299.

Coffin AW. 2007. From roadkill to road ecology: A review of the ecological effects of roads. Journal of Transport Geography. 15(5):396–406. doi:10.1016/j.jtrangeo.2006.11.006.

Cooke SC, Balmford A, Donald PF, Newson SE, Johnston A. 2020. Roads as a contributor to landscape-scale variation in bird communities. Nat Commun. 11(1):3125. doi:10.1038/s41467-020-16899-x.

Coulthard E, McCollin D, Littlemore J. 2016. The use of hedgerows as flight paths by moths in intensive farmland landscapes. J Insect Conserv. 20(2):345–350. doi:10.1007/s10841-016-9864-z.

Delattre T, Vernon P, Burel F. 2013. An agri-environmental scheme enhances butterfly dispersal in European agricultural landscapes. Agriculture, Ecosystems & Environment. 166:102–109. doi:10.1016/j.agee.2011.06.018.

Delibes-Mateos M, Farfán MÁ, Olivero J, Vargas JM. 2012. Impact of land-use changes on red-legged partridge conservation in the Iberian Peninsula. Environmental Conservation. 39(4):337–346. doi:10.1017/S0376892912000100.

Dickie M, McNay SR, Sutherland GD, Cody M, Avgar T. 2020. Corridors or risk? Movement along, and use of, linear features varies predictably among large mammal predator and prey species. Journal of Animal Ecology. 89(2):623–634. doi:10.1111/1365-2656.13130.

Dondina O, Kataoka L, Orioli V, Bani L. 2016. How to manage hedgerows as effective ecological corridors for mammals: A two-species approach. Agriculture, Ecosystems & Environment. 231:283–290. doi:10.1016/j.agee.2016.07.005.

Dudley N, Alexander S. 2017. Agriculture and biodiversity: a review. Biodiversity. 18(2–3):45–49.
doi:10.1080/14888386.2017.1351892.

Fahrig L, Rytwinski T. 2009. Effects of Roads on Animal Abundance: an Empirical Review and
Synthesis. Ecology and Society. 14(1). https://www.jstor.org/stable/26268057.

Feber RE, Johnson PJ, Gelling M, Macdonald DW. 2019. Ecology and conservation of mammals of
hedgerows and field margins. In: The Ecology of Hedgerows and Field Margins. Routledge. p. 233–
249.

Fieberg J, Signer J, Smith B, Avgar T. 2021. A 'How-to' Guide for Interpreting Parameters in
Habitat-Selection Analyses. bioRxiv.:2020.11.12.379834. doi:10.1101/2020.11.12.379834.

Foley JA, DeFries R, Asner GP, Barford C, Bonan G, Carpenter SR, Chapin FS, Coe MT, Daily GC,
Gibbs HK, et al. 2005. Global Consequences of Land Use. Science. 309(5734):570–574.
doi:10.1126/science.1111772.

Forester JD, Im HK, Rathouz PJ. 2009. Accounting for animal movement in estimation of resource
selection functions: sampling and data analysis. Ecology. 90(12):3554–3565. doi:10.1890/08-0874.1.

Forman RTT, Alexander LE. 1998. Roads and Their Major Ecological Effects. Annual Review of
Ecology and Systematics. 29(1):207–231.

Galantinho A, Santos S, Eufrázio S, Silva C, Carvalho F, Alpizar-Jara R, Mira A. 2022. Effects of
roads on small-mammal movements: Opportunities and risks of vegetation management on roadsides.
Journal of Environmental Management. 316:115272. doi:10.1016/j.jenvman.2022.115272.

Gionfriddo JP, Best LB. 1999. Grit Use by Birds, a review. In: Nolan V, Ketterson ED, Thompson
CF, editors. Current Ornithology. Boston, MA: Springer US. p. 89–148. https://doi.org/10.1007/978-
1-4757-4901-4_3.

Graham L, Gaulton R, Gerard F, Staley JT. 2018. The influence of hedgerow structural condition on
wildlife habitat provision in farmed landscapes. Biological Conservation. 220:122–131.
doi:10.1016/j.biocon.2018.02.017.

Grashof-Bokdam CJ, van Langevelde F. 2005. Green Veining: Landscape Determinants of
Biodiversity in European Agricultural Landscapes. Landscape Ecol. 20(4):417–439.
doi:10.1007/s10980-004-5646-1.

Green RE. 1984a. Double nesting of the Red-legged Partridge Alectoris rufa. Ibis. 126(3):332–346.
doi:10.1111/j.1474-919X.1984.tb00254.x.

Green RE. 1984b. The Feeding Ecology and Survival of Partridge Chicks (Alectoris rufa and Perdix
perdix) on Arable Farmland in East Anglia. Journal of Applied Ecology. 21(3):817–830.
doi:10.2307/2405049.

Grueber CE, Nakagawa S, Laws RJ, Jamieson IG. 2011. Multimodel inference in ecology and
evolution: challenges and solutions. Journal of Evolutionary Biology. 24(4):699–711.
doi:https://doi.org/10.1111/j.1420-9101.2010.02210.x.

Harrell FE. 2015. Regression Modeling Strategies: With Applications to Linear Models, Logistic and
Ordinal Regression, and Survival Analysis. Cham: Springer International Publishing (Springer Series
in Statistics). [accessed 2023 May 10]. https://link.springer.com/10.1007/978-3-319-19425-7.

Heath SK, Soykan CU, Velas KL, Kelsey R, Kross SM. 2017. A bustle in the hedgerow: Woody field
margins boost on farm avian diversity and abundance in an intensive agricultural landscape.
Biological Conservation. 212:153–161. doi:10.1016/j.biocon.2017.05.031.

Hinsley SA, Bellamy PE. 2000. The influence of hedge structure, management and landscape context
on the value of hedgerows to birds: A review. Journal of Environmental Management. 60(1):33–49.
doi:10.1006/jema.2000.0360.

Horne JS, Garton EO, Krone SM, Lewis JS. 2007. Analyzing Animal Movements Using Brownian
Bridges. Ecology. 88(9):2354–2363. doi:10.1890/06-0957.1.

Josefsson J, Berg Å, Hiron M, Pärt T, Eggers S. 2013. Grass buffer strips benefit invertebrate and
breeding skylark numbers in a heterogeneous agricultural landscape. Agriculture, Ecosystems &
Environment. 181:101–107. doi:10.1016/j.agee.2013.09.018.

Kays R, Crofoot MC, Jetz W, Wikelski M. 2015. Terrestrial animal tracking as an eye on life and
planet. Science. 348(6240):aaa2478. doi:10.1126/science.aaa2478.

Knierim TK, Marshall BM, Hayes L, Suwanwaree P, Strine CT. 2018. The movements and habitat
preferences of a Malayan krait (Bungarus candidus) in an agrarian landscape. The Herpetological
Bulletin.:5.

Lacoeuilhe A, Machon N, Julien J-F, Kerbiriou C. 2018. The Relative Effects of Local and Landscape
Characteristics of Hedgerows on Bats. Diversity. 10(3):72. doi:10.3390/d10030072.

Lagerlöf J, Stark J, Svensson B. 1992. Margins of agricultural fields as habitats for pollinating insects.
Agriculture, Ecosystems & Environment. 40(1):117–124. doi:10.1016/0167-8809(92)90087-R.

Laurance SGW, Stouffer PC, Laurance WF. 2004. Effects of Road Clearings on Movement Patterns
of Understory Rainforest Birds in Central Amazonia. Conservation Biology. 18(4):1099–1109.
doi:10.1111/j.1523-1739.2004.00268.x.

Lázaro-Lobo A, Ervin GN. 2019. A global examination on the differential impacts of roadsides on
native vs. exotic and weedy plant species. Global Ecology and Conservation. 17:e00555.
doi:10.1016/j.gecco.2019.e00555.

Maes J, Musters CJM, De Snoo GR. 2008. The effect of agri-environment schemes on amphibian
diversity   and   abundance.   Biological   Conservation.   141(3):635–645.
doi:10.1016/j.biocon.2007.12.018.

Mallet P, Béchet A, Sirami C, Mesléard F, Blanchon T, Calatayud F, Dagonet T, Gaget E, leray C,
Galewski T. 2022. Field margins as substitute habitat for the conservation of birds in agricultural
wetlands. doi:10.1101/2022.05.05.490780. http://biorxiv.org/lookup/doi/10.1101/2022.05.05.490780.

Mauvy B, Garcia A, Lartiges A, Péroux R. 1991. Occupation of area by red-legged partridge on a
cultivated plain in the Dordogne department (France). Bulletin Mensuel de l'Office National de la
Chasse.

Mayer M, Fischer C, Blaum N, Sunde P, Ullmann W. 2022. Influence of roads on space use by
European hares in different landscapes. 38:131–146. doi:10.1007/s10980-022-01552-3.

Meriggi A, Montagna D, Zacchetti D. 1991. Habitat use by partridges (perdix perdix and alectoris
rufa)   in   an   area   of   northern   apennines,   Italy.   Bollettino   di   zoologia.   58(1):85–89.
doi:10.1080/11250009109355733.

Michel VT, Naef-Daenzer B, Keil H, Grüebler MU. 2017. Reproductive consequences of farmland heterogeneity in little owls (Athene noctua). Oecologia. 183(4):1019–1029. doi:10.1007/s00442-017-3823-6.

Montgomery I, Caruso T, Reid N. 2020. Hedgerows as ecosystems: service delivery, management and restoration. Annual Review of Ecology and Systematics. doi:10.1146/annurev-ecolsys-012120-100346. https://www.annualreviews.org/doi/abs/10.1146/annurev-ecolsys-012120-100346.

Morris AJ, Gilroy JJ. 2008. Close to the edge: predation risks for two declining farmland passerines. Ibis. 150(s1):168–177. doi:10.1111/j.1474-919X.2008.00857.x.

New TR, Sands DPA, Taylor GS. 2021. Roles of roadside vegetation in insect conservation in Australia. Austral Entomology. 60(1):128–137. doi:10.1111/aen.12511.

Orłowski G. 2008. Roadside hedgerows and trees as factors increasing road mortality of birds: Implications for management of roadside vegetation in rural landscapes. Landscape and Urban Planning. 86(2):153–161. doi:10.1016/j.landurbplan.2008.02.003.

Pelletier-Guittier C, Théau J, Dupras J. 2020. Use of hedgerows by mammals in an intensive agricultural landscape. Agriculture, Ecosystems & Environment. 302:107079. doi:10.1016/j.agee.2020.107079.

Picardi S, Smith BJ, Boone ME, Frederick PC, Cecere JG, Rubolini D, Serra L, Pirrello S, Borkhataria RR, Basille M. 2020. Analysis of movement recursions to detect reproductive events and estimate their fate in central place foragers. Movement Ecology. 8(1):24. doi:10.1186/s40462-020-00201-1.

Rands MRW. 1986. Effect of Hedgerow Characteristics on Partridge Breeding Densities. Journal of Applied Ecology. 23(2):479–487. doi:10.2307/2404030.

Rantanen EM, Buner F, Riordan P, Sotherton N, Macdonald DW. 2010. Habitat preferences and survival in wildlife reintroductions: an ecological trap in reintroduced grey partridges. Journal of Applied Ecology. 47(6):1357–1364. doi:10.1111/j.1365-2664.2010.01867.x.

Ricci J-C. 1985. Variations du mode d'occupation de l'espace chez la perdrix rouge (Alectoris rufa L.) depuis la formation des couples jusqu'à la couvaison. Acta oecologica Oecologica generalis. 6(3):281–293.

Rolando A. 2002. On the ecology of home range in birds. Revue d'Ecologie, Terre et Vie. 57(1):53–73.

Šálek M, Hula V, Kipson M, Daňková R, Niedobová J, Gamero A. 2018. Bringing diversity back to agriculture: Smaller fields and non-crop elements enhance biodiversity in intensively managed arable farmlands. Ecological Indicators. 90:65–73. doi:10.1016/j.ecolind.2018.03.001.

Signer J, Fieberg J, Avgar T. 2019. Animal movement tools (amt): R package for managing tracking data and conducting habitat selection analyses. Ecology and Evolution. 9(2):880–890. doi:https://doi.org/10.1002/ece3.4823.

Souchay G, Aebischer NJ, Arroyo B, Blanco-Aguiar JA, Meriggi A, Porto M, Reino L, Ponce F. 2022. Red-Legged Partridge Monitoring and Population Trends. In: Casas F, García JT, editors. The Future of the Red-legged Partridge: Science, Hunting and Conservation. Cham: Springer International Publishing. (Wildlife Research Monographs). p. 249–273. https://doi.org/10.1007/978-3-030-96341-5_10.

Souchay G, Besnard A, Perrot C, Jakob C, Ponce F. 2018. Anthropic and natural factors drive
variation of survival in the red-legged partridge in southern France. Wildlife Biology. 2018(1):1–10.
doi:10.2981/wlb.00438.

Thurfjell H, Ciuti S, Boyce MS. 2014. Applications of step-selection functions in ecology and
conservation. Mov Ecol. 2(1):4. doi:10.1186/2051-3933-2-4.

Vickery JA, Feber RE, Fuller RJ. 2009. Arable field margins managed for biodiversity conservation:
A review of food resource provision for farmland birds. Agriculture, Ecosystems & Environment.
133(1):1–13. doi:10.1016/j.agee.2009.05.012.

Wiens JA. 2002. Central Concepts and Issues of Landscape Ecology. In: Gutzwiller KJ, editor.
Applying Landscape Ecology in Biological Conservation. New York, NY: Springer. p. 3–21.
https://doi.org/10.1007/978-1-4613-0059-5_1.

World Bank. 2017. Agricultural land (% of land area) | Data. The World Bank.
https://data.worldbank.org/indicator/AG.LND.AGRI.ZS?page=1%29.

651
652