Title: The nuclei of human adult stem cells can move within the cell and generate cellular

protrusions to contact other cells.

Authors: Carlos Bueno[1,*], David García-Bernal[1,2], Salvador Martínez[3,4,5,6], Miguel Blanquer[1]

and José M. Moraleda[1].

[1]Medicine Department and Hematopoietic Transplant and Cellular Therapy Unit, Institute of

Biomedical Research (IMIB), University of Murcia, Faculty of Medicine, Murcia, 30120,

Spain.

[2]Biochemistry, Molecular Biology and Immunology Department, University of Murcia, Faculty

of Medicine, Murcia, 30100, Spain

[3]Instituto de Neurociencias de Alicante (UMH-CSIC), Universidad Miguel Hernandez, San

Juan, Alicante, 03550, Spain.

[5]Center of Biomedical Network Research on Mental Health (CIBERSAM), ISCIII, 28029,

Madrid, Spain.

[6]Alicante Institute for Health and Biomedical Research (ISABIAL), Alicante, 03010, Spain.

*Correspondence: Carlos Bueno, PhD., Medicine Department and Hematopoietic Transplant and

Cellular Therapy Unit, Institute of Biomedical Research (IMIB), University of Murcia, Faculty

of Medicine, Murcia, 30120, Spain. Tel.: 0034-968885013. Fax: 00-34-868884150. E-mail:

carlos.bueno@um.es

Abstract

Despite a considerable interest in understanding the mechanisms regulating nucleus structure, chromatin organization and nuclear positioning over decades, the exact significance of the variety of morphologies and positioning that cell nuclei can adopt and their relationship in cellular function is still far from being clearly understood. In this study, we examined the functional significance of the variety of morphologies and positioning that cell nuclei of human bone marrow-derived mesenchymal stem cells can adopt during neural-like differentiation. Here, we show that after neural induction, human bone marrow-derived mesenchymal stem cells enter an intermediate cellular state in which the nuclei are observed to be able to move within the cells, switching shapes and positioning and even generating cellular protrusions as they attempt to contact the cells around them. These findings suggest that changes in nuclear positioning are due to the fact that human cell nuclei somehow sensing their environment.

Introduction

Since the first observation of a nucleus in 1700[1], our knowledge of nuclear composition, organization, and positioning has continuously evolved[2,3,4]. Most textbooks depict the nucleus as a spherical or ovoid object at the center of the cell. However, different cell types have very different nuclear shapes and the position of nuclei varies dramatically from this simple view[2,3,5]. Although the cell nucleus has always been considered as the largest and most rigid organelle of eukaryotic cells, emerging views of the nucleus indicate a more dynamic organelle than expected[6].

There is increasing evidence that nuclei are frequently asymmetrically positioned depending on cell type, developmental stage, migratory state, and differentiation status[2,7,8,9,10]. It has been reported that the position of the nucleus contributes to cell mechanics, as gene regulation through relative genome segregation and the organization of cells within tissues[2,9,10,11]. Furthermore, it is important to note that changes in nuclear morphology and positioning are often associated with cellular dysfunction and disease[2,12,13]. Therefore, the nucleus must be considered not only as the

primary site for the storage of genetic material and gene transcription but also as a fundamental mechanical component of the cellular structure[6]. Despite these advances, the exact significance of nuclear positioning in cellular function and tissue physiology is still far from being clearly understood[2,5,7].

In our laboratory we focus on the differentiation of human mesenchymal stem cells (hMSCs) to generate a neuronal lineage[15,16,17,18]. hMSCs are considered as promising candidates for cell-based regenerative medicine due to their self-renewal capacity, multilineage differentiation potential, trophic effects and immunomodulatory properties[19,20]. Controlled neural differentiation of hMSCs could therefore become an important source of cells for cell therapy of neurodegenerative diseases, as autologous adult hMSCs are easily harvested and effectively expanded[21,22,23].

Over the past two decades, it has been reported that bone marrow-derived cells (BMDCs) and hMSCs can be induced to overcome their mesenchymal fate and differentiate into neural cells, both *in vitro*[24,25,26,27,28,29,30,31,32] and *in vivo*[33,34,35,36,37,38,39], a phenomenon known as transdifferentiation. The term transdifferentiation refers to the conversion of one mature cell type into another cell of different blastodermic origin[40,41]. Such interconversions may involve to regress into an intermediate step before cells differentiate into a new blastodermic potentiality and mature phenotype, or they may occur directly in a process that bypasses such intermediate phenotypes[40,41]. The actual occurrence of neuronal transdifferentiation of BMDCs and MSCs is currently much debated because the findings and their interpretation have been questioned[42,43]. The main argument against these observations in culture studies is that MSCs rapidly adopt neural-like morphologies by retraction of the cytoplasm, rather than by active neurite extension[42,44,45,46]. The *in vivo* neural transdifferentiation of BMDCs and hMSCs has also been questioned, as cell fusion could explain the development of new cell types, that are misinterpreted as transdifferentiated cells[43].

In a previous publication[18], we showed that when human bone marrow-derived MSCs (hBM-MSCs) were exposed to a neural induction medium, they rapidly reshaped from a flat to a spherical morphology. Subsequently, hBM-MSCs could maintain the spherical morphology or

adopt a new one; they gradually adopted a neural-like morphology through active neurite

extension or re-differentiated back to the mesenchymal fate. Furthermore, we found that hBM-

MSCs can rapidly and repeatedly switch lineage without cell division. Our results provide

evidence that differentiation of hBM-MSCs into neural-like cells requires a transition through a

transient and characterized intermediate state of hBM-MSCs (hBM-MSC-derived intermediate

cells) and provide a stronger basis for rejecting the idea that the rapid acquisition of a neural-like

morphology during MSC transdifferentiation is merely an artifact.

This previous work also highlights that nuclear remodeling occurs during *in vitro* neural-like

differentiation of hBM-MSCs. We found that nuclei in hBM-MSC-derived intermediate cells

moved within the cell, adopting different morphologies, and even forming two nuclei connected

by an internuclear bridge. These nuclear movements generated cellular protrusions that appeared

and disappeared from the surface of hBM-MSC-derived intermediate cells. The hBM-MSC-

derived intermediate cells positioned their nucleus at the front of the cell during migration. Our

results showed that binucleated hBM-MSCs can be formed during neural-like differentiation with

independence of any cell fusion, providing evidence that transdifferentiation may also be the

mechanism behind the presence of gene-marked binucleated neurons after gene-marked bone

marrow-derived cell transplantation. It is important to note that binucleated and polymorphic

nuclear cells have been detected in adult neurogenic niches[15,47,48,49,50].

Taken together, these findings suggest that to date there is no conclusive evidence to continue to

consider neuronal transdifferentiation of BMDCs and MSCs as a simple experimental artifact,

since it recapitulates some structural steps described *in vivo*[40,41]. Therefore, future studies are

needed to understand the mechanisms of these cell conversion processes and eventually harness

them for use in regenerative medicine.

In the present study, we investigated the sequence of biological events during neural-like

differentiation of hBM-MSCs using live-cell nuclear fluorescence labelling and time-lapse

microscopy to understand why the nuclei of hBM-MSC-derived intermediate cells move within

the cell generating the cellular protrusions that appear and disappear from the surface.

Results

Characterization of hBM-MSCs cultures

In a previous publication[18], we showed that hBM-MSCs did not express hematopoietic lineage

markers such as CD45, CD14, CD34 and CD20, and were positive for CD90, CD105, CD73,

thereby demonstrating a characteristic immunophenotype of hMSCs. Under proliferation

conditions, hBM-MSCs displayed a flat, fibroblast-like morphology with β-III-tubulin

microtubules and actin microfilaments oriented parallel to the longitudinal axis of the cell (Fig.

1a). During interphase, hBM-MSCs displayed a flattened, ellipsoidal nucleus, often located in the

center of the cell and with a nuclear volume of approximately $419'30 \pm 106'38 \ \mu m^3$ (Fig. 1b). The

dynamic localization of the nuclear lamina was analyzed by immunostaining for lamin A/C, a

nuclear lamina component[51] and the dynamic localization of the nucleoli was analyzed by

immunostaining for fibrillarin, the main component of the active transcription centers[52]. A

speckled pattern was observed distributed throughout the nucleus with heterogeneity in the

number, size, and distribution of the fibrillarin positive specks (Fig. 1c). Laser scanning confocal

microscopy revealed that the inner surface of the nuclear envelope is lined by the nuclear lamina

(Fig. 1d).

Characterization of hBM-MSC-derived intermediate cells

In a previous publication[18], we showed that following neural induction, hBM-MSCs rapidly

reshaped from a flat to a spherical morphology (hBM-MSCs intermediate cells). Subsequently,

we observed that hBM-MSC-derived intermediate cells can preserve their spherical shape, change

to that of neural-like cells through active neurite extension or revert back to their mesenchymal

morphology.

In this study, we focus on hBM-MSC-derived intermediate cells that can maintain their spherical

shape for several days without assuming new fates. To better understand why the nuclei of hBM-

MSC-derived intermediate cells move within the cell to generate the cellular protrusions, we

performed time-lapse microscopy and immunocytochemical analyses of histone H2B-GFP

transfected hBM-MSCs within the first 70 hours of neural induction. The time-lapse experiments

were performed using a higher magnification objective and a shorter image capture interval than

previously published experiments[18].

Time-lapse imaging revealed that when hBM-MSCs were exposed to a neural induction medium,

they rapidly reshaped from a flat to a spherical morphology (Fig. 2a and Supplementary Video 1).

We then observed hBM-MSC-derived intermediate cells in which nuclear movements generated

only one cell protrusion (Fig. 2a, white arrow and Supplementary Video 1) and hBM-MSC-

derived intermediate cells in which nuclear movements alternately generated one or two cellular

protrusions (Fig. 2a, yellow arrows and Supplementary Video 1). We found that when hBM-MSC-

derived intermediate cell has a nucleus without lobes, its movement within the cell generates only

one cell protrusion (Fig. 2b and Supplementary Video 2). However, if the hBM-MSC-derived

intermediate cell has a lobed nucleus it will generate one or two cellular protrusions depending

on how it moves within the cell (Fig. 2c).

Although the cell nuclei switch their morphologies while moving, time-lapse imaging and

immunocytochemical analysis revealed that hBM-MSC-derived intermediate cells have three

main different nuclear morphologies: Tail-less nuclei (Fig. 3a, white asterisk), tailed nuclei (Fig.

3a, green asterisk) and lobed nuclei (Fig. 3a, yellow asterisk). Tail-less and tailed nuclei

movements generate only a single cell protrusion (Fig. 3a, white and green arrows respectively).

However, as mentioned above, lobed nucleus movements generate one or two cellular protrusions

depending on how it moves within the cell (Fig. 3a, yellow arrows).

Confocal microscopy analysis and 3D reconstruction revealed that there are small variations in

both shape and size within the three types of nuclear morphology (Fig. 3b). Tail-less nuclei, tailed

nuclei and lobed nuclei have a volume of approximately $327'11 \pm 94'19$ µm^3, $306'89 \pm 16'50$ µm^3

and $361'75 \pm 147'44$ µm^3 respectively. It is important to note that confocal microscopic analysis

and 3D reconstruction also revealed that the lobes of the lobed nuclei can be located in different

z-planes (Fig. 3c).

We observed that tail-less nuclei and tailed nuclei contained one or two fibrillarin positive specks,

whereas lobed nuclei contained one or two fibrillarin positive specks in each lobe (Fig. 3d). No

positive fibrillarin specks have been detected either in the tail of the tailed nuclei or in the region

of nucleoplasm connecting each lobe of lobed nuclei. Laser scanning confocal microscopy also

revealed that the inner surface of the nuclear envelope is lined by the nuclear lamina (Fig. 3e).

Confocal microscopy analysis of the cytoskeletal organization of hBM-MSC-derived

intermediate cells showed that the cytoskeletal network is reorganized. Actin microfilaments and

β-III tubulin microtubules are no longer oriented parallel to the longitudinal axis of the hBM-

MSC-derived intermediate cells (Fig. 4a). Furthermore, confocal microscopy analysis and 3D

reconstruction revealed that cell protrusions are almost devoid of actin microfilaments and that

the β-III tubulin protein is concentrated at the cell protrusion rim (Fig. 4b, black arrows).

Although, it has been reported that direct connections between the actin cytoskeleton and the

nucleus govern nuclear positioning and nuclear movement during cell polarization and

migration[2,7,8], we observed that the absence of the cytoskeletal network could facilitate nuclear

movement within the cell. Future research is required to determine the feasibility of this

conjecture.

Nuclear remodeling

To further understand how the three different types of nuclei that move within the hBM-MSC-

derived intermediate cells are formed, we examined the sequence of biological events during

neural-like differentiation of histone H2B-GFP-transfected hBM-MSCs using time-lapse

microscopy. Time-lapse imaging revealed that there are variations in the form and time at which

cell nuclei adopt these different nuclear morphologies (Supplementary Videos 3-11). Although

future studies are needed to determine all the possible different nuclear remodeling sequences that

occur when hBM-MSCs reshape from a flat to a spherical morphology, below we have described

some examples of the formation of each of the three types of nuclei that move within hBM-MSC-

derived intermediate cells.

We observed that tail-less nuclei are formed by a single nuclear remodeling that occurs as the cell

reshapes from a flat to a spherical morphology, positioning the nucleus in a peripheral position

within the cell (Fig. 5, white arrows and Supplementary Videos 3,4). The duration of this

particular process is approximately 30 hours. Subsequently, the tail-less nucleus began to move

within the hBM-MSC-derived intermediate cells. (Fig. 5, yellow arrows and Supplementary

Videos 3,4).

Time-lapse images also showed that tailed nuclei are formed by one (Supplementary Video 5) or

two nuclear remodeling sequences. Below, we show an example of the formation of a tailed

nucleus generated by two nuclear remodeling sequences (Fig. 6 and Supplementary Videos 6,7).

We found that first a nuclear remodeling sequence occurs as the cell reshapes from a flat to a

spherical morphology, positioning the nucleus in a peripheral position within the cell (Fig. 6,

white arrows and Supplementary Videos 6,7). The duration of this particular process is

approximately 12 hours. Subsequently, the cell nucleus moves (Fig. 6, green arrows and

Supplementary Videos 6,7) and undergoes a second nuclear remodeling sequence in which a tailed

process in the nucleus is formed (Fig. 6, yellow arrows and Supplementary Videos 6,7). The

duration of this particular process is approximately 5 hours. Finally, the tailed nucleus began to

move within the hBM-MSC-derived intermediate cells. (Fig. 6, green arrows and Supplementary

Videos 6,7). Time-lapse imaging also revealed that as the nuclei move within the cell, the tails

can switch shape and size (Fig. 7a and Supplementary Video 8) and even appear to move in

different z-planes (Fig. 7b and Supplementary Videos 6,7). Future analyses will be needed to

determine whether the nuclei use the tails to stabilize their position within the cell.

Time-lapse imaging also revealed that lobed nuclei are formed by one (Supplementary Video 8)

or two nuclear remodeling sequences. Below, we show you an example of the formation of a

lobed nucleus generated by two nuclear remodeling sequences (Fig. 8 and Supplementary Videos

9,10). We found that first a nuclear remodeling sequence occurs as the cell reshapes from a flat to

a spherical morphology, positioning the nucleus in a peripheral position within the cell (Fig. 8,

white arrows and Supplementary Videos 9,10). The duration of this particular process is

approximately 14 hours. Subsequently, the cell nucleus moves (Fig. 8, green arrows and

Supplementary Videos 9,10) and undergoes a second nuclear remodeling sequence in which a

lobed nucleus is formed (Fig. 8, yellow arrows and Supplementary Videos 9,10). The duration of

this particular process is approximately 5 hours. The lobed nuclei then began to move within the

hBM-MSC-derived intermediate cells (Fig. 8, green arrows and Supplementary Videos 9,10).

Finally, we also found that lobed nuclei can switch shape while moving within the cell, becoming

tailed nuclei (Fig. 8, blue arrows and Supplementary Videos 9,10).

As mentioned above, confocal microscopy analysis and 3D reconstruction revealed that the lobes

of the lobed nuclei can be located in different z-planes (Fig. 3c). This result together with the

time-lapse noted in Figure 2a and Supplementary Video 1 suggest that each lobule of the lobed

nuclei can move in different z-planes.

Quantitative analysis of time-lapse imaging revealed that the nuclear speed oscillated between 0.1

and 1.3 µm/min, with an average speed of 0,66 ± 0,45 µm/min. These findings are consistent with

previous studies that reported the physical characteristics of typical nuclear movement in different

cell types[2]. Nuclear movement can generate cellular protrusions that extend up to a length similar

to the cell diameter (Fig. 2,4 and Supplementary Videos 1,2).

The hBM-MSC-derived intermediate cells use their cell nuclei to interact with other cells

Time-lapse imaging revealed that hBM-MSC-derived intermediate cell nuclei move within the

cell, generating cellular protrusions as cell migration breakthrough process to contact surrounding

cells (Fig. 2,5,8 and Supplementary Videos 1,2,4,10). Furthermore, we observed that contacts

occur between cell nuclei of hBM-MSC-derived intermediate cells, even for several hours (Fig.

9 and Supplementary Video 11). Confocal microscopy analysis (Fig. 10a) and 3D reconstruction

(Fig. 10b) revealed that tail-less nuclei, tailed nuclei and lobed nuclei contact each other. We note

that lobed nuclei can contact other cells through one lobe or both simultaneously or successively

(Fig. 2a and Supplementary Video 1).

Taken together, these findings suggest that changes in nuclear positioning are due to the fact that

cell nuclei are somehow sensing their surroundings.

Discussion

The fate of adult cells was thought to be restricted to their tissue blastodermic of origin[53].

However, there is now a large body of evidence suggesting that under physiological conditions

and certain experimental conditions, adult cells may be more plastic than we previously thought

in that they can become cells of unrelated lineages, a phenomenon known as transdifferentiation

senso stricto[40,41,54,55]. The most evident transdifferentiation events are epitelial-mesenchymal

transition (EMT) and its reverse process mesenchymal-epithelial transition (MET)[56,57,58]. These

processes are associated with implantation, embryo formation, organ development and wound-

healing[56,57,58]. Importantly, EMT and MET have been implicated in pathological conditions, such

as organ fibrosis, and in cancer, where they contributes to tumor progression and metastasis[56,57,58].

In the process of transdifferentiation, cells pass through intermediate states that are not well

understood[40,56,57,58,59]. Given the potential application of this cell conversion process, not only in

developmental and cancer studies but also in regenerative medicine, a better understanding of

intermediate states is crucial to avoid uncontrolled conversion or proliferation, which poses a risk

to patients[40,56,57,58,59].

Over the last two decades, it has been reported that BMDCs and hMSCs can be induced to

overcome their mesenchymal fate and transdifferentiate into neural cells, both *in*

vitro[24,25,26,27,28,29,30,32,32] and *in vivo*[33,34,35,36,37,38,39]. However, the neuronal transdifferentiation of

BMDCs and MSCs is still considered to be merely an artefact[42,43,46]. The main argument against

these observations in culture studies is that MSCs rapidly adopt neuronal-like morphologies by

retraction of the cytoplasm, rather than by active neurite extension[42,44,45,46]. The main argument

against neuronal transdifferentiation of BMDCs and hMSCs *in vivo* is that cell fusion could

explain the development of new cell types, that are misinterpreted as transdifferentiated cells[43].

In previous publications, we have shown that MSCs isolated from adult human tissues can

differentiate into neural-like cells, both *in vitro* and *in vivo*[14,15,16,17,18]. We found that when hBM-

MSCs were exposed to a neural induction medium, they rapidly reshaped from a flat to a spherical

morphology[18], Subsequently, hBM-MSCs could maintain the spherical morphology or adopt a

new one; they gradually adopted a neural-like morphology through active neurite extension or re-

differentiated back to the mesenchymal fate. Furthermore, we found that hBM-MSCs can rapidly

and repeatedly switch lineage without cell division. These results provide evidence that the

differentiation of hBM-MSCs into neural-like cells requires a transition through an intermediate

state, as described in the natural transdifferentiation processes[40,41,56,57,58,59].

This previous work[18] also highlights that nuclear remodeling occurs during *in vitro* neural-like

differentiation of hBM-MSCs. We found that nuclei in hBM-MSC-derived intermediate cells

moved within the cell, adopting different morphologies and even forming two nuclei connected

by an internuclear bridge, independently of any cell fusion.

These results provide a strong basis for rejecting the idea that the rapid acquisition of a neural-

like morphology during *in vitro* MSC transdifferentiation is merely an artefact and also provide

evidence that transdifferentiation may also be the mechanism behind the presence of gene-marked

binucleated neurons after gene-marked bone marrow-derived cell transplantation.

It is important to note that the studies describing the presence of binucleated Purkinje neurons

after bone marrow-derived cell transplantation suggest, but do not conclusively demonstrate, that

cell fusion is the underlying mechanism to explain the presence of binucleated neurons[43,60,61].

Futhermore, it has been reported that binucleated Purkinje neurons are also present in healthy,

unmanipulated mice and humans[43,62]. In addition, many authors have described binucleated

neurons in various central and peripheral parts of the nervous system[63,64,65,66]. What is more, many

authors have reported that many cultured hippocampal neurons[48] (their Fig. S2), and neural stem

cells located in the ventricular-subventricular zone of the anterolateral ventricular wall of the

human fetal brain[49] (their Fig. 2C) and adult mouse brain[50,51] (their Fig. 1E, 4I, 6F, S1B, S2B, S6A

and their Fig. 3A respectively) have two nuclei connected by an internuclear bridge.

Taken together, these results suggest that to date there is no conclusive evidence to date to

continue to consider the neuronal transdifferentiation of BMDCs and hMSCs as an artefact.

Therefore, future studies are needed to optimize the various neural induction protocols that have

been developed for MSCs[67], not only to understand the mechanisms of these cellular conversion

processes, but also to eventually harness them for use in regenerative medicine[40,68,69].

Our current research aims to improve our understanding of what hBM-MSCs look like at this

intermediate stage. We wanted to know why the nuclei of hBM-MSC-derived intermediate cells

move within the cell and generate the cellular protrusions that appear and disappear from the

surface.

In this study, we have shown that once the hBM-MSCs enter this intermediate stage, the cell

nuclei begin to move within the cell and generate the cellular protrusions as they try to contact

the surrounding cells. Interkinetic nuclear migration (INM) in neural progenitors underlay normal

neurogenesis during normal development[70,71], suggesting that the observed nuclear motility in

hBM-MS-derived intermediated cells recapitulates an intermediate stage before to decide

proliferative of differentiative progression in development and transdifferentiation. Importantly,

in a previous publication[18] (their Fig. S4 and Supplementary Video 4), we observed that the nuclei

of hBM-MSCs can also exhibit oscillatory movement along the axis of cell polarity, similar to

that described to occurs during INM.

We also found that contacts occur between the cell nuclei of hBM-MSC-derived intermediate

cells, even for several hours. These findings suggest that changes in nuclear positioning occur

because cell nuclei somehow sense their surroundings. Under our knowledge, we have described

for the first time the process of direct interactions between cells nuclei, which opens the possibility

of a new level of intercellular interaction. During this intermediate phase, nuclear factors may be

interchanged by neighboring cells and directly modify transcriptomes. Extensive studies are

required to determine the mechanisms and consequences of this process of direct interactions

between cells nuclei in normal and pathologic circumstances.

This work also highlights that cell nuclei moving within hBM-MSC intermediate cells have

mainly three different morphologies: tail-less nuclei, tailed nuclei, and lobed nuclei. We have

shown that there are variations in the shape and time at which the nuclei of hBM-MSC adopt the

different nuclear morphologies observed in hBM-MSC-derived intermediate cells. We also found

that tail-less and tailed nuclei movements generate only a single cell protrusion when attempting

to contact other cells. However, lobed nuclei movements generate one or two cellular protrusions

depending on how it moves within the cell. Lobed nuclei may contact other cells through one lobe

or both simultaneously or successively. Nevertheless, it is important to note that the three different

nuclear morphologies observed in hBM-MSC intermediate cells can be interchanged as the

nucleus moves within the cell.

Beyond the nervous system, the presence of lobed nuclei has been reported in most immune

cells[3,72] and in various human organs, including the heart and liver[73,74]. However, the functional

significance of multilobed nuclear structures is not yet known[3,72,73,74]. It would therefore be

interesting to investigate whether these nuclear structures are also associated with nuclear

movement within the cell.

In conclusion, our findings enrich the understanding of intermediate states in the neural-like

differentiation process of hBM-MSCs and suggest that changes in nuclear positioning are due to

the fact that human cell nuclei somehow sensing their environment. Although future experiments

will be needed to understand why there are different nuclear morphologies in hBM-MSC-derived

intermediate cells and why they are formed in different ways, our results demonstrate that the

nucleus should already be considered not only as the primary site for the storage of genetic

material and the transcription of genes, but also as a fundamental mechanical component of the

cell. Human mesenchymal stromal cells could not only help to increase our understanding of the

mechanisms underlying cellular plasticity, and eventually harness them for use in regenerative

medicine, but also facilitate an understanding of the mechanisms regulating nuclear structure and

dynamics.

Methods

Ethical conduct of research

The authors declare that all protocols used to obtain and process all human samples were approved by the local ethics committees of the Miguel Hernández University of Elche (No. UMH.IN.SM.03.16) and University of Murcia (HULP3617.05/07/2012; HUSA19/1531.17/02/2020) according to Spanish and European legislation and conformed to the ethical guidelines of the Helsinki Declaration. Donors provided written informed consent before obtaining samples.

Isolation and culture of hBM-MSCs

A standard protocol for isolation and expansion of hMB-MSCs was used as previously described[18]. Bone marrow aspirates were obtained by percutaneous direct aspiration from the iliac crest of 5 healthy volunteers at University Hospital Virgen de la Arrixaca (Murcia, Spain). Bone marrow was collected with 20 U/ml sodium heparin, followed by a Ficoll density gradient-based separation by centrifugation at 540g for 20 min. After, mononuclear cell fraction was collected, washed twice with Ca^{2+}/Mg^{2+}-free phosphate buffered saline (PBS) (Gibco Invitrogen) and seeded into 175-cm2 culture flasks (Nunc, Thermo Fisher Scientific) at a cell density $1.5x10^5$ cells/cm^2 in serum-containing media (designated as the basal media), composed of DMEM low glucose medium (Thermo Fisher Scientific) supplemented with 10% fetal bovine serum (FBS; Lonza), 1% GlutaMAX (Thermo Fisher Scientific), non-essential amino acid solution (Sigma-Aldrich) and 1% penicillin/streptomycin (Thermo Fisher Scientific). After 3 days of culture at 37ºC and 7% CO_2, non-attached cells were removed and fresh complete medium was added. Culture media were renewed every 2 days, and the isolated hMB-MSCs were passaged when cultures were 70-80% confluent. All studies were performed using hMB-MSCs expanded within culture passages 3-4.

Expression Vectors and Cell Transfection

The expression vectors used in the present study were H2B-eGFP, a gift from Geoff Wahl (Addgene plasmid # 11680; http://n2t.net/addgene:11680; RRID:Addgene_11680; Kanda et al[75]., 1998). A standard protocol for transfecting MSCs was used as previously described[18]. Isolated hMB-MSCs were transfected using the Gene Pulser-II Electroporation System (Bio-Rad Laboratories). Electroporation was performed in a sterile cuvette with a 0.4-cm electrode gap (Bio-Rad Laboratories), using a single pulse of 270 V, 500 µF. Plasmid DNA (5 µg) was added to 1.5×10^6 viable hMB-MSCs in 0.2-ml DMEM low glucose medium (Thermo Fisher Scientific) before electrical pulsing.

Time-lapse microscopy of histone H2B-GFP expressing hBM-MSCs cultured in neural induction media.

We used µ-Dish 35 mm, high Grid- 500 (Ibidi) for live cell imaging. Histone H2B-GFP transfected hBM-MSCs were plated onto collagen IV (Sigma-Aldrich) coated plastic or glass coverslips. To induce neural differentiation, cells at passage 3–4 were allowed to adhere to the plates overnight. Basal media was removed the following day and the cells were cultured for 2 days in serum-free media (designated as the neural basal media) consisting in Dulbecco's modified Eagle's medium/F12 (DMEM/F12 Glutamax, Gibco) supplemented with N2-supplement (R&D systems), 0.6% glucose (Sigma-Aldrich), 5mM HEPES (Sigma-Aldrich), 0.5% human serum albumin (Sigma-Aldrich), 0.0002% heparin (Sigma-Aldrich), non-essential amino acid solution (Sigma-Aldrich) and 100 U/ml penicillin-streptomycin (Sigma-Aldrich). On day 3, cells were cultured in neural induction media, consisting in the neural basal media supplemented with 500nM retinoic acid (Sigma-Aldrich) and 1mM dibutyryl cAMP (Sigma-Aldrich). Time-lapse analysis was carried out using a Widefield Leica Thunder-TIRF imager microscope. We perform time-lapse microscopy within the first 70 hours after neural induction media was added directly to the cells. Time-lapse images were obtained with a 40X objective every 10 min. During imaging, cells were enclosed in a chamber maintained at 37°C under a humidified atmosphere of 5% CO2 in air. Data are representative of ten independent experiments.

Immunocytochemistry

A standard immunocytochemical protocol was used as previously described[14,15,17,18]. Histone H2B-GFP transfected hBM-MSCs were plated onto collagen IV (Sigma-Aldrich) coated plastic or glass coverslips and maintained in neural induction media. Cells were rinsed with PBS and fixed in freshly prepared 4% paraformaldehyde (PFA; Sigma-Aldrich). Fixed cells were blocked for 2 h in PBS containing 10% normal horse serum (Gibco) and 0.25% Triton X-100 (Sigma) and incubated overnight at 4 °C with antibodies against β-III-tubulin (TUJ1; 1:500, Covance), Fibrillarin (1/300, Abcam) and Lamin A/C (1/300, GeneTex), in PBS containing 1% normal horse serum and 0.25% Triton X-100. On the next day, cells were rinsed and incubated with the secondary antibodies conjugated with Alexa Fluor® 488 (anti-rabbit; 1:500, Molecular Probes) and Alexa Fluor® 594 (anti-mouse; 1:500, Molecular Probes). Cell nuclei were counterstained with DAPI (0.2 mg/ml in PBS, Molecular Probes). Alexa Fluor 488® phalloidin (Molecular Probes) was used to selectively stains F-actin. Data are representative of ten independent experiments per condition.

Images and Data Analyses

Photograph of visible and fluorescent stained samples were carried out in a Widefield Leica Thunder-TIRF imager microscope equipped with a digital camera or in confocal laser scanning microscope Leica TCS-SP8. We used Leica Application Suite X and Imaris software for image analysis and Filmora Video Editor software for video editing. Photoshop software was used to improve the visibility of fluorescence images without altering the underlying data. Data are representative of ten independent experiments per condition and are expressed as mean ± SD.

Data availability statement: All data generated or analysed during this study are included in this published article (and its supplementary information files).

References

1. Osorio, D. S. & Gomes, E. R. The contemporary nucleus: a trip down memory lane. *Biol. Cell*. **105**, 430-441 (2013).

2. Gundersen, G. G. & Worman, H. J. Nuclear positioning. *Cell*. **152**, 1376-1389 (2013).

3. Skinner, B. M. & Johnson, E. E. Nuclear morphologies: their diversity and functional relevance. *Chromosoma*. **126**, 195-212 (2017).

4. Misteli, T. The Self-Organizing Genome: Principles of Genome Architecture and Function. *Cell*. **183**, 28-45 (2020).

5. Jevtić, P, Edens L. J., Vuković, L. D. & Levy, D. L. Sizing and shaping the nucleus: mechanisms and significance. *Curr. Opin. Cell Biol*. **28**, 16-27 (2014).

6. Kalukula Y, Stephens AD, Lammerding J, Gabriele S. Mechanics and functional consequences of nuclear deformations. *Nat. Rev. Mol. Cell Biol*. **23**, 583-602 (2022).

7. Dupin, I. & Etienne-Manneville, S. Nuclear positioning: mechanisms and functions. *Int. J. Biochem. Cell Biol*. **43**, 1698-1707 (2011).

8. Calero-Cuenca, F. J., Janota, C. S. & Gomes, E. R. Dealing with the nucleus during cell migration. *Curr. Opin. Cell Biol*. **50**, 35-41 (2018).

9. Almonacid, M., Terret, M. E. & Verlhac, M. H. Nuclear positioning as an integrator of cell fate. *Curr. Opin. Cell Biol*. **56**, 122-129 (2019).

10. Deshpande, O. & Telley, I. A. Nuclear positioning during development: Pushing, pulling and flowing. *Semin. Cell Dev. Biol*. **120**,10-21 (2021).

11. Bone, C. R. & Starr, D. A. Nuclear migration events throughout development. *J. Cell Sci*. **129**, 1951-1961 (2016).

12. Zwerger, M., Ho, C. Y. & Lammerding, J. Nuclear mechanics in disease. *Annu. Rev. Biomed. Eng*. **13**, 397-428 (2011).

13. Manda, N. K., Golla, U., Sesham, K., Desai, P., Joshi, S., Patel, S., Nalla, S., Kondam, S.,

Singh, L., Dewansh, D., Manda, H. & Rokana N. Tuning between Nuclear Organization and

Functionality in Health and Disease. *Cells*. 2023 Feb 23;12(5):706. doi: 10.3390/cells12050706.

14. Bueno, C., Ramirez, C., Rodríguez-Lozano, F. J, Tabarés-Seisdedos, R., Rodenas, M.,

Moraleda, J. M., Jones, J. R. & Martinez, S. Human adult periodontal ligament-derived cells

integrate and differentiate after implantation into the adult mammalian brain. *Cell Transplant*. 22,

2017-2028 (2013).

15. Bueno, C., Martínez-Morga, M. & Martínez, S. Non-proliferative neurogenesis in human

periodontal ligament stem cells. *Sci. Rep*. 2019 Dec 2;9(1):18038. doi: 10.1038/s41598-019-

54745-3.

16. Bueno, C. & Martínez, S. Neurogenesis similarities in different human adult stem cells.

Neural Regen. Res. 2021 Jan;16(1):123-124. doi: 10.4103/1673-5374.286967.

17. Bueno, C., Martínez-Morga, M., García-Bernal, D., Moraleda, J. M. & Martínez, S.

Differentiation of human adult-derived stem cells towards a neural lineage involves a

dedifferentiation event prior to differentiation to neural phenotypes. *Sci. Rep*. 2021 Jun

8;11(1):12034. doi: 10.1038/s41598-021-91566-9.

18. Bueno, C., Blanquer, M., García-Bernal, D., Martínez, S. & Moraleda, J. M. Binucleated

human bone marrow-derived mesenchymal cells can be formed during neural-like differentiation

with independence of any cell fusion events. *Sci. Rep*. 2022 Nov 30;12(1):20615. doi:

10.1038/s41598-022-24996-8.

19. Pittenger, M. F., Discher, D. E., Péault, B. M., Phinney, D. G., Hare, J. M. & Caplan, A. I.

Mesenchymal stem cell perspective: cell biology to clinical progress. *NPJ Regen. Med*. 2019 Dec

2;4:22. doi: 10.1038/s41536-019-0083-6.

20. García-Bernal, D. et al. The Current Status of Mesenchymal Stromal Cells: Controversies,

Unresolved Issues and Some Promising Solutions to Improve Their Therapeutic Efficacy. Front

Cell Dev. Biol. 2021 Mar 16;9:650664. doi: 10.3389/fcell.2021.650664.

21. Mollinari, C., Zhao, J., Lupacchini, L., Garaci, E., Merlo, D. & Pei, G. Transdifferentiation:

a new promise for neurodegenerative diseases. *Cell Death Dis*. 2018 Aug 6;9(8):830. doi:

10.1038/s41419-018-0891-4.

22. Choudhary, P., Gupta, A. & Singh, S. Therapeutic advancement in neuronal

transdifferentiation of mesenchymal stromal cells for neurological disorders. *J. Mol. Neurosci*.

https:// doi. org/ 10. 1007/ s12031- 020- 01714-5 (2020).

23. Hernández, R. et al. Differentiation of human mesenchymal stem cells towards neuronal

lineage: clinical trials in nervous system disorders. *Biomol. Ther*. **28**, 34-44 (2020).

24. Woodbury, D., Schwarz, E. J., Prockop, D. J. & Black I. B. Adult rat and human bone marrow

stromal stem cells differentiate into neurons. *J. Neurosci. Res*. **61**, 364-370 (2000).

25. Woodbury, D., Reynolds, K. & Black, I. B. Adult bone marrow stromal stem cells express

germline, ectodermal, endodermal, and mesodermal genes prior to neurogenesis. *J. Neurosci. Res*.

69, 908-917 (2002).

26. Muñoz-Elias, G., Woodbury, D. & Black, I. B. Marrow stromal cells, mitosis and neuronal

differentiation: stem cell and precursor functions. *Stem Cells*. **21**,437-448 (2003).

27. Jeong, J. A. et al. Rapid neural differentiation of human cord blood-derived mesenchymal

stem cells. *Neuroreport*. **15**, 1731-1734 (2004).

28. Suon, S. et al. Transient differentiation of adult human bone marrow cells into neuron-like

cells in culture: development of morphological and biochemical traits is mediated by different

molecular mechanisms. *Stem Cells Dev*. **13**, 625-635 (2004).

29. Hermann, A. et al. Comparative analysis of neuroectodermal differentiation capacity of

human bone marrow stromal cells using various conversion protocols. *J. Neurosci. Res*. **83**, 1502-

1514 (2006).

30. Ning, H., Lin, G., Lue, T. F. & Lin, C. S. Neuron-like differentiation of adipose tissue-derived

stromal cells and vascular smooth muscle cells. *Differentiation*. **74**, 510-518 (2006).

31. Azedi, F. et al. Comparative capability of menstrual blood versus bone marrow derived stem

cells in neural differentiation. *Mol. Biol. Rep*. **44**, 169-182 (2017).

32. Radhakrishnan, S. et al. In vitro transdifferentiation of human adipose tissue-derived stem

cells to neural lineage cells - a stage-specific incidence. *Adipocyte*. **8**, 164-177 (2019).

33. Azizi, A. S., Stokes, D., Augelli, B. J., DiGirolamo, C. & Prockop, D. J. Engraftment and

migration of human bone marrow stromal cells implanted in the brains of albino rats-Similarities

to astrocyte grafts. *Proc. Natl. Acad. Sci. U.S.A*. **95**, 3908-3913 (1998).

34. Kopen, G. C., Prockop, D. J. & Phinney, D. G. Marrow stromal cells migrate throughout

forebrain and cerebellum, and they differentiate into astrocytes after injection into neonatal mouse

brains. *Proc. Natl. Acad. Sci. U.S.A*. **96**, 10711-10716 (1999).

35. Brazelton, T. R., Rossi, F. M., Keshet, G. I. & Blau, H. M. From marrow to brain: Expression

of neuronal phenotypes in adult mice. *Science*. **290**, 1775-1779 (2000).

36. Mezey, E., Chandross, K. J., Harta, G., Maki, R. A. & McKercher, S. R. Turning blood into

brain: Cells bearing neuronal antigens generated in vivo from bone marrow. *Science* **290**, 1779-

1782 (2000).

37. Priller, J. et al. Neogenesis of cerebellar Purkinje neurons from gene-marked bone marrow

cells in vivo. *J. Cell Biol*. **155**, 733-738 (2001).

38. Mezey, E. et al. Transplanted bone marrow generates new neurons in human brains. *Proc.*

Natl. Acad. Sci. U.S.A. **100**, 1364-1369 (2003).

39. Muñoz-Elias, G., Marcus, A. J., Coyne, T. M., Woodbury, D. & Black, I. B. Adult bone

marrow stromal cells in the embryonic brain:Engraftment, migration, differentiation,

differentiation, and long-term survival. *J. Neurosci*. **24**, 4585-4595 (2004).

40. Jopling, C., Boue, S., & Izpisua Belmonte J. C. Dedifferentiation, transdifferentiation and

reprogramming: three routes to regeneration. *Nat. Rev. Mol. Cell Biol*. **12**, 79-89 (2011).

41. Merrell, A. J. & Stanger, B. Z. Adult cell plasticity in vivo: de-differentiation and

transdifferentiation are back in style. *Nat. Rev. Mol. Cell Biol*. **17**, 413-25 (2016).

42. Lu, P., Blesch, A. & Tuszynski, M. H. Induction of bone marrow stromal cells to neurons:

differentiation, transdifferentiation, or artifact? *J. Neurosci. Res*. **77**, 174-191 (2004).

43. Kemp, K., Wilkins, A. & Scolding, N. Cell fusion in the brain: Two cells forward, one cell

back. *Acta Neuropathol*. **128**, 629-638 (2014).

44. Neuhuber, B. et al. Reevaluation of in vitro differentiation protocols for bone marrow stromal

cells: Disruption of actin cytoskeleton induces rapid morphological changes and mimics neuronal

phenotype. *J. Neurosci. Res*. **77**, 192-204 (2004).

45. Bertani, N., Malatesta, P., Volpi, G., Sonego, P. & Perris, R. Neurogenic potential of human

mesenchymal stem cells revisited: Analysis by immunostaining, time-lapse video and microarray.

J. Cell Sci. **118**, 3925-3936 (2005).

46. Krabbe, C., Zimmer, J. & Meyer, M. Neural transdifferentiation of mesenchymal stem cells-

A critical review. *APMIS*. **113**, 831-844 (2005).

47. Doetsch, F., Garcia-verdugo, J. M. & Alvarez-buylla, A. Cellular composition and three-

dimensional organization of the subventricular germinal zone in the adult mammalian brain. *J.*

Neurosci. **17**, 5046-5061 (1997).

48. Wittmann, M. et al. Synaptic activity induces dramatic changes in the geometry of the cell

nucleus: Interplay between nuclear structure, histone H3 phosphorylation, and nuclear calcium

signaling. *J. Neurosci*. **29**, 14687-14700 (2009).

49. Guerrero-Cázares, H. et al. Cytoarchitecture of the lateral ganglionic eminence and rostral

extension of the lateral ventricle in the human fetal brain. *J. Comp. Neurol*. **519**, 1165-1180

(2011).

50. Cebrián-Silla, A. et al. Unique organization of the nuclear envelope in the post-natal quiescent

neural stem cells. *Stem Cell Rep*. **9**, 203–216 (2017).

51. De Leeuw, R., Gruenbaum, Y. & Medalia, O. Nuclear Lamins: Thin Filaments with Major

Functions. *Trends Cell Biol*. **28**, 34-45 (2018).

52. Ochs, R. L., Lischwe, M. A., Spohn, W. H. & Busch, H. Fibrillarin: a new protein of the

nucleolus identified by autoimmune sera. *Biol. Cell*. **54**, 123-133 (1985).

53. Waddington, C. H. The Strategy of the Genes. A Discussion of Some Aspects of Theoretical

Biology (Allen & Unwin, 1957).

54. Raff, M. Adult stem cell plasticity: fact or artifact? *Annu. Rev. Cell Dev. Biol*. **19**, 1-22 (2003).

55. Rajagopal, J. & Stanger, B. Z. Plasticity in the adult: How should the waddington diagram be

applied to regenerating tissues?. *Dev.Cell*. **36**, 133-137 (2016).

56. Kalluri, R. & Weinberg, R. A. The basics of epithelial-mesenchymal transition. *J. Clin.*

Invest. **119**, 1420-1428 (2009).

57. Pei, D., Shu, X., Gassama-Diagne, A. &. Thiery, J. P. Mesenchymal–epithelial transition in

development and reprogramming. *Nat. Cell Biol*. **21**, 44-53 (2019).

58. Yang, J. et al. EMT International Association (TEMTIA). Guidelines and definitions for

research on epithelial-mesenchymal transition. *Nat. Rev. Mol. Cell Biol*. **21**, 341-352 (2020).

59. Reid, A. & Tursun, B. Transdifferentiation: do transition states lie on the path of

development? *Curr. Opin. Syst. Biol*. **11**, 18-23 (2018).

60. Alvarez-Dolado, M. et al. Fusion of bone-marrow-derived cells with Purkinje neurons,

cardiomyocytes and hepatocytes. *Nature*. **425**, 968-973 (2003).

61. Weimann, J. M., Johansson, C. B., Trejo, A. & Blau, H. M. Stable reprogrammed

heterokaryons form spontaneously in Purkinje neurons after bone marrow transplant. *Nat. Cell*

Biol. **5**, 959-966 (2003).

62. Magrassi, L. et al. Induction and survival of binucleated Purkinje neurons by selective

damage and aging. *J. Neurosci*. **27**, 9885-9892 (2007).

63. Altman, J. Autoradiographic investigation of cell proliferation in the brains of rats and cats.

Anat. Rec. **145**, 573-591 (1963).

64. Das, G. D. Binucleated neurons in the central nervous system of the laboratory animals.

Experientia. **33**, 1179-1180 (1977).

65. Ribak, C. E. & Seress, L. Five types of basket cell in the hippocampal dentate gyrus: A

combined Golgi and electron microscopic study. *J. Neurocytol*. **12**, 577-597 (1983).

66. Portiansky, E. L., Barbeito, C. G., Flamini, M. A., Gimeno, E. J. & Goya, R. G. Presence of

binucleate neurons in the spinal cord ofyoung and senile rats. *Acta Neuropathol*. **112**, 647-649

(2006).

67. Jimenez-Acosta, M. A., Hernandez, L. J. R., Cristerna, M. L. P., Tapia-Ramirez, J. & Meraz-

Rios, M. A. Review: Neuronal differentiation protocols of mesenchymal stem cells. *Adv. Biosci.*

Biotechnol. **13**, 15-71 (2022).

68. Eguizabal, C., Montserrat, N., Veiga, A. & Izpisua-Belmonte, J. C. Dedifferentiation,

transdifferentiation, and reprogramming: future directions in regenerative medicine. *Semin*

Reprod Med. **31**, 82-94 (2013).

69. Mollinari, C., Zhao, J., Lupacchini, L., Garaci, E., Merlo, D. & Pei, G. Transdifferentiation:

a new promise for neurodegenerative diseases. *Cell Death Dis*. 2018 Aug 6;9(8):830. doi:

10.1038/s41419-018-0891-4.

70. Kosodo Y. Interkinetic nuclear migration: beyond a hallmark of neurogenesis. *Cell Mol. Life*

Sci. **69**, 2727-2738 (2012).

71. Spear, P. C. & Erickson, C. A. Interkinetic nuclear migration: a mysterious process in search

of a function. *Dev Growth Differ*. **54**, 306-316 (2012).

72. Hoffmann, K., Sperling, K., Olins, A. L. & Olins, D. E. The granulocyte nucleus and lamin B

receptor: Avoiding the ovoid. *Chromosoma*. **116**, 227-235 (2007).

73. Kachi, K. & French, S. W. The connection between the nuclei of binucleated hepatocytes: An

ultrastructural study. *J. Submicrosc. Cytol. Pathol*. **26**, 163–172 (1994).

74. Paradis, A. N., Gay, M. S. & Zhang, L. Binucleation of cardiomyocytes: The transition from

a proliferative to a terminally differentiated state. *Drug Discov. Today*. **19**, 602-609 (2014).

75. Kanda, T., Sullivan, K. F. & Wahl, G. M. Histone-GFP fusion protein enables sensitive

analysis of chromosome dynamics in living mammalian cells. *Curr. Biol*. **8**, 377-385 (1998).

Acknowledgements: This study was supported by the Spanish Ministry of Science and

Innovation, the Carlos III Health Institute (ISCIII), through The Spanish Network of Advanced

Therapies (TERAV), projects RD21/0017/0001 and RD21/0017/0017, co-funded by European

Union "NextGenerationEU. Plan de Recuperación Transformación y Resiliencia". It was also

partially supported by Research Grant of the Universidad Católica San Antonio de Murcia

(UCAM). We greatly appreciate the technical assistance of Microscopy Section and the Tissue

Culture Service of the University of Murcia.

Author contributions: C.B. conceived of the study, designed the study, carried out the molecular

lab work and drafted the manuscript; D.G-B and M.B. designed experiments, participated in data

analysis and helped draft the manuscript; S.M. and J.M.M. helped draft the manuscript and

financial support.

Competing interests: The authors declare no competing interests.

Materials & Correspondence. Carlos Bueno, PhD., Medicine Department and Hematopoietic

Transplant and Cellular Therapy Unit, Institute of Biomedical Research (IMIB), University of

Murcia, Faculty of Medicine, Murcia, 30120, Spain. Tel.: 0034-968885013. Fax: 00-34-

868884150. E-mail: carlos.bueno@um.es

Figure legends

Fig. 1. **Morphology of hBM-MSCs cultured in basal medium**. **a**, Undifferentiated hBM-MSCs

exhibited a fibroblast-like morphology with β-III tubulin microtubules and actin microfilaments

oriented parallel to the longitudinal axis of the cell. **b**, During interphase, hBM-MSCs displayed

a flattened, ellipsoidal nucleus, often located in the center of the cell. **c**, Distribution of fibrillarin-

positive specks in the nuclei of undifferentiated hBM-MSCs. **d**, Immunocytochemical analysis

revealed that the inner surface of the nuclear envelope is lined by the nuclear lamina. Scale bar:

10 μm.

Fig. 2. **Nuclear movement generated cellular protrusions that appeared and disappeared**

from the surface of hBM-MSC-derived intermediate cells. **a**, Time-lapse imaging showed that

when hBM-MSCs were exposed to a neural induction medium, they rapidly reshaped from a flat

to a spherical morphology. Subsequently, we observed hBM-MSC-derived intermediate cells

(white asterisk) in which nuclear movements generate only one cell protrusion (white arrow), and

hBM-MSC-derived intermediate cells (yellow asterisk) in which nuclear movements alternately

generate one or two cellular protrusions (yellow arrows). **b**, We found that when a hBM-MSC-

derived intermediate cell has a nucleus without lobes, its movement within the cell generates only

one cell protrusion (yellow arrow). **c**, However, if the hBM-MSC-derived intermediate cell has a

lobed nucleus it will generate one or two cellular protrusions depending on how it moves within

the cell (yellow arrows). Scale bar: 10 μm. PhC: Phase-contrast photomicrographs.

Fig. 3. **Characterization of hBM-MSC-derived intermediate cell nuclei**. **a,**

Immunocytochemical analysis revealed that hBM-MSC-derived intermediate cells primarily have

three different nuclear morphologies: Tail-less nuclei (white asterisk), tailed nuclei (green

asterisk) and lobed nuclei (yellow asterisk). Tail-less and tailed nuclei movements generate only

one cell protrusion (white and green arrows respectively). However, lobed nuclei movements

generate one or two cellular protrusions depending on how it moves within the cell (yellow

arrows). Confocal microscopy analysis and 3D reconstruction revealed that there are small

variations in both shape and size within the three types of nuclear morphology (**b**) and that the

lobes of the lobed nuclei can be located in different z-planes (**c**). **d**, Immunocytochemical analysis

revealed that tail-less nuclei and tailed nuclei contained one or two fibrillarin positive specks

while lobed nuclei contained one or two fibrillarin-positive specks in each lobe. **e**, The inner

surface of the nuclear envelope is lined by the nuclear lamina. Scale bar: 10 μm. PhC: Phase-

contrast photomicrographs.

Fig. 4. **Cytoskeletal organization of hBM-MSC-derived intermediate cells**. **a**,

Immunocytochemical analysis revealed that actin microfilaments and β-III tubulin microtubules

are no longer oriented parallel to the longitudinal axis of the hBM-MSC-derived intermediate

cells. **b**. Confocal microscopy analysis and 3D reconstruction revealed that cell protrusions are

almost devoid of actin microfilaments and that the β-III tubulin protein is concentrated at the cell

protrusion rim (black arrows). Scale bar: 10 μm.

Fig. 5. **Tail-less nuclei formation**. Time-lapse imaging revealed that tail-less nuclei are formed

by a single nuclear remodeling that occurs as the cell reshapes from a flat to a spherical

morphology, positioning the nucleus in a peripheral position within the cell (white arrows).

Subsequently, the tail-less nucleus began to move within the hBM-MSC-derived intermediate

cells (yellow arrows). Scale bar: 10 μm.

Fig. 6. **Tailed nuclei formation by two nuclear remodeling sequences**. Time-lapse images show

that first a nuclear remodeling sequence occurs as the cell reshapes from a flat to a spherical

morphology, positioning the nucleus in a peripheral position within the cell (white arrows). Next,

the cell nucleus moves (green arrows) and undergoes a second nuclear sequence in which a tailed

process in the nucleus is formed (yellow arrows). Finally, the tailed nucleus began to move within

the hBM-MSC-derived intermediate cells (green arrows). Scale bar: 10 μm.

Fig. 7. **The tails of the tailed nuclei can move within the hBM-MSC-derived intermediate cells, switching shape and size and even move in different z-planes**. **a**, Time-lapse imaging revealed that as the tailed nuclei move within the cell, the tails can switch shape and size (yellow arrows). **b**, Time-lapse images also showed that the tails even appear to move in different z-planes. Scale bar: 10 μm.

Fig. 8. **Lobed nuclei formation by two nuclear remodeling sequences**. Time-lapse imaging showed that first a nuclear remodeling sequence occurs as the cell reshapes from a flat to a spherical morphology, positioning the nucleus in a peripheral position within the cell (white arrows). Subsequently, the cell nucleus moves (green arrows) and undergoes a second nuclear sequence in which a lobed nucleus is formed (yellow arrows). Afterwards, the lobed nuclei began to move within the hBM-MSC-derived intermediate cells (green arrows). Finally, we also noted that lobed nuclei can switch shape while moving within the cell, becoming tailed nuclei (blue arrows). Scale bar: 10 μm.

Fig. 9. **hBM-MSC-derived intermediate cells contact each other through their cell nuclei**. Time-lapse imaging revealed that hBM-MSC-derived intermediate cell nuclei move within the cell generating cellular protrusions as they attempt to contact the cells around them, mainly observing contacts between the nuclei (green arrows). Scale bar: 10 μm.

Fig. 10. **Tail-less nuclei, tailed nuclei and lobed nuclei contact each other**. Confocal microscopy analysis (**a**) and 3D reconstruction (**b**) revealed that tail-less nuclei (white arrow), tailed nuclei (green arrow) and lobed nuclei (yellow arrow) contact each other. Scale bar: 10 μm.

a

PhC   DAPI   β-III-tubulin   β-III-tubulin/DAPI

b

Tail-less nuclei   Lobed nuclei   Tailed nuclei

DAPI

3D Confocal

Colour-coding Depth
12 µm
10 µm
8 µm
6 µm
4 µm
2 µm
0 µm

c

3D Confocal

DAPI   β-III-tubulin   Merge

DAPI

Colour-coding Depth
12 µm
10 µm
8 µm
6 µm
4 µm
2 µm
0 µm

d

Fibrillarin/β-III-tubulin/DAPI

Tail-less nuclei

Tailed nuclei

Lobed nuclei

e

Laminin A/C/β-III-tubulin/DAPI

Tail-less nuclei

Tailed nuclei

Lobed nuclei

a

DAPI | β-III-tubulin | PhC

b

DAPI | β-III-tubulin | Merge

Depth Coding

12 µm
10 µm
8 µm
6 µm
4 µm
2 µm
0 µm