The Mediterranean mussel, *Mytilus galloprovincialis*, a novel model for developmental
studies of mollusks

Angelica Miglioli[1$†], Marion Tredez[1$], Manon Boosten[1,2], Camille Sant[1,2], João E. Carvalho[1],
Philippe Dru[1], Laura Canesi[3], Michael Schubert[1*], and Rémi Dumollard[1*]

1- Laboratoire de Biologie du Développement (LBDV), Institut de la Mer de Villefranche
(IMEV), Sorbonne Université, CNRS, Villefranche-sur-Mer, France

2- Laboratoire d'Océanologie de Villefranche (LOV), Institut de la Mer de Villefranche (IMEV),
Sorbonne Université, CNRS, Villefranche-sur-Mer, France

3- Università degli Studi di Genova, Dipartimento di Scienze della Terra dell'Ambiente e della
Vita (DISTAV), Genova, Italy

[$] Co-First authors

[*] Co-Last authors

[†] Corresponding author: angelica.miglioli@imev-mer.fr

Keywords: bivalve mollusks, embryonic development, trochophore, shell field, transcriptome,
hybridization chain reaction

Summary statement: Resources and techniques are described for the Mediterranean
mussel *Mytilus galloprovincialis*, which, together, establish a novel model system for studying
mollusk development and animal evolution.

Running title: A novel mollusk model system: the Mediterranean mussel

Abstract

A model organism in developmental biology is defined by its experimental amenability as well as by resources created for the model system by the scientific community. For the most powerful models, the combination of both has already yielded a thorough understanding of development. However, the number of developmental model systems is still limited, and their phylogenetic distribution is heavily biased. Members of one of the largest animal *phylum*, the mollusks, for example, have long been neglected as developmental model organisms. To remedy this shortcoming, we produced a detailed developmental transcriptome for the bivalve mollusk *Mytilus galloprovincialis*. Our high-quality transcriptome allowed us to identify transcriptomic signature of developmental transitions and perform a first comparison with *Crassostrea gigas* that could further be applied for multi-species comparisons. To allow co-labelling studies, we optimized protocols for immunohistochemistry and hybridization chain reaction and combined both techniques to obtain high-resolution co-expression maps of developmental genes. The resources and protocols we describe here thus represent an enormous boost for the establishment of the Mediterranean mussel as a laboratory model in developmental biology.

1. Introduction

The development of aquatic invertebrates has inspired the curiosity of biologists for well over a century (Hunt and Maslakova, 2017; Nielsen, 2005; Nielsen, 2009; Robertis and Tejeda-mu, 2022; Wang et al., 2020). Morphological similarities between larvae have, for example, been used as taxonomic characters and comparisons between larval types have yielded highly influential hypotheses on animal evolution (Tyler J. Carrier, Adam M. Reitzel, 2018). One such larval type, the trochophore, is widely distributed amongst the lophotrochozoans and may have already been present in the last common ancestor of this large clade of bilaterian animals (De Robertis and Tejeda-Muñoz, 2022). However, despite decades of studies (Martín et al., 2021; Nielsen, 2005; Page, 2009; Rawlinson, 2010; Tyler J. Carrier, Adam M. Reitzel, 2018; Wada et al., 2020; Xu et al., 2016; Yang et al., 2020), our understanding of the evolutionary origin of the trochophore larva as well as of the biological processes underpinning its ontogeny and transition to the adult body remains extremely fragmentary (Robertis and Tejeda-mu, 2022; Tyler J. Carrier, Adam M. Reitzel, 2018; Wada et al., 2020; Wang et al., 2020; Wu et al., 2019; Xu et al., 2016; Yang et al., 2020). Although multiple genome and transcriptome sequencing projects have provided abundant data from diverse organisms, most of these datasets have yet to be exploited to provide valuable insights into the overall conservation of trochophore development between different taxa (Maureen Liu et al., 2014; Sayers et al., 2021; Wada et al., 2020). A major limitation for comparisons seems to be that lophotrochozoan development is characterized by the deployment of species-specific gene sets. This fact makes the identification of comparable larval phases, even between closely related species of the same class, quite difficult (Wada et al., 2020; Wu et al., 2019; Xu et al., 2016; Yang et al., 2020).

Within the lophotrochozoans, the mollusks are the largest and anatomically most diverse phylum (Yang et al., 2020). Yet, as of today, the scientific community still lacks an amenable laboratory model for studying the mechanisms of mollusk development. Reasons include the limited availability of genomic and transcriptomic resources and the complexity of establishing even basic techniques, such as immunocytochemistry and *in situ* hybridization (Yang et al., 2020). It has been proposed that the overall fragility of mollusk larvae as well as the presence of a shell, at least in conchiferan (i.e., shell-bearing) mollusks (Kocot et al., 2016; Leclère et al., 2019; Miglioli et al., 2019; Miglioli et al., 2021a; Sayers et al., 2021; Wada et al., 2020; Yang et al., 2020), have hindered past efforts to devise more elaborate resources and

techniques for representatives of this phylum (Sayers et al., 2021; Tyler J. Carrier, Adam M. Reitzel, 2018; Wada et al., 2020). Altogether, this has slowed research into the molecular aspects of trochophore development in mollusks and thus significantly limits our capacity for comparisons of trochophore development between mollusks and other taxa (Wada et al., 2020; Yang et al., 2020). The Mediterranean mussel, *Mytilus galloprovincialis*, is an established model in environmental studies and a species of great ecological and economic interest with a worldwide distribution (Brzozowska et al., 2012; Daguin Claire, 2000; FAO, 2022). Large scale larval cultures of *M. galloprovincialis* are easily achievable in the laboratory, following standardized culture protocols by the International Standardization Organization (ISO) and the American Society for Testing Materials (ASTM) (ASTM E724-21, 2021; ISO 17244:2015, 2020). *M. galloprovincialis* is also the first metazoan with a published pangenome (Gerdol et al., 2020), a feature of significant importance for researchers interested in genome evolution. The presence of detailed culture protocols and of a well-annotated genome make *M. galloprovincialis* a promising model organism for developmental studies. In addition, basic protocols for pharmacological treatments, quantification of shell biogenesis, quantitative polymerase chain reaction (qPCR) as well as immunohistochemistry and *in situ* hybridization have recently been described for developing *M. galloprovincialis* embryos and larvae (Miglioli et al., 2019; Miglioli et al., 2021b; Miglioli et al., 2021a). These technical advances are promising basis for using the Mediterranean mussel as model system for developmental studies.

In this work, we provide a compendium of updated protocols and novel resources to significantly facilitate access to and work with *M. galloprovincialis* embryos and larvae. We thus compiled a detailed developmental transcriptome and validated it with a standardized qPCR protocol. We further established a novel protocol for multi-color, fluorescent whole-mount *in situ* hybridization, using the hybridization chain reaction (HCR) approach (Choi et al., 2014), and successfully coupled this *in situ* hybridization approach with immunohistochemistry assays. Importantly, the developmental transcriptome of *M. galloprovincialis* allowed us to define major developmental periods and transitions, results that were further confirmed into other bivalve mollusks. These results provide a basis for comparative developmental studies of lophotrochozoans on a transcriptomic level. The novel *in situ* hybridization and immunohistochemistry protocols yielded detailed information on the expression of genes marking the developmental transitions, for example, in the early shell

field, apical sensory cells, peripheral mantle cells, and hindgut. In summary, this work represents a valuable resource for future work with *M. galloprovincialis* embryos and larvae, setting the stage for establishing this animal as a mollusk model system. As such, this work will have a significant impact on developmental biology and associated research fields.

2. Results

2.1. Morphological staging of *Mytilus galloprovincialis* early development and description of transcriptome libraries

The developmental transcriptome of *M. galloprovincialis* embryos and larvae consists of 15 developmental stages (i.e. a total of 30 libraries from two independent biological replicates).The samples were obtained from standard culture conditions (ASTM E724-21, 2021; ISO 17244:2015, 2020), from unfertilized egg to 72 hours post fertilization (hpf) at 16°C (Fig. 1A). As previously described in for other Mytilids, the *M. galloprovincialis* zygote undergoes holoblastic, unequal cleavage (Kurita et al., 2009). The micromeres form at the animal pole of the embryo at 4 hpf, and subsequent cleavage follows a spiral pattern (Kurita *et al.*, 2009). By 12 hpf, the macromeres at the vegetal pole start to invaginate, indicating the start of gastrulation. At 16 hpf, the stomodeum (or presumptive mouth) has moved anteriorly, and a rosette of dorsal ectodermal cells form an invagination indicating the onset of shell field ontogeny (Kniprath, 1980; Kniprath, 1981; Miglioli et al., 2019). At 20 hpf, the columnar cells of the shell field thicken, creating the impression of a deeper invagination. At 24 hpf, the larvae have the stereotypical morphology of a trochophore, with a ventral stomodeum and a dorsal shell field that is now completing the evagination process and secreting the organic matrix of the larval shell (Kniprath, 1980; Miglioli et al., 2019). By 28 hpf, the hinge region of the shell field is completely flat. At 32 hpf, a cavity appears in the hinge region of the larva, and, at 36 hpf, it expands and seems to connect to the former stomodeum (that has now moved anteriorly) and to the anal cavity on the posterior side. At 40 hpf, the D margin of the shell becomes clearly visible and only the velum (or prototroch) of the larva is protruding. By 44 hpf, the velum is enclosed in the D-shaped shell. Thereafter, no major morphological changes are observable, excepting the progressive outgrowth of the shell with respect to the mantle tissue along the D border (Fig. 1A). In summary, our transcriptome is representative of all developmental stages in bivalves' early development.

2.2. Transcriptome assembly and gene expression dynamics

In this section are described the described the early gene expression dynamics characterizing the early development of *M. galloprovincialis*. More than 31 million paired end clean reads per library were mapped on the published *M. galloprovincialis* genome (Gerdol et al., 2020) with an average input length of 200 bp (Supplementary Table 1). More than 85% of the reads were mapped, with 71% of them assigned to a unique locus (Supplementary Table 1). The output gene count matrix showed 52'422 expressed genes (sum of gene raw counts in all libraries > 0), of a total of 60'302, throughout the libraries (Supplementary Table 2). Further analysis of the number of genes expressed throughout development is presented in Fig. 1B. The minimum number of expressed genes (EGs) was found in unfertilized eggs (which we named as 0 hpf), around 25'000, while the highest was found at the late D-Veliger stage (at 72 hpf), around 40'000, indicating that at least 15'000 genes (that is about 25% of the genome) are transcriptionally modulated during early development of *M. galloprovincialis* (Fig. 1B). The number of EGs increased dramatically during the first 12 hours of development, with the two replicates yielding similar results. Thereafter, a slow but steady increase of EGs was observed until the last sampling point. Heatmap representation of the Euclidean distances across stages and the principal component analysis (PCA) of the most variably expressed genes validated sample reproducibility (Fig. 1C and1D). The heatmap shows that the libraries clustered by developmental time and not by biological replicate (Fig. 1C). Additionally, library clustering revealed two main clusters, one before and one after 32 hpf, indicating a potential mid-development transcriptomic shift, within the trochophore stages (Fig. 1C). The PCA showed a continuous transcriptomic progression from 0 hpf to 72 hpf, along the axis of the first principal component, indicating that the maximum variation is found between libraries from 0 to 24 hpf and from 32 to 72 (Fig. 1D), which was consistent with a potential transcriptomic shift during early trochophore development (Fig. 1A). The majority of the variance is explained by PC1 (78%) that is mainly correlated to developmental timing, lending support to the observation that morphological changes during development are driven by transcriptomic changes on a molecular level.

2.3. Differential expression analysis and developmental clustering

This section describes transcriptomic staging of *M. galloprovincialis* early development based on differential expression analyses. Differential expression analysis was performed by

pairwise comparison of samples of consecutive stages. The analysis identified 11,996 differentially expressed genes (DEGs) throughout *M. galloprovincialis* development (Supplementary File 1 and Supplementary Table 3). Hierarchical clustering of the libraries based on DEG expression values identified statistically (au, unbiased adjusted p-value > 0.95) distinct developmental phases (Fig. 2A) and divided the 30 libraries in two main clusters: one grouping all libraries from 0 hpf to 16 hpf (embryonic phase) and one regrouping all libraries from 20 hpf to 72 hpf (larval phase). In the embryonic phase of development, the libraries of 0 hpf and 4 hpf were clearly distinct from the ones covering 8 hpf to 16 hpf, corresponding to, respectively, early cleavage (zygote) and gastrulation (Fig. 2A). In the second major cluster, three sub-clusters were observable. The first one was formed by libraries corresponding to three trochophore stages, from 20 hpf to 28 hpf, with three older trochophore-like larvae (advanced trochophore), from 32 hpf to 40 hpf, establishing a second cluster. The third cluster grouped the libraries of the four stages with a D-Veliger morphology, from 44 hpf to 72 hpf. The number of DEGs varied as development proceeded, with an overall decreasing trend (Fig. 2B). About 55% of DEGs were found to be exclusively up-regulated, 25% exclusively down-regulated, and the remaining 20% alternated between up- and down-regulation. Of note, 40% of DEGs were differentially expressed more than once throughout development (Fig. 2B insert). More than 1.500 DEGs were already detected at 4 hpf, indicating an early initiation of zygotic gene expression. The highest number of DEGs was reached at 8 hpf and it was followed by a progressive decrease as development proceeded. Following the differential expression analysis, DEGs were grouped as a function of their most recurrent expression profiles (Fig. 2C and 2D, Supplementary Tables 4 and 5). The analysis identified 56 clusters with more than 15 genes and of the top 6 clusters, based on the number of DEGs, we found 5 to consist of genes with low maternal (0 hpf) expression levels. This indicates that the DEGs identified are predominantly derived from zygotic expression (Fig. 2C). An expression level-based association of gene clusters with developmental periods (Supplementary Table 4) showed that 31% of the clusters reached their highest expression levels at the zygote, 32% at the gastrula, 14% at the trochophore, 7% at the advanced trochophore, and 25% at the D-veliger stage (Fig. 2D). These results indicate that Zygote and D-Veliger stages are characterized by genes with a higher diversity of expression profiles than Trochophore and Advanced Trochophore. Moreover, the analysis shows that expression levels of individual clusters are generally skewed towards earlier developmental periods and

that cluster expression levels might be directly correlated with key morphological events
during development of both embryo and larva. Finally, the analysis shows that highly dynamic
gene expression patterns characterize the early development of *Mytilus galloprovincialis*.

2.4. qPCR and *in situ* hybridization of selected genes
To validate the transcriptomic analyses, the differential expression analyses were
corroborated by qPCR experiments carried out with material from both biological replicates.
Four target genes were selected based on expression profiles. The *Foxb2*
(MGAL_10B093191) and *Wnt8a* (MGAL_10B085403) genes were selected as members of
one of the top 6 clusters (Fig. 2C), characterized, respectively, by high and low mean
expression. *SLC46A3* (MGAL_10B020966) and *SYTL4* (MGAL_10B044489) were selected
because they show a highly dynamic expression profile (Supplementary Tables 3 and 4). For
each one of these four genes, the qPCR experiments validated both the levels and the
temporal patterns of their expression (Fig. 2E). Given their shared temporal expression
patterns, we further assessed the spatiotemporal expression of *Foxb2* and *Wnt8a* in relation
to each other and relative to immunoreactivity of serotonin (5-HT), a regional marker of the
developing apical sensory organ. To do so, we established a fluorescent double *in situ*
hybridization protocol based on the hybridization chain reaction (HCR) protocol for *M.*
galloprovincialis and combined it with an immunohistochemistry assay for the
neurotransmitter serotonin (Fig. 2E and Supplementary Figure 1). Co-labeling with serotonin
allowed a standardized orientation of embryos and larvae, which significantly facilitated the
interpretation of *in situ* hybridization signals. We thus found that, while *Wnt8a* was detectable
in a discrete group of cells that will give rise to one of the posterior neural ganglions of the
larva, *Foxb2* was detectable in the stomodeal and trochal area. These observations were
consistent with the expression patterns described for these two genes in other
lophotrochozoan species (Lartillot, *et al.*, 2002; Tomer *et al.*, 2010; Marlow *et al.*, 2014). In
summary, qPCR analysis and *in situ* hybridization confirmed the developmental trends
identified by transcriptomic analysis.

2.5. Qualitative analysis of differentially expressed genes
We next performed a qualitative analysis of the DEGs based on functional domains and gene
ontologies (GOs) (Fig. 3). DEGs were screened for transcription factor domains (Fig. 3A),
domains known to be involved in shell formation (Fig. 3B) as well as for receptors, ion

channels, and neurotransmitter transporters (Fig. 3C). In total, 711 transcription factors of 65 domains were annotated, with the most recurrent (>20 genes) being: zinc fingers, homeodomains, helix-loop-helix and fork head domains (Fig. 3A, Supplementary Table 6). With respect to shell formation, 779 genes were found to contain domains associated with shell matrix proteins and calcification (Fig. 3B) (Ramesh et al., 2019). The most recurrent domains included calcium binding proteins (EF-hand and calponin), Von Willebrand factors (WKA), extracellular matrix proteins (fibronectin), carbohydrate binding proteins (lectin and concanavalin), chitin binding proteins (CB), ion channels, and transporters (Fig. 3B, Supplementary Table 5). Regarding receptors, 225 DEGs were annotated as G protein-coupled receptors of the rhodopsin and secretin families, 92 as ephrin-A receptors, 23 belonged to the Nuclear Receptor superfamily, and other DEGs included scavenger, low density (LD) lipoprotein and TGFβII receptors (Fig. 3C). Surprisingly, as many as 125 DEGs were predicted to be involved in neurotransmitter transport and reception (Fig. 3C, Supplementary Table 3). A gene ontology enrichment analysis was subsequently performed on the DEGs to identify stage-enriched biological processes (Fig. 3D). The GO enrichment analysis for biological processes identified 6 GO clusters: transmembrane transport, actin polymerization, cell adhesion, receptor-mediated signal transduction, biosynthesis, and transcription-coupled repair. While most biological processes did not exhibit a clear association to any developmental phase, transmembrane transport and receptor-mediated signal transduction were enriched in the trochophore and advanced trochophore stages (20 hpf to 28 hpf and 32 hpf to 40 hpf). Actin polymerization and transcription-coupled repair were mildly enriched in embryonic development (zygote and gastrula) and biosynthesis in the larval phase (early trochophore onward). Altogether, the results of the quantitative analysis confirm the relevance of the identified DEGs with respect to the major transcriptional (i.e., TFs) and transcriptomic (i.e., SMPs) programs governing early development of bivalve mollusks.

2.6. Gene clustering and transcriptomic signatures of early developmental transitions
Soft clustering of expressed gene was performed to reveal potential transcriptomic dynamics during development. The analysis identified nine soft clusters (Fig. 4A), which were named according to the morphotypes previously described (Fig. 1A). While the embryonic gene clusters nicely recovered the hierarchical clusters of the embryonic phase, the larval soft clusters predominantly highlighted stages in between the developmental phases identified by

hierarchical clustering and thereby representative of potential developmental transitions. The Early Trochophore gene cluster matched the transition from the Gastrula to Trochophore hierarchical clusters (16 hpf to 20 hpf, Transition 1); the Advanced Trochophore gene clusters matched the transition from Trochophore to Advanced Trochophore hierarchical clusters (28 hpf to 32 hpf, Transition 2); the Veliger gene cluster matched the transition from Advanced Trochophore to D-Veliger hierarchical clusters (40 hpf to 44 hpf, Transition 3) (Fig. 4A, Supplementary Table 7). To identify representative genes for these three developmental transitions, we isolated the genes of the three soft clusters that were also differentially expressed at the given transition stages (Supplementary Table 8). We found 98, 11, and 8 genes for Transition 1, 2, and 3, respectively (Fig. 4B, Supplementary Table 3). To assess whether similar transitions also occurred in other bivalve species, we search for homologous genes in the pacific oyster *Crassostrea gigas* and analyzed their developmental expression with the publicly available transcriptomic data (Fig. 4C, Supplementary Table 8) (Liu et al., 2021). The results indicate that about 20 genes were highly expressed *in C. gigas* gastrula and trochophore 1, thus corresponding to Transition 1; however, the majority were expressed earlier or later in development (Fig. 4C). For Transition 2, only few genes indicated a peak in expression between *C. gigas* trochophore 2 and trochophore 4 (Fig. 4C). Concerning Transition 3, most genes were highly expressed in libraries reconducible to the passage from the *C. gigas* advanced trochophore (trochophore 5) to early Veliger (early D-veliger 1) (Fig. 4C). All in all, our unbiased clustering of genes and samples revealed three uncharted developmental transitions in *M. galloprovincialis* early larval development, hardly identifiable based on morphological observations, which might be partially co-opted in other bivalves.

2.7. Tissue localization of conserved transition genes by *in situ* hybridization

To shed light on the genes co-opted by both bivalves during developmental transitions and investigate their potential relationship to the developmental process, a set of homologous 'transition marker genes' for the two bivalves was isolated based on differential expression (Log fold change >2) in *Mytilus galloprovincialis* and expression profile (z-score value becoming positive) in *C. gigas* across the samples constituting each transition (Table 1, Supplementary Figure 2, Supplementary File 1), and their expression pattern was investigated by Hybridization chain reaction (HCR) at the given times of transition in *M. galloprovincialis* larvae. The genes were named according to the BLAST results of their

predicted proteins in the Uniprot and NCBI databases (Table 1, Supplementary Table 3). As Transition 1 contained a significantly higher number of homologous genes, we selected 8 genes for Transition 1 and 2 genes each for Transitions 2 and 3.

	*Mg*ID	*Cg*ID	*Mg* annotation
Transition 1	MGAL_10B029885	CGI_10016326	T-box transcription factor 2b (*Tbx2b*)
	MGAL_10B094294	CGI_10003121	ETS-related gene (*Erg*)
	MGAL_10B037868	CGI_10025712	Not annotated 03 (*NA03*)
	MGAL_10B039707	CGI_10009468	Tyrosinase 2 (*TYRO2*)
	MGAL_10B009273	CGI_10027254	Solute carrier family 4 member 4 (*SLC4A4*)
	MGAL_10B027895	CGI_10026985	Deleted In Malignant Brain Tumors 1 (*Dmbt1*)
	MGAL_10B078802	CGI_10013987	Peroxidasin (*Pxdn*)
	MGAL_10B078617	CGI_10021520	Hypothetical predicted protein (*HPP*)
Transition 2	MGAL_10B055887	CGI_10015018	Solute carrier family 26 member 5 (*SLC26A5*)
	MGAL_10B039344	CGI_10021966	Glutamate Ionotropic Receptor NMDA Type Subunit 3A (*Grin3a*)
Transition 3	MGAL_10B028954	CGI_10015228	*EF-hand*
	MGAL_10B070218	CGI_10024466	Collagen VI-like (*COLVI*)

Table 1. List of homologous transition genes in *M. galloprovincialis* (*Mg*ID) and *C. gigas* (*Cg*ID) selected for *in situ* hybridization in *M. galloprovincialis*

The expression of these genes was analyzed at the corresponding transition stages. At 20 hpf the transcription factor genes *Tbx2b* and *Erg* were detectable in cells surrounding the stomodeum and in the apical tuft, with *Tbx2b* also being expressed in the hindgut (Fig. 5A). Differently, a majority of the conserved markers of Transition 1 were expressed in or in association with the shell field. Even though, these genes are expressed in a similar domain, the shell field genes marked several different cell types: islets of cells that elongated from invaginated tissue (*NA03* at 20 hpf), column-like cells (*Dmbt1* at 16 and 20 hpf), groups of cells surrounding the invaginating shell field (*HPP* at 16 hpf, *TYRO2* at 16 and 20 hpf and *Pxdn* at 16hpf), and layers of cells covering the invaginated tissue (*SLC4A4* at 20 hpf) (Fig. 5A). At 28 hpf the Transition 2 marker *SLC26A5*, was detectable in two symmetric groups of ectodermal cells, while *Grin3a* was weakly expressed throughout the post-trochal larva, with a

small group of anterior cells beneath the apical tuft expressing the gene more conspicuously

(Fig. 5B). The two conserved markers of Transition 3 were both expressed symmetrically with

respect to the hinge axis, at the 44hpf. Thus, while *EF-hand* was expressed in the velum, the

esophagus, and in a dense group of cells close to the anus (Fig. 5C), *COLVI* was detectable

in individual cells of the veliger larvae along the D-border of the mantle (Fig. 5C).

3. Discussion

3.1. Fine-grained   transcriptomic   dataset   combined   with   thorough   co-labelling

techniques provide valuable insight into bivalve mollusk development

The developmental transcriptome of *M. galloprovincialis* presented in this study is currently

unmatched in terms of sampling depth and detail with respect to the embryo and larva

morphotypes represented in the dataset (Fig. 1). The sampling schedule in 4-hour intervals

prevented biased sampling and overrepresentation of stereotypical stages (such as gastrula,

trochophore or veliger), leading to the generation of a gene expression dataset whose

dynamics and variability are vastly dictated by developmental timing and consistent among

biological replicates (Fig. 1B). The results of our differentially expression analysis, indicating a

decrease in the number of DEGs as development proceeds, is seemingly in contrast with the

findings of previous studies (Foulon et al., 2019; Moreira et al., 2018; Niu et al., 2016; Núñez-

Acuña et al., 2022; Núñez-Acuña et al., 2022; Song et al., 2016a); however, it is important to

note that distinct sampling schedules (i.e. based on morphological criteria with respect to

fixed time intervals) and settings of contrasts and thresholds in pairwise comparisons can

lead to different results. Still, analysis by qPCR of differential expression corroborated the

results obtained by all computational analysis (DESeq, degReport) and thus the quality of our

dataset in highlighting the gene expression dynamics of *M. galloprovincialis* early

development. The relevance and representativeness of our transcriptome in the context of

bivalves' early development is also supported by the results of the qualitative analysis of

DEGs functional domains. Transcription factors known to function in the regulation of

metazoan development (i.e.; homeobox and fork head genes) and cohort of genes encoding

shell matrix proteins of early shell formation were highly represented among the differentially

expressed genes (Guo et al., 2023; Lartilot et al., 2002; Morino et al., 2013; Paps et al., 2015;

Ramesh et al., 2019; Setiamarga et al., 2013; Zhao et al., 2018; Zheng et al., 2019). In

addition, the analysis uncovered a strong enrichment of receptors, transporters, and channels for hormones and neurotransmitters which is consistent with previous analyses indicating that such elements as pivotal regulators of larval shell biogenesis (i.e.; dopamine), neurogenesis (i.e.; serotonin) and global larval morphogenesis in bivalve mollusks (Guo et al., 2023; Liu et al., 2020; Miglioli, 2022; Miglioli et al., 2021b; Song et al., 2016a). In agreement to that, transmembrane transport and receptor-mediated signal transduction was a highly significant biological process in the *M. galloprovincialis* trochophore and advanced trochophore (Guo et al., 2023; Ramesh et al., 2019; Zhao et al., 2018; Zheng et al., 2019). Altogether, these results demonstrate that the developmental transcriptome of *Mytilus galloprovincialis* is highly representative of the transcriptional (transcription factors) and developmental programs (i.e., shell matrix proteins, G-protein coupled receptors, neurotransmission) governing the early development of bivalve mollusks.

Given the importance of neurotransmitters in the regulation of early bivalve development, in this study we also devised a high resolution, pan-larval, co-labelling protocol for double HCR *in situ* hybridization and immunocytochemistry. Together with the developmental transcriptome, such combination of protocols permits to monitor expression dynamics of multiple genes and localization of polypeptides (i.e., neurotransmitters) at the tissue level throughout the early development of *M. galloprovincialis*. With this protocol we are the first to show the developmental expression and colocalization of *Wnt8a* and *Foxb2,* key genes in animal development, together with the immunoreactivity of serotonin (5HT), an important neurotransmitter that localizes in the apical organ of bivalve larvae (Lartilot et al., 2002; Marlow et al., 2014; Miglioli et al., 2021c; Tomer et al., 2010). The design of such protocol is unprecedented for the early larval stages of bivalve mollusks and constitutes a major step-forward to establish *M. galloprovincialis* as a key model for molluscan development.

3.2. Gene expression analyses suggest the existence of a further intermediate stage between the Trochophore and D-Veliger larva

Hierarchical clustering of libraries was carries out to identify developmental phases based on DEGs expression dynamics as previously described (Leclère et al., 2019; Suzuki and Shimodaira, 2006). The results identified the main developmental phases of bivalve development (i.e., Zygote, Gastrula, Trochophore, D-Veliger) plus an additional phase, the advanced trochophore (Fig. 2A). A similar pattern was also indicated in the principal

component analysis and heatmap of the Euclidean distances which highlighted, respectively, the summit of the 'transcriptional wave' and distinct clustering within the libraries characterized by a trochophore-like morphology (Fig. 1). The existence of an Advanced Trochophore (or Early Veliger) intermediate larval stage intercalated between the Trochophore and D-Veliger larvae has been proposed by several independent sources and our results are the first to demonstrate this developmental phase on a transcriptomic level (Kamenev et al., 2018; Miglioli, 2022; Vogeler et al., 2016). Moreover, given that the morphotypes of the trochophore and of the advanced trochophore are extremely similar, our results suggest that developmental staging of bivalve mollusks by visual observation needs to be carried out with great care to avoid erroneous stage assignments. The possibility of an additional transcriptomic intermediate larval stage linking the trochophore to the competent larva in mollusks and lophotrochozoans could be a significant discovery, offering new perspectives in in future studies on the evolution of the trochophore larval form.

3.3. Identification of transcriptomic transitions during bivalve mollusks development, with potential implications for shell formation and larval morphogenesis

Gene soft clustering revealed transcriptomic signatures overlapping the transitional stages identified by hierarchical clustering of the transcriptome libraries: from gastrula to trochophore (Transition 1), trochophore to advanced trochophore (Transition 2), and advanced trochophore to D-veliger (Transition 3). Taking advantage of our fine-grained developmental sampling, combined with DEG analysis and soft gene clustering, we were able to identify highly representative genes of each transition: 98 genes for Transition 1, 11 for Transition 2 and 8 for Transitions 3. The higher number of Transition 1 genes might be related to the initiation of gene regulatory networks responsible for a wide variety of ontogenetic processes necessary for the establishment of the trochophore larva, including neurogenesis, myogenesis, hematopoiesis, gut formation, and shell secretion (Arenas-Mena, 2013; Dyachuk and Odintsova, 2009; Dyrynda et al., 1997; Kniazkina and Dyachuk, 2022; Miglioli et al., 2019; Miglioli et al., 2021a; Song et al., 2016b; Yurchenko et al., 2018). Differently, trochophore to advanced trochophore and advanced trochophore to D-veliger transitions might require a less dramatic transcriptional shift, both in terms of number of genes or expression (i.e. lower fold change variation). To evaluate the potential conservation of these three transitions within mollusks, we carried out a cross-species comparison with the pacific

oyster *C. gigas* (Liu et al., 2021). This comparison revealed that homologous genes clearly
characterize Transition 1 (gastrula to trochophore) and 3 (advanced trochophore to D-Veliger)
but not Transition 2 (trochophore to advanced trochophore) in the *C. gigas* available
transcriptome. It might be that inconsistent sampling designs of the two datasets blurred the
transcriptional signature of this transition in *C. gigas*. Alternatively, the overlap identified for
Transition 1 and Transition 3 in contrast to Transition 2 might indicate a relatively smaller
number of shared homologous transcriptomic traits during the trochophore development of
bivalve mollusks. In fact, phylostratigraphic analyses from different sources identify the late
trochophore stage of bivalves as the one characterized by the youngest transcriptome which
largely derives from the fast-evolving phylum and class specific phylostrata (Wu et al., 2019;
Xu et al., 2016). In addition, it has been suggested that specific adaptations of the larval body
plan, such as the shell field of bivalves, originate from developmental programs initiated
during the trochophore stage that rely on lineage and species-specific gene sets (Kemkemer
and Long, 2014; Salamanca-Díaz et al., 2022; Xu et al., 2016).

To infer which tissues and, potentially, biological processes could be co-opted in the
transitions of the two bivalves, we studied the expression pattern of a subset of 'transition
genes' by Hybridization chain reaction in *M. galloprovincialis* embryos and larvae. For
Transition 1, we characterized the expression of 8 genes conserved between *M.*
galloprovincialis and *C. gigas* (*Erg, Tbx2b, Dmbt1, Pxdn, SLC4A4, TYRO2, NA03,* and *HPP*)
and found evidence that they might be involved in different developmental processes defining
the transition from embryo to larva: ectomesoderm formation, dorsal specification, and shell
field ontogeny. *Erg* and *Tbx2b* are respectively described as pivotal regulators of
ectomesoderm formation and dorsal specification in other lophotrochozoans and the tissue
localization in *M. galloprovincialis* early larvae is suggestive for a similar developmental role
(Arenas-Mena, 2013; Osborne et al., 2018). With respect to shell field ontogeny, *Pxdn* and
TYRO2 are well known regulators of molluscan shell biogenesis, *SLC4A4* is an important
player in shell calcification while, by homology, *Dmbt1* might be involved in the terminal
differentiation of the shell field columnar cells (Miglioli et al., 2019; Nayshool et al., 2022;
Ramesh et al., 2019; Takito and Al-Awqati, 2004). The remaining two conserved genes of
Transition 1 (*NA03* and *HPP*) could not be annotated, although similar sequences exist in
other mollusks. However, the expression of both genes is suggestive of a potential role in
larval shell biogenesis.

We selected 2 genes conserved between *M. galloprovincialis* and *C. gigas* at Transition 2 (from 28 hpf to 32 hpf), which marks the completion of a fully calcified larval shell (Miglioli et al., 2019). Coherently, we identified *SLC26A5*, a member of the second family of bicarbonate transporters that provides substrate for calcification (Ramesh et al., 2019). The second gene, *Grin3a*, encodes the NMDA-type receptor subunit NR3. NMDA subunits NR1 and NR2 have recently been characterized in nerves and ganglions of *C. gigas* but their function in early trochophore development of bivalves is yet to be documented (Vogeler et al., 2021).

Transition 3 marks the internalization of the velum into the D-shaped shell which relies on a developed muscle system (Dyachuk and Odintsova, 2009). Interestingly, the first gene, *EF-hand*, has been linked to myosin regulation in muscle fibers in scallops and the expression in *M. galloprovincialis* D-Veligers shows similarities to the ventral distribution of paramyosin immunoreactivity in the D-veliger larva of the mussel *Mytilus trossulus* (Dyachuk and Odintsova, 2009; Fromherz and Szent-Györgyi, 1995). The second gene, *COLVI*, is expressed in mantle cells on both sides of the D-border of the larval shell which is coherent with its function in nacre formation described for the pearl oyster *Pinctada martensii* (Zheng et al., 2017).

Altogether, the analysis of the co-opted 'transition genes' corroborates for bivalve mollusks the potential conserved function and tissue distribution of well-known regulators of the gastrula to trochophore transition of lophotrochozoans. What is more, it also revealed several genes potentially involved in shell field ontogeny, muscle function, shell secretion and calcification that are likely common in the developmental transitions of bivalve mollusks. Finally, the transcriptomic staging approach we used for the characterization of *M. galloprovincialis* development could be applied to other bivalve, mollusk, and lophotrochozoan species and in cross-species comparisons for (1) the definition of the developmental stages at a transcriptomic level and (2) the identification of major developmental transitions in transcriptome datasets. Altogether, this approach enables comparisons of homologous gene sets characteristic of a given developmental phase or transition between molluscan, and potentially lophotrochozoan, species.

3.4. The Mediterranean mussel, *Mytilus galloprovincialis*, a novel model for developmental studies of mollusks and evolution of the Trochophore larva

The results reported in this study represent the first thorough transcriptome analysis of the
development of the Mediterranean mussel *M. galloprovincialis*. It further presents a collection
of protocols crucial for the validation and exploitation of transcriptomic data, including
experimental tools, such as qPCR, immunohistochemistry, and *in situ* hybridization. The
combination of fluorescent immunohistochemistry and *in situ* hybridization is particularly
powerful for assessing the co-localization of crucial developmental factors in mussel embryos
and larvae. Potential applications for this transcriptome, and associated experimental
protocols, include developmental studies, cross-species comparisons, and ecotoxicological
surveys. Taking advantage of these approaches, we are the first to document the
developmental expression of a suite of known and unknown genes in developing *M.*
galloprovincialis, revealing the full potential of this animal system for becoming a reference
model for developmental studies on bivalve mollusks and evolution of the trochophore larva.
480

4. Materials and Methods

4.1. Spawning, fertilization, and rearing of embryos and larvae
Sexually mature *M. galloprovincialis* adults were harvested from the natural population in the
bay of Villefranche-sur-Mer, France (43.682°N, 7.319°E) during the spawning seasons of
2021, 2022, and 2023 (January through April). Animals were maintained and acclimated to
laboratory conditions, as previously described (Miglioli et al., 2021b).Thirty adult mussels
were exposed to a heat shock to induce gamete release (Miglioli et al., 2021b; Miglioli et al.,
2021a). Briefly, epibionts, dirt, and algae were scrubbed off the shells under running filtered
natural seawater. Byssus and threads were carefully cut off. Clean mussels were placed on
ice for 10 minutes and then immersed, in individual 200 mL containers and on a rocking
shaker, in filtered natural seawater kept at 28°C until gamete release. Spawning mussels
were immediately moved to filtered natural seawater at 16°C. Prior to fertilization, gametes
were collected, washed, and maintained in filtered natural seawater (using a 0.2 µm Millipore
filter). Fertilization was carried out as previously described with a 1:10 ratio of eggs to sperm
(Miglioli et al., 2021b; Miglioli et al., 2021a). After 30 minutes, the fertilization success rate
(i.e., the ratio of the number of fertilized eggs over the total number eggs x 100) was
established by microscopic observation. Two parental pairs (P1 to P4) with a high fertilization
success rate (>90%) were selected for the following experimental manipulations. 2

experimental replicates were set up, each consisting of a homogeneous pool of fertilized eggs from the 2 parental pairs. Embryos from each replicate were distributed into 200 mL flasks for suspended cultures and brought to a density of 100 embryos per mL with filtered natural seawater. A total of 14 flasks per replicate were prepared, corresponding to the pre-defined sampling strategy: in addition to sampling before fertilization, sample collection every 4 hours from 4 hpf to 52 hpf, with an additional sampling timepoint at 72 hpf. Embryo cultures were maintained at 16°C throughout the experiment. Samples for immunohistochemistry and *in situ* hybridization were produced following the same culture protocol. After filtration, larvae were fixed in 4% paraformaldehyde in 1X PBS overnight at 4°C, washed 3 times for 15 minutes in 1X PBS with 0.01% Tween 20, and then stored in 100% methanol at -20°C (Miglioli et al., 2021b).

4.2. Sampling, total RNA extraction, and high-throughput sequencing

RNA extraction and purification were carried out with the RNAqueous-Micro Kit (Invitrogen) and the RNeasy MinElute Cleanup Kit (Qiagen) following published protocols (Leclère et al., 2019). Unfertilized eggs were directly collected from spawning females, and embryos and larvae were sampled with a 60 μm mesh filter and concentrated in RNAse free 1.5 mL tubes. The tubes were then placed on ice and gently centrifuged multiple times to remove the seawater from the samples. The pellet was then immediately resuspended in lysis buffer, homogenized by energetic pipetting, and flash-frozen in liquid nitrogen. One replicate was lost during extraction procedures. For each sample, 10 μL of the 1.5 mL filtrate were placed in a separate tube and fixed overnight in 4% paraformaldehyde and 1X PBS at 4°C (Miglioli et al., 2021a). The fixed larvae were then stained with Hoechst (Invitrogen) and imaged with a SP8 confocal microscope (Leica Microsystems) to validate proper developmental progression (Miglioli et al., 2021a). Lysed samples were stored at -80□°C until RNA extraction and purification (Leclère et al., 2019). RNA concentration and quality was assessed with a Nanodrop (Thermo Scientific) and the Agilent 2100 Bioanalyzer. Samples were shipped on dry ice to BGI (Hong Kong) for high-throughput sequencing. At least thirty million clean paired-end reads of 200-bp each were produced using the BGISEQ-500 Platform. Reads cleaning and trimming was performed by BGI (Hong Kong). After sequencing, the raw reads were filtered. Data filtering includes removing adaptor sequences, contamination and low-quality reads from raw reads. This step was completed by SOAP nuke software developed by

530   BGI: 1) Filter adapter: Delete the entire read if more than 28% match the adapter sequence;

531   2) Filter low-quality data: Delete the entire read if there are more than 40% bases having a

532   quality value lower than 20; 3) Remove N: Delete the entire read if there are more than 3% N

533   in the read. Libraries were not subjected to further trimming (Dobin et al., 2013).

4.3. Genome-guided transcriptome assembly and differential expression analysis

Sequencing quality of the libraries was assessed with FastQC (v.0.11.7) (Andrews, 2016).

The libraries were mapped on the indexed *M. galloprovincialis* reference genome with STAR

(v.2.7.10a) using default parameters (Dobin et al., 2013; Gerdol et al., 2020). Read counts

were obtained using StringTie (v2.2.1) in counting mode with the genome annotation file

(Pertea et al., 2015; Pertea et al., 2016). The resulting gene count matrix was exported using

prepDE.py and processed in Rstudio (v.4.2.2) with the DESeq2 (v.1.24.0) package using

default parameters and a design file indicating replicate number and developmental time for

each library (Love et al., 2014). The dataset was then pre-filtered by keeping all the genes

with at least 10 reads in at least 2 replicated libraries to reduce the potential batch effect

deriving from the variability in gene presence-absence characterizing the species (Gerdol et

al., 2020) . The resulting matrix was normalized using the regularized logarithms of counts

(DESeq2::rlog) and used to perform principal component analyses and heatmaps (Kolde,

2019). Stage-by-stage differential expression analysis was performed with DESeq2 using the

pairwise comparison function, with the reference stage for comparisons relevelled for each

contrast (Love et al., 2014). Pairwise comparisons were performed on samples of consecutive

stages (e.g.; 0 hpf vs 4hpf; 4hpf vs 8hpf). DEGs were filtered by adjusted p-value

(padj.cutoff<0.05) and log2 fold change (lfc.cutoff>0.58).

4.4. Developmental staging and gene clustering

To discriminate stages of *M. galloprovincialis* development from a transcriptomic perspective,

bootstrapped hierarchical clustering with pvclust (v. 2.2-0) default parameters was performed

on regularized logarithms (rlog) of DEG counts as previously described (Leclère et al., 2019;

Love et al., 2014; Suzuki and Shimodaira, 2006). DEGs (rlog normalized counts) were

clustered with DegReport (v.3.16) using default parameters to identify groups of DEGs with

similar expression dynamics during *M. galloprovincialis* development (Pantano, 2023).

Soft clustering with Mfuzz (v2.56.0) package was performed on the transcripts per million

normalized matrix to identify representative gene clusters of different developmental phases

and potential transcriptomic waves (M. Futschik, 2005). The number of soft clusters was determined by the elbow method to the minimum centroid distance (M. Futschik, 2005). Genes were assigned to the cluster with the highest membership probability.

4.5. Functional annotation, gene ontologies, and enrichment analyses

DEGs were annotated as follows. DEGs predicted proteins were annotated using BLASTp (v. 2.11.0+) against the UniProt /SwissProt database, conserved domains were identified with hmmerscan from HMMER 3.3 against the Pfam database, and transcription factor domains were screened with AnimalTFDB (v2.0) (Finn et al., 2015; Finn et al., 2017; Jones et al., 2014; Mahram and Herbordt, 2015; Sayers et al., 2021; Zhang et al., 2015). Gene ontology enrichment analysis was performed on the list of DEGs per stage and soft clusters. The protein sequence corresponding to genes of interest were extracted from the *M. galloprovincialis* proteome and annotated with a Gene Ontology (GO) identifiers by Interproscan (v. 95.0.) analysis (Finn et al., 2017; Gerdol et al., 2020; Jones et al., 2014). Enrichment analysis of Biological Processes (BP) was conducted with topGO R (v3.17), and significance of the enrichment of each GO was computed with Fisher's exact test (Alexa and Rahnenführer, 2007). Significant lists of GOs were then clustered by k-means using semantic similarity of SimplifyEnrichment (v.1.2.0) to identify potential sets of genes responsible for stage-enriched biological processes (Gu and Hübschmann, 2022).

4.6. Transcriptome validation by real time quantitative polymerase chain reaction

Complementary first strand DNAs (cDNAs) were synthesized from total RNA extracts of both biological replicates at all sample timepoints using the SuperScript VILO Kit (Invitrogen). cDNA synthesis was performed using random primers following the manufacturer's instructions: 10 minutes at 25°C, 60 minutes at 42°C, and 5 minutes at 85°C. To validate the differential gene expression analysis and the cluster analysis, 4 target genes were selected: *Wnt8a* (MGAL_10B085403), *Foxb2* (MGAL_10B093191), *SLC46A3* (MGAL_10B020966), and *SYTL4* (MGAL_10B044489). *EF-α1* was used for data normalization as one of the best-performing reference gene products (Balbi et al., 2016). Primer pairs were designed within the open reading frame of each gene using Primer3 (Koressaar and Rem, 2007). Sequences, efficiency, and amplicon sizes are available in the supplementary material (Supplementary Table 9). qPCR reactions were carried out for each set of primers, in experimental triplicates. The amplification was followed in real time using LightCycler 480 SYBR Green 1 master mix

(Roche) in 45 cycles (initial 5 minutes at 95ºC, followed by 45 cycles of 20 seconds at 95ºC,

20 seconds at 60ºC, and 30 seconds at 72ºC) in a LightCycler 480 (Roche). Quantification

was carried out in parallel for each of the target genes and the reference gene. The

quantification of the expression at a given stage was determined in relation to the expression

of the reference gene using the comparative CT method (Schmittgen and Livak, 2008). To

compare qPCR and transcriptome results, expression of the target genes from both datasets

were normalized with respect to their maximum value. Statistical differences in qPCR results

were assessed with using a Student's t-test (de Winter, 2013).

4.7. *In situ* hybridization using the hybridization chain reaction and co-labeling by
immunohistochemistry

Hybridization probe sets were designed using a user-friendly python interface

(https://github.com/rwnull/insitu_probe_generator) and synthesized by OligoPools (Twist

Bioscience). Amplifiers in red (B2: 546 nm) and far red (B1: 647 nm) were purchased from

Molecular Instruments (https://www.molecularinstruments.com/). All probe sets with relative

amplifier pairings are reported in Supplementary Table 10. The larval samples previously

collected and stored in methanol were re-hydrated with 4 washes of 5 minutes followed by a

10-minute incubation in 1X PBS with 0.01% Triton X-100 at room temperature.

Permeabilization was carried out by a 30-minute incubation in the detergent solution (1%

SDS, 0.005% Tween 20, 50 mM HCl pH 7.5, 1 mM EDTA, pH 8, 0.15 M NaCl) at room

temperature. After 2 washes of 1 minute and 2 washes of 5 minutes in 1X PBS with 0.01%

Triton X-100 at room temperature, the specimens were placed in pre-hybridization for 30

minutes at 37°C with gentle shaking in hybridization buffer (HB) (30% formamide, 5X SSC, 9

mM citric acid, pH 6, 0.01% Tween 20, 2 mg heparin, 1X Denhardt's solution, 10% dextran

sulphate). After addition of the probes (at a final concentration of 4 mM), hybridization was

carried out overnight. Subsequent washes were performed at 37°C: starting with an initial

addition of 1.5 volumes of washing solution (30% formamide, 5X SSC, 9 mM citric acid, pH 6,

0.01% Tween 20, 2 mg Heparin) with respect to the volume of the hybridization solution, 4

washes of 15 minutes were performed with washing solution. Specimens were then washed,

at room temperature, twice for 5 minutes in 5X SSC with 0.01% Tween 20, once for 5 minutes

in 1X SSC with 0.01% Tween 20, and 30 minutes in amplification buffer (5X SSC, 0.01%

Tween 20, 10% dextran sulphate), before amplifiers were added (at a final concentration of

60 mM) to start the amplification chain reaction. Amplifiers were prepared as follows: the

required volume of hairpins 1 and 2 from each amplifier were placed in a 0.6 ml tube and

incubated for 90 seconds at 95°C. After the heat shock, amplifiers were placed in the dark at

room temperature for 30 minutes. The amplifiers were then resuspended in amplification

buffer and distributed to the samples. Amplification was carried out at room temperature in the

dark overnight. Excess hairpins were removed in two washes of 5 minutes and two washes of

30 minutes in 1X SSC with 0.01% Tween 20, followed by two washes of 5 minutes in 1X PBS

with 0.01% Tween 20. Co-labeling by immunocytochemistry with Serotonin (5-HT) anti-rabbit

antibody (1:10'000; #20080 Immunostar, Hudson, WI, USA) and Goat Anti-Rabbit IgG (H+L)

F(ab')$_2$ Alexa Fluor 488 conjugated (1:500; #4412S Cell Signaling Technology, Leiden, The

Netherlands) secondary antibody was performed immediately after HCR amplification using

the previously described conditions (Miglioli et al., 2021a; Miglioli et al., 2021b). Specimens

were then prepared for imaging by incubation in 1 □g/L of Hoechst (Invitrogen) in 1X PBS for

30 minutes at room temperature. Following 3 washes of 5 minutes 1X PBS with 0.01% Tween

20, specimens were mounted with the antifading agent CitiFluor AF1 (Agar Scientific) and

imaged on a SP8 confocal microscope (Leica), as previously described (Miglioli et al., 2021b).

Images were analyzed with ImageJ and fluorescent signals were corrected as follows:

autofluorescence from tissue and shell was subtracted with "math", the signal was

homogenized with the "despeckle" or "smooth" functions, the stacks (z) were merged by

either "maximum intensity" or "sum slices".

4.8. Developmental transition gene selection and comparisons with the developmental
transcriptome of the Pacific oyster *Crassostrea gigas*

Representative genes of the developmental transitions were identified in *M. galloprovincialis*

following this strategy. First, developmental transitions were defined based on the hierarchical

clustering of the stages. Three transitions were identified: from 16 to 20 hpf, 28 to 32 hpf and

40 to 44 hpf. Secondly, genes were clustered according to their expression profile by soft

clustering (section 4.4) allowing the identification of three gene clusters corresponding to

transitions. From this clusters, only genes showing a differential expression at the time of

transition were selected. For transition 1, genes were selected if they showed a differential

expression (log fold change ≥ 2, approximated by excess) between 16hpf and 20hpf. For

transition 2, genes were selected if they showed a differential expression (log fold change ≥ 2,

approximated by excess) between 28hpf and 32hpf. For transition 3, genes were selected if they showed a differential expression (log fold change ≥ 2, approximated by excess) between 40hpf and 44hpf. To identify potential homologous genes, a BLASTp (Mahram and Herbordt, 2015) of the *M. galloprovincialis* 'transition genes' against the proteome of the Pacific oyster *C. gigas* (Liu et al., 2021) was performed. The expression profile of selected genes in *C. gigas* was visualized with a heatmap (Kolde, 2019) to investigate the potential presence of similar developmental transitions. HCR was performed on *M. galloprovincialis* 'transition genes' that had a homolog in *C. gigas* whose z-score value turned to positive (>0) at the time of the transition.

5. Acknowledgments

The authors are indebted to Guy Lhomond and Lucas Leclère for technical assistance and fruitful discussions. The authors would also like to thank Sébastien Schaub of the Plateforme d'Imagerie par Microscopie (PIM) as well as Laurent Gilletta and Axel Duchene of the Service Aquariologie (SA) of the Institut de la Mer de Villefranche (IMEV), France, that is supported by EMBRC-France and funded by the Agence Nationale de la Recherche (ANR) (ANR-10-INBS-02). Thank you to the members of the Ascidian BioCell and EvoInSiDe teams at the Laboratoire de Biologie du Développement de Villefranche-sur-Mer, France, as well as to the members of the Canesi team of Environmental Physiology at the Università Degli Studi Di Genova, Italy, for their continuing support.

6. Competing Interests

The authors declare that they have no competing financial interests or personal relationships that could have influenced the work reported in this paper.

7. Funding

This work was funded by the CNRS and the ANR (ANR-21-CE34-0006-02), both of which are French granting agencies, to Michael Schubert and Rémi Dumollard.

8. Data Availability

All libraries analyzed in this article have been deposited in the NCBI SRA database (accession: PRJNA996031 and ID: 996031).

686
9. References

Alexa, A. and Rahnenführer, J. (2007). Gene set enrichment analysis with topGO. *Bioconductor*
Improv. 27.

Andrews, S. (2016). Data, FASTQC. A quality control tool for high throughput sequence.

Arenas-Mena, C. (2013). Brachyury, Tbx2/3 and sall expression during embryogenesis of the
indirectly developing polychaete Hydroides elegans. *Int. J. Dev. Biol.* **57**, 73–83.

ASTM E724-21, A. S. for T. and M. (2021). ASTM E724-21. Standard Guide for Conducting Static
Short-Term Chronic Toxicity Tests Starting with Embryos of Four Species of Saltwater Bivalve
Molluscs.

Balbi, T., Franzellitti, S., Fabbri, R., Montagna, M., Fabbri, E. and Canesi, L. (2016). Impact of
bisphenol A (BPA) on early embryo development in the marine mussel Mytilus galloprovincialis:
Effects on gene transcription. *Environ. Pollut.* **218**, 996–1004.

Brzozowska, R., Sui, Z. and Kang, K. H. (2012). Testing the usability of sea mussel (Mytilus sp.) for
the improvement of seawater quality-An experimental study. *Ecol. Eng.* **39**, 133–137.

Choi, H. M. T., Beck, V. A. and Pierce, N. A. (2014). Next-generation in situ hybridization chain
reaction: Higher gain, lower cost, greater durability. *ACS Nano* **8**, 4284–4294.

Daguin Claire, B. P. (2000). Genetic relationships of Mytilus galloprovincialis Lamarck populations
worldwide: evidence from nuclear-DNA markers. *Geol. Soc. London, Spec. Publ.*

De Robertis, E. M. and Tejeda-Muñoz, N. (2022). Evo-Devo of Urbilateria and its larval forms. *Dev.*
Biol. **487**, 10–20.

de Winter, J. C. F. (2013). Using the student's t-test with extremely small sample sizes. *Pract.*
Assessment, Res. Eval. **18**, 1–12.

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M.
and Gingeras, T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–
21.

Dyachuk, V. and Odintsova, N. (2009). Development of the larval muscle system in the mussel
Mytilus trossulus (Mollusca, Bivalvia): Original Article. *Dev. Growth Differ.* **51**, 69–79.

Dyrynda, E. A., Pipe, R. K. and Ratcliffe, N. A. (1997). Sub-populations of haemocytes in the adult
and developing marine mussel, Mytilus edulis, identified by use of monoclonal antibodies. *Cell*
Tissue Res. **289**, 527–536.

FAO (2022). *The State of World Fisheries and Aquaculture 2022. Towards Blue Transformation.*

Finn, R. D., Clements, J., Arndt, W., Miller, B. L., Wheeler, T. J., Schreiber, F., Bateman, A. and
Eddy, S. R. (2015). HMMER web server: 2015 Update. *Nucleic Acids Res.* **43**, W30–W38.

Finn, R. D., Attwood, T. K., Babbitt, P. C., Bateman, A., Bork, P., Bridge, A. J., Chang, H. Y.,
Dosztanyi, Z., El-Gebali, S., Fraser, M., et al. (2017). InterPro in 2017-beyond protein family

and domain annotations. *Nucleic Acids Res.* **45**, D190–D199.

Foulon, V., Boudry, P., Artigaud, S., Guérard, F. and Hellio, C. (2019). In silico analysis of pacific
          oyster (Crassostrea gigas) transcriptome over developmental stages reveals candidate genes for
          larval settlement. *Int. J. Mol. Sci.* **20**,.

Fromherz, S. and Szent-Györgyi, A. G. (1995). Role of essential light chain EF hand domains in
          calcium binding and regulation of scallop myosin. *Proc. Natl. Acad. Sci. U. S. A.* **92**, 7652–7656.

**Gerdol, M., Moreira, R., Cruz, F., Gómez-Garrido, J., Vlasova, A., Rosani, U., Venier, P., Naranjo-
          Ortiz, M. A., Murgarella, M., Greco, S., et al.** (2020). Massive gene presence-absence variation
          shapes an open pan-genome in the Mediterranean mussel. *Genome Biol.* **21**, 275.

Gu, Z. and Hübschmann, D. (2022). Simplify enrichment: A bioconductor package for clustering and
          visualizing functional enrichment results. *Genomics. Proteomics Bioinformatics*.

Guo, X., Li, X., Zhao, F., Liu, D., Yang, Z., Li, M., Li, Y., Wei, H., Wang, H., Qin, Z., et al. (2023).
          Full-length transcriptome analysis provides insights into larval shell formation in Mulinia lateralis.
          Front. Mar. Sci. **9**, 1–14.

Hunt, M. K. and Maslakova, S. A. (2017). Development of a lecithotrophic pilidium larva illustrates
          convergent evolution of trochophore-like morphology. *Front. Zool.* **14**, 1–18.

ISO 17244:2015, I. O. for S. (2020). ISO 17244:2015. Water quality — Determination of the toxicity of
          water samples on the embryo-larval development of Japanese oyster (Crassostrea gigas) and
          mussel (Mytilus edulis or Mytilus galloprovincialis).

**Jones, P., Binns, D., Chang, H. Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J.,
          Mitchell, A., Nuka, G., et al.** (2014). InterProScan 5: Genome-scale protein function
          classification. *Bioinformatics* **30**, 1236–1240.

Kamenev, Y. O., Eliseikina, M. G., Maslennikov, S. I. and Dolmatov, I. Y. (2018). The
          ultrastructural features of embryonic and early larval development in Yesso scallop,
          Mizuhopecten yessoensis. *Tissue Cell* **53**, 76–86.

Kemkemer, C. and Long, M. (2014). New genes important for development. *EMBO Rep.* **15**, 460–
          461.

Kniazkina, M. and Dyachuk, V. (2022). Neurogenesis of the scallop Azumapecten farreri: from the
          first larval sensory neurons to the definitive nervous system of juveniles. *Front. Zool.* **19**, 1–19.

Kniprath, E. (1980a). Roux's Archives of Developmental Biology. **204**, 201–204.

Kniprath, E. (1980b). Larval development of the shell and the shell gland in Mytilus (Bivalvia).
          Wilhelm Roux's Arch. Dev. Biol. **188**, 201–204.

KNIPRATH, E. (1981). Ontogeny of the Molluscan Shell Field: a Review. *Zool. Scr.* **10**, 61–79.

Kocot, K. M., Aguilera, F., McDougall, C., Jackson, D. J. and Degnan, B. M. (2016). Sea shell
          diversity and rapidly evolving secretomes: Insights into the evolution of biomineralization. *Front.
          Zool.* **13**, 1–10.

Kolde, R. (2019). pheatmap: Pretty Heatmaps. R package version 1.0.12.

Kurita, A., Kurita, Y., Deguchi, R. and Wada, H. Early Development and Cleavage Pattern of the
Japanese Purple Mussel , Septifer virgatus Early Development and Cleavage Pattern of the
Japanese Purple Mussel , Septifer virgatus. **26**, 814–820.

Kurita, Y., Deguchi, R. and Wada, H. (2009). Early development and cleavage pattern of the
Japanese purple mussel, septifer virgatus. *Zoolog. Sci.* **26**, 814–820.

Lartilot, N., Le Gouar, M. and Adoutte, A. (2002). Expression pattern of fork head and goosecoid
homologues in the mollusc Patella vulgata supports the ancestry of the anterior mesendoderm
across Bilateria. *Dev. Genes Evol.* **212**, 551–561.

Leclère, L., Horin, C., Chevalier, S., Lapébie, P., Dru, P., Peron, S., Jager, M., Condamine, T.,
Pottin, K., Romano, S., et al. (2019). The genome of the jellyfish Clytia hemisphaerica and the
evolution of the cnidarian life-cycle. *Nat. Ecol. Evol.* **3**, 801–810.

Liu, Z., Zhou, Z., Zhang, Y., Wang, L., Song, X., Wang, W., Zheng, Y., Zong, Y., Lv, Z. and Song,
L. (2020). Ocean acidification inhibits initial shell formation of oyster larvae by suppressing the
biosynthesis of serotonin and dopamine. *Sci. Total Environ.* **735**, 139469.

Liu, F., Li, Y., Yu, H., Zhang, L., Hu, J., Bao, Z. and Wang, S. (2021). MolluscDB: An integrated
functional and evolutionary genomics database for the hyper-diverse animal phylum Mollusca.
Nucleic Acids Res. **49**, D988–D997.

Love, M. I., Huber, W. and Anders, S. (2014). Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol.* **15**, 1–21.

M. Futschik, B. C. (2005). Noise robust clustering of gene expression time-course data. 965–988.

Mahram, A. and Herbordt, M. C. (2015). NCBI BLASTP on high-performance reconfigurable
computing systems. *ACM Trans. Reconfigurable Technol. Syst.* **7**,.

Marlow, H., Tosches, M. A., Tomer, R., Steinmetz, P. R., Lauri, A., Larsson, T. and Arendt, D.
(2014). Larval body patterning and apical organs are conserved in animal evolution. *BMC Biol.*
12, 1–17.

Martín, A., Baltodano, C., Seudre, O., Guynes, K., María, J. and Durán, M. (2021). Early
embryogenesis and organogenesis in the annelid Owenia fusiformis. *Evodevo* 1–18.

Maureen Liu, M., Davey, J. W., Jackson, D. J., Blaxter, M. L. and Davison, A. (2014). A conserved
set of maternal genes? Insights from a molluscan transcriptome. *Int. J. Dev. Biol.* **58**, 501–511.

Miglioli, A. (2022). Pathways of endocrine disruption in the larval development of the mediterranean
mussel Mytilus galloprovincialis To cite this version□: HAL Id□: tel-03696706 PhD thesis of the
Università degli studi di Genova and Sorbonne Université Presented by PhD candi. 203.

Miglioli, A., Dumollard, R., Balbi, T., Besnardeau, L. and Canesi, L. (2019). Characterization of the
main steps in first shell formation in Mytilus galloprovincialis: Possible role of tyrosinase. *Proc. R.*
Soc. B Biol. Sci. **286**,.

Miglioli, A., Balbi, T., Besnardeau, L., Dumollard, R. and Canesi, L. (2021a). Bisphenol A

interferes with first shell formation and development of the serotoninergic system in early larval
stages of Mytilus galloprovincialis. *Sci. Total Environ.* **758**, 144003.

Miglioli, A., Balbi, T., Montagna, M., Dumollard, R. and Canesi, L. (2021b). Tetrabromobisphenol A
acts a neurodevelopmental disruptor in early larval stages of Mytilus galloprovincialis. *Sci. Total*
Environ. **793**, 148596.

Miglioli, A., Balbi, T., Besnardeau, L., Dumollard, R. and Canesi, L. (2021c). Bisphenol A interferes
with fi rst shell formation and development of the serotoninergic system in early larval stages of
Mytilus galloprovincialis. *Sci. Total Environ.* **758**, 144003.

Moreira, R., Pereiro, P., Balseiro, P., Milan, M., Pauletto, M., Bargelloni, L., Novoa, B. and
Figueras, A. (2018). Revealing Mytilus galloprovincialis transcriptomic profiles during ontogeny.
Dev. Comp. Immunol. **84**, 292–306.

Morino, Y., Okada, K., Niikura, M., Honda, M., Satoh, N., & Wada, H. (2013). Genome-Wide Survey
of Genes Encoding Transcription Factors in the Japanese Pearl Oyster, Pinctada fucata: I.
Homeobox Genes. *Zoolog. Sci.* **30**, 851–857.

Nayshool, O., Kol, N., Javaski, E., Amariglio, N. and Rechavi, G. (2022). SurviveAI: Long Term
Survival Prediction of Cancer Patients Based on Somatic RNA-Seq Expression. *Cancer Inform.*
21,.

Nielsen, C. (2005). Trochophora larvae: Cell-lineages, ciliary bands and body regions. 2. Other
groups and general discussion. *J. Exp. Zool. Part B Mol. Dev. Evol.* **304**, 401–447.

Nielsen, C. (2009). How did indirect development with planktotrophic larvae evolve? *Biol. Bull.* **216**,
203–215.

Niu, D., Wang, F., Xie, S., Sun, F., Wang, Z., Peng, M. and Li, J. (2016). Developmental
Transcriptome Analysis and Identification of Genes Involved in Larval Metamorphosis of the
Razor Clam, Sinonovacula constricta. *Mar. Biotechnol.* **18**, 168–175.

Núñez-Acuña, G., Fernandez, C., Sanhueza-Guevara, S. and Gallardo-Escárate, C. (2022).
Transcriptome profiling of the early developmental stages in the giant mussel Choromytilus
chorus exposed to delousing drugs. *Mar. Genomics* **65**,.

Osborne, C. C., Perry, K. J., Shankland, M. and Henry, J. Q. (2018). Ectomesoderm and epithelial–
mesenchymal transition-related genes in spiralian development. *Dev. Dyn.* **247**, 1097–1120.

Page, L. R. (2009). Molluscan larvae: Pelagic juveniles or slowly metamorphosing larvae? *Biol. Bull.*
216, 216–225.

Pantano, L. (2023). DEGreport: Report of DEG analysis. R package version.

Paps, J., Xu, F., Zhang, G. and Holland, P. W. H. (2015). Reinforcing the egg-timer: Recruitment of
novel Lophotrochozoa homeobox genes to early and late development in the Pacific oyster.
Genome Biol. Evol. **7**, 677–688.

Pertea, M., Pertea, G. M., Antonescu, C. M., Chang, T. C., Mendell, J. T. and Salzberg, S. L.
(2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat.*
Biotechnol. **33**, 290–295.

Pertea, M., Kim, D., Pertea, G. M., Leek, J. T. and Salzberg, S. L. (2016). Transcript-level
expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. *Nat. Protoc.*
11, 1650–1667.

Ramesh, K., Yarra, T., Clark, M. S., John, U. and Melzner, F. (2019). Expression of calcification-
related ion transporters during blue mussel larval development. *Ecol. Evol.* **9**, 7157–7172.

Rawlinson, K. A. (2010). Embryonic and post-embryonic development of the polyclad flatworm
Maritigrella crozieri; implications for the evolution of spiralian life history traits. *Front. Zool.* **7**, 1–
25.

Robertis, E. M. De and Tejeda-mu, N. (2022). Evo-Devo of Urbilateria and its larval forms ~ oz. **487**,
10–20.

Salamanca-Díaz, D. A., Ritschard, E. A., Schmidbaur, H. and Wanninger, A. (2022). Comparative
Single-Cell Transcriptomics Reveals Novel Genes Involved in Bivalve Embryonic Shell Formation
and Questions Ontogenetic Homology of Molluscan Shell Types. *Front. Cell Dev. Biol.* **10**, 1–10.

Sayers, E. W., Beck, J., Bolton, E. E., Bourexis, D., Brister, J. R., Canese, K., Comeau, D. C.,
Funk, K., Kim, S., Klimke, W., et al. (2021). Database resources of the National Center for
Biotechnology Information. *Nucleic Acids Res.* **49**, D10–D17.

Schmittgen, T. D. and Livak, K. J. (2008). Analyzing real-time PCR data by the comparative CT
method. *Nat. Protoc.* **3**, 1101–1108.

Setiamarga, D. H. E., Shimizu, K., Kuroda, J., Inamura, K., Sato, K., Isowa, Y., Ishikawa, M.,
Maeda, R., Nakano, T., Yamakawa, T., Hatori, R., Ishio, A., Kaneko, K., Matsumoto, K.,
Sarashina, I., Teruya, S., Zhao, R., Satoh, N., Sasaki, T., … Endo, K. (2013). An In-silico
Genomic Survey to Annotate Genes Coding for Early Development-Relevant Signaling Molecules
in the Pearl Oyster, Pinctada fucata. *Zoolog. Sci.* **30**, 877–888.

Song, H., Yu, Z. L., Sun, L. N., Gao, Y., Zhang, T. and Wang, H. Y. (2016a). De novo transcriptome
sequencing and analysis of Rapana venosa from six different developmental stages using Hi-seq
2500. *Comp. Biochem. Physiol. - Part D Genomics Proteomics* **17**, 48–57.

Song, X., Wang, H., Chen, H., Sun, M., Liang, Z., Wang, L. and Song, L. (2016b). Conserved
hemopoietic transcription factor Cg-SCL delineates hematopoiesis of Pacific oyster Crassostrea
gigas. *Fish Shellfish Immunol.* **51**, 180–188.

Suzuki, R. and Shimodaira, H. (2006). Pvclust: An R package for assessing the uncertainty in
hierarchical clustering. *Bioinformatics* **22**, 1540–1542.

Takito, J. and Al-Awqati, Q. (2004). Conversion of ES cells to columnar epithelia by hensin and to
squamous epithelia by laminin. *J. Cell Biol.* **166**, 1093–1102.

Tomer, R., Denes, A. S., Tessmar-Raible, K. and Arendt, D. (2010). Profiling by Image Registration
Reveals Common Origin of Annelid Mushroom Bodies and Vertebrate Pallium. *Cell* **142**, 800–
809.

Tyler J. Carrier, Adam M. Reitzel, A. H. (2018). *Evolutionary Ecology of Marine Invertebrate Larvae.*
Oxford University Press.

Vogeler, S., Bean, T. P., Lyons, B. P. and Galloway, T. S. (2016). Dynamics of nuclear receptor
gene expression during Pacific oyster development. *BMC Dev. Biol.* **16**, 1–13.

Vogeler, S., Carboni, S., Li, X., Ireland, J. H., Miller-Ezzy, P. and Joyce, A. (2021). Cloning and
characterisation of NMDA receptors in the Pacific oyster, Crassostrea gigas (Thunberg, 1793) in
relation to metamorphosis and catecholamine synthesis. *Dev. Biol.* **469**, 144–159.

Wada, H., Phuangphong, S., Hashimoto, N. and Nagai, K. (2020). Establishment of the novel
bivalve body plan through modification of early developmental events in mollusks. *Evol. Dev.* **22**,
463–470.

Wang, J., Zhang, L., Lian, S., Qin, Z., Zhu, X., Dai, X., Huang, Z., Ke, C., Zhou, Z., Wei, J., et al.
(2020). Evolutionary transcriptomics of metazoan biphasic life cycle supports a single
intercalation origin of metazoan larvae. *Nat. Ecol. Evol.* **4**, 725–736.

Wu, L., Ferger, K. E. and David Lambert, J. (2019). Gene expression does not support the
developmental hourglass model in three animals with Spiralian development. *Mol. Biol. Evol.* **36**,
1373–1383.

Xu, F., Domazet-Lošo, T., Fan, D., Dunwell, T. L., Li, L., Fang, X. and Zhang, G. (2016). High
expression of new genes in trochophore enlightening the ontogeny and evolution of trochozoans.
Sci. Rep. **6**, 1–10.

Yang, Z., Zhang, L., Hu, J., Wang, J., Bao, Z. and Wang, S. (2020). The evo-devo of molluscs:
Insights from a genomic perspective. *Evol. Dev.* **22**, 409–424.

Yurchenko, O. V., Skiteva, O. I., Voronezhskaya, E. E. and Dyachuk, V. A. (2018). Nervous system
development in the Pacific oyster, Crassostrea gigas (Mollusca: Bivalvia). *Front. Zool.* **15**, 1–21.

Zhang, H. M., Liu, T., Liu, C. J., Song, S., Zhang, X., Liu, W., Jia, H., Xue, Y. and Guo, A. Y.
(2015). AnimalTFDB 2.0: A resource for expression, prediction and functional study of animal
transcription factors. *Nucleic Acids Res.* **43**, D76–D81.

Zhao, R., Takeuchi, T., Luo, Y. J., Ishikawa, A., Kobayashi, T., Koyanagi, R., Villar-Briones, A.,
Yamada, L., Sawada, H., Iwanaga, S., et al. (2018). Dual gene repertoires for larval and adult
shells reveal molecules essential for molluscan shell formation. *Mol. Biol. Evol.* **35**, 2751–2761.

Zheng, Z., Du, X., Xiong, X., Jiao, Y., Deng, Y., Wang, Q. and Huang, R. (2017). PmRunt regulated
by Pm-miR-183 participates in nacre formation possibly through promoting the expression of
collagen VI-like and Nacrein in pearl oyster Pinctada martensii. *PLoS One* **12**, 1–14.

Zheng, Z., Hao, R., Xiong, X., Jiao, Y., Deng, Y. and Du, X. (2019). Developmental characteristics of
pearl oyster Pinctada fucata martensii: Insight into key molecular events related to shell
formation, settlement and metamorphosis. *BMC Genomics* **20**, 1–11.

10. Figure Captions

Figure 1: **Morphological and transcriptomic staging of *Mytilus galloprovincialis* early**

development. A) Development of *M. galloprovincialis* under standard culture conditions.

Orientation is as shown by arrows in the top right of each picture, hours post fertilization (hpf)

(at 16°C) is indicated on the top left. Red and black arrowheads highlight
stomodeum/presumptive mouth and anus, respectively, blue dotted line indicates the shell
field outline (16 hpf to 24 hpf) and the hinge region (28 hpf). At 40 hpf, the green line indicates
the D margin of the calcified shell. Abbreviations: An: animal; Ve: vegetal; A: anterior; P:
posterior; D: dorsal; V: ventral; a: anus; at: apical tuft; h: hinge; m: mantle; sf: shell field; st:
stomodeum/presumptive mouth; ve: velum. Scalebar: 20 µm. B) Dot plot of the number of
genes expressed in each library before DESeq2 normalization. The black dotted line
represents the mean trend throughout development. C) Heatmap of the Euclidean distance
matrix of the libraries clustered with default hclust parameters. D) Principal component
analysis (PCA) of DESeq2 normalized libraries, dotted line indicates the axis of Principal
Component 2 within the graph area.

Figure 2: **Differential expression analysis highlights stage-specific transcriptomic**
signatures. A) Hierarchical clustering of differentially expressed gene expression values
across all libraries. Bootstrap values are shown in red on each branch. Names for each
cluster are assigned according to morphology and developmental timepoint. B) Histogram
showing the number of up- (red) and down- (blue) regulated genes at each assayed stage.
Color code is according to log fold change values. Insert: percentage (%) of differentially
expressed gene up-, down- or up-/down-regulated, once (1X) or multiple times. C) Top six
gene expression profiles by number of genes. D) Pie graph depicting the percentage (%) of
profiles reaching their highest expression levels at different developmental stages. E)
Normalized expression of *Wnt8a*, *Foxb2*, *SLC46A3* and *SYTL4* from the transcriptomic (left)
and qPCR (right) data. F) Maximum projection superimpositions of *Foxb2* (yellow) and *Wnt8a*
(red) fluorescent *in situ* hybridization with serotonin (5-HT) immunohistochemistry (green) with
Hoechst (grey) or bright field in trochophore (24 hpf) and D-veliger (48 hpf) larvae.
Abbreviations: an: anus; ao/gc: apical organ/cerebral ganglion; at: apical tuft; aso: apical
sensory organ; eso: esophagus; h: hinge; sf: shell field; st: stomodeum/presumptive mouth;
ve: velum. Scale bar: 20 µm.

Figure 3: **Qualitative analysis of functional domains and gene ontologies (GOs) of**
differentially expressed genes (DEGs). A) Transcription factors. B) Domains associated to
shell biogenesis: shell matrix proteins in blue and calcification proteins in green. C) Receptors
and neurotransmitter transporters. D) Composite image showing the heatmap of significant

GO terms paired with GO cluster analysis at different stages of development. Colors indicate
the -$\log_{10}$(p-value). Column breaks in the heatmap divide the different developmental stages
identified by hierarchical clustering, and row breaks separate the GO clusters. Row color
annotation (on the left) highlights different GO clusters, and the most significant terms for
each cluster are shown in their respective box (on the right).

Figure 4: **Soft clustering reveals transcriptomic signatures of early developmental**
transitions. A) Left panel: expression patterns of single soft clusters. The green lines depict
the expression pattern in scaled transcripts per million (z-scores) of each gene belonging to a
soft cluster, the red line indicates the mean trend of the expression data, vertical grey lines
indicate developmental transitions. Right panel: heatmap of all soft clusters in scaled
transcripts per million (z-scores) arranged by developmental time of appearance. White lines
identify developmental transitions identified by hierarchical clustering (Fig. 2A). B) Expression
of developmental 'transition genes' in scaled transcripts per million (z-scores) in the *Mytilus*
galloprovincialis transcriptome. C) Expression of developmental 'transitions genes' in scaled
transcripts per million (z-scores) in the *Crassostrea gigas* developmental transcriptome.
Column breaks indicate the three transitions. Color annotations on the left, in B) and C), refer
to developmental transitions: blue, gold, and red for Transition 1, 2, and 3, respectively.

Figure 5: **Developmental expression of homologous transition genes in** *Mytilus*
galloprovincialis **larvae.** A) Genes selected for Transition 1 (gastrula to trochophore). B)
Genes selected for Transition 2 (trochophore to advanced trochophore). C) Genes selected
for Transition 3 (advanced trochophore to D-veliger). Left and right valves are presented
parallel to each other with respect to the hinge region. The *in situ* hybridization signal is
shown in red, Hoechst-stained nuclei are shown in grey. Gene names are indicated in red in
the top left corner. Developmental timing in hours post fertilization (hpf) (at 16°C) is indicated
in white under the gene name. Abbreviations: D: dorsal view; S: sagittal view; V: ventral view;
a: anus; at: apical tuft; eso: esophagus; h: hinge; sf: shell field; st: stomodeum/presumptive
mouth. Scale bar: 20 µm.

11. Supplementary Figure Captions

Supplementary Figure 1 – **Fluorescent *in situ* hybridization protocol validation with *Foxb2* and *Wnt8a* at different stages of *Mytilus galloprovincialis* development.** A) Expression pattern of *Foxb2*. Upper panel shows maximum z-projections of the *in situ* hybridization signal (red) with Hoechst nuclear staining (grey). Lower panel shows the *in situ* hybridization signal (red) in brightfield. B) Expression pattern of *Wnt8a*. Upper panel shows maximum z-projections of the *in situ* hybridization signal (red) with Hoechst nuclear staining (grey). Lower panel shows the *in situ* hybridization signal (red) in brightfield. Abbreviations: at: apical tuft; h: hinge; sf: shell field; st: stomodeum/presumptive mouth; ve: velum. Scale bar: 20µm.

Supplementary Figure 2 – **Patterns of developmental expression of transition genes in a developmental transcriptome of the Pacific oyster *Crassostrea gigas*.** A) Transition1. B) Transition 2. C) Transition 3.

12. Supplementary Files and Tables

Supplementary File 1 – **DESeq2 output of differentially expressed genes by pairwise comparison.** Every tab corresponds to a pairwise comparison.

Supplementary File 2 – *Supplementary Table 1:* Summary of mapping results for each library. *Supplementary Table 2:* Matrix of raw gene counts**.** *Supplementary Table 3***:** Dynamics of differentially expressed genes with Uniprot protein annotation, Pfam domain annotation, DEG cluster annotation, developmental transition membership, and soft clustering membership**.** *Supplementary Table 4***:** DegReport analysis. The mean expression value of each cluster is reported for each timepoint, and the expression trend is presented as a sparkline histogram. The green bar indicates the highest and the red one the lowest value. *Supplementary Table 5***:** Binary matrix of the DegReport output. *Supplementary Table 6***:** List of differentially expressed transcription factors identified with AnimalTFDB**.** *Supplementary Table 7:* Binary matrix of the Mfuzz soft clustering analysis**.** *Supplementary Table 8***:** Best blastp hits of *Mytilus galloprovincialis* 'transition genes' in the *Crassostrea gigas* genome**.** *Supplementary Table 9:* Primer pairs used for qPCR analyses. *Supplementary Table 10*: Probe sets for hybridization chain reaction used in the study.

A

Number of DEGs

zf-C2H2	127
Homeodomain	113
BTB	64
zf-H2C2_2	58
HLH	56
TF_bZIP	31
THAP	30
Forkhead	27
HMG_box	16
Ets	16
TRAM_LAG1_CLN8	14
zf-LITAF-like	10
Myb_DNA-binding	10
HTH_psq	9
Thyroid_Hormon.	9
T-box	8
SNF2-rel_dom	6
PAX	6
RFX_DNA_binding	6
Retinoid_X_Rece.	6
ARID	5
GATA	4
RHD_DNA_bind	4
E2F_TDP	4
zf-C4	4
CUT	3
NFYC	3
SRF-TF	3
Pou	3
STAT_bind	3
MH1	3
Estrogen_Recept.	3
MBD	2
zf-MIZ	2
Nrf1_DNA-bind	2
zf-BED	2
Tub	2
GTF2I	2
CP2	2
NFYB	2
CSD	2
IRF	2
TSC22	2
CBFB_NFYA	2
TFr_COE	2
zf-CCCH	1
HPD	1
AF-4	1
TIP_N	1
CBF_beta	1
CG-1	1
HSF_DNA-bind	1
DM	1
Runt	1
P53	1
TEA	1
SAND	1
zf-C2HC	1
TF_AP-2	1
GCFC	1
BTD	1
DACH	1
Miscellaneous	1
NGF_IB-like	1
Steroidogenic_Fa.	1
MOZ_SAS	1

B

Number of DEGs

EF-hand	192
WKA	89
Fibronectin III	88
Lectin Type C	73
CB	49
Concanavalin_A	31
Calponin	31
Kazal	21
Beta lactamase	20
Haem Peroxidase	19
Filamin	13
Tyrosinase	11
TGF Beta	10
Chitin Synthase	9
CB & WKA	7
Kunitz	7
Chitinase	6
Cyclophilin	5
Peptidase C1A	4
Thymosin beta	3
Peroxiredoxin	2
Amine Oxidase	2
Ion Chan/Trans	52
SLCs	19
Carbinic Anhydrase	15
RUNX	1

C

Number of DEGs

7TM: Rhodopsin	194
NTs ion channels	95
Ephrin A	92
LD lipoprotein	59
7TM: Secretin	31
NTs symporters (Na+)	30
Nuclear Receptors	23
Scavenger	21
TGF beta II	2

D

A

Egg

Zygote

Blastula

Gastrula

Early Trochophore

Advanced Trochophore

Veliger

D-Veliger

Late D-Veliger

00 04 | 08 12 16 | 20 24 28 | 32 36 40 | 44 48 52 72
Zygote | Gastrula | Trochophore | Advanced Trochophore | D-Veliger

Hours post fertilization
& Hierarchical clusters

B *Mytilus galloprovincialis*

00 04 | 08 12 16 | 20 24 28 | 32 36 40 | 44 48 52 72
Zygote | Gastrula | Trochophore | Advanced Trochophore | D-Veliger

Hours post fertilization
& Hierarchical clusters

C *Crassostrea gigas*

Egg | Morula | Blastula | Gastrula | Trochophore 1 | Trochophore 2 | Trochophore 3 | Trochophore 4 | Trochophore 5 | EarlyD_1 | EarlyD_2 | DV1 | DV2 | DV3 | DV4

Libraries