Differences in sperm swimming speed and morphology between the three genetic morphs in the ruff sandpiper

Martin Bulla[1,2]✉, Clemens Küpper[1], David B Lank[3], Jana Albrechtová[4], Jasmine L Loveland[1,5], Katrin Martin[1], Kim Teltscher[1], Margherita Cragnolini[1], Michael Lierz[6], Tomáš Albrecht[4,7], Wolfgang Forstmeier[1] & Bart Kempenaers[1]✉

[1]Max Planck Institute for Biological Intelligence, Eberhard-Gwinner-Str., 82319 Seewiesen, Germany
[2]Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Kamýcká 129, 165 21 Prague 6 – Suchdol, Czech Republic
[3]Department of Biological Sciences, Simon Fraser University, Burnaby, BC V5A 1S6, Canada
[4]Institute of Vertebrate Biology, Czech Academy of Sciences, Brno, Czech Republic
[5]Department of Cognitive and Behavioral Biology, University of Vienna, Vienna, Austria
[6]Clinic for Birds, Reptiles, Amphibians and Fish, Justus Liebig University Giessen, Giessen, Germany
[7]Department of Zoology, Faculty of Science, Charles University, Prague, Czech Republic
✉bulla.mar@gmail.com, b.kempenaers@orn.mpg.de

Abstract

Background. The ruff sandpiper *Calidris pugnax* is a Palearctic lekking shorebird with three genetic morphs determined by an autosomal inversion. Male morphs differ strikingly in body size, ornaments, endocrinology and mating behavior. Aggressive Independents represent the ancestral haplotype, while female-mimicking Faeders and semi-cooperative Satellites are the inverted haplotypes. Because one inversion breakpoint is homozygous lethal, the inverted haplotypes cannot recombine and are expected to accumulate deleterious mutations. The inversion regions also harbor genes involved in spermatogenesis. However, it remains unknown whether the genetic differences between the morphs also translate into differences in sperm traits. Here, we use a captive-bred population of ruffs to compare sperm swimming speed and morphology among the morphs.

Results. Ruff sperm morphologically resembled those of passerines, but moved differently, vibrating from side to side while slowly moving forward, rather than rotating while moving forward. Faeder sperm moved the slowest, which is consistent with the prediction of genetic deterioration over time. However, against expectation, sperm of Independents did not seem to be of the highest quality, i.e., their sperm were not the fastest nor the least variable, and they had sperm with the shortest tail and midpiece. Although the midpiece contains the energy-producing mitochondria, sperm midpiece length was not associated with sperm swimming speed. Instead, two of three velocity metrics weakly positively correlated with head length (absolute and relative).

Conclusions. The three genetically determined ruff morphs showed subtle differences in swimming speed and in the length of some sperm components. However, the between-morph differences in sperm swimming speed were not linked to the differences in morphology. We conclude that there is at best limited evidence for lower-quality sperm in the morphs that carry the inversion, and suggest that the potential for the evolution of morph-specific sperm adaptations may be limited in this system.

Lay Summary

The ruff sandpiper is a shorebird that exhibits three genetically distinct types of males, which differ markedly in body size, ornaments, hormones, and mating behavior. Aggressive Independents represent the type that evolved first. Semi-cooperative Satellites and female-mimicking Faeders evolved later through a specific type of genetic rearrangement called an 'inversion', in which a segment of a chromosome reversed in orientation. Due to the nature of this inversion, Satellite and Faeder chromosomes are expected to deteriorate over time. However, it remains unclear whether the genetic differences between these morphs, which affect physiological and behavioral traits, also translate into differences in sperm traits. We used a captive population of ruffs to compare sperm swimming speed and length measurements between the three types of males. Faeder sperm was the slowest, which is consistent with expectations based on genetic deterioration over time. However, against our expectations, the sperm of Independents does not appear to have better performance characteristics. Although the midpiece of a sperm is responsible for energy production, the length of the midpiece did not relate to sperm swimming speed.

Keywords: alternative mating strategies, autosomal inversion, genetic morphs, spermatozoa, sperm quality, sperm morphology, sperm velocity, testes

Background

In many animal species, males use different behavioral strategies to obtain fertilizations (Dougherty et al., 2022; Kustra & Alonzo, 2020; Mank, 2023). For example, dominant males may display, while other males "steal" copulations (Gross, 1996) by pretending to be a female or by positioning themselves in between the dominant male and the female that is about to copulate, a tactic referred to as "sneaking" (Jukema & Piersma, 2006; Shuster & Wade, 1991). Consequently, males that use alternative mating tactics often experience different levels of sperm competition, i.e., competition between the sperm from different males to fertilize one or more eggs (Parker, 1970). For example, if females prefer dominant males, it is likely that sneaker males more often experience sperm competition than dominant males. Comparative studies have shown that species experiencing higher levels of sperm competition typically produce sperm that swim faster, are longer (including longer sperm components), and less variable in length as a result of intense postcopulatory selection on optimal sperm length (Laskemoen et al., 2012; Lifjeld et al., 2010; Lipshutz et al., 2022; Lüpold et al., 2020; Simmons & Fitzpatrick, 2012). Within species, sperm fertilization ability has been linked to sperm viability (Gage et al., 2004; Garcia-Gonzalez & Simmons, 2005), swimming speed (Birkhead et al., 1999; Gage et al., 2004), and length (Bennison et al., 2014; Garcia-Gonzalez & Simmons, 2007; Lüpold et al., 2012). Thus, selection may have favored different sperm traits in males that consistently differ in mating behavior (Dougherty et al., 2022; Kustra & Alonzo, 2020). Despite the general expectation that sneaker males may have to produce more or more competitive sperm (higher quality or performance) to sire offspring, existing theory and data do not make clear predictions about how sperm traits of sneaker males should differ from those of other males (Dougherty et al., 2022; Kustra & Alonzo, 2020). Moreover, the evolutionary potential for sneakers to evolve superior sperm to compensate for their lower chances to copulate may be limited, because if sneaker males are rare in the population, the probability that a sneaker-specific beneficial mutation arises is low.

In the ruff *Calidris pugnax,* a lekking shorebird, males occur in three distinct morphs with striking differences in body size, ornaments, endocrinology and reproductive behavior (Hogan-Warburg, 1966; Höglund & Lundberg, 1989; Jukema & Piersma, 2006; Küpper et al., 2016; Loveland et al., 2021a; van Rhijn, 1991; Widemo, 1998). (1) Aggressive Independents make up 80-90% of all males and show a spectacular diversity of predominantly dark ornamental plumages. The Independents hold small display courts on a lek. Among these, the dominant male obtains the majority of matings (Vervoort & Kempenaers, 2019; Widemo, 1997). (2) Submissive and slightly smaller Satellites make up 10-20% of males, show predominantly white ornamental plumage, and co-display with court-holders on leks (Hogan-Warburg, 1966; Höglund & Lundberg, 1989; van Rhijn, 1991; Widemo, 1998). Although Satellites may "steal" some copulations, Independents may benefit from having Satellites on the lek because their presence helps attract females to the display court of that male (Hugie & Lank, 1997; Tolliver et al., 2023). (3) Faeders are rare (~1%), mimic females in appearance (lack of ornamental plumage and smaller body size), and attempt to sneak copulations when a female solicits an ornamented displaying male (Jukema & Piersma, 2006). Despite striking differences in body size, the three morphs have similar testes size (Figure 1; Jukema & Piersma, 2006; Küpper et al., 2016), and hence presumably can produce similar numbers of sperm. However, the morphs differ in their reproductive endocrinology. During the lekking season, Independent males have higher levels of circulating testosterone, whereas Satellites and Faeders have higher levels of androstenedione, a testosterone precursor (Küpper et al., 2016; Loveland et al., 2021a).

Two observations suggest that sperm competition in ruff is intense. First, more than half of all clutches are sired by multiple males (Lank et al., 2002; Thuman & Griffith, 2005). Second, ruffs have the longest sperm measured among shorebirds (Johnson & Briskie, 1999). Female ruff visit one or multiple leks, and copulate during 3-5 days starting about 5 days before laying their first egg, and they produce a clutch of 3 or 4 eggs (Lank et al., 2002). As in other birds, females store sperm in sperm storage tubules prior to fertilization (Briskie & Montgomerie, 1993). Females of a given morph are approximately 40% smaller than males of the same morph (Giraldo-Deck et al., 2020; Lank et al., 2013). Presumably because of their smaller size, Faeder females have markedly lower reproductive success compared to Independent and Satellite females (Giraldo-Deck et al., 2022).

124
The phenotypic differences between the morphs are entirely genetically determined by a 4.5 Mb
autosomal inversion region (Küpper et al., 2016; Lamichhaney et al., 2016). Faeders and Satellites carry
unique inversion haplotypes. The Faeder haplotype arose first (about 4 million years ago), whereas the
Satellite haplotype originated from a rare recombination event between the Independent and Faeder
genotypes several hundred thousand generations later, about 70,000 years ago (Hill et al., 2023;
Lamichhaney et al., 2016). Because one inversion breakpoint is homozygous lethal, the inversion
haplotypes cannot recombine and thus only occur in hemizygous state (i.e. always combined with one
Independent haplotype). As a result, the inversion haplotypes are expected to accumulate deleterious
mutations. Although the genetic differences between the morphs affect numerous phenotypic traits, it is
unknown whether they also affect sperm traits.

135
Intuitively, one might predict that the three morphs should differ in ejaculate size and in sperm traits due
to differences in copulation opportunity and frequency. When a female visits a lek and solicits a
copulation, the dominant Independent male is more likely to copulate once or multiple times with that
female compared to a Satellite male (Lank et al., 2002). Although behavioral evidence is lacking, it is
likely that copulation opportunities for Faeders are even more limited. If true, and given that the male
morphs do not differ in testes size (Figure 1), sperm allocation theory predicts that Satellites and Faeders
will transfer larger ejaculates per copulation than the dominant Independents (Pizzari et al., 2003).
Differences in ejaculate size and in the presence of multiple ejaculates will produce different cloacal and
oviductal environmental conditions in which sperm competition occurs. Such differences may have
selected for different optimal sperm traits in the three morphs. Specifically, selection pressure on
Faeders may be more intense as their ejaculate is more likely to compete with ejaculates of other males.
Although an evolutionary scenario that leads to more competitive sperm in Faeders and Satellites may
seem appealing, we argue that it is unlikely given the genetic make-up of the ruff morphs.

149
150 First, sperm traits and testicular gene expression may already have differed between morphs from the
151 moment the morphs originated. (i) Two genes regulating spermatogenesis reside within the autosomal
152 inversion that drives the Satellite and Faeder phenotypes (Küpper et al., 2016). The inversion contains
153 the gene *GAS8*, which has been linked to sperm motility in mice, and the gene *SPATA2L*, a paralog of

SPATA2, which is involved in spermatogenesis (Graziotto et al., 1999; Küpper et al., 2016; Onisto et al., 2000; Yeh et al., 2002). Moreover, expression of *SPATA2L* in testes of both Satellites and Faeders is biased toward the inversion allele (Loveland et al., 2021b). (ii) The inversion also influences expression of genes located outside of the inversion (Maney & Küpper, 2022). For example, *STAR,* a gene responsible for providing the cholesterol substrate required for sex hormone synthesis, is located outside of the inversion, but is overexpressed in the testes of Satellites and Faeders (Loveland et al., 2021a). (iii) Some genes or regulatory elements within the inversion region that have strong pleiotropic effects may also affect spermatogenesis.

Second, sperm trait differences between the morphs could also have arisen by *de novo* mutation after the origin of the three morphs, but through a different evolutionary scenario than the one described above. The genetic background outside of the inversion has been selected to function well in the context of the most abundant morph, i.e., the Independents (Giraldo-Deck et al., 2022). In contrast, even under strong selection due to intense sperm competition, Faeders, and to a lesser extent Satellites, have limited evolutionary potential due to (i) small population size, (ii) a small part (0.3%) of the genome within which a beneficial mutation would have to arise and (iii) a small number of genes within the inversion (only two of the ~125 genes identified) have known effects on sperm performance or morphology (Küpper et al., 2016). Under these conditions, it seems less likely that evolution will have led to more competitive sperm in Faeders and Satellites. In fact, if anything, it seems more likely that the two inversion morphs will have sperm of inferior quality, because the inversions do not recombine and deleterious mutations can thus accumulate inside the inversion haplotypes (Berdan et al., 2021). Under this scenario, we predict that Faeders have the lowest quality sperm because, (i) a larger number of genes may have accumulated potentially harmful mutations in the Faeder haplotype, given that it arose several million years earlier than the rather recent Satellite haplotype (Hill et al., 2023; Küpper et al., 2016; Lamichhaney et al., 2016) and (ii) the recombination through which Satellites arose may have re-instated "good" copies of a gene that (directly or indirectly) affects sperm quality.

Here, we use sperm sampled from a captive breeding population of ruffs to address three issues. First, we assess differences in sperm swimming speed and sperm (component) length between the three morphs. If longer sperm is better at fertilization (see references above) and the sperm of inversion morphs has been disadvantaged due to mutation accumulation, we expect the sperm of Independents to be (1) faster swimming, (2) longer, and (3) less variable in length. We would also expect the sperm of Independents to have (4) a longer midpiece (energetic component; Cardullo & Baltz, 1991; Knief et al., 2017; Míčková et al., 2023), (5) a longer tail and flagellum (midpiece + tail, kinetic component; Cardullo & Baltz, 1991), and possibly (6) a longer midpiece and flagellum length relative to total sperm length (Humphries et al., 2008).

Second, we test whether variation in ruff sperm morphology predicts variation in sperm swimming speed. Although a positive relationship between overall sperm length, midpiece or flagellum length and swimming speed can be predicted, previous studies, including those in birds, have shown inconsistent results (reviewed in Bennison et al., 2016; Cramer et al., 2021; Fitzpatrick & Lüpold, 2014; Kim et al., 2017; Knief et al., 2017; Míčková et al., 2023; Støstad et al., 2018).

Third, we describe the motility and shape of ruff sperm in comparison with that of other bird species. Sperm motility has not yet been described in scolopacids and only rarely in other species of shorebirds, and sperm shape has only been sketchily described in a few species in this group (Johnson & Briskie, 1999; Retzius, 1909). Furthermore, scolopacid sperm superficially resemble the sperm of passerines, but differ from those of all other (so far evaluated) avian lineages (Retzius, 1909).

Methods
We collected and analyzed the data based on *a priori* designed protocols. For any deviations from the (see Supplement Methods S0; Bulla et al., 2023).

Captive population
A population of ~300 ruffs is housed at the Max Planck Institute for Biological Intelligence, Seewiesen, Germany (for details see Supplement Methods S1). The population was founded in 2018 with 23 ruffs obtained from Dutch breeders, 5 ruffs from German breeders and 194 ruffs from Simon Fraser

University, Vancouver, Canada. The latter were individuals from a captive-bred population founded with 110 ruffs hatched from wild eggs collected in Finland in 1985, 1989 and 1990, plus two Faeder males brought from the Netherlands in 2006 (Lank et al., 2013; Lank et al., 1995). Each individual of the population has been genotyped for its morph using a set of six single-nucleotide polymorphism markers located in the inversion region (Giraldo-Deck et al., 2020).

Sperm sampling
In May and June 2021, we collected sperm by abdominal massage (for a detailed protocol of sperm collection – including video - and sample preparation see https://s.gwdg.de/UlejLot) or by electro-stimulation (Lierz et al., 2013). The length (15 mm) and diameter (4 mm) of the electro-stimulation probe, as well as the electric current and the number of electric impulses was adapted to the sampled individuals. The probe was inserted into the urodeum of the cloaca to stimulate the ampullae ductus deferens using three 1 s electric current impulses. The voltage was increased gradually from 0.09 V to a maximum of 1.0 V until contractions of the cloaca and the muscles of the tail were observed; each impulse was followed by a 2-3 s break. In June, we only sampled sperm using abdominal massage. The morphs were sampled in a haphazard order to avoid a potential confounding effect of sampling order. In both months, we attempted to obtain sperm by abdominal massage from all males. We obtained at least one ejaculate sample from 92 males (59 Independents, 25 Satellites and 8 Faeders).

Ejaculates (~0.5–3µl) were pipetted from the cloaca and immediately diluted and gently mixed in 50µl of preheated (40°C) Dulbecco's Modified Eagle's Medium (Advanced D-MEM, Invitrogen™). To record sperm swimming speed, we then pipetted an aliquot of 2.5µl onto a standard 20µm two-chamber count slide (Leja, The Netherlands) placed on a thermal plate (Tokai Hit, Tokai Hit Co., Ltd.) kept at 40 °C. When sperm densities on the slide were too high, we took a new aliquot and further diluted the sample. We then recorded sperm movements. By limiting the time between ejaculate collection and recording, we avoided that changes in sperm swimming speed over time (Cramer et al., 2016a) would influence the results. For morphology measurements, we pipetted an aliquot of 20µl into 50µl of a phosphate buffered saline solution (PBS; Sigma P-4417) containing 1% formalin.

Measurement of sperm swimming speed
For each sperm sample we recorded sperm swimming speed at 25 frames per second for approximately 45s in eight different fields of the Leja slide under a 100x magnification, using phase contrast and a digital camera (UI-1540-C, Olympus) mounted on a microscope (CX41, Olympus) fitted with a thermal plate (Tokai Hit, Tokai Hit Co., Ltd.) kept at a constant temperature of 40°C. We confirmed that this temperature was appropriate based on cloacal temperature measurements of five female ruffs. For each female, we inserted a high-resolution temperature-probe connected to a MSR145 data logger (MSR Electronics GmbH, https://www.msr.ch/en/) into the cloaca, and logged the temperature every 5s for 3 min, i.e. until an asymptotic value was reached. Cloacal temperatures ranged between 40.6°C and 42.3°C, and are similar to the body temperatures measured in shorebirds (Charadriiformes; range: 40.4-41.8°C, N = 10 species) and in passerines (39.2-43.5°C, N = 16 species; (McNab, 1966).

A few weeks later, a single person (JA) analyzed each recorded field using the CEROS computer-assisted sperm analysis system (Hamilton Thorne Inc.). JA visually inspected the tracked objects, excluded non-sperm objects and static spermatozoa from the analysis (Cramer et al., 2016b; Laskemoen et al., 2010; Opatová et al., 2016), and noted the quality of the recording (e.g., presence of faeces). We recorded a median of 192 sperm cells per sample (mean = 206, range: 5 – 562; N = 134 recordings from a total of 92 males, 46 recorded in May and 88 in June) and for each sample the software estimated the mean curvilinear, straight-line, and average-path velocity (for details see Supplement Methods S2 and Figure S1 and S2; for within-male seasonal repeatability see Figure S3 and S4, Table S1).

Measurements of sperm morphology
Within 5 days of sample collection, we pipetted (i) 20µl of the PBS-1% formalin fixed sample into 50µl of 5% formalin for long-term storage and (ii) smeared 10µl onto a microscope slide, and let it dry at room temperature or on a heating block set at 37°C. The next day, each slide was rinsed gently with deionized water to remove salt crystals and dried at room temperature.

To each dried slide we added a drop of Bisbenzimide (Hoechst 33342; Molecular Probes) to stain the
sperm nucleus, and a drop of MitoTracker™ Green FM Dye (Invitrogen™ M7514) to stain the sperm
midpiece (mitochondria; Picture 1). Within 48h of staining, we inspected the microscope slides under
200x magnification with a Zeiss Axio Imager.M2 light microscope fitted with a Zeiss Axiocam 512 color
camera (12 megapixel; 4250 × 2838, pixel size of 3.1µm × 3.1µm) and DAPI 465nm and green 519nm
filters using Zeiss ZEN blue 3.1 imaging software. For each male we used a single sample (30 from
May, 62 from June) and photographed at least 10 intact, normal-looking spermatozoa under 400x
magnification (objective size 40, ocular size 10). We then selected the 10 best, single-sperm images
per male for measurements. Previous studies also measured ten sperm per male to estimate the
coefficient of variation in sperm traits (Kleven et al., 2008; Lifjeld et al., 2010). We then randomized and
renamed all pictures, such that we measured sperm blind to the morph and the identity of the individuals.
280

281
282 **Picture 1 | Stained ruff sperm.** From left to right, faint greenish tail, green midpiece (mitochondria) stained by MitoTracker™
283 Green FM Dye, turquois nucleus stained by Hoechst 33342 and greenish acrosome.
284
For each sperm, we measured the length of the acrosome, the nucleus, the midpiece and the tail to the
nearest 0.1µm using the open software Sperm Sizer 1.6.6 (McDiarmid et al., 2021;
https://github.com/wyrli/sperm-sizer; for our Sperm Sizer protocol see https://s.gwdg.de/iGFz4F). To
assess within- and between-observer repeatability of Sperm Sizer measurements, we selected 40
single-sperm pictures and remeasured them, once by the same, once by a different person. The
measurements were highly repeatable, both within- and between persons (mean = 97%; Table S2).
Nevertheless, all measurements used in the analyses were taken by the same person (KT). Each
measured sperm and sperm part were numbered and referenced in the database and hence can be
linked to the original picture as well as to the pictures of each part generated by Sperm Sizer, which
contain the lines and measurement values (see Data in Bulla et al., 2023). This ensures transparency
and allows re-measurement of the same sperm by the same or a different person.
296
We calculated (i) total sperm length as the sum of all parts, (ii) head length as the sum of acrosome and
nucleus length, (iii) flagellum length as the sum of midpiece and tail length, (iv) relative midpiece length
as midpiece length divided by total length and (v) relative flagellum length as flagellum length divided
by total length. We then computed coefficients of variation within each male for each absolute trait as
the standard deviation divided by the mean. In the Supplement, we report correlations between sperm
components (Methods S3 and Figure S5), and estimates of within-male repeatability (Figure S3 and
Table S1).
304
Statistical analyses
All analyses were performed in R (R-Core-Team, 2022) using the 'lm' function to fit linear models and
the 'lmer' function of the 'lme4' R-package to fit linear mixed-effects models (Gelman & Hill, 2007;
Gelman & Su, 2021). We used the 'sim' function from the 'arm' R-package and a non-informative prior
distribution (Gelman & Hill, 2007; Gelman & Su, 2021) to create a sample of 5,000 simulated values for
each model parameter (i.e., posterior distribution). We report effect sizes and model predictions as
medians, and the uncertainty of the estimates and predictions as Bayesian 95% credible intervals
represented by the 2.5 and 97.5 percentiles (95% CI) from the posterior distribution of 5,000 simulated
or predicted values. Unless stated differently, plots were created with 'ggplot' function from the 'ggplo2'
R-package (Wickham, 2016).
315
316 To investigate whether sperm traits differ between the morphs we fitted linear models with each sperm
317 trait as a response variable, and with male morph as a predictor (three-level factor). For analyses of
318 velocity, we primarily used June values because the breeding season was at its peak in June and we
319 obtained data from all but four males (for those four we used May values). We added the number of

tracked sperm cells per sample (ln-transformed) as a covariate to the model to control for potential effects of sperm density or sample quality. For analyses of morphology, we used average male values. Alternative models using all velocity recordings, using individual sperm measurements, or controlling for sampling month or aviary gave similar results (Figure S6). Because correlations between sperm traits and inbreeding were weak (mean r = -0.09, range: -0.26 to 0.07; Figure S7, Methods S4), we did not include the inbreeding coefficient in the main models. Furthermore, the results were not confounded by relatedness (estimated based on microsatellite genotyping data) of the sampled individuals (see Methods S4, Table S3, Figure S8).

To investigate whether sperm morphology explained variation in sperm swimming speed, we fitted a set of linear models with each velocity measure as the dependent variable (using predominantly June values), and with the male average of each sperm morphology trait as a predictor. All models were controlled for the number of tracked sperm (ln-transformed) and morph (three-level factor). As the relationship between sperm swimming speed and morphology might be non-linear (Bennison et al., 2016), we also fitted a set of linear models that included the quadratic term of each morphology trait (2^{nd} order polynomial). We then tested whether the quadratic term improved the model fit, using Akaike's Information Criterion corrected for sample size (Anderson, 2008), with the 'AICc' function from the 'MuMIn' R-package (Bartoń, 2022). Because this was not the case for any trait (Table S4), we report results from the simple model without the quadratic term. To investigate whether the effect of one sperm component (head, midpiece and tail length) was influenced by the other components, we also ran a multivariate model that contained the three sperm components as predictors. Correlations between the three predictors were low (Figure S5). The effect sizes and uncertainties for head, midpiece and tail length were similar in the multivariate and the univariate models (Figure S9) and we report the results from the univariate models in the main text.

Results and Discussion
Between-morph differences in sperm traits
Sperm swimming speed in vitro in a standard medium, measured as curvilinear velocity, did not differ between the three morphs (Figure 2 and Supplement Figure S6 and Table S5; Bulla et al., 2023). However, when sperm swimming speed was measured as straight-line or average-path velocity, according to expectations based on mutation accumulation, Faeder sperm moved more slowly than the sperm of both Independents and Satellites, whereby the latter two morphs did not differ in sperm velocity (Figure 2 and S6, Table S5). If true, the slower sperm of Faeders appeared either with the origin of the Faeder genotype (i.e., the inversion) or later due to accumulation of deleterious mutations. In the latter case, recombination through which Satellites arose may have re-instated a "good" copy of a gene related to sperm velocity.

Sperm morphology differed weakly between the three morphs, mostly by less than one standard deviation (Figure 2 and S6, Table S5). Sperm of Independents were the shortest with the shortest midpiece and tail, and hence flagellum (midpiece and tail length combined). Acrosome and nucleus length, and hence head length, did not differ between Faeders and Independents. Satellite sperm had the shortest nucleus and head, but the longest tail. The midpiece was longest in Faeders and 1.19 standard deviations longer than the midpiece from Independents (95%CI: 0.46 - 1.88; Figure 2, Table S5). Total sperm length and flagellum length were similar for Faeders and Satellites (Figure 2 and S6). Consequently, midpiece length relative to total sperm length was largest for Faeders, but similar for Satellites and Independents; flagellum length relative to total sperm length was largest in Satellites and similar in the other two morphs (Figure S6). The coefficients of variation in sperm traits were noisy with unclear trends and sperm of Independents was not less variable than that of the Faeders and Satellites (Figure 2 and S8). These findings do not support the prediction of mutation accumulation that sperm in Satellites and particularly in Faeders should be of lower quality. Specifically, when compared to Independents, the sperm of the two derived morphs were not shorter or more variable in terms of midpiece, tail, and flagellum. Thus, sperm of Independents seemed of similar or lower (not better) quality than sperm of the derived morphs (Figure 2), although what constitutes a high quality sperm, especially within species, is still debated (Dougherty et al., 2022).

Figure 2 | Differences in sperm traits between ruff morphs. (**a**) Model predictions of between-morph differences with their 95% CI (Table S5). (**b**) Morph-specific sperm traits, shown as individual data points with dot color highlighting morph (black: Independents, white: Satellites, beige: Faeders) and summarized as boxplots (see Fig. 1 for definition). To aid visualization, the outlier coefficients of variation for tail (18.7) and total length (9.8) in Independents are depicted as 8 and 6, respectively. The red dots with bars show model predictions and their 95% CI (Table S5). Dots were stacked using the 'geom_dotplot' ggplot2 R-function (Wickham, 2016). (**a, b**) Velocity represents June values (with the exception of four males with May values only). Morphology represents average trait length and coefficients of variation are based on measurements of 10 sperm cells per male. For estimates of head, flagellum (absolute and relative) and relative midpiece length, see Fig. S6 and S8. Ruff morph illustrations by Yifan Pei under Creative Commons Attribution (CC BY 4.0).

The lack of major differences in sperm morphology among the three male morphs suggests that the two genes related to spermatogenesis that reside within the inversion have little impact on the measured sperm traits. Such finding might be unsurprising given that tens to hundreds of genes influence spermatogenesis (Linn et al., 2021). Furthermore, the lack of major differences in sperm traits accords with previous studies showing no clear and consistent differences in sperm traits between males that use alternative mating tactics (Dougherty et al., 2022; Kustra & Alonzo, 2020). Although one expects a steeper selection gradient for Faeder and Satellite males that have a lower probability to copulate (reviewed in Dougherty et al., 2022; Kustra & Alonzo, 2020), selection will act on all male morphs to optimize sperm traits. Moreover, independent of the strength of sexual selection, Faeder-beneficial mutations are about 30,000 times less likely to arise than Independent-beneficial mutations, given a Faeder population size of 1% and the inversion being 0.3% of the diploid genome ($1/(0.01 \times 0.003)$).

It remains unclear whether and how the success of Faeder males would compensate for the lower reproductive success of Faeder females (Giraldo-Deck et al., 2022). The measured sperm traits seem to give the Faeder males little advantage over Independent males. However, there is a strong mating skew with only few Independents obtaining most copulations on a given lek (Tolliver et al., 2023; Vervoort & Kempenaers, 2019; Widemo & Owens, 1995), such that the average reproductive success of an Independent will be low. Thus, the rare Faeder males might only need moderate mating success through sneaking, facilitated by their female mimicking appearance and behavior (Jukema & Piersma, 2006) to compensate. Possibly, Faeders do better, because morphs differ in other, non-measured, but

fitness-relevant sperm traits. For example, the longer midpiece of Faeder sperm might correlate with longer sperm survival inside the female reproductive tract. However, this would need further study.

Relationship between sperm swimming speed and morphology
Sperm morphology had only weak and unclear effects on variation in sperm swimming speed (Figure 3, S9 and S10, Table S6). Longer sperm, sperm with a longer midpiece (absolute and relative) or sperm with a longer flagellum did not swim faster (r varied between 0.05 and 0.18; Figure S10). The lack of an association between sperm midpiece length and swimming speed may have several explanations. First, our method to measure sperm velocity might not reflect the swimming speed of the sperm within the seminal fluid or within the female reproductive tract. Note, however, that the way the sperm moved did not change after we placed the sperm into a solution that contained cloacal fluid of a ruff female (details not shown). Also, our method is the same as the one used to measure swimming speed in passerine sperm, and we used the exact same equipment and observers as in some of the studies on passerines (e.g., Knief et al., 2017; Míčková et al., 2023; Opatová et al., 2016). Second, although the midpiece contains the mitochondria, midpiece volume rather than length might be relevant for sperm swimming speed (Cardullo & Baltz, 1991; Mendonca et al., 2018), but see Cramer et al. (2022), or midpiece length might be associated with other sperm performance parameters such as longevity (which we did not measure). Third, it remains unclear whether oxidative phosphorylation in the midpiece is the primary source of energy for moving sperm (Turner, 2003).

Against the predictions (Humphries et al., 2008), sperm with longer heads (nucleus) or smaller relative flagellum length swam faster, when measured as curvilinear and average-path velocity (but not straight-line velocity), although these effects were small (< 0.25 of standard deviation, Figure 3; maximum r = 0.22, Figure S10). Theory predicts that sperm with a shorter head relative to flagellum should swim faster because a relatively shorter head reduces drag (Humphries et al., 2008). The few existing within-species studies provide equivocal evidence for this prediction (Cramer et al., 2015; Cramer et al., 2021; Helfenstein et al., 2009; Mossman et al., 2009), but the prediction has been supported in a between-species study in mammals (Tourmente et al., 2011) and in passerine birds (Lüpold et al., 2009). A recent study on passerines showed that sperm with a long acrosome, a short nucleus, a wide helical membrane, and a more pronounced waveform along the sperm head "core" swam faster (Støstad et al., 2018).

Sperm motility and sperm shape in ruffs compared to other taxa
We observed that ruff sperm seemed to move in a different manner to that previously described for passerine and likely also for non-passerine sperm ([video examples](#)). Specifically, ruff sperm seem to 'vibrate' from side to side while slowly moving forward, whereas passerine sperm typically swim straight by rapidly rotating around their longitudinal axis (Ballowitz, 1888; Humphreys, 1972; Vernon & Woolley, 1999). We can exclude that the unusual head movements of ruff sperm were caused by a lack of depth in the standard 20μm two-chamber count Leja-slide, because using a deeper Leja-slide (100μm) resulted in similar movements (not recorded). However, our observations need further verification using other methods (Vernon & Woolley, 1999). If confirmed, the difference in movement may not be due to differences in general morphology, because ruff sperm have a helical, screw-like head (Picture 1), similar to that observed in passerine sperm (Retzius, 1909). Previous studies described that non-passerine sperm moves with regular helical (dextral) waves (Ballowitz, 1888; Humphreys, 1972; Vernon & Woolley, 1999) or via counter-clockwise turning of the entire sperm body (Bird & Laguë, 1977). Given the few descriptions and the lack of video recordings of non-passerine sperm movements (e.g., Cheng et al., 2002; Denk et al., 2005; Dogliero et al., 2017; Fischer et al., 2014; Gloria et al., 2014), it remains to be seen whether the way ruff sperm propel is unique to this species, or typical for other Scolopacids, shorebirds or Charadriiformes in general. Clearly, the directional movement of ruff sperm is distinct from the smooth, snake-like, movements of sperm of non-avian taxa (e.g., human or sea-urchin *Psammechinus miliaris* (Cosson et al., 2015; Gillies et al., 2009; Gray, 1955; Saggiorato et al., 2017; Smith et al., 2009), but see squid sperm (Bishop, 1958)).

The screw-like head of ruff sperm is similar to the head shape reported for other Scolopaci (sandpipers), but differs from the straight heads observed in other Charadriiformes, e.g., genus *Larus* or *Vanellus* (Retzius, 1909). Our results confirm that ruff sperm is remarkably long (median and mean = 139.5 μm, range: 105.7 - 150 μm) compared to other shorebirds, with species means ranging between 57 and 100 μm (Johnson & Briskie, 1999).

Figure 3 | Effects of sperm morphology on sperm swimming speed. Shapes with horizontal bars represent estimated standardized effect sizes with their 95%CIs based on the joint posterior distribution of 5,000 simulated values generated from linear models, controlled for the number of tracked sperm (ln-transformed) and morph (Table S6). The results are based on single June-values for velocity (with the exception of four males with May-values only) and average trait length of 10 sperm cells per male. For raw data and predicted relationships see Figure S10.

Conclusions

Using sperm collected from a population of captive ruffs, we found between-morph differences in sperm swimming speed and length measures. As expected from mutation accumulation in the inversion, sperm of Faeders were the slowest, but – against expectation – sperm of Independents were not the fastest, longest or least variable. Furthermore, we found that sperm of ruffs moved differently than sperm of passerines, despite having a similar head shape (Ballowitz, 1888; Humphreys, 1972; Retzius, 1909; Vernon & Woolley, 1999). Our study shows at best weak associations between sperm swimming speed and length measurements that are inconsistent with general expectations. Our results corroborate other comparative work that revealed a lack of clear and consistent differences in sperm traits of males using alternative mating tactics (Dougherty et al., 2022; Kustra & Alonzo, 2020). Our study leaves open the question about whether and how Faeder males can achieve higher reproductive success to compensate for the lower reproductive output of Faeder females (Giraldo-Deck et al., 2022), but emphasizes little potential for the evolution of strong morph-specific sperm adaptations in this system.

Availability of data and materials

All data, including sperm recordings and pictures, and computer code used to generate the results and Supplement (figures, tables and documents) of this study are freely available at GitHub: https://github.com/MartinBulla/ruff_sperm_v2 (Bulla et al., 2023).

References

Ballowitz, E. (1888). Untersuchungen über die Struktur der Spermatozoën, zugleich ein Beitrag zur Lehre vom feineren Bau der contraktilen Elemente. *Archiv für Mikroskopische Anatomie, 32*(1), 401-473.

Bartoń, K. (2022). MuMIn: Multi-Model Inference. *R package version 1.46.0.* doi:https://CRAN.R-project.org/package=MuMIn

Bennison, C., Hemmings, N., Brookes, L., Slate, J., & Birkhead, T. (2016). Sperm morphology, adenosine triphosphate (ATP) concentration and swimming velocity: unexpected relationships in a passerine bird. *Proc Biol Sci, 283*(1837). doi:https://doi.org/10.1098/rspb.2016.1558

Bennison, C., Hemmings, N., Slate, J., & Birkhead, T. (2014). Long sperm fertilize more eggs in a bird. *Proceedings of the Royal Society B: Biological Sciences, 282*(1799), 20141897-20141897. doi:https://doi.org/10.1098/rspb.2014.1897

Berdan, E. L., Blanckaert, A., Butlin, R. K., & Bank, C. (2021). Deleterious mutation accumulation and the long-term fate of chromosomal inversions. *PLoS Genetics, 17*(3), e1009411. doi:https://doi.org/10.1371/journal.pgen.1009411

Bird, D. M., & Laguë, P. C. (1977). Semen production of the American kestrel. *Canadian Journal of Zoology, 55*(8), 1351-1358. doi:https://doi.org/10.1139/z77-175

Birkhead, T. R., J. G. Martínez, Burke, T., & Froman, D. P. (1999). Sperm mobility determines the outcome of sperm competition in the domestic fowl. *Proceedings of the Royal Society B: Biological Sciences, 266*, 1759-1764. doi:https://doi.org/10.1098/rspb.1999.0843

Bishop, D. W. (1958). Motility of the sperm flagellum. *Nature, 182*, 1638-1640. doi:https://www.nature.com/articles/1821638a0

Briskie, J. V., & Montgomerie, R. (1993). Patterns of sperm storage in relation to sperm competition in passerine birds. *The*
Condor, 95, 442-454.
Bulla, M., Küpper, C., Lank, D. B., Albrechtová, J., Loveland, J. L., Martin, K., . . . Kempenaers, B. (2023). Supporting
information for '"Sperm traits of the three genetic morphs in the ruff sandpiper ": GitHub. Retrieved from
https://github.com/MartinBulla/ruff_sperm_v2
Cardullo, R. A., & Baltz, J. M. (1991). Metabolic Regulation in Mammalian Sperm: Mitochondrial Volume Determines Sperm
Length and Flagellar Beat Frequency. *Cell Motility and the Cytoskeleton, 19*, 180-188. doi:
https://doi.org/10.1002/cm.970190306
Cheng, F. P., Guo, T. J., Wu, J. T., Lin, T. E., Ursem, P. J., Colenbrander, B., & Fung, H. P. (2002). Annual variation in semen
characteristics of pigeons (*Columba livia*). *Poultry Science, 81*(7), 1050-1056. doi:https://doi.org/10.1093/ps/81.7.1050
Cosson, J. J., Prokopchuk, G., & Inaba, K. (2015). The Flagellar Mechanics of Spermatozoa and its Regulation. In J. J. Cosson
(Ed.), *Flagellar Mechanics and Sperm Guidance* (pp. 3-134). doi:https://doi.org/10.2174/97816810812811150101
Cramer, E. R., Alund, M., McFarlane, S. E., Johnsen, A., & Qvarnstrom, A. (2016a). Females discriminate against heterospecific
sperm in a natural hybrid zone. *Evolution, 70*(8), 1844-1855. doi:10.1111/evo.12986
Cramer, E. R., Alund, M., McFarlane, S. E., Johnsen, A., & Qvarnstrom, A. (2016b). Females discriminate against heterospecific
sperm in a natural hybrid zone. *Evolution*. doi:https://doi.org/10.1111/evo.12986
Cramer, E. R., Laskemoen, T., Stensrud, E., Rowe, M., Haas, F., Lifjeld, J. T., . . . Johnsen, A. (2015). Morphology-function
relationships and repeatability in the sperm of Passer sparrows. *Journal of Morphology, 276*(4), 370-377.
doi:10.1002/jmor.20346
Cramer, E. R. A., Garcia-Del-Rey, E., Johannessen, L. E., Laskemoen, T., Marthinsen, G., Johnsen, A., & Lifjeld, J. T. (2021).
Longer Sperm Swim More Slowly in the Canary Islands Chiffchaff. *Cells, 10*(6).
doi:https://doi.org/10.3390/cells10061358
Cramer, E. R. A., Gronstol, G., & Lifjeld, J. T. (2022). Flagellum tapering and midpiece volume in songbird spermatozoa. *Journal*
of Morphology, 283(12), 1577-1589. doi:https://doi.org/10.1002/jmor.21524
Denk, A. G., Holzmann, A., Peters, A., Vermeirssen, E. L. M., & Kempenaers, B. (2005). Paternity in mallards: effects of sperm
quality and female sperm selection for inbreeding avoidance. *Behavioral Ecology, 16*(5), 825-833.
doi:https://doi.org/10.1093/beheco/ari065
Dogliero, A., Rota, A., Lofiego, R., von Degerfeld, M. M., & Quaranta, G. (2017). Semen Collection and Spermatozoa
Characteristics in the Kea Parrot (*Nestor notabilis*). *Journal of Avian Medicine and Surgery, 31*(1), 24-28.
doi:https://doi.org/10.1647/2015-132
Dougherty, L. R., Skirrow, M. J. A., Jennions, M. D., & Simmons, L. W. (2022). Male alternative reproductive tactics and sperm
competition: a meta-analysis. *Biol Rev Camb Philos Soc*. doi:https://doi.org/10.1111/brv.12846
Fischer, D., Neumann, D., Wehrend, A., & Lierz, M. (2014). Comparison of conventional and computer-assisted semen analysis
in cockatiels (*Nymphicus hollandicus*) and evaluation of different insemination dosages for artificial insemination.
Theriogenology, 82(4), 613-620. doi:https://doi.org/10.1016/j.theriogenology.2014.05.023
Fitzpatrick, J. L., & Lüpold, S. (2014). Sexual selection and the evolution of sperm quality. *Mol Hum Reprod, 20*(12), 1180-1189.
doi:https://doi.org/10.1093/molehr/gau067
Gage, M. J. G., Macfarlane, C. P., Yeates, S., Ward, R. G., Searle, J. B., & Parker, G. A. (2004). Spermatozoal Traits and
Sperm Competition in Atlantic Salmon. *Current Biology, 14*(1), 44-47. doi:https://doi.org/10.1016/j.cub.2003.12.028
Garcia-Gonzalez, F., & Simmons, L. W. (2005). Sperm viability matters in insect sperm competition. *Current Biology, 15*(3), 271-
275. doi:https://doi.org/10.1016/j.cub.2005.01.032
Garcia-Gonzalez, F., & Simmons, L. W. (2007). Shorter sperm confer higher competitive fertilization success. *Evolution, 61*(4),
816-824. doi:10.1111/j.1558-5646.2007.00084.x
Gelman, A., & Hill, J. (2007). *Data analysis using regression and multilevel/hierarchical models*. Cambridge: Cambridge
University Press.
Gelman, A., & Su, Y.-S. (2021). arm: Data Analysis Using Regression and Multilevel/Hierarchical Models. R package version
1.12-2. (Version R package version 1.12-2.). Retrieved from http://CRAN.R-project.org/package=arm
Gillies, E. A., Cannon, R. M., Green, R. B., & Pacey, A. A. (2009). Hydrodynamic propulsion of human sperm. *Journal of Fluid*
Mechanics, 625, 445-474. doi:https://doi.org/10.1017/s0022112008005685
Giraldo-Deck, L. M., Loveland, J. L., Goymann, W., Tschirren, B., Burke, T., Kempenaers, B., . . . Küpper, C. (2022). Intralocus
conflicts associated with a supergene. *Nat Commun, 13*(1), 1384. doi:https://doi.org/10.1038/s41467-022-29033-w
Giraldo-Deck, L. M., Goymann, W., Safari, I., Dawson, D. A., Stocks, M., Burke, T., . . . Küpper, C. (2020). Development of
intraspecific size variation in black coucals, white-browed coucals and ruffs from hatching to fledging. *Journal of Avian*
Biology, 51(8). doi:https://doi.org/10.1111/jav.02440
Gloria, A., Contri, A., Carluccio, A., Parrillo, S., Cicconi, M., & Robbe, D. (2014). The breeding management affects fresh and
cryopreserved semen characteristics in *Melopsittacus undulatus*. *Anim Reprod Scis, 144*(1-2), 48-53.
doi:https://doi.org/10.1016/j.anireprosci.2013.11.009
Gray, J. (1955). The Movement of Sea-Urchin Spermatozoa. *Journal of Experimental Biology, 32*(4), 775–801.
doi:https://doi.org/10.1242/jeb.32.4.775
Graziotto, R., Foresta, C., Scannapieco, P., Zeilante, P., Russo, A., Negro, A., . . . Onisto, M. (1999). cDNA Cloning and
Characterization of PD1: A Novel Human Testicular Protein with Different Expressions in Various Testiculopathies.
Experimental Cell Research, 248(2), 620-626. doi:https://doi.org/10.1006/excr.1999.4449
Gross, M. R. (1996). Alternative reproductive strategies and tactics: diversity within sexes. *Trends in Ecology & Evolution, 11*(2),
92-98. doi:https://doi.org/10.1016/0169-5347(96)81050-0
Helfenstein, F., Podevin, M., & Richner, H. (2009). Sperm morphology, swimming velocity, and longevity in the house sparrow
Passer domesticus. Behavioral Ecology and Sociobiology, 64(4), 557-565. doi:10.1007/s00265-009-0871-x
Hill, J., Enbody, E. D., Bi, H., Lamichhaney, S., Lei, W., Chen, J., . . . Andersson, L. (2023). Low Mutation Load in a Supergene
Underpinning Alternative Male Mating Strategies in Ruff (Calidris pugnax). *Molecular Biology and Evolution, 40*(12).
doi:10.1093/molbev/msad224
Hogan-Warburg, A. J. (1966). Social Behavior of the Ruff, *Philomachus Pugnax* (L.). *Ardea, 54*(3-4), 109-229.
doi:https://doi.org/10.5253/arde.v54.p109

Höglund, J., & Lundberg, A. (1989). Plumage Color Correlates with Body Size in the Ruff *The Auk,* **106**(2), 336–338.
doi:https://doi.org/10.1093/auk/106.2.336
Hugie, D. M., & Lank, D. B. (1997). The resident's dilemma: a female choice model for the evolution of alternative mating
strategies in lekking male ruffs (*Philomachus pugnax*). *Behavioral Ecology,* **8**(2), 218-225.
doi:https://doi.org/10.1093/beheco/8.2.218
Humphreys, P. (1972). Brief observations on the semen and spermatozoa of certain passerine and non-passerine birds. *Journal
of Reproduction and Fertility,* **29**(3), 327-336. doi:https://doi.org/10.1530/jrf.0.0290327
Humphries, S., Evans, J. P., & Simmons, L. W. (2008). Sperm competition: linking form to function. *BMC Evolutionary Biology,*
8, 319. doi:https://doi.org/10.1186/1471-2148-8-319
Johnson, D. D. P., & Briskie, J. V. (1999). Sperm competition and sperm length in shorebirds. *Condor,* **101**(4), 848-854.
doi:https://doi.org/10.2307/1370074
Jukema, J., & Piersma, T. (2006). Permanent female mimics in a lekking shorebird. *Biology Letters,* **2**(2), 161-164.
doi:https://doi.org/10.1098/rsbl.2005.0416
Kim, K.-W., Bennison, C., Hemmings, N., Brookes, L., Hurley, L. L., Griffith, S. C., . . . Slate, J. (2017). A sex-linked supergene
controls sperm morphology and swimming speed in a songbird. *Nature Ecology & Evolution,* **1**(8), 1168-1176.
doi:https://doi.org/10.1038/s41559-017-0235-2
Kleven, O., Laskemoen, T., Fossøy, F., Robertson, R. J., & Lifjeld, J. T. (2008). Intraspecific variation in sperm length is
negatively related to sperm competition in passerine birds. *Evolution,* **62**(2), 494-499.
doi:https://doi.org/10.1111/j.1558-5646.2007.00287.x
Knief, U., Forstmeier, W., Pei, Y., Ihle, M., Wang, D., Martin, K., . . . Kempenaers, B. (2017). A sex-chromosome inversion
causes strong overdominance for sperm traits that affect siring success. *Nature Ecology & Evolution,* **1**(8), 1177-
1184. doi:https://doi.org/10.1038/s41559-017-0236-1
Küpper, C., Stocks, M., Risse, J. E., Dos Remedios, N., Farrell, L. L., McRae, S. B., . . . Burke, T. (2016). A supergene
determines highly divergent male reproductive morphs in the ruff. *Nat Genet,* **48**, 79-83.
doi:https://doi.org/10.1038/ng.3443
Kustra, M. C., & Alonzo, S. H. (2020). Sperm and alternative reproductive tactics: a review of existing theory and empirical data.
Philosophical Transactions of the Royal Society of London B Biological Sciences, **375**(1813), 20200075.
doi:https://doi.org/10.1098/rstb.2020.0075
Lamichhaney, S., Fan, G., Widemo, F., Gunnarsson, U., Thalmann, D. S., Hoeppner, M. P., . . . Andersson, L. (2016). Structural
genomic changes underlie alternative reproductive strategies in the ruff (*Philomachus pugnax*). *Nat Genet,* **48**, 84-88.
doi:https://doi.org/10.1038/ng.3430
Lank, D. B., Farrell, L. L., Burke, T., Piersma, T., & McRae, S. B. (2013). A dominant allele controls development into female
mimic male and diminutive female ruffs. *Biology Letters,* **9**(6). doi:https://doi.org/10.1098/rsbl.2013.0653
Lank, D. B., Smith, C. M., Hanotte, O., Burke, T., & Cooke, F. (1995). Genetic polymorphism for alternative mating behaviour in
lekking male ruff *Philomachus pugnax*. *Nature,* **378**(6552), 59-62. doi:https://doi.org/10.1038/378059a0
Lank, D. B., Smith, C. M., Hanotte, O., Ohtonen, A., Bailey, S., & Burke, T. (2002). High frequency of polyandry in a lek mating
system. *Behavioral Ecology,* **13**(2), 209-215. doi:https://doi.org/10.1093/beheco/13.2.209
Laskemoen, T., Albrecht, T., Bonisoli-Alquati, A., Cepak, J., Lope, F., Hermosell, I., . . . Lifjeld, J. (2012). Variation in sperm
morphometry and sperm competition among barn swallow (*Hirundo rustica*) populations. *Behavioral Ecology and
Sociobiology*, 1-9. doi:https://doi.org/10.1007/s00265-012-1450-0
Laskemoen, T., Kleven, O., Fossøy, F., Robertson, R. J., Rudolfsen, G., & Lifjeld, J. T. (2010). Sperm quantity and quality
effects on fertilization success in a highly promiscuous passerine, the tree swallow *Tachycineta bicolor*. *Behavioral
Ecology and Sociobiology*, 1-11. doi: https://doi.org/10.1007/s00265-010-0962-8
Lierz, M., Reinschmidt, M., Muller, H., Wink, M., & Neumann, D. (2013). A novel method for semen collection and artificial
insemination in large parrots (Psittaciformes). *Sci Rep,* **3**, 2066. doi:https://doi.org/10.1038/srep02066
Lifjeld, J. T., Laskemoen, T., Kleven, O., Albrecht, T., & Robertson, R. J. (2010). Sperm Length Variation as a Predictor of
Extrapair Paternity in Passerine Birds. *PLoS ONE,* **5**(10), e13456. doi:https://doi.org/10.1371/journal.pone.0013456
Linn, E., Ghanem, L., Bhakta, H., Greer, C., & Avella, M. (2021). Genes Regulating Spermatogenesis and Sperm Function
Associated With Rare Disorders. *Front Cell Dev Biol,* **9**, 634536. doi:https://doi.org/10.3389/fcell.2021.634536
Lipshutz, S. E., Torneo, S. J., & Rosvall, K. A. (2022). How female-female competition affects male-male competition: insights
on post-copulatory sexual selection from socially polyandrous species. *The American Naturalist*.
doi:https://doi.org/10.1086/722799
Loveland, J. L., Giraldo-Deck, L. M., Lank, D. B., Goymann, W., Gahr, M., & Küpper, C. (2021a). Functional differences in the
hypothalamic-pituitary-gonadal axis are associated with alternative reproductive tactics based on an inversion
polymorphism. *Hormones and Behavior,* **127**, 104877. doi:https://doi.org/10.1016/j.yhbeh.2020.104877
Loveland, J. L., Lank, D. B., & Küpper, C. (2021b). Gene Expression Modification by an Autosomal Inversion Associated With
Three Male Mating Morphs. *Front Genet,* **12**, 641620. doi:https://doi.org/10.3389/fgene.2021.641620
Lüpold, S., Calhim, S., Immler, S., & Birkhead, T. R. (2009). Sperm morphology and sperm velocity in passerine birds. *Proc Biol
Sci,* **276**(1659), 1175-1181. doi:https://doi.org/10.1098/rspb.2008.1645
Lüpold, S., de Boer, R. A., Evans, J. P., Tomkins, J. L., & Fitzpatrick, J. L. (2020). How sperm competition shapes the evolution
of testes and sperm: a meta-analysis. *Philosophical Transactions of the Royal Society of London B Biological
Sciences,* **375**(1813), 20200064. doi:https://doi.org/10.1098/rstb.2020.0064
Lüpold, S., Manier, Mollie K., Berben, Kirstin S., Smith, Kyle J., Daley, Bryan D., Buckley, Shannon H., . . . Pitnick, S. (2012).
How Multivariate Ejaculate Traits Determine Competitive Fertilization Success in *Drosophila melanogaster*. *Current
biology : CB*. doi:https://doi.org/10.1016/j.cub.2012.06.059
Maney, D. L., & Küpper, C. (2022). Supergenes on steroids. *Philosophical Transactions of the Royal Society of London B
Biological Sciences,* **377**(1855), 20200507. doi:https://doi.org/10.1016/10.1098/rstb.2020.0507
Mank, J. E. (2023). Sex-specific morphs: the genetics and evolution of intra-sexual variation. *Nature Reviews Genetics,* **24**(1),
44-52. doi:https://doi.org/10.1038/s41576-022-00524-2
McDiarmid, C. S., Li, R., Kahrl, A. F., Rowe, M., & Griffith, S. C. (2021). Sperm Sizer: a program to semi-automate the
measurement of sperm length. *Behavioral Ecology and Sociobiology,* **75**(5). doi:https://doi.org/10.1007/s00265-021-
03013-4

McNab, B. K. (1966). An analysis of the body temperatures of birds. *The Condor, 68*(1), 47-55.
doi:https://doi.org/10.2307/1365174
Mendonca, T., Birkhead, T. R., Cadby, A. J., Forstmeier, W., & Hemmings, N. (2018). A trade-off between thickness and length
in the zebra finch sperm mid-piece. *Proc Biol Sci, 285*(1883). doi:10.1098/rspb.2018.0865
Míčková, K., Tomášek, O., Jelínek, V., Šulc, M., Pazdera, L., Albrechtová, J., & Albrecht, T. (2023). Age-related changes in
sperm traits and evidence for aging costs of sperm production in a sexually promiscuous passerine. *Frontiers in
Ecology and Evolution, 11*. doi:https://doi.org/10.3389/fevo.2023.1105596
Mossman, J., Slate, J., Humphries, S., & Birkhead, T. (2009). Sperm morphology and velocity are genetically codetermined in
the zebra finch. *Evolution, 63*(10), 2730-2737. doi:10.1111/j.1558-5646.2009.00753.x
Onisto, M., Graziotto, R., Scannapieco, P., Marin, P., Merico, M., Slongo, M. L., & Foresta, C. (2000). A novel gene (PD1) with a
potential role on rat spermatogenesis. *Journal of Endocrinological Investigation volume, 23*, 605–608.
doi:https://doi.org/10.1007/BF03343783
Opatová, P., Ihle, M., Albrechtová, J., Tomášek, O., Kempenaers, B., Forstmeier, W., & Albrecht, T. (2016). Inbreeding
depression of sperm traits in the zebra finch *Taeniopygia guttata*. *Ecol Evol, 6*(1), 295-304.
doi:https://doi.org/10.1002/ece3.1868
Parker, G. A. (1970). Sperm Competition and Its Evolutionary Consequences in the Insects. *Biological Reviews (Cambridge),
45*, 525-567. doi: https://doi.org/10.1111/j.1469-185X.1970.tb01176.x
Pizzari, T., Cornwallis, C. K., Lovlie, H., Jakobsson, S., & Birkhead, T. R. (2003). Sophisticated sperm allocation in male fowl.
Nature, 426(6962), 70-74. Retrieved from http://dx.doi.org/10.1038/nature02004
R-Core-Team. (2022). R: A language and environment for statistical computing (Version Version 4.2.0). Vienna (Austria): R
Foundation for Statistical Computing. Vienna, Austria. Retrieved from http://www.R-project.org/
Retzius, G. (1909). Die Spermien der Vögel. *Biol. Unters. New Ser, 14*, 89-122.
Saggiorato, G., Alvarez, L., Jikeli, J. F., Kaupp, U. B., Gompper, G., & Elgeti, J. (2017). Human sperm steer with second
harmonics of the flagellar beat. *Nat Commun, 8*(1), 1415. doi:https://doi.org/10.1038/s41467-017-01462-y
Shuster, S. M., & Wade, M. J. (1991). Equal mating success among male reproductive strategies in a marine isopod. *Nature,
350*, 608–610. doi:https://doi.org/10.1038/350608a0
Simmons, L. W., & Fitzpatrick, J. L. (2012). Sperm wars and the evolution of male fertility. *Reproduction, 144*(5), 519-534.
doi:https://doi.org/10.1530/REP-12-0285
Smith, D. J., Gaffney, E. A., Gadelha, H., Kapur, N., & Kirkman-Brown, J. C. (2009). Bend propagation in the flagella of
migrating human sperm, and its modulation by viscosity. *Cell Motility and the Cytoskeleton, 66*(4), 220-236.
doi:https://doi.org/10.1002/cm.20345
Støstad, H. N., Johnsen, A., Lifjeld, J. T., & Rowe, M. (2018). Sperm head morphology is associated with sperm swimming
speed: A comparative study of songbirds using electron microscopy. *Evolution, 72*(9), 1918-1932.
doi:https://doi.org/10.1111/evo.13555
Thuman, K. A., & Griffith, S. C. (2005). Genetic similarity and the nonrandom distribution of paternity in a genetically highly
polyandrous shorebird. *Animal Behaviour, 69*(4), 765-770. doi:https://doi.org/10.1016/j.anbehav.2004.10.003
Tolliver, J. D. M., Kupán, K., Lank, D. B., Schindler, S., & Küpper, C. (2023). Fitness benefits from co-display favour
subdominant male–male partnerships between phenotypes. *Animal Behaviour, 197*, 131-154.
doi:https://doi.org/10.1016/j.anbehav.2022.12.004
Tourmente, M., Gomendio, M., & Roldan, E. R. (2011). Sperm competition and the evolution of sperm design in mammals. *BMC
Evolutionary Biology, 11*(12). doi:https://doi.org/10.1186/1471-2148-11-12
Turner, R. M. (2003). Tales from the tail: what do we really know about sperm motility? *J Androl, 24*(6), 790-803.
doi:https://doi.org/10.1002/j.1939-4640.2003.tb03123.x
van Rhijn, J. G. (1991). *The Ruff*: T. & A.D. Poyser, London.
Vernon, G. G., & Woolley, D. M. (1999). Three-Dimensional Motion of Avian Spermatozoa. *Cell Motility and the Cytoskeleton,
42*, 149-161. doi:https://doi.org/10.1002/(sici)1097-0169(1999)42:2%3C149::aid-cm6%3E3.0.co;2-0
Vervoort, R., & Kempenaers, B. (2019). Variation in lek attendance and copulation success of Independent and Satellite male
Ruffs *Calidris pugnax*. *Ardea, 107*. doi:https://doi.org/10.5253/arde.v107i3.a9
Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*: Springer-Verlag New York.
Widemo, F. (1997). The social implications of traditional use of lek sites in the ruff *Philomachus pugnax*. *Behavioral Ecology,
8*(2), 211-217. doi:https://doi.org/10.1093/beheco/8.2.211
Widemo, F. (1998). Alternative reproductive strategies in the ruff, *Philomachus pugnax*: a mixed ESS? *Animal Behavior, 56*,
329-336. doi:https://doi.org/10.1006/anbe.1998.0792
Widemo, F., & Owens, I. (1995). Measuring sexual selection. *Nature, 376*, 471-472. doi:https://doi.org/10.1038/376471b0
Yeh, S. D., Chen, Y. J., Chang, A. C., Ray, R., She, B. R., Lee, W. S., . . . Lin-Chao, S. (2002). Isolation and properties of Gas8,
a growth arrest-specific gene regulated during male gametogenesis to produce a protein associated with the sperm
motility apparatus. *Journal of Biological Chemistry, 277*(8), 6311-6317. doi:https://doi.org/10.1074/jbc.M106941200
710

Acknowledgements

We thank the many people who took care of the captive ruffs, Melanie Schneider for genotyping, Callum
S. McDiarmid for help with his Sperm Sizer software, Esteban Botero-Delgadillo for the script on
homozygosity by locus, Jarrod Hadfield for advice on a genomic relationship matrix, Mihai Valcu for
advice on statistics, Yifan Pei for ruff illustrations, and two anonymous reviewers for the comments.

Author contributions

MB initiated the project with support of BK and DL. BK secured funding. MB, BK, TA, DL, CK, JL
conceptualized the project. JL collected and measured the testes. MB, KT, TA, ML collected the sperm
samples. JA measured velocity. KT and MB with support of TA optimized the photographing of the
sperm. MB, MC and KT developed and tested the sperm measuring protocol. KT and MC prepared

sperm samples, took sperm pictures and KT measured sperm morphology. MB curated and analyzed the data with input from BK, WF and TA. MB wrote the first draft and revised it with the help of BK. All authors commented and MB with BK, WF, CK and DL finalized the paper.

Funding

Max Planck Society (to BK and CK), the Czech Science Foundation project 19-22538S (to TA and JA).

Ethics declarations
Ethics approval and consent to participate
The data were collected under license (311.5-5682.1/1-2014-020) of the Landratsamt Starnberg and in accordance with Directive 2010/63/EU.

Consent for publication
Not applicable.

Competing interests
The authors have no competing interests.