TITLE: Genetic context drives age-related disparities in synaptic maintenance and

structure across cortical and hippocampal neuronal circuits.

Sarah E. Heuer[1,2], Emily W. Nickerson[1,2], Gareth R. Howell[1,2,3,*], Erik B. Bloss[1,2,3,*]

[1]The Jackson Laboratory, Bar Harbor, ME 04609, USA

[2]Tufts University Graduate School of Biomedical Sciences, Boston, MA 02111, USA

[3]Graduate School of Biomedical Sciences and Engineering, University of Maine, Orono, Maine

04469, USA

[*]Correspondence: erik.bloss@jax.org, gareth.howell@jax.org

25   **KEYWORDS**

26   Aging, dendritic spines, frontal cortex, neuronal circuits, genetic diversity

ABSTRACT

The disconnection of neuronal circuits through synaptic loss is presumed to be a major driver of age-related cognitive decline. Age-related cognitive decline is heterogeneous, yet whether genetic mechanisms differentiate successful from unsuccessful cognitive decline through synaptic structural mechanisms remains unknown. Previous work using rodent and primate models leveraged various techniques to suggest that age-related synaptic loss is widespread on pyramidal cells in prefrontal cortex (PFC) circuits but absent on those in area CA1 of the hippocampus. Here, we examined the effect of aging on synapses on projection neurons forming a hippocampal-cortico-thalamic circuit important for spatial working memory tasks from two genetically distinct mouse strains that exhibit susceptibility (C57BL/6J) or resistance (PWK/PhJ) to cognitive decline during aging. Across both strains, synapses on the CA1-to-PFC projection neurons appeared completely intact with age. In contrast, we found synapse loss on PFC-to-nucleus reuniens (RE) projection neurons from aged C57BL/6J but not PWK/PhJ mice. Moreover, synapses from aged PWK/PhJ mice but not from C57BL/6J exhibited morphological changes that suggest increased synaptic efficiency to depolarize the parent dendrite. Our findings suggest resistance to age-related cognitive decline results in part by age-related synaptic adaptations, and identification of these mechanisms in PWK/PhJ mice could uncover new therapeutic targets for promoting successful cognitive aging and extending human health span.

INTRODUCTION

Synapses are the fundamental computational subunits of the, and the building blocks of neuronal circuits that vary in complexity based on brain region and mammalian organism (1, 2). Synapses enable cell-type specific forms of communication and are often targeted to defined spatial subcellular locations on postsynaptic cell types (3), which constrains neural circuit operations (4, 5). Synapses act as plastic, tunable biochemical and electrical signaling compartments, a feature that allows internal states to shape neural circuit function (6, 7), and may be the primary substrate in the nervous system mediating learning and memory recall (8, 9).

Individual excitatory synapses on many neuronal cell types take the form of small protrusions along the dendrites (i.e., dendritic spines (10)), which can be visualized by multiple imaging methods (8, 11, 12). Spine structure reflects the functional strength of each connection: spine head size correlates with synapse strength (8, 12), and the morphology of the spine neck determines the degree of electrical and biochemical filtering that occurs between the synapse and the parent dendritic cable (13, 14). During normal aging, mammalian organisms experience progressive, yet variable, cognitive decline, even in the absence of a disease, which corresponds to a loss in synaptic plasticity(15). Relative to other synapses, spine synapses seem to be the primary type that are lost or altered during aging, and these changes in the aging brain appear circuit specific (16, 17). For example, the density of spine synapses decrease over time on cortical neurons in the frontal but not visual cortex, and decrease on CA3 pyramidal neurons but not on neighboring neurons in CA1 (16, 18, 19). The loss of synapse number and the changes in synaptic structure on frontal cortical neurons correlate with progressive decreases in cognitive function with age (20–22), yet reports of the degree to which synaptic loss occurs suggests significant heterogeneity (16, 23–25). The causal factors that underlie this variability could be what determines successful versus unsuccessful neural aging (16, 26, 27), and there is significant interest in understanding how genetic components and environmental influences coordinately shape the trajectory of age-related cognitive changes and synaptic connectivity.

Non-human primates (NHPs) (22, 28, 29) and rodents (rats (25, 30, 31) and mice (32))

are commonly used model systems to study the effect of aging on synapses. Although NHPs are

the most translatable model for relating age-related neural circuit changes to cognitive decline

(22, 33), they have long lifespans (>30 years), require complex housing systems, and are not

amendable to genetic-based experimentation at scale. However, to a large extent, data from

rodent models recapitulate the magnitude of age-related synaptic loss in vulnerable brain regions

(including the prefrontal cortex) that has been observed in NHPs (24, 25, 34). Studies in mice

have shown loss in synaptic plasticity and number during aging and in age-related diseases (35,

36). Unlike NHP experiments, which necessarily use genetically diverse non-inbred subjects, the

work in mice has largely been performed using the inbred C57BL/6J (B6) mouse strain. The lack

of genetic diversity limits translatability of mouse work to humans, and led us to evaluate the

potential of incorporating genetically distinct wild-derived inbred mouse strains in our own work

(37). Our previous results show wild-derived strains exhibit diverse cognitive and behavioral

phenotypes during aging and when carrying Alzheimer's disease (AD)-linked transgenes,

suggesting genetic background strongly influences neuronal circuitry and function (38, 39). In

particular, the PWK/PhJ (PWK) mouse strain did not show amyloid-dependent cognitive decline

or synaptic loss, suggesting PWK is an ideal strain to study resilience to AD (40). However, the

neuronal circuits vulnerable in aging are different than those vulnerable to AD, and it remains

unknown whether PWK is also resistant to normal age-related synaptic loss.

To determine the effect of genetic context on age-related synapse dynamics (and how

they might relate to their observed cognitive susceptibility/resilience (38, 41)), we examined aging-

related synapse loss in the traditionally studied B6 mouse strain and the wild-derived PWK. PWK

mice differ from B6 by approximately 20 million single nucleotide polymorphisms (SNPs) and

structural variants (37). To target our analyses to specific projection neurons, we leveraged

retrograde viral strategies to label two circuits known to have different vulnerabilities to aging or

AD: PFC-projecting neurons to nucleus reuniens (RE; PFC-to-RE), and CA1-projecting neurons

to PFC (CA1-to-PFC) (17). Spine synapses on CA1-to-PFC neurons showed marked stability

throughout aging in both B6 and PWK mice. In contrast, PFC-to-RE neurons were susceptible to

age-related spine loss in B6 but not PWK. Further, there were morphological changes suggestive

of synapse strengthening in PWK but not B6 mice. Collectively, these results provide evidence

that PWK are a model of successful synaptic aging.

RESULTS

Labeling strategies to visualize synapses from two neural circuits in genetically distinct

mice.

Our previous studies suggest PWK are a model of cognitive, neuronal and synaptic

resilience to Aβ pathology (38, 40). However, since age is the leading risk factor for developing

AD, we tested the prediction that PWK would also be a model of successful synaptic aging when

compared to traditionally used inbred B6 mice. We leveraged previously established viral

approaches (42, 43) using two different fluorophores to label excitatory projection neurons forming

two arms of a connected hippocampal-cortico-thalamic loop important for spatial working memory

tasks (44). Each set of projection neurons expressed either EGFP or tdTomato across cohorts of

female B6 and PWK mice ranging from 4-29 months of age (see **Table S1**). Three unilateral

intracranial injection sites were targeted to deliver recombinant adenoassociated virus (AAV, see

Methods) to drive EGFP in PFC neurons that project to RE (PFC-to-RE) and tdTomato

expression in CA1 neurons that project to PFC (CA1-to-PFC). After allowing 3-4 weeks for proper

retrograde movement and fluorophore expression each mouse was perfused, coronal slices were

made with a vibratome, slices containing EGFP+ or tdTomato+ neurons were imaged, and

dendritic spine density and morphologies were analyzed (**Figure 1A**). Three age groups were

assessed: young (4-7 months of age), middle-aged (11-17 months of age) and aged (22-30

months of age) groups. Neurons making up the two projection circuits in B6 and PWK mice were

labeled with equivalent efficiency, suggesting a similar overall density of neurons comprising

these pathways from the compared strains (**Figure 1B-C**).

Aging spines on proximal CA1 dendrites exhibit stable densities but altered morphologies

in PWK but not B6 mice

The CA1 region of the hippocampus is well-studied on account of its anatomy, physiology,

role in episodic memory, and vulnerability to both age- and AD-related pathological insults (27).

While generally thought to be relatively resistant to normal age-related synaptic loss (45–47), it is

unclear if age-related structural changes occur on specific CA1 projections, and if they are

modulated in subjects from genetic contexts that exhibit resistance to cognitive aging. The

majority of excitatory inputs onto CA1 pyramidal cells are made onto the proximal apical oblique

and basal branches (48, 49) which both receive inputs from hippocampal area CA3 (50).

Therefore, we first analyzed dendritic spine synapses on branches from this proximal

compartment specifically on CA1-to-PFC neurons (**Figure 2A**). Consistent with the prediction

from previous reports (17, 45), we observed no significant age-related changes in proximal spine

density in either B6 or PWK mice (**Figure 2B-C**).

Dendritic spines alter their morphologies in response to repeated artificial synaptic

stimulation (6, 9), under conditions of homeostasis (7, 51), and in reaction to disease pathologies

and during aging (52, 53). Spine head morphology is a reliable predictor of synaptic stability and

strength (8, 12), so we analyzed the maximum head diameter of each reconstructed spine. Our

analysis found no overt age-related changes in B6 mice, but the same analysis in PWK mice

revealed a subtle yet significant age-related shift to spines with smaller head diameters (**Figure**

2D). This effect was evident when we examined the densities of spines across the smallest (Q1)

and largest (Q4) quartiles of head diameters (**Figure S1B**).

The morphology of dendritic spine necks (e.g. neck width and length) are similarly plastic

(13) and define the compartmentalization of spines much more critically than spine head size (13,

14, 54, 55). Dendritic spine necks act as strong electrical resistors, filtering currents from the

synapse to the parent dendrite branch (54). Necks also contain receptors for neuromodulatory

systems that regulate local excitability (56). Spine necks are often below 50 nm (49) making them

hard to resolve with light microscopy techniques, thus we analyzed putative spine neck lengths

to determine whether this aspect of spine neck morphology is readily modified during aging. Neck

lengths in B6 mice were slightly lengthened with age, complementing the stability observed in

spine heads. In contrast, PWK spine neck lengths showed a biphasic response, first increasing

from young to middle-aged, then decreasing significantly in aged mice compared to the younger

groups (**Figure 2E**, **Figure S1C**).

Spine changes on distal CA1 dendrites mirror those found on proximal compartments

Distal tuft dendrites in CA1 pyramidal cells are thought to be the most vulnerable to AD-

related synaptic changes due to their afferent inputs originating in the EC (**Figure 3A**) (57), where

associated neuropathology is observed very early in the disease process (58, 59). Synapses on

distal dendrites are understudied relative to those on more proximal compartments due to

limitations in traditional techniques used to evaluate plasticity at remote locations relative to the

soma. Consistent with our results from the proximal dendrites, we observed no significant age-

related changes to distal tuft spine densities in B6 or PWK mice (**Figure 3B-C**). Like spines from

proximal branches, we observed few age-related changes in head diameter and neck length in

spines from B6 distal branches (**Figure 3D-E, Figure S2B-C**). However, spine head diameters

were larger in aged PWK mice compared to young and middle-aged cohorts (**Figure 3D**, **Figure

S2B**), and spine neck lengths were significantly shorter in aged compared to young and middle-

aged mice (**Figure 3E**, **Figure S2C**).

Collectively, these results show spine densities on proximal and distal dendrites from CA1-

to-PFC pyramidal cells do not change with age in both B6 and PWK mice. Furthermore, B6 spines

remained morphologically stable whereas PWK spines exhibited changes indicative of increased

excitability at the synapse and to the parent dendritic cable, which support the notion that these

adaptations could play a role in maintaining information processing in the aging hippocampus.

Differential vulnerability to age-related synaptic changes on PFC-to-RE neurons across B6

and PWK mice

Previous studies in humans, NHPs, rats and mice report that pyramidal neurons in the

prefrontal cortex are susceptible to synaptic loss with age (21, 22, 25, 29, 60). Moreover,

preventing this degeneration could be key to promoting successful cognitive aging (21, 61, 62).

To test the prediction that PWK mice are resistant to age-dependent synaptic loss in the PFC, we

used AAVs to also label deep-layer neurons (in the exact mice described above) in the PFC that

project to the RE (**Figure 1**) that appear to mediate behavioral flexibility and working memory

(63–65). Like the CA1-to-PFC neurons, PFC-to-RE neurons have a similar proximal-distal

dendritic architecture, and each compartment receives a distinct set of inputs. Basal and apical

oblique dendrites located in the proximal regions receive afferent inputs from CA1 (among other

regions), whereas distal tuft receive a distinct set of inputs from the thalamus (43) (**Figure 4A**).

Therefore, we analyzed the spines on proximal and distal dendrites separately.

We first examined spine density on proximal dendrites of PFC-to-RE neurons. Consistent

with previous findings, B6 proximal dendritic spine density was significantly lower in middle-aged

and aged compared to young mice (66). Interestingly, we found no differences in spine densities

on proximal PFC dendrites across age cohorts in PWK mice (**Figure 4B-C**). When we examined

spine morphologies, we found statistically significant changes in spine head diameter with age on

both B6 and PWK proximal PFC dendrites, evidenced by the cumulative frequency analyses and

by quartile analysis of the largest and smallest spines (**Figure 4D**, **Figure S3B**). In both B6 and

PWK mice, PFC proximal spine head diameters became subtly larger with aging (**Figure 4D**).

Interestingly, spines on proximal dendrites from B6 mice showed minimal age-related changes in

spine neck length. However, spines on proximal branches from PWK mice were significantly

shortened with age, changes that should reduce neck-related filtering of synaptic currents,

mimicking the same patterns found on aging CA1 dendrites from PWK mice (**Figure 4E**, **Figure**

S3C).

Age-related dendritic spine loss and morphological remodeling on distal PFC dendrites

differs between B6 and PWK mice

Distal tuft branches from PFC-to-RE projection neurons receive distinct afferent inputs

compared to proximal branches, coming from the thalamus instead of CA1 (43) (**Figure 5A**).

Similar to the age-related changes found on proximal dendrites of PFC-to-RE neurons, distal

dendritic spine densities were lower in middle-aged and aged compared to young B6 mice but did

not change across age groups in PWK mice (**Figure 5B-C**).

Spine head diameters on distal branches of PFC-to-RE neurons from B6 mice showed a

biphasic response, first increasing in size between young and middle-aged cohorts, then

decreasing in size between middle-aged to aged groups. PWK distal PFC spine heads displayed

the opposite pattern: decreasing in size between young and middle-aged groups, and then

increasing in size between middle-aged and aged cohorts (**Figure 5D, Figure S4B**). Spine neck

lengths on the distal dendrites of PFC-to-RE neurons from B6 mice slightly lengthened with age,

while those from PWK decreased more dramatically with age (**Figure 5E**). These changes in

spine neck lengths were observed in quartile analyses, with a significant gain of long spine necks

in aged B6 mice, and an overt loss of long spine necks in aged PWK mice (**Figure S4C**). These

results from distal PFC-to-RE dendrites mirrored those from proximal dendrites: B6 branches

showed age-related synaptic loss while those from PWK did not. In contrast, PWK distal PFC

spines showed spine neck remodeling while those from B6 were markedly more stable.

Collectively, these data support our hypothesis that, like resilience to AD-related Aβ

pathology (38), PWK mice show resistance to age-related cortical synapse loss. This resistance

was evident across both proximal and distal compartments and accompanied by specific forms

of spine head and neck remodeling. Decreases in spine neck length should serve to lower the

spine neck resistor and effectively increase the efficiency by which current generated at the

synapse reaches the parent dendrite. Given the growing links between dendritic mechanisms of

integration and the plasticity that mediates learning and memory (67–70), the putative adaptive

spine responses could serve as mechanisms to promote successful maintenance of neuronal

health and cognition during aging.

DISCUSSION

Here, we asked whether the effects of aging on synapse number and structure in two

specific neural circuits was equivalent across two distinct strains of mice. We chose to investigate

B6 and PWK strains as our past work has identified them as susceptible or resilient to AD-related

cognitive and synaptic deficits, respectively (38, 40, 41). To rigorously compare across defined

neural circuits, we used multisite viral injections to label CA1-to-PFC and PFC-to-RE neurons and

systematically sampled ~90,000 spines in proximal and distal dendritic compartments across

multiple ages (ranging from 5-30 months of age). We found that PFC-projecting CA1 pyramidal

cells from both strains maintain their dendritic spine synaptic densities with age, consistent with

a body of work examining randomly labeled neurons or synapses in aging NHPs, rats and B6

mice (16, 18, 27, 28, 45). Conversely, we observed a differential pattern of synapse loss on RE-

projecting PFC neurons between B6 and PWK mice, with synapse loss restricted to B6 but not

PWK mice. Spine reconstructions suggested that spines from B6 showed either stability or

patterns of remodeling putatively associated with decreased synaptic efficacy, whereas PWK

spines tended to show morphological changes consistent with increased synaptic efficacy during

aging (**Table S6**). These results suggest that previously observed cognitive characteristics from

these strains (38), including PWK resistance to aging-related decline in cognition or resilience to

AD-related neuropathology (38) and differential microglia activation (71) may be due in part to an

adaptive forms of synaptic plasticity that maintain connectivity and excitability in cortical circuitry.

The PWK mouse strain, then, represents an opportunity to identify and harness adaptive
mechanisms to lengthen cognitive resistance to the normal process of aging.

The high-resolution spine reconstructions performed here and viral-based strategies
allowed for specific populations of projection neurons to be compared across mice (42). Past work
has relied on disparate techniques, including strategies to label random neurons with dye-filling
methods or counting random synapses by EM, to build a coherent picture of synaptic vulnerability
of the aging brain (22, 25, 29, 35). By taking advantage of multisite viral strategies to gain
projection neuron specificity within the same aging subjects, our results are the beginning of
identifying the neural circuits that are vulnerable to age-related disconnection in terms of
information flow. Interplay between CA1 of the hippocampus, PFC and RE is thought to be critical
for spatial working memory tasks, a cognitive domain affected in aged individuals (44, 65, 72).
Since our most striking finding is the strain-dependent loss on aging PFC-to-RE neurons, our
results suggest the PFC may be the critical locus of aging in terms of synapse loss. Reductions
in PFC-to-RE spine densities in B6 mice was most dramatic from young to middle-aged, which is
consistent with previous findings of synapses on PFC neurons from rats (25, 30), as well as with
behavioral findings that animals and humans start to become impaired in working memory
function tasks as early as middle-age (31, 65, 73, 74). Although CA1 projection neurons appear
to maintain their numbers during aging, our results do not imply that CA1 function is intact during
aging. Other processes, including activity-dependent spine plasticity at CA1 synapses, might also
contribute to age-related deficits in hippocampal-dependent learning or memory function.

At the level of individual synapses, spine structure is a strong determinant of function (8,
12, 13). Specifically, spine head diameter is a correlate of synapse strength, and spine neck of
the level of compartmentalization between the synapse and parent dendrite. Past work has
focused on age-related modification of head diameter with little evidence for modulation of spine
neck morphology (62). In contrast, our data suggest that both spine head diameter and neck
length are readily modified during aging in a way that depends on genetic context. In B6 mice,

CA1-to-PFC neurons exhibited very few morphological spine changes with age, consistent with a

remarkable population stability during aging. Conversely, PWK mice showed consistent evidence

for morphological spine remodeling on CA1-to-PFC and PFC-to-RE neurons that are suggestive

of an overall increase in efficacy between the spine synapse and parent dendrite (**Table S6**).

These data are compelling because both spine head and spine neck changes move in unison

towards greater synapse efficiency, and their synergism might potentially drive large changes in

how excitatory postsynaptic potentials from multiple synapses are integrated in the dendrites.

Recent work from our group has found that PWK mice are resilient to AD-related synaptic

changes in CA1-to-PFC neurons during early exponential phases of pathology spread (40). As

biological age is the primary risk factor for developing neurodegenerative disorders including AD

(75, 76), the data presented here suggest that the PWK mouse strain can be both a model of

resilience to cognitive and synaptic changes seen in AD (38, 40), and resistance to cortical

synapse loss seen during normal brain aging. Since the molecular and cellular mechanisms that

differentiate successful from unsuccessful agers are still unknown (27), the findings here could

drive the identification of cell-specific processes that maintain synapses throughout aging. Future

work should seek to advance this goal by testing additional PFC projection neurons to understand

the circuit specificity of resistance, by molecular strategies to uncover the gene(s) required for this

resistance, and by using functional studies that interrogate how behavior relates to pattens of

neuronal activity in these circuits (64, 77). Ultimately, such work would set the stage for developing

precise therapeutic strategies to promote healthy cognitive aging and resilience to AD across our

genetically diverse and aging human population.

MATERIALS AND METHODS

Ethics Statement

All research was approved by the Institutional Animal Care and Use Committee (IACUC) at The

Jackson Laboratory (approval number 12005 and 20006). Animals were humanely euthanized

with 4% tribromoethanol (800 mg/kg). Authors performed their work following guidelines

established by "The Eight Edition of the Guide for the Care and Use of Laboratory Animals" and

euthanasia using methods approved by the American Veterinary Medical Association.

316   Animal husbandry

All mice were bred and housed in a 12/12 hour light/dark cycle on aspen bedding and fed a

standard 6% Purina 5K52 Chow diet. Experiments were performed using two mouse strains:

C57BL/6J (B6, JAX stock #000664) and PWK/PhJ (PWK, JAX stock #003715). Mice were group

housed for entirety of experiments. Experimental cohorts were generated through intercrossing

B6 or PWK mice to produce 3-8 female mice per age group. Full mouse information is reported

in **Table S1**.

324   Intracranial viral injections

Recombinant adenoassociated viral (AAV) vectors were used to drive Cre-recombinase

(AAVretro-Cre)(42), Cre-dependent EGFP (serotype 2/1, AAV-FLEX-*rev*-EGFP) (42, 43), Flp-

recombinase (AAVretro-CAG-FLPo), and FLP-dependent tdTomato (serotype 2/1 AAV-CAG-

FRT-*rev*-TdTomato) (78). AAVretro-CAG-FLPo and AAV-CAG-FRT-*rev*-TdTomato were gifts

from Janelia Viral Tools (Addgene plasmid #183412, http://n2t.net/addgene:183412,

RRID:Addgene_183412; Addgene plasmid #191203, http://n2t.net/addgene:191203,

RRID:Addgene_191203). The titers of each virus were as follows (in genomic copies/mL):

AAVretro-Cre, $1x10^{12}$, AAV-FLEX-*rev*-GFP, $1x10^{13}$, AAVretro-CAG-FLPo, $1x10^{13}$; AAV-CAG-

FRT-*rev*-TdTomato, $1x10^{13}$. A 1:1 ratio of AAVretro-CAG-FLPo and AAV-FLEX-*rev*-GFP was

mixed and 45nL injected into ventral prefrontal cortex (PFC) over 5 minutes; 45-50 nL (per each

D/V coordinate) of AAV-CAG-FRT-*rev*-TdTomato was injected in CA1 (CA1) over 10 minutes;

AAVretro-Cre was diluted 1:10 with sterile saline and 45 nL injected into nucleus reunions (RE)

over 5 minutes. Since PWK brain volumes are smaller than B6, injection coordinates were

adjusted based on pilot experiments to determine injection sites. The coordinates for each

injection were as follows (in mm: posterior relative to bregma, lateral relative to midline, and

ventral relative to pial surface): B6 PFC (+1.75, -0.95, and -2.6), B6 CA1 (-3.5, -3.4, and -2.7/-

2.5/2.0), B6 RE(-1.1, -1.2, -4.15); PWK PFC (+1.45, -0.9, and -2.3), PWK CA1 (-3.5, -3.3, and -

2.75/-2.5/-2.0), PWK RE(-1.0, -1.0, -3.2). RE and PFC injections were performed with the mouse

tilted at a 15° angle. At each site the injection pipette was left in place for 3-5 minutes then slowly

retracted at a rate of 10 μm/s from the brain. After surgery mice were singly housed, monitored

for 5 days, and euthanized approximately 3-4 weeks post injection.

Tissue harvest and brain sectioning

Mice were euthanized with an intraperitoneal lethal dose of Tribromoethanol (800 mg/kg),

followed by transcardial perfusion with 45 mL ice-cold 4% paraformaldehyde (PFA) in 0.1M

phosphate-buffered saline, in accordance with IACUC protocols (12005 and 20008). Brains were

removed and placed in 5mL ice cold 4% PFA at 4°C for 24 hours, then placed into storage buffer

(1XPBS + 0.1% Sodium Azide) for long-term storage at 4°C. Brains were coronally sectioned at

50μm thickness and placed in storage buffer at 4°C until needed for imaging. Approximately 6

frontal PFC sections containing EGFP+ dendrites, and 6 sections with distal CA1 TdTomato+

dendrites were mounted on slides and coverslipped using Vectashield Hardset mounting media

(Vector Laboratories #H140010).

Dendritic spine imaging and analysis

Slices were imaged on an SP8 confocal microscope equipped with a 63X objective (oil

immersion), images collected at 50 nm pixel sizes with 0.1 μm z-steps, and stacks deconvolved

using Leica Lightning software. 10 dendrites per compartment (e.g. basal, apical oblique, tuft)

were captured per mouse. Each image was exported as TIFF format (in ImageJ version

2.9.0/1.53t) and imported into NeuronStudio(79) for analysis of dendritic spine densities and

morphologies. Since the effectiveness of a synaptic input on the dendritic tree is influenced by

the diameter of the parent dendrite(80), we ensured all dendritic calibers incorporated into this

analysis were thin (i.e. under 1 μm in diameter) (**Figure S1A, S2A, S3A, S4A**). Density

measurements were acquired by first reconstructing the dendritic cable followed by semi-

automated spine identification. Cumulative distributions of assigned spine head diameters

(HEAD.DIAMETER) and neck lengths (MAX.DTS) were analyzed by Kolmogorov-Smirnov tests,

and through a quartile-based analysis. In this latter analysis, spines within dendritic compartments

from each strain were pooled across treatment groups to create a population, and the first and

last quartiles determined. From each branch, spines belonging to the first quartile (Q1, smallest)

and last quartile (Q4, largest) were identified and the density for each quartile per branch was

calculated. Data were analyzed with each dendrite representing an individual data point.

Statistical analysis

Data were analyzed blinded to strain and age group. All statistical analyses were performed in

GraphPad Prism software (v9.5.1) except for Kolmogorov-Smirnov tests which were performed

using R (v4.2.2). Results are reported in table form in the Supplemental Information **Table S2-**

S5). Data from B6 and PWK mouse strains were analyzed separately. To assess age effects

within each strain, one-way ANOVAs were computed followed by Bonferroni post hoc tests.

Within-group differences between Q1 and Q4 densities for spine head diameter and neck length

were determined using nonparametric two-tailed t-tests. Bonferroni corrections for multiple

comparisons were performed on nominal p-values from resultant K-S tests.

Data and resource sharing plan

All mouse strains are available through The Jackson Laboratory. All reagents in this study are

commercially available. Raw data (DOI: 10.6084/m9.figshare.23620752) and images from the

figures (DOI: 10.6084/m9.figshare.23620755) are available via Figshare (made public upon

acceptance of publication).

Author contributions

S.E.H., E.B.B. and G.R.H. designed the study. S.E.H. generated and maintained mouse cohorts,

and S.E.H. and E.B.B. performed the intracranial AAV injections. S.E.H. euthanized mice and

collected tissues. S.E.H. and E.W.N. sectioned brains and prepared slides for imaging. S.E.H.

and E.W.N. imaged slices and analyzed final data. E.B.B. provided training in intracranial

injections, confocal microscopy imaging of dendrites and dendritic spines, and NeuronStudio

analysis. E.B.B. and G.R.H. advised on all data analysis, data interpretation and manuscript

preparation. S.E.H., E.W.N., E.B.B. and G.R.H. wrote and edited the manuscript. All authors

approved the final version.

Competing Interest statement

The authors have no conflicts or competing interests to declare.

ACKNOWLEDGEMENTS AND FUNDING SOURCES

Acknowledgements

We thank Dr. Kristen Onos and Kelly Keezer for providing aged PWK/PhJ mouse cohorts, and

Amanda Hewes and Melanie-Maddox Goodrich for providing critical laboratory support for the

initiation of these experiments. We also thank Dr. Philipp Henrich at the Microscopy core at The

Jackson Laboratory for training and assistance on the confocal microscope.

Funding Sources

This study was supported by the National Institute on Aging (NIA) AG079877 (E.B.B.), AG055104

(G.R.H.), program funds from The Jackson Laboratory (G.R.H.), and by the NIA T32 training

program AG062409 in the Precision Genetics of Aging, Alzheimer's Disease and Related Dementias at The Jackson Laboratory (S.E.H., G.R.H., E.B.B.).

REFERENCES

1. T. N. Lerner, L. Ye, K. Deisseroth, Communication in Neural Circuits: Tools, Opportunities, and Challenges. *Cell* **164**, 1136–1150 (2016).

2. T. C. Südhof, R. C. Malenka, Understanding Synapses: Past, Present, and Future. *Neuron* **60**, 469–476 (2008).

3. M. Sheng, E. Kim, The Postsynaptic Organization of Synapses. *Cold Spring Harb. Perspect. Biol.* **3**, a005678 (2011).

4. R. R. LLINÁS, K. D. WALTON, E. J. LANG, "Cerebellum" in *The Synaptic Organization of the Brain*, G. M. Shepherd, Ed. (Oxford University Press, 2004), p. 0.

5. I. Ballesteros-Yáñez, R. Benavides-Piccione, G. N. Elston, R. Yuste, J. DeFelipe, Density and morphology of dendritic spines in mouse neocortex. *Neuroscience* **138**, 403–409 (2006).

6. M. Matsuzaki, N. Honkura, G. C. R. Ellis-Davies, H. Kasai, Structural basis of long-term potentiation in single dendritic spines. *Nature* **429**, 761–766 (2004).

7. G. G. Turrigiano, The self-tuning neuron: synaptic scaling of excitatory synapses. *Cell* **135**, 422–435 (2008).

8. H. Kasai, M. Matsuzaki, J. Noguchi, N. Yasumatsu, H. Nakahara, Structure–stability–function relationships of dendritic spines. *Trends Neurosci.* **26**, 360–368 (2003).

9. C. D. Harvey, K. Svoboda, Locally dynamic synaptic learning rules in pyramidal neuron dendrites. *Nature* **450**, 1195–1200 (2007).

10. S. R. y Cajal, *Estructura de los centros nerviosos de las aves* (1888).

11. M. E. Serrano, E. Kim, M. M. Petrinovic, F. Turkheimer, D. Cash, Imaging Synaptic Density: The Next Holy Grail of Neuroscience? *Front. Neurosci.* **16**, 796129 (2022).

12. M. Matsuzaki, *et al.*, Dendritic spine geometry is critical for AMPA receptor expression in hippocampal CA1 pyramidal neurons. *Nat. Neurosci.* **4**, 1086–1092 (2001).

13. J. Tønnesen, G. Katona, B. Rózsa, U. V. Nägerl, Spine neck plasticity regulates compartmentalization of synapses. *Nat. Neurosci.* **17**, 678–685 (2014).

14.   Å. Grunditz, N. Holbro, L. Tian, Y. Zuo, T. G. Oertner, Spine Neck Plasticity Controls
Postsynaptic Calcium Signals through Electrical Compartmentalization. *J. Neurosci.* **28**,
13457–13466 (2008).

15.   S. N. Burke, C. A. Barnes, Neural plasticity in the ageing brain. *Nat. Rev. Neurosci.* **7**, 30–40
(2006).

16.   J. H. Morrison, M. G. Baxter, The ageing cortical synapse: hallmarks and implications for
cognitive decline. *Nat. Rev. Neurosci.* **13**, 240–250 (2012).

17.   D. L. Dickstein, C. M. Weaver, J. I. Luebke, P. R. Hof, Dendritic spine changes associated
with normal aging. *Neuroscience* **251**, 21–32 (2013).

18.   X. Fan, E. G. Wheatley, S. A. Villeda, Mechanisms of Hippocampal Aging and the Potential
for Rejuvenation. *Annu. Rev. Neurosci.* **40**, 251–272 (2017).

19.   M. E. Young, D. T. Ohm, D. Dumitriu, P. R. Rapp, J. H. Morrison, Differential effects of aging
on dendritic spines in visual cortex and prefrontal cortex of the rhesus monkey.
Neuroscience **274**, 33–43 (2014).

20.   Y. Hara, P. R. Rapp, J. H. Morrison, Neuronal and morphological bases of cognitive decline
in aged rhesus monkeys. *Age Dordr. Neth.* **34**, 1051–1073 (2012).

21.   A. R. Brennan, *et al.*, Protein kinase C activity is associated with prefrontal cortical decline
in aging. *Neurobiol. Aging* **30**, 782–792 (2009).

22.   J. I. Luebke, Y.-M. Chang, T. L. Moore, D. L. Rosene, Normal aging results in decreased
synaptic excitation and increased synaptic inhibition of layer 2/3 pyramidal cells in the
monkey prefrontal cortex. *Neuroscience* **125**, 277–288 (2004).

23.   S. W. Scheff, D. A. Price, D. L. Sparks, Quantitative assessment of possible age-related
change in synaptic numbers in the human frontal cortex. *Neurobiol. Aging* **22**, 355–365
(2001).

24.   S. W. Scheff, S. A. Scott, S. T. DeKosky, Quantitation of synaptic density in the septal nuclei
of young and aged Fischer 344 rats. *Neurobiol. Aging* **12**, 3–12 (1991).

25.   E. B. Bloss, *et al.*, Evidence for Reduced Experience-Dependent Dendritic Spine Plasticity in
the Aging Prefrontal Cortex. *J. Neurosci.* **31**, 7831–7839 (2011).

26.   M. B. Moss, T. L. Moore, S. P. Schettler, R. Killiany, D. Rosene, "Successful vs. Unsuccessful
Aging in the Rhesus Monkey" in *Brain Aging: Models, Methods, and Mechanisms*, Frontiers
in Neuroscience., D. R. Riddle, Ed. (CRC Press/Taylor & Francis, 2007) (June 27, 2023).

27.   J. H. Morrison, P. R. Hof, Life and death of neurons in the aging brain. *Science* **278**, 412–
419 (1997).

28.   P. R. Hof, J. H. Morrison, The aging brain: morphomolecular senescence of cortical circuits.
478   *Trends Neurosci.* **27**, 607–613 (2004).

479   29.   A. PETERS, C. SETHARES, J. I. LUEBKE, SYNAPSES ARE LOST DURING AGING IN THE PRIMATE
480   PREFRONTAL CORTEX. *Neuroscience* **152**, 970–981 (2008).

481   30.   E. B. Bloss, *et al.*, Morphological and molecular changes in aging rat prelimbic prefrontal
482   cortical synapses. *Neurobiol. Aging* **34**, 200–210 (2013).

483   31.   C. A. Barnes, B. L. McNaughton, An age comparison of the rates of acquisition and
484   forgetting of spatial information in relation to long-term enhancement of hippocampal
485   synapses. *Behav. Neurosci.* **99**, 1040–1048 (1985).

486   32.   M. Cizeron, *et al.*, A brainwide atlas of synapses across the mouse life span. *Science* **369**,
487   270–275 (2020).

488   33.   D. Dumitriu, *et al.*, Selective Changes in Thin Spine Density and Morphology in Monkey
489   Prefrontal Cortex Correlate with Aging-Related Cognitive Impairment. *J. Neurosci.* **30**,
490   7507–7515 (2010).

491   34.   T. Rasmussen, T. Schliemann, J. C. Sørensen, J. Zimmer, M. J. West, Memory impaired aged
492   rats: no loss of principal hippocampal and subicular neurons. *Neurobiol. Aging* **17**, 143–
493   147 (1996).

494   35.   A. R. Ouellette, *et al.*, Cross-Species Analyses Identify Dlgap2 as a Regulator of Age-Related
495   Cognitive Decline and Alzheimer's Dementia. *Cell Rep.* **32**, 108091 (2020).

496   36.   C. L. Ackert-Bicknell, *et al.*, Aging Research Using Mouse Models. *Curr. Protoc. Mouse Biol.*
497   **5**, 95–133 (2015).

498   37.   T. M. Keane, *et al.*, Mouse genomic variation and its effect on phenotypes and gene
499   regulation. *Nature* **477**, 289–294 (2011).

500   38.   K. D. Onos, *et al.*, Enhancing face validity of mouse models of Alzheimer's disease with
501   natural genetic variation. *PLoS Genet.* **15**, e1008155 (2019).

502   39.   C. Fernandes, *et al.*, Behavioral Characterization of Wild Derived Male Mice (Mus musculus
503   musculus) of the PWD/Ph Inbred Strain: High Exploration Compared to C57BL/6J. *Behav.*
504   *Genet.* **34**, 621–630 (2004).

505   40.   S. E. Heuer, *et al.*, Genetic context controls early microglia-synaptic interactions in mouse
506   models of Alzheimer's disease. 2023.04.28.538728 (2023).

507   41.   S. Gregorová, J. Forejt, PWD/Ph and PWK/Ph inbred mouse strains of Mus m. musculus
508   subspecies--a valuable resource of phenotypic variations and genomic polymorphisms.
509   *Folia Biol. (Praha)* **46**, 31–41 (2000).

510   42.   D. G. R. Tervo, *et al.*, A Designer AAV Variant Permits Efficient Retrograde Access to
511   Projection Neurons. *Neuron* **92**, 372–382 (2016).

512   43.   K. Graham, N. Spruston, E. B. Bloss, Hippocampal and thalamic afferents form distinct
513   synaptic microcircuits in the mouse infralimbic frontal cortex. *Cell Rep.* **37**, 109837 (2021).

514   44.   H. T. Ito, S.-J. Zhang, M. P. Witter, E. I. Moser, M.-B. Moser, A prefrontal-thalamo-
515   hippocampal circuit for goal-directed spatial navigation. *Nature* **522**, 50–55 (2015).

516   45.   E. W. Buss, *et al.*, Cognitive aging is associated with redistribution of synaptic weights in
517   the hippocampus. *Proc. Natl. Acad. Sci.* **118**, e1921481118 (2021).

518   46.   D. A. Nicholson, R. Yoshida, R. W. Berry, M. Gallagher, Y. Geinisman, Reduction in size of
519   perforated postsynaptic densities in hippocampal axospinous synapses and age-related
520   spatial learning impairments. *J. Neurosci. Off. J. Soc. Neurosci.* **24**, 7648–7653 (2004).

521   47.   Y. Geinisman, *et al.*, Aging, spatial learning, and total synapse number in the rat CA1
522   stratum radiatum. *Neurobiol. Aging* **25**, 407–416 (2004).

523   48.   M. Megías, Z. Emri, T. F. Freund, A. I. Gulyás, Total number and distribution of inhibitory
524   and excitatory synapses on hippocampal CA1 pyramidal cells. *Neuroscience* **102**, 527–540
525   (2001).

526   49.   E. B. Bloss, *et al.*, Single excitatory axons form clustered synapses onto CA1 pyramidal cell
527   dendrites. *Nat. Neurosci.* **21**, 353–363 (2018).

528   50.   R. M. Empson, U. Heinemann, The perforant path projection to hippocampal area CA1 in
529   the rat hippocampal-entorhinal cortex combined slice. *J. Physiol.* **484**, 707–720 (1995).

530   51.   T. A. Blanpied, M. D. Ehlers, Microanatomy of dendritic spines: emerging principles of
531   synaptic pathology in psychiatric and neurological disease. *Biol. Psychiatry* **55**, 1121–1127
532   (2004).

533   52.   K. Runge, C. Cardoso, A. de Chevigny, Dendritic Spine Plasticity: Function and Mechanisms.
534   *Front. Synaptic Neurosci.* **12** (2020).

535   53.   G. Turrigiano, Homeostatic synaptic plasticity: local and global mechanisms for stabilizing
536   neuronal function. *Cold Spring Harb. Perspect. Biol.* **4**, a005736 (2012).

537   54.   R. Araya, J. Jiang, K. B. Eisenthal, R. Yuste, The spine neck filters membrane potentials.
538   *Proc. Natl. Acad. Sci.* **103**, 17961–17966 (2006).

539   55.   K. E. Sorra, K. M. Harris, Overview on the structure, composition, function, development,
540   and plasticity of hippocampal dendritic spines. *Hippocampus* **10**, 501–511 (2000).

56. M. Wang, *et al.*, Alpha2A-adrenoceptors strengthen working memory networks by
inhibiting cAMP-HCN channel signaling in prefrontal cortex. *Cell* **129**, 397–410 (2007).

57. K. M. Neuman, *et al.*, Evidence for Alzheimer's disease-linked synapse loss and
compensation in mouse and human hippocampal CA1 pyramidal neurons. *Brain Struct.*
Funct. **220**, 3143–3165 (2015).

58. S. K. Kaufman, K. Del Tredici, T. L. Thomas, H. Braak, M. I. Diamond, Tau seeding activity
begins in the transentorhinal/entorhinal regions and anticipates phospho-tau pathology in
Alzheimer's disease and PART. *Acta Neuropathol. (Berl.)* **136**, 57–67 (2018).

59. H. Hampel, *et al.*, The Amyloid-β Pathway in Alzheimer's Disease. *Mol. Psychiatry* **26**,
5481–5503 (2021).

60. Z. Petanjek, *et al.*, Extraordinary neoteny of synaptic spines in the human prefrontal
cortex. *Proc. Natl. Acad. Sci.* **108**, 13281–13286 (2011).

61. S. A. Villeda, *et al.*, Young blood reverses age-related impairments in cognitive function
and synaptic plasticity in mice. *Nat. Med.* **20**, 659–663 (2014).

62. J. Hao, *et al.*, Interactive effects of age and estrogen on cognition and pyramidal neurons
in monkey prefrontal cortex. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 11465–11470 (2007).

63. H. Bayer, L. J. Bertoglio, Infralimbic cortex controls fear memory generalization and
susceptibility to extinction during consolidation. *Sci. Rep.* **10**, 15827 (2020).

64. C. C. Kaczorowski, S. J. Davis, J. R. Moyer, Aging redistributes medial prefrontal neuronal
excitability and impedes extinction of trace fear conditioning. *Neurobiol. Aging* **33**, 1744–
1757 (2012).

65. J. L. Bizon, *et al.*, Spatial reference and working memory across the lifespan of male Fischer
344 rats. *Neurobiol. Aging* **30**, 646–655 (2009).

66. A. Shimada, H. Keino, M. Satoh, M. Kishikawa, M. Hosokawa, Age-related loss of synapses
in the frontal cortex of SAMP10 mouse: A model of cerebral degeneration. *Synapse* **48**,
198–204 (2003).

67. N. Spruston, Pyramidal neurons: dendritic structure and synaptic integration. *Nat. Rev.*
Neurosci. **9**, 206–221 (2008).

68. T. Branco, M. Häusser, Synaptic integration gradients in single cortical pyramidal cell
dendrites. *Neuron* **69**, 885–892 (2011).

69. J. C. Magee, Dendritic integration of excitatory synaptic input. *Nat. Rev. Neurosci.* **1**, 181–
190 (2000).

70. M. E. Larkum, T. Nevian, M. Sandler, A. Polsky, J. Schiller, Synaptic integration in tuft
dendrites of layer 5 pyramidal neurons: a new unifying principle. *Science* **325**, 756–760
(2009).

71. H. S. Yang, *et al.*, Natural genetic variation determines microglia heterogeneity in wild-
derived mouse models of Alzheimer's disease. *Cell Rep.* **34** (2021).

72. C. Pliatsikas, *et al.*, Working memory in older adults declines with age, but is modulated by
sex and education. *Q. J. Exp. Psychol. 2006* **72**, 1308–1327 (2019).

73. T. L. Moore, R. J. Killiany, J. G. Herndon, D. L. Rosene, M. B. Moss, Executive system
dysfunction occurs as early as middle-age in the rhesus monkey. *Neurobiol. Aging* **27**,
1484–1493 (2006).

74. T. A. Salthouse, When does age-related cognitive decline begin? *Neurobiol. Aging* **30**, 507–
514 (2009).

75. L. J. Launer, *et al.*, Rates and risk factors for dementia and Alzheimer's disease: results
from EURODEM pooled analyses. EURODEM Incidence Research Group and Work Groups.
European Studies of Dementia. *Neurology* **52**, 78–84 (1999).

76. R. A Armstrong, Risk factors for Alzheimer's disease. *Folia Neuropathol.* **57**, 87–105 (2019).

77. W. Göbel, F. Helmchen, In Vivo Calcium Imaging of Neural Network Function. *Physiology*
22, 358–365 (2007).

78. J. Winnubst, *et al.*, Reconstruction of 1,000 Projection Neurons Reveals New Cell Types
and Organization of Long-Range Connectivity in the Mouse Brain. *Cell* **179**, 268-281.e13
(2019).

79. A. Rodriguez, D. B. Ehlenberger, D. L. Dickstein, P. R. Hof, S. L. Wearne, Automated Three-
Dimensional Detection and Shape Classification of Dendritic Spines from Fluorescence
Microscopy Images. *PLoS ONE* **3**, e1997 (2008).

80. W. R. Holmes, The role of dendritic diameters in maximizing the effectiveness of synaptic
inputs. *Brain Res.* **478**, 127–137 (1989).

MAIN FIGURES

Figure 1: Viral labeling of PFC-to-RE and CA1-to-PFC neurons across C57BL/6J and PWK/PhJ mice.

(A) Experimental outline (see *Methods* for additional details)

(B-C) Example 10X images of tdTomato+ CA1 neurons (B) from B6 (left) and PWK (right) demonstrating consistent CA1-to-PFC projection circuit labeling, and EGFP+ PFC neurons (C) from B6 (left) and PWK (right) demonstrating consistent PFC-to-RE projection circuit labeling. Summary metadata for individual mice used in this study reported in **Table S1**.

Figure 2: CA1-to-PFC proximal dendrites are resistant to synaptic density changes across both strains, but PWK/PhJ spines change morphologies.

(A) Example schematic of tdTomato+ proximal CA1-to-PFC dendrites analyzed here and projection input origins in hippocampal area CA3.

(B) Proximal CA1-to-PFC dendrite spine densities (spines/μm) comparing young (Y), middle (M) and aged (A) within each strain. Data points represent individual branches (n=20/mouse); error bars are ± SD; one-way ANOVA identified no significant effects within B6 or PWK (**Table S2**).

(C) Representative 63X images of proximal CA1-to-PFC dendrites from each strain/age group.

(D) CA1-to-PFC proximal spine head diameter example image (left) and cumulative distributions from B6 (middle) and PWK (right), across age groups. Kolmogorov-Smirnov tests were used to evaluate statistical significance (Bonferroni adj. $p<0.05$) of pairwise comparisons: young vs middle (YM), middle vs aged (MA), and young vs. aged (YA), and p-values reported on each graph (see **Table S2**). Data points are representative of measures from individual spines.

(E) Same as **(D)** for CA1-to-PFC proximal spine neck length.

Summary statistics for data points represented in each graph reported in **Table S2.**

Figure 3: CA1-to-PFC distal tuft dendrites do not lose synapses but are morphologically altered in PWK/PhJ mice.

(A) Example schematic of tdTomato+ distal tuft CA1-to-PFC dendrites analyzed here and projection input origins in entorhinal cortex (EC).

(B) Distal tuft CA1-to-PFC dendrite spine densities (spines/μm) comparing young (Y), middle (M) and aged (A) within each strain. Data points represent individual branches (n=10/mouse); error bars are ± SD; one-way ANOVA identified no significant effects within B6 or PWK (**Table S3**).

(C) Representative 63X images of distal tuft CA1-to-PFC dendrites from each strain/age group.

(D) CA1-to-PFC distal tuft spine head diameter example image (left) and cumulative distributions from B6 (middle) and PWK (right), across age groups. Kolmogorov-Smirnov tests were used to evaluate statistical significance (Bonferroni adj. p<0.05) of pairwise comparisons: young vs middle (YM), middle vs aged (MA), and young vs. aged (YA), and p-values reported on each graph (see **Table S3**). Data points are representative of measures from individual spines.

(E) Same as **(D)** for CA1-to-PFC distal tuft spine neck length.

Summary statistics for data points represented in each graph reported in **Table S3**.

Figure 4: C57BL/6J mice are susceptible to age-related synaptic loss in proximal PFC-to-RE dendrites, with PWK/PhJ demonstrating dynamic morphological changes to increase synaptic strength.

(A) Example schematic of EGFP+ proximal PFC-to-RE dendrites analyzed here and projection input origins in hippocampal area CA1.

(B) Proximal PFC-to-RE dendrite spine densities (spines/μm) comparing young (Y), middle-aged (M) and aged (A) within each strain. Data points represent individual branches (n=20/mouse); error bars are ± SD; one-way ANOVA identified significant effect within B6 but not PWK; asterisks denote significant (Bonferroni adjusted p<0.05) comparisons identified using post-hoc analysis within B6 (**Table S4**).

(C) Representative 63X images of proximal PFC-to-RE dendrites from each strain/age group.

(D) PFC-to-RE proximal spine head diameter example image (left) and cumulative distributions from B6 (middle) and PWK (right), across age groups. Kolmogorov-Smirnov tests were used to evaluate statistical significance (Bonferroni adj. p<0.05) of pairwise comparisons: young vs middle

661   (YM), middle vs aged (MA), and young vs. aged (YA), and p-values reported on each graph (see

662   **Table S4**). Data points are representative of measures from individual spines.

663   **(E)** Same as **(D)** for PFC-to-RE proximal spine neck length.

664   Summary statistics for data points represented in each graph reported in **Table S4.**

Figure 5: PFC-to-RE distal tuft dendrites exhibit age-induced synaptic loss in C57BL/6J

mice but are morphology dynamic in PWK/PhJ mice.

(A) Example schematic of EGFP+ distal tuft PFC-to-RE dendrites analyzed here and thalamic

projection input origins.

(B) Distal tuft PFC-to-RE dendrite spine densities (spines/μm) comparing young (Y), middle (M)

and aged (A) within each strain. Data points represent individual branches (n=10/mouse); error

bars are ± SD; one-way ANOVA identified significant effect within B6 but not PWK; asterisks

denote significant (Bonferroni adjusted p<0.05) comparisons identified using post-hoc analysis

within B6 (**Table S5**).

(C) Representative 63X images of distal tuft PFC-to-RE dendrites from each strain/age group.

(D) PFC-to-RE distal tuft spine head diameter example image (left) and cumulative distributions

from B6 (middle) and PWK (right), across age groups. Kolmogorov-Smirnov tests were used to

evaluate statistical significance (Bonferroni adj. p<0.05) of pairwise comparisons: young vs middle

(YM), middle vs aged (MA), and young vs. aged (YA), and p-values reported on each graph (see

Table S5). Data points are representative of measures from individual spines.

(E) Same as **(D)** for PFC-to-RE distal tuft spine neck length.

Summary statistics for data points represented in each graph reported in **Table S5.**