Title: Chemogenetic manipulation of endogenous proteins in fission yeast using a self-

localizing ligand-induced protein translocation (SLIPT) system

Authors:

Akinobu Nakamura[1], Yuhei Goto[1,2,3,§], Hironori Sugiyama[1], Shinya Tsukiji[4,5], and Kazuhiro Aoki[1,2,3,§]

Affiliations:

[1]Quantitative Biology Research Group, Exploratory Research Center on Life and Living Systems

(ExCELLS), National Institutes of Natural Sciences, 5-1 Higashiyama, Myodaiji-cho, Okazaki, Aichi

444-8787, Japan.

[2]Division of Quantitative Biology, National Institute for Basic Biology, National Institutes of Natural

Sciences, 5-1 Higashiyama, Myodaiji-cho, Okazaki, Aichi 444-8787, Japan.

[3]Department of Basic Biology, School of Life Science, SOKENDAI (The Graduate University for

Advanced Studies), 5-1 Higashiyama, Myodaiji-cho, Okazaki, Aichi 444-8787, Japan.

[4]Department of Nanopharmaceutical Sciences, Graduate School of Engineering, Nagoya Institute of

Technology, Gokiso-cho, Showa-ku, Nagoya 466-8555, Japan.

[5]Department of Life Science and Applied Chemistry, Nagoya Institute of Technology, Gokiso-cho,

Showa-ku, Nagoya 466-8555, Japan

§Author for correspondence (y-goto@nibb.ac.jp; k-aoki@nibb.ac.jp)

Abstract

Cells sense extracellular stimuli through membrane receptors and process the information through an intracellular signaling network. Protein translocation triggers intracellular signaling, and techniques such as chemically induced dimerization (CID) have been used to manipulate signaling pathways by altering the subcellular localization of signaling molecules. However, in the fission yeast *Schizosaccharomyces pombe*, the commonly used FKBP-FRB system has technical limitations, and therefore perturbation tools with low cytotoxicity and high temporal resolution are needed. We here applied our recently developed self-localizing ligand-induced protein translocation (SLIPT) system to *S. pombe* and successfully perturbed several cell cycle-related proteins. The SLIPT system utilizes self-localizing ligands to recruit binding partners to specific subcellular compartments, such as the plasma membrane or nucleus. We optimzed the self-localizing ligands to maintain long-term recruitment of target molecules to the plasma membrane. By knocking in genes encoding the binding partners for self-localizing ligands, we observed changes in the localization of several endogenous molecules and found perturbations in the cell cycle and associated phenotypes. This study demonstrates the effectiveness of the SLIPT system as a chemogenetic tool for rapid perturbation of endogenous molecules in *S. pombe*, providing a valuable approach for studying intracellular signaling and cell cycle regulation with improved temporal resolution.

Keywords. fission yeast, chemogenetics, protein translocation, DHFR.

Introduction

Cells sense extracellular stimuli through their receptors on the plasma membrane (PM) and process the information through an intracellular signaling network [1]. The intracellular signaling network consists of multiple biophysical diffusions and reactions, including protein-protein interactions and enzymatic reactions. It is essential to understand the mechanisms of intracellular signaling because several diseases, such as malignant tumors, arise from genetic dysfunction of intracellular signaling molecules [2,3]. Although gene knockout has been widely used to study the physiological function of signaling proteins, it is difficult to analyze the gene when the gene knockout causes lethality. Furthermore, because the effects of gene knockout are long-lasting, the changes caused by gene knockout may be masked by adaptations in the signal transduction system [4,5]. Therefore, tools are needed to rapidly control the activity of signaling proteins with high temporal resolution.

Protein translocation is known to trigger intracellular signaling. For example, Akt, a Ser/Thr protein kinase, is recruited to and activated at the PM by binding to phosphatidyl inositol trisphosphate PI(3,4,5)P3, leading to the initiation of downstream signaling [6]. In addition, protein translocation restricts signaling to specific subcellular compartments. Upon growth factor stimulation, ERK, a classical MAP kinase, is phosphorylated and activated in the cytoplasm, followed by nuclear translocation to phosphorylate nuclear substrates such as transcription factors [7]. Several techniques have been reported to rapidly control signaling pathways by altering the subcellular localization of intracellular signaling molecules. Among them, chemically induced dimerization (CID) has been widely used to manipulate various signaling pathways, such as the Ras-ERK MAP kinase pathway, the PI3K-Akt pathway, and the Rho family G protein pathway [8–12]. CID is a chemical process in which two proteins bind only in the presence of small chemical dimerizers [13], and in many cases the signaling system is manipulated by inducing heterodimerization of the proteins. The most widely used CID system is the heterodimerization of FKBP-FRB by rapamycin [8,14]. However, chemically induced heterodimerization requires the expression of two types of proteins, namely, a localizer and an actuator. Recently, we have developed a self-localizing ligand-induced protein translocation (SLIPT) system, which requires only an actuator and a small compound with a subcellular localization tag (self-localizing ligand). By using the SLIPT system, we have successfully altered the subcellular localization of actuators and manipulated the Grb2/SOS/Ras/ERK, PI3K, Tiam/Rac, Gqα/Ca^{2+}, Gsα/cAMP, and PKCδ signaling pathways in mammalian cells [15–22] (Figure 1A).

The fission yeast *Schizosaccharomyces pombe* is a widely used eukaryotic model organism.

Molecular genetic analysis of *S. pombe* has allowed the identification of genes that regulate cellular

processes such as the cell cycle [23]. However, analysis using gene knockout mutants can only yield

results with low temporal resolution, in addition to the difficulty of studying lethal genes. Therefore, it

remains a technical challenge to precisely analyze when and how cell cycle-related proteins function in

the cell cycle. Although the CID method is expected to provide a way to rapidly perturb intracellular

signaling, the FKBP-FRB system in fission yeast has some technical problems, such as the inhibition of

sexual development by TOR inhibition with rapamycin [24–27]. In addition, to use the FKBP-FRB system

in fission yeast, the *fkh1* gene must be knocked out due to the high expression levels of endogenous

Fkh1 (the FKBP12 homolog) [24]. For these reasons, there is a need for perturbation tools to control cell

signaling with low cytotoxicity and high temporal resolution in fission yeast.

In this study, we applied the SLIPT system to *S. pombe* and successfully demonstrated rapid

perturbation of several cell cycle-related proteins. The self-localizing ligands recruited their binding

partners to the PM or nucleus in fission yeast. We further optimized the self-localizing ligands for *S.*

pombe to maintain recruitment of target molecules to the PM over a long period of time. In addition,

we observed the localization changes of dozens of endogenous molecules by knocking in a gene of the

binding partner for self-localizing ligands, along with analysis of the associated cell-cycle perturbations

and phenotypes. Taken together, we demonstrate for the first time that the SLIPT system serves as a

useful chemogenetic tool for rapid perturbation of endogenous molecules in *S. pombe*.

Results and Discussion

Introduction of the SLIPT system to fission yeast

In this study, we sought to develop a functional SLIPT system optimized for the fission yeast *S. pombe*. We have previously reported a self-localizing ligand, m^DcTMP, which is composed of myristoylated D-cysteine (m^Dc) and trimethoprim (TMP) [17] (Figure 1B). Within a cell, the cysteine side chain of m^DcTMP is palmitoylated by palmitoyl acyltransferase at the Golgi apparatus, followed by delivery of the palmitoylated m^DcTMP to the PM and Golgi [17,28] (Figure 1A). *Escherichia coli* dihydrofolate reductase (eDHFR), known to bind specifically to TMP, is redistributed to the PM and Golgi by binding to m^DcTMP (Figure 1A). Notably, *dfr1*, a dihydrofolate reductase gene in *S. pombe*, is insensitive to TMP [29], supporting the orthogonality of this SLIPT system in fission yeast. In addition, we have recently developed ^{iK6}DHFR, in which six lysine residues are inserted into the loop region adjacent to the PM of the eDHFR-m^DcTMP complex [21] (Figure 1A). The insertion of the six lysine residues facilitates the selective localization of ^{iK6}DHFR to the PM, but not to the Golgi, through electrostatic interactions with the inner leaflet of the PM.

We investigated whether m^DcTMP could induce relocalization of ^{iK6}DHFR from the cytoplasm to the PM in *S. pombe*. The cells expressing ^{iK6}DHFR fused to green fluorescent protein mNeonGreen (^{iK6}DHFR-mNG) were placed in a microfluidic chamber (Figure S1) and imaged with a spinning disk confocal microscope. ^{iK6}DHFR-mNG was diffusely distributed in the cytoplasm and nucleus before the addition of m^DcTMP (Figure 1C). Addition of 10 μM m^DcTMP from the microfluidic inlet induced redistribution of ^{iK6}DHFR-mNG to the PM within 30 min (Figure 1C and 1D, Movie S1). However, ^{iK6}DHFR-mNG did not remain at the PM for a long time, but rather gradually returned to the cytoplasm within approximately 4 h, even in the presence of m^DcTMP. These results indicate that the m^DcTMP-eDHFR-based SLIPT system is functional in the fission yeast *S. pombe*. Since eDHFR without an iK6 tag is largely translocated to the endoplasmic reticulum (ER) and Golgi upon addition of m^DcTMP (Figure S2), the iK6 tag is essential for m^DcTMP-induced eDHFR translocation to the PM. In addition, this approach would be useful for the transient manipulation of intracellular signaling systems because m^DcTMP-induced PM translocation of ^{iK6}DHFR is transient.

In addition, we tested whether hoeTMP, a self-localizing ligand that recruits DHFR fusion proteins to the nucleus, works in fission yeast. hoeTMP is a small chemical compound consisting of

Hoechst 33258 linked to TMP [15], and is designed so that the Hoechst 33258 moiety binds selectively to

DNA and the DHFR fusion protein is captured in the nucleus via the TMP moiety. We therefore treated

fission yeast cells expressing [iK6]DHFR-mNG with hoeTMP and found that [iK6]DHFR-mNG localized to

the nucleus as expected (Figure S3). These results indicate that hoeTMP functions as a nuclear

localizing ligand in fission yeast.

Figure 1. Self-localizing ligand-induced protein translocation (SLIPT) using m^{D}cTMP-^{iK6}DHFR

in *S. pombe*.

(A) Schematic illustration of the translocation of an mNeonGreen (mNG)-fused hexalysin (K6 tag)-

inserted eDHFR (^{iK6}DHFR-mNG) by m^{D}cTMP from the cytoplasm to the plasma membrane (PM). The

m^DcTMP and ^{iK6}DHFR-mNG complex localizes to the PM through hydrophobic lipid interaction and

electrostatic interaction between the K6 tag and phosphatidylserine on the PM. (B) Molecular structure

of m^DcTMP. (C) Confocal fluorescence images of fission yeast cells expressing ^{iK6}DHFR-mNG were

taken every 5 min. Representative images at the indicated times after addition of m^DcTMP are shown.

Scale bar, 10 μm. (D) Time course of m^DcTMP-induced ^{iK6}DHFR translocation to the PM. Normalized

fluorescence intensity in the cytoplasm was plotted as a function of time. Data are presented as mean ±

SD (n = 7 cells).

Improvement of the SLIPT system for fission yeast

We have previously reported that the amide bonds of the original self-localizing ligand, myristoylated

glycine-cysteine TMP (mgcTMP), are degraded by proteases within 40 min in mammalian cultured

cells, resulting in the spontaneous dissociation of eDHFR from the PM [17]. m^DcTMP, an improved

version of mgcTMP, is able to maintain plasma membrane localization for more than 24 h in cultured

mammalian cells [17], whereas in fission yeast, m^DcTMP-induced PM localization of ^{iK6}DHFR was

diminished within 1-2 h (Figure 1D). These results suggest that m^DcTMP is further degraded by

protease(s) expressed in fission yeast. To address this issue, we newly synthesized m^DcTMP$_{4Me}$, in

which four amides of m^DcTMP are *N*-methylated, in order to render it resistant to the protease(s)

(Figure 2A), as *N*-methylation of amide bonds is known to increase the stability of compounds [19].

Next, we examined whether m^DcTMP$_{4Me}$ could induce plasma membrane localization of

^{iK6}DHFR-mNG in fission yeast and mammalian cells in the manner of m^DcTMP. ^{iK6}DHFR-mNG-

expressing fission yeast cells were applied to a microfluidic chamber and m^DcTMP$_{4Me}$ was added to the

cells. ^{iK6}DHFR-mNG translocated from the cytoplasm to the PM as expected (Figure 2B, Movie S2).

The membrane localization of ^{iK6}DHFR-mNG by m^DcTMP$_{4Me}$ persisted for more than 12 h, though the

signal of ^{iK6}DHFR-mNG on the endomembrane gradually became prominent approximately 4 h after

the addition of m^DcTMP$_{4Me}$. (Figure 2C, Movie S2). The endomembrane localization of the

m^DcTMP$_{4Me}$-bound ^{iK6}DHFR-mNG might be explained by the phosphoatidyl serine-rich surface of the

endomembrane in fission yeast [30]. The membrane localization of ^{iK6}DHFR-mNG by m^DcTMP$_{4Me}$ were

reproduced in yeast cells covered with low melting temperature agarose [31] (Figure S4). Previous studies

have shown that palmitoylation of the cysteine in m^DcTMP is critical for the plasma membrane

selective localization of the m^DcTMP-^{iK6}DHFR-mNG complex [21]. Therefore, we investigated whether

palmitoylation of m^DcTMP$_{4Me}$ is also important for the plasma membrane localization of ^{iK6}DHFR-

mNG by m^DcTMP$_{4Me}$. Fission yeast cells were pretreated with a palmitoylation inhibitor, 2-

bromopalmitate (2-BP), followed by the addition of m^DcTMP_{4Me}. ^{iK6}DHFR-mNG was localized not

only at the plasma membrane but also at endomembranes such as the ER (Figure 2D). This could be

due to the fact that m^DcTMP_{4Me} is localized to the PM and ER only when the myristoyl group of

m^DcTMP_{4Me} remains [17]. Thus, palmitoylation of m^DcTMP_{4Me} is essential for PM-selective localization

of m^DcTMP_{4Me}-^{iK6}DHFR-mNG. Furthermore, treatment of HeLa cells expressing ^{iK6}DHFR-mNG with

m^DcTMP_{4Me} resulted in rapid translocation of ^{iK6}DHFR-mNG to the PM (Figure 2E), indicating that

m^DcTMP_{4Me} is also functional in cultured cells.

Finally, we attempted to restore ^{iK6}DHFR-mNG localization from the PM to the cytoplasm by

adding free TMP in fission yeast, as previously accomplished in cultured cells (Figure 2F) [21]. Fission

yeast cells expressing ^{iK6}DHFR-mNG were applied to a microfluidic chamber and treated with

m^DcTMP_{4Me} for 1 h, followed by the addition of an excess amount of TMP for 3 h (Figure 2G). We

found that ^{iK6}DHFR-mNG localized at the plasma membrane gradually returned to the cytoplasm

(Figure 2G and 2H). These results demonstrate a potential application of the SLIPT system for transient

manipulation of protein localization using free TMP.

Figure 2. Improvement of the SLIPT system for *S. pombe*.

(A) Chemical structure of m^DcTMP_{4Me}. (B and C) Fission yeast cells expressing $^{iK6}DHFR$-mNG were loaded into a microfluidic device and treated with 10 μM m^DcTMP_{4Me}. (B) Confocal images taken at the indicated time points. Scale bar, 10 μm. (C) Time course of m^DcTMP_{4Me}-induced $^{iK6}DHFR$ translocation to the PM. Normalized fluorescence intensity in the cytoplasm was plotted as a function of time. Data are presented as mean ± SD (n = 7 cells). (D) PM recruitment of $^{iK6}DHFR$-mNG in the presence of the palmitoylation inhibitor 2-bromopalmitate (2-BP). Confocal images of fission yeast cells expressing $^{iK6}DHFR$-mNG were taken in the absence of compounds (left) or after treatment with 10 μM m^DcTMP_{4Me} (middle) or both 10 μM m^DcTMP_{4Me} and 100 μM 2-BP (right). Scale bar, 10 μm. (E) HeLa cells stably expressing $^{iK6}DHFR$-mNG were treated with 10 μM m^DcTMP_{4Me}. Confocal fluorescence images were obtained before (left) and 10 min after the addition of m^DcTMP_{4Me}. Scale bar, 10 μm. (F) Schematic of the translocation of PM-localized $^{iK6}DHFR$-mNG to the cytoplasm by competition with free TMP treatment. (G and H) Fission yeast cells expressing $^{iK6}DHFR$-mNG were loaded into a microfluidic device and treated with 10 μM m^DcTMP_{4Me} for 1 h, followed by treatment of 100 μM TMP for 3 h. (G) Confocal fluorescence images of the cells at the indicated time points are shown (lower). Scale bar, 10 μm. (H) Time course of m^DcTMP_{4Me}-induced $^{iK6}DHFR$ translocation to the PM. Normalized fluorescence intensity in the cytoplasm was plotted as a function of time. Data are presented as mean ± SD (n = 10 cells).

Perturbation of endogenous Cdc2 by using the SLIPT system

To demonstrate the potential of the SLIPT system for manipulating signaling pathways in *S. pombe*, we

fused [iK6]DHFR-mNG to endogenous proteins and attempted to perturb cell signaling by tethering the

protein to the plasma membrane with $m^D cTMP_{4Me}$. We focused on cyclin-dependent kinase (CDK)

Cdc2, a key molecule in the cell cycle [32]. Cell cycle progression is regulated by CDK, which is a

nuclear-localized protein and is thought to function primarily in the nucleus [32]. Therefore, we

hypothesized that when endogenous Cdc2 is forced to relocalize to the plasma membrane using

$m^D cTMP_{4Me}$, Cdc2 is excluded from the nucleus, phosphorylation of nuclear substrates is inhibited, and

consequently cell cycle progression is blocked.

We established a fission yeast strain in which the [iK6]DHFR-mNG gene and a selection marker

gene were knocked in at the 3' end of the *cdc2* gene (Cdc2-[iK6]DHFR-mNG) (Figure S5A). The Cdc2-

[iK6]DHFR-mNG strain showed no aberrant cell morphology, and the cell cycle duration was nearly 3

h/cycle in the absence of $m^D cTMP_{4Me}$, similar to that in wild-type cells (Figure 3A and 3B). The cell

length at division was approximately 14.5 μm (Figure 3C), which is almost identical to that of the wild-

type strain, indicating that the knock-in of the [iK6]DHFR-mNG gene into the *cdc2* locus had little effect

on the cell cycle. Next, $m^D cTMP_{4Me}$ was added to the Cdc2-[iK6]DHFR-mNG strain, and as expected,

almost all the Cdc2-[iK6]DHFR-mNG was exported to from the nucleus and localized to the plasma

membrane within 2-4 h (Figure 3D, Movie S3). After the addition of $m^D cTMP_{4Me}$, the cell length

increased in accordance with the prolonged cell cycle duration. Furthermore, several hours after the

addition of $m^D cTMP_{4Me}$, cells with either longer or shorter cell lengths accumulated (Figure 3D and

3E). Notably, longer cells formed multiple septa during cell division (Figure S5B). Finally, the cell

cycle was disrupted, resulting in the accumulation of both cells longer and cells shorter than control

cells (Figure 3F). As a control experiment, we added $m^D cTMP_{4Me}$ to a fission yeast strain stably

expressing [iK6]DHFR-mNG and found that the cell cycle duration and cell length at division were not

altered (Figure S5C and S5D).

When endogenous Cdc2 is forced to be recruited to the plasma membrane using the SLIPT

system, fission yeast cells eventually exhibit two phenotypes: one group of cells that continue to

elongate and another group of cells that maintain a short cell length (Figure 3F). In addition, the

recruitment of Cdc2 to the plasma membrane often caused the formation of multiple septations in

longer cells (Figure S5B). This may have been because the SLIPT system dysregulated the Septation

Initiation Network (SIN) mechanism by recruiting Cdc2 proteins [33]. Of note, the kinetics of plasma

membrane translocation of Cdc2-^{iK6}DHFR-mNG by m^{D}cTMP$_{4Me}$ was slower (2-4 h) than that of

^{iK6}DHFR-mNG (15-30 min) (Figures 2C and 3D). This could be due to the fact that Cdc2-^{iK6}DHFR-

mNG must first be translocated from the nucleus to the cytoplasm, before it can be translocated to the

plasma membrane by m^{D}cTMP$_{4Me}$.

Figure 3. Plasma membrane recruitment of endogenous Cdc2 with the SLIPT system induces cell

cycle arrest in *S. pombe*.

(A-C) Fission yeast cells expressing ^{iK6}DHFR-mNG-fused endogenous Cdc2 were imaged without

m^{D}cTMP$_{4Me}$ in a microfluidic chamber. (A) Confocal fluorescence images (top) and differential

interference contrast (DIC) images (bottom) of fission yeast cells expressing ^{iK6}DHFR-mNG-fused

endogenous Cdc2 were taken at the indicated time points. (B) A heat map showing the temporal profile

of the cell length of fission yeast cells expressing ^{iK6}DHFR-mNG-fused endogenous Cdc2 in the

absence of m^{D}cTMP$_{4Me}$ (n = 10 cells). (C) Cell length in fission yeast cells expressing ^{iK6}DHFR-mNG-

fused endogenous Cdc2 is plotted as a function of time (n = 10 cells). (D-F) Fission yeast cells

expressing ^{iK6}DHFR-mNG-fused endogenous Cdc2 were imaged with m^{D}cTMP$_{4Me}$ in a microfluidic

chamber. (D) Confocal fluorescence images (top) and DIC images (bottom) of fission yeast cells

expressing ^{iK6}DHFR-mNG-fused endogenous Cdc2 were taken at the indicated time points after

m^{D}cTMP$_{4Me}$ treatment. (E) A heat map showing the temporal profile of the cell length of fission yeast

cells expressing ^{iK6}DHFR-mNG-fused endogenous Cdc2 in the presence of m^{D}cTMP$_{4Me}$ (n = 22 cells).

(F) Cell length in fission yeast cells expressing ^{iK6}DHFR-mNG-fused endogenous Cdc2 is plotted as a

function of time after m^{D}cTMP$_{4Me}$ treatment (n = 22 cells). .

Perturbation of cell cycle-related proteins with the SLIPT system in fission yeast.

Finally, we used the SLIPT system to perturb cell cycle-associated proteins and examine their

phenotypes. The genes we targeted for SLIPT are listed in Table 1. We established fission yeast strains

in which the ^{iK6}*DHFR-mNG* gene was knocked in at the 3' end of each target gene. In the absence of

m^DcTMP_{4Me}, no noticeable phenotypes were observed for any of the strains, with the exception that the

Sfi1-^{iK6}DHFR-mNG strain exhibited a slower growth rate than the WT strain. These cell lines were

treated with m^DcTMP_{4Me} (10 μM), incubated for 12 h with shaking, and observed under a fluorescence

microscope (Figure 4A). The phenotypes are summarized in Table 1.

Upon m^DcTMP_{4Me} treatment, Cdc13-^{iK6}DHFR-mNG and Suc1-^{iK6}DHFR-mNG cells showed

shorter cell lengths than WT cells, and multiple septations were observed in some cells (Figure 4A,

white arrowheads). These results were close to the phenotype of m^DcTMP_{4Me}-treated Cdc2-^{iK6}DHFR-

mNG cells, suggesting that endogenous Cdc2 was also recruited to the membrane and caused the

dysregulation of the SIN mechanism. In contrast, no remarkable phenotype was observed by the SLIPT

system for the other Cdc2 regulators, *i.e.*, Cdc25 and Pom1 (Figure 4A). Mcm4, Orc2, Orc4, and

Cdc45, which are associated with S phase progression, showed little phenotype by the SLIPT system.

These results might have been due to insufficient membrane recruitment of these factors as shown in

the mNG fluorescence images (Figure 4A), which implyied that these factors are tightly bound to

histones and/or DNA and therefore are not affected by the SLIPT system. Finally, genes related to

mitosis and cytokinesis were perturbed by SLIPT. m^DcTMP_{4Me} treatment killed Spg1-^{iK6}DHFR-mNG

cells with elongated cell length (Figure 4A). To confirm when and how the Spg1-^{iK6}DHFR-mNG cells

died, we performed time-lapse imaging. We found that m^DcTMP_{4Me} treatment rapidly induced

membrane translocation of Spg1-^{iK6}DHFR-mNG from the spindle pole body (SPB) and cytoplasm,

resulting in sustained cell length elongation, multiple nuclear divisions without cytokinesis, and gradual

cell death (Figure 4B, Movie S4). This result is consistent with the phenotype of the Spg1ts mutant

[34,35]. Upon m^DcTMP_{4Me} treatment, Ark1-^{iK6}DHFR-mNG and Cut2-^{iK6}DHFR-mNG cells exhibited a

mixture of long and short cells, whereas Mal3-^{iK6}DHFR-mNG and Sfi1-^{iK6}DHFR-mNG cells showed a

banana shape and morphological defects, respectively (Figure 4A). On the other hand, no significant

phenotype was observed for the perturbation of Pol1, Alp7, or Alp14 (Figure 4A), due to there being

only partial or no membrane translocation of these proteins by the SLIPT system.

A

Cdc13-
^{iK6}DHFR-mNG   DIC

Cdc25-
^{iK6}DHFR-mNG   DIC

Suc1-
^{iK6}DHFR-mNG   DIC

Pom1-
^{iK6}DHFR-mNG   DIC

Mcm4-
^{iK6}DHFR-mNG   DIC

Orc2-
^{iK6}DHFR-mNG   DIC

Orc4-
^{iK6}DHFR-mNG   DIC

Cdc45-
^{iK6}DHFR-mNG   DIC

Plo1-
^{iK6}DHFR-mNG   DIC

Spg1-
^{iK6}DHFR-mNG   DIC

Ark1-
^{iK6}DHFR-mNG   DIC

Mal3-
^{iK6}DHFR-mNG   DIC

Sfi1-
^{iK6}DHFR-mNG   DIC

Cut2-
^{iK6}DHFR-mNG   DIC

Alp7-
^{iK6}DHFR-mNG   DIC

Alp14-
^{iK6}DHFR-mNG   DIC

B

Figure 4. Perturbation of cell cycle-related proteins using the SLIPT system

(A) Confocal (left) and DIC (right) images of ^{iK6}DHFR-mNG knock-in strains 12 h after treatment with

10 μM m^{D}cTMP$_{4Me}$. (B) Fission yeast cells expressing ^{iK6}DHFR-mNG-fused endogenous Spg1 were

time-lapse imaged with m^DcTMP_{4Me} in a microfluidic chamber. Confocal fluorescence images (top)

and DIC images (bottom) of fission yeast cells expressing [iK6]DHFR-mNG-fused endogenous Spg1

were taken at the indicated time points.

314   In this study, we succeeded in altering the localization of the [iK6]DHFR fusion protein by a self-

localizing ligand in fission yeast cells, and demonstrated that this SLIPT system enables perturbation of

the localization and functions of several molecules associated with the cell cycle. Temperature-

sensitive mutants, which have been widely used in genetic studies of fission yeast, take a long time to

show a phenotype after a change in temperature, whereas the SLIPT system could rapidly perturb the

localization and function of molecules. In addition, the SLIPT system does not require excitation light,

allowing simultaneous fluorescence imaging and multiple perturbations in combination with

optogenetic tools. Compared to other CID systems, such as the FKBP-FRB system, the SLIPT system

in fission yeast has several advantages. First, the FKBP-FRB system requires fusion of FKBP and FRB

to the target and localized molecules, respectively, whereas the SLIPT system only requires fusion of

[iK6]DHFR to the target molecule. Second, unlike the FKBP-FRB system, the SLIPT system can

reversibly manipulate localization [36].

327   Several technical challenges remain for the fission yeast SLIPT system. First, the [iK6]DHFR

fusion molecule is completely localized to the plasma membrane by m^DcTMP in approximately 5 min

in cultured mammalian cells [21], but it took approximately 15-30 min for the [iK6]DHFR-mNG to

translocate to the plasma membrane in fission yeast (Figure 1D and 2C). This result may have been be

due to the lower permeability of localized ligands through the cell wall and/or plasma membrane and

the higher activity of multidrug transporters in fission yeast [37]. This issue could be addressed by

synthesizing ligands with higher cell wall and/or plasma membrane permeability. Second, self-

localizing ligands for intracellular organelles other than the plasma membrane and nucleus have not

been tested in fission yeast. In this study, we developed and tested a localizing ligand that targeted the

DHFR fusion protein to the plasma membrane and nucleus in fission yeast. In cultured cells, self-

localizing ligands for Golgi apparatus, endoplasmic reticulum, and microtubules have been reported

[15,18,19]. Reuse of these ligands will make them more useful chemical and genetic tools in fission yeast.

Third, we found that SLIPT-induced localization changes could not be observed for the factors that

were tightly bound to intracellular organelles and components, such as Mcm4 (Figure 4A). This may be

because the affinity between the self-localizing ligand and the DHFR-fused endogenous molecule was

not sufficient to dissociate the endogenous molecule from the intracellular organelle. Indeed, a previous

report showed that the localization of endogenous Mcm4 was altered in a CID system using FKBP-

FRB [24], which induced irreversible heterodimerization.

Finally, we would like to suggest some future directions for SLIPT systems using fission yeast.

First, more rapid and localized perturbation could be achieved by adding caged moieties to self-

localizing ligands. m^{D}cTMP^{NVOC}, a photoactivatable self-localizing ligand that can uncage m^{D}cTMP

with UV, has already been reported [20,38]. The introduction of four *N*-methylations into m^{D}cTMP^{NVOC}

will allow the molecule to be perturbed with high temporal and spatial resolution by local light

illumination in fission yeast. Secondly, genome-wide perturbation and its phenotypic analysis could be

useful direction for SLIPT. A library of fission yeast strains with the *GFP* gene knocked in at the 3' end

of the endogenous gene is available [39], and thus the expression of DHFR-fused GFP nanobodies [40] with

self-localizing ligands will enable efficient genome-wide manipulation of endogenous molecules. The

fission yeast SLIPT system will provide new opportunities for understanding various biological

phenomena.

Methods

Plasmid construction

All plasmids used in this study are listed in Supplemental Table S1 with Benchling links that provides the sequences and plasmid maps. pNATZA11 [National Bio-Resource Project (NBRP) plasmid FYP2875, provided by Dr. Watanabe (The University of Tokyo) and pFA6a-mNG-kan [NBRP plasmid FYP4997] [41] were used as vector backbones to construct plasmids for the establishment of stable cell lines of the fission yeast *S. pombe*. pCSIIpuro-mNG-eDHFR(69K6) [21] was used as a PCR template and lentiviral vector for the establishment of a stable mammalian cell line. All plasmids were generated using standard cloning procedures and the NEBuilder HiFi DNA assembly system (New England Biolabs).

Culture and transformation of fission yeast *S. pombe*

A wild-type fission yeast (L972) was obtained from the NBRP (FY7507, provided by Dr. Shimoda, Osaka City University). The growth medium and other techniques for fission yeast were based on the protocol described previously [42], unless otherwise noted. The transformation protocol was modified from the previous report [43]. In brief, L972 cells were precultured in 2 mL SD medium [0.67% (w/v) bacto yeast nitrogen base without amino acids (Becton, Dickinson and Co.), 0.5% (w/v) D(+)-glucose] at 32°C. After centrifugation at $860 \times g$, 1 μL of sheared salmon sperm DNA (BioDynamics Lab. Inc.), 10-100 ng of linear DNA encoding the left/right homology arms, and the protein expression cassettes were added. The protein expression cassettes include the cDNA of the protein of interest (POI) and a drug-resistance gene. After resuspending the cells in a mixture of 50 μL LiOAc/TE buffer (100 mM LiOAc, 10 mM Tris-HCl, 1 mM EDTA, pH 7.6) containing 30% (v/v) glycerol and 145 μL LiOAc/TE buffer containing 50% (w/v) polyethylene glycol 4000 (FUJIFILM Wako Pure Chemical Corp.), the suspension was incubated at 42°C for 10 min. After centrifugation at $860 \times g$, the cells were resuspended in 20 μL distilled water and plated on the YEA agar plate [0.5% (w/v) yeast extract (Nacalai Tesque), 3% (w/v) D(+)-glucose, 1.67 mM adenine (FUJIFILM Wako Pure Chemical Corp.), 2% (w/v) agar (Nacalai Tesque)] and incubated at 32°C for at least 12 h. The cells were then transferred to the new YEA agar plate containing antibiotics [100 μg/mL G418 (InvivoGen) or 100 μg/mL nourseothricin (Gold Biotechnology)] and incubated at 32°C for at least 3 days to select the transformants. The expected protein expression in the obtained single clones was confirmed by fluorescence imaging. All cell lines used in this study are listed in Supplemental Table S2.

Live cell imaging

Confocal fluorescence imaging was performed on an IX83 equipped with a UPALXAPO60XO/1.42
NA oil objective lens (Olympus), a UPALXAPO100XO/1.45 NA oil objective lens (Olympus), a Z-
drift compensator system (IX3-ZDC2, Olympus), a sCMOS camera (Orca-Fusion BT; Hamamatsu
Photonics), and a spinning disk confocal unit (CSU-W1; Yokogawa Electric Corp.). For fission yeast
time-lapse imaging, a CellASIC ONIX2 Microfluidic System (CAX2-S0000, Millipore) and a
CellASIC ONIX2 Manifold XT (CAX2-MXT20, Millipore) were used. The microscope was controlled
by MetaMorph software (Molecular Devices). A 488 nm laser was used for excitation of mNeonGreen.
An excitation dichroic mirror (DM405/488/561/640) and emission filter (525/50 for mNeonGreen)
were used for confocal fluorescence imaging (Yokogawa Electric Corp.). Fluorescence images were
analyzed using Fiji/ImageJ [44].

SLIPT assay in fission yeast *S. pombe*

Established cell lines were precultured in YEA medium [0.5%(w/v) yeast extract (Nacalai Tesque),
3%(w/v) D(+)-glucose, 1.67 mM adenine] at 32°C. For time-lapse imaging, the precultured cells were
diluted with fresh YEA medium and loaded into a PDMS-based microfluidic chamber (Y04C-02 or
Y04T-04, Millipore). Cells were monitored by time-lapse imaging before and after starting continuous
loading with m^DcTMP (10 μM), m^DcTMP$_{4Me}$ (10 μM), or TMP (100 μM) at 68.9 kPa.

For perturbation of cell cycle-related proteins, established cell lines were precultured in 2 mL
YEA medium at 32°C, followed by the addition of m^DcTMP$_{4Me}$ (10 μM) and incubation at 32°C for 12
h. For live cell imaging, the cells were seeded on 35 mm glass-base dishes (D11531H, Matsunami
Glass). For the SLIPT assay using m^DcTMP, fission yeast cells expressing eDHFR-mNG were
precultured in 2 mL YEA medium, followed by the addition of m^DcTMP (10 μM) and incubation at
32°C for 30 min. For live cell imaging, the cells were seeded on 35 mm glass-base dishes (Matsunami
Glass). For the SLIPT assay using hoeTMP, fission yeast cells expressing ^{iK6}DHFR-mNG were
precultured in 2 mL YEA medium, followed by the addition of hoeTMP (10 μM) and incubation at
32°C for 60 min. For live cell imaging, the cells were seeded on 35 mm glass-base dishes (Matsunami
Glass).

Palmitoylation assay

Fission yeast cells expressing ^{iK6}DHFR-mNG were precultured in 2 mL YEA medium at 32°C,
followed by the addition of m^DcTMP$_{4Me}$ (10 μM) and incubation at 32°C for 40 min. For

palmitoylation inhibition, cells were treated with 100 μM 2-bromopalmitate (FUJIFILM Wako Pure

Chemical Corp.) and additionally cultured at 32°C for 40 min. For live cell imaging, the cells were

seeded on 35 mm glass-base dishes (Matsunami Glass).

SLIPT assay in fission yeast *S. pombe* on an agarose pad.

Fission yeast cells were preincubated in YEA medium at 32°C and plated at 4.0×10^5 cells in the

central area of 35 mm glass-base dishes coated with poly-L-lysine (Matsunami Glass). After incubation

for 15 min at room temperature, the supernatant was removed, and 100 μL of YEA medium containing

1% (w/v) microwave-heated low-melting point agarose (SeaPlaque agarose, 50101; Lonza) was gently

added to the central area. After resting for 15 minutes at room temperature, the cells were observed by

time-lapse imaging, and 10 μM $m^D cTMP_{4Me}$ in 2 mL YEA was added to the dish.

Mammalian cell culture and establishment of stable cell lines

HeLa cells were obtained from the Cell Resource Center for Biomedical Research, Institute of

Development, Aging and Cancer, Tohoku University. Lenti-X-293T cells were purchased from

Clontech Laboratories. All cells were maintained in Dulbecco's modified Eagle's medium (DMEM)

(Nacalai Tesque) supplemented with 10% heat-inactivated fetal bovine serum (FBS) (Sigma), 100

U/mL penicillin, and 100 μg/mL streptomycin (Nacalai Tesque) at 37°C under a humidified 5% CO_2

atmosphere. For live cell imaging, cells were seeded on 35 mm glass-base dishes (Iwaki) coated with

collagen type I-C (Nitta Gelatin).

For lentiviral production, Lenti-X-293T cells were cotransfected with the pCSIIpuro-mNG-

eDHFR(69K6), psPAX2 (Addgene plasmid #12260, a gift from D. Trono), and pCMV-VSV-G-RSV-

Rev (RIKEN BioResource Research Center plasmid RDB04393, provided by Dr. Miyoshi, Keio

University) plasmids [45] using a polyethlenimine "Max" (Polysciences Inc.). Virus-containing media

were collected 48 h after transfection, filtered, and used to infect target HeLa cells with 8 μg/mL

polybrene (Nacalai Tesque). Infected HeLa cells were selected with 0.5 μg/mL puromycin (InvivoGen)

for at least 7 days. Bulk populations of selected cells were used.

SLIPT assay in mammalian cells

HeLa cells stably expressing mNG-^{iK6}DHFR were plated at 1.5×10^5 cells in collagen-coated 35 mm

glass-base dishes and cultured at 37°C in 5% CO_2 for 24 h. The medium was changed to serum-free

DMEM (Gibco) supplemented with 100 U/mL penicillin and 100 μg/mL streptomycin, and the cells

were observed by time-lapse imaging before and after the addition of $m^D cTMP_{4Me}$ (10 μM).

Author Information

Corresponding Authors

Kazuhiro Aoki - Quantitative Biology Research Group, Exploratory Research Center on Life and Living Systems (ExCELLS), National Institutes of Natural Sciences, 5-1 Higashiyama, Myodaiji-cho, Okazaki, Aichi 444-8787, Japan; Division of Quantitative Biology, National Institute for Basic Biology, National Institutes of Natural Sciences, 5-1 Higashiyama, Myodaiji-cho, Okazaki, Aichi 444-8787, Japan; Department of Basic Biology, School of Life Science, SOKENDAI (The Graduate University for Advanced Studies), 5-1 Higashiyama, Myodaiji-cho, Okazaki, Aichi 444-8787, Japan.

Yuhei Goto - Quantitative Biology Research Group, Exploratory Research Center on Life and Living Systems (ExCELLS), National Institutes of Natural Sciences, 5-1 Higashiyama, Myodaiji-cho, Okazaki, Aichi 444-8787, Japan; Division of Quantitative Biology, National Institute for Basic Biology, National Institutes of Natural Sciences, 5-1 Higashiyama, Myodaiji-cho, Okazaki, Aichi 444-8787, Japan; Department of Basic Biology, School of Life Science, SOKENDAI (The Graduate University for Advanced Studies), 5-1 Higashiyama, Myodaiji-cho, Okazaki, Aichi 444-8787, Japan.

Authors

Akinobu Nakamura - Quantitative Biology Research Group, Exploratory Research Center on Life and Living Systems (ExCELLS), National Institutes of Natural Sciences, 5-1 Higashiyama, Myodaiji-cho, Okazaki, Aichi 444-8787, Japan.

Hironori Sugiyama - Quantitative Biology Research Group, Exploratory Research Center on Life and Living Systems (ExCELLS), National Institutes of Natural Sciences, 5-1 Higashiyama, Myodaiji-cho, Okazaki, Aichi 444-8787, Japan.

Shina Tsukiji - Department of Life Science and Applied Chemistry, Nagoya Institute of Technology, Gokiso-cho, Showa-ku, Nagoya 466-8555, Japan; Department of Nanopharmaceutical Sciences, Nagoya Institute of Technology, Gokiso-cho, Showa-ku, Nagoya 466-8555, Japan.

Author Contributions

A.N.,Y.G., and K.A conceived the study. A.N., and S.T. synthesized m^{D}cTMP, m^{D}cTMP$_{4Me}$, and hoeTMP. A.N., H.S., and Y.G. performed the plasmid construction, transformation and establishment of fission yeast strains, time-lapse imaging, and data analysis. A.N. graphed and analyzed the data. A.N., Y.G., S.T., and K.A. drafted the manuscript. All authors approved the manuscript.

Conflict of Interest

The authors declare no competing financial interest.

Acknowledgments

We thank Dr. Keiko Kuwata (Nagoya University) for HRMS measurement of m^{D}cTMP$_{4Me}$. We thank all members of the Aoki laboratory for their helpful discussions and assistance. Some fission yeast strains were provided by the National Bio-Resource Project (NBRP), Japan. A.N. was supported by JSPS Research Fellowship for Young Scientists (no. JP19J01341). K.A. was supported by a CREST, JST grant (JPMJCR1654) and JSPS KAKENHI grants (nos. JP19H05798 and JP22H02625). Y.G. was supported by a JST, ACT-X Grant Number JPMJAX22B8, JSPS KAKENHI grants (nos. JP19K16050 and JP22K15110), and a Sumitomo Research grant. H.S. was supported by JSPS KAKENHI Grants (nos. JP21J01354 and JP22K15115) and the Dr. Yoshifumi Jigami Memorial Fund, The Society of Yeast Scientists. S.T. was supported by JSPS KAKENHI grants (nos. JP18H04546 and JP20H04706 for the innovative area "Chemistry for Multimolecular Crowding Biosystems", and JP21H05226 Bottom-up Biotech).

References

(1) Grecco, H. E.; Schmick, M.; Bastiaens, P. I. H. Signaling from the Living Plasma Membrane. *Cell* **2011**, *144* (6), 897–909.

(2) Hanahan, D.; Weinberg, R. A. Hallmarks of Cancer: The next Generation. *Cell* **2011**, *144* (5), 646–674.

(3) Sever, R.; Brugge, J. S. Signal Transduction in Cancer. *Cold Spring Harb. Perspect. Med.* **2015**, *5* (4). https://doi.org/10.1101/cshperspect.a006098.

(4) Saveliev, A.; Tybulewicz, V. L. J. Lymphocyte Signaling: Beyond Knockouts. *Nat. Immunol.* **2009**, *10* (4), 361–364.

(5) Viswanathan, R.; Hartmann, J.; Pallares Cartes, C.; De Renzis, S. Desensitisation of Notch Signalling

through Dynamic Adaptation in the Nucleus. *EMBO J.* **2021**, *40* (18), e107245.

(6) Manning, B. D.; Cantley, L. C. AKT/PKB Signaling: Navigating Downstream. *Cell* **2007**, *129* (7), 1261–
1274.

(7) Lavoie, H.; Gagnon, J.; Therrien, M. ERK Signalling: A Master Regulator of Cell Behaviour, Life and Fate.
Nat. Rev. Mol. Cell Biol. **2020**, *21* (10), 607–632.

(8) Inoue, T.; Heo, W. D.; Grimley, J. S.; Wandless, T. J.; Meyer, T. An Inducible Translocation Strategy to
Rapidly Activate and Inhibit Small GTPase Signaling Pathways. *Nat. Methods* **2005**, *2* (6), 415–418.

(9) Inoue, T.; Meyer, T. Synthetic Activation of Endogenous PI3K and Rac Identifies an AND-Gate Switch for
Cell Polarization and Migration. *PLoS One* **2008**, *3* (8), e3068.

(10) Aoki, K.; Yamada, M.; Kunida, K.; Yasuda, S.; Matsuda, M. Processive Phosphorylation of ERK MAP
Kinase in Mammalian Cells. *Proceedings of the National Academy of Sciences* **2011**, *108* (31), 12675–
12680.

(11) Putyrski, M.; Schultz, C. Switching Heterotrimeric G Protein Subunits with a Chemical Dimerizer. *Chem.
Biol.* **2011**, *18* (9), 1126–1133.

(12) Suh, B.-C.; Inoue, T.; Meyer, T.; Hille, B. Rapid Chemically Induced Changes of PtdIns(4,5)P2 Gate
KCNQ Ion Channels. *Science* **2006**, *314* (5804), 1454–1457.

(13) Kopytek, S. J.; Standaert, R. F.; Dyer, J. C. D.; Hu, J. C. Chemically Induced Dimerization of Dihydrofolate
Reductase by a Homobifunctional Dimer of Methotrexate. *Chemistry and Biology* **2000**, *7* (5), 313–321.

(14) Fegan, A.; White, B.; Carlson, J. C. T.; Wagner, C. R. Chemically Controlled Protein Assembly:
Techniques and Applications. *Chem. Rev.* **2010**, *110* (6), 3315–3336.

(15) Ishida, M.; Watanabe, H.; Takigawa, K.; Kurishita, Y.; Oki, C.; Nakamura, A.; Hamachi, I.; Tsukiji, S.
Synthetic Self-Localizing Ligands That Control the Spatial Location of Proteins in Living Cells. *J. Am.
Chem. Soc.* **2013**, *135* (34), 12684–12689.

(16) Nakamura, A.; Oki, C.; Kato, K.; Fujinuma, S.; Maryu, G.; Kuwata, K.; Yoshii, T.; Matsuda, M.; Aoki, K.;
Tsukiji, S. Engineering Orthogonal, Plasma Membrane-Specific SLIPT Systems for Multiplexed Chemical
Control of Signaling Pathways in Living Single Cells. *ACS Chem. Biol.* **2020**, *15* (4), 1004–1015.

(17) Nakamura, A.; Oki, C.; Sawada, S.; Yoshii, T.; Kuwata, K.; Rudd, A. K.; Devaraj, N. K.; Noma, K.;
Tsukiji, S. Designer Palmitoylation Motif-Based Self-Localizing Ligand for Sustained Control of Protein
Localization in Living Cells and Caenorhabditis Elegans. *ACS Chem. Biol.* **2020**, *15* (4), 837–843.

(18) Nakamura, A.; Katahira, R.; Sawada, S.; Shinoda, E.; Kuwata, K.; Yoshii, T.; Tsukiji, S. Chemogenetic
Control of Protein Anchoring to Endomembranes in Living Cells with Lipid-Tethered Small Molecules.
Biochemistry **2020**, *59* (2), 205–211.

(19) Sawada, S.; Nakamura, A.; Yoshii, T.; Kuwata, K.; Nakatsu, F.; Tsukiji, S. Protein-Recruiting Synthetic
Molecules Targeting the Golgi Surface. *Chem. Commun.* **2020**, *56* (98), 15422–15425.

(20) Yoshii, T.; Oki, C.; Watahiki, R.; Nakamura, A.; Tahara, K.; Kuwata, K.; Furuta, T.; Tsukiji, S. Chemo-
Optogenetic Protein Translocation System Using a Photoactivatable Self-Localizing Ligand. *ACS Chem.*

558   *Biol.* **2021**, *16* (8), 1557–1565.

559 (21) Suzuki, S.; Nakamura, A.; Hatano, Y.; Yoshikawa, M.; Yoshii, T.; Sawada, S.; Atsuta-Tsunoda, K.; Aoki,

560   K.; Tsukiji, S. A Chemogenetic Platform for Controlling Plasma Membrane Signaling and Synthetic Signal

561   Oscillation. *Cell Chem Biol* **2022**, *29* (9), 1446–1464.e10.

562 (22) Suzuki, S.; Hatano, Y.; Yoshii, T.; Tsukiji, S. Chemogenetic Control of Protein Localization and

563   Mammalian Cell Signaling by SLIPT. In *Mammalian Cell Engineering: Methods and Protocols*; Kojima,

564   R., Ed.; Springer US: New York, NY, 2021; pp 237–251.

565 (23) Hagan, I. M.; Grallert, A.; Simanis, V. Analysis of the Schizosaccharomyces Pombe Cell Cycle. *Cold*

566   *Spring Harb. Protoc.* **2016**, *2016* (9). https://doi.org/10.1101/pdb.top082800.

567 (24) Ding, L.; Laor, D.; Weisman, R.; Forsburg, S. L. Rapid Regulation of Nuclear Proteins by Rapamycin-

568   Induced Translocation in Fission Yeast. *Yeast* **2014**, *31* (7), 253–264.

569 (25) Weisman, R.; Choder, M.; Koltin, Y. Rapamycin Specifically Interferes with the Developmental Response

570   of Fission Yeast to Starvation. *J. Bacteriol.* **1997**, *179* (20), 6325–6334.

571 (26) Weisman, R.; Finkelstein, S.; Choder, M. Rapamycin Blocks Sexual Development in Fission Yeast through

572   Inhibition of the Cellular Function of an FKBP12 Homolog. *J. Biol. Chem.* **2001**, *276* (27), 24736–24742.

573 (27) Otsubo, Y.; Nakashima, A.; Yamamoto, M.; Yamashita, A. TORC1-Dependent Phosphorylation Targets in

574   Fission Yeast. *Biomolecules* **2017**, *7* (3). https://doi.org/10.3390/biom7030050.

575 (28) Rocks, O.; Gerauer, M.; Vartak, N.; Koch, S.; Huang, Z.-P.; Pechlivanis, M.; Kuhlmann, J.; Brunsveld, L.;

576   Chandra, A.; Ellinger, B.; Waldmann, H.; Bastiaens, P. I. H. The Palmitoylation Machinery Is a Spatially

577   Organizing System for Peripheral Membrane Proteins. *Cell* **2010**, *141* (3), 458–471.

578 (29) Bertani, L. E.; Campbell, J. L. The Isolation and Characterization of the Gene (dfr1) Encoding

579   Dihydrofolate Reductase (DHFR) in Schizosaccharomyces Pombe. *Gene* **1994**, *147* (1), 131–135.

580 (30) Haupt, A.; Minc, N. Gradients of Phosphatidylserine Contribute to Plasma Membrane Charge Localization

581   and Cell Polarity in Fission Yeast. *Mol. Biol. Cell* **2017**, *28* (1), 210–220.

582 (31) Okada, S.; Wloka, C.; Bi, E. Analysis of Protein Dynamics during Cytokinesis in Budding Yeast. In

583   *Methods in Cell Biology*; Echard, A., Ed.; Academic Press, 2017; Vol. 137, pp 25–45.

584 (32) Hochegger, H.; Takeda, S.; Hunt, T. Cyclin-Dependent Kinases and Cell-Cycle Transitions: Does One Fit

585   All? *Nat. Rev. Mol. Cell Biol.* **2008**, *9* (11), 910–916.

586 (33) Guertin, D. A.; Trautmann, S.; McCollum, D. Cytokinesis in Eukaryotes. *Microbiol. Mol. Biol. Rev.* **2002**,

587   *66* (2), 155–178.

588 (34) Schmidt, S.; Sohrmann, M.; Hofmann, K.; Woollard, A.; Simanis, V. The Spg1p GTPase Is an Essential,

589   Dosage-Dependent Inducer of Septum Formation in Schizosaccharomyces Pombe. *Genes Dev.* **1997**, *11*

590   (12), 1519–1534.

591 (35) Li, C.; Furge, K. A.; Cheng, Q. C.; Albright, C. F. Byr4 Localizes to Spindle-Pole Bodies in a Cell Cycle-

592   Regulated Manner to Control Cdc7 Localization and Septation in Fission Yeast. *J. Biol. Chem.* **2000**, *275*

593   (19), 14381–14387.

(36) Lin, Y.-C.; Nihongaki, Y.; Liu, T.-Y.; Razavi, S.; Sato, M.; Inoue, T. Rapidly Reversible Manipulation of
Molecular Activity with Dual Chemical Dimerizers. *Angew. Chem. Int. Ed Engl.* **2013**, *52* (25), 6450–6454.

(37) Wolfger, H.; Mamnun, Y. M.; Kuchler, K. Fungal ABC Proteins: Pleiotropic Drug Resistance, Stress
Response and Cellular Detoxification. *Res. Microbiol.* **2001**, *152* (3-4), 375–389.

(38) Yoshii, T.; Oki, C.; Tsukiji, S. A Photoactivatable Self-Localizing Ligand with Improved Photosensitivity
for Chemo-Optogenetic Control of Protein Localization in Living Cells. *Bioorg. Med. Chem. Lett.* **2022**, *72*,
128865.

(39) Hayashi, A.; Ding, D.-Q.; Tsutsumi, C.; Chikashige, Y.; Masuda, H.; Haraguchi, T.; Hiraoka, Y.
Localization of Gene Products Using a Chromosomally Tagged GFP-Fusion Library in the Fission Yeast
Schizosaccharomyces Pombe. *Genes Cells* **2009**, *14* (2), 217–225.

(40) Kubala, M. H.; Kovtun, O.; Alexandrov, K.; Collins, B. M. Structural and Thermodynamic Analysis of the
GFP:GFP-Nanobody Complex. *Protein Sci.* **2010**, *19* (12), 2389–2401.

(41) Sakai, K.; Kondo, Y.; Fujioka, H.; Kamiya, M.; Aoki, K.; Goto, Y. Near-Infrared Imaging in Fission Yeast
Using a Genetically Encoded Phycocyanobilin Biosynthesis System. *J. Cell Sci.* **2021**, *134* (24).
https://doi.org/10.1242/jcs.259315.

(42) Moreno, S. Molecular Genetic Analyses of Fission Yeast Schizosaccharomyces Pombe. *Methods Enzymol.*
1991, *194*, 773.

(43) Suga, M.; Hatakeyama, T. A Rapid and Simple Procedure for High-Efficiency Lithium Acetate
Transformation of Cryopreserved Schizosaccharomyces Pombe Cells. *Yeast* **2005**, *22* (10), 799–804.

(44) Schindelin, J.; Arganda-Carreras, I.; Frise, E.; Kaynig, V.; Longair, M.; Pietzsch, T.; Preibisch, S.; Rueden,
C.; Saalfeld, S.; Schmid, B.; Tinevez, J. Y.; White, D. J.; Hartenstein, V.; Eliceiri, K.; Tomancak, P.;
Cardona, A. Fiji: An Open-Source Platform for Biological-Image Analysis. *Nat. Methods* **2012**, *9* (7), 676–
682.

(45) Miyoshi, H.; Blömer, U.; Takahashi, M.; Gage, F. H.; Verma, I. M. Development of a Self-Inactivating
Lentivirus Vector. *J. Virol.* **1998**, *72* (10), 8150–8157.

Table 1. Phenotypes induced by the SLIPT system.

Target gene	Product	SLIPT phenotype
cdc13	B-type cyclin	Shorter cell length, multi-septum
cdc25	Phosphatase, M-phase inducer	No obvious phenotype
suc1	CDK regulatory subunit	Shorter cell length, multi-septum
mcm4	MCM complex subunit	No obvious phenotype
orc2	Origin recognition complex subunit	Slightly longer cell length

orc4	Origin recognition complex subunit	Slightly longer cell length
cdc45	DNA replication pre-initiation complex subunit	No obvious phenotype
plo1	Polo kinase	No obvious phenotype
pom1	DYRK family cell polarity protein kinase	No obvious phenotype
spg1	Septum Promoting GTPase	Longer cell length, cell death
ark1	Aurora-B kinase	Longer cell length
mal3	Microtubule plus-end binding protein, EB1 family	Banana shape
sfi1	Spindle pole body half bridge protein	Morphological defect
cut2	Sister chromatid separation inhibitor, securin	Shorter and longer cell length, multi septum
alp7	TACC protein	No obvious phenotype
alp14	TOG/XMAP215 microtubule plus-end tracking polymerase	No obvious phenotype

623